

 Open access • Posted Content • DOI:10.1101/2020.11.18.20233932

REACT-1 round 6 updated report: high prevalence of SARS-CoV-2 swab positivity with reduced rate of growth in England at the start of November 2020

— [Source link](#) 

[Steven Riley](#), [Kylie E. C. Ainslie](#), [Oliver Eales](#), [Caroline E. Walters](#) ...+18 more authors

Institutions: [Imperial College London](#), [Lancaster University](#), [University of Oxford](#), [Imperial College Healthcare](#) ...+1 more institutions

Published on: 12 Nov 2020 - [medRxiv](#) (Cold Spring Harbor Laboratory Press)

Related papers:

- [High prevalence of SARS-CoV-2 swab positivity and increasing R number in England during October 2020: REACT-1 round 6 interim report](#)
- [REACT-1 round 15 interim report: High and rising prevalence of SARS-CoV-2 infection in England from end of September 2021 followed by a fall in late October 2021](#)
- [High prevalence of SARS-CoV-2 swab positivity in England during September 2020: interim report of round 5 of REACT-1 study](#)
- [REACT-1 round 9 interim report: downward trend of SARS-CoV-2 in England in February 2021 but still at high prevalence](#)
- [REACT-1 round 7 interim report: fall in prevalence of swab-positivity in England during national lockdown](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/react-1-round-6-updated-report-high-prevalence-of-sars-cov-2-419gys5ac2>

REACT-1 round 6 updated report: high prevalence of SARS-CoV-2 swab positivity with reduced rate of growth in England at the start of November 2020

Steven Riley^{1,2,*}, Kylie E. C. Ainslie^{1,2}, Oliver Eales^{1,2}, Caroline E. Walters^{1,2}, Haowei Wang^{1,2}, Christina Atchison¹, Claudio Fronterre³, Peter J. Diggle³, Deborah Ashby¹, Christl A. Donnelly^{1,2,4}, Graham Cooke^{5,6,7}, Wendy Barclay⁵, Helen Ward^{1,6,7}, Ara Darzi^{6,7,8}, Paul Elliott^{1,6,7,9,10,11 *}

¹ School of Public Health, Imperial College London, UK

² MRC Centre for Global infectious Disease Analysis and Abdul Latif Jameel Institute for Disease and Emergency Analytics, Imperial College London, UK

³ CHICAS, Lancaster Medical School, Lancaster University, UK and Health Data Research, UK

⁴ Department of Statistics, University of Oxford, UK

⁵ Department of Infectious Disease, Imperial College London, UK

⁶ Imperial College Healthcare NHS Trust, UK

⁷ National Institute for Health Research Imperial Biomedical Research Centre, UK

⁸ Institute of Global Health Innovation at Imperial College London, UK

⁹ MRC Centre for Environment and Health, School of Public Health, Imperial College London, UK

¹⁰ Health Data Research (HDR) UK London at Imperial College

¹¹ UK Dementia Research Institute at Imperial College

*Corresponding authors: Steven Riley and Paul Elliott, s.riley@imperial.ac.uk, p.elliott@imperial.ac.uk, School of Public Health, Imperial College London, Norfolk Place, London, W2 1PG

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

Abstract

Background

England is now in the midst of its second wave of the COVID-19 pandemic. Multiple regions of the country are at high infection prevalence and all areas experienced rapid recent growth of the epidemic during October 2020.

Methods

REACT-1 is a series of community surveys of SARS-CoV-2 RT-PCR swab-positivity in England designed to monitor the spread of the epidemic and thus increase situational awareness. Round 6 of REACT-1 commenced swab-collection on 16th October. A prior interim report included data from 16th to 25th October for 85,971 participants. Here, we report data for the entire round on 160,175 participants with swab results obtained up to 2nd November 2020.

Results

Overall weighted prevalence of infection in the community in England was 1.3% or 130 people per 10,000 infected, up from 60 people per 10,000 in the round 5 report (18th September to 5th October 2020), doubling every 24 days on average since the prior round. The corresponding R number was estimated to be 1.2. Prevalence of infection was highest in North West (2.4%, up from 1.2%), followed by Yorkshire and The Humber (2.3% up from 0.84%), West Midlands (1.6% up from 0.60%), North East (1.5% up from 1.1%), East Midlands (1.3% up from 0.56%), London (0.97%, up from 0.54%), South West (0.80% up from 0.33%), South East (0.69% up from 0.29%), and East of England (0.69% up from 0.30%). Rapid growth in the South observed in the first half of round 6 was no longer apparent in the second half of round 6. We also observed a decline in prevalence in Yorkshire and The Humber during this period. Comparing the first and second halves of round 6, there was a suggestion of decline in weighted prevalence in participants aged 5 to 12 years and in those aged 25 to 44 years. While prevalence remained high, in the second half of round 6 there was suggestion of a slight fall then rise that was seen nationally and also separately in both the North and the South.

Conclusion

The impact of the second national lockdown in England is not yet known. We provide here a detailed description of swab-positivity patterns at national, regional and local scales for the period immediately preceding lockdown, against which future trends in prevalence can be evaluated.

Introduction

England is now in the second wave of the COVID-19 pandemic [1]. Since May 2020, we have been carrying out near real-time surveillance of the epidemic in England through successive rounds of the REal-time Assessment of Community Transmission-1 (REACT-1) study, based on RT-PCR of self-administered swabs [2–7]. We recently reported a rapidly rising prevalence of infections during the first half of round 6, with a doubling time of 9.0 (6.1, 18) days, for swabs collected between 16th and 25th October 2020 [7]. Here, we report results for the complete period of data collection in round 6. This includes swabs obtained up until 2nd November 2020, three days before England entered a second national lockdown [8].

Methods

We have described REACT-1 methods elsewhere [3]. Briefly, we are using RT-PCR to analyse self-administered nose and throat swabs (or parent/guardian administered for children ages 5 to 12 years) obtained from random samples of the population of England from the age of five years upwards. Swabs were maintained on a cold chain until analysed in a single laboratory. The sample was designed to obtain approximately equal numbers of people in each of the 315 lower-tier local authorities (LTLAs) in England, using the National Health Service (NHS) list of GP registered patients to obtain the sample. We have aimed for a sample size of between 120,000 and 160,000 people for each round of data collection, with response rates varying between 22% and 31%.

Once participants provided their swab they were invited to complete a brief health and lifestyle questionnaire. SARS-CoV-2 prevalence (with 95% confidence intervals) was estimated nationally and regionally and by various socio-demographic characteristics, e.g. age. Both unweighted and weighted estimates were obtained, the latter by adjusting for region, deprivation and ethnicity, as well as differential response, so as to be representative of the population in England. We analysed time-trends in swab positivity both between and within rounds using exponential growth and decay models and we carried out multivariable logistic regression to investigate associations of socio-demographic variables and symptoms with swab positivity.

As well as quantifying regional trends, we have investigated SARS-CoV-2 prevalence at sub-regional level, using a geospatial model. The goal of these analyses was to estimate the England-wide geographical variation in the swab-positive prevalence $P(x)$ where x is a lower layer super output area (LSOA) population-weighted centroid. We model the log-odds of $P(x)$ as the sum of a region-level mean and an unobserved, zero-mean spatially correlated stochastic process $S(x)$ that captures local variation around each region-wide mean.

Conditional on $P(x)$ the LSOA-level numbers of positive swab tests are independent and binomially distributed. We estimate the model parameters by Monte Carlo maximum likelihood, then draw samples from the joint predictive distribution of $P(x)$ over all LSOAs, i.e. the joint distribution conditional on the observed numbers of positive swab tests. We then scale-up to LTLA-level by converting the LSOA-level samples to LTLA-level population-weighted averages. All computations used the PrevMap package [9] within the R computing environment [10].

We obtained research ethics approval from the South Central-Berkshire B Research Ethics Committee (IRAS ID: 283787).

Results

We found 1,732 positives from 160,175 swabs giving an unweighted prevalence of 1.08% (95% CI, 1.03%, 1.13%) and a weighted prevalence of 1.30% (1.21%, 1.39%) (Table 1). The weighted prevalence estimate was more than double that of 0.60% (0.55%, 0.71%) obtained in the prior round 5 of the study [4].

The increase in prevalence between this study and the prior round 5 represents a national doubling time of 24 (22, 27) days with a corresponding R estimate of 1.19 (1.17, 1.21) (Figure 1, Table 2). This R estimate from sequential rounds is similar to that reported for the prior round 5 alone of 1.16 (1.05, 1.27). We do not report an R estimate for the current round 6 overall because, using maximum likelihood logistic regression, we found strong evidence in favour of a non-linear prevalence function during this period ($\Delta AIC > 50$, Table 3).

Following the rapid growth reported during the first half of round 6 [7], here we describe little evidence of growth during the second half. Weighted prevalence for the earlier part of round 6 [7] was 1.28% (1.16%, 1.42%) compared with 1.32% (1.20%, 1.45%) for the most recent data with estimates of R of 1.56 (1.27, 1.89) and 0.85 (0.73, 0.99) for each half of round 6 respectively. We investigated the possibility of systematic differences in the participant characteristics between the two halves of round 6 in comparison with the first and second halves of rounds 4 and 5. While we did find differences in characteristics within rounds, patterns were similar across rounds 4, 5 and 6 (not shown).

There appears to be additional temporal structure in the most recent round 6 data not captured by the average trend described above. A p-spline [11] model suggests a fall and then rise in prevalence during this most recent period, with the lowest point around October 30th (Figure 2). A similar pattern was seen when fitting separate curves to data for the North (including Midlands) and the South (Figure 3). Similar patterns were also seen when holding out each region in turn and refitting the p-spline (Figure 4).

At regional level, based on analysis of rounds 5 and 6 together, R was increasing (Figure 5, Table 4, Table 5a). However, consistent with the national trend, the rapid growth in the South during the first half of round 6 was no longer apparent in the more recent data (Table 4, Figure 6). While the highest prevalence in the first half of round 6 was in Yorkshire and The Humber, since then, there has been a decline in prevalence in that region (Figure 6). However, the fall in Yorkshire and The Humber in the most recent period does not explain the national pattern (Figure 4).

Differences in local prevalence between rounds 5 and 6, and between the first and second halves of round 6 (averaged at the level of lower tier local authority, LTLA) reveal sub-regional patterns of growth and decline (Figure 7, Figure 8). For example, decline is suggested in a block of LTLAs in the south of Yorkshire and The Humber and the north of East Midlands, while growth is suggested in contiguous blocks of LTLAs across parts of the North West region.

In the most recent round 6 data, there is suggestion of decline in weighted prevalence of swab positivity in participants aged 5 to 12 years and to a lesser extent in those aged 25 to 44 years (Table 5b, Figure 9). Again in the most recent data, only age and region were robustly associated with increased odds of testing positive among the covariates, with higher odds of swab positivity in 13-17 and 18-24 year olds and in the North and Midlands compared with the reference groups (Figure 10, Table 6). The estimated odds ratios associated with Black and Asian ethnicity and large household size (six or more individuals per household) were closer to unity in round 6 compared with 5.

Discussion

During this sixth round of data collection in the REACT-1 study of SARS-CoV-2 virus prevalence in England, we reported high and rapidly increasing prevalence during the first half of the round (16th to 25th October 2020), with highest prevalence in the North of the country [7]. Here we extend these findings to swabs obtained up to 2nd November, three days before a second national lockdown in England. The data presented here therefore give an assessment of community prevalence prior to the second lockdown in England against which to assess the progression of the epidemic during the lockdown period starting 5th November.

In contrast with our findings for mid- to late-October, we found evidence for a slowdown in the epidemic during the final days of October and beginning of November 2020, with suggestion of a fall and then rise in prevalence during that period. This slowdown was seen across the country, both North and South, and was not being driven by any one region. The

largest falls in prevalence were seen during late October to beginning November in Yorkshire and The Humber, which had previously had the highest prevalence in the country. We also saw reductions in prevalence at the sub-regional level in that region. Falls in prevalence during this period were also observed at the youngest ages in our study (5 to 12 years).

During this period there was evidence of a downturn in daily infections from the national surveillance data on symptomatic cases (“Pillar 1 and 2”) [1] and from the coronavirus symptom app (Zoe app) [12], and a plateau in data from the Office for National Statistics Coronavirus (COVID-19) Infection Survey [13]. Despite differences between these data streams in their recruitment strategy including whether this is influenced by symptom status [1, 12], all four are broadly consistent in identifying an inflection point towards the end of our study period.

The period leading up to the second national lockdown was fluid, with a number of local initiatives being brought in to curb the virus in the highest prevalence areas, mainly in the North of England and the Midlands. At the same time, there was considerable uncertainty and speculation about a possible national lockdown or whether a “circuit-breaker” might be brought in during half-term week which was also at the end of October for most English local authorities. This reflected concerns that the NHS might not cope with increasing hospital admissions that were already being seen in parts of the country. In addition, October 2020 was one of the wettest Octobers on record in England during which the number of sunshine hours was well below average [14] which may have contributed to changes in behaviour and hence transmission. Overall though, it is difficult to ascribe the patterns of prevalence in the latter part of round 6 to any single cause.

Our national prevalence estimate of 1.3% translates to around 1 million infections in England on any one day, assuming sensitivity to detect the virus from a nose and throat swab of around 75% [15]. If we assume that shedding of the virus is detectable for 10 days on average, this would translate to around 100,000 new infections per day at the end of October with a range from 90,000 to 104,000 (reflecting the 95% confidence intervals in weighted national prevalence).

Our study has a number of limitations. In order to estimate trends over time, we assume that the individuals taking part are broadly representative of the base population by LTLA at each time point in the study. We did find evidence for limited differences in population characteristics between the first and second halves of rounds four to six, but not of sufficient size or direction to have materially affected within- or between-round trends. Also it is possible that issues with swab sample transport or changes in laboratory procedures or reagents may have affected either the integrity of the samples or the detection thresholds on

RT-PCR. However, data are collected and analysed according to strict protocols and quality control (QC) procedures, and review of both the delivery chain and laboratory QC did not reveal any differences or discrepancies that might have materially affected positivity rates.

Underlying the national trends, we have described a complex spatial pattern of growth and decline of the epidemic at sub-regional scales in the most recent data. Increased restrictions in west Yorkshire prior to the start of the national lockdown [8] appear to have been successful. In the North West, while there were signs of slowing of the epidemic in the worst affected areas, we did not see substantial groupings of LTLAs with lower prevalence in the second half of round 6 compared with the first. These differences suggest variation in the efficacy of the tiered local interventions which may reflect differences in timing, implementation or population acceptability and adherence to the restrictions.

The impact of the second national lockdown in England is not yet known. We provide here a detailed description of swab-positivity patterns at national, regional and local scales for the period immediately preceding lockdown, against which future trends in prevalence can be evaluated.

Declaration of interests

We declare no competing interests.

Funding

The study was funded by the Department of Health and Social Care in England.

Acknowledgements

SR, CAD acknowledge support: MRC Centre for Global Infectious Disease Analysis, National Institute for Health Research (NIHR) Health Protection Research Unit (HPRU), Wellcome Trust (200861/Z/16/Z, 200187/Z/15/Z), and Centres for Disease Control and Prevention (US, U01CK0005-01-02). GC is supported by an NIHR Professorship. PE is Director of the MRC Centre for Environment and Health (MR/L01341X/1, MR/S019669/1). PE acknowledges support from Health Data Research UK (HDR UK), the NIHR Imperial Biomedical Research Centre and the NIHR HPRUs in Environmental Exposures and Health and Chemical and Radiation Threats and Hazards, the British Heart Foundation Centre for Research Excellence at Imperial College London (RE/18/4/34215) and the UK Dementia Research Institute at Imperial (MC_PC_17114) . We thank The Huo Family Foundation for their support of our work on COVID-19.

We thank key collaborators on this work -- Ipsos MORI: Kelly Beaver, Sam Clemens, Gary Welch, Nicholas Gilby, Andrew Cleary and Kelly Ward; Institute of Global Health Innovation

at Imperial College: Gianluca Fontana, Dr Hutan Ashrafian, Sutha Satkunrajah and Lenny Naar; MRC Centre for Environment and Health, Imperial College London: Daniela Fecht; North West London Pathology and Public Health England for help in calibration of the laboratory analyses; NHS Digital for access to the NHS register; and the Department of Health and Social Care for logistic support. SR acknowledges helpful discussion with attendees of meetings of the UK Government Office for Science (GO-Science) Scientific Pandemic Influenza – Modelling (SPI-M) committee.

References

1. UK Government. UK government Covid-19 dashboard. In: UK government Covid-19 dashboard [Internet]. Available: <https://coronavirus.data.gov.uk/>
2. Riley S, Ainslie KEC, Eales O, Walters CE, Wang H, Atchison CJ, et al. Transient dynamics of SARS-CoV-2 as England exited national lockdown. doi:10.1101/2020.08.05.20169078
3. Riley S, Atchison C, Ashby D, Donnelly CA, Barclay W, Cooke G, et al. REal-time Assessment of Community Transmission (REACT) of SARS-CoV-2 virus: Study protocol. Wellcome Open Research. 2020;5: 200. Available: <https://wellcomeopenresearch.org/articles/5-200/v1?src=rss>
4. Riley S, Ainslie KEC, Eales O, Walters CE, Wang H, Atchison C, et al. High and increasing prevalence of SARS-CoV-2 swab positivity in England during end September beginning October 2020: REACT-1 round 5 updated report. doi:10.1101/2020.10.12.20211227
5. Riley S, Ainslie KEC, Eales O, Jeffrey B, Walters CE, Atchison CJ, et al. Community prevalence of SARS-CoV-2 virus in England during May 2020: REACT study. doi:10.1101/2020.07.10.20150524
6. Riley S, Ainslie KEC, Eales O, Walters CE, Wang H, Atchison C, et al. High prevalence of SARS-CoV-2 swab positivity in England during September 2020: interim report of round 5 of REACT-1 study. doi:10.1101/2020.09.30.20204727
7. Riley S, Ainslie KEC, Eales O, Walters CE, Wang H, Atchison CJ, et al. High prevalence of SARS-CoV-2 swab positivity and increasing R number in England during October 2020: REACT-1 round 6 interim report. medRxiv. 2020. Available: <https://www.medrxiv.org/content/10.1101/2020.10.30.20223123v1.abstract>
8. Government's approach to managing local coronavirus outbreaks. [cited 10 Nov 2020]. Available: <https://www.gov.uk/guidance/governments-approach-to-managing-local-coronavirus-outbreaks>
9. Giorgi E, Diggle PJ, Others. PrevMap: an R package for prevalence mapping. J Stat Softw. 2017;78: 1–29. Available: http://www.lancaster.ac.uk/staff/giorgi/article_revised.pdf
10. R Core Team. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing; 2020. Available: <https://www.R-project.org/>
11. Lang S, Brezger A. Bayesian P-Splines. J Comput Graph Stat. 2004;13: 183–212.

doi:10.1198/1061860043010

12. Sudre CH, Murray B, Varsavsky T, Graham MS, Penfold RS, Bowyer RC, et al. Attributes and predictors of Long-COVID: analysis of COVID cases and their symptoms collected by the Covid Symptoms Study App. *Infectious Diseases (except HIV/AIDS)*. medRxiv; 2020. p. 782. doi:10.1101/2020.10.19.20214494
13. Kara Steel And. Coronavirus (COVID-19) Infection Survey, UK - Office for National Statistics. Office for National Statistics; 6 Nov 2020 [cited 10 Nov 2020]. Available: <https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/conditionsanddiseases/bulletins/coronaviruscovid19infectionsurveypilot/6november2020>
14. Met Office Press Office. UK sees fifth wettest October since 1862. 3 Nov 2020 [cited 11 Nov 2020]. Available: <https://blog.metoffice.gov.uk/2020/11/03/uk-sees-fifth-wettest-october-since-1862/>
15. Ren X, Liu Y, Chen H, Liu W, Guo Z, Zhang Y, et al. Application and Optimization of RT-PCR in Diagnosis of SARS-CoV-2 Infection. 2020. doi:10.2139/ssrn.3546086

Tables and Figures

Supporting data to support tables and figures are available [here](#).

Table 1. Unweighted and weighted prevalence of swab-positivity across six rounds of REACT-1.

Round	Tested swabs	Positive swabs	Unweighted prevalence (95% CI)	Weighted prevalence (95% CI)	First sample	Last sample
1	120,620	159	0.13% (0.11%, 0.15%)	0.16% (0.13%, 0.19%)	1/5/2020	1/6/2020
2	159,199	123	0.077% (0.065%, 0.092%)	0.088% (0.068%, 0.11%)	19/6/2020	7/7/2020
3	162,821	54	0.033% (0.025%, 0.043%)	0.040% (0.027%, 0.053%)	24/7/2020	11/8/2020
4	154,325	137	0.089% (0.075%, 0.11%)	0.13% (0.096%, 0.15%)	20/8/2020	8/9/2020
5	174,949	824	0.47% (0.44%, 0.50%)	0.60% (0.55%, 0.71%)	18/9/2020	5/10/2020
6**	160,175	1732	1.08% (1.03%, 1.13%)	1.30% (1.21%, 1.39%)	16/10/2020	2/11/2020
6a**	85,971	863	1.00% (0.94%, 1.07%)	1.28% (1.16%, 1.42%)	16/10/2020	25/10/2020
6b**	74,210	869	1.17% (1.10%, 1.25%)	1.32% (1.20%, 1.45%)	26/10/2020*	2/11/2020

*Includes small number of samples from prior period

**Total tested swabs for round 6 is less than the sum of rounds 6a and 6b because of participant withdrawal.

Table 2. Estimates of growth rate, doubling time and reproduction number for rounds 5 and 6 together and for rounds 6a and 6b alone.

Data used	Outcome	Growth rate r (1/days)	R number	$p(R>1)$	Doubling (+) / halving (-) time (days)
Rounds 5 and 6	All positives	0.029 (0.026 , 0.032)	1.19 (1.17 , 1.21)	1.000	24.0 (26.5 , 21.8)
	Non-symptomatics	0.030 (0.026 , 0.035)	1.20 (1.17 , 1.23)	1.000	22.9 (26.5 , 20.0)
	Positive for both E and N genes	0.025 (0.021 , 0.028)	1.16 (1.14 , 1.19)	1.000	28.0 (32.4 , 24.7)
	Positive for both E and N genes or positive only for N gene with CT 35 or less	0.028 (0.025 , 0.031)	1.19 (1.16 , 1.21)	1.000	25.0 (28.0 , 22.5)
Round 6a	All positives	0.077 (0.040 , 0.115)	1.56 (1.27 , 1.89)	1.000	9.0 (17.3 , 6.0)
	Non-symptomatics	0.138 (0.082 , 0.191)	2.10 (1.60 , 2.65)	1.000	5.0 (8.4 , 3.6)
	Positive for both E and N genes	0.054 (0.006 , 0.099)	1.38 (1.04 , 1.75)	0.987	12.8 (114.8 , 7.0)
	Positive for both E and N genes or positive only for N gene with CT 35 or less	0.057 (0.017 , 0.097)	1.40 (1.11 , 1.73)	0.997	12.1 (41.7 , 7.1)
Round 6b	All positives	-0.024 (-0.047 , -0.001)	0.85 (0.73 , 0.99)	0.020	-28.4 (-14.8 , *)
	Non-symptomatics	-0.074 (-0.110 , -0.039)	0.59 (0.43 , 0.77)	<0.001	-9.3 (-6.3 , -17.9)
	Positive for both E and N genes	0.000 (-0.028 , 0.028)	1.00 (0.83 , 1.18)	0.507	* (-24.9 , 25.2)
	Positive for both E and N genes or positive only for N gene with CT 35 or less	-0.009 (-0.034 , 0.016)	0.95 (0.80 , 1.11)	0.250	-79.9 (-20.2 , 42.8)

* Doubling and halving times not given when growth rates are very close to 0.

Table 3. Comparison of parsimony of maximum likelihood linear and smooth maximum likelihood logistic regression models of daily swab positivity.

Round	AIC(linear) - AIC(smooth)*
1	2.3
2	1.1
3	0.0
4	7.3
5	0.0
6	54.8

* Akaike information criterion

Table 4. Estimates of growth rate, doubling time and reproduction number for rounds 5 and 6 together and for rounds 6a and 6b for english regions.

Data used	Region	Growth rate r (1/days)	R number	$p(R>1)$	Doubling (+) / halving (-) time (days)
Rounds 5 and 6	South East	0.035 (0.027 , 0.043)	1.24 (1.18 , 1.29)	1.000	19.8 (25.4 , 16.1)
	North East	0.019 (0.008 , 0.030)	1.12 (1.05 , 1.20)	0.999	37.0 (89.7 , 23.2)
	North West	0.025 (0.020 , 0.031)	1.17 (1.13 , 1.21)	1.000	27.2 (35.1 , 22.2)
	Yorkshire and The Humber	0.034 (0.025 , 0.043)	1.23 (1.17 , 1.29)	1.000	20.5 (27.8 , 16.2)
	East Midlands	0.030 (0.023 , 0.038)	1.20 (1.15 , 1.26)	1.000	22.7 (30.5 , 18.1)
	West Midlands	0.041 (0.031 , 0.050)	1.28 (1.21 , 1.35)	1.000	17.0 (22.1 , 13.9)
	East of England	0.026 (0.017 , 0.035)	1.17 (1.11 , 1.24)	1.000	26.7 (41.1 , 19.7)
	London	0.023 (0.014 , 0.033)	1.16 (1.09 , 1.22)	1.000	29.6 (48.7 , 21.1)
	South West	0.029 (0.018 , 0.040)	1.19 (1.12 , 1.28)	1.000	24.1 (38.6 , 17.2)
Round 6a	South East	0.162 (0.052 , 0.271)	2.34 (1.36 , 3.61)	0.999	4.3 (13.4 , 2.6)
	North East	-0.079 (-0.248 , 0.085)	0.57 (0.07 , 1.63)	0.170	-8.8 (-2.8 , 8.1)
	North West	0.031 (-0.049 , 0.110)	1.21 (0.72 , 1.84)	0.774	22.6 (-14.3 , 6.3)
	Yorkshire and The Humber	0.075 (-0.026 , 0.174)	1.54 (0.84 , 2.47)	0.927	9.2 (-27.0 , 4.0)
	East Midlands	0.057 (-0.033 , 0.151)	1.40 (0.80 , 2.23)	0.894	12.1 (-20.9 , 4.6)
	West Midlands	0.107 (-0.007 , 0.220)	1.81 (0.96 , 2.98)	0.968	6.5 (-104.6 , 3.1)
	East of England	0.146 (0.023 , 0.278)	2.18 (1.15 , 3.70)	0.990	4.7 (30.0 , 2.5)
	London	0.210 (0.066 , 0.359)	2.86 (1.47 , 4.87)	0.998	3.3 (10.5 , 1.9)
	South West	0.133 (-0.018 , 0.284)	2.06 (0.89 , 3.79)	0.957	5.2 (-38.3 , 2.4)
Round 6b	South East	-0.018 (-0.074 , 0.037)	0.89 (0.59 , 1.25)	0.264	-38.3 (-9.4 , 18.6)
	North East	0.023 (-0.076 , 0.118)	1.15 (0.58 , 1.91)	0.682	30.0 (-9.1 , 5.9)
	North West	-0.028 (-0.083 , 0.022)	0.83 (0.55 , 1.15)	0.140	-24.8 (-8.4 , 30.9)
	Yorkshire and The Humber	-0.019 (-0.095 , 0.063)	0.88 (0.50 , 1.45)	0.322	-36.6 (-7.3 , 11.0)
	East Midlands	-0.038 (-0.106 , 0.028)	0.77 (0.45 , 1.19)	0.128	-18.2 (-6.5 , 24.6)
	West Midlands	-0.037 (-0.104 , 0.028)	0.78 (0.46 , 1.18)	0.132	-18.7 (-6.6 , 25.1)
	East of England	0.018 (-0.058 , 0.092)	1.12 (0.67 , 1.69)	0.681	38.5 (-12.0 , 7.5)
	London	-0.038 (-0.110 , 0.034)	0.77 (0.43 , 1.23)	0.158	-18.3 (-6.3 , 20.4)
	South West	-0.008 (-0.091 , 0.073)	0.95 (0.51 , 1.53)	0.423	-84.5 (-7.6 , 9.5)

Table 5a. Unweighted and weighted prevalence of swab-positivity by variable and category for rounds 5 and 6.

Variable	Category	Round 5				Round 6			
		Positive	Total	Unweighted Prevalence	Weighted Prevalence	Positive	Total	Unweighted Prevalence	Weighted Prevalence
Gender	Male	387	78,565	0.49% (0.45%, 0.54%)	0.65% (0.56%, 0.74%)	769	72,316	1.06% (0.99%, 1.14%)	1.29% (1.17%, 1.43%)
	Female	437	96,380	0.45% (0.41%, 0.50%)	0.54% (0.47%, 0.61%)	963	87,853	1.10% (1.03%, 1.17%)	1.31% (1.19%, 1.44%)
Age	05-12	59	13,771	0.43% (0.33%, 0.55%)	0.60% (0.40%, 0.80%)	113	11,245	1.00% (0.84%, 1.21%)	1.27% (0.96%, 1.68%)
	13-17	59	10,480	0.56% (0.44%, 0.73%)	0.70% (0.45%, 0.95%)	119	8,616	1.38% (1.16%, 1.65%)	1.72% (1.33%, 2.21%)
	18-24	83	6,852	1.21% (0.98%, 1.50%)	1.59% (1.12%, 2.05%)	120	6,308	1.90% (1.59%, 2.27%)	2.38% (1.79%, 3.16%)
	25-34	86	15,683	0.55% (0.44%, 0.68%)	0.69% (0.50%, 0.87%)	206	15,242	1.35% (1.18%, 1.53%)	1.56% (1.30%, 1.86%)
	35-44	121	22,954	0.53% (0.44%, 0.63%)	0.58% (0.45%, 0.71%)	214	21,004	1.02% (0.89%, 1.16%)	1.25% (1.05%, 1.48%)
	45-54	157	29,380	0.53% (0.46%, 0.62%)	0.58% (0.47%, 0.69%)	314	26,898	1.17% (1.05%, 1.30%)	1.26% (1.10%, 1.44%)
	55-64	116	31,885	0.36% (0.30%, 0.44%)	0.37% (0.29%, 0.45%)	326	29,752	1.10% (0.98%, 1.22%)	1.20% (1.05%, 1.38%)
	65+	143	43,944	0.33% (0.28%, 0.38%)	0.35% (0.27%, 0.42%)	320	41,110	0.78% (0.70%, 0.87%)	0.82% (0.71%, 0.94%)
Region	South East	104	39,496	0.26% (0.22%, 0.32%)	0.29% (0.23%, 0.36%)	244	36,164	0.67% (0.60%, 0.76%)	0.69% (0.60%, 0.80%)
	North East	61	6,702	0.91% (0.71%, 1.17%)	1.06% (0.76%, 1.37%)	89	5,846	1.52% (1.24%, 1.87%)	1.49% (1.17%, 1.89%)
	North West	199	19,468	1.02% (0.89%, 1.17%)	1.21% (0.99%, 1.43%)	372	17,866	2.08% (1.88%, 2.30%)	2.39% (2.10%, 2.73%)
	Yorkshire and The Humber	74	11,658	0.63% (0.51%, 0.80%)	0.84% (0.56%, 1.12%)	189	10,697	1.77% (1.53%, 2.03%)	2.32% (1.90%, 2.81%)
	East Midlands	101	21,897	0.46% (0.38%, 0.56%)	0.56% (0.42%, 0.70%)	233	20,458	1.14% (1.00%, 1.29%)	1.25% (1.07%, 1.45%)
	West Midlands	70	16,343	0.43% (0.34%, 0.54%)	0.60% (0.36%, 0.83%)	196	14,861	1.32% (1.15%, 1.52%)	1.59% (1.18%, 2.13%)
	East of England	81	25,625	0.32% (0.25%, 0.39%)	0.30% (0.22%, 0.37%)	150	23,231	0.65% (0.55%, 0.76%)	0.69% (0.57%, 0.83%)
	London	79	17,500	0.45% (0.36%, 0.56%)	0.54% (0.40%, 0.68%)	141	15,757	0.89% (0.76%, 1.05%)	0.97% (0.79%, 1.18%)
	South West	55	16,260	0.34% (0.26%, 0.44%)	0.33% (0.22%, 0.44%)	118	15,295	0.77% (0.64%, 0.92%)	0.80% (0.64%, 0.99%)
		Health care or care home worker	51	9,620	0.53% (0.40%, 0.70%)	0.66% (0.46%, 0.94%)	124	9,257	1.34% (1.12%, 1.59%)
	Other essential/key worker	154	28,846	0.53% (0.46%, 0.62%)	0.70% (0.55%, 0.88%)	357	26,894	1.33% (1.20%, 1.47%)	1.64% (1.41%, 1.92%)
	Other worker	338	72,175	0.47% (0.42%, 0.52%)	0.56% (0.48%, 0.65%)	677	64,811	1.04% (0.97%, 1.13%)	1.22% (1.10%, 1.37%)
	Not full-time, part-time, or self-employed	256	59,762	0.43% (0.38%, 0.48%)	0.58% (0.49%, 0.69%)	520	55,445	0.94% (0.86%, 1.02%)	1.11% (0.98%, 1.25%)
Ethnic group	White	707	157,645	0.45% (0.42%, 0.48%)	0.54% (0.48%, 0.59%)	1550	145,336	1.07% (1.02%, 1.12%)	1.24% (1.16%, 1.33%)
	Asian	71	7,907	0.90% (0.71%, 1.13%)	1.11% (0.79%, 1.43%)	84	6,712	1.25% (1.01%, 1.55%)	2.05% (1.45%, 2.88%)
	Black	16	2,194	0.73% (0.45%, 1.18%)	1.17% (0.48%, 1.86%)	23	1,760	1.31% (0.87%, 1.95%)	1.23% (0.76%, 1.98%)
	Mixed	11	3,080	0.36% (0.20%, 0.64%)	0.44% (0.14%, 0.74%)	28	2,697	1.04% (0.72%, 1.50%)	1.01% (0.66%, 1.53%)
	Other	7	1,439	0.49% (0.24%, 1.00%)	0.47% (0.09%, 0.85%)	17	1,184	1.44% (0.90%, 2.29%)	1.81% (1.00%, 3.24%)
Household size	1	88	24,916	0.35% (0.29%, 0.43%)	0.44% (0.30%, 0.57%)	228	23,639	0.96% (0.85%, 1.10%)	1.08% (0.91%, 1.29%)
	2	227	61,717	0.37% (0.32%, 0.42%)	0.44% (0.36%, 0.52%)	553	59,221	0.93% (0.86%, 1.01%)	1.13% (0.99%, 1.28%)
	3	172	32,395	0.53% (0.46%, 0.62%)	0.61% (0.50%, 0.73%)	359	30,043	1.20% (1.08%, 1.32%)	1.47% (1.25%, 1.73%)
	4	211	37,231	0.57% (0.50%, 0.65%)	0.75% (0.59%, 0.91%)	349	31,911	1.09% (0.99%, 1.21%)	1.30% (1.11%, 1.52%)
	5	77	13,145	0.59% (0.47%, 0.73%)	0.76% (0.53%, 0.99%)	172	10,795	1.59% (1.37%, 1.85%)	1.73% (1.44%, 2.08%)
	6	31	3,845	0.81% (0.57%, 1.14%)	0.85% (0.51%, 1.19%)	50	3,159	1.58% (1.20%, 2.08%)	2.16% (1.49%, 3.12%)
	7+	18	1,700	1.06% (0.67%, 1.67%)	1.42% (0.64%, 2.21%)	21	1,407	1.49% (0.98%, 2.27%)	1.12% (0.67%, 1.86%)
COVID case contact	No	556	139,853	0.40% (0.37%, 0.43%)	0.50% (0.44%, 0.55%)	1131	128,070	0.88% (0.83%, 0.94%)	1.02% (0.94%, 1.11%)
	Yes, contact with a confirmed/tested COVID-19 case	102	1,566	6.51% (5.39%, 7.85%)	8.20% (6.08%, 10.32%)	301	3,621	8.31% (7.46%, 9.26%)	8.92% (7.67%, 10.35%)
	Yes, contact with a suspected COVID-19 case	17	1,487	1.14% (0.72%, 1.82%)	1.09% (0.52%, 1.66%)	50	1,669	3.00% (2.28%, 3.93%)	3.55% (2.26%, 5.53%)
Symptom status	Classic COVID symptoms	187	8,376	2.23% (1.94%, 2.57%)	2.75% (2.20%, 3.30%)	423	6,284	6.73% (6.14%, 7.38%)	7.51% (6.69%, 8.49%)
	Other symptoms	146	25,832	0.57% (0.48%, 0.66%)	0.70% (0.55%, 0.86%)	266	20,739	1.28% (1.14%, 1.45%)	1.35% (1.17%, 1.57%)
	No symptoms	344	108,910	0.32% (0.28%, 0.35%)	0.39% (0.34%, 0.45%)	798	106,439	0.75% (0.70%, 0.80%)	0.90% (0.81%, 1.01%)
Deprivation	1 Most deprived	135	17,641	0.77% (0.65%, 0.91%)	1.00% (0.80%, 1.26%)	254	15,736	1.61% (1.43%, 1.82%)	2.09% (1.74%, 2.50%)
	2	129	27,483	0.47% (0.40%, 0.56%)	0.53% (0.43%, 0.65%)	289	24,957	1.16% (1.03%, 1.30%)	1.27% (1.11%, 1.46%)
	3	146	37,427	0.39% (0.33%, 0.46%)	0.48% (0.38%, 0.59%)	354	34,379	1.03% (0.93%, 1.14%)	1.09% (0.96%, 1.24%)
	4	190	43,109	0.44% (0.38%, 0.51%)	0.49% (0.41%, 0.58%)	407	39,792	1.02% (0.93%, 1.13%)	1.10% (0.98%, 1.25%)
	5 Least deprived	224	49,289	0.45% (0.40%, 0.52%)	0.52% (0.44%, 0.61%)	428	45,311	0.94% (0.86%, 1.04%)	1.02% (0.91%, 1.15%)

Table 5b. Unweighted and weighted prevalence of swab-positivity by variable and category for rounds 6a and 6b.

Variable	Category	Round 6a				Round 6b			
		Positive	Total	Unweighted Prevalence	Weighted Prevalence	Positive	Total	Unweighted Prevalence	Weighted Prevalence
Gender	Male	387	39,319	0.98% (0.89%, 1.09%)	1.31% (1.13%, 1.53%)	382	33,000	1.16% (1.05%, 1.28%)	1.27% (1.11%, 1.46%)
	Female	476	46,648	1.02% (0.93%, 1.12%)	1.25% (1.10%, 1.43%)	487	41,208	1.18% (1.08%, 1.29%)	1.37% (1.20%, 1.57%)
Age	05-12	55	5,513	1.00% (0.77%, 1.30%)	1.49% (0.98%, 2.26%)	58	5,732	1.01% (0.78%, 1.31%)	1.07% (0.77%, 1.48%)
	13-17	60	5,169	1.16% (0.90%, 1.49%)	1.58% (1.08%, 2.31%)	59	3,447	1.71% (1.33%, 2.20%)	1.92% (1.40%, 2.63%)
	18-24	53	3,091	1.71% (1.31%, 2.24%)	2.25% (1.47%, 3.42%)	67	3,217	2.08% (1.64%, 2.64%)	2.51% (1.71%, 3.67%)
	25-34	97	7,186	1.35% (1.11%, 1.64%)	1.62% (1.25%, 2.08%)	109	8,056	1.35% (1.12%, 1.63%)	1.50% (1.18%, 1.92%)
	35-44	98	10,079	0.97% (0.80%, 1.18%)	1.29% (0.98%, 1.68%)	116	10,925	1.06% (0.89%, 1.27%)	1.22% (0.99%, 1.51%)
	45-54	137	13,009	1.05% (0.89%, 1.24%)	1.22% (0.99%, 1.51%)	177	13,889	1.27% (1.10%, 1.47%)	1.29% (1.08%, 1.54%)
	55-64	167	15,768	1.06% (0.91%, 1.23%)	1.20% (0.99%, 1.46%)	159	13,984	1.14% (0.97%, 1.33%)	1.21% (0.99%, 1.47%)
65+	196	26,156	0.75% (0.65%, 0.86%)	0.81% (0.68%, 0.96%)	124	14,960	0.83% (0.70%, 0.99%)	0.83% (0.67%, 1.03%)	
Region	South East	106	19,367	0.55% (0.45%, 0.66%)	0.55% (0.45%, 0.68%)	138	16,797	0.82% (0.70%, 0.97%)	0.84% (0.70%, 1.02%)
	North East	45	3,323	1.35% (1.01%, 1.81%)	1.17% (0.84%, 1.64%)	44	2,524	1.74% (1.30%, 2.33%)	1.88% (1.33%, 2.66%)
	North West	192	10,035	1.91% (1.66%, 2.20%)	2.27% (1.90%, 2.72%)	180	7,831	2.30% (1.99%, 2.65%)	2.53% (2.09%, 3.06%)
	Yorkshire and The Humber	113	6,179	1.83% (1.52%, 2.19%)	2.72% (2.12%, 3.50%)	76	4,518	1.68% (1.35%, 2.10%)	1.80% (1.33%, 2.44%)
	East Midlands	128	11,461	1.12% (0.94%, 1.33%)	1.20% (0.98%, 1.47%)	105	8,997	1.17% (0.97%, 1.41%)	1.31% (1.04%, 1.63%)
	West Midlands	90	8,005	1.12% (0.92%, 1.38%)	1.62% (1.05%, 2.49%)	106	6,856	1.55% (1.28%, 1.87%)	1.56% (1.06%, 2.30%)
	East of England	74	12,612	0.59% (0.47%, 0.74%)	0.64% (0.48%, 0.85%)	76	10,620	0.72% (0.57%, 0.89%)	0.74% (0.57%, 0.96%)
London	London	61	7,299	0.84% (0.65%, 1.07%)	0.89% (0.66%, 1.19%)	80	8,459	0.95% (0.76%, 1.18%)	1.03% (0.79%, 1.35%)
	South West	54	7,690	0.70% (0.54%, 0.92%)	0.72% (0.52%, 0.99%)	64	7,608	0.84% (0.66%, 1.07%)	0.88% (0.64%, 1.20%)
Employment type	Health care or care home worker	54	4,069	1.33% (1.02%, 1.73%)	2.19% (1.35%, 3.54%)	70	5,212	1.34% (1.06%, 1.69%)	1.47% (1.09%, 1.98%)
	Other essential/key worker	169	12,503	1.35% (1.16%, 1.57%)	1.86% (1.46%, 2.36%)	188	14,427	1.30% (1.13%, 1.50%)	1.46% (1.21%, 1.77%)
	Other worker	366	34,262	1.07% (0.96%, 1.18%)	1.30% (1.13%, 1.50%)	345	33,018	1.04% (0.94%, 1.16%)	1.14% (0.97%, 1.35%)
	Not full-time, part-time, or self-employed	256	33,136	0.77% (0.68%, 0.87%)	0.91% (0.76%, 1.08%)	228	19,547	1.17% (1.03%, 1.33%)	1.41% (1.18%, 1.68%)
Ethnic group	White	793	79,406	1.00% (0.93%, 1.07%)	1.24% (1.12%, 1.37%)	757	65,936	1.15% (1.07%, 1.23%)	1.25% (1.14%, 1.36%)
	Asian	35	2,863	1.22% (0.88%, 1.70%)	2.06% (1.22%, 3.48%)	49	3,849	1.27% (0.96%, 1.68%)	2.03% (1.29%, 3.19%)
	Black	7	603	1.16% (0.56%, 2.38%)	1.60% (0.72%, 3.52%)	16	1,157	1.38% (0.85%, 2.23%)	1.02% (0.57%, 1.84%)
	Mixed	14	1,326	1.06% (0.63%, 1.76%)	1.03% (0.57%, 1.83%)	14	1,371	1.02% (0.61%, 1.71%)	0.99% (0.53%, 1.82%)
Other	6	536	1.12% (0.51%, 2.42%)	1.34% (0.54%, 3.31%)	11	648	1.70% (0.95%, 3.01%)	2.19% (1.03%, 4.63%)	
Household size	1	106	12,368	0.86% (0.71%, 1.04%)	0.92% (0.72%, 1.17%)	122	11,273	1.08% (0.91%, 1.29%)	1.25% (0.98%, 1.61%)
	2	306	34,116	0.90% (0.80%, 1.00%)	1.12% (0.96%, 1.30%)	247	25,109	0.98% (0.87%, 1.11%)	1.14% (0.91%, 1.41%)
	3	180	15,409	1.17% (1.01%, 1.35%)	1.59% (1.24%, 2.04%)	179	14,634	1.22% (1.06%, 1.41%)	1.35% (1.11%, 1.65%)
	4	161	16,329	0.99% (0.85%, 1.15%)	1.36% (1.05%, 1.75%)	188	15,582	1.21% (1.05%, 1.39%)	1.25% (1.05%, 1.48%)
	5	76	5,438	1.40% (1.12%, 1.75%)	1.42% (1.08%, 1.85%)	96	5,357	1.79% (1.47%, 2.18%)	2.03% (1.58%, 2.61%)
	6	27	1,632	1.65% (1.14%, 2.40%)	2.37% (1.40%, 4.00%)	23	1,527	1.51% (1.01%, 2.25%)	1.94% (1.16%, 3.22%)
	7+	7	679	1.03% (0.50%, 2.11%)	0.96% (0.39%, 2.34%)	14	728	1.92% (1.15%, 3.20%)	1.27% (0.69%, 2.30%)
COVID case contact	No	529	67,864	0.78% (0.72%, 0.85%)	1.23% (1.10%, 1.37%)	541	54,514	0.99% (0.91%, 1.08%)	1.38% (1.24%, 1.54%)
	Yes, contact with a confirmed/tested COVID-19 case	147	1,787	8.23% (7.04%, 9.59%)	9.56% (7.71%, 11.80%)	139	1,666	8.34% (7.11%, 9.77%)	8.40% (6.70%, 10.49%)
	Yes, contact with a suspected COVID-19 case	22	842	2.61% (1.73%, 3.92%)	4.25% (2.16%, 8.19%)	28	754	3.71% (2.58%, 5.31%)	3.08% (1.95%, 4.84%)
Symptom status	Classic COVID symptoms	201	3,381	5.95% (5.20%, 6.79%)	6.91% (5.78%, 8.25%)	196	2,646	7.41% (6.47%, 8.47%)	8.00% (6.66%, 9.59%)
	Other symptoms	119	11,430	1.04% (0.87%, 1.24%)	1.12% (0.90%, 1.41%)	138	8,623	1.60% (1.36%, 1.89%)	1.62% (1.32%, 2.00%)
	No symptoms	380	55,746	0.68% (0.62%, 0.75%)	0.90% (0.75%, 1.06%)	377	45,716	0.82% (0.75%, 0.91%)	0.93% (0.81%, 1.07%)
Deprivation	1 Most deprived	119	7,919	1.50% (1.26%, 1.80%)	2.16% (1.66%, 2.81%)	135	7,819	1.73% (1.46%, 2.04%)	2.01% (1.58%, 2.56%)
	2	143	12,750	1.12% (0.95%, 1.32%)	1.32% (1.07%, 1.61%)	146	12,209	1.20% (1.02%, 1.40%)	1.23% (1.02%, 1.49%)
	3	176	18,292	0.96% (0.83%, 1.11%)	1.11% (0.92%, 1.33%)	178	16,088	1.11% (0.96%, 1.28%)	1.08% (0.91%, 1.28%)
	4	199	21,749	0.92% (0.80%, 1.05%)	1.02% (0.86%, 1.21%)	208	18,043	1.15% (1.01%, 1.32%)	1.21% (1.02%, 1.43%)
	5 Least deprived	226	25,261	0.89% (0.79%, 1.02%)	0.97% (0.83%, 1.14%)	202	20,051	1.01% (0.88%, 1.16%)	1.09% (0.93%, 1.28%)

Table 6. Estimated odds ratios and 95% confidence intervals for jointly adjusted logistic regression model of swab-positivity for rounds 5, 6, 6a (16 to 25 October 2020), and 6b (26 October to 2 November 2020).

Variable	Category	Round 5	Round 6	Round 6a	Round 6b
Gender	Male	Reference			
	Female	0.89 [0.77,1.03]	1.01 [0.91,1.11]	1.03 [0.90,1.19]	0.98 [0.85,1.13]
Age Group	5 -12	0.72 [0.52,1.00]	0.93 [0.73,1.17]	1.00 [0.72,1.40]	0.86 [0.62,1.20]
	13-17	1.01 [0.70,1.47]	1.54 [1.18,1.99]	1.63 [1.13,2.35]	1.59 [1.10,2.30]
	18-24	2.24 [1.68,2.99]	1.86 [1.48,2.35]	1.79 [1.27,2.52]	1.93 [1.41,2.64]
	25-34	1.10 [0.83,1.47]	1.34 [1.10,1.64]	1.42 [1.07,1.88]	1.28 [0.98,1.69]
	35-44	Reference			
	45-54	1.09 [0.85,1.38]	1.14 [0.95,1.36]	1.09 [0.84,1.42]	1.18 [0.93,1.51]
	55-64	0.79 [0.60,1.03]	1.14 [0.95,1.37]	1.21 [0.93,1.58]	1.09 [0.84,1.42]
	65+	0.76 [0.57,1.02]	0.93 [0.75,1.14]	1.04 [0.78,1.39]	0.86 [0.64,1.16]
Region	North East	3.54 [2.55,4.91]	2.20 [1.70,2.83]	2.33 [1.61,3.36]	2.09 [1.46,2.98]
	North West	3.91 [3.05,5.02]	3.11 [2.62,3.68]	3.51 [2.74,4.49]	2.80 [2.21,3.54]
	Yorkshire and The Humber	2.53 [1.86,3.44]	2.69 [2.21,3.27]	3.43 [2.61,4.51]	2.09 [1.56,2.80]
	East Midlands	1.80 [1.36,2.39]	1.68 [1.39,2.02]	2.04 [1.56,2.65]	1.39 [1.06,1.81]
	West Midlands	1.65 [1.21,2.25]	1.86 [1.53,2.27]	2.09 [1.57,2.79]	1.68 [1.28,2.21]
	East of England	1.20 [0.89,1.62]	0.93 [0.75,1.15]	1.07 [0.79,1.45]	0.82 [0.61,1.10]
	London	1.48 [1.09,2.02]	1.22 [0.98,1.52]	1.36 [0.98,1.90]	1.08 [0.80,1.45]
	South East	Reference			
South West	1.34 [0.96,1.89]	1.20 [0.96,1.50]	1.36 [0.97,1.89]	1.07 [0.79,1.45]	
Key Worker Status	Health care or care home worker	1.03 [0.76,1.39]	1.14 [0.94,1.40]	1.12 [0.83,1.50]	1.15 [0.87,1.50]
	Other essential/key worker	1.03 [0.85,1.25]	1.16 [1.01,1.32]	1.17 [0.97,1.41]	1.12 [0.94,1.35]
	Other worker	Reference			
	Not full-time, part-time, or self-employed	1.02 [0.84,1.23]	0.92 [0.81,1.05]	0.71 [0.60,0.86]	1.15 [0.95,1.39]
Ethnicity	Asian	1.88 [1.44,2.45]	1.15 [0.91,1.46]	1.22 [0.85,1.75]	1.09 [0.80,1.49]
	Black	1.67 [1.00,2.78]	1.31 [0.86,1.99]	1.24 [0.58,2.64]	1.32 [0.79,2.20]
	Mixed	0.77 [0.41,1.44]	0.94 [0.64,1.40]	1.02 [0.59,1.78]	0.89 [0.51,1.55]
	Other	1.14 [0.54,2.42]	1.52 [0.94,2.48]	1.26 [0.56,2.85]	1.71 [0.93,3.14]
	White	Reference			
Household Size	1-2 People	Reference			
	3-5 People	1.28 [1.08,1.53]	1.20 [1.07,1.35]	1.17 [0.99,1.39]	1.23 [1.04,1.46]
	6+ People	1.91 [1.36,2.68]	1.41 [1.07,1.85]	1.46 [0.99,2.16]	1.33 [0.90,1.97]
Deprivation Index Quintile	1 - Most Deprived	Reference			
	2	0.80 [0.62,1.03]	0.87 [0.73,1.04]	0.88 [0.68,1.13]	0.87 [0.68,1.12]
	3	0.77 [0.60,0.98]	0.83 [0.70,0.99]	0.81 [0.64,1.03]	0.86 [0.68,1.10]
	4	0.84 [0.66,1.06]	0.82 [0.69,0.97]	0.77 [0.61,0.97]	0.88 [0.69,1.11]
	5 - Least Deprived	0.93 [0.74,1.17]	0.81 [0.69,0.96]	0.82 [0.65,1.04]	0.79 [0.62,1.00]




Figure 1. Constant growth rate models fit to REACT-1 data for sequential and individual rounds. Points show unweighted prevalence estimate and vertical lines show 95% binomial confidence intervals (CIs). Because there were few swabs taken on some days, some upper bounds to CIs are truncated. Models fit to REACT-1 data for sequential rounds; 1 and 2 (yellow), 2 and 3 (blue), 3 and 4 (green), 4 and 5 (pink), and 5 and 6 (purple). Shaded areas show 95% credible intervals. Note that of the 932,171 swab tests only 918,543 had a date and so were included in the temporal analysis (3,008 of 3,029 positives).




Figure 2. Prevalence of swab-positivity estimated using a p-spline for the full period of the study with central 50% and 95% posterior credible intervals. Points show unweighted prevalence estimate and vertical lines show 95% binomial confidence intervals (CIs). Because there were few swabs taken on some days, some upper bounds to CIs are truncated. Note that of the 932,171 swab tests only 918,543 had a date and so were included in the temporal analysis (3,008 of 3,029 positives).




Figure 3. Prevalence of swab-positivity estimated using a p-spline for the full period of the study with central 50% and 95% posterior credible intervals. Points show unweighted prevalence estimate and vertical lines show 95% binomial confidence intervals (CIs). Because there were few swabs taken on some days, some upper bounds to CIs are truncated. **A** The model fit to a subset of the data containing only the regions North East, North West, Yorkshire and The Humber, East Midlands and West Midlands. **B** The model fit to a subset of the data only containing the regions London, East of England, South East and South West. Note that of the 932,171 swab tests 918,543 had a date and so were included in the temporal analysis (3,008 of 3,029 positives).




Figure 4. Prevalence of swab-positivity estimated using a p-spline for the full period of the study with central 95% posterior credible intervals for subsets of the data leaving out one region at a time from the analysis. The black dashed line shows the p-spline fit to all available data. Points show unweighted prevalence estimate and vertical lines show 95% binomial confidence intervals (CIs). Because there were few swabs taken on some days, some upper bounds to CIs are truncated. Note that of the 932,171 swab tests 918,543 had a date and so were included in the temporal analysis (3,008 of 3,029 positives).




Figure 5. Constant growth rate models fit to regions for REACT-1 data for sequential rounds; 1 and 2 (yellow), 2 and 3 (blue), 3 and 4 (green), 4 and 5 (pink), and 5 and 6 (purple). Points show unweighted prevalence estimate and vertical lines show 95% binomial confidence intervals (CIs). Because there were few swabs taken on some days, some upper bounds to CIs are truncated. Shaded regions show 95% posterior credible intervals for growth models. Note that of the 932,171 swab tests 918,543 had a date and so were included in the temporal analysis (3,008 of 3,029 positives).




Figure 6. Weighted prevalence of swab positivity by region for rounds 5, 6, 6a and 6b. Bars show 95% confidence intervals.




Figure 7. Modelled prevalence at lower tier local authority level (see Methods) for rounds 5, 6, 6a and 6b. Regions: NE = North East, NW = North West, YH = Yorkshire and The Humber, EM = East Midlands, WM = West Midlands, EE = East of England, L = London, SE = South East, SW = South West.




Figure 8. Difference in modelled prevalence at lower tier local authority level (see Methods) between round 6 and round 5, and between rounds 6b and 6a. Regions: NE = North East, NW = North West, YH = Yorkshire and The Humber, EM = East Midlands, WM = West Midlands, EE = East of England, L = London, SE = South East, SW = South West.




Figure 9. Weighted prevalence and 95% confidence intervals of swab-positivity by age for rounds 5, 6, 6a and 6b.




Figure 10. Estimated odds ratios and 95% confidence intervals for jointly adjusted logistic regression model of swab-positivity for rounds 5, 6, 6a (16 to 25 October 2020), and 6b (26 October to 2 November 2020). Models were jointly adjusted for gender, age group, region, key worker status, ethnicity, household size, and deprivation index. The deprivation index is based on the Index of Multiple Deprivation (2019) at lower super output area. Here we group scores into quintiles, where 1 = most deprived and 5 = least deprived. HCW/CHW = health care or care home workers; Not FT, PT, SE = Not full-time, part-time, or self-employed.
*Yorkshire and The Humber