

1 **Reactive responses of zebras to lion encounters shape their predator-prey space game at**
2 **large scale**

3

4 Nicolas Courbin^{1*}, Andrew J. Loveridge², David W. Macdonald², Hervé Fritz³, Marion
5 Valeix³, Edwin T Makuwe⁴, Simon Chamaille-Jammes¹

6

7 ¹ Centre d'Ecologie Fonctionnelle et Evolutive, UMR 5175, Centre National de la Recherche
8 Scientifique (CNRS), Université de Montpellier, Université Paul Valéry Montpellier, Ecole
9 Pratique des Hautes Etudes (EPHE), 1919 Route de Mende, 34293 Montpellier Cedex 5,
10 France

11 ² Wildlife Conservation Research Unit, Department of Zoology, University of Oxford,
12 Recanati-Kaplan Centre, Tubney House, Abingdon Road, Oxford OX13 5QL, United
13 Kingdom

14 ³ Laboratoire de Biométrie et Biologie Evolutive, UMR 5558, Centre National de la
15 Recherche Scientifique (CNRS), Université Claude Bernard Lyon 1, Bât Gregor Mendel, 43
16 Boulevard du 11 Novembre 1918, 69622 Villeurbanne Cedex, France

17 ⁴ Zimbabwe Parks and Wildlife Management Authority, P.O. Box CY140, Causeway, Harare,
18 Zimbabwe

19

20 *Corresponding author, e-mail: ncourbin@gmail.com

1 Abstract

2 The predator-prey space game and the costs associated with risk effects are affected by prey
3 (i) proactive adjustments (when prey modify their behaviour in response to an a priori
4 assessment of the risk level) and (ii) reactive adjustments (when prey have detected an
5 immediate threat). Proactive adjustments are generally well-studied, whereas the frequency,
6 strength and duration of reactive adjustments remain largely unknown. We studied the space
7 use and habitat selection of GPS-collared zebras (*Equus quagga*) from 2 to 48h after an
8 encounter with lions (*Panthera leo*). Lion-zebra encounters generally occurred close to
9 artificial waterholes (<1km). Two-hours after an encounter, zebras were more likely to have
10 fled than stay when the encounter occurred in more risky bushy areas. During their flight,
11 zebras selected grasslands more than usual, getting great visibility. Regardless of their initial
12 response, zebras finally fled at the end of the night and reached areas located far from
13 waterholes where encounters with lions are less frequent. The large-scale flights (~4-5km) of
14 zebras led to a local zebra depression for lions. Zebras that had fled immediately after the
15 encounter resumed their behaviour of coming close to waterholes on the following day.
16 However, zebras that had initially stayed remained far from waterholes for an extra 24h,
17 remaining an elusive prey for longer. The delay in the flight decision had different short-term
18 consequences on the lion-zebra game. We reveal that the spatial context of the encounter
19 shapes the immediate response of prey, and that encountering predators induces strong
20 behavioural responses: prey flee towards distant, safer, areas and have a constrained use of
21 key resource areas which are at the heart of the predator-prey game at larger spatio-temporal
22 scales. Nighttime encounters were infrequent (once every 35 days on average), zebra
23 responses were short-lived (<36h) but occurred over a large spatial scale (several km).

24

25

1 Introduction

2 Prey have evolved numerous anti-predator behaviours to reduce predation risk (Lima and Dill
3 1990, Caro 2005). These behaviours are costly, both in terms of energy and time, and can
4 ultimately influence prey fitness and population dynamics (Preisser et al. 2005, Creel and
5 Christianson 2008, Zanette et al. 2011). The effects of predation risk on prey (i.e., non-
6 consumptive effects) had been largely overlooked in predator-prey studies until recently,
7 although they may sometimes be stronger than lethal (i.e., consumptive) effects (Preisser et al.
8 2005, Creel and Christianson 2008). A common prey response at the heart of these risk effects
9 is the change in space or habitat use in response to increasing risk thereby shaping the
10 predator-prey space game (Lima 2002, Sih 2005) and preventing prey from consistently using
11 the richest resource patches (Laundré 2010).

12 Prey behavioural adjustments to predation risk can be either proactive or reactive
13 (*sensu* Creel et al. 2014). Proactive adjustments occur when prey modify their behaviour in
14 response to an *a priori* assessment of the level of risk, in which the awareness of prey animals
15 to spatial and temporal distribution of predators plays a key-role. This has been well studied
16 for many taxa (invertebrates: Wojdak 2009; fish: Dupuch et al. 2009; birds: Heithaus et al.
17 2009; large mammals: Fortin et al. 2005, Latombe et al. 2014, Padié et al. in press), and it has
18 been shown that prey generally respond to an increase in perceived risk by selecting safer
19 habitats. Reactive adjustments occur when prey have detected an immediate threat, i.e.,
20 encountered either the predator or cues of its presence (e.g., Creel et al. 2005, Wirsing et al.
21 2010). In particular, prey have to choose to either flee or stay within the area where they have
22 detected a predator. It is accepted that the choice between these two immediate responses is
23 based on a decision-making process involving a cost-benefit analysis weighting the estimation
24 of the risk of being attacked and killed against the estimation of the future foraging gains
25 (Stankowich and Blumstein 2005, Cooper and Frederick 2007). The assessment of immediate

1 predation risk integrates intrinsic characteristics of predator and prey biology (e.g., hunting
2 and escape modes), environmental factors affecting prey vulnerability (e.g., presence of
3 predator-concealing cover), and contextual information (e.g., distance to and behaviour of the
4 predator, prey body condition) (Wirsing et al. 2010). The estimation of the foraging
5 opportunities integrates the prey's expectation about food quality and quantity where foraging
6 would take place, as well as its vigilance level which could ultimately reduce its foraging
7 efficiency if increased while staying in the area with the predator (Fortin et al. 2004, Creel et
8 al. 2014). Thus, a prey at a profitable patch with a high probability of escaping a predator if
9 attacked might flee immediately to avoid bearing the costs of monitoring an approaching
10 predator ("flush early and avoid the rush hypothesis", Blumstein 2010). Ultimately, the
11 dynamics of the predator-prey space game is determined by the relative frequency at which
12 the prey flee and by the spatial and temporal extents of their reactive responses to encounters.

13 In the short-run, spatially reactive adjustments are likely to be stronger than proactive
14 ones, as prey must avoid being killed. The total cost of a spatially reactive response will not
15 depend only on its intensity, however, but also on the time it takes for prey space use to return
16 to its basal level (i.e., proactive behaviour). Yet, studies generally assess spatially reactive
17 responses over very short time spans (seconds to minutes) and the full spatial and temporal
18 dynamics of the prey responses remains poorly known. Studies from a wide range of taxa (but
19 not large vertebrates) have focused on the time spent hiding in refuges after an attack
20 (reviewed in Cooper 2009). Several studies of ungulates have investigated how the distance
21 between the prey and the predator affect prey behaviour (e.g., vigilance: Liley and Creel
22 2008, Middleton et al. 2013, Creel et al. 2014), but very few have focused on its spatial
23 behaviour and/or the temporal persistence of the predator-induced behavioural changes. A
24 notable exception is Middleton et al. (2013), which revealed that elk (*Cervus elaphus*) net
25 daily displacement was slightly elevated (by 450m on average) after an encounter with wolves

1 (*Canis lupus*), and that elk movement rates returned to basal levels 24h after the encounter.
2 Therefore, further studies are required to understand the scale of the spatial response of prey
3 to encounters with predators, and its temporal persistence. This is also a pre-requisite to assess
4 to what extent the responses contribute to shaping the predator-prey space game and trophic
5 cascades (Fortin et al. 2005).

6 In Hwange National Park (Zimbabwe), we studied the spatially reactive response of
7 zebras (*Equus quagga*) from 2 to 48 hours after encounters with lions (*Panthera leo*), their
8 main predator. This zebra population seems to be under top-down control by lions and
9 currently declining due to a high predation pressure (Grange et al. 2015). Previous studies
10 (using survey data rather than individual based data) in this system have revealed that during
11 daytime zebras select areas located near artificial waterholes, which provide profitable open
12 short-grass habitats with high visibility (Valeix et al. 2009) and lower mortality risk
13 (Davidson et al. 2012). Lions select for these areas to benefit from high prey densities, but at
14 night, they hunt more successfully in the dense vegetation at the edge of these areas
15 (Davidson et al. 2012). Lions rest in their vicinity during the day too (Davidson et al. 2013).
16 However, it is not known how zebras respond to the presence of hunting lions at night, and
17 specifically whether the immediate response of zebras is to flee to distant areas or is a more
18 small-scale avoidance of lions. Equally, it is not known for how long zebras adjust their
19 habitat and space use following an encounter with lions. Using simultaneously collected GPS
20 locations for lions and zebras, we investigated the spatially reactive responses of a large
21 herbivore in highly risky situations (i.e., <500m from lions) to understand how these
22 responses can shape the predator-prey spatial game.

23 Accordingly, we addressed four questions:

24 Question 1: Do zebras stay or flee immediately after encountering lions?

1 Question 2: Is zebra response influenced by the resource/risk characteristics of the encounter
2 area?

3 Question 3: Do zebras that flee move away from the risky encounter site towards safer areas
4 of the landscape (i.e., away from waterholes), and for how long do space use adjustments
5 persist?

6 Question 4: While they are staying or moving away from lions, do zebras select grasslands
7 (where they are less vulnerable because of greater visibility) more strongly than at other
8 times, and for how long does increased selection for grasslands persist?

9

10 Methods

11 *Study site*

12 The study was conducted in the north-eastern region of Hwange National Park, Zimbabwe.

13 The vegetation is typical of dystrophic semi-arid savannahs, dominated by bushlands (>60%
14 of the landscape; mostly *Combretum* spp., *Acacia* spp. and *Terminalia sericea*) but

15 interspersed with small patches of grasslands and larger patches woodlands (mostly *Baikiaea*
16 *plurijuga* woodlands) (Rogers 1993, Chamaillé-Jammes et al. 2006). Average annual rainfall

17 is c. 600 mm, with most of the rains falling between November and April (wet season,

18 Chamaillé-Jammes et al. 2006). There are no perennial rivers in the study area and natural

19 pans progressively dry up. Water is only available during the late dry season (August to

20 October) in waterholes artificially supplied with pumped groundwater. Artificial waterholes

21 are largely situated in open areas characterized by greater availability of grasslands and open

22 bushlands (bushland-1 in supplementary material Appendix 1 Fig. A1; see also Chamaillé-

23 Jammes et al. 2009a), which are habitats favoured by zebras (Fischhoff et al. 2007, Valeix et

24 al. 2009, see Results also). Artificial waterholes also drive the space use of lions all year

25 round (Valeix et al. 2010, Davidson et al. 2012). In preliminary analyses, we found that the

1 proportions of immediate responses – staying vs fleeing – were similar between wet and dry
2 seasons, and we did not find qualitative changes in the space use of zebras after an encounter
3 with lions between seasons. We therefore favoured greater statistical power and conducted all
4 analyses without distinguishing between seasons.

5 Zebra density in the study area was estimated from road-transects driven at least twice
6 a year between 1999 and 2010 and was c. 1 individual/km² (Chamaillé-Jammes et al. 2009b,
7 Grange et al. 2015). Lion density in the study area was estimated from an individual-based
8 database where almost all lions were known and identified by whisker patterns and other
9 natural physical characteristics, and was c. 3.5 lions/100 km² over the study period (Loveridge
10 et al. unpublished).

11

12 *Environmental covariates*

13 We used Landsat-7 ETM+ satellite images (30-m resolution) taken in August 2002,
14 November 2002 and April 2003 to conduct an unsupervised classification of habitat types. We
15 first performed a k-means analysis on these images. Then, for each cluster, we identified large
16 homogeneous patches and within these patches conducted field vegetation surveys: during
17 each survey, every 2m along two 60-m transects, we recorded the presence and species
18 identity of any woody species within the <1m, 1 to 3m, 3 to 8m and >8m height-class. We
19 also recorded the presence or absence of grass. Overall, 168 transects were conducted. We
20 used these field data to classify k-means clusters in six habitat types of increasing woody
21 cover: grassland, 3 types of bushlands (bushland-1, bushland-2, bushland-3) representing a
22 decreasing gradient of openness, wooded bushland, woodland. Wooded bushland was the
23 baseline habitat type in the subsequent analyses. We found no strong correlations ($-0.30 < r <$
24 0.19) between all habitat types and distance to artificial waterholes in our study area.

25

1 *GPS data and identification of lion-zebra encounters*

2 GPS collars were fitted to zebras by qualified personnel using standard protocols for the
3 species. Collars were deployed for an average of 357 ± 43 (mean \pm SE) days per zebra and in
4 analyses we used location data collected hourly between August 2009 and November 2013
5 from 22 female zebras belonging to different herds. Zebras move more during the day than at
6 night (382 ± 2 m/h vs 224 ± 2 m/h respectively). Over the same period, 21 lions (15 males and
7 6 females) equipped with GPS collars (fitted by qualified personnel using standard protocols
8 for the species, Fahlman et al. 2005) also used the area exploited by the monitored zebras. Not
9 all lions in the study area were collared (this would be logistically impossible and ethically
10 questionable). As all females of a pride are sighted together c. 90% of the time (revealing very
11 little fusion/fission dynamics, Valeix et al. 2009), efforts were made to have at least one
12 female with a collar in each pride, and the intensive field monitoring suggests that this was
13 virtually achieved in the study area. Males are only infrequently seen with females (c. 7% of
14 the sightings, Davidson et al. 2013), suggesting that they do hunt themselves and explaining
15 why monitoring this sex required more collars to track territorial males, coalitions and
16 dispersers. Overall, only 1% of zebra locations fell outside of the GPS-tracked lion home
17 ranges. Lion GPS collars acquired locations at 1h or 2h intervals during the night and every
18 2h or 2 or 3 times during the day. Lions were monitored for an average of 630 ± 62 days per
19 individual. Lions move 596 ± 3 m/h at night and much less during the day (181 ± 5 m/h).

20 We assumed that a zebra encountered a lion when their simultaneous locations were
21 less than 500m apart at night. Although it is likely that zebras can detect and respond to lion
22 presence when these are further away (e.g., Valeix et al. 2009), this short distance threshold
23 was chosen to ensure that we studied zebra response in a highly risky situation, during which
24 it was unlikely that risk was not perceived. We focused on nighttime locations only as lions
25 mainly hunt during this period and we therefore anticipated that their presence would induce

1 spatially reactive responses from their prey. For a given zebra, if several encounters occurred
2 during the same night, we considered that the encounter had taken place when the distance
3 between the zebra and a lion was smallest. We kept 192 encounters, being 8.2 ± 1.3 (mean \pm
4 SE) encounters per zebra, interspaced by at least 48h for a given zebra, allowing assessing the
5 dynamics of space use and habitat selection of zebras during the next 48h after an encounter.
6 To evaluate whether the spatially reactive responses of zebras after encountering lions
7 departed from zebra behaviours in the absence of lion (i.e., proactive response), we used a set
8 of 'no-encounter' locations: for each encounter we randomly selected 10 (when possible) 'no-
9 encounter' locations among all locations (for the same zebra and the same hour of the night)
10 when no lions were within 500m. As for the encounter case, the 'no encounter' locations
11 retained were spaced by at least 48h for a given zebra. Choosing a small distance threshold
12 ($<500\text{m}$) to define encounter cases is a conservative approach as possibly excluding encounter
13 events occurring over larger distance decreases our ability to detect differences between the
14 'encounter' and 'no encounter' cases.

15

16 *Habitat selection of lions and zebras and habitat characteristics of encounter areas*

17 We used resource selection functions (RSF, Manly et al. 2002) to estimate habitat selection of
18 lions and zebras within their home range during both day and night periods. For each species
19 and period, the RSF models compared the different habitat types and the distance to the
20 closest artificial waterhole (referred to as distance to waterhole thereafter) at the GPS
21 locations of a given individual with those at an equal number of locations drawn randomly
22 within its home range (Courbin et al. 2014). We estimated RSF coefficients using a
23 generalized linear mixed model (GLMM) with a logit link and a binomial distribution for
24 errors. We added a random intercept to account for the unbalanced sampling design among
25 individuals of a given species (Gillies et al. 2006). To account for non-independence among

1 observations of different lions, we kept only one location (randomly chosen) at each time step
 2 when distance between individuals was <300m (as lions travelling together are most often
 3 located less than 300m apart, see supplementary material Appendix 1 Fig. A2). Following
 4 recommendations of Koper and Manseau (2009), we used GLMM with robust empirical
 5 standard errors that are robust to both among- and within-individual correlations and that
 6 provide robust estimates of significance. RSFs were fitted with the *GLIMMIX* procedure of
 7 SAS 9.2 software (SAS Institute 2008) and took the following form:

$$8 \quad w(\mathbf{x}_{ij}) = \exp(b_0 + b_h \text{habitat}_{hij} + b_{\text{waterhole}} \text{waterhole}_{ij} + g_{0j}) \quad \text{eqn 1}$$

9 where $w(\mathbf{x}_{ij})$ is the relative probability of selection for lion or zebra during the day or the
 10 night, \mathbf{x} is a vector, b_0 is the mean intercept, b_h is the estimated fixed regression coefficient
 11 for habitat type h and $b_{\text{waterhole}}$ is the coefficient for distance to waterhole, i represents the i^{th}
 12 observation, j represents the j^{th} individual, and g_{0j} is the random effect on the intercept b_0 for
 13 animal j . We used a cube-root transformed variable for distance to waterhole as 1)
 14 untransformed variable was highly right-skewed distributed implying high leverage effects,
 15 and 2) model fit was largely improved based on the Akaike information criterion ($\Delta\text{AIC} >$
 16 592 for day and night periods). No multicollinearity issue was detected (variance inflation
 17 factors [VIFs] < 1.3) in all models (Dormann et al. 2013). We evaluated model robustness
 18 using k -fold cross-validation as suggested by Boyce et al. (2002).

19 We then tested if encounters between zebras and lions occurred randomly along the
 20 zebra movement paths or if they occurred within areas with specific habitat composition and
 21 at a specific distance to water. Because GPS-sampling of zebra and lion locations was
 22 discrete, we had only an approximate idea of where the encounter occurred. We therefore
 23 calculated the proportion of each habitat type within a 500-m radius buffer of the location
 24 defined as encounter location as defined above (see details in supplementary material

1 Appendix 2 Fig. A1 for explanations about the choice of this radius). We compared the
2 characteristics of areas occupied by zebra when an encounter occurred with those occupied at
3 times when lions were absent (i.e., ‘no-encounter’ set) using a GLMM with a logit link and a
4 binomial distribution for errors and a random intercept for each zebra. We assessed the
5 relative probability that an encounter occurred along the movement paths of zebras,
6 $w(\mathbf{x}_{ij})_{\text{encounter}}$, using equation 1, where b_h is the estimated fixed regression coefficient for
7 proportion of habitat type h and $b_{\text{waterhole}}$ is the coefficient for distance to waterhole. No
8 multicollinearity issue was detected (VIFs < 1.6). The models were fitted using the R software
9 (R Core Team 2013).

10

11 *Typology of immediate spatial responses of zebras after an encounter with lions (Question 1)*

12 We ascertained that zebras could display two contrasted responses (stay or flight)
13 immediately (i.e., within 2h) after encountering lions. We hypothesized that the actual
14 proximity of lions during encounters could influence zebra decision on whether to stay or flee.
15 We therefore used an unsupervised model-based clustering analysis with two axes: distance
16 between zebra and lion at the time of the encounter, and net displacement of zebra 2h after the
17 encounter. We evaluated whether the data were best described by one or two clusters, and if
18 the best two-cluster model separated the data along the axis of net displacement after the
19 encounter, thereby indicating the existence of two contrasted spatial responses (see details in
20 supplementary material Appendix 3 Fig. A1).

21

22 *Habitat determinants of immediate response choice (Question 2)*

23 We tested whether the likelihood that zebras displayed a given immediate response (stay or
24 flight) depended on the habitat composition of the encounter area and its proximity to a
25 waterhole. We compared characteristics of encounter areas where zebras had fled with those of

1 encounter areas where zebras had stayed, using a GLMM with a logit link and a binomial
2 distribution for errors, while accounting for the unbalanced number of observations among
3 individuals with a random intercept. The model took the form:

$$4 \quad P(\mathbf{x}_{ij})_{\text{response choice}} = \frac{\exp(b_0 + b_h \text{habitat}_{hij} + b_{\text{waterhole}} \text{waterhole}_{ij} + g_{0j})}{1 + \exp(b_0 + b_h \text{habitat}_{hij} + b_{\text{waterhole}} \text{waterhole}_{ij} + g_{0j})} \quad \text{eqn 2}$$

5 where $P(\mathbf{x}_{ij})_{\text{response choice}}$ is the probability to flee immediately rather than stay, \mathbf{x} is a vector, b_h
6 is the estimated fixed regression coefficient for proportion of habitat type h and $b_{\text{waterhole}}$ is the
7 coefficient for distance to waterhole, i represents the i^{th} observation, j represents the j^{th}
8 individual, and g_{0j} is the random effect on the intercept b_0 for animal j . No multicollinearity
9 issue was detected (VIFs < 1.6). The models were fitted using the R software (R Core Team
10 2013).

11

12 *Temporal dynamics of zebra space use after an encounter (Question 3)*

13 We studied the dynamics of zebra space use after encounters for each type of response (stay
14 and flight). For each encounter we computed the zebra's net displacement from the encounter
15 site and the distance to waterhole on several occasions (encounter, 2h, ..., 3rd sunrise after the
16 encounter; see Fig. 1). We also calculated the differences in the net displacement from the
17 encounter site between these consecutive occasions. For each interval between occasions we
18 assessed whether changes in the net displacement were different from those expected if zebras
19 had not encountered a lion, and if they differed between the different immediate responses.
20 We did so by fitting linear mixed-effects models (LMM) with random intercepts for each
21 zebra and a dummy binary variable as predictor, allowing to compare stay vs. proactive, flight
22 vs. proactive or flight vs. stay. We used a similar approach to assess if the dynamics of the
23 distance to waterhole between occasions varied between proactive behaviour and the two

1 types immediate response, and between these responses. Distance to waterhole was cube root
 2 transformed. LMMs were fitted using the R software (R Core Team 2013).

3

4 *Temporal dynamics of fine-scale habitat selection of zebras after an encounter (Question 4)*

5 We used step selection functions (SSF, Fortin et al. 2005) to study the habitat selection of
 6 zebras after encounters with lions, at three consecutive occasions: immediately (2h-later), the

7 day after the encounter and the night after the encounter. At each occasion, we compared
 8 habitat selection between proactive behaviour and each type of immediate response by fitting

9 one SSF per response. For each SSF, we extracted the movement path collected during the
 10 occasion of interest, after encounters and in the 'no-encounter' data set. Movement paths were

11 decomposed into a series of steps (i.e., straight-line segments linking successive 1h locations),
 12 and each step was paired with 10 random steps to create a stratum. Random steps had the

13 same starting location as observed steps, but differed in that length and turning angle were
 14 randomly drawn from the empirical distribution of step lengths and turning angles obtained by

15 pooling steps data from all other individuals, as recommended by Fortin et al. (2005) and
 16 Forester et al. (2009). We estimated SSF parameters using conditional logistic regression

17 within a generalized estimating equation (GEE) framework. Temporal autocorrelation
 18 between the steps occurring the same day or night can bias the standard errors of parameter

19 estimates, and we therefore calculated robust standard errors after having grouped all steps
 20 occurring the same day or the same night in independent clusters (Fortin et al. 2005, Craiu et

21 al. 2008). The SSF models took the form:

22
$$w(\mathbf{x}_{ij})_{zebra,step} = \frac{\exp(b_h \text{habitat}_{hij} + b_{steplength} \text{steplength}_{ij} + b_{h,encounter} \text{habitat}_{hij} \cdot \text{encounter}_{ij})}{\sum_{h'} \exp(b_{h'} \text{habitat}_{h'ij} + b_{steplength,encounter} \text{steplength}_{ij} \cdot \text{encounter}_{ij})} \quad \text{eqn 3}$$

23 where $w(\mathbf{x}_{ij})_{zebra,step}$ is the relative probability of selecting a step for zebra, \mathbf{x} is a vector, b_h is

24 the selection coefficient for the habitat type h at the step's ending point, $b_{h,encounter}$ is the

1 selection coefficient for the interaction between habitat type h and a dummy variable allowing
2 to distinguish if the steps occurred after an encounter or a no-encounter location, i represents
3 the i^{th} observation and j represents the j^{th} individual. The interactions allowed us to test
4 whether habitat selection after an encounter differed from the one observed during proactive
5 behaviour. We added a spline function of step length, with knots at the first, second, and third
6 quartiles of observed step lengths, which have been shown to remove bias in other model
7 parameters (Forester et al. 2009). No multicollinearity issue was detected (VIFs < 2.6) in all
8 these SSFs. We tested the robustness of the models using k -fold cross-validation adapted for
9 paired designs (Fortin et al. 2009, Courbin et al. 2014).

10

11 Results

12 *Habitat selection of lions and zebras and habitat characteristics of encounter areas*

13 RSFs models had high to very high predictive power ($\bar{r}_s^2 > 0.55$; see supplementary material
14 Appendix 1 Table A1). As expected, zebras strongly selected the most open habitats
15 (grasslands, and to a lesser extent, the two most open bushlands; supplementary material
16 Appendix 1 Table A1) during both day and night within their home range. Zebras selected
17 woodlands during the day. Lions selected grasslands, and the most open bushlands, during
18 day and night, and they tended to avoid woodlands at night (Supplementary material
19 Appendix 1 Table A1). Both species strongly selected areas located near waterholes within
20 their home range (Supplementary material Appendix 1 Table A1), but zebras stayed further
21 away from waterholes at night (Fig. 1B). Relatively to areas visited along their paths, at night
22 when lions hunt, zebras were more likely to encounter lions in areas with a higher proportion
23 of grasslands and a lower proportion of bushlands (bushland-3; Table 1; mean proportion of
24 grasslands around locations was 0.18 (95% CI: 0.16-0.20) and 0.09 (95% CI: 0.08-0.10)
25 during encounter and proactive behaviour, respectively; mean proportion of bushland-3

1 around locations was 0.045 (95% CI: 0.04-0.05) and 0.085 (95% CI: 0.08-0.09) during
2 encounter and proactive behaviour, respectively).

3

4 *Typology of immediate spatial responses of zebras after an encounter with lions (Question 1)*

5 The best two-cluster model classifying zebra immediate responses after encounters was
6 superior to the model assuming only one cluster ($\Delta\text{ICL} = 78$), suggesting that two immediate
7 responses were indeed present. One two-cluster model was much better than other two-cluster
8 models ($\Delta\text{ICL} > 3.9$; see supplementary material Appendix 3 Table A1), and it discriminated
9 encounters based on the zebra net displacement only: one cluster identified a 'stay' immediate
10 response (net displacement 2h after the encounter $<900\text{m}$, $n=108$), and the other cluster
11 identified a 'flight' immediate response (net displacement 2h after the encounter $>900\text{m}$, 2567
12 ± 165 [mean \pm SE], ranging 910 to 6540m , $n=84$). The mean proportion of stay responses per
13 zebra weighted for the number of observations for each zebra had a very low standard error
14 (0.56 ± 0.05 , mean \pm SE), revealing that responses were not individual-specific.

15

16 *Habitat determinants of immediate response choice (Question 2)*

17 We found that habitat characteristics influenced the choice of immediate response of zebras.
18 Zebras were more likely to flee than stay when the encounter occurred in an area where
19 bushlands (bushland-3) were more common (Table 1; mean proportion of bushland-3 was
20 0.07 and 0.03 for flight and stay, respectively).

21

22 *Temporal dynamics of zebra space use after an encounter (Question 3)*

23 At longer time scales, all zebras left the area where the encounter with lions occurred and
24 reached areas 4km away in less than 24h , while they usually moved away less than 1.5 km
25 from where they were when lions were absent (Fig. 1A). However, the two contrasting

1 immediate responses of zebras, flight or stay, were associated with widely different space use
2 that we termed immediate-flight and delayed-flight tactics, respectively. In the immediate-
3 flight tactic, zebras moved very rapidly away from both the encounter site and waterholes
4 before the first sunrise after the encounter occurred (Fig. 1A-B; supplementary material
5 Appendix 1 Table A2). By the day following an encounter, zebras resumed their daily
6 patterns of visits to areas close to water during the day and moving away at night, as done
7 during their proactive behaviour (Fig. 1B). Zebras did not come back to where the encounter
8 occurred however (compare net displacement and distance to water in Fig. 1A-B). In the
9 delayed-flight tactic, the movement of zebras towards safer areas (i.e., away from waterholes)
10 that started later during the night went on during the day following the encounter (Fig. 1A-B;
11 supplementary material Appendix 1 Table A2). Zebras resumed visiting areas near waterholes
12 only the second day after the encounter.

13

14 *Temporal dynamics of fine-scale habitat selection of zebras after an encounter (Question 4)*

15 All SSFs had very high predictive power ($\bar{r}_s > 0.79$; see supplementary material Appendix 1
16 Table A3, A4). The habitat selection of zebras immediately after an encounter depended on
17 whether they stayed or fled immediately. The next 2 hours after the encounter zebras using a
18 delayed-flight tactic had similar selection coefficients than during proactive behaviour,
19 selecting the most open habitat types (Fig. 2A; supplementary material Appendix 1 Table
20 A3). However, because grassland availability increased in encounter areas (see section
21 'Habitat selection...' above and Fig. 2A), zebras used grasslands more. Zebras using the
22 immediate-flight tactic altered their habitat selection after an encounter. They selected
23 grasslands (and possibly woodlands) even more strongly during their flight than during
24 proactive behaviour (Fig. 2D; supplementary material Appendix 1 Table A4). The day and the
25 night following an encounter, habitat selection of zebras did not differ from the one observed

1 during proactive behaviour (Fig. 2B, C, E, F; supplementary material Appendix 1 Table A3,
2 A4).

3

4 Discussion

5 Our study is one of the first to provide a comprehensive analysis of the spatially reactive
6 response of a prey species to encounters with its primary predator over large spatial and
7 temporal scales. In particular, we reveal how short-term reactive decisions (immediate or
8 delayed-flight) were associated with different space use patterns during subsequent days that
9 likely affected the predator-prey space game at the landscape scale.

10

11 *Zebras encounter lions within waterhole areas*

12 In our system, the predator-prey space race revolves around artificial waterholes that are at the
13 heart of a spatial trade-off between food and safety. Grasslands surrounding waterholes
14 provide suitable foraging habitats with good visibility and were particularly selected by zebras
15 during the day (this study, Valeix et al. 2009), but also sometimes at night (this study,
16 Fischhoff et al. 2007), although we found that zebras were commonly away from waterholes
17 at night. Lions always selected these areas (this study, Valeix et al. 2010, Davidson et al.
18 2012, Davidson et al. 2013). Zebras were thus more likely to encounter lions at night whether
19 they reached the most open areas close to waterhole (<1km, Fig. 2B). Our study reveals that
20 lions may favour areas of the landscape where encounter rate is the greatest: areas close to
21 artificial waterholes attract many species and open habitats provide high detectability of the
22 prey.

23

24 *Zebras flee after encountering lions*

1 Lions are ambush predators that use cover to approach prey and hunt successfully (Hopcraft
2 et al. 2005, Davidson et al. 2012). This likely explains why zebras were far more likely to flee
3 immediately than stay once they had encountered lions in more bushy areas. When zebras
4 decided to leave the encounter area immediately, they did so by shifting their selection
5 towards habitat types with high visibility (mostly open areas and high canopy woodlands with
6 little understory) and habitat types where the likelihood of meeting lions was reduced
7 (woodlands), probably to maximize their safety. Zebras staying in the vicinity of lions did not
8 alter their habitat selection pattern. Note that individual zebras used a mix of both types of
9 immediate response, pointing out that the response choice seemed to rely more on the context
10 of the encounter (habitat type, but also possibly lion pride size, which we did not have
11 sufficient data to study) than on fixed innate (personality-driven) or acquired (previous
12 experience) response by individuals. The potential costs associated with the close presence of
13 lions (e.g., potential costs of monitoring, risk of injury or death) might possibly explain why,
14 irrespectively of their immediate response, zebras fled (with a delay or not) and they generally
15 left the encountered area by the end of the night towards safer areas located far from
16 waterholes. Our results support previous studies suggesting that ambush predators induce
17 strong anti-predator behavioural responses because risk is spatially predictable, particularly
18 once the presence of the predator is actually established (Preisser et al. 2007, Thaker et al.
19 2011, but see discussion in Creel et al. 2014).

20

21 *Consequences of the flight decision on the lion-zebra space game*

22 Zebras rapidly moved ca. 4-5km away from lions after an encounter, and thus became
23 unavailable prey for these lions during the subsequent days. Our results thus support Valeix et
24 al's. (2011) hypothesis that changes in prey behaviour, possibly leading to changes in prey
25 abundance (if most of individuals flee, if this is true for most prey and if lions presence

1 elsewhere does not increase the number of incoming individuals), could explain why in
2 Hwange NP lions move to another waterhole after having made a kill. This may be a general
3 phenomenon underlying the lion-zebra space game: a reduction of local zebra abundance
4 when lions are present has also been observed in the Laikipia ecosystem (Fischhoff et al.
5 2007). Fortin et al. (2013) have demonstrated theoretically and empirically for caribou
6 (*Rangifer tarandus*) that avoidance behaviour might indeed lead to prey aggregation at
7 intermediate distance from their disturbance or threat source, here logging areas. A depression
8 in the local abundance of zebra at disturbed waterholes is likely to induce a temporary peak of
9 zebra abundance at the surrounding waterholes (i.e., ~4-5km away from the disturbed
10 waterhole), which could affect future prey search patterns of lions involved in the encounter,
11 but could also benefit other lions and other carnivores occupying neighbouring waterholes.
12 Our results emphasize that prey avoidance behaviour is critical to the understanding of the
13 predator-prey space game. They further highlight how it likely also creates indirect
14 relationships between predators that use different areas of the landscape. In Hwange NP,
15 zebras make a significant contribution to the lion diet (~10%) but are not, however, a species
16 highly selected by lions (Davidson et al. 2013). It is therefore unclear how changes in zebra
17 distribution within the landscape might actually affect lion hunting strategies and their shifts
18 towards other prey species. Further studies should also investigate whether the large-scale
19 movements after an encounter with lions is a common response of prey, and more generally
20 whether local prey depression may drive the predator-prey space game in landscapes with
21 strong spatial anchors of the prey (Sih 2005). Ultimately, what needs to be clarified is under
22 what conditions will prey display large-scale reactive avoidance of predators to complement,
23 or replace, other anti-predator strategies (e.g., increasing vigilance, changing grouping
24 patterns), and how the costs of staying may vary with the environmental context or prey and
25 predator ecology (Wirsing et al. 2010).

1 The fact that zebras either immediately fled or delayed their decision to flee by a few
2 hours had different consequences for the predator-prey space game. The day following the
3 night of the encounter, zebras using immediate-flight tactic resumed the proactive behaviour
4 of coming closer to waterholes during the day and moving away at night. Their behaviour
5 thus caused an immediate movement of prey towards areas of the landscape where other lions
6 could be present. Conversely, zebras using delayed-flight tactic mainly moved across the
7 landscape, away from waterholes, the day after the encounter. They were also further away
8 from water than usual on the two consecutive nights. This behaviour therefore led to a wider –
9 but still short – temporal window during which these zebras were unavailable to lions.

10

11 *On the importance of the spatially reactive responses of zebras*

12 For zebras, the overall fitness consequences of their reactive responses depend on the
13 frequency of the encounters, on the duration of the response and on the strength of the
14 response (i.e., on cost of a time unit spent in reactive response). Very few studies have
15 estimated these three properties of reactive responses *in natura*. In a rare example, Middleton
16 et al. (2013) showed that in the Great Yellowstone Ecosystem, elk have infrequent (every 9
17 and 50 days on average for migratory and resident elk, respectively), short-lived (<24h) and
18 probably weakly costly reactive responses (no reduction in feeding rate and no shift in habitat
19 use) after encounters with wolves. Ultimately, the frequency of encounters with wolves was
20 not associated with reduced body fat or pregnancy. In Hwange NP, nighttime encounters
21 between zebras and lions are probably also infrequent: zebras encountered lions at night every
22 35 days on average in the dataset used here. The actual encounter rate (that cannot be
23 calculated using the available data) is probably higher. First, zebras are more likely to
24 encounter lions during the day as they use waterhole areas more often. Daytime encounters
25 are likely to be far less risky and may not lead to a large-scale spatial response however.

1 Second, the GPS-based measures of encounter rates could be underestimated (e.g., lions were
2 not all monitored, an encounter could occur between GPS fixes, see details in Creel et al.
3 2013), although these effects are likely to be low here given the coverage and frequency of the
4 lion tracking. The frequency of risky encounter is therefore low, and we have also showed
5 that the reactive response of zebras to these encounters is short-lived (<36h). Thus, each
6 response would have to be particularly costly for these responses to significantly reduce zebra
7 fitness in this system. The actual cost of a single response is complex to estimate. The large-
8 scale displacement induced by lions probably bears some energetic costs, as zebras travelled
9 three times more kilometers than usual in a few hours to leave the encounter area. Walking is,
10 however, not so energetically costly and the significance of this increase should be low. A few
11 recent studies points towards foraging costs for zebras having been in the vicinity of lions
12 (Barnier et al. 2014, Creel et al. 2014), but again, the significance of a small reduction in
13 intake (Creel. et al. 2014) or forage quality (Barnier et al. 2014) is unknown. Altogether, this
14 suggests that currently no obvious link could be made between the strong large-scale
15 behavioural response observed here and zebra fitness in this population. This would only be
16 formally demonstrated by monitoring the lion-encounter history and demography of zebra
17 herds over time. This will be logistically challenging, but will be required to deepen our
18 understanding of risk effects, disentangling the relative contribution of proactive and reactive
19 responses on prey fitness.

20

21 Acknowledgements

22 This research was partly funded by the grants ANR-08-BLAN-0022, ANR-11-CEPS-003,
23 Darwin Initiative for Biodiversity Grants 162-09-015 and EIDPO002, Mitsubishi Fund for
24 Europe and Africa, R.G. Frankenberg Foundation, Boesak and Kruger Foundation, Rufford
25 Maurice Laing Foundation, SATIB Trust, Eppley Foundation, Riv and Joan Winant and

1 Recanati-Kaplan Foundation. M. Muzamba, J. Hunt, B. Stapelkamp, N. Elliot and H. Valls
2 Fox contributed to fieldwork. The late N. Ganzin obtained and did some initial work on the
3 Landsat images used for the vegetation map. We also thank Thierry Duchesne for his
4 statistical advices. This research was authorized by the Zimbabwe Parks and Wildlife
5 Management Authority under permits (permit numbers: REF:DM/Gen/(T) 23(1)(c)(ii):
6 03/2009, 01/2010, 25/2010, 05/2011, 06/2011, 12/2012, 15/2012, 08/2013). Comments from
7 two anonymous reviewers greatly improved the manuscript.

8

9 References

- 10 Barnier, F. et al. 2014. Diet quality in a wild grazer declines under the threat of an ambush
11 predator. – Proc. R. Soc. B 281: 20140446.
- 12 Blumstein, D. T. 2010. Flush early and avoid the rush: a general rule of antipredator
13 behavior? – Behav. Ecol. 21: 440-442.
- 14 Boyce, M. S. et al. 2002. Evaluating resource selection functions. – Ecol. Model. 157: 281-
15 300.
- 16 Caro, T. M. 2005. Antipredator defenses in birds and mammals. – University of Chicago
17 Press.
- 18 Chamaillé-Jammes, S. et al. 2006. Spatial patterns of the NDVI-rainfall relationship at the
19 seasonal and interannual time scales in an African savanna. – Int. J. Remote Sens. 27: 5185-
20 5200.
- 21 Chamaillé-Jammes, S. et al. 2009a. Piosphere contribution to landscape heterogeneity: a case
22 study of remote-sensed woody cover in a high elephant density landscape. – Ecography 32:
23 871-880.
- 24 Chamaillé-Jammes, S. et al. 2009b. Seasonal density estimates of common large herbivores in
25 Hwange National Park, Zimbabwe. – Afr. J. Ecol. 47: 804-808.

- 1 Cooper Jr, W. E. and Frederick, W. G. 2007. Optimal flight initiation distance. – J. Theor.
2 Biol. 244: 59-67.
- 3 Cooper Jr, W. E. 2009. Theory successfully predicts hiding time: new data for the lizard
4 *Sceloporus virgatus* and a review. – Behav. Ecol. 20: 585-592.
- 5 Courbin, N. et al. 2014. Logging-induced changes in habitat network connectivity shape
6 behavioral interactions in the wolf-caribou-moose system. – Ecol. Monogr. 84: 265-285.
- 7 Craiu, R. V. et al. 2008. Inference methods for the conditional logistic regression model with
8 longitudinal data. – Biometrical J. 50: 97-109.
- 9 Creel, S. et al. 2005. Elk alter habitat selection as an antipredator response to wolves. –
10 Ecology 86: 3387-3397.
- 11 Creel, S. and Christianson, D. 2008. Relationships between direct predation and risk effects. –
12 Trends Ecol. Evol. 24: 194-201.
- 13 Creel, S. et al. 2013. Underestimating the frequency, strength and cost of anti-predator
14 responses with data from GPS collars: an example with wolves and elk. – Ecol. Evol. 3: 5189-
15 5200.
- 16 Creel, S. et al. 2014. Effects of predation risk on group size, vigilance, and foraging behavior
17 in an African ungulate community. – Behav. Ecol. 25: 773-784.
- 18 Davidson, Z. et al. 2012. Environmental determinants of habitat and kill site selection in a
19 large carnivore: scale matters. – J. Mammal. 93: 677-685.
- 20 Davidson, Z. et al. 2013. Seasonal diet and prey preference of the African lion in a waterhole-
21 driven semi-arid savanna. – PLoS ONE 8:e55182.
- 22 Dormann., C. F. et al. 2013. Collinearity: a review of methods to deal with it and a simulation
23 study evaluating their performance. – Ecography 36: 27-46.

1 Dupuch, A. et al. 2009. Testing the effects of resource distribution and inherent habitat
2 riskiness on simultaneous habitat selection by predators and prey. – Anim. Behav. 78: 705-
3 713.

4 Fahlman, A. et al. 2005. Reversible anaesthesia of free-ranging lions (*Panthera leo*) in
5 Zimbabwe. – J. S. Afr. Vet. Assoc. 76: 187-192.

6 Fischhoff, I. R. et al. 2007. Habitat use and movements of plain zebra (*Equus burchelli*) in
7 response to predation danger from lions. – Behav. Ecol. 18: 725-729.

8 Forester, J. D. et al. 2009. Accounting for animal movement in estimation of resource
9 selection functions: sampling and data analysis. – Ecology 90: 3554-3565.

10 Fortin, D. et al. 2004. Foraging costs of vigilance in large mammalian herbivores. – Oikos
11 107: 172-180.

12 Fortin, D. et al. 2005. Wolves influence elk movements: behavior shapes a trophic cascade in
13 Yellowstone National Park. – Ecology 86: 1320-1330.

14 Fortin, D. et al. 2009. Group-size-mediated habitat selection and group fusion-fission
15 dynamics of bison under predation risk. – Ecology 90: 2480-2490.

16 Fortin, D. et al. 2013. Movement responses of caribou to human-induced habitat edges lead to
17 their aggregation near anthropogenic features. – Am. Nat. 181: 827-836.

18 Gillies, C. S. et al. 2006. Application of random effects to the study of resource selection by
19 animals. – J. Anim. Ecol. 75: 887-898.

20 Grange, S. et al. 2015. [Demography of plains zebras \(*Equus quagga*\) under heavy predation.](#) –
21 [Popul. Ecol. 57: 201-214.](#)

22 Heithaus, M. R. et al. 2009. Towards a predictive framework for predator risk effects: the
23 interaction of landscape features and prey escape tactics. – J. Anim. Ecol. 78: 556-562.

24 Hopcraft, G. J. C. et al. 2005. Planning for success: Serengeti lions seek prey accessibility
25 rather than abundance. – J. Anim. Ecol. 75: 559-566.

1 Koper, N. and Manseau, M. 2009. Generalized estimating equations and generalized linear
2 mixed-effects models for modelling resource selection. – *J. Appl. Ecol.* 46: 590-599.

3 Latombe, G. et al. 2014. Spatio-temporal dynamics in the response of woodland caribou and
4 moose to the passage of grey wolf. – *J. Anim. Ecol.* 83: 185-198.

5 Laundré, J. W. 2010. Behavioral response races, predator-prey shell games, ecology of fear,
6 and patch use of pumas and their ungulate prey. – *Ecology* 91: 2995-3007.

7 Liley, S. and Creel, S. 2008. What best explains vigilance in elk: characteristics of preys,
8 predators, or the environment? – *Behav. Ecol.* 19:245-254.

9 Lima, S. L. 2002. Putting predators back into behavioral predator-prey interactions. – *Trends*
10 *Ecol. Evol.* 17: 70-75.

11 Lima, S. L. and Dill, L. M. 1990. Behavioral decisions made under the risk of predation: a
12 review and prospectus. – *Can. J. Zoolog.* 68: 619-640.

13 Loveridge, A. J. et al. 2007. The impact of sport-hunting on the population dynamics of an
14 African lion population in a protected area. – *Biol. Conserv.* 134: 548-558.

15 Manly, B. F. J. et al. 2002. Resource selection by animals, statistical design and analysis for
16 field studies, 2nd edn. – Kluwer Academic.

17 Middleton, A. H. et al. 2013. Linking anti-predator behaviour to prey demography reveals
18 limited risk effects of an actively hunting large carnivore. – *Ecol. Lett.* 16: 1023-1030.

19 Padić, S. et al. In press. Roe deer at risk: teasing apart habitat selection and landscape
20 constraints in risk exposure at multiple scales. – *Oikos*. doi:10.1111/oik.02115.

21 Preisser, E. L. et al. 2005. Scared to death? The effects of intimidation and consumption in
22 predator-prey interactions. – *Ecology* 86: 501-509.

23 Preisser, E. L. et al. 2007. Predator hunting mode and habitat domain affect the strength of
24 non-consumptive effects in predator-prey interactions. – *Ecology* 88: 2744-2751.

1 R Core Team. 2013. R: A language and environment for statistical computing. R Foundation
2 for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>.

3 Rogers, C.M.L. 1993. A woody vegetation survey of Hwange National Park. Report for the
4 Department of National Parks and Wildlife Management Zimbabwe.

5 SAS Institute. 2008. SAS/STAT software: changes and enhancements, version 9.2. SAS
6 Institute Inc., Cary, North Carolina, USA.

7 Sih, A. 2005. Predator-prey space use as an emergent outcome of a behavioral response race.
8 – In: Barbosa, P. and Castellanos, I. (eds.), Ecology of predator-prey interactions. Oxford
9 University Press, pp. 240-255.

10 Stankowich, T. and Blumstein, D. T. 2005. Fear in animals: a meta-analysis and review of
11 risk assessment. – Proc. R. Soc. B 272: 2627-2634.

12 Thaker, M. et al. 2011. Minimizing predation risk in a landscape of multiple predators: effects
13 on the spatial distribution of African ungulates. – Ecology 92: 398-407.

14 Valeix, M. et al. 2009. Behavioral adjustments of African herbivores to predation risk by
15 lions: spatiotemporal variations influence habitat use. – Ecology 90: 23-30.

16 Valeix, M. et al. 2010. How key habitat features influence large terrestrial carnivore
17 movements: waterholes and African lions in a semi-arid savanna of north-western Zimbabwe.
18 – Landscape Ecol. 25: 337-351.

19 Valeix, M. et al. 2011. Understanding patch departure rules for large carnivores: lion
20 movements support a patch-disturbance hypothesis. – Am. Nat. 178: 269-275.

21 Wirsing, A. J. et al. 2010. Spatial responses to predators vary with prey escape mode. – Anim.
22 Behav. 79: 531-537.

23 Wojdak, J. M. 2009. Foraging and refuge use by a pond snail: effects of physiological state,
24 predators, and resources. – Acta Oecol. 35: 746-751.

1 Zанette, L. Y. et al. 2011. Perceived predation risk reduces the number of offspring songbirds
2 produce per year. – Science 334: 1398-1401.
3
4 Supplementary material (available as Appendix oik.XXXXXX at
5 <www.oikosjournal.org/readers/appendix>). Appendix 1–4.

1 Table 1. Coefficients (b) with 95% confidence intervals (CI) for mixed resource selection
 2 functions contrasting 1) features of areas occupied by zebra at the time of encounter ($n = 192$)
 3 and during proactive behaviour ($n = 1809$, baseline), and 2) features of areas occupied by
 4 zebra at the encounter time for immediate-flight ($n = 84$) and for delayed-flight tactics ($n =$
 5 108 , delayed-flight was the baseline).

6

Variable	Encounter vs. proactive		Immediate-flight vs. delayed-flight	
	b	95% CI	b	95% CI
Intercept	-2.26*	-2.79;-1.74	0.14	-0.91;1.18
Grassland	1.30*	0.50;2.11	-0.27	-1.88;1.33
Bushland-1	0.41	-0.33;1.16	-1.31†	-2.83;0.22
Bushland-2	-0.08	-0.99;0.84	-0.32	-2.35;1.70
Bushland-3	-1.78*	-3.50;-0.06	4.48*	0.19;8.77
Woodland	-1.08	-5.35;3.18	-2.27	-9.75;5.21
Distance to waterhole (km)	-0.04	-0.11;0.02	-0.05	-0.20;0.10

7 * 95% confidence intervals exclude zero.

8 † 90% confidence intervals exclude zero.

9 Notes: Variance and covariance of random intercept were <0.001 for the first model, and



10 equal to 0.04 and 0.19, respectively.

1 Figure legends.

2 Figure 1. Temporal dynamics of spatial large-scale movement patterns of zebras for each
3 reactive tactic with the median net displacement from the encounter site (A) and the median
4 distance to the closest waterhole (B) calculated at different occasions. Proactive behaviour of
5 zebras in the absence of lion is shown. The letter 'p' means that the changes in net
6 displacement between consecutive occasions or the distance to waterhole were different
7 between a given tactic and the proactive behaviour, while 't' indicates differences between
8 tactics.

9


10 Figure 2. Relative probability that zebras selected a given habitat type at the end of their step
11 after an encounter with lions (solid circle) at three occasions: immediately after an encounter
12 (A and D), during the next day (B and E), and during the next night (C and F), depending on
13 whether zebras used delayed-flight (A-C) or immediate-flight tactic (D-F). Proactive
14 behaviour of zebras in the absence of lion is shown (open circle). The 95% confidence
15 intervals are presented. Grey areas represent availability of respective habitats.



1

2

Figure 1.



1

2

Figure 2.