

Durham Research Online

Deposited in DRO:

08 September 2010

Version of attached file:

Accepted Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Krokhin, A. and Jeavons, P. and Jonsson, P. (2003) 'Reasoning about temporal relations : the maximal tractable subalgebras of Allen's interval algebra.', *Journal of the ACM.*, 50 (5). pp. 591-640.

Further information on publisher's website:

<http://dx.doi.org/10.1145/876638.876639>

Publisher's copyright statement:

© ACM, 2003. This is the author's version of the work. It is posted here by permission of ACM for your personal use. Not for redistribution. The definitive version was published in *Journal of the ACM*, Reasoning about temporal relations: The tractable subalgebras of Allen's interval algebra, 2003, <http://doi.acm.org/10.1145/380752.380868>.

Additional information:

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

Reasoning About Temporal Relations: The Tractable Subalgebras of Allen's Interval Algebra

Andrei Krokhin
Department of Computer Science
University of Warwick, UK
e-mail: Andrei.Krokhin@dcs.warwick.ac.uk

Peter Jeavons
Computing Laboratory
University of Oxford, UK
e-mail: Peter.Jeavons@comlab.ox.ac.uk

Peter Jonsson
Department of Computer and Information Science
Linköping University, Sweden
e-mail: Peter.Jonsson@ida.liu.se

September 6, 2010

Abstract

Allen's interval algebra is one of the best established formalisms for temporal reasoning. This paper is the final step in the classification of complexity in Allen's algebra. We show that the current knowledge about tractability in the interval algebra is complete, that is, this algebra contains exactly eighteen maximal tractable subalgebras, and reasoning in any fragment not entirely contained in one of these subalgebras is NP-complete. We obtain this result by giving a new uniform description of the known maximal tractable subalgebras and then systematically using an algebraic technique for identifying maximal subalgebras with a given property.

1 Introduction

Reasoning about temporal constraints is an important task in many areas of computer science and elsewhere, including scheduling [43], natural language processing [47], planning [2], database theory [31], technical diagnosis [41], circuit design [56], archaeology [29, 21], and behavioral psychology [11]; similar problems have been studied in genetics [7]. Several frameworks for formalizing this type of problem have been suggested (see [46] for a survey); for instance, the *point algebra* [52] (for expressing relations between time points), the *point-interval algebra* [54] (for expressing relations between time points and intervals) and the famous *Allen's interval algebra* [1] for expressing relations between time intervals.

Allen's algebra has also become the kernel of some other formalisms [3, 4, 13, 37], where it is extended with different types of metric or qualitative constraints. This algebra and some of its extensions are closely related to a number of interval-based temporal logics used for real-time system specification (see [6]). Reasoning within certain restricted fragments of Allen's algebra

(with additional restriction on the overall structure of problems) is equivalent to some well-known problems such as the *interval graph* recognition problem and the *interval order* recognition problem (see [44]) which play an important role in molecular biology [19, 28], namely in the construction of a physical mapping of DNA.

Throughout the paper we assume that $P \neq NP$. The basic satisfiability problem in Allen’s algebra is NP-complete [55], so it is unlikely that efficient algorithms exist for reasoning in the full algebra. This computational difficulty has motivated the study of algorithms and complexity in fragments of the algebra, e.g., [5, 13, 14, 15, 18, 21, 35, 36, 37, 40, 44, 51, 52, 55]¹, and the subsequent search for effective heuristics based on tractable fragments, e.g. [34, 39, 53]. In [40], Nebel and Bürckert presented the ‘ORD-Horn’ algebra, the first example of a maximal (assuming that $P \neq NP$) tractable subclass of Allen’s algebra. Since then, research in this direction has focused on identifying *maximal* tractable fragments, i.e., fragments which cannot be extended without losing tractability. So far, eighteen maximal tractable fragments of the algebra have been identified [13, 14, 36, 40]. In this paper we complete the analysis of complexity within Allen’s algebra by showing that these eighteen are the *only* forms of tractability in the algebra.

A complete classification of complexity within a certain large part of Allen’s algebra was previously obtained in [15]. This result (as well as most similar results, e.g. [26, 27]) was achieved by computer-assisted exhaustive search. However, it was noted in [15] that, for further progress, theoretical studies of the structure of Allen’s algebra are necessary, since using the method from that paper for a complete analysis of complexity would require dealing with more than 10^{50} individual cases, which is clearly not feasible. There have been some theoretical investigations of the structure of Allen’s algebra, (see, e.g., [23, 24, 33]); however they consider relation algebras in the sense defined by Tarski [50], that is, they generally allow more operations on relations than originally used in [1], which makes them inappropriate for classifying complexity within the interval algebra. In fact, none of the maximal tractable subalgebras of the interval algebra is a Tarski relation algebra. In this paper we systematically use algebraic methods that are similar to the approach taken in [36].

The first novel element in our approach is a new uniform description for all of the maximal tractable subalgebras of Allen’s algebra which have already been identified (Table 3). Then, we fully exploit the algebraic properties of Allen’s algebra by importing a technique from general algebra. This technique has been used in many other contexts to obtain a description of maximal subalgebras of a given algebra with a given property (e.g., [49, 58]). Here, for the first time, we systematically apply this technique to Allen’s algebra to obtain a complete classification of complexity in this algebra. Our main result (Theorem 1) shows that Allen’s algebra contains eighteen maximal tractable subalgebras and that reasoning within any subset not included in one of these is NP-complete.

In complexity theory, it is well known that if $P \neq NP$ then there exist infinitely many complexity classes between P and NP. In view of this, there has been a considerable interest in the so-called *dichotomy* theorems which state that one or another important NP-complete problem has only tractable and NP-complete natural subproblems (see, e.g., [12, ?, 22, 45]²). Thus, the main result obtained in this paper can also be considered as a new example of a dichotomy theorem.

The paper is organized as follows: in Section 2 we give the basic definitions of Allen’s algebra, present the known maximal tractable subalgebras in the new form, and state our main result. In Section 3 we apply this result to classify the complexity in Allen’s algebra extended with some metric information. In Section 4 we discuss the algebraic technique we use for obtaining results of this type and compare it with the computer-aided method used for a similar purpose in [15]. Sections 5 and 6 contain the proof of the new classification result—Section 5 considers the subalgebras of Allen’s algebra that contain non-trivial basic relations and Section 6 contains the proof for all other subalgebras. A number of NP-completeness results used in Section 6 are collected in the Appendix.

¹In [5, 21, 44], some additional restriction on the overall structure of problems is assumed.

²The problem of satisfiability from propositional logic [12, 45] should not be confused with the problem of satisfiability of temporal constraints.

Basic relation		Example	Endpoints
x precedes y	\mathbf{p}	xxx	$x^+ < y^-$
y preceded by x	\mathbf{p}^{-1}	yyy	
x meets y	\mathbf{m}	xxxx	$x^+ = y^-$
y met-by x	\mathbf{m}^{-1}	yyyy	
x overlaps y	\mathbf{o}	xxxx	$x^- < y^- < x^+$,
y overl.-by x	\mathbf{o}^{-1}	yyyy	$x^+ < y^+$
x during y	\mathbf{d}	xxx	$x^- > y^-$,
y includes x	\mathbf{d}^{-1}	yyyyyyy	$x^+ < y^+$
x starts y	\mathbf{s}	xxx	$x^- = y^-$,
y started by x	\mathbf{s}^{-1}	yyyyyyy	$x^+ < y^+$
x finishes y	\mathbf{f}	xxx	$x^+ = y^+$,
y finished by x	\mathbf{f}^{-1}	yyyyyyy	$x^- > y^-$
x equals y	\equiv	xxxx yyyy	$x^- = y^-$, $x^+ = y^+$

Table 1: The thirteen basic relations. The endpoint relations $x^- < x^+$ and $y^- < y^+$ that are valid for all relations have been omitted.

2 Allen's Interval Algebra

Allen's interval algebra [1] is based on the notion of *relations between intervals*. An interval x is represented as a pair $[x^-, x^+]$ of real numbers with $x^- < x^+$, denoting the left and right endpoints of the interval, respectively. The relations between intervals are the $2^{13} = 8192$ possible unions³ of the 13 *basic interval relations*, which are shown in Table 1. Note that the basic relations are jointly exhaustive and pairwise disjoint in the sense that any two given intervals are related by exactly one basic relation. For the sake of brevity, relations between intervals will be written as collections of basic relations. So, for instance, we write (\mathbf{pmf}^{-1}) instead of $\mathbf{p} \cup \mathbf{m} \cup \mathbf{f}^{-1}$. Allen's algebra \mathcal{A} consists of the 8192 possible relations between intervals together with the operations *converse* \cdot^{-1} , *intersection* \cap and *composition* \circ which are defined as follows:

$$\begin{aligned} \forall x, y : xr^{-1}y &\Leftrightarrow yrx \\ \forall x, y : x(r \cap s)y &\Leftrightarrow xry \ \& \ xsy \\ \forall x, y : x(r \circ s)y &\Leftrightarrow \exists z : (xrz \ \& \ zsy) \end{aligned}$$

It follows that the converse of $r = (b_1 \dots b_n)$ is equal to $(b_1^{-1} \dots b_n^{-1})$. The intersection of two relations can be expressed as the usual set-theoretic intersection. Since the basic relations are pairwise disjoint, the intersection of two relations $r_1, r_2 \in \mathcal{A}$ consists of the basic relations that are present in both r_1 and r_2 . Using the definition of composition, it can be shown that

$$(b_1 \dots b_n) \circ (b'_1 \dots b'_m) = \bigcup \{b_i \circ b'_j \mid 1 \leq i \leq n, 1 \leq j \leq m\}.$$

Hence the composition of two relations $r_1, r_2 \in \mathcal{A}$ is determined by the compositions of the basic relations they contain. The compositions of all possible pairs of basic relations are given in Table 2, and by using this table one can verify all the algebraic calculations in the forthcoming sections.

The problem of *satisfiability* (\mathcal{A} -SAT) for a set of interval variables with specified relations between them is that of deciding whether there exists an assignment of intervals on the real line for the interval variables, such that all of the relations between the intervals are satisfied. This is defined as follows.

Definition 1 *Let $X \subseteq \mathcal{A}$ be a set of interval relations. An instance I of \mathcal{A} -SAT(X) is a set, V , of variables and a set of constraints of the form xry where $x, y \in V$ and $r \in X$. The question is*

³Including the empty relation.

o	≡	p	p ⁻¹	m	m ⁻¹	o	o ⁻¹	d	d ⁻¹	s	s ⁻¹	f	f ⁻¹
≡	≡	p	p ⁻¹	m	m ⁻¹	o	o ⁻¹	d	d ⁻¹	s	s ⁻¹	f	f ⁻¹
p	p	p	⊤	p	ρ	p	ρ	ρ	p	p	p	ρ	p
p ⁻¹	p ⁻¹	⊤	p ⁻¹	λ ⁻¹	p ⁻¹	λ ⁻¹	p ⁻¹	λ ⁻¹	p ⁻¹	λ ⁻¹	p ⁻¹	p ⁻¹	p ⁻¹
m	m	p	ρ ⁻¹	p	θ	p	β	β	p	m	m	β	p
m ⁻¹	m ⁻¹	λ	p ⁻¹	σ	p ⁻¹	γ ⁻¹	p ⁻¹	γ ⁻¹	p ⁻¹	γ ⁻¹	p ⁻¹	m ⁻¹	m ⁻¹
o	o	p	ρ ⁻¹	p	β ⁻¹	α	ν	β	λ	o	γ	β	α
o ⁻¹	o ⁻¹	λ	p ⁻¹	γ	p ⁻¹	ν	α ⁻¹	γ ⁻¹	ρ ⁻¹	γ ⁻¹	α ⁻¹	o ⁻¹	β ⁻¹
d	d	p	p ⁻¹	p	p ⁻¹	ρ	λ ⁻¹	d	⊤	d	λ ⁻¹	d	ρ
d ⁻¹	d ⁻¹	λ	ρ ⁻¹	γ	β ⁻¹	γ	β ⁻¹	ν	d ⁻¹	γ	d ⁻¹	β ⁻¹	d ⁻¹
s	s	p	p ⁻¹	p	m ⁻¹	α	γ ⁻¹	d	λ	s	σ	d	α
s ⁻¹	s ⁻¹	λ	p ⁻¹	γ	m ⁻¹	γ	o ⁻¹	γ ⁻¹	d ⁻¹	σ	s ⁻¹	o ⁻¹	d ⁻¹
f	f	p	p ⁻¹	m	p ⁻¹	β	α ⁻¹	d	ρ ⁻¹	d	α ⁻¹	f	θ
f ⁻¹	f ⁻¹	p	ρ ⁻¹	m	β ⁻¹	o	β ⁻¹	β	d ⁻¹	o	d ⁻¹	θ	f ⁻¹

$$\begin{aligned}
\alpha &= (\text{pmo}) & \beta &= (\text{ods}) & \gamma &= (\text{od}^{-1}\text{f}^{-1}) & \sigma &= (\equiv \text{ss}^{-1}) & \theta &= (\equiv \text{ff}^{-1}) \\
\rho &= (\text{pmods}) & \lambda &= (\text{pmod}^{-1}\text{f}^{-1}) & \nu &= (\equiv \text{oo}^{-1}\text{dd}^{-1}\text{ss}^{-1}\text{ff}^{-1}) \\
\top &= (\equiv \text{pp}^{-1}\text{mm}^{-1}\text{oo}^{-1}\text{dd}^{-1}\text{ss}^{-1}\text{ff}^{-1})
\end{aligned}$$

Table 2: Composition table for the basic relations in Allen's algebra

whether I is satisfiable, i.e., whether there exists a function, f , from V to the set of all intervals such that $f(x) r f(y)$ holds for every constraint xry in I . Any such function f is called a model of I .

Example 1 1) The instance $\{x(\mathbf{m})y, y(\mathbf{m})z, x(\mathbf{m})z\}$ is not satisfiable because the first two constraints imply that interval x must precede interval z which contradicts the third constraint.

2) The instance $I = \{x(\mathbf{mo})y, y(\mathbf{df}^{-1})z, x(\mathbf{m}^{-1}\mathbf{s})z\}$ is satisfiable. The function f given by $f(x) = [0, 2]$, $f(y) = [1, 3]$, and $f(z) = [0, 4]$ is a model of I .

An instance of \mathcal{A} -SAT(X) can also be represented, in an obvious way, as a labelled digraph, where the nodes are the variables from V , and the labelled arcs correspond to the constraints. This way of representing instances can sometimes be more transparent.

If there exists a polynomial-time algorithm solving all instances of \mathcal{A} -SAT(X) then we say that X is tractable. On the other hand, if \mathcal{A} -SAT(X) is NP-complete then we say that X is NP-complete. Since the problem \mathcal{A} -SAT(\mathcal{A}) is NP-complete [55], there arises the question of identifying the tractable subsets of Allen's algebra.

Subsets of \mathcal{A} that are closed under the operations of intersection, converse and composition are said to be *subalgebras*. For a given subset X of \mathcal{A} , the smallest subalgebra containing X is called the subalgebra *generated* by X and is denoted by $\langle X \rangle$. It is easy to see that $\langle X \rangle$ is obtained from X by adding all relations that can be obtained from the relations in X by using the three operations of \mathcal{A} .

It is known [40], and easy to prove, that, for every $X \subseteq \mathcal{A}$, the problem \mathcal{A} -SAT($\langle X \rangle$) is polynomially equivalent to \mathcal{A} -SAT(X). Therefore, to classify the complexity of all subsets of \mathcal{A} it is only necessary to consider *subalgebras* of \mathcal{A} . Obviously, adding relations to a subalgebra can only increase the complexity of the corresponding satisfiability problem. Thus, since \mathcal{A} is finite, the problem of describing tractability in \mathcal{A} can be reduced to the problem of describing the *maximal* tractable subalgebras in \mathcal{A} , that is, subalgebras that cannot be extended without losing tractability.

The known maximal tractable subalgebras [13, 14, 40] are presented in Table 3. In this table, and in our proofs below, we use the symbol \pm , which should be interpreted as follows. A condition involving \pm means the conjunction of two conditions: one corresponding to $+$ and one corresponding to $-$. For example, condition $(\mathbf{o})^{\pm 1} \subseteq r \Leftrightarrow (\mathbf{d})^{\pm 1} \subseteq r$ means that both $(\mathbf{o}) \subseteq r \Leftrightarrow (\mathbf{d}) \subseteq r$ and $(\mathbf{o}^{-1}) \subseteq r \Leftrightarrow (\mathbf{d}^{-1}) \subseteq r$ hold. The main advantage of using the \pm symbol is conciseness: in any subalgebra of \mathcal{A} , the '+' and the '-' conditions are satisfied (or not satisfied) simultaneously, and, therefore only one of them needs to be verified.

In order to improve readability, the names of some of the subalgebras in Table 3 are changed from those used in earlier presentations, in the following way. Let $r_1 = (\mathbf{p}^{-1}\mathbf{m}^{-1}\mathbf{o}^{-1}\mathbf{dsf})$, $r_2 = (\mathbf{pmod}^{-1}\mathbf{sf}^{-1})$, $r_3 = (\mathbf{pmodsf})$, and $r_4 = (\mathbf{pmodsf}^{-1})$. Then, the subalgebras \mathcal{A}_i , $1 \leq i \leq 4$, from Table 3 correspond to the algebras $A(r_i, \mathbf{s})$, $1 \leq i \leq 4$, introduced in [14], while the subalgebras \mathcal{B}_i , $1 \leq i \leq 4$, from Table 3 correspond to $A(r_4^{-1}, \mathbf{f})$, $A(r_3, \mathbf{f})$, $A(r_1, \mathbf{f})$, and $A(r_2^{-1}, \mathbf{f})$ [14].

In previous papers, the subalgebras from Table 3 were defined in other ways. However, in all cases except for \mathcal{H} , it is very straightforward to verify that our definitions are equivalent to the original ones. The subalgebra \mathcal{H} was originally defined as the 'ORD-Horn algebra' [40], but has also been characterized as the algebra of 'pre-convex' relations [36]. Using the latter description it is not hard to show that our definition of \mathcal{H} is equivalent.

We are now ready to state our main theorem.

Theorem 1 *Any subset of Allen's algebra is either NP-complete or included in one of the eighteen tractable subalgebras in Table 3.*

The proof of this theorem is given in Sections 5 and 6.

As one interesting consequence of Theorem 1, it follows that reasoning with a single relation r , that is, the problem \mathcal{A} -SAT($\{r\}$), is NP-complete if and only if r either satisfies $r \cap r^{-1} = (\mathbf{mm}^{-1})$ or is a relation with $r \cap r^{-1} = \emptyset$ and such that neither r nor r^{-1} is contained in one of $(\mathbf{pmod}^{-1}\mathbf{sf}^{-1})$, $(\mathbf{pmod}^{-1}\mathbf{s}^{-1}\mathbf{f}^{-1})$, (\mathbf{pmodsf}) and (\mathbf{pmodsf}^{-1}) . Using this characterisation it is easy to check that there are precisely 667 individual temporal relations r such that \mathcal{A} -SAT($\{r\}$) is NP-complete.

$$\begin{aligned}
\mathcal{S}_p &= \{r \mid r \cap (\text{pmod}^{-1}f^{-1})^{\pm 1} \neq \emptyset \Rightarrow (p)^{\pm 1} \subseteq r\} \\
\mathcal{S}_d &= \{r \mid r \cap (\text{pmod}^{-1}f^{-1})^{\pm 1} \neq \emptyset \Rightarrow (d^{-1})^{\pm 1} \subseteq r\} \\
\mathcal{S}_o &= \{r \mid r \cap (\text{pmod}^{-1}f^{-1})^{\pm 1} \neq \emptyset \Rightarrow (o)^{\pm 1} \subseteq r\} \\
\mathcal{A}_1 &= \{r \mid r \cap (\text{pmod}^{-1}f^{-1})^{\pm 1} \neq \emptyset \Rightarrow (s^{-1})^{\pm 1} \subseteq r\} \\
\mathcal{A}_2 &= \{r \mid r \cap (\text{pmod}^{-1}f^{-1})^{\pm 1} \neq \emptyset \Rightarrow (s)^{\pm 1} \subseteq r\} \\
\mathcal{A}_3 &= \{r \mid r \cap (\text{pmod}f)^{\pm 1} \neq \emptyset \Rightarrow (s)^{\pm 1} \subseteq r\} \\
\mathcal{A}_4 &= \{r \mid r \cap (\text{pmod}f^{-1})^{\pm 1} \neq \emptyset \Rightarrow (s)^{\pm 1} \subseteq r\}
\end{aligned}$$

$$\begin{aligned}
\mathcal{E}_p &= \{r \mid r \cap (\text{pmods})^{\pm 1} \neq \emptyset \Rightarrow (p)^{\pm 1} \subseteq r\} \\
\mathcal{E}_d &= \{r \mid r \cap (\text{pmods})^{\pm 1} \neq \emptyset \Rightarrow (d)^{\pm 1} \subseteq r\} \\
\mathcal{E}_o &= \{r \mid r \cap (\text{pmods})^{\pm 1} \neq \emptyset \Rightarrow (o)^{\pm 1} \subseteq r\} \\
\mathcal{B}_1 &= \{r \mid r \cap (\text{pmods})^{\pm 1} \neq \emptyset \Rightarrow (f^{-1})^{\pm 1} \subseteq r\} \\
\mathcal{B}_2 &= \{r \mid r \cap (\text{pmods})^{\pm 1} \neq \emptyset \Rightarrow (f)^{\pm 1} \subseteq r\} \\
\mathcal{B}_3 &= \{r \mid r \cap (\text{pmod}^{-1}s^{-1})^{\pm 1} \neq \emptyset \Rightarrow (f^{-1})^{\pm 1} \subseteq r\} \\
\mathcal{B}_4 &= \{r \mid r \cap (\text{pmod}^{-1}s)^{\pm 1} \neq \emptyset \Rightarrow (f^{-1})^{\pm 1} \subseteq r\}
\end{aligned}$$

$$\mathcal{E}^* = \left\{ r \left| \begin{array}{l} 1) r \cap (\text{pmod})^{\pm 1} \neq \emptyset \Rightarrow (s)^{\pm 1} \subseteq r, \text{ and} \\ 2) r \cap (\text{ff}^{-1}) \neq \emptyset \Rightarrow (\equiv) \subseteq r \end{array} \right. \right\}$$

$$\mathcal{S}^* = \left\{ r \left| \begin{array}{l} 1) r \cap (\text{pmod}^{-1})^{\pm 1} \neq \emptyset \Rightarrow (f^{-1})^{\pm 1} \subseteq r, \text{ and} \\ 2) r \cap (\text{ss}^{-1}) \neq \emptyset \Rightarrow (\equiv) \subseteq r \end{array} \right. \right\}$$

$$\mathcal{H} = \left\{ r \left| \begin{array}{l} 1) r \cap (\text{os})^{\pm 1} \neq \emptyset \ \& \ r \cap (\text{o}^{-1}f)^{\pm 1} \neq \emptyset \Rightarrow (d)^{\pm 1} \subseteq r, \text{ and} \\ 2) r \cap (\text{ds})^{\pm 1} \neq \emptyset \ \& \ r \cap (\text{d}^{-1}f^{-1})^{\pm 1} \neq \emptyset \Rightarrow (o)^{\pm 1} \subseteq r, \text{ and} \\ 3) r \cap (\text{pm})^{\pm 1} \neq \emptyset \ \& \ r \not\subseteq (\text{pm})^{\pm 1} \Rightarrow (o)^{\pm 1} \subseteq r \end{array} \right. \right\}$$

$$\mathcal{A}_{\equiv} = \{r \mid r \neq \emptyset \Rightarrow (\equiv) \subseteq r\}$$

Table 3: The 18 maximal tractable subalgebras of Allen's algebra.

3 Allen's Interval Algebra Extended With Metric Information

In this section we give some applications of Theorem 1. Namely, we consider Allen's algebra combined with some forms of *disjunctive linear constraints*, a well-known framework which subsumes many different types of temporal reasoning problems. Some examples of these problems, including scheduling, planning, and indefinite temporal constraint databases, can be found in [25, 30, 48] (see also [10] for more information on tractable disjunctive constraints).

Definition 2 Let $V = \{x_1, \dots, x_n\}$ be a set of real-valued variables, and α, β linear polynomials (polynomials of degree one) over V with rational coefficients. A linear relation over V is an expression of the form $\alpha R \beta$, where $R \in \{<, \leq, =, \neq, \geq, >\}$.

A disjunctive linear relation (DLR) over V is a disjunction of a nonempty finite set of linear relations. A DLR is said to be Horn if and only if at most one of its disjuncts is not of the form $\alpha \neq \beta$.

The problem of satisfiability for finite sets D of DLRs, denoted $\text{DLRSAT}(D)$, is that of checking whether there exists an assignment f of variables in V to real numbers such that all DLRs in D are satisfied. Such an f is said to be a model of D . The satisfiability problem for finite sets H of Horn DLRs is denoted $\text{HORNDLRSAT}(H)$.

Example 2

$$x + 2y \leq 3z + 42.3$$

is a linear relation,

$$(x + 2y \leq 3z + 42.3) \vee (x + z < 4y - 8) \vee (x > \frac{3}{12})$$

is a disjunctive linear relation, and

$$(x + 2y \leq 3z + 42.3) \vee (x + z \neq 4y - 8) \vee (x \neq \frac{3}{12})$$

is a Horn disjunctive linear relation.

Proposition 1 ([25, 30]) The problem DLRSAT is NP-complete and HORNDLRSAT is solvable in polynomial time.

We can now define the general interval satisfiability problem with metric information.

Definition 3 Let I be an instance of $\mathcal{A}\text{-SAT}(X)$ over a set V of variables and let H be a finite set of DLRs over the set $\{v^+, v^- \mid v \in V\}$ of variables, v^- representing starting points and v^+ ending points of variables $v \in V$.

An instance of the problem of interval satisfiability with metric information for a set X of interval relations, denoted $\mathcal{A}^m\text{-SAT}(X)$, is a pair $Q = (I, H)$.

If f is a model for I , and $v \in V$, let $f(v^-)$ and $f(v^+)$ denote the starting point and the ending point of the interval $f(v)$, respectively.

An instance Q is said to be satisfiable if there exists a model f of I such that the DLRs in H are satisfied, with values for all v^- and v^+ given by $f(v^-)$ and $f(v^+)$, respectively.

Obviously, the $\mathcal{A}^m\text{-SAT}(X)$ problem is NP-complete for all choices of X since every relation in \mathcal{A} can be expressed in terms of DLRs. We let $\mathcal{A}^h\text{-SAT}(X)$ denote the $\mathcal{A}^m\text{-SAT}(X)$ restricted to metric constraints consisting of Horn DLRs only.

Theorem 2 $\mathcal{A}^h\text{-SAT}(X)$ is tractable if and only if $X \subseteq \mathcal{H}$. Otherwise $\mathcal{A}^h\text{-SAT}(X)$ is NP-complete.

Proof. $\mathcal{A}^h\text{-SAT}(\mathcal{H})$ is a tractable problem [25]. The interval constraint $a(\mathbf{m})b$ is equivalent to the metric constraint $a^+ = b^-$ so we can assume that $(\mathbf{m}) \in X$ and the result follows immediately from Theorem 1. \square

We can obtain another classification result if we further restrict the possible metric constraints. Define $\mathcal{A}_s^h\text{-SAT}(X)$ to be the $\mathcal{A}^h\text{-SAT}(X)$ problem where the metric constraints H are restricted in the following way: H may contain only the variables v^- , i.e., it may only relate starting points of intervals. The problem $\mathcal{A}_e^h\text{-SAT}(X)$ is defined symmetrically by exchanging starting and ending points.

Theorem 3 1) $\mathcal{A}_s^h\text{-SAT}(X)$ is tractable if and only if X is contained in one of the algebras \mathcal{H} , \mathcal{S}_p , \mathcal{S}_o , \mathcal{S}_d or \mathcal{S}^* . Otherwise $\mathcal{A}_s^h\text{-SAT}(X)$ is NP-complete.

2) $\mathcal{A}_e^h\text{-SAT}(X)$ is tractable if and only if X is contained in one of the algebras \mathcal{H} , \mathcal{E}_p , \mathcal{E}_o , \mathcal{E}_d or \mathcal{E}^* . Otherwise $\mathcal{A}_e^h\text{-SAT}(X)$ is NP-complete.

Proof. We prove only part 1); part 2) is similar. If X is contained in one of the five algebras listed in 1) then tractability of $\mathcal{A}_s^h\text{-SAT}(X)$ follows from [13]. The interval constraints $a(\equiv \mathbf{ss}^{-1})b$ and $a(\mathbf{pmod}^{-1}\mathbf{f}^{-1})b$ are equivalent to the metric constraints $a^- = b^-$ and $a^- < b^-$, respectively. Thus, we can assume that $\{(\equiv \mathbf{ss}^{-1}), (\mathbf{pmod}^{-1}\mathbf{f}^{-1})\} \subseteq X$. It follows from Theorem 1 that if $\mathcal{A}\text{-SAT}(X)$ is tractable then X is contained in one of the five algebras listed in 1), and that otherwise this problem is NP-complete. \square

4 Proof Techniques

In this section we describe the algebraic techniques used in this paper, and the methods for proving NP-completeness results.

In contrast to earlier approaches [15, 26, 27] we do not make use of computer-assisted exhaustive search. Instead, we develop an analytical method which breaks the proof down into a collection of simple cases, and makes extensive use of the algebraic operations. This approach is commonly used in general algebra to identify those substructures of a given structure that have a property ϕ which is *hereditary*, that is, if some substructure possesses ϕ then so does any substructure contained in it. Note that tractability of a subalgebra is an example of such a property in Allen's algebra. For examples of a similar approach in other algebraic contexts see [49, 57, 58].

As indicated above, it is sufficient to consider only those sets, \mathcal{S} , which are subalgebras of Allen's algebra. Furthermore, we can assume without loss of generality that each subalgebra \mathcal{S} contains the relation \top (the union of all basic relations) since we always allow pairs of variables to be unrelated. For each basic relation b of \mathcal{A} , we will write r_b to denote the least relation $r \in \mathcal{S}$ such that $(b) \subseteq r$, i.e., the intersection of all $r \in \mathcal{S}$ with this property. (Obviously, the relations r_b depend on \mathcal{S} ; however \mathcal{S} will always be clear from the context.)

We use the relations of the form r_b extensively in the algebraic proofs below to show that \mathcal{S} is contained in one or another maximal tractable subalgebra. For example, suppose we know that the relation (\mathbf{d}) is contained in r_o . Then any relation $r \in \mathcal{S}$ such that $(\mathbf{o}) \subseteq r$ satisfies also $(\mathbf{d}) \subseteq r$. To see this, note that if there is $r_1 \in \mathcal{S}$ such that $(\mathbf{o}) \subseteq r_1$, but $(\mathbf{d}) \not\subseteq r_1$, then $(\mathbf{o}) \subseteq r_1 \cap r_o$ and $r_1 \cap r_o$ is strictly contained in r_o which contradicts the definition of r_o . By a similar argument, if we know that (\mathbf{d}) is contained in all of r_p , r_m , r_o , and r_s , then we can conclude that, for every $r \in \mathcal{S}$, $(\mathbf{d}) \subseteq r$ whenever $r \cap (\mathbf{pmods}) \neq \emptyset$, which means that $\mathcal{S} \subseteq \mathcal{E}_d$.

Throughout the proofs we also use the obvious fact that if $r_1 \subseteq r_2$ then, for any r , we have $r \circ r_1 \subseteq r \circ r_2$ and $r_1 \circ r \subseteq r_2 \circ r$.

To establish NP-completeness of a set of relations we will often make use of Lemma 1 below. For any given relations $R, R_1, R_2 \in \mathcal{A}$ we define $\Gamma(a, b, c, x, y)$ to be the following problem instance over the variables $\{a, b, c, x, y\}$:

$$\{xR_1a, xR_1b, xR_2c, yR_2a, yR_1b, yR_1c\}.$$

We also define the instances $\Gamma_1 = \Gamma(a, b, c, x, y) \cup \{aRb, bRc, aRc\}$, $\Gamma_2 = \Gamma(a, b, c, x, y) \cup \{bRa, bRc, aR \cup R^{-1}c\}$, and $\Gamma_3 = \Gamma(a, b, c, x, y) \cup \{aRb, cRb, aR \cup R^{-1}c\}$. The problem instance Γ_1 is illustrated in Figure 1.




Figure 1: The problem instance Γ_1 used in Lemma 1

Lemma 1 *Let $R \in \{(p), (o), (d), (s), (f)\}$, $R_1, R_2 \in \mathcal{A}$, and let $\Gamma_1, \Gamma_2, \Gamma_3$ be as above. If Γ_1 is satisfiable while Γ_2 and Γ_3 are not, then $\{R \cup R^{-1}, R_1, R_2\}$ is NP-complete.*

Proof. Polynomial-time reduction from the NP-complete problem BETWEENNESS⁴ [16], which is defined as follows:

INSTANCE: A finite set A , a collection T of ordered triples (a, b, c) of distinct elements from A .

QUESTION: Is there a total ordering $<$ on A such that for each $(a, b, c) \in T$, we have either $a < b < c$ or $c < b < a$?

Let (A, T) be an arbitrary instance of BETWEENNESS and construct an instance I of $\mathcal{A}\text{-SAT}(\{R \cup R^{-1}, R_1, R_2\})$ as follows:

- (1) for each pair of distinct elements $a, b \in A$, add the constraint $aR \cup R^{-1}b$ to I ; and
- (2) for each triple $(a, b, c) \in T$, introduce two fresh variables x, y and add $\Gamma(a, b, c, x, y)$ to I .

We will henceforth refer to the variables in I that correspond to the set A as ‘basic’ variables and the other variables as ‘auxiliary’ variables.

Assume that I has a model f . Then, due to the constraints added in step (1), the intervals $f(a), a \in A$, are pairwise distinct. Moreover, the relation R induces a total order on the set $\{f(a) \mid a \in A\}$. Suppose now that there is a triple $(a, b, c) \in T$ such that the model f satisfies

⁴This problem is also known as the TOTAL ORDERING PROBLEM [42].

$f(b) R f(a) R f(c)$. Then the instance Γ_2 , with the auxiliary variables x and y introduced in step (2) for the triple (a, b, c) , is satisfiable, a contradiction. With the help of Γ_2 and Γ_3 , we can analogously rule out all orderings of $f(a), f(b), f(c)$ except $f(a) R f(b) R f(c)$ and $f(c) R f(b) R f(a)$. Hence there is a solution to the instance (A, T) : for all $a, b \in A$, set $a < b$ if and only if $f(a) R f(b)$.

We assume now that there exists a total order $<$ on A that has the required property and show how to construct a model f of I . For all $a, b \in A$, set $f(a) R f(b)$ if and only if $a < b$. Clearly, this satisfies all constraints added in step (1).

By assumption, for each triple $(a, b, c) \in T$, the instance Γ_1 is satisfiable, that is, it has some model g . Then, since R is a basic relation, we know precisely how the intervals $g(a), g(b)$, and $g(c)$ are related. It now follows that the model g can be adjusted by moving the intervals $g(a), g(b), g(c), g(x)$, and $g(y)$ along the real line and stretching or shrinking them (but without changing the relations between them) so that the new model assigns $f(a), f(b)$, and $f(c)$ to a, b , and c , respectively. The fact that $\Gamma_1 = \Gamma(a, b, c, x, y) \cup \{aRb, bRc, aRc\}$ is satisfiable and the symmetry of Γ imply that $\Gamma(a, b, c, x, y) \cup \{cRb, bRa, cRa\}$ is satisfiable (by exchanging the value for a by the value for c and vice versa, and doing the same for x and y). Thus, for every triple $(a, b, c) \in T$, we can find values for the auxiliary variables x and y so that all sets of constraints of the form $\Gamma(a, b, c, x, y)$ are satisfied at the same time, and there exists a model of I . \square

In order to use Lemma 1 to prove NP-completeness of some fixed set of relations, one only needs to check the satisfiability of three small instances of \mathcal{A} -SAT. One straightforward way to do this is to use B. Nebel's CSP solver [38], which is a computer program for checking satisfiability of an instance of \mathcal{A} -SAT.

As an example of the use of Lemma 1, set $R = (\circ)$, $R_1 = (\text{d})$ and $R_2 = (\circ\circ^{-1})$. In Figure 2, we show how the auxiliary variables x and y can be given consistent values in the two 'allowed' cases (corresponding to $f(a) < f(b) < f(c)$ and $f(c) < f(b) < f(a)$) and the reader is encouraged to prove that x and y cannot be chosen satisfactorily for the remaining four orderings. Thus, Lemma 1 implies that $\{(\text{d}), (\circ\circ^{-1})\}$ is NP-complete.

The second method we use to establish NP-completeness is based on the notion of *derivation*. Suppose $X \subseteq \mathcal{A}$ and I is an instance of \mathcal{A} -SAT(X). Let variables x, y be involved in I . Further, let $r \in \mathcal{A}$ be the relation defined as follows: a basic relation r' is included in r if and only if the instance obtained from I by adding the constraint $xr'y$ is satisfiable. In this case, we say that r is *derived* from X .

It should be noted that if the instance $I_1 = I \cup \{xr'y\}$ is satisfiable, then, for any two intervals i_1, j_1 such that $i_1 r' j_1$, there is a model f of I_1 such that $f(x) = i_1$ and $f(y) = j_1$. This can be established as follows: since I_1 is satisfiable, it has a model g . Denote $g(x)$ by i_2 and $g(y)$ by j_2 ; then $i_2 r' j_2$. There exists a continuous monotone injective mapping φ of the real line such that φ takes i_2 to i_1 and j_2 to j_1 . Obviously, φ maps intervals to intervals, and it does not change the qualitative relations between intervals. Therefore, combining φ and g we obtain the the required model f .

Now it can easily be checked that adding a derived relation r to X does not change the complexity of \mathcal{A} -SAT(X) because, in any instance, any constraint involving r can be replaced by the set of constraints in I (introducing fresh variables when needed), and this can be done in polynomial time.

Generally, one can derive more relations from a given $X \subseteq \mathcal{A}$ than one can generate using the three operations of Allen's algebra, that is, any relation generated from X can also be derived from X . This follows from the facts that any relation obtained by multiple derivations can also be obtained by a single derivation, and that relations r^{-1} , $r_1 \cap r_2$, and $r_1 \circ r_2$, between x and y are derived from the instances $\{y r x\}$, $\{x r_1 y, x r_2 y\}$, and $\{x r_1 z, z r_2 y\}$, respectively. However, derivation is essentially harder to manage in general, while the operations of Allen's algebra give us the advantage of employing algebraic techniques. Therefore we use derivations only in NP-completeness proofs. Note that derivations can also be calculated using B. Nebel's CSP solver [38].

Our last proof technique is a principle of *duality*, which will be used to simplify many of the forthcoming proofs. We make use of a function *reverse* which is defined on the basic relations of





Figure 2: Example of using Lemma 1.

\mathcal{A} by the following table:

b	\equiv	p	p^{-1}	m	m^{-1}	o	o^{-1}	d	d^{-1}	s	s^{-1}	f	f^{-1}
$\text{reverse}(b)$	\equiv	p^{-1}	p	m^{-1}	m	o^{-1}	o	d	d^{-1}	f	f^{-1}	s	s^{-1}

and is defined for all other elements of \mathcal{A} by setting $\text{reverse}(r) = \bigcup_{b \in r} \text{reverse}(b)$.

Let I be any instance of \mathcal{A} -SAT, and let I' be obtained from I by replacing every r with $\text{reverse}(r)$. It is easy to check that I has a model f if and only if I' has a model f' given by

$$f'(x_i) = [-f(x_i^+), -f(x_i^-)].$$

In other words, f' is obtained from f by redirecting the real line and leaving all intervals (as geometric objects) in their places. This observation leads to the following lemma.

Lemma 2 *Let $\mathcal{R} = \{r_1, \dots, r_n\} \subseteq \mathcal{A}$ and $\mathcal{R}' = \{r'_1, \dots, r'_n\} \subseteq \mathcal{A}$ be such that, for all $1 \leq k \leq n$, $r'_k = \text{reverse}(r_k)$. Then \mathcal{R} is tractable (NP-complete) if and only if \mathcal{R}' is tractable (NP-complete).*

As an example of the use of Lemma 2, note that a proof of NP-completeness for, say, $\{(\text{ods}^{-1})\}$, immediately yields a proof of NP-completeness for $\{(\text{o}^{-1}\text{df}^{-1})\}$.

5 Subalgebras With Non-trivial Basic Relations

This section and the next contain the proof of Theorem 1.

For a subalgebra \mathcal{S} of \mathcal{A} , we denote by $\text{bas}(\mathcal{S})$ the set of basic relations in \mathcal{S} . We can assume without loss of generality that \mathcal{S} contains the relation (\equiv) , since it is easy to show that \mathcal{S} and $\mathcal{S} \cup \{(\equiv)\}$ have the same complexity (up to polynomial-time equivalence). This implies that the size of $\text{bas}(\mathcal{S})$ is odd, since \mathcal{S} is closed under converse.

The following proposition is proved in [15].

Proposition 2

- 1) *Let \mathcal{S} be a subalgebra of \mathcal{A} with $|\text{bas}(\mathcal{S})| > 3$. Then \mathcal{S} is tractable if it is contained in one of the following 7 algebras: \mathcal{S}_p , \mathcal{S}_d , \mathcal{S}_o , \mathcal{E}_p , \mathcal{E}_d , \mathcal{E}_o , and \mathcal{H} . Otherwise \mathcal{S} is NP-complete.*
- 2) *Let \mathcal{S} be a subalgebra of \mathcal{A} such that $(m) \in \mathcal{S}$ or $(p) \in \mathcal{S}$. Then \mathcal{S} is tractable if $\mathcal{S} \subseteq \mathcal{S}_p$, or $\mathcal{S} \subseteq \mathcal{E}_p$, or $\mathcal{S} \subseteq \mathcal{H}$. Otherwise \mathcal{S} is NP-complete.*
- 3) *Let \mathcal{S} be a subalgebra of \mathcal{A} such that $(pp^{-1}) \in \mathcal{S}$ or $(pp^{-1}mm^{-1}) \in \mathcal{S}$. Then \mathcal{S} is tractable if $\mathcal{S} \subseteq \mathcal{S}_p$ or $\mathcal{S} \subseteq \mathcal{E}_p$. Otherwise \mathcal{S} is NP-complete.*

We shall say that a relation is *non-trivial* if it is not equal to the empty relation or the relation (\equiv) . The result to be shown in this section is the following:

Proposition 3 *Let \mathcal{S} be a subalgebra of \mathcal{A} which contains a non-trivial basic relation. Then \mathcal{S} is tractable if it is contained in one of the 18 algebras listed in Table 3. Otherwise \mathcal{S} is NP-complete.*

Note that if \mathcal{S} contains a non-trivial basic relation, then $\mathcal{S} \not\subseteq \mathcal{A}_{\equiv}$.

By combining Proposition 2(1) and the observation preceding it, it suffices to consider only the case $|\text{bas}(\mathcal{S})| = 3$. By Proposition 2(2), it suffices to consider the cases where $\text{bas}(\mathcal{S})$ is one of the following sets: $\{\equiv, d, d^{-1}\}$, $\{\equiv, o, o^{-1}\}$, $\{\equiv, s, s^{-1}\}$, and $\{\equiv, f, f^{-1}\}$. We do this in Subsections 5.1–5.3.

Given a relation r , we write r^* to denote the relation $r \cap r^{-1}$. Evidently, every subalgebra of \mathcal{A} is closed under the operation \cdot^* (of taking the symmetric part of a relation). By $\text{sym}(\mathcal{S})$ we denote the set $\{r \in \mathcal{S} \mid r^* = r\}$.

5.1 The case $bas(\mathcal{S}) = \{\equiv, d, d^{-1}\}$

In this subsection, we will show that if \mathcal{S} is a subalgebra with $bas(\mathcal{S}) = \{\equiv, d, d^{-1}\}$, then $\mathcal{S} \subseteq \mathcal{S}_d$, $\mathcal{S} \subseteq \mathcal{E}_d$ or \mathcal{S} is NP-complete.

To obtain this result we shall assume throughout this subsection that \mathcal{S} is a subalgebra of \mathcal{A} satisfying the following assumptions:

Assumption 1 $bas(\mathcal{S}) = \{\equiv, d, d^{-1}\}$.

Assumption 2 \mathcal{S} is not NP-complete.

Using these assumptions we obtain increasingly detailed information about \mathcal{S} in Lemmas 4-12, until we are able to show that in all cases $\mathcal{S} \subseteq \mathcal{S}_d$ or $\mathcal{S} \subseteq \mathcal{E}_d$. These lemmas rely on the following NP-completeness result.

Lemma 3 *The subsets $\{(d), (oo^{-1})\}$ and $\{(d^{-1}), (pp^{-1})\}$ of \mathcal{A} are NP-complete.*

Proof. Apply Lemma 1 with $R = (o)$, $R_1 = (d)$, $R_2 = (oo^{-1})$, or with $R = (p)$, $R_1 = (d^{-1})$, $R_2 = (pp^{-1})$, respectively. \square

Before we give the proofs, we note that $\nu = (d^{-1}) \circ (d) \in \mathcal{S}$, where $\nu = (\equiv oo^{-1}dd^{-1}ss^{-1}ff^{-1})$, as defined in Table 2.

Lemma 4 *With the assumptions above,*

$$sym(\mathcal{S}) \subseteq \{r^* | (dd^{-1}) \subseteq r\} \cup \{r^* | r \subseteq (\equiv ss^{-1})\} \cup \{r^* | r \subseteq (\equiv ff^{-1})\}.$$

Proof. We will show that if $sym(\mathcal{S})$ is not included in the above set then \mathcal{S} is NP-complete, which contradicts Assumption 2.

Suppose first that $(\equiv ss^{-1}ff^{-1})$ or $(ss^{-1}ff^{-1})$ belongs to $sym(\mathcal{S})$. Then

$$r = (\equiv ss^{-1}ff^{-1}) \circ (d) = (ss^{-1}ff^{-1}) \circ (d) = (oo^{-1}dsf) \in \mathcal{S},$$

so we have $(oo^{-1}) = r^* \in \mathcal{S}$, which implies that \mathcal{S} is NP-complete, by Lemma 3.

Suppose now there is $r^* \in sym(\mathcal{S})$ such that $r^* \subseteq (\equiv pp^{-1}mm^{-1}oo^{-1}ss^{-1}ff^{-1})$ and $r^* \not\subseteq (\equiv ss^{-1}ff^{-1})$. Consider $r_1 = (d) \circ r^* \subseteq (pp^{-1}mm^{-1}oo^{-1}dsf)$. It is easy to check that r_1^* is non-empty and $r_1^* \subseteq (pp^{-1}mm^{-1}oo^{-1})$. If $(oo^{-1}) \subseteq r_1^*$, then $r_1^* \cap \nu = (oo^{-1}) \in \mathcal{S}$, so \mathcal{S} is NP-complete, by Lemma 3. Otherwise $r_1^* \subseteq (pp^{-1}mm^{-1})$, so $(d) \circ r_1^* = (pp^{-1}) \in \mathcal{S}$, which again implies that \mathcal{S} is NP-complete, by Lemma 3. \square

Lemma 5 *With the assumptions above,*

$$\mathcal{S} \subseteq \{r | r \cap (dd^{-1}) \neq \emptyset\} \cup \{r | r \subseteq (\equiv ss^{-1})\} \cup \{r | r \subseteq (\equiv ff^{-1})\}.$$

Proof. Assume for contradiction that \mathcal{S} contains a relation r with $r \cap (dd^{-1})$ empty, $r \not\subseteq (\equiv ss^{-1})$ and $r \not\subseteq (\equiv ff^{-1})$.

Among such relations, choose r to be minimal with respect to inclusion. Then, since $\nu \in \mathcal{S}$, we have either $r \subseteq (\equiv oo^{-1}ss^{-1}ff^{-1})$, or $r \subseteq (\equiv pp^{-1}mm^{-1}ss^{-1})$, or $r \subseteq (\equiv pp^{-1}mm^{-1}ff^{-1})$.

Case 1. $r \subseteq (\equiv oo^{-1}ss^{-1}ff^{-1})$

Assume first that $(o) \subseteq r$ (the argument for (o^{-1}) is dual by using Lemma 2). Consider $r_1 = (d^{-1}) \circ r \subseteq (oo^{-1}d^{-1}s^{-1}f^{-1})$. By the minimality of r , $r_1 \cap r = r$ or empty, but $(o) \subseteq r$, so $(o) \subseteq r_1$, so $r_1 \cap r$ is not empty and hence $r \subseteq (oo^{-1}s^{-1}f^{-1})$. Now consider $r_2 = (d) \circ r \subseteq (pp^{-1}mm^{-1}oo^{-1}dsf)$. By a similar argument we get $r \subseteq (oo^{-1}sf)$. Combining these two results gives $r \subseteq (oo^{-1})$, and then by Lemma 4 we get $r = (o)$, which contradicts Assumption 1.

Hence, we must have $r \subseteq (\equiv ss^{-1}ff^{-1})$. If $(sf) \subseteq r$ then $((d^{-1}) \circ r)^* = (oo^{-1}) \in \mathcal{S}$, which contradicts Lemma 4. If $r = (\equiv sf^{-1})$ or $r = (sf^{-1})$ then $r \cap ((d) \circ r) = (s) \in \mathcal{S}$, which contradicts Assumption 1. Hence, case 1 is impossible.

Case 2. $r \subseteq (\equiv \mathbf{pp}^{-1}\mathbf{mm}^{-1}\mathbf{ss}^{-1})$.

If $r \cap (\mathbf{pp}^{-1}) \neq \emptyset$ then $r_1 = r \cap ((\mathbf{d}) \circ r) \subseteq (\mathbf{pp}^{-1}\mathbf{m}^{-1})$. Furthermore, $(\mathbf{d}) \circ r_1$ is a nonempty subrelation of (\mathbf{pp}^{-1}) , which contradicts Assumption 1 or Lemma 3. If $(\mathbf{mm}^{-1}) \subseteq r$ then $((\mathbf{d}) \circ r)^* = (\mathbf{pp}^{-1})$, a contradiction again. It remains to consider the case $(\mathbf{m}) \subseteq r \subseteq (\equiv \mathbf{mss}^{-1})$. If $(\mathbf{s}^{-1}) \subseteq r$ then, again, $((\mathbf{d}) \circ r)^* = (\mathbf{pp}^{-1}) \in \mathcal{S}$. Otherwise $r_2 = r \circ r$ satisfies $(\mathbf{p}) \subseteq r_2 \subseteq (\equiv \mathbf{pms})$, and then $(\mathbf{p}) = r_2 \cap ((\mathbf{d}) \circ r_2) \in \mathcal{S}$, which contradicts Assumption 1.

Case 3. $r \subseteq (\equiv \mathbf{pp}^{-1}\mathbf{mm}^{-1}\mathbf{ff}^{-1})$.

Dual to Case 2. □

Lemma 6 *With the assumptions above, if $(\mathbf{pmod}) \in \mathcal{S}$ then $\mathcal{S} \subseteq \mathcal{E}_d$; if $(\mathbf{pmod}^{-1}\mathbf{f}^{-1}) \in \mathcal{S}$ then $\mathcal{S} \subseteq \mathcal{S}_d$.*

Proof. We prove only the first statement, the second one is dual.

First note that if $r_s \subseteq (\equiv \mathbf{ss}^{-1})$ then if $(\mathbf{pmod}) \in \mathcal{S}$ we have $r_s = (\mathbf{s})$, which contradicts Assumption 1. Furthermore, each of the relations r_p, r_m, r_o, r_s must be contained in (\mathbf{pmod}) so, by Lemma 5, each of them contains (\mathbf{d}) . This implies that for every $r \in \mathcal{S}$, we have $(\mathbf{d}) \subseteq r$ whenever $r \cap (\mathbf{pmod})$ is non-empty. This condition precisely means that $\mathcal{S} \subseteq \mathcal{E}_d$. □

Lemma 7 *With the assumptions above, if \mathcal{S} contains a non-trivial relation r such that $r \subseteq (\equiv \mathbf{ss}^{-1})$, then $\mathcal{S} \subseteq \mathcal{S}_d$; if \mathcal{S} contains a non-trivial relation r such that $r \subseteq (\equiv \mathbf{ff}^{-1})$, then $\mathcal{S} \subseteq \mathcal{E}_d$.*

Proof. The two cases are dual so we consider only the first one.

Observe that $r \circ r^{-1} = (\equiv \mathbf{ss}^{-1})$ so we can assume that $r = (\equiv \mathbf{ss}^{-1})$. We have $(\mathbf{d}) \circ (\equiv \mathbf{ss}^{-1}) = (\mathbf{p}^{-1}\mathbf{m}^{-1}\mathbf{o}^{-1}\mathbf{df})$, so the inverse relation $(\mathbf{pmod}^{-1}\mathbf{f}^{-1}) \in \mathcal{S}$ and the result follows from Lemma 6. □

In view of Lemma 7 and Lemma 5, it is sufficient to consider only cases such that for any non-trivial $r \in \mathcal{S}$, $r \cap (\mathbf{dd}^{-1})$ is non-empty.

Lemma 8 *With the assumptions above, if $(\mathbf{dd}^{-1}) \not\subseteq r_o$ then $\mathcal{S} \subseteq \mathcal{S}_d$ or $\mathcal{S} \subseteq \mathcal{E}_d$.*

Proof. Suppose first that $r_o \cap (\mathbf{dd}^{-1}) = (\mathbf{d})$. Then, by Lemma 4, we have $(\mathbf{oo}^{-1}) \not\subseteq r_o$. By the minimality of r_o , $r_o \cap \nu = r_o$, so we have $r_1 = (\mathbf{od}) \subseteq r_o \subseteq (\equiv \mathbf{odss}^{-1}\mathbf{ff}^{-1}) = r_2$. Consequently, $r_1 \circ (\mathbf{d}) \subseteq r_o \circ (\mathbf{d}) \subseteq r_2 \circ (\mathbf{d})$, that is, $(\mathbf{ods}) \subseteq r_o \circ (\mathbf{d}) \subseteq (\mathbf{oo}^{-1}\mathbf{dsf})$. Then, by minimality of r_o , we have $(\mathbf{od}) \subseteq r_o \subseteq (\mathbf{ods})$. Finally,

$$(\mathbf{pmod}) = (\mathbf{d}) \circ (\mathbf{od}) \subseteq (\mathbf{d}) \circ r_o \subseteq (\mathbf{d}) \circ (\mathbf{ods}) = (\mathbf{pmod}),$$

that is, $(\mathbf{pmod}) = (\mathbf{d}) \circ r_o \in \mathcal{S}$, which implies, by Lemma 6, that $\mathcal{S} \subseteq \mathcal{E}_d$.

Dual calculations show that if $r_o \cap (\mathbf{dd}^{-1}) = (\mathbf{d}^{-1})$ then $(\mathbf{pmod}^{-1}\mathbf{f}^{-1}) \in \mathcal{S}$, and hence, by Lemma 6, $\mathcal{S} \subseteq \mathcal{S}_d$. □

Lemma 9 *With the assumptions above, if $r_s \cap (\mathbf{dd}^{-1}) = (\mathbf{d}^{-1})$ or $r_f \cap (\mathbf{dd}^{-1}) = (\mathbf{d}^{-1})$, then $\mathcal{S} \subseteq \mathcal{S}_d$ or $\mathcal{S} \subseteq \mathcal{E}_d$.*

Proof. The two cases are dual so we prove only the first one.

If $r_s \cap (\mathbf{dd}^{-1}) = (\mathbf{d}^{-1})$, then, by Lemma 4, we have $(\mathbf{ss}^{-1}) \not\subseteq r_s$. By the minimality of r_s , $r_s \cap \nu = r_s$, so we have $(\mathbf{d}^{-1}\mathbf{s}) \subseteq r_s \subseteq (\equiv \mathbf{oo}^{-1}\mathbf{d}^{-1}\mathbf{sff}^{-1})$.

Suppose $r_s \cap (\mathbf{oo}^{-1})$ is non-empty. Then $(\mathbf{dd}^{-1}) \not\subseteq r_o$ so $\mathcal{S} \subseteq \mathcal{S}_d$ or $\mathcal{S} \subseteq \mathcal{E}_d$, by Lemma 8. Suppose to the contrary that $r_s \cap (\mathbf{oo}^{-1}) = \emptyset$. Then, $(\mathbf{d}^{-1}\mathbf{s}) \subseteq r_s \subseteq (\equiv \mathbf{d}^{-1}\mathbf{sff}^{-1})$ and

$$(\mathbf{od}^{-1}\mathbf{f}^{-1}) = (\mathbf{d}^{-1}) \circ (\mathbf{d}^{-1}\mathbf{s}) \subseteq (\mathbf{d}^{-1}) \circ r_s \subseteq (\mathbf{d}^{-1}) \circ (\equiv \mathbf{d}^{-1}\mathbf{sff}^{-1}) = (\mathbf{od}^{-1}\mathbf{s}^{-1}\mathbf{f}^{-1}).$$

Therefore we have $(\mathbf{od}^{-1}\mathbf{f}^{-1}) \in \mathcal{S}$ or $(\mathbf{od}^{-1}\mathbf{s}^{-1}\mathbf{f}^{-1}) \in \mathcal{S}$, which implies $(\mathbf{dd}^{-1}) \not\subseteq r_o$, so the result follows from Lemma 8. □

Lemma 10 *With the assumptions above, if $(dd^{-1}) \not\subseteq r_{\mathbf{p}}$ then $\mathcal{S} \subseteq \mathcal{S}_{\mathbf{d}}$ or $\mathcal{S} \subseteq \mathcal{E}_{\mathbf{d}}$.*

Proof. We consider only the case $r_{\mathbf{p}} \cap (dd^{-1}) = (\mathbf{d})$ since the case $r_{\mathbf{p}} \cap (dd^{-1}) = (\mathbf{d}^{-1})$ is dual. By Lemma 4, we have $(pp^{-1}) \not\subseteq r_{\mathbf{p}}$. So, we have

$$(\mathbf{pd}) \subseteq r_{\mathbf{p}} \subseteq (\equiv \text{pmm}^{-1}\text{oo}^{-1}\text{dss}^{-1}\text{ff}^{-1}).$$

If $r_1 = r_{\mathbf{p}} \cap (\text{oo}^{-1}\text{s}^{-1}\text{f}^{-1})$ is non-empty then using Lemmas 8 and 9 we easily get the required result; for example, $r_{\mathbf{p}} = (\mathbf{pod})$ implies $(dd^{-1}) \not\subseteq r_{\mathbf{o}}$ and $r_{\mathbf{p}} = (\mathbf{pds}^{-1})$ implies $r_{\mathbf{s}} \cap (dd^{-1}) = (\mathbf{d}^{-1})$. So we may assume that $(\mathbf{pd}) \subseteq r_{\mathbf{p}} \subseteq (\equiv \text{pmm}^{-1}\text{dsf})$. Then

$$(\mathbf{pd}) = (\mathbf{d}) \circ (\mathbf{pd}) \subseteq (\mathbf{d}) \circ r_{\mathbf{p}} \subseteq (\mathbf{d}) \circ (\equiv \text{pmm}^{-1}\text{dsf}) = (\mathbf{pp}^{-1}\mathbf{d}).$$

Since $(\mathbf{pp}^{-1}\mathbf{d}) \notin \mathcal{S}$ (otherwise its symmetric part (\mathbf{pp}^{-1}) belongs to \mathcal{S} which contradicts Lemma 4), we get $(\mathbf{pd}) = (\mathbf{d}) \circ r_{\mathbf{p}} \in \mathcal{S}$. This implies that $(\mathbf{pmods}) = (\mathbf{pd}) \circ (\mathbf{pd}) \in \mathcal{S}$ and hence $\mathcal{S} \subseteq \mathcal{E}_{\mathbf{d}}$, by Lemma 6. \square

Lemma 11 *With the assumptions above, if $(dd^{-1}) \not\subseteq r_{\mathbf{m}}$ then $\mathcal{S} \subseteq \mathcal{S}_{\mathbf{d}}$ or $\mathcal{S} \subseteq \mathcal{E}_{\mathbf{d}}$.*

Proof. We consider only the case $r_{\mathbf{m}} \cap (dd^{-1}) = (\mathbf{d})$, the case $r_{\mathbf{m}} \cap (dd^{-1}) = (\mathbf{d}^{-1})$ is dual.

As in the proof of the previous lemma, if $r_{\mathbf{m}} \cap (\text{pp}^{-1}\text{oo}^{-1}\text{s}^{-1}\text{f}^{-1})$ is non-empty then we get the required result by Lemmas 8, 9 and 10. So we may assume that $(\mathbf{md}) \subseteq r_{\mathbf{m}} \subseteq (\equiv \text{mdsf})$. Then

$$(\mathbf{pd}) = (\mathbf{d}) \circ (\mathbf{md}) \subseteq (\mathbf{d}) \circ r_{\mathbf{m}} \subseteq (\mathbf{d}) \circ (\equiv \text{mdsf}) = (\mathbf{pd}).$$

Thus, $(\mathbf{pd}) \in \mathcal{S}$. This implies that $(\mathbf{pmods}) = (\mathbf{pd}) \circ (\mathbf{pd}) \in \mathcal{S}$ and hence $\mathcal{S} \subseteq \mathcal{E}_{\mathbf{d}}$, by Lemma 6. \square

Lemma 12 *With the assumptions above, $\mathcal{S} \subseteq \mathcal{S}_{\mathbf{d}}$ or $\mathcal{S} \subseteq \mathcal{E}_{\mathbf{d}}$.*

Proof. If a subalgebra \mathcal{S} satisfies none of the conditions of Lemmas 7-11 then, by Lemma 5, (\mathbf{d}) is contained in all of the minimal relations $r_{\mathbf{p}}$, $r_{\mathbf{m}}$, $r_{\mathbf{o}}$, and $r_{\mathbf{s}}$. Therefore, every $r \in \mathcal{S}$ satisfies $r \cap (\mathbf{pmods}) \neq \emptyset \Rightarrow (\mathbf{d}) \subseteq r$, which precisely means that $\mathcal{S} \subseteq \mathcal{E}_{\mathbf{d}}$. \square

5.2 The case $\text{bas}(\mathcal{S}) = \{\equiv, \mathbf{o}, \mathbf{o}^{-1}\}$

In this subsection, we will show that if \mathcal{S} is a subalgebra with $\text{bas}(\mathcal{S}) = \{\equiv, \mathbf{o}, \mathbf{o}^{-1}\}$, then $\mathcal{S} \subseteq \mathcal{S}_{\mathbf{o}}$, $\mathcal{S} \subseteq \mathcal{E}_{\mathbf{o}}$ or \mathcal{S} is NP-complete.

To obtain this result we shall assume throughout this subsection that \mathcal{S} is a subalgebra of \mathcal{A} satisfying the following assumptions:

Assumption 1 $\text{bas}(\mathcal{S}) = \{\equiv, \mathbf{o}, \mathbf{o}^{-1}\}$.

Assumption 2 \mathcal{S} is not NP-complete.

Using these assumptions we obtain increasingly detailed information about \mathcal{S} in Lemmas 14-20, until we are able to show that in all cases $\mathcal{S} \subseteq \mathcal{S}_{\mathbf{o}}$ or $\mathcal{S} \subseteq \mathcal{E}_{\mathbf{o}}$. (The structure of this proof is quite similar to the proof of the case when $\text{bas}(\mathcal{S}) = \{\equiv, \mathbf{d}, \mathbf{d}^{-1}\}$, above.) These lemmas rely on the following NP-completeness result.

Lemma 13 *The subset $\{(\mathbf{o}), (\mathbf{dd}^{-1})\}$ of \mathcal{A} is NP-complete.*

Proof. Apply Lemma 1 with $R = (\mathbf{d})$, $R_1 = (\mathbf{o})$, $R_2 = (\mathbf{dd}^{-1})$. \square

In the proofs below, we will make frequent use of the fact that $\nu = (\mathbf{o}) \circ (\mathbf{o}^{-1}) \in \mathcal{S}$. Note also that $(\mathbf{pmo}) = (\mathbf{o}) \circ (\mathbf{o}) \in \mathcal{S}$ and the relation (\mathbf{pm}) does not belong to \mathcal{S} since, otherwise, $(\mathbf{p}) = (\mathbf{pm}) \circ (\mathbf{pm}) \in \mathcal{S}$, which contradicts Assumption 1. Therefore $r_{\mathbf{p}} = r_{\mathbf{m}} = (\mathbf{pmo})$ which implies that, for every $r \in \mathcal{S}$, if $r \cap (\mathbf{pmo}) \neq \emptyset$ then $(\mathbf{o}) \subseteq r$.

Lemma 14 *With the assumptions above,*

$$\text{sym}(\mathcal{S}) \subseteq \{r^* | (\text{oo}^{-1}) \subseteq r\} \cup \{r^* | r \subseteq (\equiv \text{ss}^{-1})\} \cup \{r^* | r \subseteq (\equiv \text{ff}^{-1})\}.$$

Proof. We will show that if $\text{sym}(\mathcal{S})$ is not included in the above set then \mathcal{S} is NP-complete, which contradicts Assumption 2.

Suppose first that $(\equiv \text{ss}^{-1}\text{ff}^{-1})$ or $(\text{ss}^{-1}\text{ff}^{-1})$ belongs to $\text{sym}(\mathcal{S})$. Then

$$r = (\text{o}) \circ (\equiv \text{ss}^{-1}\text{ff}^{-1}) = (\text{o}) \circ (\text{ss}^{-1}\text{ff}^{-1}) = (\text{pmodd}^{-1}\text{sf}^{-1}) \in \mathcal{S},$$

so we have $(\text{dd}^{-1}) = r^* \in \mathcal{S}$, which implies that \mathcal{S} is NP-complete, by Lemma 13.

Suppose now that there exists $r^* \in \text{sym}(\mathcal{S})$ such that $r^* \subseteq (\equiv \text{pp}^{-1}\text{mm}^{-1}\text{dd}^{-1}\text{ss}^{-1}\text{ff}^{-1})$ and $r^* \not\subseteq (\equiv \text{ss}^{-1}\text{ff}^{-1})$. If $r_1 = r^* \cap (\text{pp}^{-1}\text{mm}^{-1})$ is non-empty then $r' = r^* \cap (\text{pmo}) \subseteq (\text{pm})$ implying that $(\text{p}) = r' \circ r'$ belongs to \mathcal{S} , which contradicts Assumption 1. Therefore $(\text{dd}^{-1}) \subseteq r^*$ since $r^* \not\subseteq (\equiv \text{ss}^{-1}\text{ff}^{-1})$. Now, it is easy to check that if $r_2 = ((\text{o}) \circ r^*) \cap ((\text{o}^{-1}) \circ r^*)$ then $r_2^* = (\text{dd}^{-1}) \in \mathcal{S}$, which implies that \mathcal{S} is NP-complete, by Lemma 13. \square

Lemma 15 *With the assumptions above,*

$$\mathcal{S} \subseteq \{r | r \cap (\text{oo}^{-1}) \neq \emptyset\} \cup \{r | r \subseteq (\equiv \text{ss}^{-1})\} \cup \{r | r \subseteq (\equiv \text{ff}^{-1})\}.$$

Proof. Assume for contradiction that \mathcal{S} contains a relation r with $r \cap (\text{oo}^{-1})$ empty, $r \not\subseteq (\equiv \text{ss}^{-1})$ and $r \not\subseteq (\equiv \text{ff}^{-1})$.

Among such relations, choose r to be minimal with respect to inclusion. Then, since, as noted above, $r \cap (\text{pmo}) \neq \emptyset$ implies $(\text{o}) \subseteq r$, we have $r \subseteq (\equiv \text{dd}^{-1}\text{ss}^{-1}\text{ff}^{-1})$.

Assume first that $(\text{d}) \subseteq r$ (the argument for (d^{-1}) is dual). Consider $r_1 = (\text{o}) \circ r \subseteq (\text{pmodd}^{-1}\text{sf}^{-1})$. By the minimality of r , $r_1 \cap r = r$ or empty, but $(\text{d}) \subseteq r$, so $(\text{d}) \subseteq r_1$, so $r_1 \cap r$ is not empty and hence $r \subseteq (\text{dd}^{-1}\text{sf}^{-1})$. Now consider $r_2 = (\text{o}^{-1}) \circ r \subseteq (\text{p}^{-1}\text{m}^{-1}\text{o}^{-1}\text{dd}^{-1}\text{s}^{-1}\text{f})$. By a similar argument we get $r \subseteq (\text{dd}^{-1}\text{s}^{-1}\text{f})$. Combining these two results gives $r \subseteq (\text{dd}^{-1})$, and then by Lemma 14 we get $r = (\text{d})$, which contradicts Assumption 1.

Hence, we must have $r \subseteq (\equiv \text{ss}^{-1}\text{ff}^{-1})$. If $(\text{s}^{-1}\text{f}) \subseteq r$ then $((\text{o}) \circ r)^* = (\text{dd}^{-1}) \in \mathcal{S}$, which contradicts Lemma 14. If $r = (\equiv \text{sf})$ or $r = (\text{sf})$ then $r \cap ((\text{o}) \circ r) = (\text{s}) \in \mathcal{S}$, which contradicts Assumption 1. \square

The proofs of the following two lemmas are omitted since they are very similar to the proofs of Lemmas 6 and 7, respectively.

Lemma 16 *With the assumptions above, if $(\text{pmods}) \in \mathcal{S}$ then $\mathcal{S} \subseteq \mathcal{E}_0$; and if $(\text{pmod}^{-1}\text{f}^{-1}) \in \mathcal{S}$ then $\mathcal{S} \subseteq \mathcal{S}_0$.*

Lemma 17 *With the assumptions above, if \mathcal{S} contains a non-trivial relation r such that $r \subseteq (\equiv \text{ss}^{-1})$, then $\mathcal{S} \subseteq \mathcal{S}_0$; if \mathcal{S} contains a non-trivial relation r such that $r \subseteq (\equiv \text{ff}^{-1})$, then $\mathcal{S} \subseteq \mathcal{E}_0$.*

In view of Lemma 17 and Lemma 15, it is sufficient to consider only cases such that for any non-trivial $r \in \mathcal{S}$, $r \cap (\text{oo}^{-1})$ is non-empty.

Lemma 18 *With the assumptions above, if $(\text{oo}^{-1}) \not\subseteq r_{\text{d}}$ then $\mathcal{S} \subseteq \mathcal{S}_0$ or $\mathcal{S} \subseteq \mathcal{E}_0$.*

Proof. Suppose first that $r_{\text{d}} \cap (\text{oo}^{-1}) = (\text{o})$. Then $(\text{dd}^{-1}) \not\subseteq r_{\text{d}}$, since the opposite would contradict Lemma 14. By the minimality of r_{d} , $r_{\text{d}} \cap \nu = r_{\text{d}}$ so $r_1 = (\text{od}) \subseteq r_{\text{d}} \subseteq (\equiv \text{odss}^{-1}\text{ff}^{-1}) = r_2$. Consequently, $(\text{o}) \circ r_1 \subseteq (\text{o}) \circ r_{\text{d}} \subseteq (\text{o}) \circ r_2$, that is, $(\text{pmods}) \subseteq (\text{o}) \circ r_{\text{d}} \subseteq (\text{pmodd}^{-1}\text{sf}^{-1})$. Then, by minimality of r_{d} , we have $(\text{od}) \subseteq r_{\text{d}} \subseteq (\text{odsf}^{-1})$. Finally,

$$(\text{pmods}) = (\text{o}) \circ (\text{od}) \subseteq (\text{o}) \circ r_{\text{d}} \subseteq (\text{o}) \circ (\text{odsf}^{-1}) = (\text{pmods}),$$

that is, $(\text{pmods}) = (\text{o}) \circ r_{\text{d}} \in \mathcal{S}$, which implies, by Lemma 16, that $\mathcal{S} \subseteq \mathcal{E}_0$.

Dual calculations show that if $r_{\text{d}} \cap (\text{oo}^{-1}) = (\text{o}^{-1})$ then $(\text{pmod}^{-1}\text{f}^{-1}) \in \mathcal{S}$, and hence $\mathcal{S} \subseteq \mathcal{S}_0$, by Lemma 16. \square

Lemma 19 *If $r_{\mathcal{S}} \cap (\text{oo}^{-1}) = (\text{o}^{-1})$ then $\mathcal{S} \subseteq \mathcal{S}_{\mathcal{O}}$ or $\mathcal{S} \subseteq \mathcal{E}_{\mathcal{O}}$.*

Proof. We have $(\text{ss}^{-1}) \not\subseteq r_{\mathcal{D}}$, since the opposite would contradict Lemma 14. By the minimality of $r_{\mathcal{D}}$, $r_{\mathcal{D}} \cap \nu = r_{\mathcal{D}}$ so $(\text{o}^{-1}\text{s}) \subseteq r_{\mathcal{S}} \subseteq (\equiv \text{o}^{-1}\text{dd}^{-1}\text{sff}^{-1})$.

Suppose first that $r_{\mathcal{S}} \cap (\text{dd}^{-1})$ is non-empty. Then, $(\text{oo}^{-1}) \not\subseteq r_{\mathcal{D}}$ and $\mathcal{S} \subseteq \mathcal{S}_{\mathcal{O}}$ or $\mathcal{S} \subseteq \mathcal{E}_{\mathcal{O}}$ by Lemma 18.

Suppose to the contrary that $r_{\mathcal{S}} \cap (\text{dd}^{-1}) = \emptyset$. Then $(\text{o}^{-1}\text{s}) \subseteq r_{\mathcal{S}} \subseteq (\equiv \text{o}^{-1}\text{sff}^{-1})$ and

$$(\text{o}^{-1}) \circ (\text{o}^{-1}\text{s}) \subseteq (\text{o}^{-1}) \circ r_{\mathcal{S}} \subseteq (\text{o}^{-1}) \circ (\equiv \text{o}^{-1}\text{sff}^{-1}),$$

that is,

$$(\text{p}^{-1}\text{m}^{-1}\text{o}^{-1}\text{df}) \subseteq (\text{o}^{-1}) \circ r_{\mathcal{S}} \subseteq (\text{p}^{-1}\text{m}^{-1}\text{o}^{-1}\text{dd}^{-1}\text{s}^{-1}\text{f}).$$

Hence $(\text{oo}^{-1}) \not\subseteq r_{\mathcal{D}}$, so the result follows by Lemma 18. \square

Lemma 20 *With the assumptions above, $\mathcal{S} \subseteq \mathcal{S}_{\mathcal{O}}$ or $\mathcal{S} \subseteq \mathcal{E}_{\mathcal{O}}$.*

Proof. If \mathcal{S} satisfies none of the conditions of Lemmas 17-19, then, by Lemma 15, (o) is contained in both $r_{\mathcal{D}}$ and $r_{\mathcal{S}}$. Since, as we noted in the beginning of this subsection, (o) is also contained in both $r_{\mathcal{P}}$ and $r_{\mathcal{M}}$, we conclude that \mathcal{S} is contained in $\mathcal{E}_{\mathcal{O}}$. \square

5.3 The case $\text{bas}(\mathcal{S}) = \{\equiv, \text{s}, \text{s}^{-1}\}$ or $\text{bas}(\mathcal{S}) = \{\equiv, \text{f}, \text{f}^{-1}\}$

In this subsection, we will show that if $\text{bas}(\mathcal{S}) = \{\equiv, \text{s}, \text{s}^{-1}\}$, then either \mathcal{S} is NP-complete or \mathcal{S} is contained in one of the subalgebras \mathcal{H} , $\mathcal{S}_{\mathcal{D}}$, $\mathcal{S}_{\mathcal{O}}$, $\mathcal{S}_{\mathcal{P}}$, \mathcal{E}^* , or in one of \mathcal{A}_i , $1 \leq i \leq 4$. By using the obvious symmetry between the relations (s) and (f) , it immediately follows that if $\text{bas}(\mathcal{S}) = \{\equiv, \text{f}, \text{f}^{-1}\}$, then either \mathcal{S} is NP-complete or contained in one of \mathcal{H} , $\mathcal{E}_{\mathcal{D}}$, $\mathcal{E}_{\mathcal{O}}$, $\mathcal{E}_{\mathcal{P}}$, \mathcal{S}^* or \mathcal{B}_i , $1 \leq i \leq 4$.

To obtain this result we shall assume throughout this subsection that \mathcal{S} is a subalgebra of \mathcal{A} satisfying the following assumptions:

Assumption 1 $\text{bas}(\mathcal{S}) = \{\equiv, \text{s}, \text{s}^{-1}\}$.

Assumption 2 \mathcal{S} is not NP-complete.

Using these assumptions we obtain increasingly detailed information about \mathcal{S} in Lemmas 22-29, until we are able to obtain the result. These lemmas rely on the following NP-completeness result.

Lemma 21 *The subset $\{r\}$ of \mathcal{A} is NP-complete whenever $(\text{ods}^{-1}) \subseteq r \subseteq (\text{pmods}^{-1}\text{f}^{-1})$.*

Proof. Let r_3 be the union of all basic relations except for \equiv and s^{-1} , and consider the instance $\Gamma_4 = \{xra, xrb, yrb, ary, bra\}$ over the variables x, y, a, b . In the cases when $r = (\text{ods}^{-1})$ or $r = (\text{pmods}^{-1}\text{f}^{-1})$, it can be shown that $\Gamma_4 \cup \{xr'y\}$ is satisfiable for every basic relation $r' \subseteq r_3$ but not satisfiable for any other choice of r' . It follows that, for every r such that $(\text{ods}^{-1}) \subseteq r \subseteq (\text{pmods}^{-1}\text{f}^{-1})$, we can derive r_3 from r . Further, we can derive the relation $r_4 = r \cap r_3$ satisfying $(\text{od}) \subseteq r_4 \subseteq (\text{pmodf}^{-1})$, and the relation $r_5 = r_4 \circ r_4$. It is easy to check that

$$(\text{pmods}) \subseteq r_5 \subseteq (\text{pmodsf}^{-1})$$

which implies that $r_5 \cap r^{-1} = (\text{s})$. Furthermore, $(\text{s}) \circ (\text{s}^{-1}) = (\equiv \text{ss}^{-1})$ and $r \circ r$ is the disequality relation, so the relation (ss^{-1}) can be obtained from the relation r . If $R = (\text{s})$, $R_1 = r^{-1}$ and $R_2 = r$, then these relations satisfy the conditions of Lemma 1 so $\{(\text{ss}^{-1}), r^{-1}, r\}$ is NP-complete. Since all of these relations can be derived from the single relation r , it follows that $\{r\}$ is NP-complete. \square

Lemma 22 *With the assumptions above, if \mathcal{S} contains the relation (od) , then every $r \in \mathcal{S}$ satisfies condition 3) of \mathcal{H} .*

Proof. Arbitrarily choose $r \in \mathcal{S}$. Since $(\text{od}) \in \mathcal{S}$, it follows that $r_{\text{d}} = r_{\text{o}} = (\text{od})$ and $(\text{o})^{\pm 1} \subseteq r \Leftrightarrow (\text{d})^{\pm 1} \subseteq r$. Furthermore, $r_1 = (\text{s}) \circ (\text{od}) = (\text{pmod}) \in \mathcal{S}$ so $r_{\text{p}} \subseteq r_1$ and $r_{\text{m}} \subseteq r_1$. Since $(\text{pm}) \circ (\text{pm}) = (\text{p})$, we have $r_{\text{p}} \not\subseteq (\text{pm})$ and $r_{\text{m}} \not\subseteq (\text{pm})$, and therefore $r_{\text{p}} \cap (\text{od}) \neq \emptyset$ and $r_{\text{m}} \cap (\text{od}) \neq \emptyset$. Now it follows that $(\text{od}) \subseteq r_{\text{p}}$ and $(\text{od}) \subseteq r_{\text{m}}$. Thus, if $r \cap (\text{pm}) \neq \emptyset$, then $(\text{od}) \subseteq r$ which means that r satisfies condition 3) of \mathcal{H} . \square

Lemma 23 *With the assumptions above, if \mathcal{S} contains a non-trivial relation r' with $r' \subseteq (\equiv \text{pp}^{-1}\text{mm}^{-1}\text{ff}^{-1})$ then \mathcal{S} is included in one of \mathcal{H} , \mathcal{E}^* , or \mathcal{S}_{p} .*

Proof. Case 1. $r' = (\text{ff}^{-1})$

Since $(\text{ff}^{-1}) \circ (\text{s}) = (\text{od}) \in \mathcal{S}$, it follows that any $r \in \mathcal{S}$ satisfies condition 3) of \mathcal{H} by Lemma 22. Note also that now, for every $r \in \mathcal{S}$, we have

$$(\text{o})^{\pm 1} \subseteq r \Leftrightarrow (\text{d})^{\pm 1} \subseteq r \text{ and } (\text{f}) \subseteq r \Leftrightarrow (\text{f}^{-1}) \subseteq r. \quad (1)$$

Suppose that $\mathcal{S} \not\subseteq \mathcal{H}$, i.e. some $r \in \mathcal{S}$ fails to satisfy condition 1) or condition 2) of \mathcal{H} . Then, using the conditions (1) from the previous paragraph, it is not hard to check that the relation r can be chosen so that

$$(\text{s}^{-1}\text{ff}^{-1}) \subseteq r \subseteq (\equiv \text{pmodss}^{-1}\text{ff}^{-1}) \text{ or } (\text{ods}^{-1}) \subseteq r \subseteq (\equiv \text{pmodss}^{-1}\text{ff}^{-1}).$$

In both cases, multiplying the relations by (s) from the left we get

$$(\equiv \text{pmodss}^{-1}) \subseteq (\text{s}) \circ r \subseteq (\equiv \text{pmodss}^{-1}).$$

Therefore $(\equiv \text{pmodss}^{-1}) \in \mathcal{S}$, and

$$(\text{pmodss}^{-1}) = (\equiv \text{pmodss}^{-1}) \cap ((\equiv \text{pmodss}^{-1})^{-1} \circ (\text{od})^{-1}) \in \mathcal{S},$$

so \mathcal{S} is NP-complete by Lemma 21, which contradicts Assumption 2.

Case 2. $r' \subseteq (\equiv \text{ff}^{-1})$.

Multiplying r' and its inverse we get $(\equiv \text{ff}^{-1})$, so we may assume that $r' = (\equiv \text{ff}^{-1})$. If some relation $r_2 \in \mathcal{S}$ fails to satisfy condition 2) of \mathcal{E}^* then $r_2 \cap r'$ is either $(\text{f})^{\pm 1}$, which is impossible, or (ff^{-1}) going back to Case 1. Suppose now that each $r \in \mathcal{S}$ satisfies condition 2) of \mathcal{E}^* .

We have $r' \circ (\text{s}) = (\text{ods}) \in \mathcal{S}$. If $(\text{od}) \in \mathcal{S}$ then $r' \cap ((\text{s}^{-1}) \circ (\text{od})) = (\text{ff}^{-1}) \in \mathcal{S}$, which implies $\mathcal{S} \subseteq \mathcal{H}$ by Case 1. Suppose $(\text{od}) \notin \mathcal{S}$. Then we have $r \cap (\text{od})^{\pm 1} \neq \emptyset \Rightarrow (\text{s})^{\pm 1} \subseteq r$ for every $r \in \mathcal{S}$. Assume a relation $r_3 \in \mathcal{S}$ does not satisfy condition 1) of \mathcal{E}^* , that is, $r_3 \cap (\text{pmod}) \neq \emptyset$ and $(\text{s}) \not\subseteq r_3$. Since $(\text{pmodss}) = (\text{s}) \circ r' \in \mathcal{S}$, we have $r_4 = r_3 \cap (\text{pmodss}) \in \mathcal{S}$ and $r_4 \subseteq (\text{pm})$. This implies $r_4 \circ r_4 = (\text{p})$, which contradicts Assumption 1. Therefore the relations in \mathcal{S} must satisfy both conditions of \mathcal{E}^* , that is, $\mathcal{S} \subseteq \mathcal{E}^*$.

Case 3. $r' \subseteq (\equiv \text{mm}^{-1}\text{ff}^{-1})$ and $r' \cap (\text{mm}^{-1}) \neq \emptyset$.

Without loss of generality we may assume that $(\text{m}) \subseteq r$. Then we have $(\text{m}) \subseteq r_1 = (r' \cap (r' \circ (\text{s}^{-1}))) \subseteq (\text{mm}^{-1})$ so $(\text{m}) = r_1 \cap (r_1 \circ (\text{s}^{-1})) \in \mathcal{S}$, which contradicts Assumption 1.

Case 4. $r' \cap (\text{pp}^{-1}) \neq \emptyset$.

Assume without loss of generality that $(\text{p}) \subseteq r'$. Then $(\text{p}) \subseteq (\text{s}) \circ r' \subseteq (\text{pp}^{-1}\text{mm}^{-1}\text{ods})$. Further, $(\text{p}) \subseteq r_5 = r' \cap ((\text{s}) \circ r') \subseteq (\text{pp}^{-1}\text{mm}^{-1})$ and $(\text{p}) \subseteq r_6 = r_5 \circ (\text{s}^{-1}) \subseteq (\text{pp}^{-1}\text{m})$. If $(\text{p}^{-1}) \not\subseteq r_6$ then $r_6 \circ r_6 = (\text{p}) \in \mathcal{S}$, which contradicts Assumption 1. Otherwise $(\text{pp}^{-1}) = r_6^* \in \mathcal{S}$. Then $(\text{p}) \subseteq r \Leftrightarrow (\text{p}^{-1}) \subseteq r$ holds for every $r \in \mathcal{S}$.

We have $(\text{s}^{-1}) \circ (\text{pp}^{-1}) = (\text{pp}^{-1}\text{mod}^{-1}\text{f}^{-1}) \in \mathcal{S}$. For every non-empty $r_7 \subseteq (\text{mod}^{-1}\text{f}^{-1})$, we have $((\text{s}) \circ r_7) \cap (\text{pp}^{-1}) = (\text{p})$. Therefore no such r_7 belongs to \mathcal{S} . We conclude that, for any $r \in \mathcal{S}$, if $r \cap (\text{mod}^{-1}\text{f}^{-1}) \neq \emptyset$ then $(\text{p}) \subseteq r$, which means that $\mathcal{S} \subseteq \mathcal{S}_{\text{p}}$. \square

In view of Lemma 23, it is now sufficient to consider cases where the following additional property holds:

Assumption 3 For every non-trivial $r \in \mathcal{S}$, we have $r \cap (\text{oo}^{-1}\text{dd}^{-1}\text{ss}^{-1}) \neq \emptyset$.

Lemma 24 *With the assumptions above, if $r_{\mathbf{d}} \cap (\text{oo}^{-1}\text{dd}^{-1}\text{ss}^{-1}) = (\mathbf{d})$ or $(\text{dd}^{-1}) \subseteq r_{\mathbf{d}}$ then $\mathcal{S} \subseteq \mathcal{S}_{\mathbf{d}}$.*

Proof. Case 1. $r_{\mathbf{d}} \cap (\text{oo}^{-1}\text{dd}^{-1}\text{ss}^{-1}) = (\mathbf{d})$.

By assumption, the relation $r_{\mathbf{d}}$ satisfies condition $(\mathbf{d}) \subseteq r_{\mathbf{d}} \subseteq (\equiv \text{pp}^{-1}\text{mm}^{-1}\text{dff}^{-1})$. Let r_1 be calculated as $r_{\mathbf{d}} \cap ((\mathbf{s}) \circ r_{\mathbf{d}})$. Then we have $(\mathbf{d}) \subseteq r_1 \subseteq (\text{pp}^{-1}\text{mm}^{-1}\mathbf{d})$. By minimality of $r_{\mathbf{d}}$ we get $(\mathbf{d}) \subseteq r_{\mathbf{d}} \subseteq (\text{pp}^{-1}\text{mm}^{-1}\mathbf{d})$. Calculating r_1 again, we get $(\mathbf{d}) \subseteq r_{\mathbf{d}} \subseteq (\text{pp}^{-1}\mathbf{m}^{-1}\mathbf{d})$. By Assumption 3, we have $(\text{pp}^{-1}) \notin \mathcal{S}$. Furthermore, since we have $(\mathbf{m}^{-1}\mathbf{d}) \subseteq r_{\mathbf{d}} \circ (\mathbf{s})$ only if $(\mathbf{p}^{-1}) \subseteq r_{\mathbf{d}}$, we may assume that $r_{\mathbf{d}}$ is either (pd) , or $(\mathbf{p}^{-1}\mathbf{d})$, or $(\mathbf{p}^{-1}\mathbf{m}^{-1}\mathbf{d})$. In the first case we get $(\text{pd}) \cap ((\mathbf{p}^{-1}\mathbf{d}^{-1}) \circ (\mathbf{s})) = (\mathbf{d})$, which contradicts Assumption 1. Let $(\mathbf{p}^{-1}\mathbf{d}) \subseteq r_{\mathbf{d}} \subseteq (\mathbf{p}^{-1}\mathbf{m}^{-1}\mathbf{d})$. Then $(\mathbf{p}^{-1}\mathbf{m}^{-1}\mathbf{o}^{-1}\mathbf{df}) = r_{\mathbf{d}} \circ (\mathbf{s}) \in \mathcal{S}$. Suppose \mathcal{S} contains a non-empty subrelation r_2 of $(\mathbf{p}^{-1}\mathbf{m}^{-1}\mathbf{o}^{-1}\mathbf{f})$. Then $r_3 = r_{\mathbf{d}} \cap (r_2 \circ (\mathbf{s}^{-1}))$ is a non-empty subrelation of $(\mathbf{p}^{-1}\mathbf{m}^{-1})$ implying that $(\mathbf{p}^{-1}) = r_3 \circ r_3 \in \mathcal{S}$, which contradicts Assumption 1. Therefore, for every $r \in \mathcal{S}$, we have $r \cap (\mathbf{p}^{-1}\mathbf{m}^{-1}\mathbf{o}^{-1}\mathbf{f}) \neq \emptyset \Rightarrow (\mathbf{d}) \subseteq r$, which means that $\mathcal{S} \subseteq \mathcal{S}_{\mathbf{d}}$.

Case 2. $(\text{dd}^{-1}) \subseteq r_{\mathbf{d}}$.

Suppose a relation $r_4 \in \mathcal{S}$ satisfies $(\text{dd}^{-1}) \not\subseteq r_4$. Then it is not hard to verify that if $r_4 \not\subseteq (\equiv \text{ss}^{-1})$ then either $((\mathbf{s}^{-1}) \circ r_4) \cap r_{\mathbf{d}}$ or $(r_4 \circ (\mathbf{s})) \cap r_{\mathbf{d}}$ contains exactly one of (\mathbf{d}) and (\mathbf{d}^{-1}) which contradicts the minimality of $r_{\mathbf{d}}$. Thus, for every $r \in \mathcal{S}$ such that $r \not\subseteq (\equiv \text{ss}^{-1})$, we have $(\text{dd}^{-1}) \subseteq r$. This implies that $\mathcal{S} \subseteq \mathcal{S}_{\mathbf{d}}$. \square

Lemma 25 *With the assumptions above, if $r_{\mathbf{o}} \cap (\text{oo}^{-1}\text{dd}^{-1}\text{ss}^{-1}) = (\mathbf{o})$ or $(\text{oo}^{-1}) \subseteq r_{\mathbf{o}}$ then $\mathcal{S} \subseteq \mathcal{S}_{\mathbf{o}}$.*

Proof. Similar to the previous lemma. \square

Lemma 26 *With the assumptions above, if $r_{\mathbf{d}} = r_{\mathbf{o}^{-1}}$ and $r_{\mathbf{d}} \cap (\text{oo}^{-1}\text{dd}^{-1}\text{ss}^{-1}) = (\mathbf{o}^{-1}\mathbf{d})$, then $\mathcal{S} \subseteq \mathcal{S}_{\mathbf{d}}$.*

Proof. As in the proof of Lemma 24, we can obtain $(\mathbf{o}^{-1}\mathbf{d}) \subseteq r_{\mathbf{d}} \subseteq (\text{pp}^{-1}\mathbf{m}^{-1}\mathbf{o}^{-1}\mathbf{df})$. If $(\text{po}^{-1}\mathbf{d}) \subseteq r_{\mathbf{d}}$ then $(r_{\mathbf{d}} \circ (\mathbf{s}^{-1}))^* = (\text{pp}^{-1}) \in \mathcal{S}$, and the result follows from Lemma 23. Therefore we have $(\mathbf{o}^{-1}\mathbf{d}) \subseteq r_{\mathbf{d}} \subseteq (\mathbf{p}^{-1}\mathbf{m}^{-1}\mathbf{o}^{-1}\mathbf{df})$, and $(\mathbf{p}^{-1}\mathbf{m}^{-1}\mathbf{o}^{-1}\mathbf{df}) = r_{\mathbf{d}} \circ (\mathbf{s}^{-1}) \in \mathcal{S}$. By Assumption 3, no non-empty subrelation r_1 of $(\mathbf{p}^{-1}\mathbf{m}^{-1}\mathbf{f})$ belongs to \mathcal{S} , so, for every $r \in \mathcal{S}$, we have $r \cap (\mathbf{p}^{-1}\mathbf{m}^{-1}\mathbf{o}^{-1}\mathbf{f}) \neq \emptyset \Rightarrow (\mathbf{d}) \subseteq r$, which means that $\mathcal{S} \subseteq \mathcal{S}_{\mathbf{d}}$. \square

Lemma 27 *With the assumptions above, if $r_{\mathbf{d}} = r_{\mathbf{o}}$ and $r_{\mathbf{d}} \cap (\text{oo}^{-1}\text{dd}^{-1}\text{ss}^{-1}) = (\text{od})$, then $\mathcal{S} \subseteq \mathcal{H}$.*

Proof. As in the previous lemmas, it can be shown that $(\text{od}) \subseteq r_{\mathbf{d}} \subseteq (\text{pp}^{-1}\text{mm}^{-1}\text{od})$. Then $r_{\mathbf{d}}^* = \emptyset$ and $(\text{pmod}) = (\mathbf{s}) \circ r_{\mathbf{d}} \in \mathcal{S}$. Further, $(\text{oo}^{-1}\text{dd}^{-1}\text{ff}^{-1}) = ((\mathbf{s}^{-1}) \circ (\text{pmod}))^* \in \mathcal{S}$ and $(\text{od}) = (\text{pmod}) \cap (\text{oo}^{-1}\text{dd}^{-1}\text{ff}^{-1}) \in \mathcal{S}$. By Lemma 22, we know that every $r \in \mathcal{S}$ satisfies condition 3) of \mathcal{H} . Suppose some $r_1 \in \mathcal{S}$ does not satisfy condition 1) of \mathcal{H} . Then r_1 can be chosen so that

$$(\mathbf{s}^{-1}\mathbf{f}^{-1}) \subseteq r_1 \subseteq (\equiv \text{pmodss}^{-1}\text{ff}^{-1}) \text{ or } (\text{os}^{-1}) \subseteq r_1 \subseteq (\equiv \text{pmodss}^{-1}\text{ff}^{-1}).$$

If $(\mathbf{f}) \subseteq r_1$ then $(\text{ff}^{-1}) = (r_1 \cap (\text{oo}^{-1}\text{dd}^{-1}\text{ff}^{-1}))^* \in \mathcal{S}$, which contradicts Assumption 3. Further, if $(\text{od}) \not\subseteq r_1$ then $r_1 \cap (\text{oo}^{-1}\text{dd}^{-1}\text{ff}^{-1}) = (\mathbf{f}^{-1}) \in \mathcal{S}$, which contradicts Assumption 1. Therefore we may assume that $(\text{ods}^{-1}\mathbf{f}^{-1}) \subseteq r_1 \subseteq (\equiv \text{pmodss}^{-1}\mathbf{f}^{-1})$. Now it can be checked that $r_2 = r_1 \cap (r_1^{-1} \circ (\mathbf{o}^{-1}\mathbf{d}^{-1}))$ satisfies $(\text{ods}^{-1}\mathbf{f}^{-1}) \subseteq r_2 \subseteq (\text{pmodss}^{-1}\mathbf{f}^{-1})$, so $\{r_2\}$ is NP-complete by Lemma 21, which contradicts Assumption 2.

One can proceed similarly if condition 2) of \mathcal{H} fails in \mathcal{S} . \square

Lemma 28 *With the assumptions above, if $r \cap (ss^{-1}) \neq \emptyset$ for each non-trivial $r \in \mathcal{S}$ then $\mathcal{S} \subseteq \mathcal{A}_i$ for some $1 \leq i \leq 4$.*

Proof. Case 1. $r_p \cap (ss^{-1}) = (s^{-1})$.

We have $(ps^{-1}) \subseteq r_p$ and $(s) \not\subseteq r_p$. Let $r_1 = (s^{-1}) \circ r_p \in \mathcal{S}$. Then it is easy to check that $(p\text{mod}^{-1}s^{-1}f^{-1}) \subseteq r_1$, and that $r_1 \cap (\equiv s) = \emptyset$. It follows that $r_1 = (p\text{mod}^{-1}s^{-1}f^{-1})$, since otherwise r_1^* is non-empty and $r_1^* \cap (ss^{-1}) = \emptyset$. No non-empty subrelation of $(p\text{mod}^{-1}f^{-1})$ can belong to \mathcal{S} . Therefore, for any $r \in \mathcal{S}$, we have $r \cap (p\text{mod}^{-1}f^{-1}) \neq \emptyset \Rightarrow (s^{-1}) \subseteq r$. Hence $\mathcal{S} \subseteq \mathcal{A}_1$.

Case 2. $r_d \cap (ss^{-1}) = (s^{-1})$.

The proof is similar to Case 1; the only change is that $r_1 = r_d \circ (s^{-1})$, and we deduce that $r_1 = (p^{-1}m^{-1}o^{-1}ds^{-1}f)$, and, hence, $\mathcal{S} \subseteq \mathcal{A}_2$.

Case 3. $r_o \cap (ss^{-1}) = (s^{-1})$.

In view of Cases 1 and 2 we may assume that $(os^{-1}) \subseteq r_o \subseteq (\equiv p^{-1}mm^{-1}od^{-1}s^{-1}ff^{-1})$. Let $r_1 = (s^{-1}) \circ r_o \in \mathcal{S}$. It is easy to check that r_1 satisfies $(od^{-1}s^{-1}f^{-1}) \subseteq r_1 \subseteq (p^{-1}m^{-1}oo^{-1}d^{-1}s^{-1}f^{-1})$. Since $r_1^* \neq (oo^{-1})$, we obtain $r_1 \subseteq (p^{-1}m^{-1}od^{-1}s^{-1}f^{-1})$. It can straightforwardly be verified that if $r_1 \neq (od^{-1}s^{-1}f^{-1})$, then $r_2 = (r_1 \circ r_1)^*$ contains (pp^{-1}) and $r_2 \cap (ss^{-1}) = \emptyset$, which contradicts the assumptions made. Therefore $r_1 = (od^{-1}s^{-1}f^{-1})$, and $(p\text{mod}^{-1}s^{-1}f^{-1}) = r_1 \circ r_1 \in \mathcal{S}$. Therefore, for any $r \in \mathcal{S}$, $r \cap (p\text{mod}^{-1}f^{-1}) \neq \emptyset$ implies $(s^{-1}) \subseteq r$, that is, $\mathcal{S} \subseteq \mathcal{A}_1$.

Case 4. $r_m \cap (ss^{-1}) = (s^{-1})$.

Similarly to Case 3, we infer that $\mathcal{S} \subseteq \mathcal{A}_1$.

Case 5. (s) is contained in each of r_p, r_d, r_o , and r_m .

We have $r_f \cap (ss^{-1}) \neq \emptyset$. Then it follows that if $(s) \subseteq r_f$ then $\mathcal{S} \subseteq \mathcal{A}_3$. Otherwise, $(s^{-1}) \subseteq r_f$ and we have $\mathcal{S} \subseteq \mathcal{A}_4$. \square

Lemma 29 *With the assumptions above, \mathcal{S} is contained in one of the subalgebras $\mathcal{H}, \mathcal{S}_d, \mathcal{S}_o, \mathcal{S}_p, \mathcal{E}^*$, or in one of $\mathcal{A}_i, 1 \leq i \leq 4$.*

Proof. By Lemma 28 it suffices to consider the case when \mathcal{S} contains a non-trivial relation r with $r \cap (ss^{-1}) = \emptyset$.

By Lemma 23 we can assume that, for every non-trivial $r \in \mathcal{S}$, we have $r \cap (oo^{-1}dd^{-1}ss^{-1}) \neq \emptyset$, so it suffices to consider cases when $r_d \cap (ss^{-1}) = \emptyset$ or $r_o \cap (ss^{-1}) = \emptyset$.

We claim that the result now follows from Lemmas 24-27. To establish this claim suppose that $r_d \cap (ss^{-1}) = \emptyset$, but r_d does not satisfy the conditions of Lemma 24. Then $(d^{-1}) \not\subseteq r_d$ and $r_d \cap (oo^{-1}) \neq \emptyset$. If $(oo^{-1}) \subseteq r_d$ then $r_d^* \cap (oo^{-1}dd^{-1}) = (oo^{-1})$, which implies that r_o satisfies the conditions of Lemma 25 and $\mathcal{S} \subseteq \mathcal{S}_o$. Otherwise we have $r_o \subseteq r_d$ or $r_{o^{-1}} \subseteq r_d$. If both of these inclusions are proper then it is easy to see that $r_o \cap (oo^{-1}dd^{-1}) = (o)$ and, hence, $\mathcal{S} \subseteq \mathcal{S}_o$ by Lemma 25. Thus we only need to consider two cases: $r_d = r_{o^{-1}}$ and $r_d = r_o$ which are dealt with in Lemmas 26 and 27. The case when $r_o \cap (ss^{-1}) \neq \emptyset$ but r_o does not satisfy the conditions of Lemma 25 is similar and it is sufficient to consider the same two cases. \square

6 Subalgebras With Only Trivial Basic Relation

In this section, we consider subalgebras \mathcal{S} of \mathcal{A} such that $\text{bas}(\mathcal{S}) = \{(\equiv)\}$. We can assume that \mathcal{S} contains a relation r' such that $(\equiv) \not\subseteq r'$; otherwise $\mathcal{S} \subseteq \mathcal{A}_{\equiv}$.

A relation r is *symmetric* if $r^* = r$ and it is *asymmetric* if $r^* = \emptyset$. If we choose the relation r' to be minimal, then this implies that r' is either asymmetric or symmetric. We consider these two cases in Subsections 6.1 and 6.2, respectively.

6.1 Asymmetric relations

In this subsection we prove the following proposition.

Proposition 4 *Let \mathcal{S} be a subalgebra of \mathcal{A} such that $\text{bas}(\mathcal{S}) = \{\equiv\}$, which contains an asymmetric relation. Then \mathcal{S} is tractable if it is contained in one of the 18 algebras listed in Table 3. Otherwise \mathcal{S} is NP-complete*

To obtain this result we shall assume throughout this subsection that \mathcal{S} is a subalgebra of \mathcal{A} satisfying the following assumptions:

Assumption 1 $\text{bas}(\mathcal{S}) = \{\equiv\}$.

Assumption 2 \mathcal{S} is not NP-complete.

Assumption 3 $r' \in \mathcal{S}$ is an asymmetric relation.

We first show that r' must have a very restricted form, and then show that the result holds for all possible cases in Lemmas 32-37.

A relation $r \in \mathcal{A}$ is said to be *acyclic* if, for every $k > 1$, the instance

$$x_1 r x_2, x_2 r x_3, \dots, x_{k-1} r x_k, x_k r x_1$$

has no model. The acyclic relations are characterised in [14].

Lemma 30 ([14]) *A relation $r \in \mathcal{A}$ is acyclic if and only if r or r^{-1} is a subset of one of the relations $(\text{pmod}^{-1}\text{sf}^{-1})$, $(\text{pmod}^{-1}\text{s}^{-1}\text{f}^{-1})$, (pmodsf) or (pmodsf^{-1}) .*

Proposition 5 *If r is asymmetric, but not acyclic, then $\{r\}$ is NP-complete.*

The proof of this proposition can be found in Subsection A.2. By using this result, we can now restrict our attention to cases where r' is an acyclic relation. To complete the proof of Proposition 4, we will now consider all acyclic relations. The proofs rely on the following NP-completeness results.

Lemma 31 *The following sets of relations are NP-complete:*

- 1) $\{(\text{oo}^{-1}), r\}$ where $(\text{d}) \subseteq r \subseteq (\text{dsf})$;
- 2) $\{(\text{dd}^{-1}), r\}$, where $(\text{o}) \subseteq r \subseteq (\text{pmosf}^{-1})$.

Proof. Set $R = (\text{o})$, $R_1 = r$, $R_2 = (\text{oo}^{-1})$ in the first case, $R = (\text{d})$, $R_1 = r$ and $R_2 = (\text{dd}^{-1})$ in the second case, and apply Lemma 1. \square

Lemma 32 *With the assumptions above, if $(\text{pmods}) \subseteq r'$ or $(\text{pmod}^{-1}\text{f}^{-1}) \subseteq r'$ then \mathcal{S} is contained in one of the 18 maximal tractable subalgebras.*

Proof. We will consider only the first case, the second one is dual.

By Lemma 30, there are only three possible choices for r' : (pmods) , (pmodsf) and (pmodsf^{-1}) . The relations r_{p} , r_{m} , r_{o} , r_{d} , and r_{s} must all be contained in r' . We now consider how they are related to each other.

Assume first that one of these five sets is contained in the other four. If r_{m} is contained in all of r_{p} , r_{o} , r_{d} , r_{s} , then either r_{m} is one of (mf^{-1}) and (mf) , and in this case $\mathcal{S} \subseteq \mathcal{B}_1$ or $\mathcal{S} \subseteq \mathcal{B}_2$, respectively, or else r_{m} coincides with one of r_{p} , r_{o} , r_{d} , r_{s} because $(\text{m}) \notin \mathcal{S}$, by Assumption 1. Further, if r_{p} , r_{o} , or r_{d} is contained in the other four relations, then \mathcal{S} is contained in \mathcal{E}_{p} , \mathcal{E}_{o} , or \mathcal{E}_{d} , respectively. If r_{s} is contained in all of r_{p} , r_{m} , r_{o} , r_{d} , then $\mathcal{S} \subseteq \mathcal{B}_1$ if $(\text{f}^{-1}) \subseteq r_{\text{s}}$, $\mathcal{S} \subseteq \mathcal{B}_2$ if $(\text{f}) \subseteq r_{\text{s}}$, and $\mathcal{S} \subseteq \mathcal{E}^*$ if $(\equiv \text{f}) \subseteq r_{\text{f}}$. In the remaining case, we have $r_{\text{s}} = r_{\text{m}} = (\text{ms})$ is contained in r_{p} , r_{o} and r_{d} , so if r_{f} contains (s) then $\mathcal{S} \subseteq \mathcal{A}_3$ and if r_{f} contains (s^{-1}) then $\mathcal{S} \subseteq \mathcal{A}_4$, otherwise $r_{\text{f}} \subseteq (\text{ff}^{-1})$, and we have $(\text{ms}) \cap (r_{\text{f}} \circ (\text{ms})) = (\text{m}) \in \mathcal{S}$, which contradicts Assumption 1.

Now assume to the contrary that there are two relations r_1 and r_2 amongst r_{p} , r_{m} , r_{o} , r_{d} and r_{s} which are both minimal in the inclusion ordering. Note that both r_1 and r_2 are contained in

one of (pmodsf) and (pmodsf^{-1}) , since they are both subsets of r' . We consider the first case, the second one is dual.

By the choice of r_1, r_2 , $r_1 \cap r_2$ must be r_f or empty. If r_f is contained in every possible choice of r_1 and r_2 then $\mathcal{S} \subseteq \mathcal{B}_2$, so we consider the case when $r_1 \cap r_2$ is empty.

Assume first that $r_1 \subseteq (\text{odsf})$. If $r_1 \neq (\text{sf})$ then it can be checked that $r_1^{-1} \circ r_1 = \nu \in \mathcal{S}$. If $r_1 = (\text{sf})$ then $r_3 = r_1 \circ r_1 = (\text{dsf})$ and $\nu = r_3^{-1} \circ r_3$ belongs to \mathcal{S} anyway. By the minimality of r_2 , we have either $r_2 \cap \nu = \emptyset$ or $r_2 \subseteq \nu$. If $r_2 \cap \nu = \emptyset$ then $r_2 \subseteq (\text{pm})$ and $r_2 \circ r_2 = (\text{p}) \in \mathcal{S}$, which contradicts Assumption 1. If $r_2 \subseteq \nu$ then both r_1 and r_2 are contained in (odsf) . We may assume without loss of generality that $(o) \subseteq r_1$. It can then be checked that for all possible choices of r_1, r_2 , either $(r_1 \circ r_2) \cap r_2$ or $(r_1^{-1} \circ r_2) \cap r_2$ is a non-empty proper subset of r_2 , which contradicts the minimality of r_2 .

This completes the analysis of the case when $r_1 \subseteq (\text{odsf})$.

Now we only need to consider the case when the two distinct minimal relations r_1 and r_2 are r_p and r_m . Then we have $(\text{m}) \subseteq r_m \subseteq (\text{modsf})$, and it can be checked that, unless $r_m = (\text{mf})$, $r_4 = r_m^{-1} \circ r_m$ is either ν or $\nu \cup (\text{mm}^{-1})$. In the former case $r_4 \cap r_m$ is a non-empty proper subrelation of r_m , while in the latter one $r_4 \cap r_p$ is a non-empty proper subrelation of r_p , which contradicts the choice of r_1 and r_2 . If $r_m = (\text{mf})$ then $((\text{mf}) \circ (\text{m}^{-1}\text{f}^{-1})) \cap r_p = (\text{p})$, which contradicts Assumption 1. \square

Lemma 33 *With the assumptions above, if $(\text{pmosf}^{-1}) \in \mathcal{S}$ then \mathcal{S} is contained in one of the 18 maximal tractable subalgebras.*

Proof. Consider the relation r_d . If $r_d = (\equiv d)$ then $r_d \circ (\text{pmosf}^{-1})$ satisfies the conditions of Lemma 32. If $(\text{dd}^{-1}) \subseteq r_d \subseteq (\equiv \text{dd}^{-1})$ then, using $(\equiv \text{dd}^{-1}) \cap ((\equiv \text{dd}^{-1}) \circ (\text{pmosf}^{-1})) = (\text{dd}^{-1})$, we get \mathcal{S} is NP-complete by Lemma 31(2), which contradicts Assumption 2. Hence, $r_d \cap (\text{pp}^{-1}\text{mm}^{-1}\text{oo}^{-1}\text{ss}^{-1}\text{ff}^{-1}) \neq \emptyset$.

The relations r_p, r_m, r_o, r_s , and $r_{f^{-1}}$ must all be contained in (pmosf^{-1}) . Assume first that one of these five relations is contained in the other four. If r_m is contained in all of $r_p, r_o, r_s, r_{f^{-1}}$, then r_m coincides with one of $r_p, r_o, r_s, r_{f^{-1}}$ because $(\text{m}) \notin \mathcal{S}$, by Assumption 1. Further, if r_p, r_o, r_s or $r_{f^{-1}}$ is contained in the other four relations, then it is also contained in r_d or $r_{d^{-1}}$. Hence, \mathcal{S} is contained in $\mathcal{S}_p, \mathcal{S}_o, \mathcal{A}_2, \mathcal{A}_4, \mathcal{E}_p, \mathcal{E}_o, \mathcal{B}_1$, or \mathcal{B}_4 .

Now assume to the contrary that there are two relations r_1 and r_2 amongst r_p, r_m, r_o, r_s and $r_{f^{-1}}$ which are both minimal in the inclusion ordering. By the choice of r_1, r_2 , $r_1 \cap r_2$ must be empty.

Assume first that $r_1 \subseteq (\text{osf}^{-1})$. If $(o) \subseteq r_1$ then it can be checked that $r_1^{-1} \circ r_1 = \nu \in \mathcal{S}$. Then, by the minimality of r_2 , we have either $r_2 \cap \nu = \emptyset$ or $r_2 \subseteq \nu$. If $r_2 \cap \nu = \emptyset$ then $r_2 \subseteq (\text{pm})$ and $r_2 \circ r_2 = (\text{p}) \in \mathcal{S}$, which contradicts Assumption 1. If $r_2 \subseteq \nu$ then both r_1 and r_2 are contained in (osf^{-1}) , which contradicts Assumption 1. Hence we may assume that $r_1 = (\text{sf}^{-1})$, which implies that $(o) \subseteq r_2 \subseteq (\text{pmo})$ and hence $(r_1^{-1} \circ r_2)^* = (\text{dd}^{-1})$, so \mathcal{S} is NP-complete by Lemma 31(2), which contradicts Assumption 2. This completes the analysis of the case when $r_1 \subseteq (\text{osf}^{-1})$.

Now we only need to consider the case when the two distinct minimal relations r_1 and r_2 are r_p and r_m . Then we have $(\text{m}) \subseteq r_m \subseteq (\text{mosf}^{-1})$, and it can be checked that in all cases either $r_m^{-1} \circ r_m$ or $r_m \circ r_m^{-1}$ is a non-empty proper subrelation of r_p , which contradicts the choice of r_2 . \square

Lemma 34 *With the assumptions above, if $r' \not\subseteq (\text{dsf})^{\pm 1}$, $r' \not\subseteq (\text{pmos})^{\pm 1}$, and $r' \not\subseteq (\text{pmof}^{-1})^{\pm 1}$, then \mathcal{S} is contained in one of the 18 maximal tractable subalgebras.*

Proof. It follows from Lemma 30 that if $r' \cap (\text{pp}^{-1}\text{mm}^{-1}\text{oo}^{-1}) = \emptyset$ then r' (or its converse) is one of $(\text{ds}^{-1}), (\text{df}^{-1}), (\text{sf}^{-1}), (\text{dsf}^{-1})$, or (ds^{-1}f) . In all of these cases $r' \circ r'$ (or its converse) satisfies the conditions of Lemma 32 or Lemma 33.

Hence we may assume that $r' \cap (\text{pmo}) \neq \emptyset$. Now, using Lemma 30, it can straightforwardly be checked that, except for $r' = (\text{md})$ and (md^{-1}) , the relation $r' \circ r'$ satisfies the conditions

of Lemma 32 or Lemma 33. For $r' = (\text{md})$ or (md^{-1}) , the relation $(r' \circ r') \circ r'$ is (pmods) or $(\text{pmod}^{-1}\text{f}^{-1})$, respectively. Once again, Lemma 32 can be applied. \square

Lemma 35 *With the assumptions above, if \mathcal{S} contain relations r_1 and r_2 such that $r_1 \subseteq (\text{pmos})$ or $r_1 \subseteq (\text{pmof}^{-1})$, and $r_2 \subseteq (\text{dsf})$, then \mathcal{S} is contained in one of the 18 maximal tractable subalgebras.*

Proof. Let $r_3 = r_1 \circ r_1$ and $r_4 = r_2 \circ r_2$. Then we have $(\text{p}) \subseteq r_3 \subseteq (\text{pmos})$ or $(\text{p}) \subseteq r_3 \subseteq (\text{pmof}^{-1})$, and $(\text{d}) \subseteq r_4 \subseteq (\text{dsf})$. Now it is easy to check that either $r_3 \circ r_4$ or $r_4^{-1} \circ r_3$ satisfies the conditions of Lemma 32. \square

Lemma 36 *With the assumptions above, if $r' \subseteq (\text{dsf})$ then \mathcal{S} is contained in one of the 18 maximal tractable subalgebras.*

Proof. As noted in the proof of Lemma 32, we have $\nu \in \mathcal{S}$.

We consider the case when $r' = (\text{ds})$; the other cases are similar. Then, for every $r \in \mathcal{S}$, we have $r \cap (\text{ds})^{\pm 1} \neq \emptyset \Rightarrow (\text{s})^{\pm 1} \subseteq r$.

Consider the relation r_{O} . If it is asymmetric then we get the required result by Lemmas 34 and 35, and by Proposition 5, so assume it is not. It is clear that $r_{\text{O}} \subseteq \nu$. Suppose that $(\text{ss}^{-1}) \not\subseteq r_{\text{O}}$, say $(\text{s}^{-1}) \not\subseteq r_{\text{O}}$. Then $(\text{o}) \subseteq r_{\text{O}} \subseteq (\equiv \text{oo}^{-1}\text{dsff}^{-1})$. It is easy to check that if $r_1 = (\text{ds}) \circ r_{\text{O}}$ then $(\text{o}) \subseteq r_1$ and $(\equiv) \not\subseteq r_1$. This implies that $(\equiv) \not\subseteq r_{\text{O}}$. Since r_{O} is not asymmetric, we conclude that $(\text{oo}^{-1}) \subseteq r_{\text{O}}$ or $(\text{ff}^{-1}) \subseteq r_{\text{O}}$. If $(\text{oo}^{-1}) \not\subseteq r_{\text{O}}$ then $(\text{ff}^{-1}) = r_{\text{O}}^* \in \mathcal{S}$ and $(\text{pmods}) = (\text{ds}) \circ (\text{ff}^{-1}) \in \mathcal{S}$, that is, we can make use of Lemma 32. Suppose now that $(\text{oo}^{-1}) \subseteq r_{\text{O}}$; then r_{O} is either (oo^{-1}) or $(\text{oo}^{-1}\text{ff}^{-1})$. The case $r_{\text{O}} = (\text{oo}^{-1}\text{ff}^{-1})$ is impossible in view of $(\text{oo}^{-1}\text{ff}^{-1}) \cap ((\text{ds}) \circ (\text{oo}^{-1}\text{ff}^{-1})) = (\text{oo}^{-1}\text{f})$. If $r_{\text{O}} = (\text{oo}^{-1})$ then \mathcal{S} is NP-complete by Lemma 31(1), which contradicts Assumption 2. Therefore we may consider further in this proof that, for every $r \in \mathcal{S}$, $r \cap (\text{oo}^{-1}) \neq \emptyset$ implies $(\text{ss}^{-1}) \subseteq r$.

Consider the relation r_{P} . If it is asymmetric then we get the required result by Lemmas 34-35, so assume it is not. Suppose that $(\text{ss}^{-1}) \not\subseteq r_{\text{P}}$. Then $r_{\text{P}}^* \subseteq (\equiv \text{pp}^{-1}\text{mm}^{-1}\text{ff}^{-1})$. If $(\text{ff}^{-1}) \subseteq r_{\text{P}}^*$ then $r_1 = r_{\text{P}}^* \cap \nu (\in \mathcal{S})$ is either $(\equiv \text{ff}^{-1})$ or (ff^{-1}) . In both cases $(\text{ds}) \circ r_1 = (\text{pmods}) \in \mathcal{S}$, and the result follows from Lemma 32. We have $((\text{ds}) \circ (\equiv \text{pp}^{-1}\text{mm}^{-1})) \cap (\equiv \text{pp}^{-1}\text{mm}^{-1}) = (\text{pp}^{-1}\text{m}^{-1})$. Then, if $r_{\text{P}}^* \neq (\equiv)$, we get $r_{\text{P}} = (\text{pp}^{-1})$, and \mathcal{S} is NP-complete by Proposition 2(3), since $\{(\text{pp}^{-1}), (\text{ds})\}$ is contained in neither \mathcal{S}_{P} nor \mathcal{E}_{P} . The only remaining choice for r_{P}^* is (\equiv) . It is impossible, since $(\text{p}) \subseteq (\text{ds}) \circ r_{\text{P}}$, but $(\equiv) \not\subseteq (\text{ds}) \circ r_{\text{P}}$. Therefore we may consider further in this proof that, for every $r \in \mathcal{S}$, $r \cap (\text{pp}^{-1}) \neq \emptyset$ implies $(\text{ss}^{-1}) \subseteq r$.

A similar argument shows that if $(\text{ss}^{-1}) \not\subseteq r_{\text{M}}$ then the result follows. Assume that $(\text{ss}^{-1}) \subseteq r_{\text{M}}$. Now it can be easily verified that $\mathcal{S} \subseteq \mathcal{E}^*$ if $r_{\text{f}} \subseteq (\equiv \text{ff}^{-1})$; otherwise we have $r_{\text{f}} \cap (\text{ss}^{-1}) \neq \emptyset$, and $\mathcal{S} \subseteq \mathcal{B}_1$ if $(\text{s}^{-1}) \subseteq r_{\text{f}}$, while $\mathcal{S} \subseteq \mathcal{B}_2$ if $(\text{s}) \subseteq r_{\text{f}}$. \square

Lemma 37 *With the assumptions above, if $r' \subseteq (\text{pmos})$ or $r' \subseteq (\text{pmof}^{-1})$, then \mathcal{S} is contained in one of the 18 maximal tractable subalgebras.*

Proof. We shall consider the case $r' \subseteq (\text{pmos})$; the second case is dual. Note that $r' \neq (\text{ms})$ and $r' \neq (\text{mo})$; otherwise $((r')^{-1} \circ r') \cap r'$ is (s) or (o) , respectively.

Case 1. $(\text{p}) \not\subseteq r'$.

We have $(\text{o}) \subseteq r'$; otherwise $r' = (\text{ms})$ which contradicts our assumptions. If $r' = (\text{mos})$ then $((r')^{-1} \circ r') \cap r' = (\text{os}) \in \mathcal{S}$. We may therefore assume that $r' = (\text{os})$. Then, the proof is very similar to the one of Lemma 36.

Case 2. $(\text{p}) \subseteq r'$.

Since $(\text{pm}) \circ (\text{pm}) = (\text{p})$, r' cannot be (pm) and we can assume that $(\text{o}) \subseteq r'$ or $(\text{s}) \subseteq r'$. Then $r' \circ r'$ is one of (pmo) , (ps) , (pms) , and (pmos) . We can also assume that no relation satisfying the condition of Case 1 is contained in \mathcal{S} . This implies that r_{P} is a minimal relation in \mathcal{S} .

Subcase 2.1. $r' \circ r' = (\text{pmo})$.

We have $r_{\mathbf{p}} = (\text{po})$ or $r_{\mathbf{p}} = (\text{pmo})$. In both cases, every $r \in \mathcal{S}$ such that $r \cap (\text{pmo}) \neq \emptyset$ satisfies $(\mathbf{p}) \subseteq r$ because, as shown above, $(\text{mo}) \notin \mathcal{S}$.

Suppose that \mathcal{S} contains a non-trivial relation r_1 such that $r_1 \subseteq (\equiv \text{dd}^{-1}\text{ss}^{-1}\text{ff}^{-1})$. Define r_2 to be $(\text{pmo}) \circ r_1$ if $r_1 \not\subseteq (\equiv \text{sf}^{-1})$, and $(\text{pmo}) \circ r_1^{-1}$ otherwise. Then it can be easily checked that either $r_2^* = (\text{dd}^{-1})$ or else r_2 (or r_2^{-1}) satisfies the conditions of Lemma 32 or Lemma 33. If $(\text{dd}^{-1}) \in \mathcal{S}$ then $\{(\text{pmo}), (\text{dd}^{-1})\} \subseteq \mathcal{S}$, so \mathcal{S} is NP-complete by Lemma 31(2). We may assume now that every non-trivial $r \in \mathcal{S}$ satisfies $r \cap (\text{pp}^{-1}\text{mm}^{-1}\text{oo}^{-1}) \neq \emptyset$ and, consequently, $r \cap (\text{pp}^{-1}) \neq \emptyset$.

Arbitrarily choose $r \in \mathcal{S}$. If r is such that r^* is neither \emptyset nor (\equiv) , then $(\text{pp}^{-1}) \subseteq r$. Assume that $r^* = \emptyset$, that is, r is asymmetric. The required result follows from the previous lemmas if $r \not\subseteq (\text{pmos})^{\pm 1}$ and $r \not\subseteq (\text{pmof}^{-1})^{\pm 1}$. So we can assume that every asymmetric $r \in \mathcal{S}$ satisfies $(\mathbf{p}) \subseteq r \subseteq (\text{pmosf}^{-1})$ or $(\mathbf{p}) \subseteq r^{-1} \subseteq (\text{pmosf}^{-1})$.

Suppose now that $r^* = (\equiv)$; then $r \cap (\text{pp}^{-1}\text{mm}^{-1}\text{oo}^{-1})$ is contained either in (pmo) or in $(\text{p}^{-1}\text{m}^{-1}\text{o}^{-1})$. Without loss of generality, assume that $r \cap (\text{pp}^{-1}\text{mm}^{-1}\text{oo}^{-1}) \subseteq (\text{pmo})$. Then consider the relation $r_3 = (\text{pmo}) \circ r$. If $r \not\subseteq (\equiv \text{pmosf}^{-1})$ then either $r_3^* = (\text{dd}^{-1})$ (and then \mathcal{S} is NP complete by Lemma 31(2)) or r_3 is one of the relations satisfying the conditions of Lemma 32 (and then \mathcal{S} is contained in one of the 18 maximal tractable subalgebras). Now let $r \subseteq (\equiv \text{pmosf}^{-1})$. By our assumption, \mathcal{S} contains no non-trivial subrelation of $(\equiv \text{sf}^{-1})$ which implies that $(\mathbf{p}) \subseteq r$ unless $r = (\equiv)$.

We conclude that \mathcal{S} is contained in $\mathcal{S}_{\mathbf{p}}$ or in $\mathcal{E}_{\mathbf{p}}$.

Subcase 2.2. $r' \circ r' = (\text{ps})$.

We have $r' = r_{\mathbf{p}} = (\text{ps})$ so every $r \in \mathcal{S}$ such that $r \cap (\text{ps}) \neq \emptyset$ satisfies $(\mathbf{p}) \subseteq r$. Suppose that \mathcal{S} contains a non-trivial relation r_4 such that $r_4 \subseteq (\equiv \text{mm}^{-1}\text{oo}^{-1}\text{dd}^{-1}\text{ff}^{-1})$ and $r_4 \not\subseteq (\equiv \text{ff}^{-1})$. Then it can be verified that the relation $r_5 = (r' \circ r_4) \cap r_4$ is non-empty and $(\equiv) \not\subseteq r_5$. Therefore, we can assume that $(\equiv) \not\subseteq r_4$. This leads to a contradiction in view of $(r_4 \circ r') \cap r' = (\mathbf{p})$. The case $r_4 = (\text{ff}^{-1})$ is impossible because $((\text{ps}) \circ (\text{ff}^{-1})) \cap (\text{ps}) = (\mathbf{p})$. Finally, if r_4 is $(\equiv \text{ff}^{-1})$ or $(\equiv \text{f})$ then $(\text{ps}) \circ r_4 = (\text{pmods})$, and we can apply Lemma 32.

We may therefore assume that, for non-trivial every $r \in \mathcal{S}$, $r \cap (\text{pp}^{-1}\text{ss}^{-1}) \neq \emptyset$ and, consequently, $r \cap (\text{pp}^{-1}) \neq \emptyset$.

Arbitrarily choose a non-trivial $r \in \mathcal{S}$. If r is such that r^* is neither \emptyset nor (\equiv) , then $(\text{pp}^{-1}) \subseteq r$. Suppose that $r^* = \emptyset$. If $r \not\subseteq (\text{pmos})^{\pm 1}$ and $r \not\subseteq (\text{pmof}^{-1})^{\pm 1}$, then we get the required result by previous lemmas so we can assume that every asymmetric $r \in \mathcal{S}$ satisfies $(\mathbf{p}) \subseteq r \subseteq (\text{pmosf}^{-1})$ or $(\mathbf{p}) \subseteq r^{-1} \subseteq (\text{pmosf}^{-1})$.

If $r^* = (\equiv)$ then $r \cap (\text{pp}^{-1}\text{ss}^{-1})$ is either (ps) or $(\text{p}^{-1}\text{s}^{-1})$. Without loss of generality, assume that $r \cap (\text{pp}^{-1}\text{ss}^{-1}) = (\text{ps})$. Consider the relation $r_6 = (\text{ps}) \circ r$. If $r \cap (\text{m}^{-1}\text{o}^{-1}) \neq \emptyset$ or $(\text{d}^{-1}\text{f}) \subseteq r$ then r_6^* is a non-trivial subrelation of $(\text{mm}^{-1}\text{oo}^{-1}\text{dd}^{-1}\text{ff}^{-1})$ which contradicts our assumptions. Therefore, either $r \subseteq (\equiv \text{pmodsf})$, or $r \subseteq (\equiv \text{pmodsf}^{-1})$, or $r \subseteq (\equiv \text{pmod}^{-1}\text{sf}^{-1})$. Moreover, all $r \in \mathcal{S}$ such that $r \cap (\text{pp}^{-1}\text{ss}^{-1}) = (\text{ps})$ have to satisfy one (and the same) of these three conditions, since otherwise it is easy to generate a non-empty subrelation of (df) which would lead to a contradiction with our assumptions. We conclude that \mathcal{S} is contained in $\mathcal{S}_{\mathbf{p}}$ or in $\mathcal{E}_{\mathbf{p}}$.

Subcase 2.3. $r' \circ r' = (\text{pms})$ or $r' \circ r' = (\text{pmos})$.

Similar to previous subcases. □

6.2 Symmetric relations

To conclude the proof of Theorem 1, in this subsection we prove the following proposition.

Proposition 6 *Let \mathcal{S} be a subalgebra of \mathcal{A} such that $\text{bas}(\mathcal{S}) = \{(\equiv)\}$, which contains a symmetric relation r' such that $(\equiv) \not\subseteq r'$. Then \mathcal{S} is tractable if it is contained in one of the 18 algebras listed in Table 3. Otherwise \mathcal{S} is NP-complete*

To obtain this result we shall assume throughout this subsection that \mathcal{S} is a subalgebra of \mathcal{A} satisfying the following assumptions:

Assumption 1 $bas(\mathcal{S}) = \{\equiv\}$.

Assumption 2 \mathcal{S} is not NP-complete.

Assumption 3 $r' \in \mathcal{S}$ is a minimal symmetric relation such that $(\equiv) \not\subseteq r'$.

We show that the result holds for all possible choices of r' in Lemmas 40–41. These lemmas rely on the following NP-completeness results.

Proposition 7 *If r, s are symmetric relations such that $r \cap s = \emptyset$, both r and s are contained in neither $(\equiv ss^{-1})$ nor $(\equiv ff^{-1})$, then $\{r, s\}$ is NP-complete.*

The proof of this proposition can be found in Subsection A.3.

Lemma 38 *The following sets of relations are NP-complete:*

- 1) $\{(mm^{-1})\}$; and
- 2) $\{r, (ss^{-1}), (ff^{-1})\}$ when $r \in \{(oo^{-1}), (dd^{-1})\}$.

Proof. 1) We note that the set $\{(mm^{-1}), (pp^{-1}ss^{-1}ff^{-1})\}$ has been shown NP-complete in [15]. Let $r_1 = (mm^{-1}) \circ (mm^{-1}) = (\equiv pp^{-1}ss^{-1}ff^{-1})$ and $r_2 = (mm^{-1}) \circ r_1 = \top \setminus (\equiv)$. Thus, $(pp^{-1}ss^{-1}ff^{-1}) = r_1 \cap r_2$.

2) Let $r = (dd^{-1})$ and consider the following set of constraints:

$$a(dd^{-1})b, x(ss^{-1})a, x(ff^{-1})b, y(ff^{-1})a, y(ss^{-1})b.$$

The derived relation between x and y is (oo^{-1}) and NP-completeness of $\{(oo^{-1}), (dd^{-1})\}$ follows from Proposition 7.

The case when $r = (oo^{-1})$ is similar: $a(dd^{-1})b$ is replaced by $a(oo^{-1})b$, the derived relation is then (dd^{-1}) , and again we have NP-completeness by Proposition 7. \square

Lemma 39 *With the assumptions above, if \mathcal{S} contains $(\equiv ss^{-1})$, $(\equiv ff^{-1})$, and a non-trivial relation r_1 such that $r_1 \cap (\equiv ss^{-1}ff^{-1}) = \emptyset$ then \mathcal{S} is contained in one of the 18 maximal tractable subalgebras.*

Proof. Choose r_1 to be minimal. If $r_1 = (mm^{-1})$ then \mathcal{S} is NP-complete by Lemma 38(1). Hence we shall assume that $r_1 \neq (mm^{-1})$.

We have $(\equiv ss^{-1}) \circ (\equiv ff^{-1}) = (\equiv pmoo^{-1}dd^{-1}ss^{-1}ff^{-1}) \in \mathcal{S}$. Then either r_1 is asymmetric or $r_1 = r_1^* \subseteq (oo^{-1}dd^{-1})$. In the former case, we get the required result by Proposition 4, so we shall assume that r_1 is symmetric.

If $r_1 = (oo^{-1})$ or $r_1 = (dd^{-1})$, then $(ss^{-1}) = (\equiv ss^{-1}) \cap (r_1 \circ (\equiv ff^{-1})) \in \mathcal{S}$. Similarly, $(ff^{-1}) \in \mathcal{S}$. Hence, \mathcal{S} is NP-complete by Lemma 38(2), which contradicts Assumption 2.

It follows that $r_1 = (oo^{-1}dd^{-1})$. Since r_1 is minimal, we have $(oo^{-1}dd^{-1}) \subseteq r$ for every $r \in \mathcal{S}$ such that $(oo^{-1}dd^{-1}) \cap r \neq \emptyset$. Further, we have

$$(pmoo^{-1}dd^{-1}) = ((r_1 \circ (\equiv ff^{-1})) \cap ((\equiv ss^{-1}) \circ r_1)) \in \mathcal{S}.$$

In view of $(pm) \circ (pm) = (p)$, no non-empty subrelation of (pm) can belong to \mathcal{S} . This implies that $(oo^{-1}dd^{-1}) \subseteq r$ for every $r \in \mathcal{S}$ such that $(pmoo^{-1}dd^{-1}) \cap r \neq \emptyset$. Further, neither (sf) nor (sf^{-1}) can belong to \mathcal{S} because they give (s) being intersected with $(\equiv ss^{-1}) \in \mathcal{S}$. If $(\equiv sf) \in \mathcal{S}$ or $(\equiv sf^{-1}) \in \mathcal{S}$ then we can obtain $(\equiv s)$ and $(\equiv f)$, and, further, $(\equiv dsf)$. However, this contradicts the fact that every relation containing (d) must also contain $(oo^{-1}dd^{-1})$. Now it is straightforward to check that $\mathcal{S} \subseteq \mathcal{H}$. \square

By examining the composition table given in Table 2, it is easy to verify that $(\equiv) \subseteq r_1 \circ r_2$ if and only if there exists a non-trivial basic relation b such that $(b) \subseteq r_1$ and $(b^{-1}) \subseteq r_2$. In the next two lemmas, we shall make use of this fact.

Lemma 40 *With the assumptions above, if $r' \not\subseteq (ss^{-1}ff^{-1})$, then \mathcal{S} is contained in one of the 18 maximal tractable subalgebras.*

Proof. We may assume that $r' \not\subseteq (pp^{-1}mm^{-1})$; otherwise the result follows from Proposition 2(3).

By the minimality of r' , for every non-trivial relation r_1 in \mathcal{S} we have $r' \cap r_1 = r'$ or empty. Obviously, if every non-trivial relation in \mathcal{S} contains r' then \mathcal{S} is contained in one of the 18 maximal tractable subalgebras. Hence we shall assume that \mathcal{S} contains a non-trivial relation r_1 such that $r' \cap r_1 = \emptyset$.

Case 1. r_1 is asymmetric.

Apply Proposition 4.

Case 2. r_1 is symmetric.

It follows from Proposition 7 that if $r_1 \not\subseteq (\equiv ss^{-1})$ and $r_1 \not\subseteq (\equiv ff^{-1})$ then \mathcal{S} is NP-complete, which contradicts Assumption 2. We shall consider the case $r_1 \subseteq (\equiv ss^{-1})$; the second case is dual.

Note that we have $r' \cap (\equiv ss^{-1}) = \emptyset$. Also, $r' \neq (mm^{-1}ff^{-1})$, since otherwise $r' \cap (r' \circ (\equiv ss^{-1})) = (mm^{-1}f) \in \mathcal{S}$, which contradicts the minimality of r' . Hence, r' contains (bb^{-1}) where b is one of p , o , and d .

We have $r_1 \circ r_1^{-1} = (\equiv ss^{-1}) \in \mathcal{S}$. Let $r_2 = r' \circ (\equiv ss^{-1})$. Obviously, $r' \subseteq r_2$. Moreover, it can be easily checked by examining Table 2 that $r_2 \cap (\equiv ss^{-1}) = \emptyset$ and that, for every basic relation b_1 such that $b_1 \notin \{\equiv, s, s^{-1}\}$, at least one of (b_1) and (b_1^{-1}) is contained in r_2 . If $(bb^{-1}) \subseteq r$ for every $r \in \mathcal{S}$ such that $r \not\subseteq (\equiv ss^{-1})$ then \mathcal{S} is contained in one of \mathcal{S}_p , \mathcal{S}_o , and \mathcal{S}_d . Otherwise \mathcal{S} contains a non-empty relation r_3 such that $r_3 \not\subseteq (\equiv ss^{-1})$ and $r_3 \cap r' = \emptyset$ (because r' is minimal). Then $r_3 \cap r_2$ or $r_3 \cap r_2^{-1}$ is non-empty. Denote this non-empty relation by r_4 . Then we have $r_4 \cap (\equiv ss^{-1}) = \emptyset$ and $r_4 \cap r' = \emptyset$. Consider a minimal relation r_5 contained in r_4 . This minimal relation must be either symmetric or asymmetric. Therefore, unless $r_5 = (ff^{-1})$, we get the required result by Proposition 7 or 4, respectively. If $r_5 = (ff^{-1})$ then $r_5 \circ r_5 = (\equiv ff^{-1}) \in \mathcal{S}$ and $r' \cap (\equiv ss^{-1}ff^{-1}) = \emptyset$, and we can apply Lemma 39.

Case 3. r_1 is neither symmetric nor asymmetric.

We may assume that r_1 is minimal. Then, by minimality, we have $r_1^* = (\equiv)$. Since r' is symmetric and $r' \cap r_1 = \emptyset$, we obtain that $(\equiv) \not\subseteq r' \circ r_1$ and that $(\equiv) \not\subseteq r' \circ r_1^{-1}$. It follows that if one of $r' \circ r_1$ and $r' \circ r_1^{-1}$ has a non-empty intersection with r_1 or with r_1^{-1} then we get a contradiction with minimality of r_1 .

Now it can be checked that we indeed get this contradiction except when r_1 (or r_1^{-1}) is one of the relations $(\equiv m)$, $(\equiv s)$, and $(\equiv f)$. If $r_1 = (\equiv m)$ then $r_6 = (\equiv pm) = r_1 \circ r_1 \in \mathcal{S}$, and arguing as in the previous paragraph we can obtain a non-empty subrelation r_7 of pm which leads to a contradiction because $r_7 \circ r_7 = (p)$. If $r_1 = (\equiv s)$ then $r' \cap (\equiv ss^{-1}) = \emptyset$. Further $(\equiv ss^{-1}) = r_1 \circ r_1^{-1} \in \mathcal{S}$, and Case 2 applies. If $r_1 = (\equiv f)$ then the argument is dual. \square

Lemma 41 *With the assumptions above, if $r' \subseteq (ss^{-1}ff^{-1})$, then \mathcal{S} is contained in one of the 18 maximal tractable subalgebras.*

Proof. Case 1. $r' = (ss^{-1}ff^{-1})$.

We have $(ss^{-1}ff^{-1}) \subseteq r$ for every $r \in \mathcal{S}$ such that $r \cap (ss^{-1}ff^{-1}) \neq \emptyset$. If every non-trivial $r \in \mathcal{S}$ satisfies $(ss^{-1}ff^{-1}) \subseteq r$ then $\mathcal{S} \subseteq \mathcal{A}_1$. Suppose that \mathcal{S} has a non-trivial relation r_1 such that $r_1 \cap (ss^{-1}ff^{-1}) = \emptyset$. Then consider the relation $r_2 = r_1 \cap (r_1 \circ (ss^{-1}ff^{-1}))$. It is easy to check that r_2 is a non-empty subrelation of r_1 and that $(\equiv) \not\subseteq r_2$ (in fact, $r_2 = r_1 \setminus (\equiv)$). The relation r_2 contains some minimal relation that must be either symmetric or asymmetric. Now we obtain the required result by Lemma 40 or Proposition 4.

Case 2. Both (ss^{-1}) and (ff^{-1}) belong to \mathcal{S} .

We have $(oo^{-1}dd^{-1}) = ((ss^{-1}) \circ (ff^{-1}))^* \in \mathcal{S}$. Furthermore, $(ss^{-1}) \circ (ss^{-1}) = (\equiv ss^{-1})$ and $(ff^{-1}) \circ (ff^{-1}) = (\equiv ff^{-1})$ both belong to \mathcal{S} , and the result follows from Lemma 39.

Case 3. Exactly one of (ss^{-1}) and (ff^{-1}) belongs to \mathcal{S} .

Assume that $(ss^{-1}) \in \mathcal{S}$ and $(ff^{-1}) \notin \mathcal{S}$, the second case is dual.

If (ss^{-1}) is the only minimal relation in \mathcal{S} then every non-trivial relation in \mathcal{S} contains (ss^{-1}) , and we have $\mathcal{S} \subseteq \mathcal{A}_1$. Suppose that there exists a minimal relation $r_3 \in \mathcal{S}$ such that $r_3 \cap (ss^{-1}) = \emptyset$. Then we may assume that $(\equiv) \subseteq r_3$; otherwise we get the required result by Proposition 4 or by Lemma 40. Let $r_4 = r_3 \circ (ss^{-1})$. We have $(\equiv) \not\subseteq r_4$.

It is easy to verify that if $r_3 \not\subseteq (\equiv ff^{-1})$ then, for every basic relation b_1 such that $b_1 \notin \{\equiv, s, s^{-1}, f, f^{-1}\}$ and $(b_1) \subseteq r_3$, at least one of (b_1) and (b_1^{-1}) is contained in r_4 . This leads to a contradiction with minimality of r_3 .

If $r_3 \subseteq (\equiv ff^{-1})$ then every $r \in \mathcal{S}$ such that $r \cap (ff^{-1}) \neq \emptyset$ also satisfies $(\equiv) \subseteq r$. Further on, we have $(\equiv ff^{-1}) = r_3 \circ r_3^{-1} \in \mathcal{S}$ and $(pmoo^{-1}dd^{-1}ss^{-1}) = (ss^{-1}) \circ (\equiv ff^{-1}) \in \mathcal{S}$. If some non-empty subrelation of $(pmoo^{-1}dd^{-1})$ belongs to \mathcal{S} then we get the required result by Lemma 40 or Proposition 4. Else, every $r \in \mathcal{S}$ such that $r \cap (pmoo^{-1}dd^{-1}) \neq \emptyset$ also satisfies $(ss^{-1}) \subseteq r$, and we have $\mathcal{S} \subseteq \mathcal{E}^*$. \square

7 Conclusion

We have now completed the classification of complexity in Allen's algebra and shown that there exist exactly eighteen forms of tractability in this algebra. We did this by applying a technique from general algebra which has not been previously used in this context.

Both the result and the method can be used to classify the complexity in other temporal and spatial formalisms; a first application is given in Section 3. There are also strong connections with the analysis of complexity in temporal logics ([6]) which deserve further investigation.

It has already been established that the maximal tractable subalgebra, \mathcal{H} , can be used to speed up backtracking algorithms [39]. We believe that the complete description of tractability in Allen's algebra which is presented here may lead to new methods in approximate temporal reasoning, as one can uniquely loosen, in a minimal way, any set of interval constraints to obtain an instance of a given tractable case.

In this paper, we considered the problem of satisfiability of temporal constraints. However, there are other important tasks in temporal (and spatial) reasoning, for example, the task of answering queries in different types of constraint networks (see, e.g., [32]). The method and the results presented in this paper can contribute to further progress in tackling such tasks.

Finally, we note that many other constraint formalisms (not just temporal ones) are based on manipulating objects with intrinsic structural properties which can be captured by an appropriate algebra. This prompts us to conjecture that algebraic approaches to constraint manipulation, such as the one taken in this paper, or those presented in [8, 9], provide the appropriate reasoning tools across many different areas of constraint reasoning and artificial intelligence.

Acknowledgements

The research is partially supported by the UK EPSRC under grants GR/R22704 and GR/R29598, and the Swedish Research Council for Engineering Sciences (TFR) under grants 97-301 and 2000-361.

The authors are grateful to Andrei Bulatov for many helpful comments on an earlier version of the paper.

A Appendix

This appendix contains the proofs of Propositions 5 and 7. In the sequel, we will make frequent (but implicit) use of Proposition 3 in the following way. With the help of Lemma 30, it is easy to check that the relations r mentioned in Proposition 5 and pairs of relations $\{r, s\}$ mentioned in Proposition 7 are not contained in one of the 18 tractable subalgebras in Table 3. Therefore, we

conclude that $\{r\}$ or $\{r, s\}$ is NP-complete whenever we can derive, from this set, either a non-trivial basic relation or some set of relations whose NP-completeness was shown before. Recall that derivation is introduced in Section 4. B. Nebel's CSP solver [38] can considerably simplify calculating the derivations. Recall also that if a relation can be obtained from a given set by several derivations then it can be obtained by using a single derivation.

A.1 Model Transformations

This subsection contains the basics of *model transformations* which is a method for proving NP-completeness results. It is based on transforming a solution of one problem to a solution of a related problem. This method will be used many times in the proofs of Propositions 5 and 7.

Suppose T is a mapping on models of \mathcal{A} -SAT-instances with the same set of variables and let f_T be a function from the set of all basic relations to \mathcal{A} such that the following holds: for any model f of an \mathcal{A} -SAT-instance over a set V of variables, for any $x, y \in V$, and for any basic relation b , if $f(x)$ is related to $f(y)$ by (b) then $T(f(x))$ is related to $T(f(y))$ by $f_T(b)$. Then we say that T is a *model transformation* with *description* f_T . A description f_T can be extended to handle all relations $r \in \mathcal{A}$ in the obvious way: $f_T(r) = \bigcup_{b \in r} f_T(b)$.

The following lemma gives us a way of proving NP-completeness by using model transformations.

Lemma 42 *Let $\mathcal{R} = \{r_1, \dots, r_n\} \subseteq \mathcal{A}$ and $\mathcal{R}' = \{r'_1, \dots, r'_n\} \subseteq \mathcal{A}$ be such that $r'_k \subseteq r_k$ for all $1 \leq k \leq n$. If there exists a model transformation T with description f_T such that $f_T(r_k) \subseteq r'_k$ for every $1 \leq k \leq n$, then \mathcal{R} is NP-complete if and only if \mathcal{R}' is NP-complete.*

Proof. The proof of the only-if direction can be found in [15] and the proof of the other direction is analogous. \square

Let S be a finite set of real numbers. The *minimal distance in S* , $\text{MD}(S)$, is defined as

$$\text{MD}(S) = \min\{x - y \mid x, y \in S \wedge x > y\}.$$

For a model f of an \mathcal{A} -SAT-instance over a set V of variables, we define

$$\text{MD}(f) = \text{MD}(\{f(x^-), f(x^+) \mid x \in V\}),$$

where $f(x^-)$ and $f(x^+)$ denote the starting and the ending point of the interval $f(x)$, as in Section 3.

We continue by defining a number of model transformations. We shall use them with fixed descriptions which can be found in Table 4. We define the model transformation **shrink** as follows. Let f be a model of an \mathcal{A} -SAT-instance over $\{x_1, \dots, x_n\}$ and let $\epsilon = \text{MD}(f)/3$. Then $\text{shrink}(f) = f'$ where, for $1 \leq i \leq n$,

$$f'(x_i) = [f(x_i^-) + \epsilon, f(x_i^+) - \epsilon].$$

We can analogously define a model transformation **expand** by subtracting ϵ from $f(x^-)$ and adding ϵ to $f(x^+)$.

By ordering the intervals with respect to their length, we can obtain a number of useful model transformations. We define the model transformation **ordshrink** as follows. Let f be a model of an \mathcal{A} -SAT-instance over $\{x_1, \dots, x_n\}$, let $\epsilon = \text{MD}(f)/(2n)$ and rename the variables so that $|f(x_1)| \geq \dots \geq |f(x_n)|$. Then $\text{ordshrink}(f) = f'$ where, for $1 \leq i \leq n$,

$$f'(x_i) = [f(x_i^-) + i\epsilon, f(x_i^+) - i\epsilon].$$

We analogously define a model transformation **ordexpand**.

	\equiv	p	p ⁻¹	m	m ⁻¹	o	o ⁻¹	d	d ⁻¹	s	s ⁻¹	f	f ⁻¹
shrink	\equiv	p	p ⁻¹	p	p ⁻¹	o	o ⁻¹	d	d ⁻¹	s	s ⁻¹	f	f ⁻¹
expand	\equiv	p	p ⁻¹	o	o ⁻¹	o	o ⁻¹	d	d ⁻¹	s	s ⁻¹	f	f ⁻¹
ordshrink	dd ⁻¹	p	p ⁻¹	p	p ⁻¹	o	o ⁻¹	d	d ⁻¹	d	d ⁻¹	d	d ⁻¹
ordexpand	dd ⁻¹	p	p ⁻¹	o	o ⁻¹	o	o ⁻¹	d	d ⁻¹	o	o ⁻¹	o ⁻¹	o
leftordshrink	ff ⁻¹	p	p ⁻¹	p	p ⁻¹	o	o ⁻¹	d	d ⁻¹	o	o ⁻¹	f	f ⁻¹
leftordexpand	ff ⁻¹	p	p ⁻¹	o	o ⁻¹	o	o ⁻¹	d	d ⁻¹	d	d ⁻¹	f	f ⁻¹

Table 4: Model transformations.

We will also use model transformations that only change one of the endpoints of an interval. We define the model transformation `leftordshrink` as follows. Let f be a model of an \mathcal{A} -SAT-instance over $\{x_1, \dots, x_n\}$, let $\epsilon = \text{MD}(f)/(n+1)$ and rename the variables so that $|f(x_1)| \leq \dots \leq |f(x_n)|$. Then `leftordshrink`(f) = f' , where, for $1 \leq i \leq n$,

$$f'(x_i) = [f(x_i^-) + i\epsilon, f(x_i^+)].$$

A.2 Proof of Proposition 5

We will now show that if r is asymmetric, but not acyclic, then $\{r\}$ is NP-complete. The proof is largely based on the use of different derivations. Let $C_9(r)$ denote the relation (between x and y) derived from the following set of constraints:

$$\left\{ \begin{array}{l} xra_1 \quad a_1ra_2 \quad a_1ry \\ xra_2 \quad a_2ra_3 \quad a_2ry \\ xra_3 \quad a_3ra_1 \quad a_3ry \end{array} \right\}$$

$C_{9b}(r)$ is the relation derived from the same set of constraints but with xra_2 replaced by $xr^{-1}a_2$ and a_2ry replaced by $a_2r^{-1}y$. Finally, $C_{14}(r)$ denotes the relation derived from

$$\left\{ \begin{array}{l} xr^{-1}a_1 \quad a_1r^{-1}a_2 \quad a_1ry \\ xr^{-1}a_2 \quad a_1ra_3 \quad a_2r^{-1}y \\ xr^{-1}a_3 \quad a_1r^{-1}a_4 \quad a_3r^{-1}y \\ xra_4 \quad a_2r^{-1}a_3 \quad a_4r^{-1}y \\ \quad \quad a_2ra_4 \\ \quad \quad a_3r^{-1}a_4 \end{array} \right\}$$

Before the proof, we need one auxiliary lemma.

Lemma 43 $\{(pp^{-1}dd^{-1}), (oo^{-1})\}$ is NP-complete.

Proof. Let $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ be two graphs such that $V_1 = V_2$ and $E_1 \subseteq E_2$; in this case, we say that G_2 is a *supergraph* of G_1 . The GRAPH SANDWICH PROBLEM FOR PROPERTY Π is defined as follows:

INSTANCE: Two graphs, $G_1 = (V, E_1)$ and $G_2 = (V, E_2)$, such that G_2 is a supergraph of G_1 .

QUESTION: Is there a graph $G = (V, E)$ such that $E_1 \subseteq E \subseteq E_2$ and (V, E) has property Π ?

Two intervals *overlap*⁵ if their intersection is non-empty but neither one of them properly contains the other. An *overlap* graph (also known as a *circle* graph [17]) is an undirected graph $G = (V, E)$ for which there is an assignment of an interval to each vertex such that two vertices are adjacent iff the corresponding intervals overlap. The graph sandwich problem for overlap graphs (SP-OVERLAP) is NP-complete [20]. Clearly, this problem can be transformed to the satisfiability problem for $\{(\equiv \text{mm}^{-1}\text{oo}^{-1}), (\text{pp}^{-1}\text{dd}^{-1}\text{ss}^{-1}\text{ff}^{-1})\}$: given an instance $G_1 = (V, E_1)$, $G_2 = (V, E_2)$ of SP-OVERLAP, construct an instance as follows:

1. for each node $v \in V$, introduce a variable v ;
2. for each $(v, w) \in E_1$, add the constraint $v(\equiv \text{mm}^{-1}\text{oo}^{-1})w$; and
3. for each $(v, w) \notin E_2$, add the constraint $v(\text{pp}^{-1}\text{dd}^{-1}\text{ss}^{-1}\text{ff}^{-1})w$.

It is easy to see that the resulting instance has a model if and only if the given instance of SP-OVERLAP has a solution. Thus, $S = \{(\equiv \text{mm}^{-1}\text{oo}^{-1}), (\text{pp}^{-1}\text{dd}^{-1}\text{ss}^{-1}\text{ff}^{-1})\}$ is NP-complete. By applying Lemma 42 to S and the model transformation *expand*, we get that $S' = \{(\equiv \text{oo}^{-1}), (\text{pp}^{-1}\text{dd}^{-1}\text{ss}^{-1}\text{ff}^{-1})\}$ is NP-complete.

Now define the model transformation T_1 as follows: $T_1(f) = f'$, where f' is obtained from f by first setting $\epsilon = \text{MD}(f)/((n+1) \cdot c)$ where $c = \max\{|f(x_i)|; 1 \leq i \leq n\}$ and defining

$$f'(x_i) = [f(x_i^-) - |f(x_i)| \cdot \epsilon, f(x_i^+) + |f(x_i)| \cdot \epsilon].$$

Then, T_1 has the description f_{T_1} with the following properties: $f_{T_1}(\equiv) = (\equiv)$, $f_{T_1}(\text{mm}^{-1}) = (\text{pp}^{-1}\text{oo}^{-1})$, $f_{T_1}(\text{ss}^{-1}) = f_{T_1}(\text{ff}^{-1}) = (\text{dd}^{-1})$ and $f_{T_1}(b) = b$ for $b \in \{\text{p}, \text{p}^{-1}, \text{o}, \text{o}^{-1}, \text{d}, \text{d}^{-1}\}$. By applying Lemma 42 to this transformation and S' , we see that $S'' = \{(\equiv \text{oo}^{-1}), (\text{pp}^{-1}\text{dd}^{-1})\}$ is NP-complete. Note that we cannot replace T_1 with the very similar *ordexpand* transformation since *ordexpand* changes the (\equiv) relation.

Finally, define the model transformation T_2 as follows: $T_2(f) = f'$, where f' is obtained from f by first setting $\epsilon = \text{MD}(f)/(n+1)$ and defining

$$f'(x_i) = [f(x_i^-) + i\epsilon, f(x_i^+) + i\epsilon].$$

It is easy to see that T_2 has the description f_{T_2} with the following properties: $f_{T_2}(\equiv) = (\text{oo}^{-1})$, $f_{T_2}(\text{mm}^{-1}) = (\text{pp}^{-1}\text{oo}^{-1})$, $f_{T_2}(\text{ss}^{-1}) = f_{T_2}(\text{ff}^{-1}) = (\text{oo}^{-1}\text{dd}^{-1})$ and $f_{T_2}(b) = b$ for $b \in \{\text{p}, \text{p}^{-1}, \text{o}, \text{o}^{-1}, \text{d}, \text{d}^{-1}\}$. By applying Lemma 42 to this transformation and S'' , we have shown that $\{(\text{oo}^{-1}), (\text{pp}^{-1}\text{dd}^{-1})\}$ is NP-complete. \square

Proof. (of Proposition 5) It is clear that sets $\{r\}$ and $\{r^{-1}\}$ are NP-complete simultaneously, so it suffices to consider only one of them.

We assume first that $r \cap (\text{p}^{-1}\text{m}^{-1}\text{o}^{-1}) = \emptyset$. If $r \cap (\text{pmo}) = \emptyset$, then it follows from Lemma 30 that r or r^{-1} equals $(\text{ds}^{-1}\text{f}^{-1})$; however, $C_9((\text{ds}^{-1}\text{f}^{-1})) = (\text{d})$. Assume now that $r \cap (\text{pmo}) \neq \emptyset$. By using Lemma 30 once again, one of the following holds:

1. $(\text{ds}^{-1}) \subseteq r$;
2. $(\text{d}^{-1}\text{f}) \subseteq r$; or
3. $(\text{s}^{-1}\text{f}) \subseteq r$.

Case 1: $(\text{ds}^{-1}) \subseteq r$.

If $r \notin \{(\text{mds}^{-1}), (\text{mds}^{-1}\text{f}), (\text{mds}^{-1}\text{f}^{-1})\}$, then $C_9(r) = (\text{pmod})$ and $C_9(r) \cap r^{-1} = (\text{s})$. If $r = (\text{mds}^{-1}\text{f})$ or $r = (\text{mds}^{-1}\text{f}^{-1})$, then $C_9(r) = (\text{d})$ or (pd) , respectively, and $C_9(r) \cap r = (\text{d})$. Finally, if

⁵Note that this notion is different from the notion ‘‘One interval overlaps another one’’ used in definition of the basic relation o .

$r = (\text{mds}^{-1})$, then $((\text{m}^{-1}\text{d}^{-1}\text{s}) \circ (\text{mds}^{-1})) \cap (\text{mds}^{-1}) = (\text{ds}^{-1})$ and $((\text{ds}^{-1}) \circ (\text{ds}^{-1})) \cap (\text{m}^{-1}\text{d}^{-1}\text{s}) = (\text{m}^{-1})$.

Case 2: $(\text{d}^{-1}\text{f}) \subseteq r$.

Dual to case 1.

Case 3: $(\text{s}^{-1}\text{f}) \subseteq r$.

We can assume that $(\text{dd}^{-1}) \cap r = \emptyset$ since one of the previous two cases applies otherwise. If $r \neq (\text{ms}^{-1}\text{f})$, then one of the following holds:

1. $(\text{o}) \not\subseteq r$. This implies that $C_9(r) = (\text{p})$ and NP-completeness follows immediately.
2. $(\text{o}) \subseteq r$. Then, $C_9(r) = (\text{pmo})$, $(\text{pmo}) \circ r = (\text{pmodd}^{-1}\text{sf}^{-1})$ and we can obtain the relation (dd^{-1}) . Let $R = (\text{d})$, $R_1 = (\text{pmo})$, $R_2 = (\text{dd}^{-1})$ and NP-completeness of $\{r\}$ follows from Lemma 1.

Finally, if $r = (\text{ms}^{-1}\text{f})$, then consider the following set of constraints:

$$\left\{ \begin{array}{l} xra_1 \\ a_1ra_2 \\ a_2rx \\ yra_1 \\ yra_2 \end{array} \right\}$$

The relation between x and y derived from these constraints is $(\text{pp}^{-1}\text{dd}^{-1})$. Next, the relation (oo^{-1}) is derived from the following set of constraints:

$$\left\{ \begin{array}{lll} xra_1 & a_1r_1a_2 & a_1ry \\ xr_1a_2 & a_2r_1a_3 & a_2r_1y \\ xra_3 & a_3r_1a_1 & a_3ry \end{array} \right\}$$

Consequently, NP-completeness of $\{(\text{ms}^{-1}\text{f})\}$ follows from Lemma 43.

Finally, we consider the case when $r \cap (\text{pmo}) \neq \emptyset$ and $r \cap (\text{p}^{-1}\text{m}^{-1}\text{o}^{-1}) \neq \emptyset$; then Lemma 30 implies that r is not acyclic. The proof considers four cases depending on the value of $r \cap (\text{pp}^{-1}\text{oo}^{-1}\text{mm}^{-1})$. For every $r_1 \in \mathcal{A}$, let r_1^2 denote the relation $r_1 \circ r_1$. Recall, that \top denotes the union of all basic relations.

In the following, Lemma 2 significantly reduces the number of cases to be considered. For instance, by showing that $\{(\text{mo}^{-1}\text{ds})\}$ is NP-complete, we also know that $\{(\text{m}^{-1}\text{odf})\}$ is NP-complete.

Case 1: $r \cap (\text{pp}^{-1}\text{oo}^{-1}\text{mm}^{-1}) \in \{(\text{po}^{-1}), (\text{pmo}^{-1}), (\text{pm}^{-1}\text{o}^{-1})\}$.

Suppose first that $r = r' \cup (\text{po}^{-1})$.

r'	$C_9(r)$	NP-completeness of $\{r\}$
\emptyset	(p)	$C_9(r) = (p)$
(d)	(pmods)	$C_9(r) \cap r^{-1} = (o)$
(s)	(pds)	$C_9(r)^2 \cap r^{-1} = (o)$
(s^{-1})	(p)	$C_9(r) = (p)$
(ds)	(pmods)	$C_9(r) \cap r^{-1} = (o)$
(ds^{-1})	(pmods)	$(C_9(r) \cap r^{-1})^2 \cap r = (p)$
(df)	(pmods)	$C_9(r) \cap r^{-1} = (o)$
(df^{-1})		$C_{14}(r)^* \cap r = (o^{-1}df^{-1})$
(sf)	(pds)	$C_9(r)^2 \cap r^{-1} = (o)$
(sf^{-1})	$\top \setminus (\equiv s^{-1}f)$	$C_9(r)^* \cap r = (po^{-1})$
($s^{-1}f$)	(p)	$C_9(r) = (p)$
(dsf)	(pmods)	$C_9(r) \cap r^{-1} = (o)$
(dsf^{-1})	$\top \setminus (\equiv s^{-1})$	$C_9(r)^* \cap r = (po^{-1}df^{-1})$
($ds^{-1}f$)	(pmods)	$(C_9(r) \cap r^{-1})^2 \cap r = (p)$
($ds^{-1}f^{-1}$)	$\top \setminus (\equiv s^{-1})$	$C_9(r) \cap r = (po^{-1}df^{-1})$

Note that, in the table above, if $r' = (df^{-1})$, that is, if $r = (po^{-1}df^{-1})$ then $C_{14}(r) = \top \setminus (\equiv s^{-1})$, and we can obtain $(o^{-1}df^{-1})$ as shown in the table; NP-completeness of $(o^{-1}df^{-1})$ follows from Lemma 21 by using Lemma 2. If $r \cap (pp^{-1}mm^{-1}oo^{-1})$ is (pmo^{-1}) or $(pm^{-1}o^{-1})$, then NP-completeness of $\{r\}$ follows from the proof above by using Lemma 42 with model transformations shrink and expand, respectively.

Case 2: $r \cap (pp^{-1}oo^{-1}mm^{-1}) = (pm^{-1})$. Let $r = r' \cup (pm^{-1})$.

r'	$C_9(r)$	NP-completeness of $\{r\}$
\emptyset	(p)	$C_9(r) = (p)$
(d)	(pmods)	$C_9(r) \cap r^{-1} = (m)$
(s)	(p)	$C_9(r) = (p)$
(s^{-1})	(p)	$C_9(r) = (p)$
(ds)	(pmods)	$C_9(r) \cap r^{-1} = (m)$
(ds^{-1})	(pmods)	$C_9(r) \cap r^{-1} = (ms)$
(df)	(pmods)	$C_9(r) \cap r^{-1} = (m)$
(df^{-1})	(pmods)	$C_9(r) \cap r^{-1} = (m)$
(sf)	(pds)	$C_9(r)^2 \cap r^{-1} = (m)$
(sf^{-1})	(pmo)	$C_9(r) \cap r^{-1} = (m)$
($s^{-1}f$)	(p)	$C_9(r) = (p)$
(dsf)	(pmods)	$C_9(r) \cap r^{-1} = (m)$
(dsf^{-1})	(pmods)	$C_9(r) \cap r^{-1} = (m)$
($ds^{-1}f$)	(pmods)	$C_9(r) \cap r^{-1} = (ms)$
($ds^{-1}f^{-1}$)	($pp^{-1}modd^{-1}sf^{-1}$)	$C_9(r) \cap r^{-1} = (p^{-1}md^{-1}s)$

The relation (ms) generates (s) since $((m^{-1}s^{-1}) \circ (ms)) \cap (ms) = (s)$.

Case 3: $r \cap (pp^{-1}oo^{-1}mm^{-1}) = (pm^{-1}o)$. Let $r = r' \cup (pm^{-1}o)$.

r'	$C_9(r)$	NP-completeness of $\{r\}$
\emptyset	(p)	$C_9(r) = (p)$
(d)	(pmods)	$C_9(r) \cap r^{-1} = (m)$
(s)	(p)	$C_9(r) = (p)$
(s^{-1})	(pmo)	$C_9(r) \cap r^{-1} = (m)$
(ds)	(pmods)	$C_9(r) \cap r^{-1} = (m)$
(ds^{-1})	($pp^{-1}moo^{-1}dd^{-1}s$)	$(C_9(r) \cap r)^2 \cap r^{-1} = (ms)$
(df)	(pmods)	$C_9(r) \cap r^{-1} = (m)$
(df^{-1})	(pmods)	$C_9(r) \cap r^{-1} = (m)$
(sf)	(pmods)	$C_9(r) \cap r^{-1} = (m)$
(sf^{-1})	(pmo)	$C_9(r) \cap r^{-1} = (m)$
($s^{-1}f$)	($pp^{-1}moo^{-1}dd^{-1}$)	$C_9(r) \cap r^{-1} = (p^{-1}mo^{-1})$
(dsf)	(pmods)	$C_9(r) \cap r^{-1} = (m)$
(dsf^{-1})	(pmods)	$C_9(r) \cap r^{-1} = (m)$
($ds^{-1}f$)	($pp^{-1}moo^{-1}dd^{-1}sf$)	$C_9(r) \cap r^{-1} = (p^{-1}mo^{-1}d^{-1}s)$
($ds^{-1}f^{-1}$)	($pp^{-1}moo^{-1}dd^{-1}sf^{-1}$)	$C_9(r) \cap r^{-1} = (p^{-1}mo^{-1}d^{-1}s)$

Case 4: $r \cap (pp^{-1}oo^{-1}mm^{-1}) = (mo^{-1})$. Let $r = r' \cup (mo^{-1})$.

r'	$C_9(r)$	NP-completeness of $\{r\}$
\emptyset	\emptyset	$r^2 \cap r = (o^{-1})$
(d)	\emptyset	$(r^2 \cap r)^2 \cap r^{-1} = (m^{-1})$
(s)	\emptyset	$r^2 \cap r^{-1} = (m^{-1}o)$
(s^{-1})	\emptyset	$((r^{-1} \circ r) \cap r)^2 \cap r^{-1} = (m^{-1})$
(ds)	\emptyset	$((r^{-1} \circ r) \cap r)^2 \cap r^{-1} = (m^{-1})$
(ds^{-1})	\emptyset	$((r^{-1} \circ r) \cap r)^2 \cap r^{-1} = (m^{-1})$
(df)	(d)	$C_9(r) = (d)$
(df^{-1})	($pp^{-1}m^{-1}oo^{-1}f$)	$(C_9(r) \cap r)^2 \cap r^{-1} = (m^{-1}f)$
(sf)	(d)	$C_9(r) = (d)$
(sf^{-1})	($pp^{-1}m^{-1}o^{-1}dd^{-1}$)	$C_9(r) \cap r = (o^{-1})$
($s^{-1}f$)	\emptyset	$C_{9b}(r) = (mo^{-1})$
(dsf)	(d)	$C_9(r) = (d)$
(dsf^{-1})	($pp^{-1}m^{-1}oo^{-1}dd^{-1}sf$)	$C_9(r) \cap r^{-1} = (m^{-1}od^{-1}f)$
($ds^{-1}f$)	(d)	
($ds^{-1}f^{-1}$)	($pp^{-1}m^{-1}oo^{-1}dd^{-1}f$)	$C_9(r) \cap r^{-1} = (m^{-1}od^{-1}f)$

A.3 Proof of Proposition 7

This subsection contains the proof of Proposition 7. We will use a short-hand notation for relations of the type (bb^{-1}) by writing (O) to denote (oo^{-1}) , (S) to denote (ss^{-1}) and so on. We will also use combinations of these macro relations—for instance, by writing (PO), we mean the relation $(pp^{-1}oo^{-1})$.

Let s be a symmetric relation such that $s \not\subseteq (\equiv S)$ and $s \not\subseteq (\equiv F)$. We write “ \hat{s} is NP-complete” to denote that, for every symmetric relation r such that $r \cap s = \emptyset$, $r \not\subseteq (\equiv S)$ and $r \not\subseteq (\equiv F)$, $\{r, s\}$ is NP-complete. When we show results of the form “ \hat{s} is NP-complete”, we will tacitly assume

that r' is an arbitrary symmetric relation satisfying the requirements stated above. Thus, we can formulate Proposition 7 as follows:

if s is a symmetric relation such that $s \not\subseteq (\equiv \text{S})$ and $s \not\subseteq (\equiv \text{F})$, then \hat{s} is NP-complete.

To prove this result, we begin by showing that \hat{s} is NP-complete for all $s \subseteq (\equiv \text{MSF})$ such that $s \not\subseteq (\equiv \text{S})$ and $s \not\subseteq (\equiv \text{F})$; this proof can be found in Subsection A.3.1.

Next, we show that we do not have to care about the (\equiv) relation. More precisely, assume that $\{s, s'\}$ is NP-complete for all choices of s, s' such that

1. s and s' are symmetric relations;
2. $s \cap s' = \emptyset$;
3. $(\equiv) \not\subseteq s$ and $(\equiv) \not\subseteq s'$;
4. $s \not\subseteq (\text{S})$ and $s' \not\subseteq (\text{S})$; and
5. $s \not\subseteq (\text{F})$ and $s' \not\subseteq (\text{F})$.

We show that $X = \{s \cup (\equiv), s'\}$ is NP-complete for all choices of s, s' satisfying the requirements above. If $s \subseteq (\text{MSF})$ or $s' \subseteq (\text{MSF})$, then X is NP-complete by Lemma 52. Hence, we assume that each of s and s' contains at least one of the relations P, D, and O. Let $s_{\equiv} = s \cup (\equiv)$. It is easy to realize that $(\equiv) \not\subseteq s_{\equiv} \circ s'$ by inspecting the composition table. Furthermore, the following result can easily be shown: if $B, B' \in \{\text{P}, \text{D}, \text{O}\}$, $B \neq \text{D}$ and $B' \neq \text{P}$, then $(B) \subseteq B \circ B'$. If $(\text{D}) \not\subseteq s_{\equiv}$, then $s_1 = s_{\equiv} \cap (s_{\equiv} \circ s')^*$ is a non-trivial symmetric relation not containing (\equiv) and it can be checked that the set $\{s_1, s'\}$ satisfies the conditions above and implying that it is NP-complete. Hence, $\{s_{\equiv}, s'\}$ is NP-complete. Now assume $(\text{D}) \subseteq s_{\equiv}$. If $s' \cap (\text{OSF}) \neq \emptyset$, then $(\text{D}) \subseteq s_{\equiv} \circ s'$ and we can reason as above to show that $\{s_{\equiv}, s'\}$ is NP-complete. Otherwise, $s' \subseteq (\text{PM})$ and NP-completeness follows from Proposition 2(3), since $\{s_{\equiv}, s'\}$ is contained in neither \mathcal{S}_{p} nor \mathcal{E}_{p} , or from Lemma 38(1).

Hence, we can now restrict our attention to pairs of relations s, s' satisfying conditions 1-5 and such that $s \cap (\text{POD}) \neq \emptyset$ and $s' \cap (\text{POD}) \neq \emptyset$. These proofs are collected in Subsection A.3.2.

A.3.1 Below $(\equiv \text{MSF})$

Lemma 44 $\{(\text{OD}), (\text{SF})\}$ and $\{(\text{OD}), (\equiv \text{SF})\}$ are NP-complete.

Proof. We note that $((\text{OD}) \circ (\equiv \text{SF})) \cap (\equiv \text{SF}) = (\text{SF})$ so it is sufficient to give a proof for the set $\{(\text{OD}), (\text{SF})\}$. The proof is by a polynomial-time reduction from the NP-complete problem NOT-ALL-EQUAL SATISFIABILITY [16].

An instance of NOT-ALL-EQUAL SATISFIABILITY consists of a set U of Boolean variables, and a collection C of *clauses* over U , where each clause is a set of three *literals*, and a literal is a variable or a negated variable. The question is whether there is an assignment of truth variables to the variables such that each clause contains at least one true literal and at least one false literal.

To obtain the reduction from NOT-ALL-EQUAL SATISFIABILITY, we design three “gadgets”, that is, small sets of constraints with convenient properties. The first corresponds to a Boolean variable, the second corresponds to a clause, and the third ensures that the variables are connected to the clauses in the appropriate way. Hence there are three parts to the construction.

Let P be an instance of NOT-ALL-EQUAL SATISFIABILITY. We construct a corresponding instance I of $\mathcal{A}\text{-SAT}(\{(\text{OD}), (\text{SF})\})$ as follows:

1. For each variable $u \in U$,
 - introduce variables v_{u_1}, v_{u_2} and v_{u_3} in V ;
 - impose the constraint (SF) on the edges (v_{u_1}, v_{u_2}) and (v_{u_2}, v_{u_3}) ;

- impose the constraint (OD) on the edge (v_{u_1}, v_{u_3}) .
2. For each clause $c \in C$,
 - introduce variables $v_{c_1}, v_{c_2}, v_{c_3}$ and v_{c_4} in V ;
 - impose the constraint (SF) on the edges (v_{c_1}, v_{c_2}) , (v_{c_2}, v_{c_3}) and (v_{c_3}, v_{c_4}) ;
 - impose the constraint (OD) on the edge (v_{c_1}, v_{c_4}) .
 3. For each literal c_i , $i = 1, 2, 3$ in each clause c ,
 - introduce variables v'_{c_i} and v''_{c_i} in V and impose the constraint (SF) on the edge (v'_{c_i}, v''_{c_i}) ;
 - impose the constraint (SF) on the edge (v_{c_i}, v'_{c_i}) and impose the constraint (OD) on the edges (v_{c_i}, v'_{c_i}) and $(v_{c_{i+1}}, v''_{c_i})$;
 - if c_i is the (unnegated) variable u , then impose the constraint (SF) on the edge (v'_{c_i}, v_{u_2}) and impose the constraint (OD) on the edges (v'_{c_i}, v_{u_1}) and (v''_{c_i}, v_{u_2}) ;
 - If c_i is the negated variable u , then impose the constraint (SF) on the edge (v'_{c_i}, v_{u_3}) and impose the constraint (OD) on the edges (v'_{c_i}, v_{u_2}) and (v''_{c_i}, v_{u_3}) .

Clearly, this construction can be carried out in polynomial time and we will now show that I has a solution if and only if P has a solution. First, assume that I has a solution. We will use this to construct a corresponding solution to P . Consider a variable $u \in U$. Because of the constraints imposed in part (1) of the above construction, exactly one of the pairs (v_{u_1}, v_{u_2}) and (v_{u_2}, v_{u_3}) must be related by f or f^{-1} . If it is the pair (v_{u_1}, v_{u_2}) , then we assign the value T (true) to u , otherwise we assign the value F (false) to u .

Now consider each clause $c \in C$. Because of the constraints imposed in part (3) of the construction above, the relation between v_{c_i} and $v_{c_{i+1}}$ is f or f^{-1} if and only if the corresponding literal is assigned the value T . Finally, because of the constraints imposed in part (2) of the above construction, v_{c_i} and $v_{c_{i+1}}$ must be related by f or f^{-1} for at least one and at most two of the 3 possibilities $i = 1, 2, 3$. Hence the chosen assignment gives at least one true literal and one false literal in each clause, and so is a solution to P .

Conversely, assume that P has a solution σ . We will use this to construct a corresponding solution to I .

Consider the variables v_{u_1}, v_{u_2} and v_{u_3} in V , which are associated with the variable $u \in U$. Assign v_{u_1} the interval $[5, 8]$, and assign v_{u_3} the interval $[6, 7]$. If u is assigned the value T in σ , then assign v_{u_2} the interval $[6, 8]$, otherwise assign v_{u_2} the value $[5, 7]$.

Now consider the variables $v_{c_1}, v_{c_2}, v_{c_3}$ and v_{c_4} in V , which are associated with the clause $c \in C$. Since σ is a solution, it must assign values to the literals c_1, c_2, c_3 which contain at least one true value and at least one false value. There are therefore 6 possibilities, and we assign intervals to the variables $v_{c_1}, v_{c_2}, v_{c_3}$ and v_{c_4} in each case according to the following table:

c_1	c_2	c_3	v_{c_1}	v_{c_2}	v_{c_3}	v_{c_4}
T	T	F	$[1, 12]$	$[2, 12]$	$[3, 12]$	$[3, 10]$
T	F	T	$[1, 12]$	$[2, 12]$	$[2, 10]$	$[3, 10]$
F	T	T	$[1, 12]$	$[1, 10]$	$[2, 10]$	$[3, 10]$
F	F	T	$[1, 12]$	$[1, 11]$	$[1, 10]$	$[3, 10]$
F	T	F	$[1, 12]$	$[1, 11]$	$[3, 11]$	$[3, 10]$
T	F	F	$[1, 12]$	$[3, 12]$	$[3, 11]$	$[3, 10]$

It is easy to check that each of these assignments satisfies all constraints on the variables $v_{c_1}, v_{c_2}, v_{c_3}$ and v_{c_4} .

It remains to show that each of these assignments can be extended to a complete solution to I by assigning the remaining variables $v'_{c_1}, v''_{c_1}, v'_{c_2}, v''_{c_2}, v'_{c_3}$ and v''_{c_3} appropriately. To show this we note that, for $i = 1, 2, 3$, if the literal c_i is assigned the value T in σ , then in order to satisfy the

constraints on v'_{c_i} and v''_{c_i} the starting point of v'_{c_i} must equal the starting point of v_{u_1} or v_{u_2} and the starting point of v''_{c_i} must equal the starting point of v_{c_i} . In this case we simply assign the ending points of both of these intervals the value 13. Similarly, if the literal c_i is assigned the value F in σ , then the ending points of these intervals are forced to take certain (distinct) values, and we assign the starting points of these intervals the value 0. It is easy to check that this assignment satisfies all the constraints. For example, if $c = \{x, \neg y, z\}$ and $\sigma(x) = T, \sigma(y) = T$, and $\sigma(z) = F$ then these variables are assigned the values $[6, 13], [1, 13], [0, 7], [0, 12], [0, 7]$ and $[0, 11]$, respectively. \square

Lemma 45 $(\widehat{\text{SF}})$ is NP-complete.

Proof. we have $(B) \subseteq (\text{SF}) \circ (B)$ when $B \in \{\text{P}, \text{M}, \text{O}, \text{D}\}$. It can be easily verified that $(\equiv) \not\subseteq r_1 = (r' \cap ((\text{SF}) \circ r'))^*$. Consequently, we may assume that $(\equiv) \not\subseteq r'$. If $r' \subseteq (\text{PM})$ then NP-completeness follows from Proposition 2(3) or Lemma 38(1), so we assume that $r' \cap (\text{OD}) \neq \emptyset$.

To show NP-completeness of the remaining cases, we introduce the *OD-switch*. The switch is an instance Γ on five variables a, b, c, x, y :

$$\{a(\text{SF})b, a(\text{SF})c, x(\text{SF})b, x(\text{SF})c, y(\text{SF})b, y(\text{SF})c\}.$$

It has the following properties (which can easily be checked):

1. if $b(\text{PM})c$ holds, then $x(\equiv)y$;
2. if $b(\text{D})c$ holds, then $x(\equiv \text{O})y$; and
3. if $b(\text{O})c$ holds, then $x(\equiv \text{D})y$.

It can be checked that one of the relations $(\equiv \text{O}), (\equiv \text{D}), (\equiv \text{OD})$ is derived from $\Gamma \cup \{br'c\}$. Therefore we can obtain one of $(\text{O}), (\text{D}), (\text{OD})$ as described above. For (OD) , apply Lemma 44. If we obtain (O) or (D) then apply Lemma 1 with $R = (\text{o}), R_1 = (\text{SF}), R_2 = (\text{O})$ or with $R = (\text{d}), R_1 = (\text{SF}), R_2 = (\text{D})$ if $r' = (\text{D})$, respectively. \square

Lemma 46 $(\widehat{\equiv \text{SF}})$ are NP-complete.

Proof. This proof is analogous to the previous one because the properties of the OD-switch are the same if we replace the relation (SF) with $(\equiv \text{SF})$.

Lemma 47 $(\widehat{\equiv \text{M}})$ is NP-complete.

Proof. Consider the following:

$$\begin{aligned} (\equiv \text{M}) \circ (\text{P}) &= (\text{PMoD}^{-1}\text{s}^{-1}\text{f}^{-1}) & (\equiv \text{M}) \circ (\text{O}) &= (\text{POdsf}) \\ (\equiv \text{M}) \circ (\text{D}) &= (\text{PODsf}) & (\equiv \text{M}) \circ (\text{S}) &= (\text{p}^{-1}\text{mo}^{-1}\text{dSf}) \\ (\equiv \text{M}) \circ (\text{F}) &= (\text{pm}^{-1}\text{odsF}) \end{aligned}$$

If $(\text{PSF}) \cap r' \neq \emptyset$, then $(\equiv) \not\subseteq r_1 = (\equiv \text{M}) \circ r'$ but at least one of $(\text{m}), (\text{m}^{-1})$ is a member of r_1 . Consequently, the relation $r_1 \cap (\equiv \text{M})$ implies NP-completeness by either Proposition 3 or Lemma 38(1). Otherwise, $r' \subseteq (\text{OD})$ and $(\text{PO}) \subseteq ((\equiv \text{M}) \circ r')^* \subseteq (\text{POD})$. Since $(\equiv \text{M}) \circ (\equiv \text{M}) = (\equiv \text{PSF})$, we can obtain the relation (P) and NP-completeness is a consequence of Proposition 2(3) because $\{(\equiv \text{M}), (\text{P})\}$ is contained in neither \mathcal{S}_p nor \mathcal{E}_p . \square

Lemma 48 $(\widehat{\text{MS}})$ and $(\widehat{\text{MF}})$ are NP-complete.

Proof. We show the result for $\widehat{(\text{MS})}$; the other case follows by applying Lemma 2. Consider the symmetric relation $r_1 = (r' \cap ((\text{MS}) \circ r'))^*$. It is easily verified that $(\equiv) \not\subseteq r_1$. Furthermore r_1 is non-empty since $(B) \subseteq (\text{MS}) \circ (B)$ when $B \in \{\text{P}, \text{O}, \text{D}\}$ and we know that $r' \not\subseteq (\equiv \text{F})$. Consequently, we can assume that $(\equiv) \not\subseteq r'$.

Case 1: $(\text{P}) \cap r' = \emptyset$ or $(\text{F}) \cap r' = \emptyset$. We note the following:

$$\begin{aligned} (\text{MS}) \circ (\text{P}) &= (\text{PMod}^{-1}\text{s}^{-1}\text{f}^{-1}) & (\text{MS}) \circ (\text{O}) &= (\text{PmODsF}) \\ (\text{MS}) \circ (\text{D}) &= (\text{PmODsF}) & (\text{MS}) \circ (\text{F}) &= (\text{pMODs}) \end{aligned}$$

It follows that $r_1 = ((\text{MS}) \circ r') \cap (\text{MS})$ equals either (ms) , (ms^{-1}) , (mS) , (Ms) , or (Ms^{-1}) . If $r_1 = (\text{Ms})$ or $r_1 = (\text{Ms}^{-1})$, then $r_1^* = (\text{M})$ and NP-completeness follows from Lemma 38(1). If $r_1 = (\text{ms})$ or $r_1 = (\text{ms}^{-1})$, then $((\text{m}^{-1}\text{s}^{-1}) \circ (\text{ms})) \cap (\text{ms}) = ((\text{m}^{-1}\text{s}) \circ (\text{ms}^{-1})) \cap (\text{m}^{-1}\text{s}) = (\text{s})$. Finally, if $r_1 = (\text{mS})$, then $(\text{S}) = r_1^*$ and

$$\begin{aligned} (\text{S}) \circ (\text{P}) &= (\text{Pmod}^{-1}\text{f}^{-1}) & (\text{S}) \circ (\text{O}) &= (\text{pmODF}) \\ (\text{S}) \circ (\text{D}) &= (\text{pmODF}) & (\text{S}) \circ (\text{F}) &= (\text{pmOD}) \end{aligned}$$

Hence, $((\text{S}) \circ r') \cap (\text{MS}) = (\text{m})$ and NP-completeness follows immediately.

Case 2: $(\text{PF}) \subseteq r'$. Consider the following instance Γ over the variables x, y, a, b :

$$x(\text{MS})y, y(\text{MS})a, a(\text{MS})x, b(\text{MS})x, b(\text{MS})y, br'a.$$

One can show $x(\text{S})y$ is derived from Γ so $((\text{S}) \circ r') \cap (\text{MS}) = (\text{m})$ and NP-completeness follows. \square

Lemma 49 $(\widehat{(\equiv \text{MS})})$ and $(\widehat{(\equiv \text{MF})})$ are NP-complete.

Proof. We show the result for $(\widehat{(\equiv \text{MS})})$; the other case follows by applying Lemma 2. It can be shown that $r_1 = ((\equiv \text{MS}) \circ r') \cap (\equiv \text{MS}) \in \{(\text{m}), (\text{ms}), (\text{Ms}), (\text{Ms}^{-1})\}$. All three cases lead to NP-completeness as shown in Lemma 48. \square

Lemma 50 $(\widehat{(\text{MSF})})$ is NP-complete.

Proof. It is easy to see that $B \subseteq (\text{MSF}) \circ B$ when $B \in \{(\text{P}), (\text{O}), (\text{D})\}$. Hence, $r' \subseteq (\text{MSF}) \circ r'$. By inspecting the composition table, one can see that if (\equiv) is a member of the composition of two symmetric relations, then these two relations cannot be disjoint. It follows that $(\equiv) \not\subseteq (\text{MSF}) \circ r'$ so we can obtain the relation $r' \setminus (\equiv) = r' \cap ((\text{MSF}) \circ r')$. We assume henceforth that $(\equiv) \not\subseteq r'$ and continue by noting that $(\text{MSF}) \circ (\text{P}) = (\text{PMod}^{-1}\text{s}^{-1}\text{f}^{-1})$ and $(\text{MSF}) \circ (\text{O}) = (\text{MSF}) \circ (\text{D}) = \text{neq}$ where $\text{neq} = \top \setminus (\equiv)$.

If $r' = (\text{P})$, then NP-completeness follows from Proposition 2(3). Otherwise, $(\text{MSF}) \circ r' = \text{neq}$ and we can show NP-completeness by a polynomial-time reduction from GRAPH 4-COLOURABILITY. Let $G = (V, E)$ be an arbitrary, undirected graph and construct a set of Allen constraints as follows:

1. introduce two variables x, y and the constraint $xr'y$;
2. for each $v \in V$, introduce a variable v and the constraints $v(\text{MSF})x$ and $v(\text{MSF})y$; and
3. for each $(v, w) \in E$, add the constraint $v\text{neq}w$.

It is a routine verification to see that the resulting instance is satisfiable iff G is 4-colourable. \square

Lemma 51 $(\widehat{(\equiv \text{MSF})})$ is NP-complete.

$$\left(\begin{array}{ccc} xr_1a_1 & a_7r_1a_1 & yr_1a_1 \\ a_1r_1a_2 & a_7r_2x & yr_1a_2 \\ xr_1a_2 & a_7r_2a_2 & yr_1a_3 \\ \\ a_3r_2x & a_8r_1x & yr_1a_4 \\ a_3r_2a_1 & a_8r_2a_1 & yr_2a_5 \\ a_3r_1a_2 & a_8r_2a_2 & yr_2a_6 \\ \\ a_4r_1x & a_9r_1a_1 & yr_1a_7 \\ a_4r_2a_1 & a_9r_2x & yr_1a_8 \\ a_4r_2a_2 & a_9r_2a_2 & yr_1a_9 \\ \\ a_5r_1x & a_7r_1a_8 \\ a_5r_1a_1 & a_8r_1a_9 \\ a_5r_1a_2 & a_7r_2a_9 \\ \\ a_6r_1x \\ a_6r_2a_1 \\ a_6r_2a_2 \\ \\ a_4r_1a_5 \\ a_5r_1a_6 \\ a_4r_2a_6 \end{array} \right)$$

Table 5: Definition of $C_{39}(r_1, r_2)$.

Proof. Note that $(\equiv \text{MSF}) \circ (\text{P}) = (\text{PMod}^{-1}\mathfrak{s}^{-1}\mathfrak{f}^{-1})$ and $(\equiv \text{MSF}) \circ (\text{O}) = (\equiv \text{MSF}) \circ (\text{D}) = \text{neq}$. If $r' = (\text{P})$, then NP-completeness follows from Proposition 2(3). Otherwise, $(\equiv \text{MSF}) \circ r' = \text{neq}$ so we can generate the relation (MSF) and NP-completeness follows from Lemma 50. \square

Lemma 52 *If $s \subseteq (\equiv \text{MSF})$, then \hat{s} is NP-complete.*

Proof. Combine Lemmas 38 and 47–51. \square

A.3.2 Remaining cases

This subsection considers the relations not covered in the previous section. We will henceforth assume that $r' \not\subseteq (\equiv \text{MSF})$ and we will only consider \hat{s} where $s \not\subseteq (\equiv \text{MSF})$. In the proofs, we will make use of a derivation $C_{39}(r_1, r_2)$ that denotes the relation (between x and y) derived from the set of constraints found in Table 5.

- $(\widehat{\text{PS}})$ and $(\widehat{\text{PF}})$: We give the proof for $(\widehat{\text{PS}})$; the other case follows by applying Lemma 2. Now, $(\text{PS}) \circ (B) = (\text{PMod}\mathfrak{s}\mathfrak{F})$ when $B \in \{\text{M}, \text{O}, \text{D}\}$ and $(\text{PS}) \circ (\text{F}) = (\text{PmODs})$. Thus, $((\text{PS}) \circ r') \cap (\text{PS})^* = (\text{P})$ and the result follows from Proposition 2.

- $\widehat{(\text{PSF})}$: Recall that $r' \neq (\text{M})$ and consider the following instance Γ on $\{x, y, a, b\}$:

$$\{x(\text{PSF})a, ar'y, x(\text{PSF})b, br'y, ar'b.\}$$

Let $r_1 = \top \setminus (\equiv s^{-1}f^{-1})$. One can show that xr_1y is derived from Γ which implies that $(\text{P}) = ((\text{PSF}) \cap r_1)^*$ can be obtained. NP-completeness follows from Proposition 2(3).

- \widehat{s} when $(\text{PM}) \subseteq s \subseteq (\text{PMSF})$: Follows from previous results by applying Lemma 42 with model transformation **shrink**.

We will from now assume that r' does not satisfy the inclusion $(\text{P}) \subseteq r' \subseteq (\text{PMSF})$; the previous results together with Proposition 2(3) allows us to do this without loss of generality.

- $\widehat{(\text{D})}$: By setting $R = (\text{d})$, $R_1 = r'$ and $R_2 = (\text{D})$ and applying Lemma 1, we have that $\{(\text{D}), r'\}$ is NP-complete.

- $\widehat{(\text{O})}$: Case 1. $(\text{PD}) \subseteq r' \subseteq (\text{PMDSF})$. The case $r' = (\text{PD})$ is considered in Lemma 43. Otherwise apply Lemma 43 and Lemma 42 with model transformation **ordshrink**.

Case 2. $(\text{D}) \subseteq r' \subseteq (\text{DSF})$. Apply Lemma 42 with model transformation **ordshrink** and then Lemma 1 with $R = (\text{d})$, $R_1 = (\text{O})$, $R_2 = (\text{D})$.

Case 3. $(\text{DM}) \subseteq r' \subseteq (\text{DMSF})$. Now, $(\text{D}) \subseteq C_{39}(r', (\text{O})) \subseteq (\text{PD})$ so we can obtain the relation (D) .

- $\widehat{(\text{PM})}$ and $\widehat{(\text{MO})}$: Using Lemma 42 with model transformations **shrink** and **expand**, the result follows from the earlier results.

- $\widehat{(\text{PD})}$: Obviously, $(\text{O}) \subseteq r' \subseteq \top \setminus (\text{PD})$. Apply Lemma 42 with model transformation **ordexpand** and then use Lemma 43.

- $\widehat{(\text{PO})}$: Case 1: $(\text{D}) \subseteq r' \subseteq (\text{DSF})$. Apply Lemma 42 with model transformation **ordshrink** and then Lemma 1 with $R = (\text{d})$, $R_1 = (\text{PO})$, $R_2 = (\text{D})$.

Case 2: $(\text{DM}) \subseteq r' \subseteq (\text{DMSF})$. In this case, $(\text{DSF}) \subseteq C_{39}(r', (\text{PO})) \subseteq (\text{PODSF})$ and we can obtain a relation r_1 such that $(\text{D}) \subseteq r_1 \subseteq (\text{DSF})$. Hence, NP-completeness follows from the previous case.

- $\widehat{(\text{OS})}$ and $\widehat{(\text{OF})}$: We prove the result for $\widehat{(\text{OS})}$; the other case follows by applying Lemma 2.

Case 1. $(\text{PD}) \subseteq r' \subseteq (\text{PMDF})$. Apply Lemma 42 thrice with model transformations **shrink**, **leftordshrink**, and **ordshrink**, consecutively, and then use Lemma 43.

Case 2. $(\text{D}) \subseteq r' \subseteq (\text{DMF})$. Now, $(\text{ODS}) \subseteq C_{39}(r', (\text{OS})) \subseteq (\text{PODS})$ so we can obtain the relation (D) .

- $\widehat{(\text{DS})}$ and $\widehat{(\text{DF})}$: We prove the result for $\widehat{(\text{DS})}$; the other case follows by applying Lemma 2.

Case 1. $(\text{PO}) \subseteq r' \subseteq (\text{PMOF})$. Apply Lemma 42 thrice with model transformations **shrink**, **leftordexpand**, and **ordexpand**, consecutively, and then use Lemma 43.

Case 2. $(\text{O}) \subseteq r' \subseteq (\text{OMF})$. Use Lemma 42 with model transformation **expand** and the earlier results.

- $\widehat{(\text{MD})}$: Case 1. $(\text{O}) \subseteq r' \subseteq (\text{OSF})$. In this case, $(\text{D}) \subseteq C_{39}((\text{MD}), r') \subseteq (\text{ODSF})$ and NP-completeness follows since we can obtain the relation (D) .

Case 2. $(\text{PO}) \subseteq r' \subseteq (\text{POSF})$. Now, $(\text{DSF}) \subseteq C_{39}((\text{MD}), r') \subseteq (\text{PODSF})$ so we can obtain the relation (D) .

Now, it remains to consider pairs r, s of disjoint relations such that neither r nor s contains (\equiv) and both of them contain exactly six basic relations. There are 10 such pairs. NP-completeness of $\{(\text{MSF}), (\text{POD})\}$ was proved in Lemma 50. NP-completeness of $\{(\text{PMS}), (\text{ODF})\}$, $\{(\text{PMF}), (\text{ODS})\}$, and $\{(\text{PSF}), (\text{MOD})\}$ was shown earlier in this subsection. Here we consider the remaining six pairs.

For the pairs $\{(PMO), (DSF)\}$, $\{(PMD), (OSF)\}$, and $\{(PDF), (MOS)\}$, apply Lemma 42 with model transformations *shrink*, *leftordshrink*, and *expand*, respectively, and then use earlier results. NP-completeness of $\{(PDS), (MOF)\}$ follows from that of $\{(PDF), (MOS)\}$ by using Lemma 2.

Consider $\{(POF), (MDS)\}$. It can be verified that $C_{39}((MDS), (POF)) = (PODSF)$, and we can obtain the relation (DS) and use earlier results. Finally, NP-completeness of $\{(POS), (MDF)\}$ follows from that of $\{(POF), (MDS)\}$ by using Lemma 2 once again.

References

- [1] J.F. Allen. Maintaining knowledge about temporal intervals. *Communications of the ACM*, 26(11):832–843, 1983.
- [2] J.F. Allen. Temporal reasoning and planning. In J.F. Allen, H. Kautz, R.N. Pelavin, and J. Tenenber, editors, *Reasoning about Plans*, chapter 1, pages 1–67. Morgan Kaufmann, 1991.
- [3] O. Angelsmark and P. Jonsson. Some observations on durations, scheduling and Allen’s algebra. In *Proceedings of the 6th Conference on Constraint Programming (CP’00)*, volume 1894 of *Lecture Notes in Computer Science*, pages 484–488. Springer-Verlag, 2000.
- [4] F.D. Anger and R.V. Rodriguez. Effective scheduling of tasks under weak temporal interval constraints. In B. Bouchon-Meunier, R. R. Yager, and L. A. Zadeh, editors, *Advances in Intelligent Computing - IPMU’94*, pages 584–594. Springer, Berlin, 1994.
- [5] A. Belfer and M.C. Golumbic. A combinatorial approach to temporal reasoning. In *Proceedings of 5th Jerusalem Conference on Information Technology*, pages 774–780. IEEE Computer Society Press, October 1990.
- [6] P. Bellini, R. Mattolini, and P. Nesi. Temporal logics for real-time system specification. *ACM Computing Surveys*, 32(1):12–42, 2000.
- [7] S. Benzer. On the topology of the genetic fine structure. In *Proceedings of the National Academy of Science, U.S.A.*, volume 45, pages 1607–1620, 1959.
- [8] A.A. Bulatov, A.A. Krokhin, and P. Jeavons. Constraint satisfaction problems and finite algebras. In *Proceedings of the 27th International Colloquium on Automata, Languages, and Programming (ICALP’00)*, volume 1853 of *Lecture Notes in Computer Science*, pages 272–282. Springer, 2000.
- [9] A.A. Bulatov, A.A. Krokhin, and P. Jeavons. The complexity of maximal constraint languages. In *Proceedings of the 33rd ACM Symposium on the Theory of Computing (STOC’01)*, pages 667–674, 2001.
- [10] D. Cohen, P. Jeavons, P. Jonsson, and M. Koubarakis. Building tractable disjunctive constraints. *Journal of the ACM*, 47(5):826–853, 2000.
- [11] C.H. Coombs and J.E.K. Smith. On the detection of structures in attitudes and developmental processes. *Psychological Review*, 80:337–351, 1973.
- [12] N. Creignou. A dichotomy theorem for maximum generalized satisfiability problems. *Journal of Computer and System Sciences*, 51(3):511–522, 1995.
- [13] T. Drakengren and P. Jonsson. Eight maximal tractable subclasses of Allen’s algebra with metric time. *Journal of Artificial Intelligence Research*, 7:25–45, 1997.
- [14] T. Drakengren and P. Jonsson. Twenty-one large tractable subclasses of Allen’s algebra. *Artificial Intelligence*, 93(1-2):297–319, 1997.

- [15] T. Drakengren and P. Jonsson. A complete classification of tractability in Allen's algebra relative to subsets of basic relations. *Artificial Intelligence*, 106(2):205–219, 1998.
- [16] M. Garey and D. Johnson. *Computers and Intractability: A Guide to the Theory of NP-Completeness*. Freeman, New York, 1979.
- [17] F. Gavril. Algorithms for a maximum clique and a maximum independent set of a circle graph. *Networks*, 3:261–273, 1973.
- [18] A. Gerevini and L. Schubert. Efficient temporal reasoning through timegraphs. In *Proceedings of International Joint Conference on Artificial Intelligence (IJCAI'93)*, pages 648–654. Morgan Kaufmann, 1993.
- [19] M.C. Golumbic, H. Kaplan, and R. Shamir. On the complexity of DNA physical mapping. *Advances in Applied Mathematics*, 15(3):251–261, 1994.
- [20] M.C. Golumbic, H. Kaplan, and R. Shamir. Graph sandwich problems. *Journal of Algorithms*, 19(3):449–473, 1995.
- [21] M.C. Golumbic and R. Shamir. Complexity and algorithms for reasoning about time: A graph-theoretic approach. *Journal of the ACM*, 40(5):1108–1133, 1993.
- [22] P. Hell and J. Nešetřil. On the complexity of H-coloring. *Journal of Combinatorial Theory, Ser. B*, 48:92–110, 1990.
- [23] R. Hirsch. Relation algebras of intervals. *Artificial Intelligence*, 83(2):1–29, 1996.
- [24] R. Hirsch. Expressive power and complexity in algebraic logic. *Journal of Logic and Computation*, 7(3):309–351, 1997.
- [25] P. Jonsson and C. Bäckström. A unifying approach to temporal constraint reasoning. *Artificial Intelligence*, 102(1):143–155, 1998.
- [26] P. Jonsson and T. Drakengren. A complete classification of tractability in RCC-5. *Journal of Artificial Intelligence Research*, 6:211–221, 1997.
- [27] P. Jonsson, T. Drakengren, and C. Bäckström. Computational complexity of relating time points with intervals. *Artificial Intelligence*, 109(1-2):273–295, 1999.
- [28] R.M. Karp. Mapping the genome: some combinatorial problems arising in molecular biology. In *Proceedings of the 25th ACM Symposium on the Theory of Computing (STOC'93)*, pages 278–285, New-York, 1993. ACM Press.
- [29] D.G. Kendall. Some problems and methods in statistical archaeology. *World Archaeology*, 1:68–76, 1969.
- [30] M. Koubarakis. Tractable disjunctions of linear constraints. In *Proceedings of the 2nd Conference on Principles and Practice of Constraint Programming (CP'96)*, pages 297–307, Boston, MA, 1996.
- [31] M. Koubarakis. The complexity of query evaluation in indefinite temporal constraint databases. *Theoretical Computer Science*, 171(1-2):25–60, 1997.
- [32] M. Koubarakis and S. Skiadopoulos. Querying temporal and spatial constraint networks in PTIME. *Artificial Intelligence*, 123(1-2):223–263, 2000.
- [33] P.B. Ladkin and R.D. Maddux. On binary constraint problems. *Journal of the ACM*, 41(3):435–469, 1994.
- [34] P.B. Ladkin and A. Reinefeld. Fast algebraic methods for interval constraint problems. *Annals of Mathematics and Artificial Intelligence*, 19(3-4):383–411, 1997.

- [35] G. Ligozat. A new proof of tractability for ORD-Horn relations. In *Proceedings of the 13th National (US) Conference on Artificial Intelligence (AAAI-96)*, pages 395–401, Menlo Park, CA, 1996. AAAI Press.
- [36] G. Ligozat. “Corner” relations in Allen’s algebra. *Constraints*, 3(2-3):165–177, 1998.
- [37] I. Meiri. Combining qualitative and quantitative constraints in temporal reasoning. *Artificial Intelligence*, 87(1-2):343–385, 1996.
- [38] B. Nebel. Archive of C-programs used for obtaining the results from [39], 1997. available from the author at <http://www.informatik.uni-freiburg.de/~nebel/journals.html>.
- [39] B. Nebel. Solving hard qualitative temporal reasoning problems: Evaluating the efficiency of using the ORD-Horn class. *Constraints*, 1(3):175–190, 1997.
- [40] B. Nebel and H.-J. Bürckert. Reasoning about temporal relations: A maximal tractable subclass of Allen’s interval algebra. *Journal of the ACM*, 42(1):43–66, 1995.
- [41] K. Nökel. *Temporally Distributed Symptoms in Technical Diagnosis*, volume 517 of *Lecture Notes in Artificial Intelligence*. Springer-Verlag, 1991.
- [42] J. Opatrný. Total ordering problem. *SIAM Journal on Computing*, 8(1):111–114, 1979.
- [43] C. Papadimitriou and M. Yannakakis. Scheduling interval ordered tasks. *SIAM Journal on Computing*, 8(3):405–409, 1979.
- [44] I. Pe’er and R. Shamir. Satisfiability problems on intervals and unit intervals. *Theoretical Computer Science*, 175(2):349–372, 1997.
- [45] T.J. Schaefer. The complexity of satisfiability problems. In *Proceedings of the 10th ACM Symposium on the Theory of Computing (STOC’78)*, pages 216–226, New-York, 1978. ACM Press.
- [46] E. Schwalb and L. Vila. Temporal constraints: A survey. *Constraints*, 3(2-3):129–149, 1998.
- [47] F. Song and R. Cohen. The interpretation of temporal relations in narrative. In *Proceedings of the 7th National (US) Conference on Artificial Intelligence*, pages 745–750, St. Paul, MN, USA, 1988. AAAI Press/The MIT Press.
- [48] K. Stergiou and M. Koubarakis. Backtracking algorithms for disjunctions of temporal constraints. *Artificial Intelligence*, 120(1):81–117, 2000.
- [49] Á. Szendrei. Maximal non-affine reducts of simple affine algebras. *Algebra Universalis*, 34(1):144–174, 1995.
- [50] A. Tarski. On the calculus of relations. *Journal of Symbolic Logic*, 6:73–89, 1941.
- [51] P.G. van Beek. Reasoning about qualitative temporal information. *Artificial Intelligence*, 58(1-3):297–326, 1992.
- [52] P.G. van Beek and R. Cohen. Exact and approximate reasoning about temporal relations. *Computational Intelligence*, 6(3):132–144, 1990.
- [53] P.G. van Beek and D.W. Manchak. The design and experimental analysis of algorithms for temporal reasoning. *Journal of Artificial Intelligence Research*, 4:1–18, 1996.
- [54] M.B. Vilain. A system for reasoning about time. In *Proceedings of the 2nd National (US) Conference on Artificial Intelligence*, pages 197–201, Pittsburgh, PA, USA, 1982. AAAI Press.

- [55] M.B. Vilain, H.A. Kautz, and P.G. van Beek. Constraint propagation algorithms for temporal reasoning: A revised report. In Daniel S. Weld and Johan de Kleer, editors, *Readings in Qualitative Reasoning about Physical Systems*, pages 373–381. Morgan Kaufmann, San Mateo, CA, 1989.
- [56] S.A. Ward and R.H. Halstead. *Computation Structures*. MIT Press, Cambridge, MA, 1990.
- [57] R.A. Wilson. The maximal subgroups of the Baby Monster. I. *Journal of Algebra*, 211(1):1–14, 1999.
- [58] X. Yang. A classification of maximal subsemigroups of finite order-preserving transformation semigroups. *Communications in Algebra*, 28(3):1503–1513, 2000.