

Recent advances in single-cell sequencing technologies

Lu Wen¹ and Fuchou Tang¹

Beijing Advanced Innovation Center for Genomics (ICG), Biomedical Pioneering Innovation Center (BIOPIIC), School of Life Sciences, Peking University, Beijing 100871, China

*Correspondence: Lu Wen, wenu@pku.edu.cn; Fuchou Tang, tangfuchou@pku.edu.cn

Abstract

Single-cell omics sequencing was first achieved for the transcriptome in 2009, which was followed by fast development of technologies for profiling the genome, DNA methylome, 3D genome architecture, chromatin accessibility, histone modifications, etc., in an individual cell. In this review we mainly focus on the recent progress in four topics in the single-cell omics field: single-cell epigenome sequencing, single-cell genome sequencing for lineage tracing, spatially resolved single-cell transcriptomics and third-generation sequencing platform-based single-cell omics sequencing. We also discuss the potential applications and future directions of these single-cell omics sequencing technologies for different biomedical systems, especially for the human stem cell field.

Keywords: single-cell omics, genome sequencing, epigenome sequencing, lineage tracing, third-generation sequencing, human stem cell

Introduction

A single cell zygote at the starting point of our life develops into 37 trillion cells in our body. The huge cellular heterogeneity and complexity represent a major bottleneck and pose great challenges for research on human development, stem cell biology and many other fields. In many organs of an adult individual, stem cells are continuously self-renewing and differentiating to maintain the lifelong physiological functions of the organs. Stem cells are generally rare in organs and are located at specific positions surrounded and controlled by well-organized niche cells. They are quite often heterogeneous, containing distinct subtypes or distinct biological states, such as quiescent and actively proliferating states, and there are also intermediate cell subpopulations during their multi-lineage differentiation processes. In recent years, single-cell omics sequencing technologies have been resolving many of these issues and revolutionizing the stem cell field. This starts from the first single cell RNA-seq (scRNA-seq) technique developed in 2009¹. With rapid development of the technology, including tremendous improvement of throughput, accuracy, automation, and commercialization, scRNA-seq techniques have been widely applied to address critical biological and medical questions. In this review, we will discuss recent advances in single-cell omics sequencing technologies in four topics. First, epigenetic regulation stands at the center of the gene regulatory networks, study of which provides important insights into how transcription is regulated and how epigenetic memory is established and maintained in stem cells as well as their differentiated progenies. The epigenome of a cell is all of the epigenetic information stored and maintained in a cell. It is comprised of a variety of precisely regulated and tightly interconnected epigenetic features including chromatin states, 3D genome architecture, DNA methylation, histone modifications, as well as the specific binding of transcription factors or non-coding RNAs onto the chromatin.

Single-cell epigenome sequencing techniques for many of these features have been established and have been routinely used for stem cell biology studies. Second, another layer of information crucial for understanding stem cell biology is the trajectory or lineage 'history' of a cell during stem cell self-renewal and differentiation. This information is vital to rigorously confirm the multi-lineage differentiation potentials of stem cells *in vivo*, which is one of the defining features of stem cells. Since genetic manipulation is in general prohibited for humans, endogenous genetic variants in the genome of a cell in human bodies, including single-nucleotide variants (SNVs), copy number variations (CNVs), Indels, structure variations (SVs), variations of microsatellites or other repetitive elements, as well as mitochondrial DNA mutations, provide invaluable information for tracing the developmental trajectory of stem cells in intact human tissues *in vivo*. The rapidly improving single-cell genome sequencing technologies are satisfying such lineage-tracing requirements for human stem cell studies. Third, stem cells and their descendants, as well as the surrounding niche cells, are usually well spatially organized in the tissues. So the spatial organization and interaction information is important for understanding stem cell biology *in vivo*. This issue can be addressed by spatially resolved single-cell transcriptome sequencing techniques. Fourth, third-generation sequencing (TGS) technologies have progressed rapidly in recent years. Combining TGS and single-cell omics technologies provides novel information for alternative splicing and other crucial biological features of an individual cell.

Single-cell epigenome sequencing

Single-cell epigenome sequencing technology is particularly challenging since the epigenetic information is scattered in the genome of a diploid cell, which has only two copies of the

Received: November 9, 2021. Revised: January 11, 2022. Accepted: January 13, 2022

© The Author(s) 2022. Published by Oxford University Press on behalf of the West China School of Medicine & West China Hospital of Sichuan University. This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact <email>journals.permissions@oup.com

genomic DNA. As a comparison, single-cell transcriptome analysis is somehow easier with many expressed genes in a diploid cell having over a dozen copies of the mRNAs. A highly sensitive enzyme or chemical reaction is essential for single-cell epigenome sequencing technology. To make the loss of DNA as minimal as possible, the experimental steps should be simple. Particularly if a barcode can be linked to a cell at an early step, hundreds to thousands of individual cells can be pooled together for subsequent operations, which will greatly increase the throughput of the method. Droplet- and microchip-based methods greatly facilitate automation and a combinatorial indexing strategy can increase the throughput in an exponential manner. It is noteworthy that Tn5 transposase shows excellent performance in sensitivity, simplicity, early barcoding and throughput, and thus has stood at the center of technological improvement of single-cell epigenome sequencing in recent years. Bioinformatics tools have also been developed to tackle the challenges of the sparse nature of single-cell epigenomic data. For example, chromVAR measures motifs or functional annotations together instead of individual open chromatin regions, while ArchR uses an iterative dimensionality reduction approach^{2,3}.

Chromatin states

Chromatin states represent the active or repressive status of the regulatory genomic regions in a cell. Practically, the active chromatin states can be assessed by accessibility of an enzyme such as transposase, DNase I, micrococcal nuclease (MNase), or GpC methyltransferase, by which single-cell chromatin accessibility sequencing techniques have been established⁴⁻⁹. Among these strategies, the Tn5 transposon-based ATAC-seq (Assay for Targeting Accessible-Chromatin with high-throughout sequencing) method simultaneously inserts, fragments and adds adaptor tags to the active chromatin regions in a cell and thus is excellent for low-cost and high-throughput analysis¹⁰. Using plate-, droplet-, or combinatorial indexing-based methods, recent studies have shown that thousands to hundreds of thousands of individual cells can be analyzed in a single sample¹¹⁻¹⁵. A major shortcoming of ATAC-seq is that it cannot directly detect the repressive chromatin states that can be reliably detected by the GpC methyltransferases-based methods. The latter can also simultaneously analyze both chromatin states and endogenous DNA methylation of a cell, and have relatively higher resolution, as a GpC dinucleotide (GCH) occurs approximately every 25 bp in the human genome. Both the GpC methyltransferases- and transposon-based methods have also been shown to be able to simultaneously analyze CNVs at megabase resolution^{16,14}.

3D genome architecture

While the chromatin state provides information on where the chromatin opens, 3D genome architecture analysis provides information on how a genome is spatially and structurally organized and compartmentalized, as well as how different genomic regions interact with each other in a cell. Since the first single-cell Hi-C technique was established, which detects ~1 000 contacts in an individual cell, the methodology has been continuously improved, and the latest techniques are able to detect >1 million contacts in an individual diploid cell¹⁷⁻¹⁹. The technological improvement includes omitting the biotin enrichment step, usage of highly efficient single-cell whole genome amplification methods such as multiplex end-tagging amplification (META) and multiple displacement amplification (MDA), as well as higher sequencing depth. This leads to a resolution of ~20 kb for analyzing the 3D

structures in an individual cell, which is capable of distinguishing two parental genomes of a cell and different neuronal subtypes^{20,19}. On the other hand, the throughput of the single-cell Hi-C technique has been increased by using combinatorial indexing and Tn5 transposase/plate-based strategies^{21,22}.

Histone modifications and transcription factor bindings

Histone modifications contribute greatly to the organization of chromatin structures and regulation of gene transcription. Chromatin immunoprecipitation (ChIP) is a widely used method for detecting modifications of histones in nucleosomes of chromatin. Realizing that low sensitivity and specificity of antibody capture is the main obstacle in single-cell ChIP-seq analysis, cell-specific barcodes are added before aggregating the cells for immunoprecipitation²³⁻²⁵. Among these methods, Drop-ChIP and scChIP-seq add cell barcodes by MNase digestion and ligation with a droplet microfluidics workflow, while itChIP adds cell barcodes by Tn5 transposase tagmentation with a chromatin opening step. These methods are able to detect ~1 000 to ~10 000 unique reads per cell.

Enzyme-tethering represents a non-immunoprecipitation chromatin profiling approach that is becoming increasingly popular and has been adapted to single-cell analysis²⁶⁻³¹. In these techniques, Tn5 transposase, MNase, or adenine methyltransferase is tethered to protein A that binds to the antibody, directly to the antibody, or directly to the target chromatin protein, which allows marking of the genomic regions with specific histone marks. ChIC, CUN&RUN, and scChIC use MNase, while scCUT&Tag, COBATCH, ACT-seq, and ChIL-seq use Tn5 transposase. A key cation-activation step, Ca²⁺ for MNase and Mg²⁺ for Tn5 transposase, allows activation of the enzyme activity in a short time window after washing off the nonspecifically-bound enzyme and drastically increases the signal-to-noise ratio. As the Tn5-based method simultaneously adds tagged sequences, it is more convenient for high-throughput single-cell epigenomics analysis. The current enzyme-tethering single-cell techniques are able to detect several thousand unique reads per individual cell.

DNA methylation

DNA methylation comprises another critical epigenetic layer showing cell-type-specific patterns. Single-cell DNA methylome sequencing techniques have been established using various strategies including the reduced representation bisulfite sequencing (RRBS)- and the post-bisulfite adaptor tagging (PBAT)-based methods³²⁻³⁴. The mapping efficiency and throughput of PBAT-based methods can be increased by using 3' tagging techniques such as adaptase and TdT tailing; yet since only one round of random amplification is used, the coverage is decreased^{35,36}. A sci-MET method has applied a combinatorial indexing strategy for increasing the throughput, with the first and second rounds of barcodes being incorporated by Tn5 transposon and random priming, respectively³⁷. Conventional RRBS enriches CpG-containing regions by selecting genomic regions between a pair of MspI (CCGG) sites, but also covers the whole genome with many randomly fragmented genomic fragments. Recently, a single-cell extended-representation bisulfite sequencing (scXRBS) method uses an alternative approach by ligating the adaptor to a single MspI site, and thus achieves a new balance between coverage and enrichment of functionally relevant genomic regions³⁸.

Table 1. Single-cell genome sequencing techniques.

	Uniformity	Accuracy	Coverage	Operation	Reference
DOP-PCR	+++	+	+	+++	46
MDA	++	+++	+++	+++	47
eMDA	+++	+++	+++	++	50
SISSOR	+++	++++	++	++	49
MALBAC	+++	++	++	+++	52
LIANTI	++++	+++	++++	++	48
META-CS	+++	++++	+++	+++	51

Joint analysis of chromatin states and transcriptome

Methods for joint analysis of multiple omics in the same individual cell have been achieved by physical separation of different omics molecules, parallel indexing, or parallel capturing³⁹. In recent years, several methods have been reported for jointly detecting chromatin accessibility and transcriptome in an individual cell, which use various strategies for barcoding and separating two types of information^{40–44}. sci-CAR uses well-specific barcodes to perform *in situ* reverse transcription and Tn5 transposition separately⁴⁰. The scCAT-seq method physically separates the nucleus and cytoplasmic RNA by centrifugation⁴². Both Paired-seq and SHARE-seq use a combinatorial indexing strategy, in which two to three rounds of barcoding are performed, adding barcodes to the 5' ends of both the Tn5 and RT primers by split-and-pool. To separate the chromatin accessibility and transcriptome libraries, Paired-seq uses a restriction enzyme strategy, while SHARE-seq uses a biotin affinity pull-down strategy^{43,44}. SNARE-seq is a droplet-based method; it uses a barcoded oligo(dT)-bearing splint oligonucleotide for simultaneously performing a reverse transcription reaction for capturing the transcriptome and a ligation reaction for capturing the chromatin accessibility information⁴¹.

Single-cell lineage tracing

Single-cell lineage tracing techniques have recently been developed by using a combination of transposons or CRISPR/Cas9 genome editing and single-cell transcriptome sequencing⁴⁵. However, these genetic manipulation-based methods are not suitable for *in vivo* study of humans. Genome sequence information is continuously changing during development of the human from zygote to adult and further ageing processes due to stochastic genetic mutations. So the genetic mutations are intrinsic and ideal 'markers' for lineage tracing of a cell in the human body. In fact, they have been widely used for lineage tracing of tumor cells at bulk levels for many years and are the basis for tumorigenesis studies. Recent studies have reported the use of these endogenous changes of genome and mitochondrial DNA (mtDNA) information for lineage tracing of human stem cells in development and ageing, using clones derived from single cells or single-cell genome sequencing technologies.

Single-cell genome sequencing

Single cell whole genome amplification (scWGA) techniques such as degenerated oligonucleotide primer (DOP)-PCR⁴⁶ and MDA⁴⁷ have long been reported. More recently, several new methods, including MALBAC, eMDA, LIANTI, SISSOR, and META-CS, have been developed^{48–52}. The general characters of these methods are shown in Table 1. MDA uses Phi29 for isothermal single-cell whole genome amplification⁴⁷. As Phi29 is of high fidelity with about one

nucleotide per 10⁸ error rate, MDA has relatively high accuracy for calling SNVs, and has been applied to single-cell genome lineage tracing⁵³. One disadvantage of MDA is its exponential-like manner of amplification of genomic DNA, which results in amplification of initial extension errors and dampens coverage uniformity. LIANTI uses T4 RNA polymerase to linearly amplify the original template hundreds of times, which increases uniformity and accuracy⁴⁸. Further, META-CS and SISSOR acquire information from both strands (Watson and Crick strands, or duplex) for reciprocal corrections, which gives even higher accuracy for calling SNVs^{49,51}. The duplex information can also be recalled in MDA by using single-nucleotide polymorphism (SNP) information, though the number of informative SNVs is reduced⁵⁴. Further investigations are needed to reach the extremely challenging requirement of single-cell genome sequencing-based lineage tracing.

Uniformity is measured by the coefficient of variation of the sequence-dependent bias along the genome. At a bin size of 1 000 kb, LIANTI shows the highest uniformity with a value of ~0.03, while eMDA, MALBAC (normalized), META-CS, and DOP-PCR show a range between 0.1 and 0.15, and MDA shows a value >0.21; the value of SISSOR has not been described and is expected to be similar to eMDA.

Accuracy is measured by the false positive rate (FPR). META-CS and SISSOR, which give strand-specific information, have the highest accuracy with the lowest FPR (<2.4 × 10⁻⁸). LIANTI measures the linear amplification product many times and has the second highest accuracy with a FPR of 5.4 × 10⁻⁶. MDA and eMDA rank third with a FPR of 1.3 × 10⁻⁴. MALBAC and DOP-PCR show FPRs of 3.8 × 10⁻⁴ and 9.6 × 10⁻⁴, respectively.

Coverage of LIANTI is the highest, covering 95% of the genome of a human diploid cell by sequencing 83 Gb data, while META-CS (64% by 18 Gb data), MDA (87% by 85 Gb data), and eMDA (72% by 30 Gb data) rank second, followed by MALBAC (Yikon, 73% by 94 Gb data), SISSOR (64% by 195 Gb data), and DOP-PCR (45% by 84 Gb data).

Experimental operations of DOP-PCR, MDA, MALBAC, and META-CS are easier than those of eMDA, SISSOR, and LIANTI.

Genome sequence information in a cell for lineage tracing

Different types of genetic variants have different characters for the purpose of lineage tracing. Nucleotide substitution changes are estimated to occur at a frequency of ~1 per cell division in a human cell. As expected, the number of SNVs increases with accumulating cell divisions during development and ageing. A fetal hematopoietic stem cell has tens of somatic SNVs while a hematopoietic stem cell from a middle-aged adult has ~1 000 somatic SNVs^{55,56}. Similarly, adult stem cells of colon, small intestine, and liver have a few thousands somatic SNVs, with an accumulation of ~36 mutations per year⁵⁷. Thus SNVs represent a rich source of endogenous genetic polymorphisms for lineage tracing.

Microsatellites are expected to be an even richer source of endogenous polymorphism as the mutation rate of a microsatellite is as high as 10^{-3} to 10^{-5} /locus/cell division. It is estimated that 50 microsatellite mutations occur per cell division in humans and the complete cell lineage tree can be reconstructed using this information⁵⁸. Obstacles for full application of this information include error-prone sequencing, difficulty in capturing, and short read length.

The repetitive elements including long interspersed repeat elements (LINEs), short interspersed nuclear elements (SINEs), and long terminal repeats (LTRs) comprise a large portion of the human genome. A number of LINEs such as LINE1 and SINEs such as Alus are still active in humans, and transposition of LINE1 and Alus occurs at a rate of about 1 per 140 generation and 15 per generation, respectively⁵⁹. Low-rate somatic mutations of LINE1 have been reported in individual neurons at rates ranging from <0.1 to >10 insertions per cell^{60,61}. Also, insertions have been detected in normal gastrointestinal tissues and occur very early during the development of gastrointestinal tumors⁶². Though the frequency is low, the unique insertion sequences of new transposon elements can be definitely verified and detected, which facilitates using them as good markers for lineage tracing⁶³.

It is well known that CNVs occur in high frequency during human early development, although most of them lead to death of the embryo or are self-corrected by elimination of the aneuploid cells in chimeric embryos; how frequently the early CNVs result in human chimeras is not known. In recent years, somatic chimeric CNVs have been identified in many cell types including neurons, blood cells, and fibroblasts^{64–69}. The advantage of CNVs as an endogenous marker for lineage tracing is that they can be detected by relatively low-depth single-cell genome sequencing. However, the disadvantages include low frequency and difficulty of verification.

mtDNA, which is a ~16 kb long circular genome, is also a part of the cellular genome in addition to the nuclear genome. mtDNA mutation occurs 10–100 times more than genomic mutation, and there are hundreds of copies of mtDNAs in an individual cell. Also, mtDNA can be detected in other single-cell omics data, including scATAC-seq, scRNA-seq, and single-cell genome sequencing. These features make mtDNA a unique marker for simultaneous lineage tracing and cell-state detection⁷⁰.

Several recent studies have reported lineage tracing using single cells, clones derived from single cells, or microdissection^{71–73,63,55,53,74,56}. Lee-Six *et al.* analyzed 140 colonies derived from single hematopoietic stem cells of a 59-year-old male individual and identified an average of 1 023 SNVs and 20 small insertion/deletions in each clone. They constructed a phylogenetic tree that revealed clonal relationships among these 140 colonies. The results showed a rapid population expansion of hematopoietic stem cells during early life and estimated that there are between 50 000 and 200 000 hematopoietic stem cells actively making white blood cells at any specific time⁵⁵. In another study, Spencer Chapman *et al.* analyzed 511 colonies derived from hematopoietic stem cells of two human fetuses and identified 25.5 and 41.9 SNVs per clone in the 8 and 18 post-conception week fetuses, respectively⁵⁶. These SNVs allowed for reconstructing phylogenetic trees for early human embryonic development, which revealed several interesting findings, including an unequal contribution of each of the two-cell stage blastomeres to the blood compartment, a higher mutation rate in the first three cell divisions, and hypoblast origin of the extra-embryonic mesoderm and primitive blood.

Cells like neurons are post-mitotic and not able to proliferate, limiting the use of clone amplification and bulk sequencing strategies. Evrony *et al.* analyzed 16 cerebral cortex neurons by MDA and whole genome sequencing, which allowed for identification of two new L1 retrotranspositions and one poly-A microsatellite mutation. They designed a custom droplet digital PCR (ddPCR) assay and analyzed these mutations in various brain regions, which revealed one clone limiting to the left middle frontal gyrus and another distributing over the entire left hemisphere⁶³. In another study by the same group, 36 neurons were analyzed for SNVs and custom ddPCR was used for revealing the polyclonal architecture of the human cerebral cortex⁵³.

The field of lineage tracing at single-cell resolution is developing rapidly. The approach can be used to reconstruct the developmental history of stem cells, showing their precursors and progenies. It can also be used for estimating the number of stem cells that give rise to the differentiated cells, and whether or not they give equivalent contributions to these progeny cells. For example, Spencer Chapman *et al.* showed that two blastomeres in a two-cell stage embryo contribute unequally to the body⁵⁶. Also, during tissue injury, it is important to know which types of cells contribute to repair the injured tissue and recover its physiological functions, as has been studied in mice by genetic lineage tracing. Single-cell multiple-omics sequencing for genome and epigenome or transcriptome may help in elucidating the situation in humans.

Single-cell spatial transcriptome

Spatial localization is essential for determining cellular fate. Single cell spatially resolved transcriptome technologies have been developed and improved rapidly in recent years, and are particularly useful for human study for which genetic labeling techniques are not applicable⁷⁵. The methods of spatially resolved transcriptomics are listed in Table 2. Among them, two types of technologies, single-molecule FISH (smFISH)-based ones and *in situ* sequencing-based ones, provide single-cell resolution. The first one, including seqFISH and MERFISH, uses combinatorial barcodes for smFISH^{76,77}. seqFISH labels each RNA by a combinatorial set of colored probes, through multiple rounds of sequential hybridizations and clearance of the probes⁷⁷. MERFISH uses a similar combinatorial labeling strategy with the use of readout probes instead of direct labeling probes⁷⁶. The latest versions of both methods are able to image mRNAs for up to 10 000 genes in a single cell^{78,79}.

The second set of methods is *in situ* sequencing (ISS, FISSEQ, STARmap, ExSeq)^{80,83}. All these methods use rolling cycle amplification for signal amplification before *in situ* sequencing analysis. ISS and STARmap are targeting methods. ISS uses padlock probes for targeting⁸⁰, and STARmap uses designed nucleic acids for directly targeting RNAs that bypass the reverse transcription step and increases detection efficiency⁸³. FISSEQ and ExSeq are untargeted methods, and ExSeq is an improved technique of FISSEQ that adapts the chemistry of expansion microscopy to allow high spatial resolution mapping of RNAs^{82,81}.

Another set of spatial transcriptomics techniques use *in situ* capturing strategies^{86,85}. These techniques are approaching single-cell resolution with the high density of on-slide capturing^{87,88}.

Table 2. Spatially resolved transcriptome techniques.

	Full name	Strategy	Targeting	Single-cell resolution	References
seqFISH	Sequential fluorescence in situ hybridization (FISH)	Combinatorial barcodes for single-molecule FISH	Yes	Yes	78,77
MERFISH	Multiplexed error-robust (FISH)	Combinatorial barcodes for single-molecule FISH	Yes	Yes	76,79
ISS	<i>In situ</i> sequencing	<i>In situ</i> sequencing	Yes	Yes	80
FISSEQ	Fluorescence in situ sequencing	<i>In situ</i> sequencing	No	Yes	81
ExSeq	Expansion sequencing	<i>In situ</i> sequencing with expansion microscopy	No	Yes	82
STARmap	Spatially-resolved transcript amplicon readout mapping	<i>In situ</i> sequencing	Yes	Yes	83
TIVA	Transcriptome <i>in vivo</i> analysis	Photoactive tag	Yes	Yes	84
Spatial Transcriptomics	Spatial transcriptomics	Gene ChIPs with immobilized reverse-transcription oligo (dT) primers	No	No	85
Slide-seq	Slide-seq	Spatially resolved DNA-barcoded beads	No	Nearly	86,87
Stereo-seq	Spatial enhanced resolution omics-sequencing	Spatially resolved DNA nanoball	No	Yes/nearly	88
iTranscriptome	<i>In silico</i> spatial transcriptome	Combination of the low-input RNA sequencing with serial cryosection and laser capture microdissection	No	No	89
Tomo-Seq	RNA tomography sequencing	Combination of the low-input RNA sequencing with serial cryosection	No	No	90

Third-generation sequencing

Third-generation/real-time single molecule sequencing (TGS) methods have been developing especially fast in recent years. These include Nanopore sequencing (ONT) introduced by Oxford Nanopore Technologies, and single-molecule real-time (SMRT) sequencing by Pacific Biosciences (PacBio)⁹¹. Emerging as a new field of single-cell omics sequencing, TGS has several unique advantages and applications, some of which have been achieved (Fig. 1).

First, TGS-based scRNA-seq techniques are powerful for detecting alternative splicing or DNA rearrangements by directly sequencing the full-length intact cDNAs. Several studies including ours have recently developed TGS-based scRNA sequencing techniques including SCAN-seq, R2C2, ScISOr-Seq, ScNaUmi-seq, and RAGE-seq^{93–96}. SCAN-seq is able to detect >8 000 genes in an individual mouse embryonic stem cell (mESC), exhibiting a similar sensitivity to the next-generation sequencing (NGS) platform-based scRNA-seq techniques such as SMART-seq2, and SUP-seq. A large number of unannotated novel transcripts have been detected. SCAN-seq detected 6 487 unannotated transcripts corresponding to 3 834 genes in mESCs, and 27 250 unannotated transcripts corresponding to 9 338 genes in mouse preimplantation embryos⁹³. ScISOr-Seq detected 18 173 known and 16 872 novel isoforms in mouse cerebellum⁹⁴. In addition, RAGE-seq has shown the ability of the TGS-based approach for detecting fusion

transcripts from somatic DNA rearrangements of T-cell-receptor (TCR) and B-cell-receptor (BCR) transcripts⁹⁵.

Second, while small variants such as SNVs and short indels can be accurately detected using NGS-based short reads, larger structural variations (SVs) are more challenging to detect and characterize. TGS-based methods have been developed rapidly to increase the reliability and resolution of SV detection⁹⁷. Our group has recently developed a TGS-based single-cell genome sequencing technique SMOOTH-seq⁹⁸. For individual cells, the technique gets long sequencing reads with an average length of 6 kb, and reaches 19% genome coverage by 0.4X sequencing depth. Except for insertions, deletions, duplications, and translocations, the technique also effectively detects extra-chromosomal circular DNAs (ecDNA), being able to cover the full-length ones of <10 kb in a single read.

Third, single molecular nanopore sequencing is able to directly detect epigenetic modifications such as 5mC and 6mA. Combining such power with the enzyme accessibility of N6-methyladenosine (m6A) methyltransferase or GpC methyltransferase, three groups have recently reported TGS-based enzyme accessibility techniques for detecting chromatin state along single-molecules over a long distance (SMAC-seq, Fiber-seq, and nanoNOME)^{99–101}. Although they have not been adapted to single-cell analysis, these methods display the value of TGS for investigating coordination between the states of neighboring regulatory elements over large genomic regions, which opens a new avenue for future study.




Figure 1. TGS-based single cell sequencing technologies. Red and green indicate the aspects that have been achieved, while the red indicates those where TGS-based methods have advantages over NGS-based methods. *The Nanopore-based technology may be used for directly sequencing the protein in the future⁹². Note the differences between NGS-based and TGS-based single-cell omics sequencing technologies (compare this figure to Fig. 1 of Ref. 39)

Prospective

Single-cell omics sequencing technology has already made fruitful progresses in the stem cell biology field. However, the current techniques are still not ideal for human *in vivo* stem cell studies. We expect that the techniques will further be developed and improved within the next few years. The current single-cell omics sequencing technology has both strong (developed) and weak (developing) characteristics (Fig. 2). The technology has high universality as it is applicable in a wide range of biological research fields from plants to medicine. It gives high sensitivity and accuracy, but still requires improvement. The throughput, automation, and speed of the technology have increased greatly with the cost decreasing, but have not met clinical requirements. Particularly for stem cell studies, the temporal and spatial resolution of the technology is not satisfactory and is being improved. With its weaknesses being improved, single-cell omics sequencing technology will be routinely used to dissect the biology of stem cells, including (1) their self-renewal abilities or multiple lineage differentiation potentials under physiological conditions or under pathological conditions, (2) their premature differentiation or delayed leaving from self-renewal mode in diseased situations, (3) their microenvironments, (4) the consequences of their genetic perturbations, and (5) short-term responses to environmental changes or long-term maintenance of their fates.

Further, it will be ideal if genome, epigenome, and transcriptome can be simultaneously analyzed for an individual cell. The transcriptome will permit identifying and separating different types of cells. It will also act as a functional readout of the global transcriptional activity. Then how different layers of epigenomes regulate the organization of the genome and transcriptional activity of every gene can be delineated. Finally, the genome information can also be used to construct the lineage relationship. In addition, how genetic changes contribute to the abnormal behavior of a stem cell can be analyzed. If a genetic change perturbs the expression of a gene, it may change the physiological function and phenotype of a cell. In the future it is possible that through single-cell multi-omics sequencing we will identify genetic changes in the stem cells in our body and their potential connections to the phenotypic changes of a stem cell.

Of course, single-cell omics sequencing techniques are just a series of technologies and they cannot answer every question that arises in stem cell biology. Nevertheless, by integrating properly with other sets of powerful technologies such as gene editing tools and organoid 3D culture systems and biological concepts, they will definitely accelerate the transformation of our rich and deep knowledge of stem cells in animal models into more clinically relevant knowledge of stem cells in human.




Figure 2. Characteristics of the current single-cell omics sequencing technology. Single-cell omics sequencing technology is shown shaped like a barrel with both long (developed) and short (developing) boards (left panel), and a radar chart shows the current technical states of nine major characteristics, with the more developed state of the character indicated by its being positioned more peripherally (right panel).

Acknowledgements

This work was supported by the National Key Research and Development Program of China (Grant No. 2018YFA0107601).

Conflict of interest

None declared. In addition, as an Editorial Board Member of *Precision Clinical Medicine*, the corresponding author Fuchou Tang was blinded from reviewing and making decisions on this manuscript.

References

- Tang F, Barbacioru C, Wang Y, et al. mRNA-Seq whole-transcriptome analysis of a single cell. *Nat Methods* 2009;**6**:377–82. doi: 10.1038/nmeth.1315.
- Granja JM, Corces MR, Pierce SE, et al. ArchR is a scalable software package for integrative single-cell chromatin accessibility analysis. *Nat Genet* 2021;**53**:403–11. doi: 10.1038/s41588-021-00790-6.
- Schep AN, Wu B, Buenrostro JD, et al. chromVAR: inferring transcription-factor-associated accessibility from single-cell epigenomic data. *Nat Methods* 2017;**14**:975–8. doi: 10.1038/nmeth.4401.
- Buenrostro JD, Wu B, Litzenburger UM, et al. Single-cell chromatin accessibility reveals principles of regulatory variation. *Nature* 2015;**523**:486–90. doi: 10.1038/nature14590.
- Clark SJ, Argelaguet R, Kapourani CA, et al. scNMT-seq enables joint profiling of chromatin accessibility DNA methylation and transcription in single cells. *Nat Commun* 2018;**9**:781. doi: 10.1038/s41467-018-03149-4.
- Cusanovich DA, Daza R, Adey A, et al. Epigenetics. Multiplex single-cell profiling of chromatin accessibility by combinatorial cellular indexing. *Science* 2015;**348**:910–4. doi: 10.1126/science.aab1601.
- Guo F, Li L, Li J, et al. Single-cell multi-omics sequencing of mouse early embryos and embryonic stem cells. *Cell Res* 2017;**27**:967–88. doi: 10.1038/cr.2017.82.
- Jin W, Tang Q, Wan M, et al. Genome-wide detection of DNase I hypersensitive sites in single cells and FFPE tissue samples. *Nature* 2015;**528**:142–6. doi: 10.1038/nature15740.
- Pott S. Simultaneous measurement of chromatin accessibility, DNA methylation, and nucleosome phasing in single cells. *Elife* 2017;**6**:e23203. doi: 10.7554/eLife.23203.
- Buenrostro JD, Giresi PG, Zaba LC, et al. Transposition of native chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-binding proteins and nucleosome position. *Nat Methods* 2013;**10**:1213–8. doi: 10.1038/nmeth.2688.
- Cusanovich DA, Hill AJ, Aghamirzaie D, et al. A Single-Cell atlas of in vivo mammalian chromatin accessibility. *Cell* 2018;**174**:1309–24. e18. doi: 10.1016/j.cell.2018.06.052.
- Lake BB, Chen S, Sos BC, et al. Integrative single-cell analysis of transcriptional and epigenetic states in the human adult brain. *Nat Biotechnol* 2018;**36**:70–80. doi: 10.1038/nbt.4038.
- Preissl S, Fang R, Huang H, et al. Single-nucleus analysis of accessible chromatin in developing mouse forebrain reveals cell-type-specific transcriptional regulation. *Nat Neurosci* 2018;**21**:432–9. doi: 10.1038/s41593-018-0079-3.
- Satpathy AT, Granja JM, Yost KE, et al. Massively parallel single-cell chromatin landscapes of human immune cell development and intratumoral T cell exhaustion. *Nat Biotechnol* 2019;**37**:925–36. doi: 10.1038/s41587-019-0206-z.
- Xu W, Wen Y, Liang Y, et al. A plate-based single-cell ATAC-seq workflow for fast and robust profiling of chromatin accessibility. *Nat Protoc* 2021;**16**:4084–107. doi: 10.1038/s41596-021-00583-5.
- Bian S, Hou Y, Zhou X, et al. Single-cell multiomics sequencing and analyses of human colorectal cancer. *Science* 2018;**362**:1060–3. doi: 10.1126/science.aao3791.
- Flyamer IM, Gassler J, Imakaev M, et al. Single-nucleus Hi-C reveals unique chromatin reorganization at oocyte-to-zygote transition. *Nature* 2017;**544**:110–4. doi: 10.1038/nature21711.
- Nagano T, Lubling Y, Stevens TJ, et al. Single-cell Hi-C reveals cell-to-cell variability in chromosome structure. *Nature* 2013;**502**:59–64. doi: 10.1038/nature12593.
- Tan L, Xing D, Chang CH, et al. Three-dimensional genome structures of single diploid human cells. *Science* 2018;**361**:924–8. doi: 10.1126/science.aat5641.
- Tan L, Ma W, Wu H, et al. Changes in genome architecture and transcriptional dynamics progress independently of sensory experience during post-natal brain development. *Cell* 2021;**184**:741–58. e717. doi: 10.1016/j.cell.2020.12.032.

- 21 Nagano T, Lubling Y, Várnai C, et al. Cell-cycle dynamics of chromosomal organization at single-cell resolution. *Nature* 2017;**547**:61–7. doi: 10.1038/nature23001.
- 22 Ramani V, Deng X, Qiu R, et al. Massively multiplex single-cell Hi-C. *Nat Methods* 2017;**14**:263–6. doi: 10.1038/nmeth.4155.
- 23 Ai S, Xiong H, Li CC, et al. Profiling chromatin states using single-cell itChIP-seq. *Nat Cell Biol* 2019;**21**:1164–72. doi: 10.1038/s41556-019-0383-5.
- 24 Gosselin K, Durand A, Marsolier J, et al. High-throughput single-cell ChIP-seq identifies heterogeneity of chromatin states in breast cancer. *Nat Genet* 2019;**51**:1060–6. doi: 10.1038/s41588-019-0424-9.
- 25 Rotem A, Ram O, Shores N, et al. Single-cell ChIP-seq reveals cell subpopulations defined by chromatin state. *Nat Biotechnol* 2015;**33**:1165–72. doi: 10.1038/nbt.3383.
- 26 Carter B, Ku WL, Kang JY, et al. Mapping histone modifications in low cell number and single cells using antibody-guided chromatin tagmentation (ACT-seq). *Nat Commun* 2019;**10**:3747. doi: 10.1038/s41467-019-11559-1.
- 27 Harada A, Maehara K, Handa T, et al. A chromatin integration labelling method enables epigenomic profiling with lower input. *Nat Cell Biol* 2019;**21**:287–96. doi: 10.1038/s41556-018-0248-3.
- 28 Kaya-Okur HS, Wu SJ, Codomo CA, et al. CUT&Tag for efficient epigenomic profiling of small samples and single cells. *Nat Commun* 2019;**10**:1930. doi: 10.1038/s41467-019-09982-5.
- 29 Schmid M, Durussel T, Laemmli UK. ChIC and ChEC; genomic mapping of chromatin proteins. *Mol Cell* 2004;**16**:147–57. doi: 10.1016/j.molcel.2004.09.007.
- 30 Skene PJ, Henikoff S. An efficient targeted nuclease strategy for high-resolution mapping of DNA binding sites. *Elife* 2017;**6**:e21856. doi: 10.7554/eLife.21856.
- 31 Wang Q, Xiong H, Ai S, et al. CoBATCH for High-Throughput Single-Cell Epigenomic Profiling. *Mol Cell* 2019;**76**:206–16. e7. doi: 10.1016/j.molcel.2019.07.015.
- 32 Guo H, Zhu P, Wu X, et al. Single-cell methylome landscapes of mouse embryonic stem cells and early embryos analyzed using reduced representation bisulfite sequencing. *Genome Res* 2013;**23**:2126–35. doi: 10.1101/gr.161679.113.
- 33 Luo C, Keown CL, Kurihara L, et al. Single-cell methylomes identify neuronal subtypes and regulatory elements in mammalian cortex. *Science* 2017;**357**:600–4. doi: 10.1126/science.aan3351.
- 34 Smallwood SA, Lee HJ, Angermueller C, et al. Single-cell genome-wide bisulfite sequencing for assessing epigenetic heterogeneity. *Nat Methods* 2014;**11**:817–20. doi: 10.1038/nmeth.3035.
- 35 Gu C, Liu S, Wu Q, et al. Integrative single-cell analysis of transcriptome, DNA methylome and chromatin accessibility in mouse oocytes. *Cell Res* 2019;**29**:110–23. doi: 10.1038/s41422-018-0125-4.
- 36 Wu P, Gao Y, Guo W, et al. Using local alignment to enhance single-cell bisulfite sequencing data efficiency. *Bioinformatics* 2019;**35**:3273–8. doi: 10.1093/bioinformatics/btz125.
- 37 Mulqueen RM, Pokholok D, Norberg SJ, et al. Highly scalable generation of DNA methylation profiles in single cells. *Nat Biotechnol* 2018;**36**:428–31. doi: 10.1038/nbt.4112.
- 38 Shareef SJ, Bevil SM, Raman AT, et al. Extended-representation bisulfite sequencing of gene regulatory elements in multiplexed samples and single cells. *Nat Biotechnol* 2021;**39**:1086–94. doi: 10.1038/s41587-021-00910-x.
- 39 Wen L, Tang F. Single cell epigenome sequencing technologies. *Mol Aspects Med* 2018;**59**:62–9. doi: 10.1016/j.mam.2017.09.002.
- 40 Cao J, Cusanovich DA, Ramani V, et al. Joint profiling of chromatin accessibility and gene expression in thousands of single cells. *Science* 2018;**361**:1380–5. doi: 10.1126/science.aau0730.
- 41 Chen S, Lake BB, Zhang K. High-throughput sequencing of the transcriptome and chromatin accessibility in the same cell. *Nat Biotechnol* 2019;**37**:1452–7. doi: 10.1038/s41587-019-0290-0.
- 42 Liu L, Liu C, Quintero A, et al. Deconvolution of single-cell multi-omics layers reveals regulatory heterogeneity. *Nat Commun* 2019;**10**:470. doi: 10.1038/s41467-018-08205-7.
- 43 Ma S, Zhang B, LaFave LM, et al. Chromatin potential identified by shared single-cell profiling of RNA and chromatin. *Cell* 2020;**183**:1103–16. e20. doi: 10.1016/j.cell.2020.09.056.
- 44 Zhu C, Yu M, Huang H, et al. An ultra high-throughput method for single-cell joint analysis of open chromatin and transcriptome. *Nat Struct Mol Biol* 2019;**26**:1063–70. doi: 10.1038/s41594-019-0323-x.
- 45 Wagner DE, Klein AM. Lineage tracing meets single-cell omics: opportunities and challenges. *Nat Rev Genet* 2020;**21**:410–27. doi: 10.1038/s41576-020-0223-2.
- 46 Telenius H, Carter NP, Bebb CE, et al. Degenerate oligonucleotide-primed PCR: general amplification of target DNA by a single degenerate primer. *Genomics* 1992;**13**:718–25. doi: 10.1016/0888-7543(92)90147-k.
- 47 Dean FB, Nelson JR, Giesler TL, et al. Rapid amplification of plasmid and phage DNA using Phi 29 DNA polymerase and multiply-primed rolling circle amplification. *Genome Res* 2001;**11**:1095–9. doi: 10.1101/gr.180501.
- 48 Chen C, Xing D, Tan L, et al. Single-cell whole-genome analyses by Linear Amplification via Transposon Insertion (LIANTI). *Science* 2017;**356**:189–94. doi: 10.1126/science.aak9787.
- 49 Chu WK, Edge P, Lee HS, et al. Ultraaccurate genome sequencing and haplotyping of single human cells. *Proc Natl Acad Sci U S A* 2017;**114**:12512–7. doi: 10.1073/pnas.1707609114.
- 50 Fu Y, Li C, Lu S, et al. Uniform and accurate single-cell sequencing based on emulsion whole-genome amplification. *Proc Natl Acad Sci U S A* 2015;**112**:11923–8. doi: 10.1073/pnas.1513988112.
- 51 Xing D, Tan L, Chang CH, et al. Accurate SNV detection in single cells by transposon-based whole-genome amplification of complementary strands. *Proc Natl Acad Sci U S A* 2021;**118**:e2013106118. doi: 10.1073/pnas.2013106118.
- 52 Zong C, Lu S, Chapman AR, et al. Genome-wide detection of single-nucleotide and copy-number variations of a single human cell. *Science* 2012;**338**:1622–6. doi: 10.1126/science.1229164.
- 53 Lodato MA, Woodworth MB, Lee S, et al. Somatic mutation in single human neurons tracks developmental and transcriptional history. *Science* 2015;**350**:94–8. doi: 10.1126/science.aab1785.
- 54 Lodato MA, Rodin RE, Bohrsen CL, et al. Aging and neurodegeneration are associated with increased mutations in single human neurons. *Science* 2018;**359**:555–9. doi: 10.1126/science.aao4426.
- 55 Lee-Six H, Øbro NF, Shepherd MS, et al. Population dynamics of normal human blood inferred from somatic mutations. *Nature* 2018;**561**:473–8. doi: 10.1038/s41586-018-0497-0.
- 56 Spencer Chapman M, Ranzoni AM, Myers B, et al. Lineage tracing of human development through somatic mutations. *Nature* 2021;**595**:85–90. doi: 10.1038/s41586-021-03548-6.
- 57 Blokzijl F, de Ligt J, Jager M, et al. Tissue-specific mutation accumulation in human adult stem cells during life. *Nature* 2016;**538**:260–4. doi: 10.1038/nature19768.
- 58 Chapal-Ilani N, Maruvka YE, Spiro A, et al. Comparing algorithms that reconstruct cell lineage trees utilizing information on microsatellite mutations. *PLoS Comput Biol* 2013;**9**:e1003297. doi: 10.1371/journal.pcbi.1003297.

- 59 Kazazian HH, Moran JV. Mobile DNA in Health and Disease. *N Engl J Med* 2017;**377**:361–70. doi: 10.1056/NEJMra1510092.
- 60 Evrony GD, Cai X, Lee E, et al. Single-neuron sequencing analysis of L1 retrotransposition and somatic mutation in the human brain. *Cell* 2012;**151**:483–96. doi: 10.1016/j.cell.2012.09.035.
- 61 Upton KR, Gerhardt DJ, Jesuadian JS, et al. Ubiquitous L1 mosaicism in hippocampal neurons. *Cell* 2015;**161**:228–39. doi: 10.1016/j.cell.2015.03.026.
- 62 Ewing AD, Kazazian HH Jr. High-throughput sequencing reveals extensive variation in human-specific L1 content in individual human genomes. *Genome Res* 2010;**20**:1262–70. doi: 10.1101/gr.106419.110.
- 63 Evrony GD, Lee E, Mehta BK, et al. Cell lineage analysis in human brain using endogenous retroelements. *Neuron* 2015;**85**:49–59. doi: 10.1016/j.neuron.2014.12.028.
- 64 Cai X, Evrony GD, Lehmann HS, et al. Single-cell, genome-wide sequencing identifies clonal somatic copy-number variation in the human brain. *Cell Rep* 2015;**10**:645. doi: 10.1016/j.celrep.2015.01.028.
- 65 Jacobs KB, Yeager M, Zhou W, et al. Detectable clonal mosaicism and its relationship to aging and cancer. *Nat Genet* 2012;**44**:651–8. doi: 10.1038/ng.2270.
- 66 Knouse KA, Wu J, Amon A. Assessment of megabase-scale somatic copy number variation using single-cell sequencing. *Genome Res* 2016;**26**:376–84. doi: 10.1101/gr.198937.115.
- 67 Laurie CC, Laurie CA, Rice K, et al. Detectable clonal mosaicism from birth to old age and its relationship to cancer. *Nat Genet* 2012;**44**:642–50. doi: 10.1038/ng.2271.
- 68 McConnell MJ, Lindberg MR, Brennand KJ, et al. Mosaic copy number variation in human neurons. *Science* 2013;**342**:632–7. doi: 10.1126/science.1243472.
- 69 Zhou Y, Bian S, Zhou X, et al. Single-Cell multiomics sequencing reveals prevalent genomic alterations in tumor stromal cells of human colorectal cancer. *Cancer Cell* 2020;**38**:818–28. e815. doi: 10.1016/j.ccell.2020.09.015.
- 70 Ludwig LS, Lareau CA, Ulirsch JC, et al. Lineage tracing in humans enabled by mitochondrial mutations and single-cell genomics. *Cell* 2019;**176**:1325–39. e1322. doi: 10.1016/j.cell.2019.01.022.
- 71 Bae T, Tomasini L, Mariani J, et al. Different mutational rates and mechanisms in human cells at pregastrulation and neurogenesis. *Science* 2018;**359**:550–5. doi: 10.1126/science.aan8690.
- 72 Behjati S, Huch M, van Boxtel R, et al. Genome sequencing of normal cells reveals developmental lineages and mutational processes. *Nature* 2014;**513**:422–5. doi: 10.1038/nature13448.
- 73 Coorens THH, Moore L, Robinson PS, et al. Extensive phylogenies of human development inferred from somatic mutations. *Nature* 2021;**597**:387–92. doi: 10.1038/s41586-021-03790-y.
- 74 Park S, Mali NM, Kim R, et al. Clonal dynamics in early human embryogenesis inferred from somatic mutation. *Nature* 2021;**597**:393–7. doi: 10.1038/s41586-021-03786-8.
- 75 Asp M, Bergenstråhle J, Lundeberg J. Spatially Resolved Transcriptomes-Next Generation Tools for Tissue Exploration. *Bioessays* 2020;**42**:e1900221. doi: 10.1002/bies.201900221.
- 76 Chen KH, Boettiger AN, Moffitt JR, et al. RNA imaging. Spatially resolved, highly multiplexed RNA profiling in single cells. *Science* 2015;**348**:aaa6090. doi: 10.1126/science.aaa6090.
- 77 Lubeck E, Coskun AF, Zhiyentayev T, et al. Single-cell in situ RNA profiling by sequential hybridization. *Nat Methods* 2014;**11**:360–1. doi: 10.1038/nmeth.2892.
- 78 Eng CL, Lawson M, Zhu Q, et al. Transcriptome-scale super-resolved imaging in tissues by RNA seqFISH. *Nature* 2019;**568**:235–9. doi: 10.1038/s41586-019-1049-y.
- 79 Xia C, Fan J, Emanuel G, et al. Spatial transcriptome profiling by MERFISH reveals subcellular RNA compartmentalization and cell cycle-dependent gene expression. *Proc Natl Acad Sci U S A* 2019;**116**:19490–9. doi: 10.1073/pnas.1912459116.
- 80 Ke R, Mignardi M, Pacureanu A, et al. In situ sequencing for RNA analysis in preserved tissue and cells. *Nat Methods* 2013;**10**:857–60. doi: 10.1038/nmeth.2563.
- 81 Lee JH, Daugharthy ER, Scheiman J, et al. Highly multiplexed subcellular RNA sequencing in situ. *Science* 2014;**343**:1360–3. doi: 10.1126/science.1250212.
- 82 Alon S, Goodwin DR, Sinha A, et al. Expansion sequencing: Spatially precise in situ transcriptomics in intact biological systems. *Science* 2021;371. doi: 10.1126/science.aax2656.
- 83 Wang X, Allen WE, Wright MA, et al. Three-dimensional intact-tissue sequencing of single-cell transcriptional states. *Science* 2018;361. doi: 10.1126/science.aat5691.
- 84 Lovatt D, Ruble BK, Lee J, et al. Transcriptome in vivo analysis (TIVA) of spatially defined single cells in live tissue. *Nat Methods* 2014;**11**:190–6. doi: 10.1038/nmeth.2804.
- 85 Ståhl PL, Salmén F, Vickovic S, et al. Visualization and analysis of gene expression in tissue sections by spatial transcriptomics. *Science* 2016;**353**:78–82. doi: 10.1126/science.aaf2403.
- 86 Rodrigues SG, Stickels RR, Goeva A, et al. Slide-seq: A scalable technology for measuring genome-wide expression at high spatial resolution. *Science* 2019;**363**:1463–7. doi: 10.1126/science.aaw1219.
- 87 Stickels RR, Murray E, Kumar P, et al. Highly sensitive spatial transcriptomics at near-cellular resolution with Slide-seqV2. *Nat Biotechnol* 2021;**39**:313–9. doi: 10.1038/s41587-020-0739-1.
- 88 Xia K, Sun H, Li J, et al. Single-cell Stereo-seq enables cell type-specific spatial 1 transcriptome characterization in Arabidopsis leaves. *bioRxiv* 2021. doi: <https://doi.org/10.1101/2021.10.20.465066>.
- 89 Peng G, Suo S, Chen J, et al. Spatial Transcriptome for the Molecular Annotation of Lineage Fates and Cell Identity in Mid-gastrula Mouse Embryo. *Dev Cell* 2016;**36**:681–97. doi: 10.1016/j.devcel.2016.02.020.
- 90 Junker JP, Noel ES, Guryev V, et al. Genome-wide RNA Tomography in the zebrafish embryo. *Cell* 2014;**159**:662–75. doi: 10.1016/j.cell.2014.09.038.
- 91 Goodwin S, McPherson JD, McCombie WR. Coming of age: ten years of next-generation sequencing technologies. *Nat Rev Genet* 2016;**17**:333–51. doi: 10.1038/nrg.2016.49.
- 92 Brinkerhoff H, Kang ASW, Liu J, et al. Multiple rereads of single proteins at single-amino acid resolution using nanopores. *Science* 2021;**374**:1509–13. doi: 10.1126/science.abl4381.
- 93 Fan X, Tang D, Liao Y, et al. Single-cell RNA-seq analysis of mouse preimplantation embryos by third-generation sequencing. *PLoS Biol* 2020;**18**:e3001017. doi: 10.1371/journal.pbio.3001017.
- 94 Gupta I, Collier PG, Haase B, et al. Single-cell isoform RNA sequencing characterizes isoforms in thousands of cerebellar cells. *Nat Biotechnol* 2018. doi: 10.1038/nbt.4259.
- 95 Singh M, Al-Eryani G, Carswell S, et al. High-throughput targeted long-read single cell sequencing reveals the clonal and transcriptional landscape of lymphocytes. *Nat Commun* 2019;**10**:3120. doi: 10.1038/s41467-019-11049-4.
- 96 Volden R, Palmer T, Byrne A, et al. Improving nanopore read accuracy with the R2C2 method enables the sequencing of highly multiplexed full-length single-cell cDNA. *Proc Natl Acad Sci U S A* 2018;**115**:9726–31. doi: 10.1073/pnas.1806447115.
- 97 Sedlazeck FJ, Rescheneder P, Smolka M, et al. Accurate detection of complex structural variations using single-molecule

- sequencing. *Nat Methods* 2018;**15**:461–8. doi: 10.1038/s41592-018-0001-7.
- 98 Fan X, Yang C, Li W, et al. SMOOTH-seq: single-cell genome sequencing of human cells on a third-generation sequencing platform. *Genome Biol* 2021;**22**:195. doi: 10.1186/s13059-021-02406-y.
- 99 Lee I, Razaghi R, Gilpatrick T, et al. Simultaneous profiling of chromatin accessibility and methylation on human cell lines with nanopore sequencing. *Nat Methods* 2020;**17**:1191–9. doi: 10.1038/s41592-020-01000-7.
- 100 Shipony Z, Marinov GK, Swaffer MP, et al. Long-range single-molecule mapping of chromatin accessibility in eukaryotes. *Nat Methods* 2020;**17**:319–27. doi: 10.1038/s41592-019-0730-2.
- 101 Stergachis AB, Debo BM, Haugen E, et al. Single-molecule regulatory architectures captured by chromatin fiber sequencing. *Science* 2020;**368**:1449–54. doi: 10.1126/science.aaz1646.