

Receptor for advanced glycation end-products (RAGE) provides a link between genetic susceptibility and environmental factors in type 1 diabetes

J. M. Forbes · J. Söderlund · F. Y. T. Yap · M. Knip · S. Andrikopoulos · J. Ilonen · O. Simell · R. Veijola · K. C. Sourris · M. T. Coughlan · C. Forsblom · R. Slattery · S. T. Grey · M. Wessman · H. Yamamoto · A. Bierhaus · M. E. Cooper · P.-H. Groop

Received: 10 October 2010 / Accepted: 14 December 2010 / Published online: 6 February 2011
© Springer-Verlag 2011

Abstract

Aims/hypothesis This group of studies examines human genetic susceptibility conferred by the receptor for advanced glycation end-products (RAGE) in type 1 diabetes

and investigates how this may interact with a western environment.

Methods We analysed the *AGER* gene, using 13 tag SNPs, in 3,624 Finnish individuals from the FinnDiane study,

Electronic supplementary material The online version of this article (doi:10.1007/s00125-011-2058-z) contains supplementary material, which is available to authorised users.

J. M. Forbes (✉) · F. Y. T. Yap · K. C. Sourris · M. T. Coughlan · M. E. Cooper · P.-H. Groop
Diabetes Complications Division,
Baker IDI Heart and Diabetes Institute, St Kilda Rd Central,
P.O. Box 6492, Melbourne, VIC 8008, Australia
e-mail: Josephine.Forbes@bakeridi.edu.au

J. Ilonen
Immunogenetics Laboratory, University of Turku,
Turku, Finland

O. Simell
Department of Paediatrics, University of Turku,
Turku, Finland

J. M. Forbes · R. Slattery · M. E. Cooper
Department of Immunology and Medicine,
Monash University, Melbourne, VIC, Australia

R. Veijola
Department of Paediatrics, University of Oulu,
Oulu, Finland

J. Söderlund · M. Knip · C. Forsblom · M. Wessman · P.-H. Groop
Department of Diabetes Genetics, Folkhälsan Institute of Genetics,
University of Helsinki, Helsinki, Finland

S. T. Grey
Gene Therapy and Autoimmunity Group, Garvan Institute,
Darlinghurst, NSW, Australia

J. Söderlund · C. Forsblom · P.-H. Groop
Department of Medicine, Division of Nephrology,
Helsinki University Central Hospital,
Helsinki, Finland

H. Yamamoto
Department of Biochemistry and Molecular Vascular Biology,
Kanazawa University Graduate School of Medical Science,
Kanazawa, Japan

M. Knip
Hospital for Children and Adolescents,
University of Helsinki, Helsinki, Finland

A. Bierhaus
Department of Internal Medicine I and Clinical Chemistry,
University of Heidelberg,
Heidelberg, Germany

M. Knip
Department of Paediatrics,
Tampere University Hospital,
Tampere, Finland

S. Andrikopoulos
Department of Medicine,
(AH/NH) University of Melbourne,
Heidelberg Repatriation Hospital,
Heidelberg Heights, VIC, Australia

J. Ilonen
Department of Clinical Microbiology,
University of Eastern Finland,
Kuopio, Finland

followed by *AGER* associations with a high risk HLA genotype (*DR3*)-*DQA1**05-*DQB1**02/*DRB1**0401-*DQB1**0302 ($n=546$; *HLA-DR3/DR4*), matched in healthy newborn infants from the Finnish Type 1 Diabetes Prediction and Prevention (DIPP) Study ($n=373$) using allelic analysis. We also studied islets and circulating RAGE in *NODLt* mice.

Results The rs2070600 and rs17493811 polymorphisms predicted increased risk of type 1 diabetes, whereas the rs9469089 SNP was related to decreased risk, on a high risk HLA background. Children from the DIPP study also showed a decline in circulating soluble RAGE levels, at seroconversion to positivity for type 1 diabetes-associated autoantibodies. Islet RAGE and circulating soluble RAGE levels in prediabetic *NODLt* mice decreased over time and were prevented by the AGE lowering therapy alagebrium chloride. Alagebrium chloride also decreased the incidence of autoimmune diabetes and restored islet RAGE levels.

Conclusions/interpretation These studies suggest that inherited *AGER* gene polymorphisms may confer susceptibility to environmental insults. Declining circulating levels of soluble RAGE, before the development of overt diabetes, may also be predictive of clinical disease in children with high to medium risk HLA II backgrounds and this possibility warrants further investigation in a larger cohort.

Keywords Advanced glycation · Alagebrium chloride · Autoimmune diabetes · Children · Insulin · NOD · Polymorphism

Abbreviations

CML	Carboxymethyl lysine
DIPP	Diabetes Prediction and Prevention Study
HMGB1	High mobility group protein B1
HWE	Hardy–Weinberg equilibrium
MAF	Minor allele frequencies
RAGE	Receptor for advanced glycation end-products
SNP	Single nucleotide polymorphism

Introduction

Type 1 diabetes occurs as the result of a complex disease process where genetic and environmental factors lead to an autoimmune response which, to date, remains incompletely defined [1]. Although some major genetic determinants of type 1 diabetes, such as alleles of the major histocompatibility locus (HLA) at the *HLA-DRB1* and *DQB1* loci [2] and more recently the *HLA-B*39* locus [3], have been identified, these only account for some 40–50% of the familial clustering. In addition, approximately 70% of monozygotic twins are discordant for the development of type 1 diabetes

[4], which indicates that environmental triggers are also likely to be important. Indeed, the risk of developing type 1 diabetes is still increasing by 3–4% annually in developed nations [5], which remains unexplained.

One such environmental contributor to type 1 diabetes may be AGEs. Western diets provide excesses of these non-enzymatic modifications [6], which contribute flavour and colour to foodstuffs (e.g. roasted meat or coffee), while more recently AGE modifications also add functional properties, such as improved emulsification. Dietary AGEs are absorbed intestinally and are a prominent contributor to the body's AGE pool [7], in addition to the natural ageing process [8], redox imbalances [9], non-diabetes related diseases including renal impairment [10] and hyperglycaemia [11]. There is increasing evidence to suggest that AGEs also promote beta cell dysfunction [12, 13]. Furthermore, a reduced incidence of diabetes in a mouse model of autoimmune diabetes, the *NOD* mouse (*NODLt*) [14], has been reported following dietary restriction of AGEs. However, the specific mechanisms whereby insulin secretory pathways are damaged by AGEs are not known.

AGEs can exert their biological effects via receptors such as the receptor for advanced glycation end-products (RAGE) [15], although these compounds may have greater effects via modulation of RAGE production per se [16]. RAGE is a multi-ligand receptor involved in host-pathogen defence, in addition to cellular apoptosis [17]. Thus, a recent study has identified a blockade of the late stages of adoptively transferred autoimmune diabetes with the 'decoy' soluble RAGE receptor [18]. In addition, a small molecule inhibitor of RAGE [19] or antibodies against another RAGE ligand, HMGB1 [20], have been shown to delay islet destruction in hyperglycaemic *NODLt* mice. Furthermore, the heritability of insulin resistance [21], another factor postulated to be contributing to the increasing rate of type 1 diabetes incidence [22], is associated with specific RAGE polymorphisms.

Genetic susceptibility to type 1 diabetes remains to be fully defined, in particular, the contribution of certain genotypes in the context of our increasingly Westernised environment. It is therefore unknown if a specific RAGE genotype may play a role in susceptibility to type 1 diabetes; we have now investigated this issue using a combination of human genetics, molecular biology and integrative pathophysiology.

Methods

Human study

FinnDiane/DIPP design and recruitment Characteristics of the 2,921 unrelated type 1 diabetic patients from the nationwide Finnish Diabetic Nephropathy (FinnDiane)

study group [23] are provided in Table 1. Age of diagnosis of 30 years or below was used as a criterion for inclusion in the analysis. A total of 703 unrelated control samples were obtained from the Finnish Red Cross Blood Service. These controls represent a healthy non-diabetic Finnish population of European descent given the strict criteria that the Red Cross Blood Service practices regarding blood donors. Notably, there was no population stratification or admixture applied to their selection which might influence the genetic results (see Electronic supplementary material [ESM]).

Since *AGER* is located within the HLA class III region [24], we further analysed these *AGER* associations outlined above on a matched high risk HLA background. To achieve this, newborn infants ($n=373$), who had the highest risk HLA genotype (*DR3*)-*DQA1**05-*DQB1**02/*DRB1**0401-*DQB1**0302 (from $n=546$; *HLA-DR3/DR4* [25]), were matched with unaffected newborn infants with the identical HLA genotype, from the Finnish Type 1 Diabetes Prediction and Prevention (DIPP) Study ($n=373$) [26]. A subgroup of children was also used for circulating soluble RAGE analyses. We included all children with the high risk HLA genotype identified from the DIPP study as controls. However, unfortunately some participants with the high risk HLA background had to be excluded because it was not possible to extract sufficient quantities of DNA for the analyses from the miniscule blood samples available. All children studied were from Finland and no other children were excluded from the present study. The ethics committees of all participating centres approved the study protocol and all patients or guardians gave written consent. The studies follow the Declaration of Helsinki.

DNA isolation and genotyping DNA was isolated from whole blood with either a Puregene DNA Purification Kit (Gentra Systems, Minneapolis, MN, USA) according to the manufacturer's protocol or with a phenol-chloroform protocol [27].

Tag SNPs for genotyping were selected from the HapMap Project's NCBI build 34 and 35 for the CEU (Utah, USA residents with ancestry from northern and

western Europe populations; $n=180$), as defined by Gabriel et al. [28] to encompass the genetic variation of the entire *AGER* gene. Minor allele frequencies (MAFs) were required to be ≥ 0.05 . Genotyping was performed with TaqMan SNP Genotyping Assays (Applied Biosystems, Foster City, CA, USA) with an ABI Prism 7900HT Sequence Detection System (Perkin-Elmer, Foster City, CA, USA) and the Sequenom multiplex platform (Sequenom, Hamburg, Germany) according to the manufacturers' instructions.

HLA genotyping for the two high risk haplotypes was performed with a sequence specific hybridisation method using oligonucleotide probes specific for *DQA1**05 and *DQB1**02 (*DR3-DQ2* haplotype) and *DRB1**0401 and *DQB1**0302 (*DR4-DQ8* haplotype) as described previously [25].

Soluble RAGE analysis A series of children from the DIPP study were analysed for plasma soluble RAGE. The group comprised 15 children (12 boys) who progressed to clinical type 1 diabetes during prospective observation (patients) and 15 autoantibody-negative controls (non-progressors) who remained non-diabetic and were matched with the patients for sex, date of birth and HLA genotype. Altogether 14 children carried the high risk genotype (*DQA1**05-*DQB1**02/*DQB1**0302) and 16 patients carried moderate risk genotypes (*HLADQB1**0302/ x ; $x \neq$ *DQB1**02, *0301, *0602 or *0603). All children provided three plasma samples. Among the patients, the first sample was obtained before seroconversion to autoantibody positivity at a mean age of 1.1 (range 0.3–3.0) years, the second sample soon after the seroconversion at a mean age of 2.4 (range 0.7–5.3) years and the third sample soon after the diagnosis of clinical type 1 diabetes at a mean age of 5.2 (range 2.0–8.9) years. In the control patients, the samples were obtained at the corresponding ages. Human plasma samples were assayed undiluted according to the manufacturer's instructions contained within the Human RAGE ELISA (R&D Systems, Minneapolis, MN, USA). The inter-assay CV was 7.8% while the intra-assay CV was 5.7%. The limit of sensitivity was 4.12 pg/ml.

Statistical analysis Hardy–Weinberg equilibrium (HWE) for SNPs was tested in the control individuals ($n=703$), and which were considered to be in HWE when $p > 0.05$ using Haploview 4.0 (MIT/Harvard Broad Institute, Boston, MA, USA) [29]. Testing for allelic associations was carried out with logistic regression adjusted for sex in SPSS 15.0 (Chicago, IL, USA).

Due to the strong linkage disequilibrium between SNPs in the HLA region and the HLA class II loci, it was necessary to exclude an effect of the HLA on the associations. Therefore, we matched patients and references for the high risk HLA genotype (*DR3*)-*DQA1**05-

Table 1 Patient characteristics: patients with T1D taken from the FinnDiane study

Characteristic	Patients	Controls ^a	DIPP ^b
<i>n</i>	2,921	703	373
Sex, male (%)	52.2	48.9	N/A
Age (years)	40.3±9.9		N/A
Age at T1D diagnosis (years)	12.8±7.6	N/A	N/A

^a Finnish Red Cross Blood Donors

^b Newborn infants from the Type 1 DIPP T1D, Type 1 diabetes

*DQB1*02/DRB1*0401-DQB1*0302*. Logistic regression was used to evaluate allelic odds ratios.

The issue of sufficient power to detect association between *AGER* SNPs and type 1 diabetes was evaluated using the Genetic Power Calculator (<http://pngu.mgh.harvard.edu/~purcell/gpc/>; [30]) with the module of case–control for discrete traits. Bonferroni correction for multiple comparisons was applied to the FinnDiane cohort under the conditions of 13 SNPs, α level 0.05, and three performed tests. Human soluble RAGE concentrations were analysed by repeated measures ANOVA.

Rodent study

We next studied islet and circulating RAGE production in a spontaneous experimental model which resembles type 1 diabetes, the *NODLt* mouse, since the genetic loci thought to be responsible for susceptibility to diabetes in these mice include the *AGER* gene sequence [31, 32]. *NODLt* or *NODScid* mice fed standard mouse chow (containing the AGE, carboxymethyl lysine (CML): 93.39 nmol [mol lysine]⁻¹ [100 mg]⁻¹) were studied prediabetes at 4, 8 and 12 weeks of age ($n=6$ per group). In addition, groups of 4-week-old *NODLt* mice were randomised to receive either no treatment (vehicle) or the AGE-lowering therapy alagebrium chloride (1 mg kg⁻¹ day⁻¹ i.p.) and followed for 6 weeks (prediabetes, $n=10$ per group). For incidence studies, littermate *NODLt* mice ($n=20$ per group) were studied until the diagnosis of diabetes was confirmed or until day 200 of age, with one group randomised to receive alagebrium chloride (1 mg kg⁻¹ day⁻¹ i.p.) for the study duration from week 4 of age. Mice were given ad libitum access to food and water and maintained on 12 h dark–light cycles. Plasma glucose was measured by autoanalyser (Beckman Coulter LX20PRO, Brea, CA, USA) and plasma insulin concentrations were measured by radioimmunoassay (Linco Research, MO, USA).

Circulating RAGE and AGE ELISA Soluble RAGE in mouse serum (RAGE sandwich ELISA kit, R&D systems, Minneapolis, MN, USA) was assayed undiluted according to the manufacturer's instructions. The inter-assay CV was 6.7% and the intra-assay CV was 5.9%. The limit of sensitivity of the assay was 1 pg/ml. The AGE, CML, was assessed in serum or plasma using an in-house ELISA. The limit of detection of the assay was 8.0 nmol/mol lysine. The inter-assay CV was 7.3%. The intra-assay coefficient of variation was 5.5%.

Immunohistochemistry Formalin fixed paraffin sections [33] were incubated with primary antibody (AGE [34],

1:1,000; goat anti-RAGE 1:2,500, Chemicon, Temecula, CA, USA) overnight at 4°C. Tissue sections were consecutively stained with biotinylated IgG for 10 min and avidin–biotin horseradish peroxidase complex for 15 min (Vectastain ABC Elite kit, Vector Laboratories, Burlingame, CA, USA) before 3,3'-diaminobenzidine tetrahydrochloride treatment (DAB; Sigma Chemical Co., St Louis, MI, USA). Negative control sections omitted the primary antibody. Positive control tissues were also included. Quantitation of islet immunostaining was completed by computer-aided densitometry (Image Pro Plus 6.0, Media Cybernetics, Bethesda, MD, USA) where all islets were analysed ($\times 100$) and results expressed as proportional to the area of positive staining.

Real-time reverse transcription-polymerase chain reaction RNA obtained from pancreatic islet tissue collected immediately was later used to synthesise cDNA with the Superscript First strand synthesis system for RT-PCR (Gibco BRL, Grand Island, NY, USA). Gene expression for *AGER* was analysed by real-time quantitative RT-PCR performed with the TaqMan system based on real-time detection of accumulated fluorescence (ABI Prism 7700, Perkin-Elmer) normalised to 18 S as previously [35]. For mouse *Ager* (*NM_007425*), the probe was 6-FAM CACAGCCCGGATTG-MGB, the forward primer was 5'-GCTGTAGCTGGTGGTCAGAACA-3' and the reverse primer was 5'-CCCCTTACAGCTTAGCACAAGTG-3'.

Islet extraction Pancreatic islets were isolated as previously described [36]. Briefly, 10 ml of cold Hanks balanced salt solution (HBBS) containing 0.75 mg/ml collagenase type V (Sigma Chemical Co.) was injected into the bile duct. The pancreas was digested at 37°C for 10–20 min and then disrupted and filtered through a 500 μ m mesh. The pancreatic islets were separated from exocrine tissue by histopaque density gradient, for which the islets were suspended in histopaque 1.119 g/l, followed by layering of histopaque 1.083 g/l and histopaque 1.077 g/l (Sigma). The hand-picked islets were then used for flow cytometry as outlined below and RNA extraction [37]. For flow cytometry, purified islets were washed (10 mM EDTA in Hanks' balanced salt solution) and stained using goat anti-RAGE (Santa Cruz Biotechnology, Santa Cruz, CA, USA) followed by rabbit anti-goat IgG, F(ab')₂, FITC conjugated (Chemicon International, Temecula, CA, USA).

Statistical analysis Mouse results are expressed as mean \pm standard deviation unless otherwise specified. All analyses of rodent data with more than two groups were performed by ANOVA followed by post hoc analysis with Tukey's test. Student's *t* test or Mann–Whitney analyses were used to compare two groups where appropriate and are labelled

as such. A logrank (Mantel–Cox) test was used to analyse diabetes incidence in NOD mice. A value of $p < 0.05$ was considered to be statistically significant (GraphPad Prism; GraphPad Software, San Diego, CA, USA).

Results

AGER gene polymorphisms and type 1 diabetes susceptibility

FinnDiane Characteristics of individuals from the FinnDiane study have been previously described in detail [38] and are summarised in Table 1. We observed that all 13 SNPs for *AGER* were strongly associated with type 1 diabetes (Table 2), even after Bonferroni correction for multiple comparisons (cut-off $p = 0.0013$). Regarding the HLA frequencies, 19% of the FinnDiane type 1 diabetic patients have the (*DR3*)-*DQA1*05-DQB1*02/DRB1*0401-DQB1*0302* genotype. This percentage of the high risk *DR3/DR4* genotype is small given that this is only one of several other HLA haplotypes/genotypes which are known risk factors for the development of type 1 diabetes [39]. Therefore, in the FinnDiane cohort, 72.8% of all patients

belong to the genetic risk group, as described by Nääntö-Salonen et al. [39]. HWE was tested in the controls and all SNPs were in HWE.

DIPP We performed allelic analyses, which identified that three SNPs (rs2070600, rs9469089 and rs17493811) were still associated with type 1 diabetes ($p = 0.017$, 3.65×10^{-5} , and 0.031, respectively) in HLA-matched children from the DIPP (Table 2). In particular, the rs2070600 and rs17493811 polymorphisms predicted increased risk of type 1 diabetes (ORs 1.452 and 1.518, respectively) whereas rs9469089 was related to decreased risk (OR 0.423) in the high risk HLA setting. However, using less stringent HLA adjustment, i.e. analysing those individuals with the risk alleles *DQB1*02* and/or *DQB1*0302* (type 1 diabetes patients $n = 2,660$ and controls $n = 294$), all *AGER* SNPs apart from rs204993 remained associated with type 1 diabetes following HLA matching (data not shown).

Circulating soluble RAGE concentrations

Children from the DIPP study who progressed to type 1 diabetes (diagnosis; 5.2 ± 2.2 years) had a decline in

Table 2 SNP associations between *AGER* and type 1 diabetes: analysis of genomic DNA from patients with T1D from the FinnDiane cohort and from newborn infants with the highest risk HLA genotype

Position NCBI35 Chr 6 (bp)	SNP (alleles)	Minor ^a allele	SNP location	The FinnDiane cohort		HLA- <i>DR3/DR4</i> matched		OR (95% CI) ^b
				MAF controls/ patients	<i>p</i> value	MAF controls/ patients	<i>p</i> value	
Sample size (<i>n</i>)				703/2,921	3,624	373/546	924	924
32,269,408	rs204990 (G/T)	T	5'	0.148/0.231	1.82×10^{-11} ^c	0.468/0.45	0.465	0.932 (0.773–1.125)
32,269,374	rs3132940 (C/A)	A	5'	0.143/0.043	1.45×10^{-39} ^c	0.011/0.015	0.436	1.404 (0.598–3.298)
32,269,302	rs12663103 (T/C)	C	5'	0.025/0.005	6.30×10^{-10} ^c	0/0.002	n/a	n/a
32,263,559	rs204993 (T/C)	C	5'	0.18/0.318 [‡]	1.27×10^{-23} ^c	0.445/0.425	0.380	1.088 (0.901–1.314)
32,262,976	rs204994 (G/A)	A	5'	0.148/0.220	2.82×10^{-9} ^c	0.464/0.446	0.499	0.937 (0.777–1.131)
32,260,420	rs1800625 (T/C)	C	Promoter	0.146/0.218	2.76×10^{-9} ^c	0.463/0.444	0.474	0.933 (0.772–1.128)
32,260,365	rs1800624 (T/A)	A	Promoter	0.291/0.348	5.89×10^{-5} ^c	0.270/0.239	0.109	0.839 (0.677–1.040)
32,259,421	rs2070600 (G/A)	A	G82S	0.074/0.118	2.71×10^{-6} ^c	0.091/0.127	0.017	1.452 (1.068–1.973)
32,259,200	rs1035798 (C/T)	T	Intronic	0.295/0.348	1.53×10^{-4} ^c	0.268/0.239	0.132	0.848 (0.685–1.051)
32,254,635	rs9469089 (G/C)	C	3'	0.234/0.142	3.03×10^{-16} ^c	0.084/0.037	3.65×10^{-5}	0.423 (0.282–0.637)
32,253,685	rs2269423 (G/T)	T	3'	0.440/0.280	1.36×10^{-30} ^c	0.150/0.143	0.785	0.964 (0.741–1.254)
32,253,377	rs17493811 (C/G)	G	3'	0.026/0.048	6.34×10^{-4} ^c	0.056/0.082	0.031	1.518 (1.038–2.220)
32,253,183	rs3134947 (G/A)	A	3'	0.150/0.219	1.23×10^{-8} ^c	0.465/0.444	0.423	0.926 (0.767–1.118)

In the FinnDiane cohort, the *p* values are from logistic regression analyses, adjusted for sex, and reflect the joint effect of *AGER* and HLA. The DIPP HLA *DR3/DR4* matched set show the effect of *AGER* conditional for HLA haplotype

^a The minor allele is the opposite in the *DR3/DR4* matched data

^b The allelic ORs with 95% CIs are modelled for the respective minor allele

^c SNPs in the FinnDiane cohort that survive the stringent Bonferroni correction for multiple testing nucleotide polymorphism

MAF, minor allele frequency

circulating soluble RAGE levels with concomitant seroconversion to positivity for type 1 diabetes-associated autoantibodies (islet cell antibodies, insulin autoantibodies, antibodies to glutamic acid decarboxylase and/or antibodies to islet antigen 2; Fig. 1).

Islet RAGE and circulating soluble RAGE decline prediabetes in NODLt mice

We first identified that islet RAGE expression was decreased in NODLt mice and, to a significantly lesser extent, in NODScid mice by week 8, compared with 4-week-old NOD mice (Fig. 2 a,b and d). However, by week 12, islet RAGE expression was significantly higher in both NODLt and NODScid (Fig. 2c and d). In NODLt mice, the loss of islet RAGE expression between weeks 4 and 12 was prevented with the AGE-lowering therapy, alagebrium chloride (Fig. 2e). In addition, the islet expression of the gene encoding RAGE (*AGER*) declined over time and was below detectable levels by week 8 of age in NODLt mice (4 weeks, 1.14 ± 0.63 fold induction; $n=4$ per group; $p < 0.001$ 4 vs 8 or 12 weeks) as assessed by real-time RT-PCR [40]. There was also a significant decline in the levels of circulating soluble RAGE from 4 to 6 weeks of age in NODLt mice (Fig. 2f) that was not seen in NODScid mice (Fig. 2f). Alagebrium chloride also significantly improved circulating soluble RAGE concentrations at week 10 of age (Fig. 2g). There was a positive association between the loss of islet RAGE and declining circulating soluble RAGE in NODLt mice ($r=0.85$, $p < 0.001$). In addition, serum concentrations of the AGE,

CML, were elevated at both week 8 and 12 weeks of age in NODLt mice as compared with NODScid mice (Fig. 2h), which was also ameliorated by treatment to reduce AGE accumulation with alagebrium chloride at week 10 (NODLt, $4,826 \pm 12,61$ nmol/mol lysine vs NODLt treated with alagebrium chloride, $2,059 \pm 589$ nmol/mol lysine, $p=0.026$ Student's *t* test). There was also a decline in plasma concentrations of the RAGE ligand HMGB1 seen at week 8 in NODLt mice when compared with NODScid mice (Fig. 2i). Furthermore, alagebrium chloride markedly decreased the incidence of autoimmune diabetes in NODLt mice from 50% to 10% (Fig. 2j).

Immunohistochemistry also demonstrated a decline in islet RAGE staining in NODLt mice at weeks 8 and 12 prediabetes, which was not seen in NODScid mice (see Fig. 3a–g). RAGE immunostaining in islets was localised primarily to alpha cells in both NODLt and NODScid mice (Fig. 3). Islet AGE immunohistochemistry demonstrated increased concentrations of CML at all prediabetic time points in NODScid mice (Fig. 3h), although only a small proportion of AGE staining in NODScid mice was cell surface bound as determined by flow cytometry compared with NODLt mice (Fig. 3i). There was also a significant, but transient, increase in plasma glucose concentrations seen at week 8 in NODLt (4 week NODLt, 2.7 ± 0.5 mmol/l vs 8 week NODLt, 7.2 ± 2.1 mmol/l, $p < 0.05$; vs 12 week NODLt, 6.9 ± 1.1 mmol/l, NS). There was no difference seen in plasma glucose (8 week NODScid, 7.4 ± 4.2 mmol/l; 12 week NODScid, 6.1 ± 2.6 mmol/l) or glycated haemoglobin concentrations (Table 3) between NODLt and NODScid mice at either 8 or 12 weeks of age. NODLt mice were heavier than NODScid mice by 12 weeks of the study (Table 3).




Fig. 1 Human participants who progress to type 1 diabetes show a decline in soluble RAGE at seroconversion to autoantibody positivity. Serial plasma samples before seroconversion (white bars; mean age 1.1 ± 0.6 years), following initial seroconversion (black bars; mean age 2.4 ± 1.2 years) to positivity for diabetes-associated autoantibodies (islet cell antibodies, insulin autoantibodies, antibodies to glutamic acid decarboxylase, and/or antibodies to islet antigen 2) and at clinical diagnosis of type 1 diabetes (hatched bar; mean age 5.2 ± 2.2 years), in the progressors ($n=15$) and samples obtained at corresponding ages in autoantibody-negative non-progressors (hatched bar; $n=15$), were analysed for soluble RAGE concentrations. † $p=0.025$ vs initial sample; ‡ $p=0.02$ vs initial sample; *** $p < 0.001$ vs sample obtained at seroconversion to autoantibody positivity. sRAGE, soluble RAGE

Discussion

Many chronic diseases are likely to be polygenic, including type 1 diabetes, where less than 50% of the familial clustering can be attributed to currently reported genotypes. This highlights the importance of discovering other candidate genes and environmental contributors. Within the present study, we have identified three polymorphisms in the *AGER* gene, which encodes for the proteins RAGE and soluble RAGE, that may partly explain the genetic susceptibility to type 1 diabetes in humans, even in individuals with a high risk HLA class II background. Of particular interest is the finding that a temporal decline in circulating concentrations of the protective isoform of RAGE (termed soluble RAGE) at seroconversion to produce autoantibodies, predicted the clinical onset of type 1 diabetes in patients within high to moderate risk class II HLA genotypes. These findings are reminiscent of changes




Fig. 2 Prediabetic NOD mice have a deficiency in islet RAGE production. **a–c** Flow cytometry for islet RAGE levels; **a–c** representative histograms in prediabetic NOD Lt mouse islets ($n=5$ per group) at 4 weeks (**a**), 8 weeks (**b**) and 12 weeks (**c**). **d, e** Percentage of cell surface RAGE-positive islets in (**d**) NOD Lt (white bars) and NOD $Scid$ (black bars) mice at weeks 4, 8 and 12 and (**e**) at week 10 ($n=10$ per group, after treatment with alagebrium chloride (ALT), an inhibitor of RAGE ligand (AGE) accumulation). **f, g** Plasma concentrations of circulating soluble RAGE by ELISA; (**f**) time course in prediabetic NOD Lt and NOD $Scid$ mice (NOD Lt , white bars; NOD $Scid$, black bars) (**g**) At week 10 following ALT therapy. **h** Time-course plasma concentrations of CML in prediabetic NOD Lt and NOD $Scid$ mice

($n=5-8$ per group) (NOD Lt , white bars; NOD $Scid$, black bars). **i** Plasma concentrations of the RAGE ligand, HMGB1, in NOD Lt and NOD $Scid$ mice ($n=10$ per group) (NOD Lt , white bars; NOD $Scid$, black bars). **j** Diabetes incidence in untreated NOD Lt mice (square, dashed line) or in NOD Lt mice administered with ALT (circle, continuous line) ($n=20$ per group). $\dagger p=0.0283$ vs NOD Lt , Student's t test, two tailed; $**p<0.01$ vs 4 W NOD Lt ; $***p<0.001$ vs 4 W NOD Lt , $\dagger p=0.0238$ vs 4 W, Mann-Whitney test; $\ddagger p=0.0016$ vs untreated NOD Lt , Student's t test, two tailed; $\S p=0.009$ vs NOD Lt +ALT, logrank (Mantel-Cox) test; $\ddagger\ddagger p<0.01$ vs 8 W NOD Lt ; $\ddagger\ddagger p<0.01$ vs 12 W NOD Lt , $*p<0.05$ vs 4 W NOD $Scid$. sRAGE, soluble RAGE; W, week

in NOD Lt mice, in whom seroconversion appears concurrently with a decrease in circulating soluble RAGE concentrations [41]. There was also an inverse relationship between circulating CML (an AGE) and either soluble RAGE concentrations or islet RAGE expression seen in both NOD Lt and NOD $Scid$ mice. Interestingly, despite elevated circulating AGE levels, islet AGE concentrations were also lower in NOD Lt mice compared with NOD $Scid$ mice, suggesting that RAGE may be involved in the cellular uptake or intracellular production of AGEs in pancreatic islet cells.

A decline in circulating soluble RAGE has also been shown in children during active autoimmune disease, such as Kawasaki disease or systemic onset juvenile idiopathic arthritis [42]. Decreases in plasma soluble RAGE were also seen in samples from children who progressed to type 1 diabetes at seroconversion to autoantibodies, compared with samples obtained before seroconversion. At the

diagnosis of diabetes, those individuals who progressed to type 1 diabetes in the DIPP study had an increase in plasma soluble RAGE, probably as the result of active inflammation, as has been seen previously in individuals with type 2 diabetes [43]. Therefore, although tantalising, the changes in soluble RAGE seen within the present study in children who progressed to type 1 diabetes need to be confirmed in larger patient cohorts. In addition, since soluble RAGE concentrations were not measured in the individuals from whom we have the *AGER* genotype data, Mendelian randomisation was not possible in the present populations.

The genetic loci thought to be responsible for susceptibility to autoimmune diabetes in NOD Lt mice include the *AGER* gene sequence encoding RAGE proteins [31, 32]. The presence of the serine residue within the G82S polymorphism, identified using *AGER* rs2070600 in humans, has previously been associated with lower [44] or elevated [43] circulating soluble RAGE concentrations.




Fig. 3 AGE and RAGE localisation in pancreatic islets in NOD mice. Photomicrographs of immunohistochemistry for RAGE in (a–c) NODScid mice at (a) week 4 (b) week 8 and (c) week 12; and (d–f) NODLt mice at (d) week 4 (e) week 8 and (f) week 12. **g**

Quantification of islets. **h** Quantification of islet CML by immunohistochemistry. **i** Cell surface binding of AGEs to islets by flow cytometry. * $p < 0.05$ NODScid vs NODLt within the same time point; † $p < 0.05$ vs 4 W NODLt; ‡ $p < 0.05$ vs 4 W NODScid. W, week

Table 3 Metabolic variables for NOD mouse groups

Mouse group	GHb (%)	BW (g)
NODLt		
4 weeks	2.6±0.1	18.3±1.3
8 weeks	2.2±0.1	23.7±1.7*
12 weeks	2.4±0.1	27.7±1.0*†
NODScid		
4 weeks	2.8±0.3	18.1±1.6
8 weeks	3.1±0.1	23.4±1.6*
12 weeks	2.8±0.1	24.0±1.3*‡

Data are mean±SD

Glycated haemoglobin and final body weights at the study end-points $n=10$ mice per group

* $p < 0.05$ vs respective 4 week group; † $p = 0.05$ vs 8 week NODLt, ‡ $p = 0.05$ at 12 weeks for NODLt vs NODScid

GHb, glycated haemoglobin; BW, body weight

Interestingly, this SNP was also associated with a higher risk of type 1 diabetes in our human cohorts in the present study. In our human studies, it is difficult to increase the numbers of patients and controls studied, given the key requirement to control for high risk HLA genotypes to delineate the association between *AGER* for type 1 diabetes. It is nevertheless intriguing that, even in such a small HLA-matched sample, there were clear genotypic differences and changes in circulating soluble RAGE concentrations between those with type 1 diabetes and those without. However, this finding now needs to be replicated in larger human populations.

One cannot discount, however, the possibility that changes in the proteases responsible for the cleavage of membrane RAGE to soluble RAGE [45] may also be altering the circulating soluble RAGE pool in our diabetic individuals and mice, independent of changes in the *AGER* gene. Indeed, this possibility should be further investigated in subsequent studies, in particular given the effects of AGEs on circulating RAGE concentrations in NODLt mice observed in the present study. Interestingly, prediabetic

NODLt mice in the present study also had a transient decline in islet RAGE expression at seroconversion to autoantibody positivity which was not seen in NODScid mice. Furthermore, we have determined that circulating soluble RAGE concentrations correlate with islet RAGE expression. In the future, it would be desirable to confirm these findings using a NODLt mouse with an islet specific knockout of the gene encoding RAGE (*AGER*). It would also have been highly desirable to confirm a loss of islet RAGE expression in human participants with type 1 diabetes, although we have shown that decreases in circulating soluble RAGE concentrations in children with medium to high risk HLA II genetic backgrounds may be predictive of this loss in islet RAGE expression. Therefore, at this stage we can only speculate as to the association of RAGE with type 1 diabetes.

Some previous studies in adoptively transferred autoimmune diabetes and syngeneic islet transplants in hyperglycaemic NODLt mice have shown benefits from the interruption of RAGE ligand binding with small molecule inhibitors of RAGE [19] or exogenous soluble RAGE [18]. In adoptively transferred diabetes [18], however, it could only be concluded that RAGE ligands may be involved in the differentiation of T cells to a mature phenotype. Indeed, NODScid mice also have neither T nor B cells, providing further evidence that RAGE expression on T cells, such as Th1 [18, 19], may also be important in the pathogenesis of diabetes, given that NODScid mice do not develop autoimmune diabetes.

Prediabetic NODLt mice in the present study showed increases in circulating RAGE ligands, AGEs, that were not seen with respect to another RAGE ligand, HMGB1. This was surprising given the previous association of HMGB1 to autoimmune diabetes in NODLt mice [20]. However, another ligand of RAGE is the amylin polypeptide, which is co-produced with insulin by the pancreatic beta cells in response to elevation of plasma glucose [46]. Moreover, there is speculation that insulin may also bind to RAGE, since there is evidence of amyloidosis composed of iatrogenic A-Ins type amyloid [47]. Indeed, other investigators have suggested that proinflammatory RAGE and HLA II DRB1 polymorphisms may synergise to activate the immune response in diabetes complications [48]. However, whether this affects the immune system during the development of autoimmune diabetes or type 1 diabetes remains to be determined.

Importantly, lowering AGEs using alagebrium chloride in the present study not only increased circulating soluble RAGE concentrations in NODLt mice at seroconversion to autoantibodies, but also reduced the incidence of autoimmune diabetes. Taken together, one could speculate that it is important therapeutically to prevent ligand-induced chronic modulation of RAGE to protect islet function. Also of

interest was the fact that it is unlikely that increases in plasma glucose are driving the elevations in plasma AGEs seen within the present study, as is classically seen in diabetes complications [33, 49, 50], since elevations in circulating AGEs by 8 weeks of age were also evident in NODScid mice, which do not develop diabetes. These results raise the possibility that precipitation of diabetes could occur as the result of either genetic or environmental modulation of RAGE.

Another limitation of the present study is that so-called high risk HLA haplotypes are ‘losing power’ in the younger age groups developing type 1 diabetes [51]. However, FinnDiane is collecting data on adult type 1 diabetic patients, rather than paediatric patients, and therefore the mean duration of type 1 diabetes is relatively high (24.4 years, or median of 24.2 years), meaning that less than 10% of the patients studied were born in the late 1970s and 1980s, when the HLA trend mentioned above was primarily observed. Furthermore, given that the controls for the FinnDiane study were recruited through the Finnish blood bank, we have little epidemiological data relating to these individuals.

Thus, in summary, the data presented here show that inherited *AGER* gene polymorphisms should be considered as novel contributors to susceptibility to type 1 diabetes. This group of studies also suggests that excesses of environmental factors such as AGEs might modulate changes in islet and circulating RAGE expression, which may contribute to insulin secretory defects and ultimately the development of overt diabetes. In addition, declining circulating levels of soluble RAGE, which binds excesses of AGEs, at seroconversion before the development of overt diabetes, may warrant investigation in larger human cohorts as a predictor of clinical type 1 diabetes. Thus, we not only propose a novel mechanism for the development of diabetes, but also suggest a potential treatment, the AGE lowering therapy alagebrium chloride, currently under clinical investigation, albeit for other medical indications. Furthermore, we believe that these results challenge current thoughts in this area, by hypothesising that genetic or environmental declines in RAGE may be important for the development of type 1 diabetes. These findings are also likely to have implications for the pathogenesis of other chronic diseases in which this receptor is considered to play a pivotal role.

Acknowledgements We are indebted to M Parkkonen and R Sallinen, University of Helsinki, Helsinki, Finland, who contributed to the genotyping of the FinnDiane patients. The authors would also like to thank K Gilbert, V Thallas and M Arnstein, Baker IDI Heart and Diabetes Institute, Melbourne, Australia, for their technical expertise. Thanks also to A Blair for assistance with rat and mouse IV/IPGTTs. This work was completed with support from an innovative grant from the Juvenile Diabetes Research Foundation

(JDRF; 5-2010-163). J Forbes is a JDRF Career Development Fellow. S Andrikopoulos is a recipient of an RD Wright Fellowship from the National Health and Medical Research Council of Australia (NHMRC) of Australia. M Cooper is a JDRF Scholar and an NHMRC Australia Fellow. The FinnDiane is funded by the Folkhälsan Research Foundation, Wilhelm and Else Stockmann Foundation, Sigrid Juselius Foundation, European Commission (LSHB-CT-2003-503364, LSHB-CT-2006-037681), Academy of Finland (214335 and 124280 to M.W.), Medicinska understödsföreningen Liv och Hälsa, Signe and Ane Gyllenberg Foundation, Waldemar von Frenckell Foundation and Governmental Grants for Health Sciences Research. We acknowledge all the physicians and nurses at each participating centre for their invaluable role in patient recruitment, collection of samples and data (see ESM). The DIPP Study is supported by grants from the Juvenile Diabetes Research Foundation (grants 4-1998-274, 4-1999-731, 4-2001-435), Emil Aaltonen Foundation, Academy of Finland, Jalmari and Rauha Ahokas Foundation, Diabetes Research Foundation, Finland, Signe and Ane Gyllenberg Foundation, Yrjö Jahnnson Foundation, Sigrid Juselius Foundation, Novo Nordisk Foundation, Päivikki and Sakari Sohlberg Foundation, and Special Research Funds for the Oulu, Tampere and Turku University Hospitals.

Duality of interest The authors declare that there is no duality of interest associated with this manuscript.

References

- Mathis D, Vence L, Benoist C (2001) Beta-cell death during progression to diabetes. *Nature* 414:792–798
- Davies JL, Kawaguchi Y, Bennett ST et al (1994) A genome-wide search for human type 1 diabetes susceptibility genes. *Nature* 371:130–136
- Nejentsev S, Howson JM, Walker NM et al (2007) Localization of type 1 diabetes susceptibility to the MHC class I genes HLA-B and HLA-A. *Nature* 450:887–892
- Hyttinen V, Kaprio J, Kinnunen L, Koskenvuo M, Tuomilehto J (2003) Genetic liability of type 1 diabetes and the onset age among 22,650 young Finnish twin pairs: a nationwide follow-up study. *Diabetes* 52:1052–1055
- Harjutsalo V, Sjöberg L, Tuomilehto J (2008) Time trends in the incidence of type 1 diabetes in Finnish children: a cohort study. *Lancet* 371:1777–1782
- Maillard L (1912) Action des acides aminés sur les sucres: formation des melanoidines par voie. *CR Acad Sci* 154:66–68, article in French
- Henle T (2005) Protein-bound advanced glycation endproducts (AGEs) as bioactive amino acid derivatives in foods. *Amino Acids* 29:313–322
- Sell D, Monnier V (1989) Structure and elucidation of a senescence cross-link from human extracellular matrix. *J Biol Chem* 264:21597–21602
- Baynes JW, Thorpe SR (1999) Role of oxidative stress in diabetic complications: a new perspective on an old paradigm. *Diabetes* 48:1–9
- Miyata T, Wada Y, Cai Z et al (1997) Implication of an increased oxidative stress in the formation of advanced glycation end products in patients with end-stage renal failure. *Kidney Int* 51:1170–1181
- Brownlee M (2001) Biochemistry and molecular cell biology of diabetic complications. *Nature* 414:813–820
- Kaneto H, Fujii J, Myint T et al (1996) Reducing sugars trigger oxidative modification and apoptosis in pancreatic beta-cells by provoking oxidative stress through the glycation reaction. *Biochem J* 320:855–863
- Matsuoka T, Kajimoto Y, Watada H et al (1997) Glycation-dependent, reactive oxygen species-mediated suppression of the insulin gene promoter activity in HIT cells. *J Clin Invest* 99:144–150
- Peppas M, He C, Hattori M, McEvoy R, Zheng F, Vlassara H (2003) Fetal or neonatal low-glycotoxin environment prevents autoimmune diabetes in NOD mice. *Diabetes* 52:1441–1448
- Chavakis T, Bierhaus A, Al-Fakhri N et al (2003) The pattern recognition receptor (RAGE) is a counterreceptor for leukocyte integrins: a novel pathway for inflammatory cell recruitment. *J Exp Med* 198:1507–1515
- Bierhaus A, Illmer T, Kasper M et al (1997) Advanced glycation end product (AGE)-mediated induction of tissue factor in cultured endothelial cells is dependent on RAGE. *Circulation* 96:2262–2271
- Hofmann MA, Drury S, Fu C et al (1999) RAGE mediates a novel proinflammatory axis: a central cell surface receptor for S100/calgranulin polypeptides. *Cell* 97:889–901
- Chen Y, Yan SS, Colgan J et al (2004) Blockade of late stages of autoimmune diabetes by inhibition of the receptor for advanced glycation end products. *J Immunol* 173:1399–1405
- Chen Y, Akirav EM, Chen W et al (2008) RAGE ligation affects T cell activation and controls T cell differentiation. *J Immunol* 181:4272–4278
- Han J, Zhong J, Wei W et al (2008) Extracellular high-mobility group box 1 acts as an innate immune mediator to enhance autoimmune progression and diabetes onset in NOD mice. *Diabetes* 57:2118–2127
- Sullivan CM, Futers TS, Barrett JH, Hudson BI, Freeman MS, Grant PJ (2005) RAGE polymorphisms and the heritability of insulin resistance: the Leeds family study. *Diab Vasc Dis Res* 2:42–44
- Wilkin TJ (2001) The accelerator hypothesis: weight gain as the missing link between type I and type II diabetes. *Diabetologia* 44:914–922
- Groop PH, Thomas MC, Moran JL et al (2009) The presence and severity of chronic kidney disease predicts all-cause mortality in type 1 diabetes. *Diabetes* 58:1651–1658
- The MHC sequencing consortium (1999) Complete sequence and gene map of a human major histocompatibility complex. *Nature* 401:921–923
- Hermann R, Turpeinen H, Laine AP et al (2003) HLA DR-DQ-encoded genetic determinants of childhood-onset type 1 diabetes in Finland: an analysis of 622 nuclear families. *Tissue Antigens* 62:162–169
- Kukko M, Virtanen SM, Toivonen A et al (2004) Geographical variation in risk HLA-DQB1 genotypes for type 1 diabetes and signs of beta-cell autoimmunity in a high-incidence country. *Diab Care* 27:676–681
- Vandenplas S, Wiid I, Grobler-Rabie A et al (1984) Blot hybridisation analysis of genomic DNA. *J Med Genet* 21:164–172
- Gabriel SB, Schaffner SF, Nguyen H et al (2002) The structure of haplotype blocks in the human genome. *Science* 296:2225–2229
- Barrett JC, Fry B, Maller J, Daly MJ (2005) Haploview: analysis and visualization of LD and haplotype maps. *Bioinformatics* 21:263–265
- Purcell S, Cherny SS, Sham PC (2003) Genetic Power Calculator: design of linkage and association genetic mapping studies of complex traits. *Bioinformatics* 19:149–150
- Hattori M, Buse JB, Jackson RA et al (1986) The NOD mouse: recessive diabetogenic gene in the major histocompatibility complex. *Science* 231:733–735
- Sugaya K, Fukagawa T, Matsumoto K et al (1994) Three genes in the human MHC class III region near the junction with the class II: gene for receptor of advanced glycosylation end products, PBX2 homeobox gene and a notch homolog, human

- counterpart of mouse mammary tumor gene int-3. *Genomics* 23:408–419
33. Forbes JM, Thallas V, Thomas MC et al (2003) The breakdown of preexisting advanced glycation end products is associated with reduced renal fibrosis in experimental diabetes. *FASEB J* 17:1762–1764
 34. van der Pijl JW, Daha MR, van den Born J et al (1998) Extracellular matrix in human diabetic nephropathy: reduced expression of heparan sulphate in skin basement membrane. *Diabetologia* 41:791–798
 35. Candido R, Forbes JM, Thomas MC et al (2003) A breaker of advanced glycation end products attenuates diabetes-induced myocardial structural changes. *Circ Res* 92:785–792
 36. Kebede M, Favalaro J, Gunton JE et al (2008) Fructose-1, 6-bisphosphatase overexpression in pancreatic beta-cells results in reduced insulin secretion: a new mechanism for fat-induced impairment of beta-cell function. *Diabetes* 57:1887–1895
 37. Kooptiwut S, Kebede M, Zraika S et al (2005) High glucose-induced impairment in insulin secretion is associated with reduction in islet glucokinase in a mouse model of susceptibility to islet dysfunction. *J Mol Endocrinol* 35:39–48
 38. Pettersson-Fernholm K, Forsblom C, Hudson BI, Perola M, Grant PJ, Groop PH (2003) The functional -374 T/A RAGE gene polymorphism is associated with proteinuria and cardiovascular disease in type 1 diabetic patients. *Diabetes* 52:891–894
 39. Nanto-Salonen K, Kupila A, Simell S et al (2008) Nasal insulin to prevent type 1 diabetes in children with HLA genotypes and autoantibodies conferring increased risk of disease: a double-blind, randomised controlled trial. *Lancet* 372:1746–1755
 40. Forbes JM, Thorpe SR, Thallas-Bonke V et al (2005) Modulation of soluble receptor for advanced glycation end products by angiotensin-converting enzyme-1 inhibition in diabetic nephropathy. *J Am Soc Nephrol* 16:2363–2372
 41. Reddy S, Bibby N, Elliott RB (1990) Longitudinal study of islet cell antibodies and insulin autoantibodies and development of diabetes in non-obese diabetic (NOD) mice. *Clin Exp Immunol* 81:400–405
 42. Wittkowski H, Hirono K, Ichida F et al (2007) Acute Kawasaki disease is associated with reverse regulation of soluble receptor for advanced glycation end products and its proinflammatory ligand S100A12. *Arthritis Rheum* 56:4174–4181
 43. Gaens KH, Ferreira I, van der Kallen CJ et al (2009) Association of polymorphism in the receptor for advanced glycation end products (RAGE) gene with circulating RAGE levels. *J Clin Endocrinol Metab* 94:5174–5180
 44. Jang Y, Kim JY, Kang SM et al (2007) Association of the Gly82Ser polymorphism in the receptor for advanced glycation end products (RAGE) gene with circulating levels of soluble RAGE and inflammatory markers in nondiabetic and nonobese Koreans. *Metabolism* 56:199–205
 45. Zhang L, Bukulin M, Kojro E et al (2008) Receptor for advanced glycation end products is subjected to protein ectodomain shedding by metalloproteinases. *J Biol Chem* 283:35507–35516
 46. Yan SD, Zhu H, Zhu A et al (2000) Receptor-dependent cell stress and amyloid accumulation in systemic amyloidosis. *Nat Med* 6:643–651
 47. Shikama Y, Kitazawa J, Yagihashi N et al (2010) Localized amyloidosis at the site of repeated insulin injection in a diabetic patient. *Intern Med* 49:397–401
 48. Damiani G, Campo I, Zorzetto M et al (2007) Pro-inflammatory variants of DRB1 and RAGE genes are associated with susceptibility to pediatric type 1 diabetes: a new hypothesis on the adaptive role of autoimmunity. *Riv Biol* 100:285–304
 49. Forbes JM, Soulis T, Thallas V et al (2001) Renoprotective effects of a novel inhibitor of advanced glycation. *Diabetologia* 44:108–114
 50. Fu MX, Wells-Knecht KJ, Blackledge JA, Lyons TJ, Thorpe SR, Baynes JW (1994) Glycation, glycooxidation, and cross-linking of collagen by glucose. Kinetics, mechanisms, and inhibition of late stages of the Maillard reaction. *Diabetes* 43:676–683
 51. Fourlanos S, Varney MD, Tait BD et al (2008) The rising incidence of type 1 diabetes is accounted for by cases with lower-risk human leukocyte antigen genotypes. *Diab Care* 31:1546–1549