

**RECOMMENDED NOMENCLATURE FOR ZEOLITE MINERALS: REPORT OF THE  
SUBCOMMITTEE ON ZEOLITES OF THE INTERNATIONAL MINERALOGICAL  
ASSOCIATION, COMMISSION ON NEW MINERALS AND MINERAL NAMES**

**DOUGLAS S. COOMBS<sup>1</sup> (Chairman)**

*Geology Department, University of Otago, P.O. Box 56, Dunedin, New Zealand*

**ALBERTO ALBERTI**

*Istituto di Mineralogia, Università di Ferrara, Corso Ercole I° d'Este, 32, I-44100 Ferrara, Italy*

**THOMAS ARMBRUSTER**

*Laboratorium für chemische und mineralogische Kristallographie, Universität Bern, Freiestrasse 3, CH-3012 Bern, Switzerland*

**GILBERTO ARTIOLI**

*Dipartimento di Scienze della Terra, Università di Milano, via Botticelli, 23, I-20133 Milano, Italy*

**CARMINE COLELLA**

*Dipartimento di Ingegneria dei Materiali e della Produzione, Università Federico II di Napoli, Piazzale V. Tecchio, 80, I-80125 Napoli, Italy*

**ERMANNO GALLI**

*Dipartimento di Scienze della Terra, Università di Modena, via S. Eufemia, 19, I-41100 Modena, Italy*

**JOEL D. GRICE**

*Mineral Sciences Division, Canadian Museum of Nature, Ottawa, Ontario K1P 6P4, Canada*

**FRIEDRICH LIEBAU**

*Mineralogisch-Petrographisches Institut, Universität Kiel, Olshausenstrasse 40-60, D-2300 Kiel, Germany*

**JOSEPH A. MANDARINO (retired from Subcommittee, December, 1994)**

*Department of Mineralogy, Royal Ontario Museum, Toronto, Ontario M5S 2C6 Canada*

**HIDEO MINATO**

*5-37-17 Kugayama, Sugiyama-ku, Tokyo 168, Japan*

**ERNEST H. NICKEL**

*Division of Exploration and Mining, CSIRO, Private Bag, Wembley 6014, Western Australia, Australia*

**ELIO PASSAGLIA**

*Dipartimento di Scienze della Terra, Università di Modena, via S. Eufemia, 19, I-41100 Modena, Italy*

**DONALD R. PEACOR**

*Department of Geological Sciences, University of Michigan, Ann Arbor, Michigan 48109, U.S.A.*

**SIMONA QUARTIERI**

*Dipartimento di Scienze della Terra, Università di Modena, via S. Eufemia, 19, I-41100 Modena, Italy*

**ROMANO RINALDI**

*Dipartimento di Scienze della Terra, Università di Perugia, I-06100 Perugia, Italy*

**MALCOLM ROSS**

*U.S. Geological Survey, MS 954, Reston, Virginia 20192, U.S.A.*

**RICHARD A. SHEPPARD**

*U.S. Geological Survey, MS 939, Box 25046, Federal Center, Denver, Colorado 80225, U.S.A.*

**EKKEHART TILLMANN**

*Institut für Mineralogie und Kristallographie, Universität Wien, Althanstrasse 14, A-1090 Vienna, Austria*

**GIOVANNA VEZZALINI**

*Dipartimento di Scienze della Terra, Università di Modena, via S. Eufemia, 19, I-41100 Modena Italy*

<sup>1</sup> E-mail address: doug.coombs@stonebow.otago.ac.nz

## ABSTRACT

This report embodies recommendations on zeolite nomenclature approved by the International Mineralogical Association, Commission on New Minerals and Mineral Names. In a working definition of a zeolite mineral used for this review, structures containing an interrupted framework of tetrahedra are accepted where other zeolitic properties prevail, and complete substitution by elements other than Si and Al is allowed. Separate species are recognized in topologically distinctive compositional series in which different extra-framework cations are the most abundant in atomic proportions. To name these, the appropriate chemical symbol is attached by a hyphen to the series name as a suffix, except for the names harmotome, pollucite and wairakite in the phillipsite and analcime series. Differences in space-group symmetry and in order-disorder relationships in zeolites having the same topologically distinctive framework do not in general provide adequate grounds for recognition of separate species. Zeolite species are not to be distinguished solely on the ratio Si : Al except for heulandite (Si : Al < 4.0) and clinoptilolite (Si : Al  $\geq$  4.0). Dehydration, partial hydration, and overhydration are not sufficient grounds for the recognition of separate species of zeolites. Use of the term "ideal formula" should be avoided in referring to a simplified or averaged formula of a zeolite. Newly recognized species in compositional series are as follows: brewsterite-Sr, -Ba, chabazite-Ca, -Na, -K, clinoptilolite-K, -Na, -Ca, dachiardite-Ca, -Na, erionite-Na, -K, -Ca, faujasite-Na, -Ca, -Mg, ferrierite-Mg, -K, -Na, gmelinite-Na, -Ca, -K, heulandite-Ca, -Na, -K, -Sr, levyne-Ca, -Na, paulingite-K, -Ca, phillipsite-Na, -Ca, -K, and stilbite-Ca, -Na. Key references, type locality, origin of name, chemical data, IZA structure-type symbols, space-group symmetry, unit-cell dimensions, and comments on structure are listed for 13 compositional series, 82 accepted zeolite mineral species, and three of doubtful status. Herschelite, leonhardite, svetlozarite, and wellsite are discredited as mineral species names. Obsolete and discredited names are listed.

**Keywords:** zeolite nomenclature, herschelite, leonhardite, svetlozarite, wellsite, brewsterite, chabazite, clinoptilolite, dachiardite, erionite, faujasite, ferrierite, gmelinite, heulandite, levyne, paulingite, phillipsite, stilbite.

## SOMMAIRE

Ce rapport contient les recommandations à propos de la nomenclature des zéolites, telles qu'approuvées par l'Association minéralogique internationale, commission des nouveaux minéraux et des noms de minéraux. Dans la définition d'une zéolite retenue ici, les structures contenant une trame interrompue de tétraèdres sont acceptées dans les cas où les autres propriétés satisfont les critères de cette famille de minéraux. De plus, il peut y avoir remplacement complet de Si et Al par d'autres éléments. Des espèces distinctes font partie de séries de compositions dont l'agencement topologique est le même, le cation dominant ne faisant pas partie de la trame déterminant l'espèce. Pour en déterminer le nom, il s'agit de rattacher le symbole chimique approprié au nom de la série par un trait d'union, sauf dans les cas de harmotome, pollucite et wairakite, faisant partie des séries de la phillipsite et de l'analcime. Des différences en symétrie exprimées par le groupe spatial et en degré d'ordre Si-Al dans les zéolites ayant le même agencement topologique ne suffisent pas en général pour définir une espèce distincte. Le seul critère de rapport Si : Al ne suffit pas pour distinguer les espèces de zéolites, sauf pour la heulandite (Si : Al < 4.0) et clinoptilolite (Si : Al  $\geq$  4.0). L'état de déshydratation, d'hydratation partielle, et de sur-hydratation ne suffit pas pour reconnaître une espèce distincte de zéolite. On doit éviter d'utiliser le concept d'une "formule idéale" en parlant des formules simplifiées ou représentatives des zéolites. Les nouveaux noms d'espèces dans ces séries de compositions sont: brewsterite-Sr, -Ba, chabazite-Ca, -Na, -K, clinoptilolite-K, -Na, -Ca, dachiardite-Ca, -Na, ériomite-Na, -K, -Ca, faujasite-Na, -Ca, -Mg, ferrierite-Mg, -K, -Na, gmelinite-Na, -Ca, -K, heulandite-Ca, -Na, -K, -Sr, lévyne-Ca, -Na, paulingite-K, -Ca, phillipsite-Na, -Ca, -K, et stilbite-Ca, -Na. Nous présentons les références-clés, la localité-type, l'origine du nom, des données chimiques, le symbole structural IZA, le groupe spatial et la symétrie, les paramètres réticulaires et des commentaires sur la structure de 13 séries compositionnelles, 82 espèces homologuées, et trois dont le statut est douteux. Herschelite, léonhardite, svetlozarite, et wellsite sont discréditées comme noms d'espèces minérales. Nous dressons une liste des noms obsolètes et discrédités.

(Traduit par la Rédaction)

**Mots-clés:** nomenclature des zéolites, herschelite, léonhardite, svetlozarite, wellsite, brewsterite, chabazite, clinoptilolite, dachiardite, ériomite, faujasite, ferrierite, gmelinite, heulandite, lévyne, paulingite, phillipsite, stilbite.

## INTRODUCTION

The name "zeolite" was introduced by the Swedish mineralogist Cronstedt in 1756 for certain silicate minerals in allusion to their behavior on heating in a borax bead (Greek *zeo* = boil; *lithos* = stone). Three such minerals were listed by Haty (1801), namely stilbite, analcime, and harmotome, together with "mesotype", which has not survived. Chabazite and leucite had been named even earlier. Nineteen had been

described with their present meaning by 1842. Forty-six zeolites were listed by Gottardi & Galli (1985), and new species continue to be described. The first crystal-structure determination of a zeolite was done on analcime (Taylor 1930); following this, Hey (1930) concluded that zeolites in general have aluminosilicate frameworks with loosely bonded alkali or alkali-earth cations, or both. Molecules of H<sub>2</sub>O occupy extra-framework positions. He pointed out the consequential requirements that the molar ratio

$\text{Al}_2\text{O}_3 : (\text{Ca}, \text{Sr}, \text{Ba}, \text{Na}_2, \text{K}_2)\text{O} = 1$  and that  $\text{O} : (\text{Si} + \text{Al}) = 2$  in the empirical formula.

Zeolites have other highly characteristic features developed to varying degrees, notably the potential for reversible low-temperature dehydration, the ability of the dehydrated forms to reversibly absorb other molecules, a tendency toward more or less easy low-temperature exchange of extra-framework cations, and a lack of clear-cut, structurally controlled constraints on end-member compositions in terms of  $\text{Si} : \text{Al}$  ratios within the framework. In some cases, observed extra-framework compositions may be artefacts of cation exchange resulting from human activities in the laboratory or elsewhere, and furthermore, the compositions are not conveniently determined by traditional optical methods. Perhaps for a combination of such reasons, separate names have been given to few zeolites on the basis of the dominant extra-framework cation in solid-solution series. This conflicts with standard practice in most mineral groups and with guidelines of the Commission on New Minerals and Mineral Names (CNMMN) (Nickel & Mandarino 1987).

With intensification of research and the advent of the electron microprobe, a flood of information on compositions has become available, and with automated single-crystal X-ray diffractometers and other developments, many complexities have been investigated, including order-disorder relationships in the frameworks and associated changes in unit-cell parameters and symmetry. Thus in the case of analcime, Mazzi & Galli (1978), Teertstra *et al.* (1994), and others have demonstrated a wide range of space-group symmetries associated with different patterns of order in the framework and possible displacive transformations. Sites of extra-framework cations are commonly less well defined in an open, zeolitic structure than in most other minerals, and are variably occupied. Guidelines allowing recognition of separate species depending on the dominant ion occupying each structural site are thus compromised in the case of extra-framework sites in zeolites. Furthermore, changes in the occupancy of such sites can distort the framework to varying degrees, changing the space-group symmetry.

Some minerals meet traditional criteria for zeolites in all respects except that they contain P, Be, or other elements in tetrahedral sites, with consequent departure from the requirement of Hey (1930) that  $\text{O} : (\text{Si} + \text{Al}) = 2$ . Other structurally related minerals with zeolitic properties have all tetrahedral sites occupied by elements other than Si and Al. Certain other minerals displaying zeolitic properties depart from traditional requirements for a zeolite in having a framework that is interrupted by some (OH) groups. An example is parthéite, listed by Gottardi & Galli (1985) as a zeolite. Synthesis and structural analysis of materials having zeolitic properties have become major fields of research and have led to a voluminous literature, as has the industrial use of zeolitic materials.

The recommendations of an IMA CNMMN subcommittee set up to review zeolite nomenclature are set out below. These recommendations have been adopted by the Commission.

#### DEFINITION OF A ZEOLITE MINERAL

In arriving at its working definition of a zeolite, the Subcommittee took the view that zeolites in the historical and mineralogical sense are naturally occurring minerals, irrespective of how the term may be applied to synthetic materials and in industry. In the light of advances in mineralogy, the Hey (1930) definition is found to be too restrictive. The Subcommittee gave particular consideration to the following questions. Is more than 50% substitution of elements other than Si and Al permissible in tetrahedral sites? Is the presence of  $\text{H}_2\text{O}$  and of extra-framework cations absolutely essential? Can "interrupted" framework structures qualify as zeolite minerals? These matters are further discussed in Appendix 1.

#### Definition

*A zeolite mineral is a crystalline substance with a structure characterized by a framework of linked tetrahedra, each consisting of four O atoms surrounding a cation. This framework contains open cavities in the form of channels and cages. These are usually occupied by  $\text{H}_2\text{O}$  molecules and extra-framework cations that are commonly exchangeable. The channels are large enough to allow the passage of guest species. In the hydrated phases, dehydration occurs at temperatures mostly below about 400°C and is largely reversible. The framework may be interrupted by (OH,F) groups; these occupy a tetrahedron apex that is not shared with adjacent tetrahedra.*

*Application of the definition (see also Appendix 1)*

Relatively easy exchange of extra-framework cations at relatively low temperature is a characteristic feature of zeolites and zeolitic behavior, but varies greatly from species to species. Its extent does not provide a convenient basis for the definition of zeolites. In practice, it appears that channels must have a minimum width greater than that of 6-membered rings (*i.e.*, rings consisting of six tetrahedra) in order to allow zeolitic behavior at normal temperatures and pressures. Framework structures such as in feldspars, nepheline, sodalites, scapolites, melanophlogite, and probably leifite, in which any channels are too restricted to allow typical zeolitic behavior such as reversible dehydration, molecular sieving, or cation exchange, are not regarded as zeolites.

Framework density, defined as the number of tetrahedral sites in  $1000 \text{ \AA}^3$ , was used as the criterion for inclusion in the *Atlas of Zeolite Structure Types*

(Meier *et al.* 1996). However, this criterion provides no evidence that the channels necessary for diffusion are present, as well as cages, and it has not been adopted in the present definition.

In some minerals with a tetrahedral framework structure and other zeolitic characteristics as described, namely parthéite, roggianite, maricopaite, and chiavennite, one apex of some tetrahedra is occupied by an (OH) group or F atom instead of being occupied by an O atom. This (OH) group or F atom does not form a bridge with an adjacent tetrahedron. The framework is thus interrupted. Such minerals are here accepted as zeolites.

In terms of the definition adopted, minerals of the cancrinite group can arguably be considered as zeolites. This group has long been regarded by many or most mineralogists as distinct from the zeolites, in part, at least, because of the presence of large volatile anions (*e.g.*, Hassan 1997). They are not reviewed in the present report. Rather similarly, wenkite contains large cages and channels, but these are blocked by  $\text{SO}_4$ , Ca, and Ba ions (Wenk 1973, Merlino 1974), inhibiting zeolitic behavior. In addition, no water is lost below 500°C. Wenkite is not included as a zeolite in this report.

Leucite has seldom been regarded as a zeolite, as it does not display a full range of zeolitic behavior. Nevertheless, it has the same framework structure as analcime and conforms to the adopted definition. Ammonioleucite can be regarded as an analcime derivative, can be synthesized from analcime by cation exchange, and may have formed naturally by low-temperature replacement of analcime. Leucite and ammonioleucite are included in the list of zeolites, as is kalborsite, a derivative of the edingtonite structure.

Also conforming to the definition adopted are the beryllophosphates pahasapaite and weinebeneite. These contain neither Si nor Al and can be regarded as end-member examples of Si-free zeolites or zeolite phosphates.

#### RULES FOR NOMENCLATURE OF ZEOLITE MINERALS

In presenting the following rules for nomenclature of zeolite minerals, the Subcommittee feels strongly that they should be viewed as guidelines rather than as being rigidly prescriptive. As stated by Nickel & Mandarino (1987): "It is probably not desirable to formulate rigid rules to define whether or not a compositional or crystallographic difference is sufficiently large to require a new mineral name, and each new mineral proposal must be considered on its own merits". Explanatory notes following the proposed rules or guidelines give examples of how the Subcommittee envisages that rule being applied, but like Nickel and Mandarino, the Subcommittee urges that each case be treated on its merits. In some cases, compelling reasons may exist on grounds of historical usage for retaining an existing name, or other grounds may exist

for departing from the rules for giving a new name. Cases arising under Rule 2 are particularly difficult, and require individual consideration.

#### Rule 1

(a) One or more zeolite minerals having a topologically distinctive framework of tetrahedra, and a composition that is distinctive for zeolites having that framework, constitute separate *species*. (b) Zeolites having the same topologically distinctive framework of tetrahedra constitute a *series* when they display a substantial range in composition in which differing extra-framework cations may be the most abundant in atomic proportions. These cations may occupy different extra-framework sites. Such *series* consist of two or more *species* that are distinguished on the basis of the most abundant extra-framework cation.

#### Application of the rule

Laumontite, for example, has a topologically distinctive framework and a composition which, as far as is currently known, is distinctive in that Ca is always the dominant extra-framework cation. It is a separate zeolite species under Rule 1a. Natrolite, mesolite, and scolecite have the same topologically distinctive framework structure as each other, and have compositions that are distinctive. They also are separate species under Rule 1a.

Zeolites having the topologically distinctive chabazite structure have a range of compositions in which any one of Ca, Na, or K may be the most abundant extra-framework cation. Substantial Sr is in some cases present as well, but so far has never been reported as the most abundant in natural examples. Chabazite is a series consisting of three separate species under Rule 1b. It is known that near-end-member Na, K, Ca, and Sr compositions are readily obtainable by ion exchange from natural Ca-dominant chabazite at 110°C (Alberti *et al.* 1982a), but this is not the essential criterion for recognition of the natural series.

Mesolite may have either Na or Ca slightly in excess of the other, but the ratio Na : Ca is always close to 1 : 1. The range of its composition is not regarded as "substantial", and mesolite is not divided into more than one species on grounds of composition.

Several distinct structural sites for extra-framework cations are recognized in many zeolites, but in view of the relatively loose bonding and specialized problems in establishing the individual site-occupancies, only the total population of extra-framework cations should in general be used in defining zeolite species.

#### Rule 2

(a) Differences in space-group symmetry and in order-disorder relationships in zeolite minerals having the same topologically distinctive framework do not in

general provide adequate grounds for recognition of separate species, but each case should be treated on its merits. (b) In assessing such cases, other factors, such as relationship to chemical composition, should be taken into consideration.

#### *Application of the rule*

The Subcommittee found it to be impracticable to formulate quantified criteria for handling problems arising from this rule. Irrespective of decisions that have been made in the past, care should be taken that departures envisaged in Rule 2b from the principle enunciated in Rule 2a are based on grounds that are truly compelling.

Analcime and certain other zeolites exist with several different space-group symmetries, in some cases occurring on a very fine scale in the same hand specimen and with the same chemistry. Even though this may be related to Si,Al ordering, separate species names in these cases are in general not warranted.

Gismondine and garronite are examples of zeolites that have the same topologically distinctive framework. Both have Ca as the dominant extra-framework cation. Their differing space-group symmetry is associated with disordered Si,Al and the presence of significant Na in garronite. They are accepted as separate species. Gobbinsite and amicite have topologically the same framework structure as gismondine, but are alkali-dominant. Their different space-group symmetries appear to be related to Si,Al disorder in gobbinsite and possible chemical differences, and they are provisionally retained. Barrerite is topologically similar to stilbite and stellerite, but it has different symmetry correlated with the presence of extra cations that cause rotational displacements within the framework (Galli & Alberti 1975b); it is similarly retained.

#### **Rule 3**

Zeolite mineral species shall not be distinguished solely on the basis of the framework Si : Al ratio. An exception is made in the case of heulandite and clinoptilolite; heulandite is defined as the zeolite mineral series having the distinctive framework topology of heulandite and the ratio Si : Al < 4.0. Clinoptilolite is defined as the series with the same framework topology and Si : Al ≥ 4.0.

#### *Application of the rule*

Many zeolites have a widely variable Si : Al ratio, but this, in itself, is not regarded as providing adequate grounds for recognition of separate species. The exception is based on entrenched usage of the names *heulandite* and *clinoptilolite*, and their convenience for recognizing an important chemical feature. The cutoff value adopted (following Boles 1972) is arbitrary in a

continuous range of compositions. The usual 50% compositional rule cannot be applied, as there are no clearly defined Si,Al end-member compositions for heulandite and clinoptilolite. Thermal stability has been used by some investigators to distinguish clinoptilolite from heulandite. This is a derivative property, however, suggested by Mumpton (1960) as an aid to identification, and it is not appropriate as the basis for definition. Alietti (1972) and Boles (1972) have shown that there is no gap in composition either in framework or extra-framework cation contents between heulandite and clinoptilolite, and that samples transitional in composition show intermediate properties in terms of thermal stability.

#### **Rule 4**

Dehydration, partial hydration, and overhydration, whether reversible or irreversible, are not sufficient grounds for the recognition of separate species of zeolite minerals.

#### *Application of the rule*

If a new topologically distinctive framework arises from overhydration or partial dehydration, separate species status would result from application of Rule 1. Leonhardite, a partially and in most cases reversibly dehydrated form of laumontite, is not accepted as a separate mineral species.

#### **Rule 5**

Individual species in a zeolite mineral series with varying extra-framework cations are named by attaching to the series name a suffix indicating the chemical symbol for the extra-framework element that is most abundant in atomic proportions, *e.g.*, chabazite-Ca.

The following exceptions are made: a) On grounds of historical precedence and long-established usage, the name *harmotome* is retained for the Ba-dominant member of the phillipsite series. b) On grounds of long-established usage, pollucite is retained as the Cs-dominant zeolite of the analcime structure-type. On grounds of established usage and markedly different space-group symmetry and Si,Al order related to the extra-framework cation content (Rule 2b), wairakite is retained as the Ca-dominant zeolite of the analcime structure-type. On the other hand, herschelite is suppressed in favor of chabazite-Na (Appendix 2).

#### *Application of the rule*

New species arising from Rule 5 that are well authenticated by published data are set out in Table 1. Future proposals for additional new species under this rule should be dealt with as for any other proposal for a new mineral name.

Adoption of a Levinson-style system of suffixes avoids the proliferation of a large number of new and potentially unrelated species names, and ensures that all members of a topologically identical compositional series are indexed together. It has the great advantage that where adequate chemical data are not required or are not available, a mineral can be referred to correctly by an unambiguous series name. The system adopted here is without the brackets (parentheses) used by Levinson (1966) in suffixes for rare-earth minerals.

Substantial amounts of extra-framework cations other than the dominant one may be indicated, if desired, by the use of adjectives such as calcian and sodian, *e.g.*, calcian clinoptilolite-K. Such adjectival modifiers are not part of the formal name of a species.

Informal use is often made of descriptive terms such as calcium chabazite and Ca chabazite, in which the name or symbol of an element is used adjectivally. In conformity with general IMA guidelines, these should not appear in print as mineral names or in hyphenated form. The correct name for the mineral species in this case is chabazite-Ca. Terms such as sodium-substituted chabazite-Ca are suggested for what in effect would be a synthetic chabazite-Na prepared by cation exchange from chabazite-Ca. Chabazite remains the correct name for a member of the chabazite series that is not specifically identified on compositional grounds.

## Rule 6

(a) Space-group variants of zeolite mineral species may be indicated by placing the space-group symbol in round brackets (parentheses) after the mineral species name, *e.g.*, analcime (*Ibca*), heulandite-Ca (*C2/m*). (b) Levels of order may be indicated by adjectival use of words such as "disordered" or "fully ordered" before the mineral name.

### Application of the rule

Modifiers as suggested here are not part of the formal name of the mineral.

### ACCEPTED ZEOLITE SERIES AND SPECIES

Zeolites to be elevated to series status and the consequential new species to be recognized on the basis of the most abundant extra-framework cation (Rule 5) are set out in Table 1.

An annotated list of accepted zeolite series and species follows below. In each entry for series, and for those species that are not members of compositional series, a simplified or generalized formula is given in the first line. This is followed by *Z*, the number of these formula units per unit cell, as given later in the entry. The simplified or generalized formula should be regarded as representative only, and should not be regarded as an "ideal" composition (see next

TABLE 1. NEWLY PROPOSED ZEOLITE SPECIES WITHIN COMPOSITIONAL SERIES<sup>§</sup>

Series	Species name	Series	Species name
brewsterite	brewsterite-Sr	gmelinite	gmelinite-Na
	brewsterite-Ba		gmelinite-Ca
chabazite	chabazite-Ca	heulandite	heulandite-Ca
	chabazite-Na		heulandite-Na
	chabazite-K		heulandite-K
clinoptilolite	clinoptilolite-K	levyne	levyne-Ca
	clinoptilolite-Na		levyne-Na
	clinoptilolite-Ca		
dachardite	dachardite-Ca	paulingite	paulingite-K
	dachardite-Na		paulingite-Ca
erionite	erionite-Na	phillipsite	phillipsite-Na
	erionite-K		phillipsite-Ca
	erionite-Ca		phillipsite-K
faujasite	faujasite-Na	stilbite	stilbite-Ca
	faujasite-Ca		stilbite-Na
	faujasite-Mg		
ferrierite	ferrierite-Mg		
	ferrierite-K		
	ferrierite-Na		

<sup>§</sup> The first-named member of each series is the one to which the original type-specimen for the series seems to belong.

paragraph). Users of the list should bear in mind that the Si : Al ratio, or, more generally, occupancy of tetrahedral sites by Si, Al, P, Be, Zn, and possibly other elements, varies widely in many zeolites. The total extra-framework cation charge varies accordingly. Major variation in more-or-less exchangeable, extra-framework cations is also a feature of many natural zeolites. Contents of H<sub>2</sub>O tend to decrease with increasing number and size of extra-framework cations, as well as with increasing temperature and decreasing P(H<sub>2</sub>O). Such variations can be vital to petrological, geochemical, environmental, and experimental considerations.

Simplified or generalized formulae of zeolites, *e.g.*, NaAlSi<sub>2</sub>O<sub>6</sub>·H<sub>2</sub>O for analcime, have often been referred to as "ideal" formulae. However, the supposed ideality may be in writers' desire for simplicity, rather than in anything fundamental to the zeolites concerned, and can lead to false assumptions. There is much evidence that the composition of naturally occurring analcime is a function of the chemical environment in which it forms (*e.g.*, Coombs & Whetten 1967). In environments of low Si activity, as in altered strongly silica-deficient

alkaline rocks, natural analcime approaches a Si : Al ratio of 1.5. The composition in burial metamorphic rocks in equilibrium with quartz appears to be distinctly more Si-rich than the supposed "ideal" Si : Al value of 2. The evidently metastable equilibrium in natural environments containing siliceous volcanic glass or other source of silica yielding higher activity of Si than coexistence with quartz, leads to analcime with Si : Al approaching 3. Analogous observations apply to heulandite and other zeolites. If "ideal" is taken to imply equilibrium, it can therefore be concluded that this is a function of the chemical (and P-T) environment during crystallization, rather than simply being a function of crystal structure. Differing Si : Al ratios may in turn favor different patterns of order in the framework. Application of the term "ideal" to simplified or averaged formulae of zeolites should be avoided.

Also given in the first line of each entry is the structure-type code allocated by the Structure Commission of the International Zeolite Association (IZA) and listed in Meier *et al.* (1996). The code consists of three capital letters. A preceding hyphen indicates an interrupted framework of tetrahedra.

The second line of each relevant entry starts with the original reference in which the current name of the mineral, or a near variant of that name, is given, followed by the type locality, or, in the case of descriptions that predate the concept of type localities, the general region of origin of the material on which the name and original description are based, where this is known. The locality is followed by a note on the derivation of the name. Further information on these matters is given by Gottardi & Galli (1985), Clark (1993), and Blackburn & Dennen (1997), but in some cases the information is here revised.

Next is given information on the currently known range in composition of the mineral concerned. This includes known values, or range of values, for  $T_{Si}$ , the proportion of tetrahedron sites occupied by Si atoms, as reported in published results of acceptable analyses. For many zeolites, this value varies widely, and the values reported may not indicate the full range possible, especially in the case of the rarer zeolites.

Much information on zeolite compositions was given by Gottardi & Galli (1985). The present compilation incorporates results of further extensive searches of the literature. A widely used criterion for acceptability of zeolite compositions is that the value of the balance-error function of Passaglia (1970)


should be less than 10%, a figure that is itself arguably excessive. The calculation of E% may be modified to allow for other suspected cations, such as  $Fe^{2+}$  and  $Cs^+$ . The role of Fe causes problems that may not be resolvable. Some Fe reported in zeolites is undoubtedly a contaminant, but there are reasons to suspect that

both  $Fe^{2+}$  and  $Fe^{3+}$  may enter the structures of some zeolites in extra-framework or framework sites, or both.

Space-group symmetry and crystallographic parameters follow. Many accepted zeolite species exist with more than one known space-group symmetry, and these are listed. Variations in space-group symmetry and variations in order-disorder relationship of framework cations are not in themselves adequate evidence for establishing new species (Rule 2). Cell parameters given are as reported for material specified in key references. Cell dimensions of many species vary widely as a result of variable compositions, variable extent of order, and differing levels of hydration. Except for a few newly described species, details of structure, including size and orientation of channels, can be obtained for each structure type from Meier *et al.* (1996) and are discussed in Gottardi & Galli (1985).

The accepted series and species are as follows:

### Amicite



Alberti *et al.* (1979). Type locality: Höwenegg (a Tertiary melilite nephelinite volcano), Hegau, southwestern Germany. Named after Giovan Battista Amici (1786–1863), inventor of the Amici lens and microscope objectives with a hemispherical front lens.


Both type amicite and the only other known example (Khomyakov *et al.* 1982) include minor Ca.  $T_{Si} = 0.51, 0.49$ .

Monoclinic, *I2*, *a* 10.226(1), *b* 10.422(1), *c* 9.884(1) Å,  $\beta$  88.32(2)°.

The framework is characterized by double crankshaft chains as in gismondine (Alberti & Vezzalini 1979).

Amicite has the same framework topology as gismondine. Si, Al and Na, K distributions are ordered and lower the symmetry from topological *I4<sub>1</sub>/amd* to real symmetry *I2*.

### Ammonioleucite




Hori *et al.* (1986). Type locality: Tatarazawa, Fujioka, Gunma Prefecture, Japan. The name reflects its composition and relationship to leucite.

Material from the only known locality contains significant K.  $T_{Si} = 0.70$ .

Tetragonal, *I4<sub>1</sub>/a*, *a* 13.214(1), *c* 13.713(2) Å.

### Analcime



Haüy (1797, p. 278). Type locality: near Catanes, Cyclopean Isles, Italy (Haüy, 1801, p. 180–185). Name from Greek roots meaning "without strength", in

allusion to the weak electrical effects induced by friction. In most analyzed specimens, Na is the only substantial extra-framework cation, but analcime forms a continuous series with pollucite and possibly with wairakite (Seki & Oki 1969, Seki 1971, Cho & Liou 1987).  $T_{Si}$  varies widely, 0.59–0.73 (*e.g.*, Coombs & Whetten 1967). As Si increases, NaAl decreases and  $H_2O$  increases.

Topological symmetry is cubic,  $Ia\bar{3}d$ . Real symmetry variants include:

cubic,  $Ia\bar{3}d$ ,  $a$  13.725 Å;

tetragonal,  $I4_1/acd$ ,  $a$  13.723(7),  $c$  13.686(10) Å;  $a$  13.721(1),  $c$  13.735(1) Å (Mazzi & Galli 1978);

tetragonal,  $I4_1/a$ ;

orthorhombic,  $Ibca$ ,  $a$  13.733(1),  $b$  13.729(1),  $c$  13.712(1) Å;  $a$  13.727(2),  $b$  113.714(2),  $c$  13.740(2) Å (Mazzi & Galli 1978);

monoclinic with 2-fold axis parallel both to pseudo-cubic [100] and [110];

triclinic,  $a$  13.6824(5),  $b$  13.7044(6),  $c$  13.7063(5) Å,  $\alpha$  90.158(3)°,  $\beta$  89.569(3)°,  $\gamma$  89.543(3)° (Hazen & Finger 1979);

and probably trigonal with variable Si,Al order (*e.g.*, Hazen & Finger 1979, Teertstra *et al.* 1994).

The name applies to Na-dominant compositions with this framework structure regardless of the degree and patterns of order.

### Barrerite

$Na_2[Al_2Si_7O_{18}] \cdot 6H_2O$   $Z = 8$  STI

Passaglia & Pongiluppi (1974, 1975). Type locality: Capo Pula, Sardinia, Italy. Named after Professor Richard M. Barrer (1910–1996) of Imperial College, London, for contributions to the chemistry of molecular sieves.

Also known from Kuiu Island, Alaska (Di Renzo & Gabelica 1997).  $T_{Si} = 0.77, 0.78$ . The type example has composition:

$(Na_{2.45}K_{1.06}Ca_{0.84}Mg_{0.17})[Al_{8.19}Fe_{0.01}Si_{27.72}O_{72}] \cdot 25.78H_2O$ . Orthorhombic, *Amma* or *Ammm*,  $a$  13.643(2),  $b$  18.200(3),  $c$  17.842(3) Å (Passaglia & Pongiluppi 1974).

The structure is similar to that of stilbite and stellerite, but it has different symmetry as a result of extra cations, which cause rotational displacements within the framework (Galli & Alberti 1975b).

### Bellbergite

$(K,Ba,Sr)_2Sr_2Ca_2(Ca,Na)_4[Al_{18}Si_{18}O_{72}] \cdot 30H_2O$

$Z = 1$  EAB

Rüdinger *et al.* (1993). Type and only known locality: Bellberg (or Bellerberg) volcano, near Mayen, Eifel, Germany. Named after the locality.

Ca is overall the dominant extra-framework cation.  $T_{Si} = 0.51$ .

Hexagonal, possible space-groups  $P6_3/mmc$ ,  $\bar{P}62c$ , and  $P6_3mc$ ,  $a$  13.244(1),  $c$  15.988(2) Å.

The framework structure is as for synthetic zeolite TMA-EAB.

### Bikitaite

$Li[AlSi_2O_6] \cdot H_2O$   $Z = 2$  BIK

Hurlbut (1957). Type locality: Bikita, Zimbabwe. Named after the type locality.

Two known localities, with the bikitaite having very similar compositions.  $T_{Si} = 0.67$ .

Monoclinic,  $P2_1$ ,  $a$  8.613(4),  $b$  4.962(2),  $c$  7.600(4) Å,  $\beta$  114.45(1)° (Kocman *et al.* 1974).

Also triclinic,  $P1$ ,  $a$  8.606(1),  $b$  4.953(1),  $c$  7.599(1) Å,  $\alpha$  89.89(2)°,  $\beta$  114.42(2)°,  $\gamma$  89.96(2)° (Bissert & Liebau 1986).

The framework structure consists of 5-membered rings linked by additional tetrahedra. Its topological symmetry is  $P2_1$ . The monoclinic  $P2_1$  variant of Kocman *et al.* has partly ordered Si,Al distribution; the triclinic  $P1$  variant of Bissert and Liebau is highly ordered.

### Boggsite

$Ca_8Na_3[Al_{19}Si_{77}O_{192}] \cdot 70H_2O$   $Z = 1$  BOG

Pluth *et al.* (1989) and Howard *et al.* (1990). Type locality: Basalt above cliff, Goble Creek, south side of the Neer Road, 0.2 km north of Goble, Columbia County, Oregon, U.S.A. Named after Robert Maxwell Boggs (father) and Russell Calvin Boggs (son), mineral collectors in the Pacific Northwest.

Type boggsite approximates the above formula, with minor Fe, Mg, and K. Boggsite from Mt. Adamson, Antarctica (Galli *et al.* 1995) approximates  $Ca_6Na_5K[Al_{18}Si_{78}O_{192}] \cdot 70H_2O$ , with minor Fe, Mg, Sr, Ba.  $T_{Si} = 0.81$ .

Orthorhombic, *Imma*,  $a$  20.236(2),  $b$  23.798(1),  $c$  12.798(1) Å (Pluth & Smith 1990).

Si,Al highly disordered.

### Brewsterite (series)

$(Sr,Ba)_2[Al_4Si_{12}O_{32}] \cdot 10H_2O$   $Z = 1$  BRE

Brooke (1822). Type locality: Strontian, Argyll, Scotland. Named after Sir David Brewster (1781–1868), Scottish natural philosopher who discovered laws of polarization of light in biaxial crystals.

Monoclinic,  $P2_1/m$ ,  $P2_1$ , or triclinic (Akizuki 1987a, Akizuki *et al.* 1996).

The structure is sheet-like parallel to (010) (Perrotta & Smith 1964).

### Brewsterite-Sr

New name for the original species of the series; Sr is the most abundant extra-framework cation.  $T_{Si}$  in the range 0.74–0.75.

Monoclinic,  $P2_1/m$ ,  $a$  6.793(2),  $b$  17.573(6),  $c$  7.759(2) Å,  $\beta$  94.54(3)°, for composition  $(Sr_{1.42}Ba_{0.48}K_{0.02})$



[Al<sub>4.12</sub>Si<sub>11.95</sub>O<sub>32</sub>]*n*H<sub>2</sub>O (Schlenker *et al.* 1977a).  
On optical grounds, possibly triclinic (Akizuki 1987a).  
Refined as triclinic in three separate growth-sectors by  
Akizuki *et al.* (1996).  
Partly ordered Si,Al distribution.

### Brewsterite-Ba

New name; Ba is the most abundant extra-framework cation.

Proposed type-example: the Gouverneur Talc Company's No. 4 wollastonite mine near Harrisville, Lewis County, New York, U.S.A. (Robinson & Grice 1993). Also Cerchiara mine, Liguria, Italy (Cabella *et al.* 1993, including structure refinement).  $T_{Si} = 0.73, 0.74$ .

Monoclinic,  $P2_1/m$  or  $P2_1$ ,  $a 6.780(3), b 17.599(9), c 7.733(2)$  Å,  $\beta 94.47(3)^\circ$  for type example, containing up to 0.85 Ba per 16 O atoms.

### Chabazite (series)

(Ca<sub>0.5</sub>,Na,K)<sub>4</sub>[Al<sub>4</sub>Si<sub>8</sub>O<sub>24</sub>]*n*H<sub>2</sub>O

$Z = 1$  (trigonal) CHA

Bosc d'Antic (1792), as "chabazie". The source of the original specimen is unclear. The name is from a word "chabazion" used for an unknown substance in the story of Orpheus.

Ca-, Na-, and K-dominant species occur in that order of frequency, with Sr and Mg occasionally significant, Ba more minor.  $T_{Si}$  varies widely, 0.58 to 0.81.

Topological symmetry of the framework, trigonal ( $R\bar{3}m$ ), where  $a \approx 13.2, c \approx 15.1$  Å (pseudo-hexagonal cell). Significant deviations to triclinic,  $P\bar{1}$ ,  $a \approx 9.4, b \approx 9.4, c \approx 9.4$  Å,  $\alpha \approx 94^\circ, \beta \approx 94^\circ, \gamma \approx 94^\circ$  (Smith *et al.* 1964, Mazzi & Galli 1983).

Partial ordering leads to the lower symmetry.

### Chabazite-Ca

New name for the original and most common species; Ca is the most abundant single extra-framework cation. Other cations vary widely.  $T_{Si}$  in the range 0.58–0.80.  $a 13.790(5), c 15.040(4)$  Å, for pseudo-hexagonal cell, with composition (Ca<sub>1.86</sub>Na<sub>0.03</sub>K<sub>0.20</sub>Mg<sub>0.02</sub>Sr<sub>0.03</sub>)[Al<sub>3.94</sub>Fe<sub>0.01</sub>Si<sub>8.03</sub>O<sub>24</sub>]*n*H<sub>2</sub>O, from Col de Lares, Val di Fassa, Italy (Passaglia 1970, #13).

### Chabazite-Na

New name; Na is the most abundant single extra-framework cation. Other cations vary widely.  $T_{Si}$  in the range 0.62–0.79.

Suggested type-locality: biggest "Faraglione" facing Acì Trezza, Sicily, Italy (Passaglia 1970, #1).  $a 13.863(3), c 15.165(3)$  Å, for hexagonal cell, with composition (Na<sub>3.11</sub>K<sub>1.05</sub>Ca<sub>0.19</sub>Mg<sub>0.06</sub>Sr<sub>0.05</sub>)[Al<sub>4.53</sub>Fe<sub>0.01</sub>Si<sub>7.40</sub>O<sub>24</sub>]*n*H<sub>2</sub>O.

Although originally described as containing "silex, alumina, and potash" (Lévy 1825), the name *herschelite* has often been applied to chabazite minerals of tabular habit and high Na content. *Herschelite* should no longer be used as a species name.

### Chabazite-K

New name; K is the most abundant single extra-framework cation. Other cations vary widely.  $T_{Si}$  in the range 0.60–0.74.

Suggested type-specimen: Tufo Ercolano, Ercolano, Naples, Italy (De Gennaro & Franco 1976),  $a 13.849(3), c 15.165(3)$  Å, for hexagonal cell, with composition (K<sub>2.06</sub>Na<sub>0.98</sub>Ca<sub>0.46</sub>Mg<sub>0.10</sub>Sr<sub>0.01</sub>)[Al<sub>4.37</sub>Fe<sub>0.08</sub>Si<sub>7.60</sub>O<sub>24</sub>]*n*H<sub>2</sub>O.

### Chiavennite

CaMn[Be<sub>2</sub>Si<sub>5</sub>O<sub>13</sub>(OH)<sub>2</sub>]*n*H<sub>2</sub>O

$Z = 4$

–CHI

Bondi *et al.* (1983), Raade *et al.* (1983). Type locality: Chiavenna, Lombardy, Italy. Named after type locality. The limited data available show up to 0.72 Al and 0.15 B in tetrahedral sites, and significant extra-framework Fe and Na (Raade *et al.* 1983, Langhof & Holstam 1994).  $T_{Si}$  in the range 0.63 to 0.68. Orthorhombic,  $Pnab$ ,  $a 8.729(5), b 31.326(11), c 4.903(2)$  Å (Tazzoli *et al.* 1995).

A Ca,Mn beryllosilicate with an interrupted framework of four-connected [SiO<sub>4</sub>] and three-connected [BeO<sub>4</sub>] tetrahedra.

### Clinoptilolite (series)

(Na,K,Ca<sub>0.5</sub>,Sr<sub>0.5</sub>,Ba<sub>0.5</sub>,Mg<sub>0.5</sub>)<sub>6</sub>[Al<sub>6</sub>Si<sub>30</sub>O<sub>72</sub>]*n*H<sub>2</sub>O

$Z = 1$

HEU

Schaller (1923, 1932). Type locality: in decomposed basalt at a high point on ridge running east from Hoodoo Mountain, Wyoming, U.S.A. ("crystallized mordenite" of Pirsson 1891). The name reflects its inclined extinction and supposed similarity in composition to "ptilolite" (mordenite). Ptilo-, from Greek, alludes to the downy, finely fibrous nature of that mineral.

The cation content is highly variable. Ca-, Na-, and K-dominant compositions are known, and Sr, Ba, and Mg are in some cases substantial. Fe<sup>2+</sup> and Fe<sup>3+</sup> are possible constituents. In Pirsson's (1890) analysis, K is the most abundant single cation by a small margin. Clinoptilolite-K is therefore taken as the type species of the series.  $T_{Si}$  in the range 0.80–0.84.

Minerals with the same framework topology but with  $T_{Si} < 0.80, Si/Al < 4.0$  are classified as heulandite, with which clinoptilolite forms a continuous series.

Monoclinic,  $C2/m$ , or  $C2$ , or  $Cm$ .

Structure refinements by Alberti (1975a) and Armbruster (1993) demonstrate variations in extra-

framework cation sites compared with heulandite and as a function of the extent of dehydration.

### Clinoptilolite-K

New name for the original species; K is the most abundant single extra-framework cation. A moderately K-rich clinoptilolite-K was referred to as "potassium clinoptilolite" by Minato & Takano (1964).  $T_{Si}$  in the range 0.80–0.83.

Monoclinic,  $C2/m$ ,  $C2$ , or  $Cm$ ,  $a$  17.688(16),  $b$  17.902(9),  $c$  7.409(7) Å,  $\beta$  116.50(7)°, for  $(K_{4.72}Na_{0.85}Ca_{0.04}Sr_{0.37}Mg_{0.19}Fe_{0.03}Mn_{0.01})[Al_{6.52}Si_{29.38}O_{72}] \cdot nH_2O$ , from an off-shore borehole, Japan (Ogihara & Iijima 1990).

### Clinoptilolite-Na

New name; Na is the most abundant single extra-framework cation. Other cations vary widely.  $T_{Si}$  in the range 0.80–0.84.

Suggested type-example: Barstow Formation, about 1.6 km east of mouth of Owl Canyon, San Bernardino County, California, U.S.A., USGS Lab. no. D100594 (Sheppard & Gude 1969a).

Monoclinic,  $C2/m$ ,  $C2$ , or  $Cm$ ,  $a$  17.627(4),  $b$  17.955(4),  $c$  7.399(4) Å,  $\beta$  116.29(2)° (Boles 1972), for type material of Sheppard & Gude (1969a),  $(Na_{3.78}K_{1.31}Ca_{0.61}Ba_{0.09}Mg_{0.23}Mn_{0.01})[Al_{6.61}Fe_{0.16}Si_{29.19}O_{72}] \cdot 20.4H_2O$ .

### Clinoptilolite-Ca

New name; Ca is the most abundant single extra-framework cation. Other cations vary widely.  $T_{Si}$  in the range 0.80–0.84.

Suggested type-specimen: Kuruma Pass, Fukushima Prefecture, Japan (Koyama & Takéuchi 1977).

Monoclinic,  $C2/m$ ,  $C2$ , or  $Cm$ ,  $a$  17.660(4),  $b$  17.963(5),  $c$  7.400(3) Å,  $\beta$  116.47(3)° based on  $C2/m$  (Koyama & Takéuchi 1977), for Kuruma Pass specimen,  $(Na_{1.76}K_{1.05}Ca_{1.90}Mg_{0.17})[Al_{6.72}Si_{29.20}O_{72}] \cdot 23.7H_2O$ .

### Cowlesite

$Ca[Al_2Si_3O_{10}] \cdot 5.3H_2O$   $Z = 52$  (IZA code not assigned)

Wise & Tschernich (1975). Type locality: road cuts 0.65 km northwest of Goble, Columbia County, Oregon, U.S.A. Named after John Cowles of Rainier, Oregon, amateur mineralogist.

Minor substitution for Ca by Na and lesser K, Mg, Sr, Ba, Fe.  $T_{Si}$  in the range 0.60–0.62 (Vezzalini *et al.* 1992). Orthorhombic,  $P222_1$  or  $Pmnm$ ,  $Pmm2$ ,  $P2mm$ ,  $P222$  (Nawaz 1984),  $a$  23.249(5),  $b$  30.629(3),  $c$  24.964(4) Å (Artioli *et al.* 1987).

Structure and degree of order of framework cations have not been determined.

### Dachiardite (series)

$(Ca_{0.5}, Na, K)_{4-5}[Al_{4-5}Si_{20-19}O_{48}] \cdot 13H_2O$

$Z = 1$

DAC

D' Achiardi (1906). Type locality: San Piero in Campo, Elba, Italy. Named by the author in memory of his father, Antonio D' Achiardi (1839–1902), first full professor of Mineralogy at the University of Pisa.

May contain minor Cs and Sr.  $T_{Si}$  in the range 0.78–0.86. Monoclinic, topological symmetry  $C2/m$ , real symmetry  $Cm$ .

The structure consists of complex chains of 5-membered rings cross-linked by 4-membered rings (Gottardi & Meier 1963), but with complexities that commonly result in diffuse and streaked X-ray-diffraction maxima (Quartieri *et al.* 1990).

### Dachiardite-Ca

New name for the original species of the series; Ca is the most abundant extra-framework cation. Dachiardite from the type locality contains 0.12 Cs atoms per formula unit (*apfu*) (Bonardi 1979).  $T_{Si}$  in the range 0.78–0.83.

Monoclinic, topological symmetry  $C2/m$ , real symmetry  $Cm$ .  $a$  18.676,  $b$  7.518,  $c$  10.246 Å,  $\beta$  107.87°, for the composition  $(Ca_{1.54}Na_{0.42}K_{0.92}Cs_{0.11}Sr_{0.12}Ba_{0.01})[Al_{4.86}Fe_{0.02}Si_{18.96}O_{48}] \cdot 12.56H_2O$  from the type locality (Vezzalini 1984).

Partly ordered distribution of Si, Al.

### Dachiardite-Na

New name; Na is the most abundant extra-framework cation.

Suggested type-example: Alpe di Siusi, Bolzano, Italy (Alberti 1975b).

Available analytical results for material from seven localities, *e.g.*, Bonardi *et al.* (1981) show considerable variation in Na : K : Ca proportions.  $T_{Si}$  in the range 0.81–0.86.

Monoclinic,  $a$  18.647(7),  $b$  7.506(4),  $c$  10.296(4) Å,  $\beta$  108.37(3)°, for  $(Na_{2.59}K_{0.71}Ca_{0.53}Mg_{0.04}Ba_{0.01})[Al_{4.27}Fe_{0.11}Si_{19.61}O_{48}] \cdot 13.43H_2O$  from the type locality (Alberti 1975b).

Diffuse diffraction-spots indicate disorder.

### Edingtonite

$Ba[Al_2Si_3O_{10}] \cdot 4H_2O$   $Z = 2$

EDI

Haidinger (1825). Type locality: Kilpatrick Hills, near Glasgow, Scotland. Named after a Mr. Edington of Glasgow, in whose collection Haidinger found the mineral. Small amounts of K, Na, and Ca may replace Ba.  $T_{Si}$  in the range 0.59–0.61.


Orthorhombic,  $P2_12_12_1$ ,  $a$  9.550(10),  $b$  9.665(10),  $c$  6.523(5) Å (Böhlet mine, Westergotland, Sweden) (Galli 1976).

Also tetragonal,  $P\bar{4}2_1m$ ,  $a$  9.584(5),  $c$  6.524(3) Å (Old Kilpatrick, near Glasgow, Scotland) (Mazzi *et al.* 1984).

From optical evidence, Akizuki (1986) suggested that a triclinic true symmetry also is possible.

The structure is similar to that of natrolite, but with a distinctive cross-linking of the chains (Taylor & Jackson 1933, Mazzi *et al.* 1984). Examples of orthorhombic edingtonite have nearly perfect (Si,Al) order. The tetragonal form is disordered, and available analytical results show that slightly more Ba has been replaced by other ions.

### Epistilbite



$$Z = 4$$

EPI


Rose (1826). Type localities: "Iceland" and "Farøe Islands". Named from Greek *epi* in the sense of near, and stilbite, from its supposed similarity to the latter. Na/(Na + Ca) varies from about 0.1 to 0.3, with minor K and Ba (*e.g.*, Galli & Rinaldi 1974).  $T_{\text{Si}}$  in the range 0.72–0.77.

Monoclinic,  $C2$ ,  $a$  9.101(2),  $b$  17.741(1),  $c$  10.226(1) Å,  $\beta$  124.66(2)° (Teigarhorn, Iceland: Alberti *et al.* 1985), or

triclinic,  $C\bar{1}$ ,  $a$  9.083(1),  $b$  17.738(3),  $c$  10.209(1) Å,  $\alpha$  89.95(1)°,  $\beta$  124.58(1)°,  $\gamma$  90.00(1)° (Gibelsbach, Valais, Switzerland: Yang & Armbruster 1996).

The structural framework belongs to the mordenite group (Gottardi & Galli 1985). Earlier work suggested space-group symmetry  $C2/m$  (Perrotta 1967). Alberti *et al.* (1985) proposed a domain structure involving acentric configurations of tetrahedra, and space group  $C2$ . Yang & Armbruster (1996) indicated that the proposed domains can be modeled by (010) disorder caused by a local mirror plane, and that increased partial order of Si,Al leads to triclinic symmetry.

### Erionite (series)



$$Z = 1$$

ERI

Eakle (1898). Type locality: Durkee, Oregon, U.S.A., in rhyolitic, welded ash-flow tuff. Name from Greek root meaning wool, in reference to its appearance.

Substantial amounts of any or all of Ca, Na, and K, and subordinate Mg may be present, and there is evidence that trace Fe may enter tetrahedral and extra-framework sites. Eakle's (1898) analysis of type erionite shows Na as the most abundant extra-framework cation; Passaglia *et al.* (1998) found Ca to be the most abundant in a type-locality specimen.  $T_{\text{Si}}$  in the range 0.68–0.79.

Hexagonal,  $P6_3/mmc$ ,  $a$  13.15,  $c$  15.02 Å (Kawahara & Curien 1969).

The structure is related to those of offretite, with which it may form intergrowths with stacking faults (Schlenker *et al.* 1977b), and levyne, on which it forms

epitactic growths (Passaglia *et al.* 1998). The three minerals have 4-, 6- and 8-membered rings. They differ in the stacking of single and double 6-membered rings, resulting in different  $c$  dimensions and differently sized and shaped cages. Si,Al disordered.

### Erionite-Na

New name; Na is the most abundant extra-framework cation.

Proposed type-example: Cady Mountains, California, U.S.A. (Sheppard *et al.* 1965).  $T_{\text{Si}}$  in the range 0.74–0.79. For the type specimen,  $a$  13.214(3),  $c$  15.048(4) Å, composition  $(\text{Na}_{5.39}\text{K}_{2.00}\text{Ca}_{0.11}\text{Mg}_{0.18}\text{Fe}_{0.02})[\text{Al}_{7.57}\text{Si}_{28.27}\text{O}_{72}] \cdot 24.60\text{H}_2\text{O}$  (Sheppard & Gude 1969b).

### Erionite-K

New name; K is the most abundant extra-framework cation.

Proposed type-example: Rome, Oregon, U.S.A. in which K makes up 58% of extra-framework cations; significant Na, Ca, and Mg also are present (Eberly 1964).  $T_{\text{Si}}$  in the range 0.74–0.79.

For a specimen from Ortenberg, Germany,  $a$  13.227(1),  $c$  15.075(3) Å,  $(\text{K}_{3.32}\text{Na}_{2.31}\text{Ca}_{0.99}\text{Mg}_{0.06}\text{Ba}_{0.02})[\text{Al}_{8.05}\text{Si}_{28.01}\text{O}_{72}] \cdot 31.99\text{H}_2\text{O}$  (Passaglia *et al.* 1998).

### Erionite-Ca

New name; Ca is the most abundant extra-framework cation.

Proposed type-example: Mazé, Niigata Prefecture, Japan (Harada *et al.* 1967).  $T_{\text{Si}}$  in the range 0.68–0.79. For the type example:  $a$  13.333(1),  $c$  15.091(2) Å;  $(\text{Ca}_{2.28}\text{K}_{1.54}\text{Na}_{0.95}\text{Mg}_{0.86})[\text{Al}_{8.83}\text{Si}_{26.90}\text{O}_{72}] \cdot 31.35\text{H}_2\text{O}$  (Harada *et al.* 1967).

### Faujasite (series)



$$Z = 16$$

FAU

Damour (1842). Type locality: Sasbach, Kaiserstuhl, Germany. Named after Barthélémy Faujas de Saint Fond, noted for his work on extinct volcanoes.

Major amounts of Na, Ca, and Mg are commonly present, and in some cases, K; minor Sr is also reported. The ratio Si : Al also varies;  $T_{\text{Si}}$  in the range 0.68–0.74, with one record of 0.64. In most samples analyzed,  $x$  in the above generalized formula is in the range 3.2–3.8, with one record of 4.4 (Rinaldi *et al.* 1975a, Wise 1982, Ibrahim & Hall 1995).

Cubic,  $Fd\bar{3}m$ ,  $a$  24.65 Å (material from Sasbach: Bergerhoff *et al.* 1958).

The framework structure is very open, with complete sodalite-type cages and with very large cavities having 12-membered ring openings. Up to 260 molecules of  $\text{H}_2\text{O}$  can be accommodated per unit cell (Bergerhoff *et al.* 1958, Baur 1964).

**Faujasite-Na**

New name; Na is the most abundant extra-framework cation, as it is in the original (incomplete) and most subsequent analyses of samples from the type locality, Sasbach, Kaiserstuhl, and some other localities.  $T_{Si}$  in the range 0.70–0.74, with one report of 0.64.

Reported values of  $a$  range from 24.638(3) Å (Wise 1982) to 24.728(2) Å (Ibrahim & Hall 1995).

**Faujasite-Ca**

New name; Ca is the most abundant extra-framework cation. Reported  $T_{Si}$  in the range 0.68–0.73. Proposed type-example: drill core from Haselborn near Ilbeshausen, Vogelsberg, Hessen, Germany (Wise 1982), composition  $(Ca_{1.32}Na_{0.56}Mg_{0.26}K_{0.04})[Al_{3.83}Si_{8.19}O_{24}] \cdot nH_2O$ ,  $Z = 16$ .

Reported values of  $a$ : 24.714(4) and 24.783(3) Å (Jabal Hanoun, Jordan: Ibrahim & Hall 1995).

**Faujasite-Mg**

New name; Mg is the most abundant extra-framework cation.

Proposed type (and only) example: "Old (museum sample)" (# 32, Genth Collection, Pennsylvania State University) from Sasbach, Kaiserstuhl, Germany (anal. #15, Rinaldi *et al.* 1975a), composition  $(Mg_{15.3}Ca_{4.0}Na_{7.0}K_{6.4})[Al_{56}Si_{137}O_{384}] \cdot nH_2O$ ,  $Z = 1$ .

**Ferrierite (series)**

$(K, Na, Mg_{0.5}, Ca_{0.5})_6[Al_6Si_{30}O_{72}] \cdot 8H_2O$   
 $Z = 1$

FER

Graham (1918). Type locality: Kamloops Lake, British Columbia, Canada. Named after Dr. Walter F. Ferrier, mineralogist, mining engineer, and one-time member of the Geological Survey of Canada, who first collected it. Substantial amounts of any or all of Mg, K, Na, and Ca, may be present, and smaller amounts of Fe, Ba, and Sr.  $T_{Si}$  in the range 0.80–0.88.

Statistical symmetry, orthorhombic, *Immm*; true symmetries orthorhombic, *Pnmm*,  $a$  19.23,  $b$  14.15,  $c$  7.50 Å (Alberti & Sabelli 1987), and monoclinic,  $P2_1/n$ ,  $a$  18.89,  $b$  14.18,  $c$  7.47 Å,  $\beta$  90.0° (Gramlich-Meier *et al.* 1985).

The structure was first determined by Vaughan (1966). Framework Si, Al partially ordered (Alberti & Sabelli 1987).

**Ferrierite-Mg**

New name for the original member of the series; Mg is the most abundant single extra-framework cation.

Substantial extra-framework Na, K, and lesser Ca commonly present.  $T_{Si}$  in the range 0.80–0.84.

True symmetry orthorhombic, *Pnmm*,  $a$  19.231(2),

$b$  14.145(2),  $c$  7.499(1) Å for specimen from Monastir, Sardinia, of composition  $(Mg_{2.02}K_{1.19}Na_{0.56}Ca_{0.52}Sr_{0.14}Ba_{0.02})[Al_{6.89}Si_{29.04}O_{72}] \cdot 17.86H_2O$  (Alberti & Sabelli 1987).

**Ferrierite-K**

New name; K is the most abundant single extra-framework cation.

Proposed type-example: Santa Monica Mountains, California, U.S.A., composition  $(K_{2.05}Na_{1.14}Mg_{0.74}Ca_{0.14})[Al_{5.00}Si_{31.01}O_{72}] \cdot nH_2O$  (Wise & Tschernich 1976, #3).  $T_{Si}$  in the range 0.81–0.87.

Orthorhombic,  $a$  18.973(7),  $b$  14.140(6),  $c$  7.478(4) Å for type specimen.

**Ferrierite-Na**

New name; Na is the most abundant single extra-framework cation.

Proposed type-example: Altoona, Washington, U.S.A., composition  $(Na_{3.06}K_{0.97}Mg_{0.35}Ca_{0.05}Sr_{0.02}Ba_{0.02})[Al_5Si_{31}O_{72}] \cdot 18H_2O$  (Wise & Tschernich 1976, #1).

$T_{Si}$  in the range 0.85–0.88.

Monoclinic,  $P2_1/n$ ,  $a$  18.886(9),  $b$  14.182(6),  $c$  7.470(5) Å,  $\beta$  90.0(1)° (Gramlich-Meier *et al.* 1985, for a specimen from Altoona, Washington).

**Garronite**

$NaCa_{2.5}[Al_6Si_{10}O_{32}] \cdot 14 H_2O$   
 $Z = 1$

GIS

Walker (1962). Type locality: slopes of Glenariff Valley, County Antrim, Northern Ireland. Named after the Garron Plateau, where the type locality is sited.

Ca/(Na + K) is variable, but Ca predominates. Type-locality garronite has about 1.3 Na *apfu*, some others have (Na + K) < 0.2 *apfu*.  $H_2O$  in the range 13.0–14.0 molecules per formula unit.  $T_{Si}$  in the range 0.60–0.65.

The crystal structure has been refined in tetragonal symmetry,  $I\bar{4}m2$ ,  $a$  9.9266(2),  $c$  10.3031(3) Å, by Artioli (1992), and Na-free synthetic garronite has been refined in  $I4_1/a$ ,  $a$  9.873(1),  $c$  10.288(1) Å, by Schröpfer & Joswig (1997). Orthorhombic symmetry has been proposed on the basis of X-ray diffraction with twinned crystals (Nawaz 1983) and crystal morphology (Howard 1994). The framework topology is the same as for gismondine, but Si and Al are essentially disordered. The different space-group symmetry (Artioli 1992) is associated with disorder and the presence of significant Na. Gottardi & Alberti (1974) proposed partial ordering subsequent to growth to explain twin domains.

**Gaultite**

$Na_4[Zn_2Si_7O_{18}] \cdot 5H_2O$   $Z = 8$

VSV

Ercit & Van Velthuizen (1994). Type locality: Mont Saint-Hilaire, Quebec, Canada. Named after Robert A.

Gault, (b. 1943), mineralogist at the Canadian Museum of Nature, Ottawa, Ontario, Canada.

No other elements detected in the one reported example;  $T_{Si} = 0.78$ .

Orthorhombic,  $F2dd$ ,  $a$  10.211(3),  $b$  39.88(2),  $c$  10.304(4) Å.

The zincosilicate framework of tetrahedra is characterized by stacked sheets of edge-sharing 4- and 8-membered rings. The sheets are cross-linked by tetrahedra. Gaultite is isostructural with synthetic zeolite VPI-7 and similar in structure to lovdarite (Ercit & Van Velthuizen 1994).

### Gismondine

$Ca[Al_2Si_2O_8] \cdot 4.5H_2O$   $Z = 4$  GIS  
von Leonhard (in footnote, 1817), renaming "zeagonite" of Gismondi (1817). Type locality: Capo di Bove, near Rome, Italy. Named after Carlo Giuseppe Gismondi (1762–1824), lecturer in Mineralogy in Rome.

(K + Na) does not exceed 0.12 *apfu* with K less than 0.08 *apfu*; analyses showing high K result from intergrown phillipsite. Minor Sr may be present;  $T_{Si}$  in the range 0.51–0.54 (Vezzalini & Oberti 1984).  $H_2O$  is slightly variable (4.4–4.5 molecules per formula unit) because of mixed 6- and 7-coordination of Ca (Artioli *et al.* 1986b).

Monoclinic, originally refined in  $P2_1/a$  by Fischer & Schramm (1970); cell converted to standard  $P2_1/c$  second setting is  $a$  10.023(3),  $b$  10.616(5),  $c$  9.843(15) Å,  $\beta$  92.42(25)°. Also refined (two samples) by Rinaldi & Vezzalini (1985).

The framework topology is based on crankshaft chains of 4-membered rings as in feldspars, connected in UDD configuration.

Si,Al are strictly ordered.

### Gmelinite (series)

$(Na_2, Ca, K)_4[Al_8Si_{16}O_{48}] \cdot 22H_2O$   
 $Z = 1$  GME

Brewster (1825a). Type locality: the name was proposed for minerals occurring both at Little Deer Park, Glenarm, County Antrim, Northern Ireland, and at Montecchio Maggiore, Vicenza, Italy. Named after Christian Gottlob Gmelin, Professor of Chemistry, University of Tübingen.

Na-dominant members are the most common.  $T_{Si}$  in the range 0.65–0.72.

Hexagonal,  $P6_3/mmc$ ,  $a$  13.62–13.88,  $c$  9.97–10.25 Å. The structure is similar to that of chabazite, with which it is commonly intergrown (Strunz 1956), but gmelinite has a different stacking of the double 6-membered rings (Fischer 1966). Si,Al are disordered.

### Gmelinite-Na

New name for the most common species of the series. It occurs in at least one of the gmelinite type-localities (Montecchio Maggiore). The Ca content is commonly substantial, K is minor, and Sr is significant in a few samples analyzed.  $T_{Si}$  in the range 0.65–0.71.

Hexagonal,  $P6_3/mmc$ ,  $a$  13.756(5),  $c$  10.048(5) Å (Galli *et al.* 1982), for near-end-member material from Queensland, Australia, of composition  $(Na_{7.61}Ca_{0.03}K_{0.16})[Al_{7.41}Si_{16.49}O_{48}] \cdot 21.51H_2O$  (Passaglia *et al.* 1978a).

### Gmelinite-Ca

New name for a species that also occurs in at least one of the type localities (Montecchio Maggiore). Ca is the most abundant single extra-framework cation. Significant to substantial Sr and Na, minor K.  $T_{Si}$  in the range 0.68–0.70.

Hexagonal,  $P6_3/mmc$ ,  $a$  13.800(5),  $c$  9.964(5) Å (Galli *et al.* 1982), from Montecchio Maggiore, of composition  $(Ca_{2.06}Sr_{1.35}Na_{0.78}K_{0.11})[Al_{7.82}Si_{16.21}O_{48}] \cdot 23.23H_2O$  (Passaglia *et al.* 1978a).

### Gmelinite-K


New name; K is the most abundant single extra-framework cation. Proposed type-example: Fara Vicentina, Vicenza, Italy, composition  $(K_{2.72}Ca_{1.67}Sr_{0.39}Na_{0.22}Mg_{0.13})[Al_{7.79}Si_{16.32}O_{48}] \cdot 23.52H_2O$  (Vezzalini *et al.* 1990). Also known from the Kola Peninsula (Malinovskii 1984).

Hexagonal,  $P6_3/mmc$ ,  $a$  13.621(3),  $c$  10.254(1) Å.

### Gobbinsite

$Na_5[Al_5Si_{11}O_{32}] \cdot 12H_2O$   $Z = 1$  GIS  
Nawaz & Malone (1982). Type locality: basalt cliffs near Hills Port, south of the Gobbins area, County Antrim, Northern Ireland. Named after the locality.

Na : Ca : Mg : K variable, with Na greatly predominant, Ca < 0.6 *apfu*. Reports of high K are ascribed to intergrown phillipsite (Artioli & Foy 1994).  $T_{Si}$  in the range 0.62–0.68, substantially higher than in gismondine. Orthorhombic,  $Pmn2_1$ ,  $a$  10.108(1),  $b$  9.766(1),  $c$  10.171(1) Å for the anhydrous composition  $(Na_{2.50}K_{2.11}Ca_{0.59})[Al_{6.17}Si_{9.93}O_{32}]$  from Two-Mouth Cave, County Antrim, Northern Ireland (McCusker *et al.* 1985);  $a$  10.1027(5),  $b$  9.8016(5),  $c$  10.1682(6) Å for  $(Na_{4.3}Ca_{0.6})[Al_{5.6}Si_{10.4}O_{32}] \cdot 12H_2O$  from Magheramorne quarry, Larne, Northern Ireland (Artioli & Foy 1994). The framework topology is the same as for gismondine and is based on crankshaft chains of 4-membered rings, as in feldspars. Distortion from tetragonal topological symmetry results from the arrangement of cations in the channels. Si,Al in the framework are disordered.

**Gonnardite**


NAT

Lacroix (1896). Type locality: Chaux de Bergonne, Gignat, Puy-de-Dôme, France. Named after Ferdinand Gonnard, who had earlier described the material as "mesole" (= thomsonite).

Forms an extensive substitution series, commonly approximating  $\text{Na}_{8-3x}\text{Ca}_{2x}[\text{Al}_{8+x}\text{Si}_{12-x}\text{O}_{40}] \cdot 12\text{H}_2\text{O}$  (after Ross *et al.* 1992), with minor  $\text{Fe}^{3+}$ , Mg, Ba, Sr, and K.  $T_{\text{Si}}$  in the range 0.52–0.59 (or 0.52–0.62 if tetranatrolite = gonnardite).

Tetragonal, *I2d*, *a* 13.21(1), *c* 6.622(4) Å for material from Tvedalen, Langesund, Norway, of composition  $(\text{Na}_{6.42}\text{K}_{0.01}\text{Ca}_{1.50})[\text{Al}_{9.22}\text{Si}_{10.73}\text{O}_{40}] \cdot 12.37\text{H}_2\text{O}$  (Mazzi *et al.* 1986).

The structure is similar to that of natrolite, but with Si,Al disordered, and usually with significant to substantial Ca (Mazzi *et al.* 1986, Artioli & Torres Salvador 1991, Alberti *et al.* 1995).

**Goosecreekite**


Z = 2

GOO

Dunn *et al.* (1980). Type locality: Goose Creek quarry, Loudoun County, Virginia, U.S.A. Named after the locality. Results of the single analysis available conform closely to the formula given, with no other elements detected.  $T_{\text{Si}} = 0.75$ .

Monoclinic, *P2*<sub>1</sub>, *a* 7.401(3), *b* 17.439(6), *c* 7.293(3) Å, β 105.44(4)° (Rouse & Peacor 1986).

The framework consists of 4-, 6-, and 8-membered rings that link to form layers parallel to (010), with some similarities to the brewsterite structure. Si,Al are nearly perfectly ordered (Rouse & Peacor 1986).

**Gottardiite**

Z = 1


NES

Alberti *et al.* (1996), Galli *et al.* (1996). Mt. Adamson, Victoria Land, Antarctica. Named after Professor Glauco Gottardi (1928–1988), University of Modena, in recognition of his pioneering work on the structure and crystal chemistry of natural zeolites.

Known from the type locality only, with composition approximating the above simplified formula; minor K, and very high Si.  $T_{\text{Si}} = 0.86$ .

Orthorhombic, topological symmetry *Fmmm*, real symmetry *Cmca*, *a* 13.698(2), *b* 25.213(3), *c* 22.660(2) Å (Alberti *et al.* 1996).

The framework topology is the same as for the synthetic zeolite NU-87, which, however, has monoclinic symmetry, *P2*<sub>1</sub>/*c*. Some Si,Al order is probable.

**Harmotome**

Z = 1


PHI

Häyly (1801, p. 191–195), renaming andreasbergolite, also known as andréolite, of Delamétherie (1795, p. 393). Type locality: Andreasberg, Harz, Germany. Named from Greek words for a "joint" and "to cut", in allusion to a tendency to split along junctions (twin planes).

Ba is the most abundant extra-framework cation. Harmotome forms a continuous series with phillipsite-Ca. The name *harmotome* predates *phillipsite*; on grounds of history and usage, both are retained in spite of Rule 1 of the present report.  $T_{\text{Si}}$  in the range 0.68–0.71 (*e.g.*, Černý *et al.* 1977).

Monoclinic, refined in *P2*<sub>1</sub>/*m*, but on piezoelectric and optical grounds, the true symmetry may be noncentrosymmetric and triclinic, *P1* (*e.g.*, Akizuki 1985, Stuckenschmidt *et al.* 1990), *a* 9.879(2), *b* 14.139(2), *c* 8.693(2) Å, β 124.81(1)° for  $(\text{Ba}_{1.93}\text{Ca}_{0.46}\text{K}_{0.07})[\text{Al}_{4.66}\text{Si}_{11.29}\text{O}_{32}] \cdot 12\text{H}_2\text{O}$  from Andreasberg, Harz (Rinaldi *et al.* 1974).

The structure is the same as for phillipsite, with little or no Si,Al order.

**Heulandite (series)**

Z = 1

HEU

Brooke (1822). Type locality: none; the name was given to the more distinctly monoclinic minerals previously known as stilbite. Named after Henry Heuland, English mineral collector.

The cation content is highly variable. Ca-, Na-, K-, and Sr-dominant compositions are known, and Ba and Mg are in some cases substantial.  $T_{\text{Si}}$  in the range 0.71 to 0.80. Minerals with the same framework topology, but with  $T_{\text{Si}} \geq 0.80$ , Si/Al  $\geq 4.0$ , are distinguished as clinoptilolite.

Monoclinic, with highest possible topological symmetry *C2/m* (*I2/m*). *Cm* and *C2* also have been suggested. The sheet-like structure was solved by Merkle & Slaughter (1968). There is partial order of Si,Al.

**Heulandite-Ca**

New name for the most common species of the series, and that deduced from results of most older analyses. Ca is the most abundant single extra-framework cation.  $T_{\text{Si}}$  in the range 0.71–0.80.

Monoclinic, *C2/m*, *Cm*, or *C2*, *a* 17.718(7), *b* 17.897(5), *c* 7.428(2) Å, β 116.42(2)° from Farøe Islands, composition  $(\text{Ca}_{3.57}\text{Sr}_{0.05}\text{Ba}_{0.06}\text{Mg}_{0.01}\text{Na}_{1.26}\text{K}_{0.43})[\text{Al}_{9.37}\text{Si}_{26.70}\text{O}_{72}] \cdot 26.02\text{H}_2\text{O}$  ( $T_{\text{Si}} = 0.74$ ) (Alberti 1972).

**Heulandite-Sr**

New name; Sr is the most abundant single extra-framework cation.

One known example: Campegli, Eastern Ligurian ophiolites, Italy, of composition  $(\text{Sr}_{2.10}\text{Ca}_{1.76}\text{Ba}_{0.14}\text{Mg}_{0.02}\text{Na}_{0.40}\text{K}_{0.22})[\text{Al}_{9.19}\text{Si}_{26.94}\text{O}_{72} \cdot n\text{H}_2\text{O}]$ ,  $T_{\text{Si}} = 0.75$  (Lucchetti *et al.* 1982).

Monoclinic,  $C2/m$ ,  $Cm$ , or  $C2$ ,  $a$  17.655(5),  $b$  17.877(5),  $c$  7.396(5) Å,  $\beta$  116.65°.

**Heulandite-Na**

New name; Na is the most abundant single extra-framework cation.

Proposed type-example: Challis, Idaho, U.S.A., U.S. National Museum #94512/3 (Ross & Shannon 1924, Boles 1972, #6).

Monoclinic,  $C2/m$ ,  $Cm$ , or  $C2$ ,  $a$  17.670(4),  $b$  17.982(4),  $c$  7.404(2) Å,  $\beta$  116.40(2)° (Boles 1972) for the type example, of composition  $(\text{Na}_{3.98}\text{Ca}_{1.77}\text{K}_{0.55})[\text{Al}_{7.84}\text{Si}_{28.00}\text{O}_{72}] \cdot 21.74\text{H}_2\text{O}$ ,  $T_{\text{Si}} = 0.78$ .

**Heulandite-K**

New name; K is the most abundant single extra-framework cation.

Proposed type-example: Albergo Bassi, Vicenza, Italy (Passaglia 1969a), composition  $(\text{K}_{2.40}\text{Na}_{0.96}\text{Ca}_{1.64}\text{Mg}_{0.64}\text{Sr}_{0.56}\text{Ba}_{0.12})[\text{Al}_{9.08}\text{Fe}_{0.56}\text{Si}_{26.48}\text{O}_{72}] \cdot 25.84\text{H}_2\text{O}$ ,  $T_{\text{Si}} = 0.73$ . Monoclinic,  $C2/m$ ,  $Cm$ , or  $C2$ ,  $a$  17.498,  $b$  17.816,  $c$  7.529 Å,  $\beta$  116.07°.

A close approach to end-member  $\text{K}_9[\text{Al}_9\text{Si}_{27}\text{O}_{72}] \cdot n\text{H}_2\text{O}$  has been reported by Nørnberg (1990).

**Hsianghualite**

$\text{Li}_2\text{Ca}_3[\text{Be}_3\text{Si}_3\text{O}_{12}]_2\text{F}_2$   $Z = 8$  ANA

Huang *et al.* (1958). Type locality unclear, in metamorphosed Devonian limestone, Hunan Province, China. The name is from a Chinese word for fragrant flower. Known from the original locality only. Minor Al, Fe, Mg, Na, and 1.28% loss on ignition reported (Beus 1960).  $T_{\text{Si}} = 0.48$ .

Cubic,  $I_{23}$ ,  $a$  12.864(2) Å.

Has an analcime-type structure, with tetrahedral sites occupied alternately by Si and Be. Extra-framework Ca, Li, and F ions (Rastsvetayeva *et al.* 1991).

**Kalbarsite**

$\text{K}_6[\text{Al}_4\text{Si}_6\text{O}_{20}]\text{B}(\text{OH})_4\text{Cl}$   
 $Z = 2$  ?EDI

Khomyakov *et al.* (1980), Malinovskii & Belov (1980). Type locality: rischorrite pegmatite, Mt. Rasvumchorr, Khibina alkaline massif, Kola Peninsula, Russia. The name alludes to the composition.

Known from two localities in the Khibina massif, both with compositions close to the above formula (Pekov & Chukanov 1996).  $T_{\text{Si}}$  values are 0.59, 0.61.

Tetragonal,  $P4_2/c$ ,  $a$  9.851(5),  $c$  13.060(5) Å.

Framework of Si, Al tetrahedra, with channels along  $c$  containing  $\text{B}(\text{OH})_4$  tetrahedra and K, Cl (Malinovskii & Belov 1980). Considered by Smith (1988) to be an anhydrous analogue of the edingtonite structure-type EDI.

**Laumontite**

$\text{Ca}_4[\text{Al}_9\text{Si}_{16}\text{O}_{48}] \cdot 18\text{H}_2\text{O}$   $Z = 1$  LAU

As lomonite, Jameson (1805), who credits the name to Werner without specific reference; spelling changed to laumonite by Haiiy (1809), and to laumontite by von Leonhard (1821). Named after Gillet de Laumont, who collected material described as "zéolithe efflorescente" by Haiiy (1801, p. 410-412), from lead mines of Huelgoët, Brittany. The later spellings were applied to this material, and the Huelgoët mines are effectively the type locality.

Always Ca-dominant, with minor (K, Na). "Primary leonhardite" of Fersman (1908) is laumontite with approximately 1.5 Ca replaced by 3(K, Na) *apfu* and reduced  $\text{H}_2\text{O}$ .  $T_{\text{Si}}$  in the range 0.64-0.70.

Monoclinic,  $C2/m$  (although reported to be pyroelectric),  $a$  14.845(9),  $b$  13.167(2),  $c$  7.5414(8) Å,  $\beta$  110.34(2)° (Nasik, India: Artioli & Ståhl 1993).

Except where unusually rich in (K, Na), reversibly loses *ca.* 4 $\text{H}_2\text{O}$  at low humidity at room temperature and pressure to form the variety termed "leonhardite" (*e.g.*, Fersman 1908, Armbruster & Kohler 1992); structure refined by Bartl (1970) and others. Si, Al in the framework is highly ordered.


**Leucite**

$\text{K}[\text{AlSi}_2\text{O}_6]$   $Z = 16$  ANA

Blumenbachs (1791), who attributed the name to Werner, who had previously described the mineral as "white garnet". Type locality: Vesuvius, Italy. Named from Greek, meaning white, in reference to color.

Minor substitution of Na for K at low temperatures, and Si in excess of that in the ideal formula, are commonly reported, also significant  $\text{Fe}^{3+}$ .  $T_{\text{Si}}$  in the range 0.66-0.69. Tetragonal,  $I4_1/a$ ,  $a$  13.09,  $c$  13.75 Å (Mazzi *et al.* 1976).

At ordinary temperatures, leucite is invariably finely twinned as a result of a displacive inversion from a cubic polymorph with the structure of analcime, space group  $Ia3d$ , apparently stable above 630°C (Wyart 1938, Peacor 1968). Heaney & Veblen (1990) noted that high leucite inverts to lower symmetry at temperatures between 600° and 750°C depending on the sample, and that there is a tetragonal, metrically cubic form intermediate between high (cubic) and low (tetragonal) forms.

**Levyne (series)**

LEV

Brewster (1825b). Type locality: Dalsnypen, Farøe Islands. Named after Armand Lévy (1794–1841), mathematician and crystallographer, Université de Paris.

Extra-framework cations range from strongly Ca-dominant to strongly Na-dominant, with minor K and, in some cases, minor Sr or Ba; Si:Al is also variable (Galli *et al.* 1981).  $T_{\text{Si}}$  in the range 0.62–0.70.

Hexagonal,  $R\bar{3}m$ ,  $a$  13.32–13.43,  $c$  22.66–23.01 Å.

The stacking of single and double 6-membered rings differs from that in the related structures of erionite and offretite (Merlino *et al.* 1975).

**Levyne-Ca**

New name for the original member of the series; Ca is the most abundant extra-framework cation. Type locality: Dalsnypen, Farøe Islands. Material closely approaching end-member  $\text{Ca}_3[\text{Al}_6\text{Si}_{12}\text{O}_{36}] \cdot 17\text{H}_2\text{O}$  has been reported by England & Ostwald (1979) from near Merriwa, New South Wales, Australia.  $T_{\text{Si}}$  in the range 0.62–0.70.

Hexagonal,  $R\bar{3}m$ ,  $a$  13.338(4),  $c$  23.014(9) Å for composition  $(\text{Ca}_{2.73}, \text{Na}_{0.63}, \text{K}_{0.20})[\text{Al}_{6.31}, \text{Si}_{11.69}, \text{O}_{36}] \cdot 16.66\text{H}_2\text{O}$  from near the Nurri to Orroli road, Nuora, Sardinia (Passaglia *et al.* 1974, Merlino *et al.* 1975).

**Levyne-Na**

New name; Na is the most abundant extra-framework cation.

Proposed type-example: Chojabaru, Nagasaki Prefecture, Japan (Mizota *et al.* 1974).  $T_{\text{Si}}$  in the range 0.65–0.68.

Hexagonal,  $R\bar{3}m$ ,  $a$  13.380(5),  $c$  22.684(9) Å for

$(\text{Na}_{3.84}, \text{K}_{0.38}, \text{Ca}_{0.89}, \text{Mg}_{0.08})[\text{Al}_{6.33}, \text{Si}_{11.71}, \text{O}_{36}]$  (Mizota *et al.* 1974).

**Lovdarite**

Z = 1


LOV

Men'shikov *et al.* (1973). Type locality: alkaline pegmatites on Mt. Karnasurt, Lovozero alkaline massif, Kola Peninsula, Russia. Name means "a gift of Lovozero".

In the type and only known occurrence, approximately 1 Al atom substitutes for Si in the above structure-derived formula, with introduction of additional extra-framework Na and Ca.  $T_{\text{Si}} = 0.75$ .

Orthorhombic,  $Pma2$ , but contains  $b$ -centered domains in which  $a$  is doubled;  $a$  39.576(1),  $b$  6.9308(2),  $c$  7.1526(3) Å (Merlino 1990).

The structure consists of a three-dimensional framework of Si (with minor Al) and Be tetrahedra. It contains three-membered rings, made possible by the presence of Be instead of Si in one of the tetrahedra.

**Maricopaite**

Z = 1

Structure closely related to MOR

Peacor *et al.* (1988). Type locality: Moon Anchor mine, near Tonopah, Maricopa County, Arizona, U.S.A. Named after the locality.

Only one known occurrence.  $T_{\text{Si}} = 0.76$ .

Orthorhombic,  $Cm2m$  (pseudo- $Cmcm$ ),  $a$  19.434(2),  $b$  19.702(2),  $c$  7.538(1) Å (Rouse & Peacor 1994).

Has an interrupted, mordenite-like framework. Pb atoms form  $\text{Pb}_4(\text{O}, \text{OH})_4$  clusters with  $\text{Pb}_4$  tetrahedra within channels (Rouse & Peacor 1994).

**Mazzite**

Z = 1

MAZ

Galli *et al.* (1974). Type locality: in olivine basalt near top of Mont Semiol, south slope, near Montbrison, Loire, France. Named after Fiorenzo Mazzi, Professor of Mineralogy at the University of Pavia, Italy.

A new chemical analysis from the type and only known locality (G. Vezzalini, pers. commun., 1996) gives the above formula (*cf.* Rinaldi *et al.* 1975b).  $T_{\text{Si}} = 0.72$ .

Hexagonal,  $P6_3/mmc$ ,  $a$  18.392(8),  $c$  7.646(2) Å.

The framework is characterized by stacked gmelinite-type cages (Galli 1975), with evidence for limited Si, Al order (Alberti & Vezzalini 1981b).

**Merlinoite**

Z = 1

MER


Passaglia *et al.* (1977). Type locality: Cupaello quarry in kalsilite melilitite, near Santa Rufina, Rieti, Italy. Named after Stefano Merlino, Professor of Crystallography at the University of Pisa.

The two available reliable analyses (Passaglia *et al.* 1977, Della Ventura *et al.* 1993) show strongly K-dominant compositions, with significant Ca, and less Na and Ba;  $T_{\text{Si}}$  in the range 0.66–0.71.

Orthorhombic,  $Immm$ ,  $a$  14.116(7),  $b$  14.229(6),  $c$  9.946(6) Å (Passaglia *et al.* 1977).

The framework is built of double 8-membered rings linked with 4-membered rings (Galli *et al.* 1979). The structure is related to, but different from, that of phillipsite.



**Mesolite**


$$Z = 1$$

NAT

Gehlen & Fuchs (1813), as mesolith, for some varieties of "mesotype" (mostly natrolite) of Häuy (1801). No type locality was given. Fuchs (1816) clarified the distinctions among natrolite, scolecite, and mesolite, and gave analytical data for mesolite from the Faröe Islands, Iceland and Tyrol. The name recognizes its compositional position between natrolite and scolecite.  $(\text{Na} + \text{K})/(\text{Mg} + \text{Ca} + \text{Sr} + \text{Ba})$  varies from 0.45 to 0.52, with K, Mg, Sr, Ba very minor (Alberti *et al.* 1982b).  $T_{\text{Si}}$  in the range 0.59–0.62.

Orthorhombic, *Fdd2*, *a* 18.4049(8), *b* 56.655(6), *c* 6.5443(4) Å, for material from Poona, India (Artioli *et al.* 1986a).

Ordered Si,Al in the framework, with one natrolite-like layer alternating with two scolecite-like layers parallel to (010) (Artioli *et al.* 1986a, Ross *et al.* 1992).

**Montesommaite**

MON

Rouse *et al.* (1990). Type locality: Pollena, Monte Somma, Vesuvius, Italy. Named after the locality. Minor Na was detected in the one published analytical data-set.  $T_{\text{Si}} = 0.70$ .

Orthorhombic, *Fdd2*, *a* = *b* 10.099(1), *c* 17.307(3) Å (pseudotetragonal, *I4<sub>1</sub>lamd*).

The framework can be constructed by linking (100) sheets of five- and eight-membered rings; it has similarities to those of merlinoite and the gismondine group (Rouse *et al.* 1990).

**Mordenite**

$$Z = 1$$

MOR

How (1864). Type locality: shore of Bay of Fundy, 3–5 km east of Morden, King's County, Nova Scotia, Canada. Named after the locality.

The cation content is variable, with  $\text{Na}/(\text{Na} + \text{Ca})$  typically in the range 0.50–0.81. Some K, Mg, Fe, Ba, and Sr also may be present (Passaglia 1975, Passaglia *et al.* 1995). In some examples, K is reported as the dominant cation (Thugutt 1933, Lo *et al.* 1991, Lo & Hsieh 1991), potentially justifying the recognition of a mordenite series with Na- and K-dominant species.  $T_{\text{Si}}$  in the range 0.80–0.86.

Orthorhombic, *Cmcm*, *a* 18.052–18.168, *b* 20.404–20.527, *c* 7.501–7.537 Å (Passaglia 1975).

Structure determined by Meier (1961). Si,Al disorder in the framework is extensive, but not complete.

**Mutinaite**

$$Z = 1$$


MFI

Galli *et al.* (1997b), Vezzalini *et al.* (1997b). Type locality: Mt. Adamson, Northern Victoria Land, Antarctica. The name is for Mutina, the ancient Latin name for Modena, Italy.

Electron-microprobe analyses of mutinaite from the type and only known locality show limited departure from the simplified formula, with minor Mg (~0.21 *apfu*) and K (~0.11 *apfu*). Very high  $T_{\text{Si}} = 0.88$ .

Orthorhombic, *Pnma*, *a* 20.223(7), *b* 20.052(8), *c* 13.491(5) Å.

Mutinaite conforms closely in structure with synthetic zeolite ZSM-5.

**Natrolite**

NAT

Klaproth (1803). Type locality: Hohentwiel, Hegau, Baden-Württemberg, Germany. Name from *natro-* for sodium-bearing.

$(\text{Na} + \text{K})/(\text{Mg} + \text{Ca} + \text{Sr} + \text{Ba})$  varies from 0.97 to 1.00, with K, Mg, Sr, and Ba very minor.  $T_{\text{Si}}$  in the range 0.59–0.62 (Alberti *et al.* 1982b, Ross *et al.* 1992).

Orthorhombic, *Fdd2*, *a* 18.272, *b* 18.613, *c* 6.593 Å (Si,Al highly ordered, Dutoitspan, South Africa: Artioli *et al.* 1984); *a* 18.319(4), *b* 18.595(4), *c* 6.597(1) Å (~70% Si,Al order, Zeilberg, Germany: Hesse 1983).

Si,Al partly to highly ordered (Alberti & Vezzalini 1981a, Ross *et al.* 1992, Alberti *et al.* 1995).

**Offretite**

$$Z = 1$$

OFF

Gonnard (1890) as offrétite. Type locality: Mont Simionse (Mont Semiol), Loire, France. Named after Albert J.J. Offret, professor in the Faculty of Sciences, Lyon, France.


Ca, Mg, and K substantial, commonly in proportions approaching 1 : 1 : 1; Na commonly trace or minor. Passaglia *et al.* (1998) and W. Birch (pers. commun., 1997) show that earlier published analytical data pertaining to apparently Ca- and Na-dominant variants are compromised by identification problems, including possible mixtures.  $T_{\text{Si}}$  in the range 0.69–0.74.

Hexagonal, *P6m2*, *a* 13.307(2), *c* 7.592(2) Å for composition  $(\text{Mg}_{1.06}\text{Ca}_{0.97}\text{K}_{0.88}\text{Sr}_{0.01}\text{Ba}_{0.01})[\text{Al}_{5.26}\text{Si}_{12.81}\text{O}_{36}] \cdot 16.85\text{H}_2\text{O}$  from the type locality (Passaglia & Tagliavini 1994).

The framework is related to those of erionite and levyne, but differs in the stacking of sheets of six-membered rings, resulting in different values for *c* and differently sized and shaped cages (Gard & Tait

1972). A high degree of Si,Al order is inferred. Offretite may contain intergrown macro- or crypto-domains of erionite (*e.g.*, Rinaldi 1976). It forms epitactic intergrowths with chabazite, but epitactic associations with levyne are questionable (Passaglia *et al.* 1998).

### Pahasapaite



Z = 1 RHO


Rouse *et al.* (1987). Type locality: Tip Top mine, Black Hills, South Dakota, U.S.A. Named after Pahasapa, a Sioux Indian name for the Black Hills.

Known from the type locality only.  $T_{\text{Si}} = 0$ .

Cubic, *I23*, *a* 13.781(4) Å.

A beryllophosphate zeolite with ordered  $\text{BeO}_4$  and  $\text{PO}_4$  tetrahedra and a distorted synthetic zeolite RHO-type framework, structurally related to the faujasite series (Rouse *et al.* 1989).

### Parthéite



Z = 4 -PAR

Sarp *et al.* (1979). Type locality: in ophiolitic rocks, 7 km southeast of Doganbaba, Burdur province, Taurus Mountains, southwestern Turkey. Named after Erwin Parthé, Professor of Structural Crystallography, University of Geneva, Switzerland.

Minor Na and K.  $T_{\text{Si}} = 0.52$  and 0.495 in the only two known occurrences.

Monoclinic, *C2/c*, *a* 21.553(3), *b* 8.761(1), *c* 9.304(2) Å,  $\beta$  91.55(2)° (type locality; Engel & Yvon 1984).

The framework contains various 4-, 6-, 8-, and 10-membered rings, and is interrupted at every second  $\text{AlO}_4$  tetrahedron by hydroxyl groups. Si and Al are ordered.

### Paulingite (series)



Z = 16 PAU

Kamb & Oke (1960). Type locality: Rock Island Dam, Columbia River, Wenatchee, Washington, U.S.A. Named after Linus C. Pauling, Nobel Prize winner and Professor of Chemistry, California Institute of Technology.

Electron-microprobe analyses show K as the most abundant cation at three known localities and Ca at two. Significant Ba and Na also are reported (Tschernich & Wise 1982, Lengauer *et al.* 1997).  $T_{\text{Si}}$  in the range 0.73–0.77.

Cubic, *Im3m*, *a* 35.093(2) Å (Gordon *et al.* 1966).

The framework contains several kinds of large polyhedral cages (Gordon *et al.* 1966). The structure has been refined by Bieniok *et al.* (1996) and by Lengauer *et al.* (1997).

### Paulingite-K

New name; K is the most abundant extra-framework cation.


Average composition from five analyses of samples from Rock Island Dam, Washington, U.S.A., the suggested type-example for paulingite-K:  $(\text{K}_{4.44}\text{Na}_{0.95}\text{Ca}_{1.88}\text{Ba}_{0.18})[\text{Al}_{9.82}\text{Si}_{32.21}\text{O}_{84}]\cdot 44\text{H}_2\text{O}$  (Tschernich & Wise 1982); *a* 35.093(2) Å (Gordon *et al.* 1966).

### Paulingite-Ca

New name; Ca is the most abundant extra-framework cation. Average result of four analyses, Ritter, Oregon, U.S.A., the suggested type-locality for paulingite-Ca:  $(\text{Ca}_{3.70}\text{K}_{2.67}\text{Na}_{0.86}\text{Ba}_{0.10})[\text{Al}_{10.78}\text{Si}_{31.21}\text{O}_{84}]\cdot 34\text{H}_2\text{O}$ ; *a* 35.088(6) Å (Tschernich & Wise 1982).

Lengauer *et al.* (1997) found evidence of reduced  $\text{H}_2\text{O}$  content (27  $\text{H}_2\text{O}$  for *Z* = 16) in barian paulingite-Ca from Vinarická Hora, Czech Republic.

### Perlielite



Z = 1 LTL

Men'shikov (1984). Type locality: pegmatites of Mt. Eveslogchorr and Mt. Yukspor, Khibina alkaline massif, Kola Peninsula, Russia. Named after Lily Alekseevna Perekrest, instructor in mineralogy at Kirov Mining Technical School.

Minor substitution by Sr and Ba, but little other compositional variation in the two known occurrences.  $T_{\text{Si}}$  in the range 0.65–0.67.

Hexagonal, *P6/mmm*, *a* 18.49(3), *c* 7.51(1) Å (Men'shikov 1984).

Perlielite has the same framework topology as synthetic zeolite-L (Artioli & Kvick 1990). Structural columns have alternating cancrinite-type cages and double 6-membered rings. No Si,Al order has been detected.

### Phillipsite (series)



Z = 1 PHI

Lévy (1825). Type locality as recorded by Lévy: Aci Reale, now Acireale, on the slopes of Etna, Sicily, Italy. Contemporary literature (see Di Franco 1942) and present-day exposures suggest that the occurrence was probably in basaltic lavas at Aci Castello, nearby. Named after William Phillips (1773–1828), author of geological and mineralogical treatises and a founder of the Geological Society of London.

Either K, Na, Ca, or Ba may be the most abundant extra-framework cation, but the name *harmotome* is retained for the Ba-dominant member of the series. Minor Mg and Sr may be present. In the generalized formula above, *x* ranges from about 4 to about 7.  $T_{\text{Si}}$

varies from approximately 0.56 to 0.77.

Monoclinic,  $P2_1$  or  $P2_1/m$ ,  $a$  9.865(2),  $b$  14.300(4),  $c$  8.668(2) Å,  $\beta$  124.20(3)° (phillipsite-K with substantial Ca from Casal Brunori, Rome, Italy: Rinaldi *et al.* 1974). A pseudo-orthorhombic cell has  $a \approx 9.9$ ,  $b \approx 14.2$ ,  $c \approx 14.2$  Å,  $\beta \approx 90.0^\circ$ ,  $Z = 2$ .

Two cation sites have been identified, one, with two atoms per formula unit fully occupied by K in phillipsite-K and by Ba in harmotome, is surrounded by eight framework atoms of oxygen and four molecules of H<sub>2</sub>O; the other is partly occupied by Ca and Na in distorted octahedral coordination with two framework atoms of oxygen and four molecules of H<sub>2</sub>O (Rinaldi *et al.* 1974). Framework Si,Al largely disordered.

### Phillipsite-Na

New name; Na is the most abundant extra-framework cation.

Na forms 81% of all extra-framework cations in material from Aci Castello, Sicily, Italy, suspected to be the original locality for phillipsite (#6 of Galli & Loschi Ghittoni 1972). Known range in  $T_{Si}$ : 0.64–0.77. For pseudocell,  $a$  9.931–10.003,  $b$  14.142–14.286,  $c$  14.159–14.338 Å,  $\beta$  90°,  $Z = 2$  (e.g., Galli & Loschi Ghittoni 1972, Sheppard & Fitzpatrick 1989).

### Phillipsite-K

New name; K is the most abundant extra-framework cation. Proposed type-locality: Capo di Bove, Rome, Italy (Hintze 1897, #2 of Galli & Loschi Ghittoni 1972). Known range in  $T_{Si}$ : 0.59–0.76.

For the pseudocell,  $a$  9.871–10.007,  $b$  14.124–14.332,  $c$  14.198–14.415 Å,  $\beta$  90°,  $Z = 2$  (e.g., Galli & Loschi Ghittoni 1972, Sheppard *et al.* 1970).

### Phillipsite-Ca

New name; Ca is the most abundant extra-framework cation. Proposed type-locality: Lower Salt Lake Tuff, Puuloa Road near Moanalua Road junction, Oahu, Hawaii (Iijima & Harada 1969).

Known range in  $T_{Si}$ : 0.57–0.74.

For the pseudocell,  $a$  9.859–9.960,  $b$  14.224–14.340,  $c$  14.297–14.362 Å,  $\beta$  90°,  $Z = 2$  (e.g., Galli & Loschi Ghittoni 1972, Passaglia *et al.* 1990).

### Pollucite

(Cs,Na)[AlSi<sub>2</sub>O<sub>6</sub>] $\cdot n$ H<sub>2</sub>O, where (Cs + n) = 1

$Z = 16$  ANA

Breithaupt (1846). Type locality: Elba, Italy. Named "pollux" with coexisting mineral "castor" (a variety of petalite) for twins Castor and Pollux, of Greek mythology; name modified to pollucite by Dana (1868). Forms a series with analcime (Černý 1974) reaching

end-member compositions (Teertstra & Černý 1995).  $T_{Si}$  in the range 0.67–0.74. Minor Rb and Li may be present. Sodian pollucite commonly contains more Si than the simplified formula. The name *pollucite* applies where Cs exceeds Na in atomic proportions.

Cubic,  $Ia3d$ ,  $a$  13.69 Å for Cs<sub>11.7</sub>Na<sub>3.1</sub>Li<sub>0.25</sub>K<sub>0.4</sub>[Al<sub>15</sub>Si<sub>33</sub>O<sub>96.2</sub>] $\cdot$ H<sub>2</sub>O (Beger 1969);  $a$  in the range 3.672(1)–13.674(1) Å for 0.114–0.173 Na *apfu*,  $Z = 16$  (Černý & Simpson 1978).

Si,Al disordered.

### Roggianite

Ca<sub>2</sub>[Be(OH)<sub>2</sub>Al<sub>2</sub>Si<sub>4</sub>O<sub>13</sub>] $\cdot$ <2.5H<sub>2</sub>O

$Z = 8$

–ROG

Passaglia (1969b). Type locality: in albitite dike (E. Passaglia, pers. commun., 1998) at Alpe Rosso in Val Vigezzo about 1.5 km south of Orcesco, Novara Province, Italy. Named after Aldo G. Roggiani, a teacher of natural sciences, who first found the mineral. Contains minor Na and K.

Tetragonal,  $I4/mcm$ ,  $a$  18.33(1),  $c$  9.16(1) Å (Galli 1980).

Contains framework tetrahedrally coordinated Be (Passaglia & Vezzalini 1988) and framework-interrupting (OH) groups (Giuseppetti *et al.* 1991).

### Scolecite

Ca[Al<sub>2</sub>Si<sub>3</sub>O<sub>10</sub>] $\cdot$ 3H<sub>2</sub>O  $Z = 4$  or 8

NAT

Gehlen & Fuchs (1813), as skolezit. Clark (1993) gave the type locality as Berufjord, Iceland, but this is not apparent in the original reference. Fuchs (1816) clarified the distinctions among natrolite, scolecite, and mesolite. He listed occurrences of scolecite as Faröe Islands, Iceland and Staffa (Western Isles, Scotland), with analytical data for specimens from the Faröe Islands and Staffa. Named from Greek *skolex*, worm, for a tendency to curl when heated.

(Na + K)/(Mg + Ca) varies from 0 to 0.16, with very little K, Mg, or other elements.  $T_{Si}$  in the range 0.60–0.62 (Alberti *et al.* 1982b).

Monoclinic,  $Cc$ ,  $a$  6.516(2),  $b$  18.948(3),  $c$  9.761(1) Å,  $\beta$  108.98(1)°,  $Z = 4$  (Bombay, India: Kwick *et al.* 1985), or, by analogy with natrolite, pseudo-orthorhombic  $Fd$ , e.g.,  $a$  18.508(5),  $b$  18.981(5)  $c$  6.527(2) Å,  $\beta$  90.64(1)°,  $Z = 8$  (Berufjord, Iceland: Joswig *et al.* 1984).

The structure is similar to that of natrolite, with a well-ordered Si,Al framework, Ca instead of Na<sub>2</sub>, and an extra molecule of H<sub>2</sub>O.

### Stellerite

Ca[Al<sub>2</sub>Si<sub>7</sub>O<sub>18</sub>] $\cdot$ 7H<sub>2</sub>O  $Z = 8$

STI

Morozewicz (1909). Type locality: Commander Island, Bering Sea. Named after Wilhelm Steller (1709–1746),

natural scientist and military doctor who made important observations on Commander Island.

Variations in composition include up to about 0.2 $apfu$  Na and minor K, Mg, Fe.  $T_{Si}$  in the range 0.75–0.78.

Orthorhombic, *Fmmm*, *a* in the range 13.507–13.605, *b* in the range 18.198–18.270, *c* in the range 17.823–17.863 Å (Passaglia *et al.* 1978b).

The framework is topologically the same as for stilbite, but it has higher symmetry, correlated with fewer extra-framework cations. Only one independent extra-framework site is occupied, and the symmetry is *Fmmm* (Galli & Alberti 1975a). Na-exchanged stellerite retains the *Fmmm* symmetry, unlike the Na zeolite, barrerite, with which it is isochemical (Passaglia & Sacerdoti 1982).

Villarroel (1983) has suggested the occurrence of Na-dominant *Fmmm* stellerite from Roberts Island, South Shetland group.

#### Stilbite (series)



$$Z = 1$$

STI

Häuy (1801, p. 161–166), for minerals, apparently including heulandite, that had previously been described with informal names. He mentioned occurrences in volcanic terranes, and named Iceland, Andreasberg in Harz, Alpes Dauphinoises, and Norway, but there is no clear type-locality. Named from Greek word for mirror, in allusion to its luster (“un certain éclat”).

Ca is almost always the dominant extra-framework cation, accompanied by subordinate Na and minor K and Mg, approximating  $Ca_4(Na, K)$  per formula unit, but Na-rich members also are known.  $T_{Si}$  in the range 0.71 to 0.78.

Monoclinic, *C2/m*, *a* 13.64(3), *b* 18.24(4), *c* 11.27(2) Å,  $\beta$  128.00(25)° (Galli & Gottardi 1966, Galli 1971); an alternative setting is pseudo-orthorhombic, *F2/m*, *Z* = 2.

Increasing departure from the topological symmetry of the orthorhombic framework, *Fmmm*, tends to correlate with increasing content of monovalent cations (Passaglia *et al.* 1978b), which causes the framework to rotate (Galli & Alberti 1975a, b). However, {001} growth sectors with appreciable Na and orthorhombic *Fmmm* symmetry have been observed in crystals in which other isochemical sectors are monoclinic, *C2/m* (Akizuki & Konno 1985, Akizuki *et al.* 1993). The centrosymmetric space-group depends on statistically complete Si,Al disorder, and the true space-group may be noncentrosymmetric (Galli 1971).

#### Stilbite-Ca

New name for common stilbite in which Ca is the most abundant extra-framework cation.

For the pseudo-orthorhombic cell, *F2/m*, *a* 13.595–

13.657, *b* 8.201–18.291, *c* 17.775–17.842 Å,  $\beta$  90.06–90.91° (Passaglia *et al.* 1978b).

#### Stilbite-Na

New name; Na is the most abundant extra-framework cation.


Proposed type-locality: Capo Pula, Cagliari, Sardinia, Italy (Passaglia *et al.* 1978b, #21).

Known examples contain significant Ca and K, and minor Mg, as well as clearly predominant Na.  $T_{Si}$  in the range 0.73–0.78 (Passaglia *et al.* 1978b, Ueno & Hanada 1982, Di Renzo & Gabelica 1997).

Monoclinic, *C2/m*. Using the pseudo-orthorhombic *F2/m* setting, *a* 13.610, *b* 18.330, *c* 17.820 Å,  $\beta$  90.54° for type material, of composition  $(Na_{8.18}K_{1.94}Ca_{3.45}Mg_{0.08})[Al_{16.62}Si_{55.25}O_{144}] \cdot 53.53H_2O$  (Quartieri & Vezzalini 1987).

In spite of the high Na content, the monoclinic *C2/m* symmetry of stilbite is retained, in contrast to stellerite, *Fmmm*, and barrerite, *Amma*.

#### Terranovaite



$$Z = 4$$

TER

Galli *et al.* (1997a). Type locality: Mt. Adamson, Northern Victoria Land, Antarctica. Named after the Italian Antarctic station at Terranova Bay.

Type material contains minor amounts of K and Mg.  $T_{Si} = 0.85$ .

Orthorhombic, *Cmcm*, *a* 9.747(1), *b* 23.880(2), *c* 20.068(2) Å.

The framework topology is not known in other natural or synthetic zeolites. It contains polyhedral units found in laumontite, heulandite, and boggsite.

#### Thomsonite




$$Z = 4$$

THO

Brooke (1820). Type locality: Old Kilpatrick, near Dumbarton, Scotland. Named after Dr. Thomas Thomson (1773–1852), editor of the journal in which the name was published, and who contributed to the improvement of methods of chemical analysis.

Extensive variation in  $Na:(Ca + Sr)$  and  $Si : Al$  approximately according to the formula  $Na_{4+x}(Ca, Sr)_{3-x}[Al_{20-x}Si_{20+x}O_{80}] \cdot 24H_2O$ , where *x* varies from about 0 to 2; small amounts of Fe, Mg, Ba, and K also may be present (Ross *et al.* 1992).  $T_{Si}$  in the range 0.50–0.56. Orthorhombic, *Pncn*, *a* 13.1043(14), *b* 13.0569(18), *c* 13.2463(30) Å (Ståhl *et al.* 1990).

Chains with a repeat unit of five tetrahedra occur as in the NAT structure type, but they are cross-linked in a different way; Si,Al are highly ordered, but disorder increases with increasing Si : Al (Alberti *et al.* 1981).


**Tschernichite**

Smith *et al.* (1991), Boggs *et al.* (1993). Type locality: Goble Creek, 0.2 km north of Goble, Columbia County, Oregon, U.S.A. Named after Rudy W. Tschernich, zeolite investigator of the American Pacific Northwest, who discovered the mineral.

Na, Mg, and K are minor but variable constituents in specimens from the one known locality.  $T_{\text{Si}}$  in the range 0.74–0.78 (0.73, 0.80 in a tschernichite-like mineral from Mt. Adamson, Antarctica: Galli *et al.* 1995).

Tetragonal, possible space-group  $P4/mmm$ ,  $a$  12.880(2),  $c$  25.020(5) Å, but may consist of an intergrowth of a tetragonal enantiomorphic pair with space groups  $P4_322$  and  $P4_322$  and a triclinic polymorph  $P1$ . See also Galli *et al.* (1995).

This is a structural analogue of synthetic zeolite beta.


**Tschörtnerite**

Krause *et al.* (1997), Effenberger *et al.* (1998). Bellberg volcano, near Mayen, Eifel, Germany. Named after Jochen Tschörtner, mineral collector and finder of the mineral.

$T_{\text{Si}} = 0.50$  for the only known occurrence.

Cubic,  $Fm\bar{3}m$ ,  $a$  31.62(1) Å.

Cages in the framework include a large super-cage with 96 tetrahedra and 50 faces. A Cu(OH)-bearing cluster occupies another cage. The framework density is the lowest known for a zeolite with a non-interrupted framework.

**Wairakite**


Steiner (1955), Coombs (1955). Wairakei, Taupo Volcanic Zone, New Zealand. Named after the locality. Most analyzed samples have Na/(Na + Ca) less than 0.3, but wairakite possibly forms a continuous solid-solution series with analcime (Seki & Oki 1969, Seki 1971, Cho & Liou 1987). Other reported substitutions are very minor.  $T_{\text{Si}}$  in the range 0.65–0.69.

Monoclinic (highly ordered),  $I2/a$ ,  $a$  13.692(3),  $b$  13.643(3),  $c$  13.560(3) Å,  $\beta$  90.5(1)° for  $(\text{Ca}_{0.90}\text{Na}_{0.10})[\text{Al}_{1.92}\text{Si}_{4.07}\text{O}_{12}] \cdot 2\text{H}_2\text{O}$  (Takéuchi *et al.* 1979).

Tetragonal or near-tetragonal,  $I4_1/acd$ ,  $a$  13.72(4),  $c$  13.66(4) Å for  $(\text{Ca}_{0.92}\text{Na}_{0.10})[\text{Al}_{1.92}\text{Si}_{4.07}\text{O}_{12}] \cdot 2.11\text{H}_2\text{O}$  (Nakajima 1983).

The framework topology is similar to that of analcime, but Al is preferentially located in a pair of tetrahedral sites associated with Ca, and Ca is in one specific extra-framework site. Smaller departures from cubic symmetry are correlated with decreased Si,Al order.

The name applies to zeolites of ANA structural type in which Ca is the most abundant extra-framework cation, irrespective of the degree of order or space-group symmetry.

**Weinebeneite**

$$Z = 4$$


WEI

Walter (1992). Type locality: vein of spodumene-bearing pegmatite 2 km west of Weinebene Pass, Koralmpe, Carinthia, Austria. Named after the locality.

No elements other than those in the given formula were detected in the one known occurrence.

Monoclinic,  $Cc$ ,  $a$  11.897(2),  $b$  9.707(1),  $c$  9.633(1) Å,  $\beta$  95.76(1)°.

A calcium beryllophosphate zeolite with 3-, 4-, and 8-membered rings in the framework (Walter 1992).

**Willhendersonite**

$$Z = 2$$


CHA

Peacor *et al.* (1984). Type locality: San Venanzo quarry, Terni, Umbria, Italy. Named after Dr. William A. Henderson, of Stamford, Connecticut, U.S.A., who noted this as an unusual mineral and provided it for study.

Type willhendersonite conforms closely to  $\text{KCa}[\text{Al}_3\text{Si}_3\text{O}_{12}] \cdot 5\text{H}_2\text{O}$ . End-member  $\text{Ca}_{1.5}[\text{Al}_3\text{Si}_3\text{O}_{12}] \cdot 5\text{H}_2\text{O}$  and intermediate compositions are now known (Vezzalini *et al.* 1997a).  $T_{\text{Si}} = 0.50, 0.51$ .

Triclinic,  $P1$ ,  $a$  9.206(2),  $b$  9.216(2),  $c$  9.500(4) Å,  $\alpha$  92.34(3)°,  $\beta$  92.70(3)°,  $\gamma$  90.12(3)° (Ettringer Bellerberg, near Mayen, Eifel, Germany; Tillmanns *et al.* 1984).

The framework is the same as for chabazite, which has idealized framework topological symmetry  $R\bar{3}m$ , but with much lower Si and with Si,Al fully ordered. This reduces the topochemical framework symmetry to  $R\bar{3}$ , and the nature and ordering of the extra-framework cations further reduce the framework symmetry to  $P\bar{1}$ . The low-K variants also have fully ordered Si,Al, but are less markedly triclinic (Vezzalini *et al.* 1996).

**Yugawaralite**

YUG

Sakurai & Hayashi (1952). Type locality: Yugawara Hot Springs, Kanagawa Prefecture, Honshu, Japan. Named after the locality.

Reported compositions are close to the ideal stoichiometry, with up to 0.2 *apfu* of Na + K + Sr.  $T_{\text{Si}}$  in the range 0.74–0.76.

Monoclinic,  $Pc$ ,  $a$  6.700(1),  $b$  13.972(2),  $c$  10.039(5) Å,  $\beta$  111.07° (Kvick *et al.* 1986).

Triclinic,  $P1$ , by symmetry reduction ascribed to local Si,Al order, has been reported on the basis of optical measurements (Akizuki 1987b).

Si,Al are strictly ordered in samples from Iceland (Kerr & Williams 1969, Kvick *et al.* 1986). The partial order reported for the Yugawara sample (Leimer & Slaughter 1969) is doubtful (Gottardi & Galli 1985).

#### ZEOLITES OF DOUBTFUL STATUS AND A POSSIBLE ZEOLITE

Further work is recommended to clarify the status of paranatrolite and tetranatrolite. Essential data for these minerals and for tvedalite, which is possibly a beryllosilicate zeolite, are as follows.

#### Paranatrolite

$\text{Na}_2[\text{Al}_2\text{Si}_3\text{O}_{10}]\cdot 3\text{H}_2\text{O}$   $Z = 8$  NAT  
Chao (1980). Type locality, Mont Saint-Hilaire, Quebec, Canada. The name recognizes its association with and similarity in chemical composition to natrolite,  $\text{Na}_2[\text{Al}_2\text{Si}_3\text{O}_{10}]\cdot 2\text{H}_2\text{O}$ .

Contains additional  $\text{H}_2\text{O}$  relative to natrolite, also minor Ca and K.

Pseudo-orthorhombic,  $F^{***}$ , probably monoclinic,  $a$  19.07(1),  $b$  19.13(1),  $c$  6.580(3) Å. Gives very diffuse diffraction-spots, and a powder pattern similar to that of gonnardite (Chao 1980).

Dehydrates to tetranatrolite and could be regarded as overhydrated natrolite, tetranatrolite or gonnardite. Without further justification, separate species status is debatable according to Rule 4.

#### Tetranatrolite

$(\text{Na,Ca})_{16}[\text{Al}_{19}\text{Si}_{21}\text{O}_{80}]\cdot 16\text{H}_2\text{O}$   
 $Z = 0.5$  NAT

Chen & Chao (1980). Type locality: Mont Saint-Hilaire, Quebec, Canada. The name indicates a tetragonal analogue of natrolite. First described as "tetragonal natrolite", from Ilímaussaq, Greenland, by Krogh Andersen *et al.* (1969).

Extensive solid-solution approximating  $\text{Na}_{16-x}\text{Ca}_x\text{Al}_{16+x}\text{Si}_{24-x}\text{O}_{80}\cdot 16\text{H}_2\text{O}$ , where  $x$  varies from about 0.4 to 4, is reported by Ross *et al.* (1992). Small amounts of  $\text{Fe}^{3+}$ , Sr, Ba, and K may replace Na and Ca.  $T_{\text{Si}}$  in the range 0.50–0.59.

Tetragonal,  $I\bar{4}2d$ ,  $a$  13.141,  $c$  6.617 Å (Mont Saint-Hilaire, Quebec, Canada: Ross *et al.* 1992).

The framework is of disordered natrolite type. Tetranatrolite is considered to be a product of dehydration of paranatrolite (Chen & Chao 1980, Ross *et al.* 1992). It differs from natrolite in CaAl substitution for NaSi, as well as in space-group symmetry. These, however, are also characteristics of gonnardite, to which its relationship is debatable.

#### Tvedalite

$(\text{Ca,Mn})_4\text{Be}_3\text{Si}_6\text{O}_{17}(\text{OH})_4\cdot 3\text{H}_2\text{O}$   
 $Z = 2$

Larsen *et al.* (1992). Type locality: Vevya quarry, Tvedalen, Vestfold County, Norway. Named after the locality.

Spot analyses show a range from  $(\text{Ca}_{3.20}\text{Mn}_{0.72}\text{Fe}_{0.08})_{\Sigma 4}$  to  $(\text{Ca}_{2.00}\text{Mn}_{1.86}\text{Fe}_{0.14})_{\Sigma 4}$  for  $\text{Be}_3\text{Si}_6\text{O}_{17}(\text{OH})_4\cdot 3\text{H}_2\text{O}$ , with about 0.9 to 0.21 Al and minor Be substituting for Si in the generalized formula.

Orthorhombic ( $c$ -centered),  $a$  8.724(6),  $b$  23.14(1),  $c$  4.923(4) Å.

Considered to be structurally related to chiavennite, but in the absence of an adequate determination of its structure, it has not been listed here as an accepted zeolite species.

#### DISCREDITED, OBSOLETE, AND OTHER NON-APPROVED ZEOLITE NAMES

**Herschelite**, **leonhardite**, **svetlozarite**, and **wellsite** are discredited as names of mineral species (Appendix 2).

**Kehoeite** was regarded by McConnell (1964) as a zinc phosphate analogue of analcime, but according to White & Erd (1992), type kehoeite is a heterogeneous mixture of quartz and sphalerite with other phases including gypsum and woodhouseite, or a very similar phase. No phase present bears any relationship to analcime. It is not accepted as a valid zeolite species.

**Viséite** is shown by Di Renzo & Gabelica (1995) not to be a zeolite, as had commonly been supposed. They regard it as a defective member of the crandallite group, with composition  $\text{CaAl}_3(\text{PO}_4)_2(\text{SiO}_4)_2(\text{OH})_n\cdot m\text{H}_2\text{O}$ . Kim & Kirkpatrick (1996) showed that a specimen examined by them is very disordered, with a structure similar to that of crandallite, but contains other phases including opal. Viséite is excluded from the list of accepted zeolites.

Obsolete and discredited names are listed below, followed by the correct names or identifications. The list is based on one compiled by the late G. Gottardi, using the following references: Hintze (1897), Dana (1914), Cocco & Garavelli (1958), Davis (1958), Hey (1960, 1962), Merlini (1972), and Strunz (1978). Numerous additions and amendments have been made in the light of more recently published work and of the notes below, and of listings in Clark (1993), in which much information on the history and usages of these names can be found.

abrazite = gismondine, phillipsite

acacialite = chabazite

achiardite = dachiardite

adipite = chabazite?

aedelforsite = laumontite?, stilbite?

aedelite (of Kirwan), aedilite = natrolite

ameletite = mixtures of sodalite, analcime, phillipsite, and relict nepheline

amphigène = leucite

- analcinite = analcime  
 analcite = analcime  
 analzim = analcime  
 andreasbergolite = harmotome  
 andreolite, andréolithe = harmotome  
 antiédrite = edingtonite  
 apoanalcite = natrolite  
 arduinite = mordenite  
 aricite = gismondine  
 ashtonite = strontian mordenite  
 bagotite = thomsonite  
 barium-heulandite = barian heulandite (unless Ba is the most abundant cation)  
 barytkreuzstein = harmotome  
 beaumontite = heulandite  
 bergmannite = natrolite  
 blätterzeolith = heulandite, stilbite  
 brevicite = natrolite  
 cabasite = chabazite  
 caporcianite = laumontite  
 carphostilbite = thomsonite  
 chabasie, chabasite = chabazite  
 christianite (of des Cloizeaux) = phillipsite  
 cluthalite = analcime  
 comptonite = thomsonite  
 crocalite = natrolite  
 cubicite, cubizit = analcime  
 cubic zeolite = analcime?, chabazite  
 cuboite = analcime  
 cuboizite = chabazite  
 desmine = stilbite  
 diagonite = brewsterite  
 dollanite = analcime  
 doranite = analcime with thomsonite, natrolite, and Mg-rich clay minerals (Teertstra & Dyer 1994)  
 echellite = natrolite  
 efflorescing zeolite = laumontite  
 eisenatrolith = natrolite with other mineral inclusions  
 ellagite = a ferriferous natrolite or scolecite?  
 epidesmne = stellerite  
 epinatrolite = natrolite  
 ercinite = harmotome  
 eudnophite = analcime  
 euthalite, euthallite = analcime  
 euzeolith = heulandite  
 falkensteinite = probably plagioclase (Raade 1996)  
 fargite = natrolite  
 faröelite = thomsonite  
 fassaite (of Dolomieu) = probably stilbite  
 feugasite = faujasite  
 flokite, flockit = mordenite  
 foliated zeolite = heulandite, stilbite  
 foresite = stilbite + cookeite  
 galactite = natrolite  
 gibsonite = thomsonite  
 ginzburgite (of Voloshin *et al.*) = roggianite  
 gismondite = gismondine  
 glottalite = chabazite  
 granatite = leucite  
 grenatite (of Daubenton) = leucite  
 grodeckite = gmelinite?  
 hairzeolite (group name) = natrolite, thomsonite, mordenite  
 harmotomite = harmotome  
 harringtonite = thomsonite, mesolite mixture  
 haydenite = chabazite  
 hegautit (högauite) = natrolite  
 hercynite (of Zappe) = harmotome  
 herschelite = chabazite-Na  
 högauite = natrolite  
 hsiang-hua-shih = hsianghualite  
 hydrocastorite = stilbite, mica, petalite mixture  
 hydrolite (of Leman) = gmelinite  
 hydronatrolite = natrolite  
 hydronephelite = a mixture, probably containing natrolite  
 hypodesmine = stilbite  
 hypostilbite = stilbite or laumontite  
 idrocastorite (hydrocastorite) = stilbite, mica, petalite mixture  
 kali-harmotome, kalkharmotome = phillipsite  
 kalithomsonite = ashcroftine (not a zeolite)  
 kalkkreuzstein = phillipsite  
 karphostilbite = thomsonite  
 kehoeite = a mixture including quartz, sphalerite, gypsum, and ?woodhouseite  
 koodilite = thomsonite  
 krokolith = natrolite  
 kubizit = analcime  
 kuboite = analcime  
 laubanite = natrolite  
 laumonite = laumontite  
 ledererite, lederite (of Jackson) = gmelinite  
 lehuntite = natrolite  
 leonhardite = H<sub>2</sub>O-poor laumontite  
 leuzit = leucite  
 levyine, levynite, levyite = levyne  
 lime-harmotome = phillipsite  
 lime-soda mesotype = mesolite  
 lincolmine, lincolnite = heulandite  
 lintonite = thomsonite  
 lomonite = laumontite  
 marburgite = phillipsite  
 mesole = thomsonite  
 mesoline = levyne? chabazite?  
 mesolitine = thomsonite  
 mesotype = natrolite, mesolite, scolecite  
 metachabazite = partially dehydrated chabazite  
 metadesmine = partially dehydrated stilbite  
 metaepistilbite = partially dehydrated epistilbite  
 metaheulandite = partially dehydrated heulandite  
 metalaumontite = partially dehydrated laumontite  
 metaleonhardite = dehydrated "leonhardite" (laumontite)  
 metaleucite = leucite  
 metamesolite = mesolite  
 metanatrolite = partially dehydrated natrolite

- metascolecite, metaskolecit, metaskolezit = partially dehydrated scolecite
- metathomsonite = partially dehydrated thomsonite
- monophane = epistilbite
- mooraboolite = natrolite
- morvenite = harmotome
- natrochabazite = gmelinite
- natron-chabasit, natronchabazit (of Naumann) = gmelinite
- natronite (in part) = natrolite
- needle zeolite, needle stone = natrolite, mesolite, scolecite
- normalin = phillipsite
- orizite, oryzite = epistilbite
- ozarkite = thomsonite
- parastilbite = epistilbite
- phacolite, phakolit(e) = chabazite
- picranalcime = analcime
- picrothomsonite = thomsonite
- pollux = pollucite
- poonahlite, poonalite = mesolite
- portite = natrolite (Franzini & Perchiazzi 1994)
- potassium clinoptilolite = clinoptilolite-K
- pseudolaumontite = pseudomorphs after laumontite
- pseudomesolite = mesolite
- pseudonatrolite = mordenite
- pseudophillipsite = phillipsite
- ptilolite = mordenite
- pufferite, pufferite = stilbite
- punahlite = mesolite
- radiolite (of Esmark) = natrolite
- ranite = gonnardite (Mason 1957)
- reissite (of Fritsch) = epistilbite
- retzite = stilbite?, laumontite?
- sarcolite (of Vauquelin) = gmelinite
- sasbachite, saspachite = phillipsite?
- savite = natrolite
- schabasit = chabazite
- schneiderite = laumontite (Franzini & Perchiazzi 1994)
- schorl blanc = leucite
- scolesite, scolezit = scolecite
- scoulerite = thomsonite
- seebachite = chabazite
- skolezit = scolecite
- sloanite = laumontite?
- snaiderite (schneiderite) = laumontite
- soda-chabazite = gmelinite
- soda mesotype = natrolite
- sodium dachiardite = dachiardite-Na
- sommaite = leucite
- spangite = phillipsite
- sphaerodesmine, sphaerostilbite = thomsonite
- spreustein = natrolite (mostly)
- staurobaryte = harmotome
- steeleite, steelit = mordenite
- stellerycye = stellerite
- stilbite anamorphique = heulandite
- stilbite (of many German authors) = heulandite
- strontium-heulandite = strontian heulandite and heulandite-Sr
- svetlozarite = dachiardite-Ca
- syanhualite, syankhualite = hsianghualite
- syhadrite, syhedrite = impure stilbite?
- tetraedingtonite = edingtonite
- tonsonite = thomsonite
- tripoclase, triploklase = thomsonite
- vanadio-laumontite = vanadian laumontite
- verrucite = mesolite
- Vesuvian garnet = leucite
- Vesuvian (of Kirwan) = leucite
- viséite = disordered crandallite and other phases
- weissian = scolecite
- wellsite = barian phillipsite-Ca and calcian harmotome
- white garnet = leucite
- winchellite = thomsonite
- Würfelzeolith = analcime, chabazite
- zeagonite = gismondine, phillipsite
- zeolite mimetica = dachiardite
- zéolithe efflorescente = laumontite

## ACKNOWLEDGEMENTS

Members of the present Subcommittee, which commenced work in 1993, are grateful to a previous Subcommittee, established in 1979 under the Chairmanship of Professor W.S. Wise of the University of California, Santa Barbara, California, for work contained in a draft report completed in 1987. Members of the 1979 Subcommittee included L. P. van Reeuwijk and the late G. Gottardi and M.H. Hey, as well as D.S.C. and H.M. of the present Subcommittee. J.V. Smith, W.M. Meier, R.W. Tschernich, and the late V.A. Frank-Kamenetskii were consultants. Although recommendations in the present report differ significantly from those in the 1987 report, the existence of that report has greatly facilitated our task. We thank J.V. Smith and L.B. McCusker for advice, C.E.S. Arps and W.D. Birch, successive secretaries of CNMMN, for much help, and many other colleagues for contributions of time, advice, and specimens. Staff of the Science Library, University of Otago, and others, helped trace obscure references.

## REFERENCES

- AKIZUKI, M. (1985): The origin of sector twinning in harmotome. *Am. Mineral.* **70**, 822-828.
- \_\_\_\_\_ (1986): Al-Si ordering and twinning in edingtonite. *Am. Mineral.* **71**, 1510-1514.
- \_\_\_\_\_ (1987a): Crystal symmetry and order-disorder structure of brewsterite. *Am. Mineral.* **72**, 645-648.
- \_\_\_\_\_ (1987b): An explanation of optical variation in yugawaralite. *Mineral. Mag.* **51**, 615-620.



- \_\_\_\_\_ & KONNO, H. (1985): Order-disorder structure and the internal texture of stilbite. *Am. Mineral.* **70**, 814-821.
- \_\_\_\_\_, KUDOH, Y. & KURIBAYASHI, T. (1996): Crystal structures of the {011}, {610}, and {010} growth sectors in brewsterite. *Am. Mineral.* **81**, 1501-1506.
- \_\_\_\_\_, \_\_\_\_\_ & SATOH, Y. (1993): Crystal structure of the orthorhombic {001} growth sector of stilbite. *Eur. J. Mineral.* **5**, 839-843.
- ALBERTI, A. (1972): On the crystal structure of the zeolite heulandite. *Tschermaks Mineral. Petrogr. Mitt.* **18**, 129-146.
- \_\_\_\_\_ (1975a): The crystal structure of two clinoptilolites. *Tschermaks Mineral. Petrogr. Mitt.* **22**, 25-37.
- \_\_\_\_\_ (1975b): Sodium-rich dachiardite from Alpe di Siusi, Italy. *Contrib. Mineral. Petrol.* **49**, 63-66.
- \_\_\_\_\_, CRUCIANI, G. & DAURU, I. (1995): Order-disorder in natrolite-group minerals. *Eur. J. Mineral.* **7**, 501-508.
- \_\_\_\_\_, GALLI, E. & VEZZALINI, G. (1985): Epistilbite: an acentric zeolite with domain structure. *Z. Kristallogr.* **173**, 257-265.
- \_\_\_\_\_, \_\_\_\_\_, \_\_\_\_\_, PASSAGLIA, E. & ZANAZZI, P.F. (1982a): Position of cations and water molecules in hydrated chabazite. Natural and Na-, Ca-, Sr- and K-exchanged chabazites. *Zeolites* **2**, 303-309.
- \_\_\_\_\_, HENTSCHEL, G. & VEZZALINI, G. (1979): Amicite, a new natural zeolite. *Neues Jahrb. Mineral., Monatsh.*, 481-488.
- \_\_\_\_\_, PONGILUPPI, D. & VEZZALINI, G. (1982b): The crystal chemistry of natrolite, mesolite and scolecite. *Neues Jahrb. Mineral., Abh.* **143**, 231-248.
- \_\_\_\_\_ & SABELLI, C. (1987): Statistical and true symmetry of ferrierite: possible absence of straight T-O-T bridging bonds. *Z. Kristallogr.* **178**, 249-256.
- \_\_\_\_\_ & VEZZALINI, G. (1979): The crystal structure of amicite, a zeolite. *Acta Crystallogr.* **B35**, 2866-2869.
- \_\_\_\_\_ & \_\_\_\_\_ (1981a): A partially disordered natrolite: relationships between cell parameters and Si-Al distribution. *Acta Crystallogr.* **B37**, 781-788.
- \_\_\_\_\_ & \_\_\_\_\_ (1981b): Crystal energies and coordination of ions in partially occupied sites: dehydrated mazzite. *Bull. Minéral.* **104**, 5-9.
- \_\_\_\_\_, \_\_\_\_\_, GALLI, E. & QUARTIERI, S. (1996): The crystal structure of gottardiite, a new natural zeolite. *Eur. J. Mineral.* **8**, 69-75.
- \_\_\_\_\_, \_\_\_\_\_ & TAZZOLI, V. (1981): Thomsonite: a detailed refinement with cross checking by crystal energy calculations. *Zeolites* **1**, 91-97.
- ALIETTI, A. (1972): Polymorphism and crystal-chemistry of heulandites and clinoptilolites. *Am. Mineral.* **57**, 1448-1462.
- ARMBRUSTER, T. (1993): Dehydration mechanism of clinoptilolite and heulandite: single-crystal X-ray study of Na-poor, Ca-, K-, Mg-rich clinoptilolite at 100 K. *Am. Mineral.* **78**, 260-264.
- \_\_\_\_\_ & KOHLER, T. (1992): Re- and dehydration of laumontite: a single-crystal X-ray study at 100 K. *Neues Jahrb. Mineral., Monatsh.*, 385-397.
- ARTIOLI, G. (1992): The crystal structure of garronite. *Am. Mineral.* **77**, 189-196.
- \_\_\_\_\_ & FOY, H. (1994): Gobbinsite from Magheramorne quarry, Northern Ireland. *Mineral. Mag.* **58**, 615-620.
- \_\_\_\_\_, GOTTARDI, G., RINALDI, R., SATOH, Y., HORIUCHI, H., YE, J., SAWADA, H., TANAKA, M. & TOKONAMI, M. (1987): A single crystal diffraction study of the natural zeolite cowlesite. *Photon Factory, National Laboratory for High Energy Physics, Activity Rep.* **1987**, 316.
- \_\_\_\_\_ & KVICK, Å. (1990): Synchrotron X-ray Rietveld study of perialite, the natural counterpart of synthetic zeolite-L. *Eur. J. Mineral.* **2**, 749-759.
- \_\_\_\_\_, RINALDI, R., KVICK, Å. & SMITH, J.V. (1986b): Neutron diffraction structure refinement of the zeolite gismondine at 15 K. *Zeolites* **6**, 361-366.
- \_\_\_\_\_, SMITH, J.V. & KVICK, Å. (1984): Neutron diffraction study of natrolite, Na<sub>2</sub>Al<sub>2</sub>Si<sub>3</sub>O<sub>10</sub>•2H<sub>2</sub>O, at 20 K. *Acta Crystallogr.* **C40**, 1658-1662.
- \_\_\_\_\_, \_\_\_\_\_ & PLUTH, J.J. (1986a): X-ray structure refinement of mesolite. *Acta Crystallogr.* **C42**, 937-942.
- \_\_\_\_\_ & STÄHL, K. (1993): Fully hydrated laumontite: a structure study by flat-plate and capillary powder diffraction techniques. *Zeolites* **13**, 249-255.
- \_\_\_\_\_ & TORRES SALVADOR, M.R. (1991): Characterization of the natural zeolite gonnardite. Structure analysis of natural and cation exchanged species by the Rietveld method. *Material Science Forum* **79-82**, 845-850.
- BARTL, H. (1970): Strukturverfeinerung von Leonhardt, Ca[Al<sub>2</sub>Si<sub>4</sub>O<sub>22</sub>]•3H<sub>2</sub>O, mittels Neutronenbeugung. *Neues Jahrb. Mineral., Monatsh.*, 298-310.
- BAUR, W.H. (1964): On the cation and water positions in faujasite. *Am. Mineral.* **49**, 697-704.
- BEGER, R.M. (1969): The crystal structure and chemical composition of pollucite. *Z. Kristallogr.* **129**, 280-302.
- BERGERHOFF, G., BAUR, W.H. & NOWACKI, W. (1958): Über die Kristallstruktur des Faujasits. *Neues Jahrb. Mineral., Monatsh.*, 193-200.

- BEUS, A.A. (1960): *Geochemistry of Beryllium and the Genetic Types of Beryllium Deposits*. Akademii Nauk, SSSR, Inst. mineral., geokhim., i kristallochim. redkikh elementov, 1-329 (in Russian). Abstract in *Am. Mineral.* **46**, 244.
- BIENIOK, A., JOSWIG, W. & BAUR, W.H. (1996): A study of paulingites: pore filling by cations and water molecules. *Neues Jahrb. Mineral., Abh.* **171**, 119-134.
- BISSERT, G. & LIEBAU, F. (1986): The crystal structure of a triclinic bikitaite,  $\text{Li}[\text{AlSi}_2\text{O}_6] \cdot \text{H}_2\text{O}$ , with ordered Al/Si distribution. *Neues Jahrb. Mineral., Monatsh.*, 241-252.
- BLACKBURN, W.H. & DENNEN, W.H. (1997): Encyclopedia of Minerals Names. *Can. Mineral., Spec. Publ.* **1**.
- BLUMENBACHS, J.F. (1791): Auszüge und Kezensionen bergmanischer und mineralogischer Schriften. *Bergmannisches J.* **2**, 489-500.
- BOGGS, R.C., HOWARD, D.G., SMITH, J.V. & KLEIN, G.L. (1993): Tschernichite, a new zeolite from Goble, Columbia County, Oregon. *Am. Mineral.* **78**, 822-826.
- BOLES, J.R. (1972): Composition, optical properties, cell dimensions, and thermal stability of some heulandite group zeolites. *Am. Mineral.* **57**, 1463-1493.
- BONARDI, M. (1979): Composition of type dachiardite from Elba: a re-examination. *Mineral. Mag.* **43**, 548-549.
- \_\_\_\_\_, ROBERTS, A.C. & SABINA, A.P. (1981): Sodium-rich dachiardite from the Francon quarry, Montreal Island, Quebec. *Can. Mineral.* **19**, 285-289.
- BONDI, M., GRIFFIN, W.L., MATTIOLI, V. & MOTTANA, A. (1983): Chiavennite,  $\text{CaMnBe}_2\text{Si}_5\text{O}_{13}(\text{OH})_2 \cdot 2\text{H}_2\text{O}$ , a new mineral from Chiavenna (Italy). *Am. Mineral.* **68**, 623-627.
- BOSC D'ANTIC, L. (1792): Mémoire sur la chabazite. *J. d'Histoire Naturelle* **2**, 181-184.
- BREITHAUP, A. (1846): Pollux. (*Poggendorff's Annalen der Physik und Chemie* **69**, 439.
- BREWSTER, D. (1825a): Description of gmelinite, a new mineral species. *Edinburgh J. Sci.* **2**, 262-267.
- \_\_\_\_\_. (1825b): Description of levyne, a new mineral species. *Edinburgh J. Sci.* **2**, 332-334.
- BROOKE, H.J. (1820): On mesotype, needlestone, and thomsonite. *Annals of Philosophy* **16**, 193-194.
- \_\_\_\_\_. (1822): On the comptonite of Vesuvius, the brewsterite of Scotland, the stilbite and the heulandite. *Edinburgh Philos. J.* **6**, 112-115.
- CABELLA, R., LUCCHETTI, G., PALENZONA, A., QUARTIERI, S. & VEZZALINI, G. (1993): First occurrence of a Ba-dominant brewsterite: structural features. *Eur. J. Mineral.* **5**, 353-360.
- ČERNÝ, P. (1974): The present status of the analcime-pollucite series. *Can. Mineral.* **12**, 334-341.
- \_\_\_\_\_, RINALDI, R. & SURDAM, R.C. (1977): Wellsite and its status in the phillipsite - harmotome group. *Neues Jahrb. Mineral., Abh.* **128**, 312-320.
- \_\_\_\_\_ & SIMPSON, F.M. (1978): The Tanco pegmatite at Bernic Lake, Manitoba. X. Pollucite. *Can. Mineral.* **16**, 325-333.
- CHAO, G.Y. (1980): Paranatrolite, a new zeolite from Mont St-Hilaire, Québec. *Can. Mineral.* **18**, 85-88.
- CHEN, T.T. & CHAO, G.Y. (1980): Tetranatrolite from Mont St-Hilaire, Québec. *Can. Mineral.* **18**, 77-84.
- CHO, M. & LIOU, J.G. (1987): Prehnite-pumpellyite to greenschist facies transition in the Karmutsen metabasites, Vancouver Island, B.C. *J. Petrol.* **28**, 417-443.
- CLARK, A.M. (1993): *Hey's Mineral Index*. Chapman & Hall, London, U.K.
- COCCO, G. & GARAVELLI, C. (1958): Riesame di alcune zeoliti elbane. *Atti Soc. Toscana Sci. Naturali* **65**, 262-283.
- COOMBS, D.S. (1955): X-ray observations on wairakite and non-cubic analcime. *Mineral. Mag.* **30**, 699-708.
- \_\_\_\_\_ & WHETTEN, J.T. (1967): Composition of analcime from sedimentary and burial metamorphic rocks. *Geol. Soc. Am., Bull.* **78**, 269-282.
- CRONSTEDT, A.F. (1756): Observation and description of an unknown kind of rock to be named zeolites. *Kongl. Vetenskaps Acad. Handl. Stockholm* **17**, 120-123 (in Swedish).
- D'ACHIARDI, G. (1906): Zeoliti del filone della Speranza presso S. Piero in Campo (Elba). *Atti Soc. Toscana Sci. Naturali* **22**, 150-165.
- DAMOUR, M. (1842): Description de la faujasite, nouvelle espèce minérale. *Annales des Mines, Sér.* **4**, **1**, 395-399.
- DANA, E.S. (1914): *A System of Mineralogy of J.D. Dana* (6th ed., with Appendices I and II). John Wiley & Sons, New York, N.Y.
- DANA, J.D. (1868): *A System of Mineralogy* (5th ed.). John Wiley & Sons, New York, N.Y.
- DAVIS, R.J. (1958): Mordenite, ptilolite, flokite, and arduinite. *Mineral. Mag.* **31**, 887-888.
- DE GENNARO, M. & FRANCO, E. (1976): La K-chabazite di alcuni "Tufi del Vesuvio". *Rend. Accad. Naz. Lincei* **40**, 490-497.
- DELAMÉTHÉRIE, J.-C. (1795): *Théorie de la Terre* **1**. Chez Maradan, Paris, France.
- DELLA VENTURA, G., PARODI, G.C. & BURRAGATO, F. (1993): New data on merlinoite and related zeolites. *Rend. Lincei Sci. Fisiche Naturali, Ser.* **9**, **4**, 303-312.

- DI FRANCO, S. (1942): *Mineralogia Etna*. Zuccarello & Izzi, Catania, Italy (158-161).
- DI RENZO, F. & GABELICA, Z. (1995): New data on the structure and composition of the silicoaluminophosphate viséite and a discreditation of its status as a zeolite. In *Natural Zeolites '93: Occurrence, Properties, Use* (D.W. Ming & F.A. Mumpton, eds.). International Committee on Natural Zeolites, Brockport, New York, N.Y. (173-185).
- \_\_\_\_\_ & \_\_\_\_\_ (1997): Barrerite and other zeolites from Kuiu and Kupreanof islands, Alaska. *Can. Mineral.* **35**, 691-698.
- DUNN, P.J., PEACOR, D.R., NEWBERRY, N. & RAMIK, R.A. (1980): Goosecreekite, a new calcium aluminum silicate hydrate possibly related to brewsterite and epistilbite. *Can. Mineral.* **18**, 323-327.
- EAKLE, A.S. (1898): Erionite, a new zeolite. *Am. J. Sci.* **156**, 66-68.
- EBERLY, P.E., JR. (1964): Adsorption properties of naturally occurring erionite and its cationic-exchanged forms. *Am. Mineral.* **49**, 30-40.
- EFFENBERGER, H., GIESTER, G., KRAUSE, W. & BERNHARDT, H.-J. (1998): Tschörtnerite, a copper-bearing zeolite from the Bellberg volcano, Eifel, Germany. *Am. Mineral.* **83**, 607-617.
- ENGEL, N. & YVON, K. (1984): The crystal structure of parthéite. *Z. Kristallogr.* **169**, 165-175.
- ENGLAND, B.M. & OSTWALD, J. (1979): Levyne – offretite intergrowths from Tertiary basalts in the Merriwa district, Hunter Valley, New South Wales, Australia. *Aust. Mineral.* **25**, 117-119.
- ERCIT, T.S. & VAN VELTHUIZEN, J. (1994): Gaultite, a new zeolite-like mineral species from Mont Saint-Hilaire, Quebec, and its crystal structure. *Can. Mineral.* **32**, 855-863.
- FERSMAN, A.E. (1908): Materialien zur Untersuchung der Zeolithe Russlands. I. Leonhardt und Laumontit aus der Umgebung von Simferopol (Krim). *Trav. du Musée géol. Pierre le Grand pr. l'Acad. Imp. de Science St Pétersbourg* **2**, 103-150 (abstr. in *Z. Kristallogr.* **50**, 75-76).
- FISCHER, K. (1966): Untersuchung der Kristallstruktur von Gmelinit. *Neues Jahrb. Mineral., Monatsh.*, 1-13.
- \_\_\_\_\_ & SCHRAMM, V. (1970): Crystal structure of gismondite, a detailed refinement. In *Molecular Sieve Zeolites*. *Am. Chem. Soc., Adv. Chem. Ser.* **101**, 250-258.
- FRANZINI, M. & PERCHIAZZI, N. (1994): Portite discredited = natrolite and new data on "schneiderite" (= laumontite). *Eur. J. Mineral.* **6**, 351-353.
- FUCHS, J.N. (1816): Ueber die Zeolithe. (*Schweigger's*) *J. Chem. und Phys.* **18**, 1-29.
- GALLI, E. (1971): Refinement of the crystal structure of stilbite. *Acta Crystallogr.* **B27**, 833-841.
- \_\_\_\_\_ (1975): Crystal structure refinement of mazzite. *Rend. Soc. It. Mineral. Petrol.* **31**, 599-612.
- \_\_\_\_\_ (1976): Crystal structure refinement of edingtonite. *Acta Crystallogr.* **B32**, 1623-1627.
- \_\_\_\_\_ (1980): The crystal structure of roggianite, a zeolite-like silicate. *Proc. 5th Int. Conf. on Zeolites* (L.V.C. Rees, ed.). Heyden, London, U.K. (205-213).
- \_\_\_\_\_ & ALBERTI, A. (1975a): The crystal structure of stellerite. *Bull. Soc. fr. Minéral. Cristallogr.* **98**, 11-18.
- \_\_\_\_\_ & \_\_\_\_\_ (1975b): The crystal structure of barrerite. *Bull. Soc. fr. Minéral. Cristallogr.* **98**, 331-340.
- \_\_\_\_\_ & GOTTARDI, G. (1966): The crystal structure of stilbite. *Mineral. Petrogr. Acta (Bologna)* **12**, 1-10.
- \_\_\_\_\_, \_\_\_\_\_ & PONGLUPPI, D. (1979): The crystal structure of the zeolite merlinoite. *Neues Jahrb. Mineral., Monatsh.*, 1-9.
- \_\_\_\_\_ & LOSCHI GHITTONI, A.G. (1972): The crystal chemistry of phillipsites. *Am. Mineral.* **57**, 1125-1145.
- \_\_\_\_\_, PASSAGLIA, E., PONGLUPPI, D. & RINALDI, R. (1974): Mazzite, a new mineral, the natural counterpart of the synthetic zeolite  $\Omega$ . *Contrib. Mineral. Petrol.* **45**, 99-105.
- \_\_\_\_\_, \_\_\_\_\_ & ZANAZZI, P.F. (1982): Gmelinite: structural refinements of sodium-rich and calcium-rich natural crystals. *Neues Jahrb. Mineral., Monatsh.*, 145-155.
- \_\_\_\_\_, QUARTIERI, S., VEZZALINI, G. & ALBERTI, A. (1995): Boggsite and tschernichite-type zeolites from Mt. Adamson, Northern Victoria Land (Antarctica). *Eur. J. Mineral.* **7**, 1029-1032.
- \_\_\_\_\_, \_\_\_\_\_, \_\_\_\_\_ & \_\_\_\_\_ (1996): Gottardiite, a new high-silica zeolite from Antarctica: the natural counterpart of synthetic NU-87. *Eur. J. Mineral.* **8**, 687-693.
- \_\_\_\_\_, \_\_\_\_\_, \_\_\_\_\_, \_\_\_\_\_ & FRANZINI, M. (1997a): Terranovaite from Antarctica: a new 'pentasil' zeolite. *Am. Mineral.* **82**, 423-429.
- \_\_\_\_\_ & RINALDI, R. (1974): The crystal chemistry of epistilbites. *Am. Mineral.* **59**, 1055-1061.
- \_\_\_\_\_, \_\_\_\_\_ & MODENA, C. (1981): Crystal chemistry of levynes. *Zeolites* **1**, 157-160.
- \_\_\_\_\_, VEZZALINI, G., QUARTIERI, S., ALBERTI, A. & FRANZINI, M. (1997b): Mutinaite, a new zeolite from Antarctica: the natural counterpart of ZSM-5. *Zeolites* **19**, 318-322.
- GARD, J.A. & TAIT, J.M. (1972): The crystal structure of the zeolite offretite,  $K_{1.1}Ca_{1.1}Mg_{0.7}[Si_{12.8}Al_{5.2}O_{36}] \cdot 15.2H_2O$ . *Acta Crystallogr.* **B28**, 825-834.

- GEHLEN, A.F. & FUCHS, J.N. (1813): Ueber Werner's Zeolith, Hatly's Mesotype und Stilbite. (*Schweigger's*) *J. Chem. und Phys.* **8**, 353-366.
- GISMONDI, C.G. (1817): Osservazioni sopra alcuni fossili particolari de' contorni di Roma. *Giornale Enciclopedico di Napoli, Anno XI*, **2**, 3-15.
- GIUSEPPETTI, G., MAZZI, F., TADINI, C. & GALLI, E. (1991): The revised crystal structure of roggianite:  $\text{Ca}_2[\text{Be}(\text{OH})_2\text{Al}_2\text{Si}_4\text{O}_{13}] \cdot 2.5\text{H}_2\text{O}$ . *Neues Jahrb. Mineral., Monatsh.*, 307-314.
- GONNARD, F. (1890): Sur l'offrétite, espèce minérale nouvelle. *C.R. Acad. Sci., Paris* **111**, 1002-1003.
- GORDON, E.K., SAMSON, S. & KAMB, W.B. (1966): Crystal structure of the zeolite paulingite. *Science* **154**, 1004-1007.
- GOTTARDI, G. & ALBERTI, A. (1974): Domain structure in garronite: a hypothesis. *Mineral. Mag.* **39**, 898-899.
- \_\_\_\_\_ & GALLI, E. (1985): *Natural Zeolites*. Springer-Verlag, Berlin, Germany.
- \_\_\_\_\_ & MEIER, W.M. (1963): The crystal structure of dachiardite. *Z. Kristallogr.* **119**, 53-64.
- GRAHAM, R.P.D. (1918): On ferrierite, a new zeolitic mineral, from British Columbia; with notes on some other Canadian minerals. *Trans. R. Soc. Can., Ser. 3*, **12**, 185-201.
- GRAMLICH-MEIER, R., GRAMLICH, V. & MEIER, W.M. (1985): The crystal structure of the monoclinic variety of ferrierite. *Am. Mineral.* **70**, 619-623.
- HADINGER, W. (1825): Description of edingtonite, a new mineral species. *Edinburgh J. Sci.* **3**, 316-320.
- HARADA, K., IWAMOTO, S. & KIHARA, K. (1967): Erionite, phillipsite and gonnardite in the amygdales of altered basalt from Mazé, Niigata Prefecture, Japan. *Am. Mineral.* **52**, 1785-1794.
- HASSAN, I. (1997): Feldspathoids and their relationships to zeolites. *Kuwait J. Sci. Engineering* **24**, 163-187.
- HAÛY, R.-J. (1797): Analcime. *J. des Mines* **5**, 278-279.
- \_\_\_\_\_ (1801): *Traité de minéralogie* **3**. Chez Louis, Paris, France.
- \_\_\_\_\_ (1809): *Tableau Comparatif des Résultats de Cristallographie et de l'Analyse Chimique Relativement à la Classification des Minéraux*. Courcier, Paris, France.
- HAZEN, R.M. & FINGER, L.W. (1979): Polyhedral tilting: a common type of pure displacive phase transition and its relationship to analcite at high pressure. *Phase Transitions* **1**, 1-22.
- HEANEY, P.J. & VEBLEN, D.R. (1990): A high-temperature study of the low-high leucite phase transition using the transmission electron microscope. *Am. Mineral.* **75**, 464-476.
- HESSE, K.-F. (1983): Refinement of a partially disordered natrolite,  $\text{Na}_2\text{Al}_2\text{Si}_3\text{O}_{10} \cdot \text{H}_2\text{O}$ . *Z. Kristallogr.* **163**, 69-74.
- HEY, M.H. (1930): Studies on the zeolites. I. General review. *Mineral. Mag.* **22**, 422-437.
- \_\_\_\_\_ (1960): Glottalite is chabazite. *Mineral. Mag.* **32**, 421-422.
- \_\_\_\_\_ (1962): *An Index of Mineral Species and Varieties Arranged Chemically* (2nd ed., and Appendices, 1963 and 1974). British Museum, London, U.K.
- HINTZE, C. (1897): *Handbuch der Mineralogie* **2**. Von Veit, Leipzig, Germany.
- HORI, H., NAGASHIMA, K., YAMADA, M., MIYAWAKI, R. & MARUBASHI, T. (1986): Ammonioleucite, a new mineral from Tatarazawa, Fujioka, Japan. *Am. Mineral.* **71**, 1022-1027.
- HOW, H. (1864): On mordenite, a new mineral from the trap of Nova Scotia. *J. Chem. Soc.* **17** (new ser.), 17, 100-104.
- HOWARD, D.G. (1994): Crystal habit and twinning of garronite from Fara Vicentina, Vicenza (Italy). *Neues Jahrb. Mineral., Monatsh.*, 91-96.
- \_\_\_\_\_, TSCHERNICH, R.W., SMITH, J.V. & KLEIN, G.L. (1990): Boggisite, a new high-silica zeolite from Goble, Columbia County, Oregon. *Am. Mineral.* **75**, 1200-1204.
- HUANG, WEN-HUI, TU, SHAO-HUA, WANG, K'UNG-HAI, CHAO, CHUN-LIN & YU, CHENG-CHIH (1958): Hsiang-hua-shih, a new beryllium mineral. *Ti-chih-yueh-k'an* **7**, 35 (abstr. in *Am. Mineral.* **44**, 1327-1328; **46**, 244).
- HURLBUT, C.S., JR. (1957): Bikitaite,  $\text{LiAlSi}_2\text{O}_6 \cdot \text{H}_2\text{O}$ , a new mineral from Southern Rhodesia. *Am. Mineral.* **42**, 792-797.
- IBRAHIM, K. & HALL, A. (1995): New occurrences of diagenetic faujasite in the Quaternary tuffs of north-east Jordan. *Eur. J. Mineral.* **7**, 1129-1135.
- ILIUMA, A. & HARADA, K. (1969): Authigenic zeolites in zeolitic palagonite tuffs on Oahu, Hawaii. *Am. Mineral.* **54**, 182-197.
- JAMESON, R. (1805): *System of Mineralogy* **II**. Bell and Bradford, Edinburgh, U.K. (539).
- JOSWIG, W., BARTL, H. & FUESS, H. (1984): Structure refinement of scolecite by neutron diffraction. *Z. Kristallogr.* **166**, 219-223.
- KAMB, W.B. & OKE, W.C. (1960): Paulingite, a new zeolite, in association with erionite and filiform pyrite. *Am. Mineral.* **45**, 79-91.
- KAWAHARA, A. & CURIEN, H. (1969): La structure cristalline de l'ériónite. *Bull. Soc. fr. Minéral. Cristallogr.* **92**, 250-256.
- KERR, I.S. & WILLIAMS, D.J. (1969): The crystal structure of yugawaralite. *Acta Crystallogr.* **B25**, 1183-1190.

- KHOMYAKOV, A.P., CHEREPIVSKAYA, G.YE., KUROVA, T.A. & KAPTSOV, V.V. (1982): Amicite,  $K_2Na_2Al_4Si_4O_{16} \cdot 5H_2O$ , first find in the USSR. *Dokl. Akad. Nauk SSSR* **263**, 978-980 (in Russ.).
- \_\_\_\_\_, SANDOMIRSKAYA, S.M. & MALINOVSKII, YU.A. (1980): Kalborsite,  $K_2BaAl_4Si_4O_{20}(OH)_4Cl$ , a new mineral. *Dokl. Akad. Nauk SSSR* **252**, 1465-1468 (in Russ.).
- KIM, Y. & KIRKPATRICK, R.J. (1996): Application of MAS NMR spectroscopy to poorly crystalline materials: viséite. *Mineral. Mag.* **60**, 957-962.
- KLAPROTH, M.H. (1803): Chemische Untersuchung des Natroliths. *Ges. Naturforschender Freunde zu Berlin, Neue Schriften* **4**, 243-248.
- KOCMAN, V., GAIT, R.I. & RUCKLIDGE, J. (1974): The crystal structure of bikitaite,  $Li[AlSi_2O_6] \cdot H_2O$ . *Am. Mineral.* **59**, 71-78.
- KOYAMA, K. & TAKÉUCHI, Y. (1977): Clinoptilolite: the distribution of potassium atoms and its role in thermal stability. *Z. Kristallogr.* **145**, 216-239.
- KRAUSE, W., BERNHARDT, H.-J., EFFENBERGER, H. & GIESTER, G. (1997): Tschörnerite, a copper-bearing zeolite from the Bellberg volcano, Eifel, Germany. *Ber. Deutsch. Mineral. Ges.* **1**, 205 (abstr.).
- KROGH ANDERSEN, E., DANØ, M. & PETERSEN, O.V. (1969): A tetragonal natrolite. *Medd. om Grønland* **181**, 1-19.
- KVICK, Å., ARTIOLI, G. & SMITH, J.V. (1986): Neutron diffraction study of the zeolite yugawaralite at 13 K. *Z. Kristallogr.* **174**, 265-281.
- \_\_\_\_\_, STÄHL, K. & SMITH, J.V. (1985): A neutron diffraction study of the bonding of zeolitic water in scolecite at 20 K. *Z. Kristallogr.* **171**, 141-154.
- LACROIX, A. (1896): Sur la gonnardite. *Bull. Soc. fr. Minéral.* **19**, 426-429.
- LANGHOF, J. & HOLSTAM, D. (1994): Boron-bearing chiavennite and other late-stage minerals of the Proterozoic lithium-pegmatites of Utö, Stockholm, Sweden. *Int. Mineral. Assoc., 16th Gen. Meet. (Pisa), Abstr.*, 232.
- LARSEN, A.O., ÅSHEIM, A., RAADE, G. & TAFTØ, J. (1992): Tvedalite,  $(Ca,Mn)_4Be_3Si_6O_{17}(OH)_4 \cdot 3H_2O$ , a new mineral from syenite pegmatite in the Oslo region, Norway. *Am. Mineral.* **77**, 438-443.
- LEIMER, H.W. & SLAUGHTER, M. (1969): The determination and refinement of the crystal structure of yugawaralite. *Z. Kristallogr.* **130**, 88-111.
- LENGAUER, C.L., GIESTER, G. & TILLMANN, E. (1997): Mineralogical characterization of paulingite from Vinarická Hora, Czech Republic. *Mineral. Mag.* **61**, 591-606.
- VON LEONHARD, K.C. (1817): Die Zeagonit, ein neues Mineral vom Capo do Bove bei Rom. *Taschenbuch für die gesammte Mineralogie mit Hinsicht auf die neuesten Entdeckungen* **11**, 164-168 (extracted from Gismondi 1817).
- \_\_\_\_\_, (1821): *Handbuch der Oryktognosie*. Mohr & Winter, Heidelberg, Germany (448).
- LEVINSON, A.A. (1966): A system of nomenclature for rare-earth minerals. *Am. Mineral.* **51**, 152-158.
- LÉVY, A. (1825): Descriptions of two new minerals. *Annals of Philosophy, new ser.* **10**, 361-363.
- LO, H.-J. & HSIEH, Y.-L. (1991): High potassium natural mordenite and the chemical variation of mordenite. *Proc. Geol. Soc. China (Taiwan)* **34**, 305-312.
- \_\_\_\_\_, SONG, S.-R. & WEN, S.-B. (1991): High potassium mordenite in the andesite from the Coastal Range, eastern Taiwan. *Proc. Geol. Soc. China (Taiwan)* **34**, 293-304.
- LUCCHETTI, G., MASSA, B. & PENCO, A.M. (1982): Strontian heulandite from Campegli (eastern Ligurian ophiolites, Italy). *Neues Jahrb. Mineral., Monatsh.*, 541-550.
- MALINOVSKII, YU.A. (1984): The crystal structure of K-gmelinite. *Kristallografiya* **29**, 426-430 (in Russ.).
- \_\_\_\_\_, & BELOV, N.V. (1980): Crystal structure of kalborsite. *Dokl. Akad. Nauk SSSR* **252**, 611-615 (in Russ.).
- MASON, B.H. (1957): Gonnardite (ranite) from Langesundsfjord. *Norsk geol. Tidsskr.* **37**, 435-437.
- MAZZI, F. & GALLI, E. (1978): Is each analcime different? *Am. Mineral.* **63**, 448-460.
- \_\_\_\_\_, & \_\_\_\_\_ (1983): The tetrahedral framework of chabazite. *Neues Jahrb. Mineral., Monatsh.*, 461-480.
- \_\_\_\_\_, \_\_\_\_\_ & GOTTARDI, G. (1976): The crystal structure of tetragonal leucite. *Am. Mineral.* **61**, 108-115.
- \_\_\_\_\_, \_\_\_\_\_ & \_\_\_\_\_ (1984): Crystal structure refinement of two tetragonal edingtonites. *Neues Jahrb. Mineral., Monatsh.*, 373-382.
- \_\_\_\_\_, LARSEN, A.O., GOTTARDI, G. & GALLI, E. (1986): Gonnardite has the tetrahedral framework of natrolite: experimental proof with a sample from Norway. *Neues Jahrb. Mineral., Monatsh.*, 219-228.
- MCCONNELL, D. (1964): A zinc phosphate analogue of analcime: kehoite. *Mineral. Mag.* **33**, 799-803.
- MCCUSKER, L.B., BAERLOCHER, C. & NAWAZ, R. (1985): Rietveld refinement of the crystal structure of the new zeolite mineral gobbinsite. *Z. Kristallogr.* **171**, 281-289.
- MEIER, W.M. (1961): The crystal structure of mordenite (ptilolite). *Z. Kristallogr.* **115**, 439-450.
- \_\_\_\_\_, OLSON, D.H. & BAERLOCHER, C. (1996): Atlas of zeolite structure types. *Zeolites* **17**, 1-230.

- MEN'SHIKOV, YU.P. (1984): Perllialite,  $K_3Na(Ca,Sr)[Al_{12}Si_{24}O_{72}] \cdot 15H_2O$ , a new potassian zeolite from the Khibina Massif. *Zap. Vses. Mineral. Obshchest.* **113**, 607-612.
- \_\_\_\_\_, DENISOV, A.P., USPENSKAYA, E.I. & LIPATOVA, E.A. (1973): Lovdarite, a new hydrous beryllsilicate of alkalis. *Dokl. Akad. Nauk SSSR* **213**, 429-432 (in Russ.).
- MERKLE, A.B. & SLAUGHTER, M. (1968): Determination and refinement of the structure of heulandite. *Am. Mineral.* **53**, 1120-1138.
- MERLINO, S. (1972): Orizite discredited (= epistilbite). *Am. Mineral.* **57**, 592-593.
- \_\_\_\_\_. (1974): The crystal structure of wenkite. *Acta Crystallogr.* **B30**, 1262-1268.
- \_\_\_\_\_. (1990): Lovdarite,  $K_4Na_{12}(Be_3Si_{28}O_{72}) \cdot 18H_2O$ , a zeolite-like mineral: structural features and OD character. *Eur. J. Mineral.* **2**, 809-817.
- \_\_\_\_\_, GALLI, E. & ALBERTI, A. (1975): The crystal structure of levyne. *Tschermaks Mineral. Petrogr. Mitt.* **22**, 117-129.
- MINATO, H. & TAKANO, T. (1964): An occurrence of potassium clinoptilolite from Itaya, Yamagata Prefecture, Japan. *J. Clay Sci. Soc. Japan* **4**, 12-22 (in Japanese with English abstr.).
- MIZOTA, T., SHIBUYA, G., SHIMAZU, M. & TAKESHITA, Y. (1974): Mineralogical studies on levyne and erionite from Japan. *Geol. Soc. Japan, Mem.* **11**, 283-290.
- MOROZEWICZ, J. (1909): Über Stellerit, ein neues Zeolithmineral. *Bull. International de l'Académie des Sciences de Cracovie*, 344-359.
- MUMPTON, F.A. (1960): Clinoptilolite redefined. *Am. Mineral.* **45**, 351-369.
- NAKAJIMA, W. (1983): Disordered wairakite from Hikihara, Haga Town, Hyōgo Prefecture. *Bull. Faculty of Education, Kobe Univ.* **70**, 39-46.
- NAWAZ, R. (1983): New data on gobbinsite and garronite. *Mineral. Mag.* **47**, 567-568.
- \_\_\_\_\_. (1984): New data on cowlesite from Northern Ireland. *Mineral. Mag.* **48**, 565-566.
- \_\_\_\_\_ & MALONE, J.F. (1982): Gobbinsite, a new zeolite mineral from Co. Antrim, N. Ireland. *Mineral. Mag.* **46**, 365-369.
- NICKEL, E.H. & MANDARINO, J.A. (1987): Procedures involving the IMA Commission on New Minerals and Mineral Names, and guidelines on mineral nomenclature. *Can. Mineral.* **25**, 353-377.
- NØRNBERG, P. (1990): A potassium-rich zeolite in soil development on Danian chert. *Mineral. Mag.* **54**, 91-94.
- OGIHARA, S. & IJIMA, A. (1990): Exceptionally K-rich clinoptilolite – heulandite group zeolites from three offshore boreholes off northern Japan. *Eur. J. Mineral.* **2**, 819-826.
- PASSAGLIA, E. (1969a): Le zeoliti di Albergo Bassi (Vicenza). *Per. Mineral.* **38**, 237-243.
- \_\_\_\_\_. (1969b): Roggianite, a new silicate mineral. *Clay Minerals* **8**, 107-111.
- \_\_\_\_\_. (1970): The crystal chemistry of chabazites. *Am. Mineral.* **55**, 1278-1301.
- \_\_\_\_\_. (1975): The crystal chemistry of mordenites. *Contrib. Mineral. Petrol.* **50**, 65-77.
- \_\_\_\_\_, ARTIOLI, G. & GUALTIERI, A. (1998): The crystal chemistry of the zeolites erionite and offretite. *Am. Mineral.* **83**, 577-589.
- \_\_\_\_\_, \_\_\_\_\_, \_\_\_\_\_ & CARNEVALI, R. (1995): Diagenetic mordenite from Ponza, Italy. *Eur. J. Mineral.* **7**, 429-438.
- \_\_\_\_\_, GALLI, E., LEONI, L. & ROSSI, G. (1978b): The crystal chemistry of stilbites and stellerites. *Bull. Minéral.* **101**, 368-375.
- \_\_\_\_\_, \_\_\_\_\_ & RINALDI, R. (1974): Levynes and erionites from Sardinia, Italy. *Contrib. Mineral. Petrol.* **43**, 253-259.
- \_\_\_\_\_ & PONGILUPPI, D. (1974): Sodian stellerite from Capo Pula, Sardegna. *Lithos* **7**, 69-73.
- \_\_\_\_\_ & \_\_\_\_\_ (1975): Barrerite, a new natural zeolite. *Mineral. Mag.* **40**, 208.
- \_\_\_\_\_, \_\_\_\_\_ & RINALDI, R. (1977): Merlinoite, a new mineral of the zeolite group. *Neues Jahrb. Mineral., Monatsh.*, 355-364.
- \_\_\_\_\_, \_\_\_\_\_ & VEZZALINI, G. (1978a): The crystal chemistry of gmelinites. *Neues Jahrb. Mineral., Monatsh.*, 310-324.
- \_\_\_\_\_ & SACERDOTI, M. (1982): Crystal structural refinement of Na-exchanged stellerite. *Bull. Minéral.* **105**, 338-342.
- \_\_\_\_\_ & TAGLIAVINI, A. (1994): Chabazite – offretite epitaxial overgrowths in cornubianite from Passo Forcel Rosso, Adamello, Italy. *Eur. J. Mineral.* **6**, 397-405.
- \_\_\_\_\_ & VEZZALINI, G. (1985): Crystal chemistry of diagenetic zeolites in volcanoclastic deposits of Italy. *Contrib. Mineral. Petrol.* **90**, 190-198.
- \_\_\_\_\_ & \_\_\_\_\_ (1988): Roggianite: revised chemical formula and zeolitic properties. *Mineral. Mag.* **52**, 201-206.

- \_\_\_\_\_, \_\_\_\_\_ & CARNEVALI, R. (1990): Diagenetic chabazites and phillipsites in Italy: crystal chemistry and genesis. *Eur. J. Mineral.* **2**, 827-839.
- PEACOR, D.R. (1968): A high temperature single crystal diffractometer study of leucite,  $(K,Na)AlSi_2O_6$ . *Z. Kristallogr.* **127**, 213-224.
- \_\_\_\_\_, DUNN, P.J., SIMMONS, W.B., TILLMANN, E. & FISCHER, R.X. (1984): Willhendersonite, a new zeolite isostructural with chabazite. *Am. Mineral.* **69**, 186-189.
- \_\_\_\_\_, \_\_\_\_\_, WICKS, F.J. & RAUDSEPP, M. (1988): Maricopaite, a new hydrated Ca-Pb, zeolite-like silicate from Arizona. *Can. Mineral.* **26**, 309-313.
- PEKOV, I.V. & CHUKANOV, N.V. (1996): New data on kalborsite. *Zap. Vses. Mineral. Obshchest.* **125**(4), 55-59 (in Russ.).
- PERROTTA, A.J. (1967): The crystal structure of epistilbite. *Mineral. Mag.* **36**, 480-490.
- \_\_\_\_\_, SMITH, J.V. (1964): The crystal structure of brewsterite,  $(Sr,Ba,Ca)_2(Al_4Si_{12}O_{32}) \cdot 10H_2O$ . *Acta Crystallogr.* **17**, 857-862.
- PIRSSON, L.V. (1890): On mordenite. *Am. J. Sci.* **140**, 232-237.
- PLUTH, J.J. & SMITH, J.V. (1990): Crystal structure of boggsite, a new high-silica zeolite with the first three-dimensional channel system bounded by both 12- and 10-rings. *Am. Mineral.* **75**, 501-507.
- \_\_\_\_\_, \_\_\_\_\_, HOWARD, D.G. & TSCHERNICH, R.W. (1989): Boggsite; the first three-dimensional channel system with both 12- and 10-rings. In *Zeolites for the Nineties, Recent Research Reports* (J.C. Jansen, L. Moscou & M.F.M. Post, eds.), Eighth Int. Zeolite Conf. (Amsterdam), 111-112 (abstr.).
- QUARTIERI, S. & VEZZALINI, G. (1987): Crystal chemistry of stilbites: structure refinements of one normal and four chemically anomalous samples. *Zeolites* **7**, 163-170.
- \_\_\_\_\_, \_\_\_\_\_ & ALBERTI, A. (1990): Dachiardite from Hokiya-dake: evidence of a new topology. *Eur. J. Mineral.* **2**, 187-193.
- RAADE, G. (1996): Minerals originally described from Norway. Including notes on type material. *Norsk Bergverksmuseums Skrifiserie* **11**.
- \_\_\_\_\_, ÅMLI, R., Mladeck, M.H., DIN, V.K., LARSEN, A.O. & ÅSHEIM, A. (1983): Chiavennite from syenite pegmatites in the Oslo region, Norway. *Am. Mineral.* **68**, 628-633.
- RASTSVETAeva, R.K., REKHLOVA, O.Y., ANDRIANOV, V.I. & MALINOVSKI, YU.A. (1991): Crystal structure of hsianghualite. *Dokl. Akad. Nauk SSSR* **316**, 624-628 (in Russ.).
- RINALDI, R. (1976): Crystal chemistry and structural epitaxy of offretite - erionite from Sasbach, Kaiserstuhl. *Neues Jahrb. Mineral., Monatsh.*, 145-156.
- \_\_\_\_\_, PLUTH, J.J. & SMITH, J.V. (1974): Zeolites of the phillipsite family. Refinement of the crystal structures of phillipsite and harmotome. *Acta Crystallogr.* **B30**, 2426-2433.
- \_\_\_\_\_, \_\_\_\_\_ & \_\_\_\_\_ (1975b): Crystal structure of mazzite dehydrated at 600°C. *Acta Crystallogr.* **B31**, 1603-1608.
- \_\_\_\_\_, SMITH, J.V. & JUNG, G. (1975a): Chemistry and paragenesis of faujasite, phillipsite and offretite from Sasbach, Kaiserstuhl, Germany. *Neues Jahrb. Mineral., Monatsh.*, 433-443.
- \_\_\_\_\_, \_\_\_\_\_ & VEZZALINI, G. (1985): Gismondine; the detailed X-ray structure refinement of two natural samples. In *Zeolites; Synthesis, Structure, Technology and Application* (B. Držaj, S. Hočevar & S. Pejovnik, eds.). Elsevier, Amsterdam, The Netherlands (481-492).
- ROBINSON, G.W. & GRICE, J.D. (1993): The barium analog of brewsterite from Harrisville, New York. *Can. Mineral.* **31**, 687-690.
- ROSE, G. (1826): Ueber den Epistilbit, eine neue zur Familie der Zeolithe gehörige Mineralgattung. (*Poggendorff's Annalen der Physik und Chemie* **6**, 183-190.
- ROSS, C.S. & SHANNON, E.V. (1924): Mordenite and associated minerals from near Challis, Custer County, Idaho. *Proc. U.S. Nat. Museum* **64**(19), 1-19.
- ROSS, M., FLOHR, M.J.K. & ROSS, D.R. (1992): Crystalline solution series and order-disorder within the natrolite mineral group. *Am. Mineral.* **77**, 685-703.
- ROUSE, R.C., DUNN, P.J., GRICE, J.D., SCHLENKER, J.L. & HIGGINS, J.B. (1990): Montesommaite,  $(K,Na)_9Al_9Si_{23}O_{64} \cdot 10H_2O$ , a new zeolite related to merlinoite and the gismondine group. *Am. Mineral.* **75**, 1415-1420.
- \_\_\_\_\_, \_\_\_\_\_ & PEACOR, D.R. (1986): Crystal structure of the zeolite mineral goosecreekite,  $CaAl_2Si_6O_{16} \cdot 5H_2O$ . *Am. Mineral.* **71**, 1494-1501.
- \_\_\_\_\_, \_\_\_\_\_ & \_\_\_\_\_ (1994): Maricopaite, an unusual lead calcium zeolite with an interrupted mordenite-like framework and intrachannel  $Pb_4$  tetrahedral clusters. *Am. Mineral.* **79**, 175-184.
- \_\_\_\_\_, \_\_\_\_\_, DUNN, P.J., CAMPBELL, T.J., ROBERTS, W.L., WICKS, F.J. & NEWBURY, D. (1987): Pahasapaite, a beryllophosphate zeolite related to synthetic zeolite rho, from the Tip Top pegmatite of South Dakota. *Neues Jahrb. Mineral., Monatsh.*, 433-440.
- \_\_\_\_\_, \_\_\_\_\_ & MERLINO, S. (1989): Crystal structure of pahasapaite, a beryllophosphate mineral with a distorted zeolite rho framework. *Am. Mineral.* **74**, 1195-1202.
- RÜDINGER, B., TILLMANN, E. & HENTSCHEL, G. (1993): Bellbergite - a new mineral with the zeolite structure type EAB. *Mineral. Petrol.* **48**, 147-152.

- SAKURAI, K. & HAYASHI, A. (1952): "Yugawaralite", a new zeolite. *Sci. Rep., Yokohama Nat. Univ., Ser. II*, 1, 69-77.
- SARP, H., DEFERNE, J., BIZOUARD, H. & LIEBICH, B.W. (1979): La parthéite,  $\text{CaAl}_2\text{Si}_2\text{O}_8 \cdot 2\text{H}_2\text{O}$ , un nouveau silicate naturel d'aluminium et de calcium. *Schweiz. Mineral. Petrogr. Mitt.* 59, 5-13.
- SCHALLER, W.T. (1923): Ptilolite and related zeolites. In Proc. Societies (E.T. Wherry, ed.). *Am. Mineral.* 8, 93-94.
- \_\_\_\_\_ (1932): The mordenite - ptilolite group; clinoptilolite, a new species. *Am. Mineral.* 17, 128-134.
- SCHLENKER, J.L., PLUTH, J.J. & SMITH, J.V. (1977a): Refinement of the crystal structure of brewsterite,  $\text{Ba}_{0.3}\text{Sr}_{1.5}\text{Al}_4\text{Si}_{12}\text{O}_{32} \cdot 10\text{H}_2\text{O}$ . *Acta Crystallogr.* B33, 2907-2910.
- \_\_\_\_\_, \_\_\_\_\_ & \_\_\_\_\_ (1977b): Dehydrated natural erionite with stacking faults of the offretite type. *Acta Crystallogr.* B33, 3265-3268.
- SCHRÖPFER, L. & JOSWIG, W. (1997): Structure analyses of a partially dehydrated synthetic Ca-garronite single crystal under different T, pH<sub>2</sub>O conditions. *Eur. J. Mineral.* 9, 53-65.
- SEKI, Y. (1971): Wairakite - analcime solid solution as an indicator of water pressures in low-grade metamorphism. *J. Geol. Soc. Japan* 77, 667-674.
- \_\_\_\_\_ & OKI, Y. (1969): Wairakite - analcime solid solutions from low-grade metamorphic rocks of the Tanzawa Mountains, central Japan. *Mineral. J.* 6, 36-45.
- SHEPPARD, R.A. & FITZPATRICK, J.J. (1989): Phillipsite from silicic tuffs in saline, alkaline-lake deposits. *Clays Clay Minerals* 37, 243-247.
- \_\_\_\_\_ & GUDE, A.J., III (1969a): Diagenesis of tuffs in the Barstow Formation, Mud Hills, San Bernardino County, California. *U.S. Geol. Surv., Prof. Pap.* 634, 1-35.
- \_\_\_\_\_ & \_\_\_\_\_ (1969b): Chemical composition and physical properties of the related zeolites offretite and erionite. *Am. Mineral.* 54, 875-886.
- \_\_\_\_\_, \_\_\_\_\_ & GRIFFIN, J.J. (1970): Chemical composition and physical properties of phillipsite from the Pacific and Indian oceans. *Am. Mineral.* 55, 2053-2062.
- \_\_\_\_\_, \_\_\_\_\_ & MUNSON, E.L. (1965): Chemical composition of diagenetic zeolites from tuffaceous rocks of the Mojave Desert and vicinity, California. *Am. Mineral.* 50, 244-249.
- SMITH, J.V. (1988): Topochemistry of zeolites and related materials. 1. Topology and geometry. *Chem. Rev.* 88, 149-182.
- \_\_\_\_\_, KNAWELS, C.R. & RINALDI, R. (1964): Crystal structures with a chabazite framework. III. Hydrated Ca-chabazite at +20 and -150°C. *Acta Crystallogr.* 17, 374-384.
- \_\_\_\_\_, PLUTH, J.J., BOGGS, R.C. & HOWARD, D.G. (1991): Tschernichite, the mineral analogue of zeolite beta. *J. Chem. Soc., Chem. Commun.*, 363-364.
- STÄHL, K., KVICK, Å. & SMITH, J.V. (1990): Thomsonite, a neutron diffraction study at 13 K. *Acta Crystallogr.* C46, 1370-1373.
- STEINER, A. (1955): Wairakite, the calcium analogue of analcime, a new zeolite mineral. *Mineral. Mag.* 30, 691-698.
- STRUNZ, H. (1956): Die Zeolithe Gmelinit, Chabasit, Levyn (Phakolith, Herschelit, Seebachit, Offretit). *Neues Jahrb. Mineral., Monatsh.*, 250-259.
- \_\_\_\_\_ (1978): *Mineralogische Tabellen* (7th ed.). Akademische Verlagsges., Leipzig, Germany.
- STUCKENSCHMIDT, E., FUESS, H. & KVICK, Å. (1990): Investigation of the structure of harmotome by X-ray (293 K, 100 K) and neutron diffraction (15 K). *Eur. J. Mineral.* 2, 861-874.
- TAKÉUCHI, Y., MAZZI, F., HAGA, N. & GALLI, E. (1979): The crystal structure of wairakite. *Am. Mineral.* 64, 993-1001.
- TAYLOR, W.H. (1930): The structure of analcite ( $\text{NaAlSi}_2\text{O}_6 \cdot \text{H}_2\text{O}$ ). *Z. Kristallogr.* 74, 1-19.
- \_\_\_\_\_ & JACKSON, R. (1933): The structure of edingtonite. *Z. Kristallogr.* 86, 53-64.
- TAZZOLI, V., DOMENEGHETTI, M.C., MAZZI, F. & CANNILLO, E. (1995): The crystal structure of chiavennite. *Eur. J. Mineral.* 7, 1339-1344.
- TEERTSTRA, D.K. & ČERNÝ, P. (1995): First natural occurrences of end-member pollucite: a product of low-temperature reequilibration. *Eur. J. Mineral.* 7, 1137-1148.
- \_\_\_\_\_ & DYER, A. (1994): The informal discreditation of "doranite" as the magnesium analogue of analcime. *Zeolites* 14, 411-413.
- \_\_\_\_\_, SHERRIFF, B.L., XU, ZHI & ČERNÝ, P. (1994): MAS and DOR NMR study of Al-Si order in the analcime - pollucite series. *Can. Mineral.* 32, 69-80.
- THUGUTT, S.J. (1933): O ptylocie z Mydzka na Wolyniu - Sur la ptilolite de Mydzk en Volhynie. *Arch. Minéral., Soc. Sci. Varsovie* 9, 99-102 (Polish), 103-104 (French résumé). (*Mineral. Abstr.* 6, 129).
- TILLMANN, E., FISCHER, R.X. & BAUR, W.H. (1984): Chabazite-type framework in the new zeolite willhendersonite,  $\text{KCaAl}_3\text{Si}_3\text{O}_{12} \cdot 5\text{H}_2\text{O}$ . *Neues Jahrb. Mineral., Monatsh.*, 547-558.
- TSCHERNICH, R.D. & WISE, W.S. (1982): Paulingite: variations in composition. *Am. Mineral.* 67, 799-803.
- UENO, T. & HANADA, K. (1982): Chemical compositions and geneses of zeolites from Tsuyazaki, Fukuoka Prefecture, Japan. *J. Mineral. Soc. Japan* 15, 259-272 (in Japanese, with English abstr.).



- VAUGHAN, P.A. (1966): The crystal structure of the zeolite ferrierite. *Acta Crystallogr.* **21**, 983-990.
- VEZZALINI, G. (1984): A refinement of Elba dachiardite: opposite acentric domains simulating a centric structure. *Z. Kristallogr.* **166**, 63-71.
- \_\_\_\_\_, ARTIOLI, G. & QUARTIERI, S. (1992): The crystal chemistry of cowlesite. *Mineral. Mag.* **56**, 575-579.
- \_\_\_\_\_, & OBERTI, R. (1984): The crystal chemistry of gismondines: the non-existence of K-rich gismondines. *Bull. Minéral.* **107**, 805-812.
- \_\_\_\_\_, QUARTIERI, S. & GALLI, E. (1996): Relazioni strutturali nelle zeoliti con topologia CHA alla luce del ritrovamento di una Ca-willhendersonite. *XXVI Congresso Nazionale della Associazione Italiana di Cristallografia (Alessandria)*, 91 (abstr.).
- \_\_\_\_\_, \_\_\_\_\_ & \_\_\_\_\_ (1997a): Occurrence and crystal structure of a Ca-pure willhendersonite. *Zeolites* **19**, 75-79.
- \_\_\_\_\_, \_\_\_\_\_, \_\_\_\_\_, ALBERTI, A., CRUCIANI, G. & KVICK, Å. (1997b): Crystal structure of the zeolite mutinaite, the natural analogue of ZSM-5. *Zeolites* **19**, 323-325.
- \_\_\_\_\_, \_\_\_\_\_ & PASSAGLIA, E. (1990): Crystal structure of a K-rich natural gmelinite and comparison with the other refined gmelinite samples. *Neues Jahrb. Mineral., Monatsh.*, 504-516.
- VILLARROEL, H.S. (1983): Sobre la existencia de otras variedades de estellerita y una forma de reconocerlas. *Anais Academia Brasileira de Ciencias* **55**, 87-91.
- WALKER, G.P.L. (1962): Garronite, a new zeolite, from Ireland and Iceland. *Mineral. Mag.* **33**, 173-186.
- WALTER, F. (1992): Weinebeneite,  $\text{CaBe}_3(\text{PO}_4)_2(\text{OH})_2 \cdot 4\text{H}_2\text{O}$ , a new mineral species: mineral data and crystal structure. *Eur. J. Mineral.* **4**, 1275-1283.
- WENK, H.-R. (1973): The structure of wenkite. *Z. Kristallogr.* **137**, 113-126.
- WHITE, J.S. & ERD, R.C. (1992): Kehoeite is *not* a valid species. *Mineral. Mag.* **56**, 256-258.
- WISE, W.S. (1982): New occurrence of faujasite in southeastern California. *Am. Mineral.* **67**, 794-798.
- \_\_\_\_\_, & TSCHERNICH, R.W. (1975): Cowlesite, a new Ca-zeolite. *Am. Mineral.* **60**, 951-956.
- \_\_\_\_\_, & \_\_\_\_\_ (1976): Chemical composition of ferrierite. *Am. Mineral.* **61**, 60-66.
- WYART, J. (1938): Étude sur la leucite. *Bull. Soc. fr. Minéral.* **61**, 228-238.
- YANG, PING & ARMBRUSTER, T. (1996): (010) disorder, partial Si,Al ordering, and Ca distribution in triclinic (C1) epistilbite. *Eur. J. Mineral.* **8**, 263-271.

Received December 3, 1997.

## APPENDIX 1. NOTES ON THE DEFINITION OF A ZEOLITE

### *Is more than 50% substitution of elements other than Si and Al permissible in tetrahedral sites?*

There was complete agreement in the Subcommittee that some substitution of elements such as P and Be for Si and Al in tetrahedral sites must be permitted in the definition. Discussion in this context focussed on whether a 50% rule should be applied. The so-called 50% rule (Nickel 1992) is normally applied to split a binary solid-solution series into two species at the half-way point according to the predominant cations concerned, but not to separate members of a solid-solution series into two separate classes of minerals, as could happen if applied in the present context. Proponents of a 50% rule argued that the definition of zeolites should be on grounds of both structure and composition, zeolites being aluminosilicates or possibly Al-free silicates. The contrary opinion is that where structures are topologically equivalent and other essentially identical zeolitic characteristics prevail, irrespective of Si and Al contents in tetrahedral sites, any restrictions based on specific Si and Al contents would be arbitrary and undesirable. The Subcommittee voted by a substantial majority for this view. The beryllsilicates lovdarite and chiavennite, like the zincosilicate gaultite, have more than 50% tetrahedral sites occupied by Si, and are here accepted as zeolites in spite of having little if any Al. Also included are the beryllphosphates pahasapaite and weinebeneite, which have neither Si nor Al, but have typically zeolitic structures and other zeolitic characteristics. They can be regarded as end-member examples of Si-free zeolites or zeolite phosphates.

A compositional factor is included in the adopted definition in that the framework consists essentially of oxygen atoms together with cations that enter into tetrahedral co-ordination with oxygen.

### *Is the presence of H<sub>2</sub>O and of extra-framework cations essential?*

Reversible dehydration is a characteristic feature of zeolitic behavior, but how much H<sub>2</sub>O must be present for a mineral to be considered a zeolite? Pollucite forms a continuous series with analcime, the H<sub>2</sub>O content declining progressively with increasing Cs content such that the Na-free, Cs member is essentially anhydrous. It seems unnecessary, impractical, and illogical to prescribe some arbitrary H<sub>2</sub>O content below which pollucite (or other mineral) would be defined as anhydrous, and no longer a zeolite. Furthermore, it is not inconceivable that some typical zeolite might be reversibly dehydrated under natural conditions without essential loss of structure. If so, it has not ceased to be a zeolite. Although zeolites typically are hydrous, it is inexpedient to specify the presence of H<sub>2</sub>O in the definition.

Natural zeolites are known with up to 88% of tetrahedral sites occupied by Si, as in mutinaite, and there is no theoretical reason why this figure cannot be exceeded. If the site occupancy of tetrahedra by Si approaches 100%, the extra-framework cation content will approach zero, even though the structure and other characteristics may remain typically zeolitic. It is again considered inexpedient to word the definition so as to exclude such a hypothetical end-member case from the zeolite category. Melanophlogite, a low-density SiO<sub>2</sub> phase with large cages in its framework, would be a possible example, but is otherwise excluded by the adopted definition because it lacks appropriate channels for the passage of guest species.

## REFERENCE

- NICKEL, E.H. (1992): Solid solutions in mineral nomenclature. *Can. Mineral.* **30**, 231-234.

## APPENDIX 2. DISCREDITATIONS

### *Herschelite is chabazite-Na*

Herschelite, Na[AlSi<sub>2</sub>O<sub>6</sub>]•3H<sub>2</sub>O, was named by Lévy (1825) from material brought to him by Herschel from "Aci Reale" (now Acireale), on the flanks of Mt. Etna in Sicily. Contemporary literature and present-day exposures suggest that the actual occurrence may have been in basaltic lavas at Aci Castello, nearby. Lévy described herschelite as tabular crystals of hexagonal outline that contain "silex, alumina, and potash". It was later identified with chabazite (*e.g.*, Hausmann 1847) and relegated to synonymy, although shown to be Na-rich, not K-rich. Strunz (1956) confirmed that herschelite and chabazite give essentially identical X-ray powder patterns. Mason (1962) proposed reevaluation on the bases of a supposed compositional gap between herschelite and "normal" Ca-rich chabazite, the distinctive habit, and lower indices of refraction.

Passaglia (1970) demonstrated a continuum of compositions from Ca- to Na-dominant types, extending into the field of K-dominance in a ternary series; there is no discernible gap in composition. The lower indices of refraction reflect the Na-rich composition. Variant crystal habit is not an accepted basis for species status for minerals, and some examples of strongly Na-dominant chabazite have rhombohedral, not tabular habit, as in the case of micrometer-scale crystals aggregated into thin ragged plates, illustrated by Sheppard *et al.* (1978).

In view of its chequered history and the above considerations, the name herschelite is suppressed, and the name chabazite-Na is to be applied to those members of the chabazite series in which Na is the most abundant extra-framework cation. Herschelite may retain some use as a term for a distinctive habit.

*Leonhardite is H<sub>2</sub>O-poor laumontite*

Leonhardite  $\text{Ca}_4[\text{Al}_8\text{Si}_{16}\text{O}_{48}] \cdot \sim 14\text{H}_2\text{O}$  was described by Blum (1843) for a mineral closely related to laumontite  $\text{Ca}_4[\text{Al}_8\text{Si}_{16}\text{O}_{48}] \cdot 18\text{H}_2\text{O}$ , but with different morphology. The type locality was near Schemnitz, nowadays Banska Stiavnica, then in Hungary, now in Slovakia. Delffs (1843) showed that type-locality leonhardite has less  $\text{H}_2\text{O}$  (ca. 13 molecules of  $\text{H}_2\text{O}$  per formula unit) than laumontite. Doelter (1921) agreed that leonhardite is identical in composition to laumontite, apart from its lower content of  $\text{H}_2\text{O}$ . The name has continued to be used widely for a material that forms rapidly and reversibly by partial dehydration of laumontite under ambient conditions. This happens upon exposure in the field and in the laboratory as a function of  $\text{H}_2\text{O}$  vapor pressure or by soaking in water, giving a readily observable change in extinction angle and cell dimensions (e.g., Coombs 1952, Armbruster & Kohler 1992).

Fersman (1908) introduced the term "primary leonhardite" for a variety from Kurtsy (nowadays Ukrainka), Crimea, with 14 molecules of  $\text{H}_2\text{O}$ , which neither dehydrates nor rehydrates under ambient conditions. In it, (K,Na) substitutes for Ca, although Ca is still dominant (Pipping 1966).

Type leonhardite of Blum from Schemnitz catalogued in the Museum of Natural History, Vienna, in 1843 and type "primary leonhardite" of Fersman obtained from the Fersman Mineralogical Museum in Moscow are shown by Wuest & Armbruster (1997) and Stolz & Armbruster (1997), respectively, to have the same Si,Al ordered framework of tetrahedra as laumontite. The low  $\text{H}_2\text{O}$  content of "primary leonhardite" is attributed to space limitations resulting from the introduction of additional cations of larger size.

In conformity with Rule 4, leonhardite is discredited as the name of a separate species. It is an  $\text{H}_2\text{O}$ -poor variety of laumontite. "Primary leonhardite" is  $\text{H}_2\text{O}$ -poor sodian potassian laumontite.

*Svetlozarite is dachiardite-Ca*

Svetlozarite was described by Maleev (1976) as a high-silica zeolite occurring as spherulites in chalcedony veinlets in brecciated andesites west of Zvesdel, eastern Rhodopes, Bulgaria. Analysis showed  $\text{Ca} > \text{Na} > \text{K}$ , and minor Fe and Mg. From X-ray powder-diffraction studies, Maleev suggested an orthorhombic symmetry, with a *c*-axis repeat of 7.5 Å, which is characteristic of the mordenite group, to which he attributed the mineral.

Gellens *et al.* (1982) concluded from powder and single-crystal X-ray and transmission electron microscopy (TEM) studies, that svetlozarite, space group *Ccma* (?), is related to the ideal dachiardite structure by irregular periodic twinning and stacking

faults, and that it is not a topologically distinct member of the mordenite family. Its composition is within the range of other samples of dachiardite. It is regarded as a multiply twinned and highly faulted dachiardite (dachiardite-Ca), and is discredited as a separate species.

*Wellsite is barian phillipsite-Ca and calcian harmotome*

The mineral named wellsite by Pratt & Foote (1897) has been shown by Galli (1972) and Galli & Loschi Ghittoni (1972) to be isostructural with phillipsite and harmotome, and Černý *et al.* (1977) have shown that zoning in crystals of wellsite covers most of the range from Ca-rich phillipsite to potassian calcian harmotome. Wellsite is discredited. Most examples of wellsite are barian phillipsite-Ca, and others are calcian harmotome.

## REFERENCES

- ARMBRUSTER, T. & KOHLER, T. (1992): Re- and dehydration of laumontite: a single-crystal X-ray study at 100 K. *Neues Jahrb. Mineral., Monatsh.*, 385-397.
- BLUM, J.R. (1843): Leonhardit, ein neues Mineral. (*Poggendorff's Annalen der Physik und Chemie* **59**, 336-339.
- ČERNÝ, P., RINALDI, R. & SURDAM, R.C. (1977): Wellsite and its status in the phillipsite-harmotome group. *Neues Jahrb. Mineral., Abh.* **128**, 312-320.
- COOMBS, D.S. (1952): Cell size, optical properties and chemical composition of laumontite and leonhardite. *Am. Mineral.* **37**, 812-830.
- DELFFS, W. (1843): Analyse des Leonhardits. (*Poggendorff's Annalen der Physik und Chemie* **59**, 339-342.
- DOELTER, C. (1921): *Handbuch der Mineralchemie* **II**, 3. Verlag Theodor Steinkopff, Dresden, Germany.
- FERSMAN, A.E. (1908): Materialien zur Untersuchung der Zeolithe Russlands. I. Leonhardit und Laumontit aus der Umgebung von Simferopol (Krim). *Trav. du Musée géol. Pierre le Grand pr. l'Acad. Imp. de Science St Pétersbourg* **2**, 103-150 (abstr. in *Z. Kristallogr.* **50**, 75-76).
- GALLI, E. (1972): La phillipsite barifera ("wellsite") di M. Calvarina (Verona). *Per. Mineral.* **41**, 23-33.
- \_\_\_\_\_ & LOSCHI GITTONI, A.G. (1972): The crystal chemistry of phillipsites. *Am. Mineral.* **57**, 1125-1145.
- GELLENS, R.L., PRICE, G.D. & SMITH, J.V. (1982): The structural relation between svetlozarite and dachiardite. *Mineral. Mag.* **45**, 157-161.

- HAUSMANN, J.F.L. (1847): *Handbuch der Mineralogie* (2nd ed.). 2, 1600.
- LÉVY, A. (1825): Descriptions of two new minerals. *Annals of Philosophy, new ser.* **10**, 361-363.
- MALEEV, M.N. (1976): Svetlozarite, a new high-silica zeolite. *Zap. Vses. Mineral. Obshchest.* **105**, 449-453 (in Russ.).
- MASON, B. (1962): Herschelite – a valid species? *Am. Mineral.* **47**, 985-987.
- PASSAGLIA, E. (1970): The crystal chemistry of chabazites. *Am. Mineral.* **55**, 1278-1301.
- PIPPING, F. (1966): The dehydration and chemical composition of laumontite. *Mineral. Soc. India, IMA Vol.*, 159-166.
- PRATT, J.H. & FOOTE, H.W. (1897): On wellsite, a new mineral. *Am. J. Sci.* **153**, 443-448.
- SHEPPARD, R.A., GUDE, A.J. & EDSON, G.M. (1978): Bowie zeolite deposit, Cochise and Graham Counties, Arizona. In *Natural Zeolites – Occurrence, Properties and Use* (L.B. Sand & F.A. Mumpton, eds.). Pergamon, Oxford, U.K. (319-328).
- STOLZ, J. & ARMBRUSTER, T. (1997): X-ray single-crystal structure refinement of a Na,K-rich laumontite, originally designated "primary leonhardite". *Neues Jahrb. Mineral., Monatsh.* 131-134.
- STRUNZ, H. (1956): Die Zeolithe Gmelinit, Chabasit, Levyn (Phakolith, Herschelit, Seebachit, Offretit). *Neues Jahrb. Mineral., Monatsh.*, 250-259.
- WUEST, T. & ARMBRUSTER, T. (1997): Type locality leonhardite: a single-crystal X-ray study at 100 K. *Zeolite '97, 5th Int. Conf. on the Occurrence, Properties, and Utilization of Natural Zeolites (Ischia, Italy), Program Abstr.*, 327-328.