



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Reconstruction of the mouse extrahepatic biliary tree using primary human extrahepatic cholangiocyte organoids

Citation for published version:

Sampaziotis, F, Justin, AW, Tysoe, OC, Sawiak, S, Godfrey, EM, Upponi, SS, Gieseck, RL, de Brito, MC, Berntsen, NL, Gómez-Vázquez, MJ, Ortmann, D, Yiangou, L, Ross, A, Bargehr, J, Bertero, A, Zonneveld, MCF, Pedersen, MT, Pawlowski, M, Valestrand, L, Madrigal, P, Georgakopoulos, N, Pirmadjid, N, Skeldon, GM, Casey, J, Shu, W, Materek, PM, Snijders, KE, Brown, SE, Rimland, CA, Simonic, I, Davies, SE, Jensen, KB, Zilbauer, M, Gelson, WTH, Alexander, GJ, Sinha, S, Hannan, NRF, Wynn, TA, Karlsen, TH, Melum, E, Markaki, AE, Saeb-Parsy, K & Vallier, L 2017, 'Reconstruction of the mouse extrahepatic biliary tree using primary human extrahepatic cholangiocyte organoids', *Nature Medicine*, vol. 23, no. 8, pp. 954-963. <https://doi.org/10.1038/nm.4360>

Digital Object Identifier (DOI):

[10.1038/nm.4360](https://doi.org/10.1038/nm.4360)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Nature Medicine

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



1 **Extrahepatic cholangiocyte organoids for cell-based therapy**
2 **applications**

3

4 Fotios Sampaziotis^{1,2,3}, Alexander W Justin⁴, Olivia C. Tysoe^{1,2}, Steve
5 Sawiak^{5†}, Edmund M Godfrey^{6†}, Sara S Upponi^{6†}, Richard L. Gieseck III⁷,
6 Miguel Cardoso de Brito¹, Natalie Lie Berntsen⁸, María J Gómez-Vázquez⁹,
7 Daniel Ortmann¹, Loukia Yiangou¹, **Alexander Ross^{1,10,11}**, Johannes
8 Bargehr^{1,12}, Alessandro Bertero¹, Mariëlle CF Zonneveld¹, Marianne T
9 Pedersen¹³, Matthias Pawlowski¹, **Laura Valestrand^{7,14,15}**, **Pedro Madrigal¹**,
10 Nikitas Georgakopoulos², Negar Pirmadjid², Gregor M Skeldon^{16,17}, **John**
11 **Casey¹⁸**, Wenmiao Shu^{16,17}, Paulina M Materek¹, Kirsten E Snijders¹,
12 Stephanie E Brown¹, Casey A Rimland^{1,2,7,19}, Ingrid Simonic²⁰, Susan E
13 Davies²¹, Kim B Jensen¹³, **Matthias Zilbauer^{10,11,22}**, William TH Gelson³,
14 Graeme Alexander²³, Sanjay Sinha^{1,12}, Nicholas RF Hannan²⁴, Thomas
15 Wynn⁷, Tom H Karlsen⁸, Espen Melum⁸, Athina E. Markaki⁴, Kourosh Saeb-
16 Parsy^{2*}, Ludovic Vallier^{1,2,25*}

17

18 ¹Wellcome Trust-Medical Research Council Stem Cell Institute, Cambridge
19 Stem Cell Institute, Anne McLaren Laboratory, Department of Surgery,
20 University of Cambridge, Cambridge, UK

21 ²Department of Surgery, University of Cambridge and NIHR Cambridge
22 Biomedical Research Centre, Cambridge, UK

23 ³Department of Hepatology, Cambridge University Hospitals NHS Foundation
24 Trust, Cambridge, UK

1 ⁴Department of Engineering, University of Cambridge, Cambridge, UK

2 ⁵Department of Clinical Neurosciences, University of Cambridge, Cambridge,
3 UK

4 ⁶Department of Radiology, Cambridge University Hospitals NHS Foundation
5 Trust, Cambridge, UK

6 ⁷Immunopathogenesis Section, Laboratory of Parasitic Diseases, National
7 Institute of Allergy and Infectious Diseases, National Institutes of Health,
8 Bethesda, Maryland, USA

9 ⁸Norwegian PSC Research Center, Division of Surgery, Inflammatory
10 Medicine and Transplantation, Oslo University Hospital, Rikshospitalet, Oslo.

11 ⁹Cambridge Genomic Services, Department of Pathology, University of
12 Cambridge, Cambridge, UK

21 ¹⁰University Department of Paediatrics, University of Cambridge, Cambridge,
22 UK

23 ¹¹Department of Paediatric Gastroenterology, Hepatology and Nutrition,
24 Cambridge University Hospitals NHS Foundation Trust, Cambridge, UK

25 ¹²Division of Cardiovascular Medicine, University of Cambridge, Cambridge,
26 UK

27 ¹³Biotech Research and Innovation Centre (BRIC), University of Copenhagen,
28 Copenhagen, Denmark

29 ¹⁴Research Institute of Internal Medicine, Division of Surgery, Inflammatory
30 Medicine and Transplantation, Oslo University Hospital, Rikshospitalet, Oslo,

31 ¹⁵Institute of Clinical Medicine, University of Oslo, Oslo.

32 ¹⁶School of Engineering and Physical Sciences, Heriot-Watt University, United
33 Kingdom

1 ¹⁷Department of Biomedical Engineering, University of Strathclyde, United
2 Kingdom

3 ¹⁸Department of Surgery, University of Edinburgh, Edinburgh Royal Infirmary,
4 Edinburgh, EH16 4SU, United Kingdom

5 ¹⁹University of North Carolina, Chapel Hill School of Medicine, Chapel Hill,
6 North Carolina, USA

7 ²⁰Medical Genetics Laboratories, Cambridge University Hospitals NHS Trust,
8 Cambridge, UK

9 ²¹Department of Histopathology, Cambridge University Hospitals NHS
10 Foundation Trust, Cambridge, UK

11 ²²Wellcome Trust-Medical Research Council Stem Cell Institute, Cambridge
12 Stem Cell Institute, University of Cambridge, Cambridge, UK

13 ²³Department of Medicine, School of Clinical Medicine, University of
14 Cambridge, Cambridge, UK

15 ²⁴Center for Biomolecular Sciences, University of Nottingham, UK

16 ²⁵Wellcome Trust Sanger Institute, Hinxton, UK

17

18 **Authorship note:** * Kourosh Saeb-Parsy and * Ludovic Vallier contributed
19 equally to this work.

20 [†]Stephen Sawiak, [†]Edmund Godfrey and [†]Sara Upponi contributed equally to
21 this work.

22 **Correspondence:** Ludovic Vallier, Laboratory for Regenerative Medicine,
23 West Forvie Building, Robinson Way, University of Cambridge. Cambridge
24 CB2 0SZ, United Kingdom. Telephone: 44.1223.747489; Fax:
25 44.1223.763.350; E-mail: lv225@cam.ac.uk.

1 Disorders of the extrahepatic bile ducts carry significant morbidity and
2 mortality. Indeed, 70% of pediatric liver transplantations are performed to treat
3 biliary atresia (1), Primary Sclerosing Cholangitis (PSC) alone accounts for
4 5% of US liver transplantations (2) and biliary complications are the leading
5 cause of graft failure following deceased liver transplantation (3,4). Treatment
6 options remain limited (5,6) due to the lack of healthy tissue that can be used
7 to reconstruct and replace diseased bile ducts. In vitro expansion of native
8 cholangiocytes could address this challenge and provide cells suitable for
9 tissue engineering applications such as biliary reconstruction. However, the
10 culture of primary biliary epithelium remains problematic (7). Here we report a
11 novel method for the isolation and propagation of primary human
12 cholangiocytes from the extrahepatic biliary tree, compatible with regenerative
13 medicine applications. The resulting Extrahepatic Cholangiocyte Organoids
14 (ECOs) express key biliary markers such as CK7, CK19, GGT, CFTR and
15 maintain their functional properties in vitro including ALP, GGT activity and
16 responses to secretin and somatostatin. The potential of ECOs for tissue
17 engineering and clinical applications is further illustrated by their capacity to
18 populate biodegradable scaffolds, organize into a functional biliary epithelium
19 and rescue a murine model of extrahepatic biliary injury (EHBI).

20 To establish this system, we first focused on identifying optimal conditions to
21 isolate primary cholangiocytes from the biliary epithelium which forms a
22 monolayer covering the luminal surface of the biliary tree (8). We tested
23 several approaches for recovering these cells and mechanical dissociation by
24 brushing or scraping the bile duct lumen was associated with improved
25 survival compared to enzymatic digestion (Figure 1a, Supplementary Fig. 1a).

1 Furthermore, the majority of the resulting cells co-expressed the biliary
2 markers Cytokeratin 7 (CK7) and Cytokeratin 19 (CK19) ($94.6 \pm 2.4\%$ (SD;
3 $n=3$); while no contamination from mesenchymal cell types was detected.
4 (Supplementary Fig 2). Consequently, mechanical dissociation constitutes the
5 optimal method for harvesting extrahepatic cholangiocytes.

6 To discern appropriate conditions for the maintenance and propagation of
7 these cells, we optimized our recently established system for 3D culture of
8 human induced pluripotent stem cell-derived intrahepatic cholangiocytes (9).
9 Screening of multiple growth factors known to support expansion of
10 cholangiocytes and epithelial organoids (10,11) (Supplementary Fig 1b-1c)
11 identified that the combination of Epidermal Growth Factor (EGF), R-Spondin
12 and Dickkopf-related protein 1 (DKK-1) promoted the growth of primary
13 cholangiocytes into organoids (Figure 1b, 1c). Due to the paradoxical
14 requirement for both a Wnt potentiator (R-spondin) and an inhibitor (DKK-1),
15 we characterized the canonical and non-canonical/PCP Wnt pathway activity
16 in ECOs. Our results demonstrate increased levels of β -catenin
17 phosphorylation (Supplementary Fig. 1d-1e), signifying reduced WNT
18 canonical pathway activity in ECOs, as well as increased Rho Kinase activity
19 (Supplementary Fig. 1f), **which could be** consistent with enhanced non-
20 canonical/PCP signaling in ECOs. Thus, **it is possible that** non-canonical Wnt
21 signaling controls ECO expansion marking a significant difference with
22 previous organoid culture conditions (11).

23 Under these conditions, we derived 8 different ECO lines (Supplementary
24 Table 1) from a variety of deceased donors aged from 33 to 77 years.
25 Importantly, we obtained similar results by using cholangiocytes isolated from

1 the gallbladder or by harvesting common bile duct cholangiocytes using an
2 Endoscopic Retrograde Cholangio-Pancreatography (ERCP) brush instead of
3 scrapping the lumen (Supplementary Fig 3). Consequently, ECOs can be
4 derived from different areas of the extra-hepatic biliary tree and harvested
5 using peri-operative (dissection and scrapping) or minimally invasive (ERCP
6 brushings) approaches.

7 The resulting cells were expanded in vitro for prolonged periods of time
8 (Supplementary Fig 4a) while maintaining their genetic stability
9 (Supplementary Fig 4b-4c). Electron microscopy revealed the presence of
10 characteristic ultrastructural features including cilia, microvilli and tight
11 junctions (12) (Figure 1d, Supplementary Fig 5), while QPCR and
12 immunofluorescence (IF) analyses established the expression of key biliary
13 markers such as CK7, CK19, Hepatocyte Nuclear Factor-1-beta (HNF1B),
14 Gamma Glutamyl-Transferase (*GGT*), Secretin Receptor (SCR), sodium-
15 dependent bile acid transporter (ASBT/SLC10A2), Cystic fibrosis
16 transmembrane conductance regulator (CFTR) and Sox9 (9) (Figure 1e-1f,
17 Supplementary Fig 3c-3d, 6a-6b). Importantly, stem cell markers (POU5F1,
18 NANOG, PROM1, *LGR-4/5/6*), markers of non-biliary lineages (albumin, α 1-
19 antitrypsin, CK18, PDX1, insulin and glucagon) and EMT markers (vimentin,
20 SNA1L and S100A4) were not detected (Supplementary Fig 7a-7c). On the
21 other hand, 98.1% \pm 0.9% (SD; n=3) of the cells co-expressed CK7 and CK19
22 following 20 passages (Supplementary Fig 2) thereby confirming the presence
23 of a near homogeneous population of cholangiocytes.

24 Transcriptomic analyses (Figure 1g, Supplementary Fig. 8, Supplementary
25 Table 2) revealed that ECOs maintain a stable gene expression profile over

1 multiple passages (Pearson correlation coefficient for Passage 1 (P1) vs.
2 Passage 20 (P20) $r=0.99$, Supplementary Fig 8a-b), express key biliary
3 markers and cluster closely to freshly isolated cholangiocytes (Pearson
4 correlation coefficient for Primary Cholangiocytes (PCs) vs. Passage 20 (P20)
5 $r=0.92$; Figure 1g, Supplementary Fig 8b-8d). Gene ontology analyses
6 confirmed enrichment of pathways characteristic for the biliary epithelium
7 (Supplementary Fig 8e). Considered collectively, these results demonstrate
8 that primary cholangiocytes derived from the extrahepatic biliary tree can be
9 expanded in vitro without losing their original characteristics.

10 We then decided to further characterize ECOs by focusing on their function
11 following long term culture (20 passages). The biliary epithelium regulates the
12 homeostasis of bile through the transport of ions, water and bile acids (8,13).
13 The secretory capacity of ECOs was interrogated using Rhodamine-123, a
14 fluorescent substrate for the cholangiocyte surface glycoprotein Multidrug
15 Resistance protein-1 (MDR1) (14,15) (Figure 2a-2c). Rhodamine-123
16 accumulated in the ECO lumen only in the absence of the MDR-1 antagonist
17 verapamil, thereby confirming active secretion through MDR-1 (Figure 2a-2c).
18 Luminal extrusion of bile acids (16) was also demonstrated by showing that
19 the fluorescent bile acid Cholyl-Lysyl-Fluorescein (CLF) was actively exported
20 from ECOs (Figure 2d-2f). Furthermore, ECO ALP and GGT activity was
21 comparable to freshly plated primary cholangiocytes (Figure 2g-2h,
22 Supplementary Fig. 3e-3f). The response of ECOs to secretin and
23 somatostatin was also assessed. Secretin promotes water secretion,
24 distending the bile duct lumen, while somatostatin negates the effects of
25 secretin (17–19). Accordingly, organoids exposed to secretin increased their

1 diameter compared to untreated controls, while somatostatin inhibited the
2 effect of secretin (Figure 2i-2j). Our data, therefore, demonstrate that ECOs
3 maintain their functional properties after long term culture.

4 These results prompted us to investigate the interest of ECOs for in vivo use,
5 especially regenerative medicine applications. We first characterized the
6 potential of ECOs for in vivo engraftment and survival by transplanting cells
7 under the kidney capsule of NOD.Cg-Prkdc^{scid} Il2rg^{tm1Wjl} (NSG) mice
8 (Supplementary Fig 9a) for 12 weeks (20). ECOs successfully engrafted
9 forming tubular structures expressing biliary markers such as CK19
10 (Supplementary Fig 9b-d). Importantly, no tumour formation or markers of
11 differentiation to other lineages were detected (Supplementary Fig 9d). Thus,
12 ECOs appear to maintain their basic characteristics even after prolonged
13 engraftment in vivo **under the kidney capsule**.

14 Following these encouraging results, we decided to define the capacity of
15 ECOs to repair the biliary epithelium. For that, we developed a mouse model
16 of extrahepatic biliary injury (EHBI). More specifically, to simulate biliary tree
17 wall defects requiring biliary reconstruction (21), the biliary tree of healthy
18 NSG mice was compromised through a longitudinal incision in the gallbladder
19 wall (Figure 3a). We then generated a bioengineered tissue patch to repair
20 this injury by populating Polyglycolic Acid (PGA) biodegradable scaffolds with
21 either GFP-expressing ECOs (Supplementary Fig 10-12, Supplementary
22 Note) or GFP-expressing fibroblasts (Supplementary Fig 13a-13d). The
23 resulting bioengineered tissue was subsequently transplanted into the injured
24 animals to close the wall defect created by surgery. Animals receiving
25 acellular scaffolds died within 24 hours of the operation (Figure 3b) and post-

1 mortem examination revealed yellow pigmentation of the peritoneal cavity and
2 seminal vesicles consistent with bile leak (Supplementary Fig 14a); while all
3 animals in fibroblast-scaffold group failed to reconstruct their gallbladder
4 which was replaced by fibrotic tissue incompatible with bile transport or
5 storage (Supplementary Fig 13e-13g). In contrast, animals transplanted with
6 scaffolds containing ECOs survived for up to 104 days without complications
7 and were culled electively (Figure 3b). Importantly, the reconstructed
8 gallbladders in the ECO group were fully remodeled resembling the
9 morphology of their native counterparts (Figure 3c, Supplementary Fig 14b).
10 Histology (Figure 3d), IF and QPCR analyses of the ECO-reconstructed
11 gallbladders (Figure 3e, Supplementary Fig 14c-14d) unveiled integration of
12 GFP-positive ECOs expressing biliary markers, such as CK19, CK7, HNF1B,
13 Sox9, CFTR and a human-specific epitope for Ku80 (Figure 3e,
14 Supplementary Fig 14c). Importantly, these IF analyses also showed the
15 presence of mouse mesenchymal cells expressing vimentin and endothelial
16 cells expressing CD31 in the reconstructed biliary epithelium suggesting that
17 the scaffold is colonized by endogenous cells after transplantation
18 (Supplementary Fig 14c). Interestingly, we also identified a population of
19 GFP+/vimentin+/CK19- cells, suggesting that ECOs may also contribute to
20 the scaffold stroma; possibly through epithelial to mesenchymal transition
21 (EMT; Supplementary Fig 14c, 14e). The integrity of the reconstructed
22 gallbladder lumen and its exposure to bile through continuity with the biliary
23 tree were demonstrated using Magnetic Resonance Cholangio-
24 Pancreatography (MRCP) imaging prior to removal of the organ and was
25 further confirmed with FITC cholangiograms (Figure 3f-3g, Supplementary Fig

1 14f, Supplementary Video 1). Post mortem surgical examination and full body
2 Magnetic resonance Imaging 104 days post transplantation revealed no
3 evidence of tumor formation (Supplementary Fig 14f, Supplementary Video 2)
4 while IF analyses revealed no GFP+ cells in the adjacent liver tissue (data not
5 shown). On the contrary, gallbladders reconstituted with fibroblasts controls
6 exhibited obliteration of the gallbladder lumen (Supplementary Fig 13h-13i)
7 and replacement of the lumen and biliary epithelium by fibroblasts expressing
8 Fibroblast Specific Antigen S100A4 (Supplementary Fig. 13i-13j). Considered
9 collectively, our findings demonstrate the capacity of ECOs to colonize their
10 physiological niche and regenerate part of the biliary tree without any
11 complications.

12 Reconstruction of the gallbladder wall provided proof-of-principle for the
13 capacity of ECOs to regenerate the biliary epithelium after injury; however, the
14 majority of extrahepatic bile duct disorders affect the common bile duct (CBD).
15 Therefore we decided to explore the possibility to replace the native CBD of
16 NSG mice with a bioengineered duct consisting of an ECO-populated
17 densified collagen tube. (Supplementary results, Supplementary Fig 15, 16,
18 Fig 4a). A mid-portion of the native CBD was removed and an ECO-populated
19 collagen tube was anastomosed end-to-end to the proximal and distant duct
20 remnants (n=4 animals). Fibroblast populated tubes were used as a negative
21 control (n=4). Biliary reconstruction was achieved in all animals transplanted
22 with ECO-populated tubes (Figure 4b-4c, Supplementary Fig 17a-17d), which
23 were followed up for up to a month post transplantation (Supplementary Fig
24 17d). Histology and IF analyses revealed a patent lumen, with formation of a
25 biliary epithelium by the transplanted GFP+ cells (Figure 4e-4f,

1 **Supplementary Fig 17a-17b**). IF and QPCR analyses confirmed the
2 expression of biliary markers, such as CK19, CK7, HNF1B, CFTR, Sox9
3 (Figure 4d, 4f, **Supplementary Fig 17b**) by the engrafted cells but also
4 illustrated the presence of mouse stromal and endothelial cells
5 (**Supplementary Fig 17b**). **Moreover, we observed minimal apoptosis and**
6 **proliferation in the transplanted tubes 1 month after transplantation,**
7 **confirming the stability and integrity of the reconstituted biliary epithelium**
8 **(Supplementary Fig 17b-17c)**. Lumen patency was further confirmed by FITC-
9 cholangiogram **and MRCP** (Figure 4g, **Supplementary Fig 17f**) **and all the**
10 **animals receiving ECO-populated tubes exhibited no increase in serum**
11 **cholestasis markers (Bilirubin, ALP; Supplementary Fig 17e) accordingly;**
12 while the bio-artificial common bile ducts retained their ALP activity in vivo
13 (Figure 4h). On the contrary, all the fibroblast-populated collagen tubes failed
14 due to lumen occlusion (Figure 4b-4c, 4e-4g, **Supplementary Fig 17d**),
15 **resulting in increased biliary pressures and bile leak through the site of**
16 **anastomosis (Figure 4b)**. In conclusion, our results demonstrate the capacity
17 of ECO-populated collagen tubes to replace the native CBD in vivo.

18 In summary, we have demonstrated that epithelial cells from the extrahepatic
19 biliary tree can be expanded and propagated in vitro while maintaining their
20 cholangiocyte transcriptional signature and functional characteristics. In
21 addition, our results show that primary cholangiocytes expanded in vitro as
22 organoids have a unique potential for organ regeneration. Indeed, our system
23 provides the first proof-of-principle for the application of regenerative medicine
24 in the context of common bile duct pathology. The capacity to replace a
25 diseased common bile duct with an in vitro bio-engineered ECO-tube could

1 have a significant impact for the management of disorders such as biliary
2 atresia, which constitutes the leading cause for pediatric liver transplantation
3 (1); or ischemic strictures which are one of the most common complications
4 following transplantation (3). Consequently ECO-populated scaffolds
5 constitute a novel system with increased clinical relevance in the field of
6 cholangiopathies.

7 Furthermore, studies of the extrahepatic biliary epithelium have been limited
8 by technical challenges in long-term culture and significant expansion of
9 primary cholangiocytes. These challenges have so far precluded large scale
10 experiments such as transcriptomic and genome-wide analyses which are
11 urgently needed to better understand bile duct diseases, such as PSC and
12 cholangiocarcinoma. The capacity of ECOs for large scale expansion, could
13 address this challenge. Indeed, we demonstrate that starting from 10^5
14 extrahepatic cholangiocytes we can generate between 10^{20} – 10^{25} cells after
15 20 passages. Therefore, ECOs not only represent a novel source of cells for
16 cell based therapy but also provide a unique model system for studying the
17 physiology and modeling disorders of the extrahepatic biliary tree in vitro.

18 Access to human tissue constitutes a significant limitation for systems based
19 on primary cells. However, we show that ECOs can be obtained not only from
20 the common bile duct but also from the gallbladder. Gallbladder tissue is
21 easily accessible and routinely discarded following liver transplantation and
22 cholecystectomy, one of the most common surgical procedures performed.
23 Furthermore, in patients not having surgery the common bile duct can be
24 accessed using minimally invasive procedures, such as Endoscopic
25 Retrograde Cholangio-Pancreatography (ERCP) and we demonstrate that

1 cholangiocytes can be obtained through brushings, which are routinely
2 performed to acquire histology specimens. Importantly, no morphological or
3 functional differences were observed between organoids obtained with these
4 different methods. Moreover, due to the scalability of our system only a small
5 amount of starting material is required. **Finally, recent progress in replacing**
6 **Matrigel by custom made hydrogels to grow gut organoids (22) suggest that**
7 **translating our system from Matrigel to Good Manufacturing Practice (GMP)**
8 **could be feasible.** Considered together, these approaches address any issues
9 of tissue availability and open the possibility of autologous as well as
10 allogeneic cell based therapy.

11 Importantly, the derivation of primary hepatic stem cells using an organoid
12 culture system has been reported previously (11). However, the capacity of
13 the resulting cells to differentiate into functional cholangiocytes and populate
14 the biliary tree in vivo remains to be demonstrated. Furthermore, in vivo
15 applications of such platforms could be restricted by contaminating stem cells
16 with a capacity to proliferate inappropriately after transplantation and /or
17 differentiate into non-biliary cell types. Importantly, despite the association
18 between organoids and adult stem cells (23), we never observed the
19 expression of hepatocyte or pancreatic markers during our experiments either
20 in vitro or after transplantation, suggesting that the differentiation capacity of
21 ECOs is limited to their lineage of origin. Moreover, canonical WNT signaling,
22 which is crucial for the expansion of adult stem cell organoids (24) is blocked
23 in our culture conditions through the use of DKK-1 **and further studies may be**
24 **required to fully elucidate the role of R-spondin in our system.** Considered
25 together, these observations suggest that our culture system does not include

1 a stem cell population. However, we cannot completely exclude that these
2 cells could represent a biliary progenitor population based on their ability to
3 self-propagate and generate organoids from single cells (Supplementary
4 Video 3).

5 In conclusion, our results open up novel avenues for the use of extrahepatic
6 primary biliary tissue as a novel platform for in vitro studies, disease modeling
7 and cell based therapy applications.

8

1 **Online Methods**

2 **Primary biliary tissue**

3 Primary biliary tissue (bile duct or gallbladder) was obtained from deceased
4 organ donors from whom organs were being retrieved for transplantation. The
5 gallbladder or a section of the bile duct was excised during the organ retrieval
6 operation after obtaining informed consent from the donor's family (REC
7 reference numbers: 09/H0306/73, NRES Committee East of England –
8 Norfolk, 12/EE/0253, NRES Committee East of England - Cambridge Central
9 and 15/EE/0152 NRES Committee East of England - Cambridge South).

10 **Isolation of primary cholangiocytes**

11 Excised bile duct segments were placed in a 10cm plate and washed once
12 with William's E medium (Gibco, Life Technologies). A longitudinal incision
13 was made along the wall of the excised bile duct segment exposing the lumen
14 and 10-15ml of William's E medium were added to cover the tissue. The
15 luminal epithelium was subsequently scraped off using a surgical blade, while
16 submerged in medium. The supernatant was collected and the tissue and
17 plate were washed 2-3 times with William's E medium to harvest any
18 remaining cells. The supernatant and washes were centrifuged at 444g for 4
19 minutes. The pellet was washed with William's E, re-centrifuged and the
20 supernatant was discarded (Figure 1a).

21 Excised gallbladders were placed in a 15cm plate, a longitudinal incision was
22 made along the wall of the excised gallbladder and the lumen was washed
23 once with William's E medium (Gibco, Life Technologies). Cholangiocytes

1 were isolated and harvested following the method described above
2 (Supplementary Fig. 3a).

3 For isolation through brushings, an excised bile duct segment was placed in a
4 10cm plate and cannulated using an ERCP brush. The lumen was brushed
5 10-20 times and the cells were harvested by washing the brush several times
6 in a falcon tube containing 40-50 ml of William's E medium (Supplementary
7 Fig. 3b).

8 **Generation and culture of ECOs**

9 Isolated primary cholangiocytes were centrifuged at 444g for 4 minutes and
10 re-suspended in a mixture of 66% matrigel (BD Biosciences, catalogue
11 number: 356237) and 33% William's E medium (Gibco, Life Technologies)
12 supplemented with 10mM nicotinamide (Sigma-Aldrich), 17mM sodium
13 bicarbonate (Sigma Aldrich), 0.2mM 2-Phospho-L-ascorbic acid trisodium salt
14 (Sigma-Aldrich), 6.3mM sodium pyruvate (Invitrogen), 14mM glucose (Sigma-
15 Aldrich), 20mM HEPES (Invitrogen), ITS+ premix (BD Biosciences), 0.1µM
16 dexamethasone (R&D Systems), 2mM Glutamax (Invitrogen), 100U/ml
17 penicillin per 100µg/ml streptomycin, 20ng/ml EGF (R&D Systems), 500ng/ml
18 R-Spondin (R&D Systems) and 100ng/ml DKK-1 (R&D Systems). The cell
19 suspension was plated in 24-well plate format, at 50µl/well, so that a small
20 dome of matrigel was formed in the centre of each well and then incubated at
21 37°C for 10-30 minutes until it solidified. Subsequently, 1ml of William's E
22 medium with supplements was added. The culture medium was changed
23 every 48 hours.

1 To split the cells, the matrigel was digested by adding Cell Recovery Solution
2 (Corning) for 30 minutes at 4°C. The resulting cell suspension was harvested,
3 centrifuged at 444g for 4 minutes, washed once with William's E medium and
4 re-suspended in 66% matrigel and 33% William's E medium with
5 supplements, as described above.

6 All experiments were performed using passage 20 ECOs unless otherwise
7 stated.

8 **Cell line identity**

9 Demographic data for donor corresponding to the each ECO lines is provided
10 in supplementary table 1. Following derivation ECO lines were authenticated
11 by matching their karyotype (Supplementary Fig. 2c) to the sex of the donor of
12 origin. The lines were tested on a regular basis and found to be negative for
13 mycoplasma contamination.

14 **Immunofluorescence, RNA extraction and Quantitative Real Time PCR**

15 IF, RNA extraction and QPCR were performed as previously described (9). A
16 complete list of the primary and secondary antibodies used is provided in
17 supplementary table 3. A complete list of the primers used is provided in
18 supplementary table 4.

19 All QPCR data are presented as the median, interquartile range (IQR) and
20 range (minimum to maximum) of four independent biological replicates.
21 Values are relative to the housekeeping gene Hydroxymethylbilane Synthase
22 (*HMBS*).

23 All IF images were acquired using a Zeiss Axiovert 200M inverted microscope
24 or a Zeiss LSM 700 confocal microscope. Imagej 1.48k software (Wayne

1 Rasband, NIHR, USA, <http://imagej.nih.gov/ij>) was used for image
2 processing. IF images are representative of at least 3 different experiments.
3 IF images of reconstructed gallbladder sections are representative of 5
4 different animals.

5 **Microarrays**

66 RNA for microarray analysis was collected from 3 different ECO lines (n=3).
67 The RNA was assessed for concentration and quality using a SpectroStar
68 (BMG Labtech, Aylesbury, UK) and a Bioanalyser (Agilent Technologies,
69 Cheadle, UK). Microarray experiments were performed at Cambridge
70 Genomic Services, University of Cambridge, using the HumanHT-12 v4
71 Expression BeadChip (Illumina, Chesterford, UK) according to the
72 manufacturer's instructions. Briefly, 200ng of Total RNA underwent linear
73 amplification using the Illumina TotalPrep RNA Amplification Kit (Life
74 Technologies, Paisley, UK) following the manufacturer's instructions. The
75 concentration, purity and integrity of the resulting cRNA were measured by
76 SpectroStar and Bioanalyser. Finally cRNA was hybridised to the HumanHT-
77 12 v4 BeadChip overnight followed by washing, staining and scanning using
78 the Bead Array Reader (Illumina). The microarray data are available on
79 ArrayExpress (Accession number: E-MTAB-4591). For reviewer access,
80 please use the following login details Username: Reviewer_E-MTAB-4591
81 Password: rtlImbi0

82 **Microarrays analysis**

83 Raw data was loaded into R using the lumi package from bioconductor (25)
84 and divided into subsets according to the groups being compared; only the
85 samples involved in a given comparison are used. Subsets were then filtered

1 to remove any non-expressed probes using the detection p-value from
2 Illumina. Across all samples probes for which the intensity values were not
3 statistically significantly different ($p > 0.01$) from the negative controls were
4 removed from the analysis. Following filtering the data was transformed using
5 the Variance Stabilization Transformation (26) from lumi and then normalised
6 to remove technical variation between arrays using quantile normalisation.
7 Comparisons were performed using the limma package (27) with results
8 corrected for multiple testing using False Discovery Rate (FDR) correction.
9 Finally the quality of the data was assessed along with the correlations
10 between samples within groups.

11 Probes differentially expressed between HEP and ECOs representing the
12 aggregate transcriptional “signature” of ECOs were selected for Euclidean
13 hierarchical clustering using Perseus software (MaxQuant). Standard scores
14 (z-scores) of the log₂ normalized probe expression values across the different
15 conditions were calculated and used for this analysis.

16 **Western Analysis**

17 Total protein was extracted with lysis buffer (50mM Tris pH 8, 150mM NaCl,
18 0.1% SDS, 0.5% sodium deoxycholate, 1% Triton X-100 and protease and
19 phosphatase inhibitors). Protein concentrations were determined by BCA
20 Protein Assay Kit (Thermo Fisher Scientific) according to the manufacturer’s
21 instructions. Samples were prepared for Western blot by adding 1x NuPAGE
22 LDS Sample Buffer with 1% β -mercaptoethanol and incubated for 5 minutes
23 at 95°C. Protein (25 μ g) was separated by 4-12% NuPAGE Bis-Tris protein
24 gels (Invitrogen) and transferred onto PVDF membranes (Bio-Rad). Proteins
25 were detected by probing with antibodies specific to Phospho- β -catenin

1 (Ser33/37/Thr41) (Cell Signalling Technology), Active- β -catenin (Millipore),
2 Total- β -catenin (R&D), α -tubulin (Sigma) followed by incubation with
3 horseradish peroxidase anti-mouse, anti-goat or anti-rabbit secondary
4 antibodies. Membranes were developed using Pierce ECL Western blotting
5 substrate (Thermo Scientific) according to the manufacturer's instructions.

6 **Rho Kinase activity analyses**

7 Rho Kinase activity was measured using a commercially available kit (Cell
8 Biolabs, STA-416) according to the manufacturer's instructions

9 **Flow cytometry analyses**

10 ECO organoids were harvested using Cell Recovery Solution (Corning) for 30
11 minutes at 4°C, centrifuged at 444g for 4 minutes and dissociated to single
12 cells using TrypLE™ Express (Gibco). The cells were subsequently fixed
13 using 4% PFA for 20 minutes at 4°C. Cell staining and flow cytometry
14 analyses were performed as previously described (9,28).

15 **Karyotyping**

16 ECO organoids were harvested using Cell Recovery Solution (Corning),
17 dissociated to single cells as described above, plated in gelatin coated plates
18 and cultured using William's E medium with supplements. When the cells
19 were sub-confluent, usually after 72hrs, the cultures were incubated for 3-4
20 hours with William's E medium with supplements containing 0.1 μ g/ml
21 colcemid (Karyomax®, Gibco). The cells were then harvested using Trypsin-
22 EDTA (0.05%) (Gibco) for 4-5 minutes at 37°C, centrifuged at 344g for 5
23 minutes and re-suspended in 5mls of KCl hypotonic solution (0.055M). The
24 suspension was re-centrifuged at 344g for 5 minutes, 2 mls of a 3:1 100%

1 methanol:glacial acetic acid solution were added and slides were prepared as
2 previously described (29)

3 **Comparative Genomic Hybridization analyses**

4 Genomic DNA was labeled using the BioPrime DNA Labeling Kit (Invitrogen),
5 according to the manufacturer's instructions and samples were hybridised to
6 Agilent Sureprint G3 unrestricted CGH ISCA 8x60K human genome arrays
7 following the manufacturer's protocol, as previously described (30). The data
8 was analysed using the Agilent CytoGenomics Software.

9 **Rhodamine123 transport assay**

10 The Rhodamine 123 transport assay was performed as previously described
11 (9) and images were acquired using a Zeiss LSM 700 confocal microscope.
12 Fluorescence intensity was measured between the organoid interior and
13 exterior and luminal fluorescence was normalized over the background of the
14 extraluminal space. Each experiment was repeated in triplicate. Error bars
15 represent SD. Mean fluorescence intensity comparisons were performed
16 using a two sided student's t-test.

17 **Cholyl-Lysyl-Fluorescein transport assay**

18 To achieve loading with Cholyl-Lysyl-Fluorescein (CLF, Corning
19 Incorporated), ECO organoids were split in 5 μ M of CLF and incubated at 37°C
20 for 30 minutes. Images were acquired using a Zeiss LSM 700 confocal
21 microscope and fluorescence intensity was measured between the organoid
22 interior and exterior as described for the Rhodamine 123 transport assay. To
23 demonstrate that the changes in CLF fluorescence intensity observed were
24 secondary to active export of CLF from the organoid lumen, the experiment
25 was repeated with 5 μ M of unconjugated Fluorescein Isothiocyanate (FITC)

1 (Sigma-Aldrich) as a control. Fluorescence intensity measurements were
2 performed as described for the Rhodamine 123 transport assay. Each
3 experiment was repeated in triplicate. Error bars represent SD. Mean
4 fluorescence intensity comparisons were performed using a two sided
5 student's t-test.

6 **GGT activity**

7 GGT activity was measured in triplicate using the MaxDiscovery™ gamma-
8 Glutamyl Transferase (GGT) Enzymatic Assay Kit (Bioo scientific) based on
9 the manufacturer's instructions. Error bars represent SD. Mean absorbance
10 was compared using a two sided student's t-test.

11 **Alkaline Phosphatase staining**

12 Alkaline phosphatase was carried out using the BCIP/NBT Color
13 Development Substrate (5-bromo-4-chloro-3-indolyl-phosphate/nitro blue
14 tetrazolium) (Promega) according to the manufacturer's instructions.

15 **Response to Secretin and Somatostatin**

16 Responses to secretin and somatostatin were assessed as previously
17 described (9).

18 **Generation of ECOs expressing Green Fluorescent Protein**

19 EGFP expressing VSV-G pseudotyped, recombinant HIV-1 lentiviral particles
20 were produced with an optimized second generation packaging system by
21 transient co-transfection of three plasmids into HEK 293T cells (ATCC CRL-
22 11268). EGFP expression is under control of a core EF1 α -promoter. All
23 plasmids were a gift from Didier Trono and obtained from addgene (pWPT-
24 GFP #12255, psPAX2 #12260, pMD2.G, #12259). Viral infection of organoids
25 was performed as previously described (31). Infected ECOs were expanded

1 for 2 passages, harvested as described above for flow cytometry analyses
2 and cell sorting by flow cytometry for GFP positive cells was performed. GFP
3 expressing single cells were plated using our standard plating method and
4 cultured in William's E medium with supplements for 1-2 weeks until fully
5 grown ECO organoids developed.

6 **Generation of ECO populated PGA scaffolds**

7 1mm thick PolyGlycolic Acid (PGA) scaffolds with a density of 50mg/cc were
8 used for all experiments. Prior to seeding cells, the PGA scaffolds were pre-
9 treated with a 1M NaOH for 10-30 seconds washed 3 times, decontaminated
10 in a 70% ethanol solution for 30 minutes and then air-dried for another 30
11 minutes until all the ethanol had fully evaporated. All scaffolds were a gift from
12 Dr Sanjay Sinha and obtained from Biomedical Structures (Biofelt).

13 ECOs were harvested and dissociated to single cells as previously described
14 for flow cytometry analyses. $5-10 \times 10^6$ cells were re-suspended in 100 μ l of
15 William's E medium with supplements, seeded on a scaffold surface area of
16 1cm^2 and incubated at 37°C for 30-60 minutes to allow the cells to attach to
17 the scaffold. The scaffolds were placed in wells of a 24-well plate and
18 checked at regular intervals during this period to ensure the medium did not
19 evaporate. If necessary, 10-20 μ l of William's E medium with supplements
20 were added. After 1 hour, 2-3 mls of William's E medium with supplements
21 were added to the wells and the medium was changed twice weekly.

22 **Generation of densified collagen tubes**

23 Densified collagen tubes were prepared using a novel approach. A 3D printed
24 chamber was fabricated, consisting of a funnel piece and a base plate. A
25 $250\mu\text{m}$ thick metallic wire was mounted into the base plate and fed through

1 the centre of the funnel. Absorbent paper towels were compacted between
2 the two 3D printed parts, which were then screwed together. 5 mg mL⁻¹
3 collagen gel solution, loaded with cells, was poured into the funnel and gelled
4 at 37°C for 30 min. After that time, the screws were loosened and, by placing
5 the 3D printed chambers at 37°C for 2-4h, water was drawn out of the
6 collagen gel. A cell-loaded densified collagen tube was thus formed with a
7 250µm lumen and a wall thickness of 30-100 µm, determined by the duration
8 of the drying phase. Upon removal from the chamber, the tube was trimmed
9 for excess collagen and cut to the required length.

10 **Culture of Human Mammary Epithelial Cells (HMECs)**

11 HMECs and the required tissue culture consumables were purchased as a kit
12 from Lonza (cat no. cat no. CC-2551B) and the cells were cultured according
13 to the supplier's instructions

14 **Animal experiments**

15 All animal experiments were performed in accordance with UK Home Office
16 regulations. Immunodeficient NOD.Cg-Prkdc^{scid} Il2rg^{tm1Wjl}/SzJ (NSG) mice
17 which lack B, T and NK lymphocytes (32) were bred in-house with food and
18 water available ad libitum pre- and post-procedures. A mix of male and female
19 animals were used, aged approximately 6-8 weeks. All the ECO-constructs
20 used were populated with ECOs derived from the common bile duct.

21 **Generation of ExtraHepatic Biliary Injury mouse model**

22 To generate a model of extrahepatic biliary injury, midline laparotomy was
23 performed and the gallbladder was first mobilized by dividing the ligamentous
24 attachment connecting its fundus to the anterior abdominal wall under
25 isoflurane general anesthesia. A longitudinal incision was then made along

1 2/3 of the length of the gallbladder, from the fundus towards Hartmann's
2 pouch (neck of gallbladder).

3 **Biliary reconstruction in EHBI mice**

4 To reconstruct the gallbladder, a scaffold section measuring approximately 1 x
5 1 mm (seeded with ECOs or without ECOs in controls) was sutured as a
6 'patch' to close the defect using 4 – 6 interrupted 10'0 non-absorbable nylon
7 sutures under 40x magnification. The laparotomy was closed in two layers
8 with continuous 5'0 absorbable Vicryl sutures. The animals were given
9 buprenorphine (temgesic 0.1 mg/kg) analgesia as a bolus and observed every
10 15 minutes in individual cages until fully recovered.

11 8 animals underwent biliary reconstruction using an ECO-populated scaffold.
12 All animals survived up to 104 days without complications and were culled
13 electively for further analyses. Two control experiments were performed,
14 where the animals underwent biliary reconstruction using acellular scaffolds.
15 Both animals died within 24 hours from bile leak, therefore no further control
16 experiments were performed to minimize animal discomfort.

17 **Bile duct replacement**

18 The native common bile duct was divided and a short segment excised. The
19 populated densified collagen tube was anastomosed end-to-end, using
20 interrupted 10'0 nylon sutures, between the divided proximal and distal
21 common bile duct. A length of 5'0 nylon suture material (diameter 100 µm)
22 was inserted into the collagen tube and fed into the proximal and distal
23 common bile duct to ensure patency of the lumen during the anastomosis.
24 After the anastomosis was complete, the 5'0 suture was pushed into the
25 duodenum through the distal bile duct and was removed through an incision in

1 the duodenum, which was then closed with interrupted 10'0 nylon sutures.
2 Lumen patency was assessed at the time of transplantation through light
3 microscopy and cannulation of the lumen with a 5'0 non-absorbable suture.
4 Transplantation was abandoned as futile in case of fully occluded tubes due
5 to cell infiltration. These events were considered construct/tube failure rather
6 than surgical complications and therefore were not censored in the survival
7 analysis.

8 **Bile duct ligation**

9 C57BL/6 mice were purchased from The Jackson Laboratory (Bar Harbor,
10 ME). The mice were housed and bred in a Minimal Disease Unit at the animal
11 facility at Oslo University Hospital, Rikshospitalet, Oslo. All experiments were
12 performed on male mice between 8 and 12 weeks of age. We performed a
13 median laparotomy exposed the bile duct and ligated the common bile duct
14 close to the junction of the hepatic bile ducts. Sham operated mice underwent
15 the exact same procedure without ligation. Serum was harvested after 5 days.
16 Alanine transaminase (ALT), aspartate transaminase (AST) and alkaline
17 phosphatase (ALP) were measured in serum using an ADVIA 1800 (Siemens)
18 at The Central Laboratory, Norwegian School of Veterinary Science. All
19 animal experiments were approved by the Norwegian Food Safety Authority
20 (project license no FOTS 8210/15) and all animals received human care in
21 line with "Guide for the Care and Use of Laboratory Animals" (National
22 Institutes of Health Publication, 8th Edition, 2011).

23 **Blood sample collection**

1 Blood was taken using a 23g needles directly from the inferior vena cava
2 under terminal anaesthesia at the time the animals were electively culled and
3 transferred into 1.5ml Eppendorf tubes for further processing.

4 **Blood sample processing**

5 The blood samples were routinely processed by the University of Cambridge
6 Core biochemical assay laboratory (CBAL). All of the sample analysis was
7 performed on a Siemens Dimension EXL analyzer using reagents and assay
8 protocols supplied by Siemens.

9 **Light microscopy imaging**

10 Light microscopy images of excised reconstructed gallbladders were acquired
11 using a Leica MZFLIII fluorescence dissecting microscope. The images are
12 representative of 5 animals.

13 **Cryosectioning and Histology**

14 Excised gallbladders were fixed in 4% PFA, immersed in sucrose solution
15 overnight, mounted in optimal cutting temperature (OCT) compound and
16 stored at -80°C until sectioning. Sections were cut to a thickness of 10µm
17 using a cryostat microtome and mounted on microscopy slides for further
18 analysis

19 **Haematoxylin and Eosin Staining**

20 H&E staining was performed using Sigma-Aldrich reagents according to the
21 manufacturer's instructions. Briefly, tissue sections were hydrated, treated
22 with Meyer's Haematoxylin solution for 5 minutes (Sigma-Aldrich), washed
23 with warm tap water for 15 minutes, placed in distilled water for 30-60
24 seconds and treated with eosin solution (Sigma-Aldrich) for 30-60 seconds.

1 The sections were subsequently dehydrated and mounted using the Eukitt®
2 quick-hardening mounting medium (Sigma-Aldrich). Histology sections were
3 reviewed by an independent histopathologist with a special interest in
4 hepatobiliary histology (SD).

5 **TUNEL assay**

6 The TUNEL assay was performed using a commercially available kit (abcam,
7 ab66110) according to the manufacturer's instructions.

8 **Fluorescein Isothiocyanate (FITC) cholangiogram**

9 In situ FITC cholangiogram was performed in sacrificed animals after
10 dissection of the gallbladder free from the adherent liver lobes, but before
11 surgical interruption of the extrahepatic biliary tree. The distal bile duct was
12 cannulated with a 23½ gauge needle and FITC injected retrogradely into the
13 gallbladder and images taken under a fluorescent microscope.

14 **Magnetic resonance cholangiogram (MRCP)**

15 Magnetic resonance cholangiogram (MRCP) was performed after sacrifice of
16 the animals. MRCP was performed at 4.7T using a Bruker BioSpec 47/40
17 system. A rapid acquisition with relaxation enhancement sequence was used
18 with an echo train length of 40 echoes at 9.5ms intervals, a repetition time of
19 1000ms, field of view $5.84 \times 4.18 \times 4.18 \text{cm}^3$ with a matrix of $256 \times 180 \times 180$
20 yielding an isotropic resolution of $230 \mu\text{m}$. The actively-decoupled four-channel
21 mouse cardiac array provided by Bruker was used for imaging.

22 For the second mouse imaged, for higher signal to noise ratio to give
23 improved visualisation of the biliary ducts a two-dimensional sequence was
24 used with slightly varied parameters (24 spaced echoes at 11ms intervals to
25 give an effective echo time of 110ms; repetition time 5741ms; matrix size of

1 256×256; field of view of 4.33×5.35cm² yielding a planar resolution of
2 170×200µm²). Fifteen slices were acquired coronally through the liver and gall
3 bladder with a thickness of 0.6mm. For this acquisition, a volume coil was
4 used to reduce the impact of radiofrequency inhomogeneity.

5 To examine the biliary ducts and gall bladder, images were prepared by
6 maximum intensity projections. Structural imaging to rule out neoplastic
7 growths was performed using a T1-weighted 3D FLASH (fast low-angle shot)
8 sequence with a flip angle of 25°, repetition time of 14ms and an echo time of
9 7ms. The matrix was 512×256×256 with a field of view of 5.12×2.56×2.56cm³
10 for a final isotropic resolution of 100µm.

11 The MRCP images were reviewed by 2 independent radiologists with a
12 special interest in hepatobiliary radiology (EMG, SU).

13 **Statistical analyses**

14 All statistical analyses were performed using GraphPad Prism 6. For small
15 sample sizes where descriptive statistics are not appropriate, individual data
16 points were plotted. For comparison between 2 mean values a 2-sided
17 student's t-test was used to calculate statistical significance. The normal
18 distribution of our values was confirmed using the D'Agostino & Pearson
19 omnibus normality test where appropriate. Variance between samples was
20 tested using the Brown-Forsythe test. For comparing multiple groups to a
21 reference group one-way ANOVA with Dunnett correction for multiple
22 comparisons was used between groups with equal variance, while the
23 Kruskal-Wallis test with Dunn's correction for multiple comparisons was
24 applied for groups with unequal variance. Survival was compared using log-
25 rank (Mantel-Cox) tests. Where the number of replicates (n) is given this

1 refers to biological replicates or number of different animals unless otherwise
2 stated.

3 For animal experiments, initial group sizes were estimated based on previous
4 study variance. No statistical methods were used to calculate sample size. No
5 formal randomization method was used to assign animals to the experimental
6 group. However; animals were randomly picked from a cage by a technician
7 not involved in the study. No animals were excluded from the analysis. No
8 blinding was used when reviewing radiology images or histopathology slides.

9

10 **Author Contributions:** FS: Design and concept of study, execution of
11 experiments and data acquisition, development of protocols and validation,
12 collection and interpretation of data, production of figures, manuscript writing,
13 editing and final approval of manuscript; AWJ: Conception of the technique,
14 scaffold design and generation of densified collagen tubular scaffolds; OCT:
15 animal experiments including kidney capsule injections; cell culture, provision
16 and harvesting of mouse tissue; StS: Magnetic Resonance Imaging (MRI);
17 EMG, SSU: MRI review and reporting; RLG: Animal experiments, IF, tissue
18 histology; MCDB: Cell culture, generation of viral particles, viral transduction,
19 generation of GMP-ECOs; NLB, LV: Animal experiments; MJGV, PM:
20 Bioinformatics analyses; DO: Flow cytometry analyses; LY: Western blot
21 analyses; AB: Flow cytometry analyses, bioinformatics support; JB: Tissue
22 histology, IF; MarZ: Scaffold preparation; MTP: Generation of viral particles,
23 viral transduction, generation of GMP-ECOs; MP: Generation of viral particles;
24 GMS: scaffold generation; PMM,KES: maintenance and provision of fibroblast
25 controls; NP: tissue culture; NG, CAR: Harvesting and preparation of primary

1 tissue; IS: Karyotyping, CGH analyses; SD: Histology review and reporting;
2 WS, **JC**, KBJ, **MatZ**, SaS, WTHG, GJA, NRFH, SEB, TW, THK, EM: critical
3 revision of the manuscript for important intellectual content. AEM: Scaffold
4 design, critical revision of the manuscript for important intellectual content.
5 KSP: Primary tissue provision, animal experiments, design and concept of
6 study, study supervision, interpretation of data, editing and final approval of
7 manuscript. LV: Design and concept of study, study supervision, interpretation
8 of data, editing and final approval of manuscript.

9

10 **Acknowledgements:** This work was funded by ERC starting grant Relieve
11 IMDs (LV, NRFH), the Cambridge Hospitals National Institute for Health
12 Research Biomedical Research Center (LV, NRFH, FS.), the Evelyn trust
13 (NH) and the EU Fp7 grant TissuGEN (MCDB) and supported in part by the
14 Intramural Research Program of the NIH/NIAID (RLG, CAR). FS has been
15 supported by an Addenbrooke's Charitable Trust Clinical Research Training
16 Fellowship and a joint MRC-Sparks Clinical Research Training Fellowship.
17 AWJ and AEM acknowledge the support from EPSRC (EP/L504920/1) and an
18 Engineering for Clinical Practice Grant from the Department of Engineering,
19 University of Cambridge.

20 The authors would like to thank Dr Jeremy Skepper, Ms Lyn Carter and the
21 University of Cambridge Advanced Imaging Centre for their help with electron
22 microscopy; Dr Edward Farnell and the University of Cambridge, Cambridge
23 Genomic Services for their help with microarray data processing and analysis;
24 Professor Anna Petrunina and the NIHR Cambridge BRC Cell Phenotyping
25 Hub for their help with cell sorting; **Dr Keith Burling and the MRC MDU Mouse**

1 **Biochemistry Laboratory [MRC_MC_UU_12012/5];** the Cambridge
2 Biorepository for Translational Medicine for the provision of human tissue
3 used in the study and Mr Barlow McLeod for IT support. The monoclonal
4 antibody TROMA-III developed by R. Kemler was obtained from the
5 Developmental Studies Hybridoma Bank, created by the NICHD of the NIH
6 and maintained at The University of Iowa, Department of Biology, Iowa City,
7 IA 52242.

8

9 **Competing interests:** LV is a founder and shareholder of DefiniGEN. The
10 remaining authors have nothing to disclose.

1 **References**

- 2 1. Murray, K. F. & Carithers, R. L. AASLD practice guidelines: Evaluation
3 of the patient for liver transplantation. *Hepatology* **41**, 1407–1432
4 (2005).
- 5 2. Perkins, J. D. Are we reporting the same thing?: Comments. *Liver*
6 *Transplant.* **13**, 465–466 (2007).
- 7 3. Skaro, A. I. *et al.* The impact of ischemic cholangiopathy in liver
8 transplantation using donors after cardiac death: The untold story.
9 *Surgery* **146**, 543–553 (2009).
- 10 4. Enestvedt, C. K. *et al.* Biliary complications adversely affect patient and
11 graft survival after liver retransplantation. *Liver Transpl.* **19**, 965–72
12 (2013).
- 13 5. Gallo, A. & Esquivel, C. O. Current options for management of biliary
14 atresia. *Pediatr. Transplant.* **17**, 95–98 (2013).
- 15 6. Felder, S. I. *et al.* Hepaticojejunostomy using short-limb Roux-en-Y
16 reconstruction. *JAMA Surg* **148**, 253–7-8 (2013).
- 17 7. Sampaziotis, F., Segeritz, C.-P. & Vallier, L. Potential of human induced
18 pluripotent stem cells in studies of liver disease. *Hepatology* **62**, 303–
19 311 (2015).
- 20 8. Kanno, N., LeSage, G., Glaser, S., Alvaro, D. & Alpini, G. Functional
21 heterogeneity of the intrahepatic biliary epithelium. *Hepatology* **31**, 555–
22 61 (2000).
- 23 9. Sampaziotis, F. *et al.* Cholangiocytes derived from human induced

- 1 pluripotent stem cells for disease modeling and drug validation. *Nat.*
2 *Biotechnol.* 1–11 (2015). doi:10.1038/nbt.3275
- 3 10. LeSage, G., Glaser, S. & Alpini, G. Regulation of cholangiocyte
4 proliferation. *Liver* **21**, 73–80 (2001).
- 5 11. Huch, M. *et al.* Long-Term Culture of Genome-Stable Bipotent Stem
6 Cells from Adult Human Liver. *Cell* **160**, 299–312 (2014).
- 7 12. Masyuk, A. I., Masyuk, T. V & LaRusso, N. F. Cholangiocyte primary
8 cilia in liver health and disease. *Dev. Dyn.* **237**, 2007–12 (2008).
- 9 13. Tabibian, J. H., Masyuk, A. I., Masyuk, T. V., O’Hara, S. P. & LaRusso,
10 N. F. Physiology of cholangiocytes. *Compr. Physiol.* **3**, 541–565 (2013).
- 11 14. Cízková, D., Morký, J., Micuda, S., Osterreicher, J. & Martínková, J.
12 Expression of MRP2 and MDR1 transporters and other hepatic markers
13 in rat and human liver and in WRL 68 cell line. *Physiol. Res.* **54**, 419–28
14 (2005).
- 15 15. Gigliozzi, A. *et al.* Molecular identification and functional
16 characterization of Mdr1a in rat cholangiocytes. *Gastroenterology* **119**,
17 1113–22 (2000).
- 18 16. Xia, X., Francis, H., Glaser, S., Alpini, G. & LeSage, G. Bile acid
19 interactions with cholangiocytes. *World J. Gastroenterol.* **12**, 3553–63
20 (2006).
- 21 17. Caperna, T. J., Blomberg, L. A., Garrett, W. M. & Talbot, N. C. Culture
22 of porcine hepatocytes or bile duct epithelial cells by inductive serum-
23 free media. *In Vitro Cell. Dev. Biol. Anim.* **47**, 218–33 (2011).

- 1 18. Marinelli, R. A. *et al.* Secretin induces the apical insertion of aquaporin-
2 1 water channels in rat cholangiocytes. *Am. J. Physiol.* **276**, G280-6
3 (1999).
- 4 19. Gong, A.-Y. *et al.* Somatostatin stimulates ductal bile absorption and
5 inhibits ductal bile secretion in mice via SSTR2 on cholangiocytes. *Am.*
6 *J. Physiol. Cell Physiol.* **284**, C1205-14 (2003).
- 7 20. Ito, M. *et al.* NOD/SCID/gamma(c)(null) mouse: an excellent recipient
8 mouse model for engraftment of human cells. *Blood* **100**, 3175–82
9 (2002).
- 10 21. Jabłonska, B. End-to-end ductal anastomosis in biliary reconstruction:
11 indications and limitations. *Can. J. Surg.* **57**, 271–277 (2014).
- 12 22. Gjorevski, N. *et al.* Designer matrices for intestinal stem cell and
13 organoid culture. *Nature* **539**, 560–564 (2016).
- 14 23. Koo, B. K. & Clevers, H. Stem cells marked by the r-spondin receptor
15 LGR5. *Gastroenterology* **147**, 289–302 (2014).
- 16 24. Farin, H. F., Van Es, J. H. & Clevers, H. Redundant sources of Wnt
17 regulate intestinal stem cells and promote formation of paneth cells.
18 *Gastroenterology* **143**, 1518–1529.e7 (2012).
- 19 25. Du, P., Kibbe, W. a & Lin, S. M. lumi: a pipeline for processing Illumina
20 microarray. *Bioinformatics* **24**, 1547–8 (2008).
- 21 26. Lin, S. M., Du, P., Huber, W. & Kibbe, W. A. Model-based variance-
22 stabilizing transformation for Illumina microarray data. *Nucleic Acids*
23 *Res.* **36**, (2008).

- 1 27. Smyth, G. K. Linear models and empirical bayes methods for assessing
2 differential expression in microarray experiments. *Stat Appl Genet Mol*
3 *Biol* **3**, Article3-Article3 (2004).
- 4 28. Bertero, A. *et al.* Activin/Nodal signaling and NANOG orchestrate
5 human embryonic stem cell fate decisions by controlling the H3K4me3
6 chromatin mark. *Genes Dev.* **29**, 702–17 (2015).
- 7 29. Campos, P. B., Sartore, R. C., Abdalla, S. N. & Rehen, S. K.
8 Chromosomal spread preparation of human embryonic stem cells for
9 karyotyping. *J. Vis. Exp.* 4–7 (2009). doi:10.3791/1512
- 10 30. Hannan, N. R. F. *et al.* Generation of multipotent foregut stem cells from
11 human pluripotent stem cells. *Stem Cell Reports* **1**, 293–306 (2013).
- 12 31. Koo, B.-K., Sasselli, V. & Clevers, H. Retroviral gene expression control
13 in primary organoid cultures. *Curr. Protoc. Stem Cell Biol.* **27**, Unit 5A.6.
14 (2013).
- 15 32. Shultz, L. D. *et al.* Human lymphoid and myeloid cell development in
16 NOD/LtSz-scid IL2R gamma null mice engrafted with mobilized human
17 hemopoietic stem cells. *J. Immunol.* **174**, 6477–6489 (2005).

18

19

1 **Figure Legends**

2

3 **Figure 1**

4 Derivation and characterization of Extrahepatic Cholangiocyte Organoids
5 (ECOs). **(a)** Schematic representation of the method used for the derivation of
6 ECOs. **(b)** Photograph of a 24-well plate well containing ECO organoids under
7 3D culture conditions. Scale bar: 1cm **(c)** Light microscopy image of ECO
8 organoids. Scale bar: 1 mm **(d)** Transmitted electron microscopy picture of
9 ECOs demonstrating the presence of cilia (black arrowhead). Scale bar:
10 500nm **(e)** Quantitative real time PCR (QPCR) confirming the expression of
11 biliary markers in Passage 1 (P1), Passage 10 (P10) and Passage 20 (P20)
12 ECOs compared to freshly isolated Primary Cholangiocytes (PCs) and
13 Embryonic Stem (ES) cells used as a negative control, n=4 biological
14 replicates. Center line, median; box, interquartile range (IQR); whiskers, range
15 (minimum to maximum). Values are relative to the housekeeping gene
16 Hydroxymethylbilane Synthase (*HMBS*) **(f)** Immunofluorescence (IF) analyses
17 confirming the expression of biliary markers in ECO organoids. Scale bars:
18 100 μ m. Single channel and higher magnification images are provided in
19 Supplementary Figure 6a. **(g)** Euclidian hierarchical clustering analysis
20 comparing the transcriptome of primary cholangiocytes (Primary), passage 20
21 ECOs (ECO), iPS-derived intrahepatic cholangiocyte-like-cells (iChoLC), ES
22 cells (ES) and hepatocytes (HEP). For each probe, standard scores (z-
23 scores) indicate the differential expression measured in number of standard
24 deviations from the average level across all the samples. Clusters of genes

1 expressed in ECOs, primary cholangiocytes or both cell types are indicated.
2 GO analyses for each cluster are provided in Supplementary Figure 7e. The
3 data corresponds to biological triplicates.

4

5 **Figure 2**

6 Functional characterization of ECO organoids. **(a)** Fluorescence images
7 demonstrating secretion of the MDR1 fluorescent substrate rhodamine 123 in
8 the lumen of ECOs. Luminal accumulation of rhodamine is inhibited the MDR1
9 inhibitor verapamil, confirming MDR1 activity. Scale bars: 100 μm . **(b)**
10 Fluorescence intensity along the red line in **(a)**. **(c)** Mean intraluminal
11 fluorescence intensity normalized to background in freshly plated Primary
12 Cholangiocytes (Rho PCs), Passage 20 ECOs (Rho P20) and P20 ECOs
13 treated with verapamil (Ver). Error bars, Standard Deviation (SD); n=1565
14 measurements in total. Asterisks (****) indicate statistical significance
15 ($P<0.001$, Kruskal-Wallis test with Dunn's correction for multiple comparisons)
16 **(d)** Luminal extrusion of the fluorescent bile acid CLF compared to controls
17 loaded with Fluorescein Isothiocyanate (FITC), confirming active bile acid
18 transfer. Scale bars: 100 μm . **(e)** Fluorescence intensity along the red line in
19 **(d)**. **(f)** Mean intra-luminal fluorescence intensity normalized over background,
20 n=1947 total measurements. Error bars, SD; asterisks as in **(c)**. **(g)** ALP
21 staining of ECOs. Scale bars: Light microscopy: 500 μm , Whole well images:
22 1cm. **(h)** Mean GGT activity of P20 ECOs vs. PCs; error bars, SD; n=3,
23 asterisks as in **(c)**. **(i,j)** Mean diameter measurements **(i)** and live images **(j)**
24 of ECOs treated with secretin or secretin and somatostatin, n=8. Error bars,

1 SD; *** $P < 0.001$; # $P > 0.05$ (Kruskal-Wallis test with Dunn's correction for
2 multiple comparisons). **(a-j)** Data representative of 3 different experiments.

3

4 **Figure 3**


5 Biliary reconstruction in an extrahepatic biliary injury (EHBI) mouse model
6 using ECOs. **(a)** Schematic representation of the method used for biliary
7 reconstruction. **(b)** Kaplan–Meier survival analysis, demonstrating rescue of
8 EHBI mice following biliary reconstruction with ECO-populated scaffolds.
9 ** $P < 0.01$ (log-rank test). **(c)** Images of gallbladders reconstructed with
10 acellular PGA scaffolds (scaffold only), PGA scaffolds populated with ECOs
11 (transplanted) and native not reconstructed gallbladder controls (not
12 transplanted), demonstrating full reconstruction with ECO populated scaffolds.
13 CD: cystic duct, CBD: common bile duct, CHD: common hepatic duct, F:
14 fundus, A: anterior surface, P: posterior surface. Scale bars: 500 μm . **(d)** H&E
15 staining demonstrating physiological architecture of the reconstructed
16 gallbladders. L: lumen. Scale bars: 100 μm **(e)** IF analyses demonstrating the
17 presence of GFP-positive ECOs expressing biliary markers in the
18 reconstructed gallbladders. L: lumen Scale bars: 100 μm . Higher
19 magnification images are provided in supplementary figure 14 **(f,g)** FITC
20 cholangiogram (n=1) **(f)** and Magnetic Resonance Cholangio-
21 Pancreatography (MRCP) images (n=2) **(g)** of reconstructed (transplanted)
22 vs. native control (not transplanted) gallbladders (GB) demonstrating a patent
23 lumen and unobstructed communication with the rest of the biliary tree. Scale
24 bars: 1mm


1

2 **Figure 4**


3 Bile duct replacement using ECO-populated densified collagen tubes. **(a)**
4 Schematic representation of the method used. **(b)** Postmortem images of
5 mice receiving ECO-populated collagen tubes (ECOs) vs. mice receiving
6 fibroblast-populated tubes (fibroblasts). Bile flow results in yellow
7 pigmentation of ECO-tubes. The white color of the fibroblast conduit
8 combined with a dilated bile-filled (yellow color) proximal bile duct (PDB)
9 suggests luminal occlusion, resulting in bile leak (yellow peritoneal
10 pigmentation; white dashed line). SC: Collagen tubes/scaffolds; DBD: Distal
11 Bile duct; scale bars 500 μ m. **(c)** Images of a thin walled construct resembling
12 the native bile duct in animals receiving ECO-populated tubes vs. a thickened
13 construct with no distinguishable lumen in animals receiving fibroblast tubes.
14 Scale bars 500 μ m. **(d)** QPCR using human-specific primers confirming the
15 expression of biliary markers by transplanted ECO-populated tubes (ECOs in
16 vivo) compared to cultured ECOs (ECOs in vitro) and mouse biliary tissue
17 used as a negative control, n=4 replicates. Center line, median; box,
18 interquartile range (IQR); whiskers, range (minimum to maximum). Values are
19 relative to *HMBS* expression. **(e)** H&E staining demonstrating the presence of
20 a biliary epithelium and a patent lumen in ECO-tubes but not fibroblast
21 constructs. Scale bars 100 μ m. **(f)** IF analyses demonstrating a GFP+/ CK19+
22 epithelium lining the lumen of ECO-constructs, vs. obliteration of the lumen by
23 fibroblasts in fibroblast constructs. Scale bars 100 μ m. **(g)** FITC
24 cholangiogram, demonstrating lumen patency in ECO-tubes vs. lumen

- 1 occlusion in fibro-constructs. Scale bars: 500 μ m **(h)** ALP activity is observed
- 2 only in ECO-tubes, but not in fibroblast constructs. Scale bars: 500 μ m






a




b

Survival curve




Number of mice at risk						
Days	0	1	17	22	75	104
Control	2	2	0	0	0	0
Scaffold	8	8	8	5	4	3


c




d




e




f



g





1 **Supplementary Note**

2 **Supplementary Results**

3 To assess the potential of ECOs for tissue engineering, we first interrogated their
4 capacity for populating Polyglycolic Acid (PGA) biodegradable scaffolds commonly
5 used to provide the structural and mechanical support required for tissue
6 reconstruction (1). Indeed, PGA is one of the most widely used synthetic polymers
7 since it does not induce inflammatory responses in the surrounding tissue; it is
8 biodegradable; and it is more flexible and easier to process compared to natural
9 polymers such as collagen (2). To facilitate tracking of the cells, ECOs expressing
10 Green Fluorescent Protein (GFP) were generated through viral transduction
11 (Supplementary Fig. 10a-10b). The resulting cells were seeded on PGA scaffolds,
12 attached to the PGA fibers after 24-48 hours and continued to grow for 4 weeks until
13 the scaffold was confluent (Supplementary Fig 11a-11d). Primary cholangiocytes
14 plated in 2D conditions demonstrated limited expansion potential and failed to reach
15 confluency when seeded on the scaffolds (Supplementary Fig 12a-b), suggesting
16 that the proliferative capacity of ECOs is crucial for successful scaffold colonization.
17 The populated PGA scaffolds (Supplementary Fig 11b-11c), could easily be handled
18 with forceps and divided into smaller pieces with a surgical blade. Furthermore, the
19 cells populating the scaffolds retained expression of biliary markers such as CK7 and
20 CK19 (Supplementary Fig 11e-11f), **demonstrated no evidence of EMT**
21 **(Supplementary Fig. 11e, 11g)** and maintained their functional properties including
22 ALP and GGT activity (Supplementary Figure **11h-11i**). Therefore, ECOs can
23 successfully populate PGA scaffolds, while maintaining their functionality and marker
24 expression.

1 We then focused on the generation of a tubular ECO-populated scaffold, which could
2 be used in the context of bile duct replacement surgery. The internal diameter of the
3 mouse CBD is approximately 100µm with a wall thickness of less than 50µm, which
4 precluded the use of a PGA scaffold due to mechanical properties. Instead, we
5 generated densified collagen tubular scaffolds (Supplementary Fig 15a-15b) which
6 were populated with GFP-expressing ECOs (supplementary Fig 15c-15e). The use
7 of densified collagen enabled the generation of constructs with an external diameter
8 ranging from 250 to 600µm and adequate strength to maintain a patent lumen
9 (Supplementary Fig 15d). Importantly, the cells populating the collagen scaffolds
10 maintained expression of biliary markers such as CK19, CK7, HNF1b, Sox9 and
11 CFTR (Supplementary Fig 15f-15g) and exhibited GGT and ALP enzymatic activity
12 (Supplementary Fig 15h-15i). **Primary epithelial cells of different origin (human
13 mammary epithelial cells; HMEC) failed to survive and adequately populate densified
14 collagen tubes under the same conditions (Supplementary Fig. 16a). Moreover,
15 plated HMECs failed to survive in a 10% (vol/vol) bile solution compared to ECOs
16 (Supplementary Fig. 16b), further confirming that ECOs constitute the only cell type
17 capable of generating bile resistant bio-engineered bile ducts. Collectively, these
18 results demonstrate the capacity of ECOs for populating tubular densified collagen
19 scaffolds without losing their original characteristics.**

20

21 **Supplementary discussion**

22 Our system provides proof-of-principle for the application of primary cells in
23 regenerative medicine; however, the use of stem cells has been suggested as an
24 alternative for cell based therapy. Although we have recently established a system

1 for the generation of stem cell-derived cholangiocyte-like cells (CLCs) (3), there are
2 significant differences between ECOs and CLCs that render ECOs better suited to
3 regenerative therapies for extrahepatic biliary injury. CLCs correspond to intrahepatic
4 cholangiocytes, while ECOs represent extrahepatic biliary epithelium. These two cell
5 types are distinct in terms of embryological origin and disease involvement (4).
6 Furthermore, CLCs still express fetal markers and therefore are more immature
7 compared to ECOs derived from primary cells (3). Therefore, CLCs may require a
8 period of adjustment and further maturation in vivo, while mature, functional cells,
9 such as ECOs, are required for coping with biliary injury in the acute setting. Finally,
10 although hPSCs provide a very good source of cells capable of generating almost
11 any tissue, fully differentiated CLCs cannot be expanded; initial derivation/
12 characterization of hPSC lines remains time consuming; while variability in capacity
13 of differentiation still constitutes a challenge. ECOs can be derived in less than 24
14 hours with a very high efficiency and can be expanded for multiple passages without
15 losing their original characteristics. Consequently, ECOs are comparable to CLCs in
16 terms of scalability, while their mature phenotype provides a unique advantage for
17 regenerative medicine applications in the context of tissue repair.

18

19 **Supplementary References**

- 20 1. Chan, B. P. & Leong, K. W. Scaffolding in tissue engineering: general
21 approaches and tissue-specific considerations. *Eur. Spine J.* **17**, 467–479
22 (2008).
- 23 2. Cheung, H.-Y., Lau, K.-T., Lu, T.-P. & Hui, D. A critical review on polymer-
24 based bio-engineered materials for scaffold development. *Compos. Part B*

1 *Eng.* **38**, 291–300 (2007).

2 3. Sampaziotis, F. *et al.* Cholangiocytes derived from human induced pluripotent
3 stem cells for disease modeling and drug validation. *Nat. Biotechnol.* 1–11
4 (2015). doi:10.1038/nbt.3275

5 4. Tabibian, J. H., Masyuk, A. I., Masyuk, T. V., O’Hara, S. P. & LaRusso, N. F.
6 Physiology of cholangiocytes. *Compr. Physiol.* **3**, 541–565 (2013).

7

1 **Supplementary Figure 1**

2 Screening for optimal conditions for the isolation and propagation of extra-hepatic
3 cholangiocytes **(a)** Mean viability ratio following isolation of primary cholangiocytes
4 with different methods. C+D: Collagenase + Dispase. Error bars, SD; n=3. Asterisks
5 represent statistically significant differences in viability ratio between mechanical
6 dissociation and other isolation methods; *** $P < 0.001$, **** $P < 0.0001$; one-way
7 ANOVA with Dunnett correction for multiple comparisons **(b)** Mean number of
8 resulting cells following 7 days of culture with various growth factors. E: Epidermal
9 Growth Factor, R: R-spondin, D: DKK-1, IL6: Interleukin-6, HGF: Hepatocyte Growth
10 Factor, VEGF: Vascular Endothelial Growth Factor, FBS: Foetal Bovine serum, F2:
11 Fibroblast Growth Factor (FGF) 2, F7: FGF-7, F10: FGF-10, A: Activin-A, SB: Activin
12 inhibitor SB-431542. Error bars, SD; n=3. Asterisks represent statistically significant
13 differences in the number of resulting cells between E+R+D and other culture
14 conditions; **** $P < 0.0001$; one-way ANOVA with Dunnett correction for multiple
15 comparisons **(c)** Representative live images of freshly isolated primary
16 cholangiocytes grown under different culture conditions for 7 days. EGF: Epidermal
17 Growth Factor. Scale bars: 500 μm . **(d)** Western blot analyses demonstrating
18 increased levels of phosphorylated β -catenin in ECOs treated with R-spondin and
19 DKK vs. R-spondin alone or R-spondin and the GSK-3 inhibitor CHIR 99021 (CHIR),
20 used as a positive control. **(e)** Quantification of the western blot demonstrated in
21 panel **(d)**. **(f)** ECOs treated with R-spondin and DKK exhibit increased Rho Kinase
22 activity, consistent with non-canonical Wnt signalling/ PCP pathway activation. Error
23 bars: SD; **** $P < 0.0001$; one way anova with Dunnett correction for multiple
24 comparisons; n=3; Y-27632: Rho Kinase inhibitor Y-27632.

25

1 **Supplementary Figure 2**

2 ECOs represent a highly homogeneous population of epithelial CK19+/CK7+ cells,
3 with minimal mesenchymal contamination. Flow cytometry analyses demonstrating
4 the expression of the biliary markers CK7 and CK19 but not the mesenchymal
5 marker vimentin (VIM) in freshly isolated primary cholangiocytes (Primary) and
6 Passage 20 (P20) ECOs, n=3. Fibroblasts are used as a positive control for
7 mesenchymal markers.

8

9 **Supplementary Figure 3**

10 Characterization of ECO lines derived from the gallbladder and common bile duct
11 brushings. **(a)** Schematic representation of the method for the derivation of ECOs
12 from the gallbladder. **(b)** Schematic representation of the method for the derivation of
13 ECOs from common bile duct brushings. **(c)** Quantitative real time PCR (QPCR)
14 confirming the expression of biliary markers in Passage 20 (P20) organoids of 2
15 ECO lines derived from common bile duct brushings (ECO_BR) and the gallbladder
16 (ECO_GB). Freshly isolated cholangiocytes (Primary) are used as a positive control,
17 n=4 biological replicates. Center line, median; box, interquartile range (IQR);
18 whiskers, range (minimum to maximum). Values are relative to the housekeeping
19 gene Hydroxymethylbilane Synthase (*HMBS*). **(d)** Immunofluorescence (IF) analyses
20 confirming the expression of biliary markers in P20 ECO_BR and ECO_GB
21 organoids. Scale bars: 100 μ m. **(e)** ALP staining of P20 ECO_BR and ECO_GB
22 organoids. Scale bars: 500 μ m. **(f)** Mean GGT activity of P20 ECO_BR and ECO_GB
23 vs. primary cholangiocytes; n=3; error bars, SD. These data are complementary to
24 the data shown in Figures 1 and 2.

1

2 **Supplementary Figure 4**

3 Genetic stability of ECOs following long-term in vitro culture for 20 passages. **(a)**
4 Growth curves of 3 different ECO lines cultured in vitro for 20 passages. **(b)** Normal
5 Karyotype of ECOs at P20; n=3. **(c)** Comparative Genomic Hybridization analyses
6 comparing ECOs at Passage 1 (red) and P20 (blue), demonstrating genomic stability
7 *in vitro*; n=3.

8

9 **Supplementary Figure 5**

10 ECOs exhibit ultra-structural features characteristic of cholangiocytes. Transmission
11 electron microscopy image demonstrating the presence of microvilli (white
12 arrowheads) and tight junctions (black arrowheads) in passage 20 ECO cells. **Scale**
13 **bar: 2 μ m.**

14

15 **Supplementary Figure 6**

16 ECOs express markers biliary markers. **(a-b)** Single channel **(a)** and high
17 magnification **(b)** images of the IF analyses demonstrated in Figure 1f demonstrating
18 the expression of biliary markers by ECOs. Scale bars: 100 μ m. These data are
19 complementary to the data shown in Figure 1.

20

21 **Supplementary Figure 7**

1 ECOs do not express markers of other lineages. **(a)** IF images demonstrating the
2 lack of expression of stem cell markers or markers of other lineages (liver, pancreas)
3 by ECOs. Scale bars: 100 μ m **(b)** IF images demonstrating appropriate positive
4 controls expressing the markers demonstrated in **(a)**. **(c)** QPCR analyses
5 demonstrating the lack of expression of stem cell markers or markers of other
6 lineages (liver, pancreas) by ECOs. Stel, activated stellate cells; Fibro, fibroblasts;
7 HEP, primary human freshly plated hepatocytes; Panc, primary human pancreatic
8 cells; ES, embryonic stem cells; HB, hIPSC-derived hepatoblasts; SC, ECO
9 populated scaffolds; **Int. Org., Intestinal Organoids**; n=4 biological replicates; n=3
10 biological replicates for POU5F1, NANOG, PROM1. Center line, median; box,
11 interquartile range (IQR); whiskers, range (minimum to maximum). Values are
12 relative to the housekeeping gene Hydroxymethylbilane Synthase (*HMBS*).

13

14 **Supplementary Figure 8**

15 Transcriptomic profile analyses of ECOs. **(a)** Euclidean hierarchical clustering
16 analysis comparing the transcriptome of ECOs across multiple passages (P1, P10,
17 P20) and Embryonic Stem (ES) cells used as a negative control, focusing on genes
18 that define the transcriptional signature of ECOs (4513 genes differentially
19 expressed between ECOs and ES cells). For each probe, standard scores (z-scores)
20 indicate the differential expression measured in number of standard deviations from
21 the average level across all the samples, n=3. **(b)** Heatmap showing the Pearson
22 correlation coefficient (r) of the global gene expression between ECOs across
23 multiple passages (P1, P10, P20), freshly isolated primary cholangiocytes (PCs),
24 hIPSC-derived intrahepatic Cholangiocyte Like Cells (iChoLC), primary hepatocytes

1 (HEPs) and ES used as a negative control, n=3. The analysis was performed
2 focusing on all probes detected in all 3 **biological** replicates. **(c)** Heatmap showing
3 the expression of representative biliary, hepatic and stem cell markers between
4 passage 20 ECOs (ECOs), PCs, HEPs and ES cells; n=3; z-scores as in **(a)**. **(c)**
5 Principal Component Analysis (PCA) between ECOs, PCs, iChoLC, HEPs and ES
6 cells, n=3. The analysis was performed using all probes detected in all 3 replicates.
7 ECOs cluster closely with primary cholangiocytes for component 1 accounting for
8 41% of total variability. **(d)** Gene ontology (GO) analyses focusing on the genes
9 differentially expressed between PCs and ECOs in Figure 1g; n=3. Characteristic
10 GO terms from each cluster are provided ($P<0.05$). Terms associated with the biliary
11 epithelium are identified only in the common cluster.

12

13 **Supplementary Figure 9**

14 ECOs injected under the kidney capsule of immune compromised mice form tubular
15 structures expressing biliary markers, but do not demonstrate spontaneous
16 differentiation in other lineages. **(a)** Schematic representation of the method used for
17 the injection of ECOs. **(b)** Image of an excised kidney transplanted with ECOs; scale
18 bar: 500 μ m. **(c)** Cross-section of a transplanted kidney demonstrating the formation
19 of tubular structures; scale bar: 500 μ m. **(d)** IF images demonstrating the formation of
20 tubular structures by the engrafted cells expressing the biliary marker CK7 and a
21 human-specific Ku80 epitope, but not expressing stem cell markers or markers of
22 other lineages (liver, pancreas). Scale bars: 100 μ m.

23

1 **Supplementary Figure 10**

2 Generation of ECOs expressing Green Fluorescent Protein (GFP). **(a)** Flow
3 cytometry analyses demonstrating the generation of a homogeneous population of
4 GFP-ECOs following lentiviral transduction and flow sorting. **(b)** Confocal microscopy
5 images demonstrating the generation of fully grown GFP-ECO organoids from a
6 single GFP-expressing cell. Scale bars: 100 μ m. Images are representative.

7

8 **Supplementary Figure 11**

9 ECOs dissociated to single cells (ECO-SCs) can populate biodegradable PGA
10 scaffolds. **(a,b)** Photographs of a PGA scaffold before **(a)** and after **(b)** treatment with
11 ECOs. Scale bars: 1cm. **(c)** Light microscopy images of a PGA scaffold populated
12 with ECO-SCs. Red arrowheads: Fully populated scaffold; black arrowheads: cells
13 recruiting new PGA fibers; white arrowheads: PGA fibers. Scale bars: 100 μ m. **(d)**
14 Confocal microscopy images demonstrating cell expansion at different time-points
15 after seeding of GFP-positive ECO-SCs on a PGA scaffold. White lines indicate the
16 position of PGA fibers. Scale bars: 100 μ m. **(e)** IF demonstrating the expression of
17 biliary markers and lack of EMT markers in ECO-SCs seeded on PGA scaffolds.
18 Scale bars: 50 μ m **(f)** QPCR analyses demonstrating the expression of biliary
19 markers in ECOs before (ECOs) and after (scaffold) seeding on PGA scaffolds, n=4
20 biological replicates. Center line, median; box, interquartile range (IQR); whiskers,
21 range (minimum to maximum). Values are relative to the housekeeping gene
22 Hydroxymethylbilane Synthase (*HMBS*). **(g)** Ratio of CK7+/CK19+ and
23 CK19+/Vimentin (VIM)+ cells in randomly selected IF images similar to the image
24 shown in **(e)**; n=6. **(h)** Mean GGT activity of ECO-SCs populating a PGA scaffold,

1 n=4. Error bars represent SD. **** $P<0.001$ (two-tailed t-test). **(i)** ALP staining of PGA
2 scaffolds populated by ECO-SCs. Scale bars: 500 μ m.

3

4 **Supplementary Figure 12**

5 2D plated cholangiocytes fail to populate PGA scaffolds. **(a)** Growth curves of
6 primary cholangiocytes plated as monolayer (2D) vs. primary cholangiocytes grown
7 as ECOs (ECO), demonstrating that 2D cholangiocytes stop proliferating after a few
8 passages. Starting from the same number of cells (5×10^5) 2D cholangiocytes fail to
9 provide the number of cells required to seed a PGA scaffold (10^7 cells). **(b)**
10 Brightfield images demonstrating that 2D cholangiocytes fail to expand and populate
11 PGA scaffolds and remain limited to the site of injection. A brightfield image of a
12 scaffold populated by ECOs dissociated to single cells (ECO-SC) is provided as a
13 positive control. The scaffold was seeded with same number of cells and cultured for
14 the same period of time as the 2D cholangiocyte scaffold. Scale bars: 100 μ m.

15

16 **Supplementary Figure 13**

17 Fibroblast-populated scaffolds fail to reconstruct the gallbladder following
18 transplantation in EHBI mice. **(a)** Flow cytometry analyses demonstrating the
19 generation of a homogeneous population of GFP-expressing Fibroblasts (GFP-Fibro)
20 following lentiviral transduction. **(b)** IF images demonstrating the expression of GFP
21 in fibroblasts following lentiviral transduction. Scale bars: 100 μ m. **(c, d)** Brightfield **(c)**
22 and confocal microscopy **(d)** images demonstrating the generation of fibroblast
23 populated PGA scaffolds. Scale bars: 100 μ m. **(e,f)** Postmortem images of

1 transplanted EHBI mice demonstrating failure to reconstruct the gallbladder with
2 fibroblast-populated scaffolds. The site of reconstruction can only be identified by the
3 presence of non-absorbable sutures (white arrowheads). Scale bars: 1mm. **(g)**
4 Image of a gallbladder transplanted with a fibroblast-populated scaffold
5 demonstrating failure to reconstruct the organ, abnormal morphology more
6 consistent with a fibrotic mass and lack of a bile filled lumen. Scale bars: 1mm. **(h)**
7 T2 phase Magnetic Resonance Images (MRI; axial plane) demonstrating the
8 absence of a bile filled lumen in a gallbladder transplanted with a fibroblast
9 populated scaffold. The white signal consistent with bile is identified up to the cystic
10 duct but not at the site of the transplanted gallbladder. **(i)** H&E images demonstrating
11 replacement of the gallbladder epithelium by fibroblasts and obliteration of the
12 gallbladder lumen. **(j)** IF analyses demonstrating the absence of GFP-expressing
13 cells in the transplanted gallbladders, the lack of epithelial markers and the presence
14 of connective tissue markers, Scale bars: 100µm.

15

16 **Supplementary Figure 14**

17 Characterization of ECO-reconstructed gallbladders in extrahepatic biliary injury
18 (EHBI) mice. **(a)** Postmortem images of mice receiving acellular PGA scaffolds
19 (scaffold only), healthy control mice (not transplanted) and mice receiving ECO-
20 populated PGA scaffolds (transplanted). The yellow pigmentation of the peritoneal
21 cavity (top, white dashed line) and seminal vesicles (bottom, black dashed line) is
22 consistent with bile leak in the 'scaffold only' group. Scale bars: 5mm **(b)** Post
23 mortem images demonstrating the reconstructed gallbladders in situ (n=3). Black
24 arrows: sutures; L: liver; scale bars: 1mm. **(c)** IF images of the transplanted

1 scaffolds, corresponding to figure 3e, demonstrating GFP-positive cells expressing
2 human (Ku80) and biliary markers (CK19, CK7, CFTR) integrated in the
3 reconstructed biliary epithelium; the presence of mouse stromal cells expressing
4 vimentin and mouse endothelial cells expressing CD31; and the presence of GFP+,
5 CK19-, Vimentin+ cells integrated in connective tissue of the reconstructed organ.
6 Scale bars: 100µm. **(d)** QPCR analyses using human-specific primers confirming
7 expression of biliary markers by transplanted ECO-populated scaffolds (ECOs in
8 vivo) compared ECOs in vitro and mouse gallbladder used as a negative control, n=4
9 replicates. Center line, median; box, interquartile range (IQR); whiskers, range
10 (minimum to maximum). Values relative to *HMBS* expression. **(e)** Ratio of
11 CK19+/CK7+, CK19+/GFP+ and Vimentin (VIM)/GFP+ cells quantified in randomly
12 selected sections of transplanted ECO-populated scaffolds; n=18. **(f)** T1&T2 vs. T2
13 phase Magnetic Resonance Images (MRI; axial plane) demonstrating bile in the
14 lumen of a reconstructed gallbladder (white signal) and a patent cystic duct relative
15 to the surrounding liver (n=2). GB: Gallbladder; CD: Cystic Duct; L: Liver
16 parenchyma.

17

18 **Supplementary Figure 15**

19 ECOs can populate densified collagen tubular scaffolds. **(a)** Schematic
20 representation of the method used. **(b)** Image of a densified collagen construct prior
21 to tube excision. Scale bar, 500µm. **(c)** Maximum intensity projection image
22 demonstrating a GFP+ ECO-populated tube after its generation. Scale bar; 10µm **(d)**
23 Confocal microscopy image demonstrating lumen patency of an ECO-populated
24 collagen tube. Scale bar; 10µm. **(e)** Images of a near confluent GFP+ ECO-tube.
25 Scale bar; 100µm. **(f)** IF analyses demonstrating the expression of biliary markers by

1 ECOs following the generation of ECO-tubes. Scale bar; 100 μ m. **(g)** QPCR analyses
2 demonstrating the expression of biliary markers before (ECOs) and after (Scaffold)
3 the generation of ECO-populated collagen tubes. ES cells are used as a negative
4 control, n=4 replicates. Center line, median; box, interquartile range (IQR); whiskers,
5 range (minimum to maximum). Values are relative to *HMBS* expression. **(h, i)** ECO-
6 tubes exhibit ALP **(h)** and GGT **(i)** activity. Scale bars, 500 μ m; MEFs, Mouse
7 Embryonic feeders used as negative control; Scaffold, ECO-populated, densified
8 collagen tubes; error bars, SD; n=3.

9

10 **Supplementary Figure 16**

11 Primary human mammary epithelial cells (HMECs) lack the capacity to adequately
12 populate densified collagen scaffolds or survive in the environment of primary
13 cholangiocytes. **(a)** Light microscopy images demonstrating failure of HMECs survive
14 and adequately populate collagen tubular scaffolds. Scale bar: 100 μ m. **(b)** Flow
15 cytometry analysis demonstrating survival of ECOs vs. plated HMECs following
16 exposure to a 10% (vol/vol) bile solution.

17

18 **Supplementary Figure 17**

19 Survival analysis and characterization of ECO-populated densified collagen tubes.
20 **(a)** H&E staining demonstrating the presence of a biliary epithelium and a patent
21 lumen in ECO-tubes. The images are complementary to Figure 4e **(b)** IF analyses
22 demonstrating the expression of human specific (Ku80), biliary (CK19, CK7, CFTR),
23 stromal (Vimentin, VIM) and vascular (CD31) markers in transplanted bio-tubes
24 populated with GFP⁺ ECOs. Apoptosis (TUNEL) and proliferation (Ki67) markers are

1 also demonstrated. Scale bars; 100 μ m. **(c)** IF images following TUNEL staining of
2 transplanted fibroblast-populated collagen tubes, demonstrating increased cell
3 death. Scale bars: 100 μ m. **(d)** Kaplan–Meier survival analysis demonstrating a
4 survival benefit in NSG mice following biliary reconstruction with ECO-populated
5 densified collagen tubes vs. fibroblast-populated densified collagen tube controls and
6 untreated controls. **** P <0.0001 (log-rank test). **(e)** Liver function tests of untreated
7 control animals (CTRL) vs. animals transplanted with ECO-populated tubes (ECOs)
8 and Bile Duct Ligation (BDL) animals used as a positive control, demonstrating
9 levels of serum cholestasis markers comparable to untreated controls following long-
10 term transplantation (n=5, 31 days; n=1, 27 days). Error bars, SD; ALP, Bilirubin
11 (Bili), one-way ANOVA with Dunnett correction for multiple comparisons; Alanine
12 aminotransferase (ALT), Kruskal-Wallis test; CTRL, n=11; ECOs, n=6; BDL, n=5.
13 ** P <0.01, **** P <0.0001, # P <0.05 (not statistically significant difference). **(f)** T2 phase
14 Magnetic Resonance Images (MRI; coronal plane) 1 month following transplantation
15 demonstrating bile in the lumen of a reconstructed bile duct (white signal) and a
16 patent construct lumen (n=2). SC: ECO-populated collagen tubular scaffold; L: Liver
17 parenchyma.

18

19 **Supplementary Video 1**

20 Magnetic Resonance Cholangio-Pancreatography (MRCP, sagittal plane) of an
21 Extrahepatic Biliary Injury (EHBI) mouse 104 days following biliary reconstruction
22 with an ECO-populated scaffold. The T2 weighed image sequence demonstrates the
23 presence of bile in the lumen of a reconstructed gallbladder (high intensity white
24 signal) and a patent cystic duct. The surrounding tissues are characterized by low
25 intensity signal and appear dark. To assess the anatomy of the surrounding organs

1 please refer to supplementary video 2 for a T1 weighed Magnetic Resonance
2 Imaging sequence.

3

4 **Supplementary Video 2**

5 T1 weighed Magnetic Resonance Imaging (MRI) (coronal plane) of an Extrahepatic
6 Biliary Injury (EHBI) mouse 104 days following biliary reconstruction with an ECO-
7 populated scaffold. This image sequence is optimal for demonstrating the anatomy
8 of the tissue surrounding the reconstructed gallbladder and biliary tree. However,
9 water, water-rich tissues and bile are characterized by low intensity signal, appear
10 dark and may be difficult to identify

11

12 **Supplementary video 3**

13 Time lapse images demonstrating the generation of a fully grown organoid from a
14 single cell.

15

16

1 **Supplementary table 1**

2 Donor demographics, corresponding to 8 different ECO lines. DCD: Donation after
3 Circulatory Death, DBD: Donation after Brain Death, F: Female, M: male, CBD:
4 Common Bile Duct, GB: Gallbladder, BR: CBD Brushings

5

Donor type	Blood Group	Age (years)	Gender	Site
DCD	O-	33	F	CBD
DBD	O+	56	F	CBD
DCD	A-	77	M	CBD
DBD	O+	57	M	CBD, GB
DCD	O-	44	M	CBD, GB
DCD	O+	48	M	CBD
DBD	A+	36	M	BR
DBD	B+	48	F	BR

6

7

8

9 **Supplementary table 2**

10 Microarray gene expression data corresponding to the heat map in Figure 1g

1 **Supplementary Table 3: List of antibodies used**

Target protein	Dilution	Company	Cat Number
CYTOKERATIN 19	1:100	ABCAM	ab7754
CYTOKERATIN 19 (KRT19)	1:50	DSHB	TROMA-III
SOX9 H-90	1:100	SANTA CRUZ	sc-20095
HNF1B (c-20)	1:100	SANTA CRUZ	sc-7411
CYTOKERATIN 7 (RCK105)	1:100	ABCAM	ab9021
CYTOKERATIN 7	1:100	ABCAM	ab68459
CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR (CFTR)	1:100	SANTA CRUZ	sc-10747
GAMMA GLUTAMYL TRANSPEPTIDASE (GGT)	1:100	ABCAM	ab55138
SOMATOSTATIN RECEPTOR 2	1:100	ABCAM	ab134152
SECRETIN RECEPTOR (C-20)	1:100	SANTA CRUZ	sc-26633
SECRETIN RECEPTOR	1:100	ABCAM	ab85565
ASBT (C14)	1:100	SANTA CRUZ	sc-27493
STEM101 HUMAN KU80 CELL NUCLEUS MARKER	1:100	STEM CELLS INC.	Y40400
ALEXA FLUOR DONKEY ANTI-Rabbit 568	1:1000	INVITROGEN	A10042
ALEXA FLUOR DONKEY ANTI-Rabbit 488	1:1000	INVITROGEN	A21206
ALEXA FLUOR DONKEY ANTI-Rabbit 647	1:1000	INVITROGEN	A31573
ALEXA FLUOR DONKEY ANTI-goat 568	1:1000	INVITROGEN	A11057
ALEXA FLUOR DONKEY ANTI-goat 488	1:1000	INVITROGEN	A11055
ALEXA FLUOR DONKEY ANTI-goat 647	1:1000	INVITROGEN	A21447
ALEXA FLUOR DONKEY ANTI-mouse 568	1:1000	INVITROGEN	A10037
ALEXA FLUOR DONKEY ANTI-mouse 488	1:1000	INVITROGEN	A21202
ALEXA FLUOR DONKEY ANTI-mouse 647	1:1000	INVITROGEN	A31571

2


3


4


1 **Supplementary table 4: List of primers used**




Gene	Primer sequence (5' à 3')	Gene	Primer sequence (5' à 3')
HNF1B	F TCACAGATACCAGCAGCATCAGT R GGGCATCACCAGGCTTGTA	CK7	F GATTGCTGGCCTTCGGGGT R TCATCACAGAGATATTCACGGCTC
PBGD	F GGAGCCATGTCTGGTAACGG R CCACGCGAATCACTCTCATCT	GGT	F GTGAGAGCAGTTGGCTGTGC R GTTGA ACTCTGCTGTGGGGC
SOX9	Hs_SOX9_1_SG QuantiTect Primer Assay (Quiagen, Cat Number: QT00001498)	CFTR	F AGTTGCAGATGAGGTTGGGC R AAAGAGCTTCACCCTGTCCG
CK19	F ACGACCATCCAGGACCTGCGG R TCCCACTTGGCCCCTCAGCGTA	SCR	F TGCTCACCAGCAGAAATGGT R AGGTAGGAGTGCCGCTTCTC
AQPR1	F GGCCAGCGAGTTCAAGAAGAA R TCACACCATCAGCCAGGTCAT	SSTR	F GAAAAGCAAAGATGTCACACTGGA R TTGGCATAGCGGAGGATGAC


2











a

CK19 / CK7

GGT / CK7


SCR / ASBT

CK7 / Sox9

CK7 / HNF1B

CK7 / CFTR

SCR / CK7



b

CK19 / CK7 /
DAPI

GGT / CK7 / DAPI


SCR / ASBT /
DAPI




CK7 / Sox9 /
DAPI




CK7 / HNF1B /
DAPI





CK7 / CFTR /
DAPI



SCR / CK7 /
DAPI
















a



b

2D cholangiocyte populated scaffold



ECO-populated scaffold

