

Recruitment of coat-protein-complex proteins on to phagosomal membranes is regulated by a brefeldin A-sensitive ADP-ribosylation factor

Walter BERÓN^{*1}, Luis S. MAYORGA^{*}, María I. COLOMBO^{*} and Philip D. STAHL[†]

^{*}Instituto de Histología y Embriología-CONICET, Universidad Nacional de Cuyo, Casilla de Correo 56, (5500) Mendoza, Argentina, and [†]Department of Cell Biology and Physiology, Washington University School of Medicine, St Louis, Missouri 63110, U.S.A.

Particle internalization in macrophages is followed by a complex maturation process. We have previously observed that proteins bound to phagocytosed particles are sorted from phagosomes into a heterogeneous population of vesicles that fuse with endosomes. However, the mechanism and the protein machinery involved in the formation of these phagosome-derived vesicles are largely unknown. It has been shown that vesicles coated with coat protein complex type I (COPI) have a role in both secretion and endocytosis. To address the possibility that COPI proteins might participate in the formation of phagosome-derived vesicles we studied the recruitment of β -COP to highly purified phagosomes. The binding of β -COP to phagosomal membranes was regulated by nucleotides and inhibited by brefeldin A (BFA). An ADP-ribosylation factor 1 (ARF1) mutant defective in GTP

hydrolysis supported the binding of β -COP to phagosomes independently of added nucleotide. AIF₄ and G $\beta\gamma$ subunits, agents known to modulate heterotrimeric G-protein activity, were tested in the β -COP binding assay. AIF₄ increased β -COP association, whereas binding was inhibited by the addition of G $\beta\gamma$ subunits. Our results suggest that COP proteins are recruited to phagosomal membranes by a mechanism that involves heterotrimeric GTP-binding proteins and a BFA-sensitive ARF. In addition, our findings indicate that COPI proteins are involved in the recycling of components from phagosomes to the cell surface.

Key words: phagocytosis, phagosomes, recycling, vesicular transport.

INTRODUCTION

Phagocytosis is responsible for the internalization of microorganisms, damaged cells or inert particles [1,2]. During this process, phagosomes actively exchange material with other intracellular compartments, mainly from the endocytic pathway, finally acquiring hydrolytic enzymes that eventually degrade the internalized particle. The mechanisms responsible for recycling from the phagosomal compartment are still largely unknown. However, they seem to involve the formation of tubules and transport vesicles that selectively include molecules to be recycled back to the plasma membrane [3].

Transport vesicles forming from a given compartment are known to be coated by specific sets of peripheral membrane-associated proteins such as clathrin, adaptor proteins (reviewed in [4]) and coat protein complex type I (COPI) and COPII [5,6]. Both COPI and COPII are involved in transport through the biosynthetic pathway. These coats participate in anterograde and retrograde transport between the endoplasmic reticulum and the Golgi apparatus [7,8]. COPI is a hetero-oligomeric complex composed of seven subunits (α , β , β' , γ , δ , ϵ and ζ) that is recruited to the membrane after the activation of the small GTPase ADP-ribosylation factor 1 (ARF1) (reviewed in [9]). Binding of ARF to the membranes depends on GDP-GTP exchange on the protein. The drug brefeldin A (BFA) inhibits the ARF exchange factor [10–14], blocking the assembly of COPI and altering normal Golgi transport [15].

Biochemical and immunocytochemical evidence has implicated both clathrin and coatamer coats in endocytic sorting and vesicle budding [16]. It has been shown that one of the subunits of the COPI complex, β -COP, is present on endosomes and is required

for the formation of vesicles that mediate transport to late endosomes [17,18]. Microinjection of antibodies against β -COP blocks infection by vesicular stomatitis virus, a virus internalized via the endocytic pathway [17]. In addition, a mutant Chinese hamster ovary (CHO) cell line with a temperature-sensitive defect in ϵ -COP has a defect in the sorting and recycling functions of endosomes [19,20]. There is also evidence that ARF is involved in the endocytic pathway. Studies *in vitro* have indicated that an ARF protein regulates endosome fusion [21,22]. Additionally, overexpression of an ARF6 mutant defective in GTP binding altered transferrin recycling and causes the accumulation of non-clathrin-coated structures on peripheral tubes [23,24]. Moreover, BFA, a drug that inhibits the Golgi-associated ARF1 exchange factor [10–12], changes the morphology of endosomes [15]. Indeed, it was shown recently that ARF1 regulates the binding of COP proteins to endosomes [25].

Given the similarities between the phagocytic and endocytic pathways, it is very likely that related COP-coated vesicles also participate in transport from the phagosomal compartment. To address the possibility that COPI-proteins might participate in the formation of phagosome-derived vesicles, we studied the recruitment of β -COP and β' -COP to highly purified phagosomes. The binding of β -COP to phagosomes was regulated by monomeric and heterotrimeric GTPases. The binding was sensitive to BFA.

With the use of a recycling assay to study the efflux of phagosomal components into the extracellular medium, we observed that recycling from phagosomes was partly inhibited in cells treated with BFA. Our findings indicate that COPI is involved in the recycling of components from the phagosomal compartment.

Abbreviations used: ARF, ADP-ribosylation factor; BFA, brefeldin A; CHO, Chinese hamster ovary; COP, coat protein complex; N₂ph, 2,4-dinitrophenol; GTP[S], guanosine 5'-[γ -thio]triphosphate; PNS, postnuclear supernatant; [S]ppG, guanosine 5'-[γ -thio]diphosphate; WT, wild-type ARF1.

¹ To whom correspondence should be addressed (e-mail wberon@fmed2.uncu.edu.ar).

EXPERIMENTAL

Biological materials

Clone J774-E, a mannose-receptor-positive macrophage cell line, was grown at 37 °C in minimal essential medium containing Earle's salts and supplemented with 10% (v/v) fetal bovine serum. Macrophage cytosol aliquots were gel-filtered through a 1 ml Sephadex G-25 spin column just before use in the binding assay.

Proteins were measured with BSA as a standard. Purified $\beta\gamma$ subunits of G-proteins were kindly provided by P. Casey (Duke University, Durham, NC, U.S.A.) and R. J. Lefkowitz (Duke University). M3A5, a mouse monoclonal antibody against β -COP, was kindly provided by T. Kreis (Universite of Geneva, Geneva, Switzerland). An affinity-purified polyclonal antibody directed against β' -COP was a gift from Felix Wieland (Universität Heidelberg, Heidelberg, Germany). BFA analogues were kindly provided by J. Donaldson (Laboratory of Cell Biology, NHLBI, National Institutes of Health, Bethesda, MD, U.S.A.). All other chemicals were obtained from Sigma Chemical Co. (St Louis, MO, U.S.A.).

Phagocytic probes

Formaldehyde-fixed *Staphylococcus aureus* (IgG-sorb; The Enzyme Center) were washed with HBSA [Hanks balanced salt solution buffered with 10 mM Hepes and 10 mM Tes, pH 7.4, and supplemented with 1% (w/v) BSA]. After incubation of 200 μ l of *S. aureus* suspension (10% cell suspension, approx. 4×10^7 particles/ml, 2 mg/ml IgG binding capacity) with 200 μ g of rabbit anti-mouse IgG polyclonal antibody (IgG fraction; Organon Teknika Corporation) for 1 h at 20 °C, the particles were washed three times with HBSA and incubated with 25 μ g of IgG raised against 2,4-dinitrophenol (N_2 ph) for 1 h at 20 °C. Coated *S. aureus* was washed three times and resuspended in HBSA.

Preparation and isolation of phagosomes

J774-E clone macrophages (10^8 cells) were incubated with antibody-coated *S. aureus* (200 μ l) for 1 h at 4 °C. Uptake was initiated by the addition of prewarmed HBSA. After 5 min at 37 °C, the incubation was stopped by the addition of ice-cold HBSA. Cells were subsequently washed with HBSA, HBSA supplemented with 5 mM EDTA and finally HBE buffer [250 mM sucrose/0.5 mM EGTA/20 mM Hepes/KOH (pH 7.2)] by centrifugation (300 g for 3 min). Cells were resuspended in HBE buffer to a final concentration of 2×10^8 cells/ml and homogenized in a homogenizer using ball bearings. Homogenates were centrifuged at 300 g for 3 min to eliminate nuclei and intact cells. The postnuclear supernatants (PNSs) were quickly frozen in liquid nitrogen and stored at -80 °C. Phagosomal fractions were prepared by diluting a quickly thawed PNS aliquot in 1 ml of HBE buffer and centrifuging at 12000 g in a Microfuge for 10 s at 4 °C. The supernatant was kept at 4 °C and the pellet was resuspended in HBE buffer and centrifuged again. The resulting supernatants were combined and centrifuged at 12000 g for 1.5 min at 4 °C. The pellet (phagosomal fraction) contained 70% of the total phagosomes in the sample. To isolate the phagosomes further, the phagosomal fraction was loaded on a discontinuous sucrose-step gradient [0.2 ml of 50% (w/v) sucrose and 0.4 ml of 65% (w/v) sucrose] and centrifuged at 128000 g for 45 min at 4 °C. The purified phagosomes were harvested from the pellet after resuspension in HBE buffer. To measure the phagosomal membrane protein concentration, purified phagosomes were resuspended with solubilization buffer [1% (v/v) Triton

X-100/0.2% methylbenzethonium chloride/1 mM EDTA/150 mM NaCl/10 mM Tris/HCl (pH 7.4)]. After incubation for 15 min on ice, the phagosomes were centrifuged for 1.5 min at 12000 g and the protein concentration was measured in the resulting supernatant (Triton-extracted proteins).

Labelling of plasma membrane

J774-E clone macrophages (10^8 cells) were incubated with 10 μ g/ml anti- N_2 ph IgG and 10 μ g/ml N_2 ph- β -glucuronidase for 1 h at 4 °C. Cells were subsequently washed with HBSA, HBSA supplemented with 5 mM EDTA and finally HBE buffer by centrifugation (300 g for 3 min at 4 °C). Cells were resuspended in HBE buffer to 2×10^8 cells/ml and homogenized in a homogenizer using ball bearings. Homogenates were centrifuged at 300 g for 3 min to eliminate nuclei and intact cells. The PNSs were quickly frozen in liquid nitrogen and stored at -80 °C.

The PNSs containing the plasma membrane markers were processed after the same steps used in the phagosome isolation protocol. To measure the plasma membrane markers, samples were mixed with solubilization buffer and the immune complex N_2 ph- β -glucuronidase-anti- N_2 ph IgG was immunoprecipitated in multiwell plates coated with rabbit anti-mouse IgG. After 30–45 min of incubation at room temperature, the wells were washed three times with solubilization buffer and the β -glucuronidase activity was measured with 4-methylumbelliferyl β -D-glucuronide as a substrate.

Preparation of Golgi membranes

A Golgi-enriched membrane fraction was obtained from J774-E clone macrophages by sucrose-gradient centrifugation as described previously [26]. Membranes were collected at the 0.8/1.2 M sucrose interface, frozen in liquid nitrogen and stored at -80 °C until use. Galactosyltransferase activity, a Golgi marker, was measured with an assay modified from Brew et al. [27].

β -COP binding assay

Isolated phagosomes (5–10 μ g of membrane proteins) were resuspended in a buffer containing 250 mM sucrose, 0.5 mM EGTA, 20 mM Hepes/KOH, pH 7.0, 1 mM dithiothreitol, 1.5 mM $MgCl_2$, 50 mM KCl, 1 mM ATP, 8 mM creatine phosphate and 31 units/ml creatine phosphokinase, supplemented with gel-filtered cytosol (1.2 mg/ml) in a final volume of 50 μ l, and incubated at 37 °C for 15 min. The reaction was stopped by chilling on ice and the addition of cold HBE buffer. The phagosomes were recovered by centrifugation (12000 g for 1.5 min at 4 °C) and resuspended in a buffer containing 1% (v/v) Triton X-100, 0.2% methylbenzethonium chloride, 1 mM EDTA, 150 mM NaCl and 10 mM Tris/HCl, pH 7.4, incubated for 15 min on ice and centrifuged again. The resulting supernatant was mixed with reducing sample buffer, boiled for 5 min and analysed by SDS/PAGE.

Western blotting

After resolution by SDS/PAGE, the proteins were transferred to nitrocellulose membranes with standard procedures. Membranes were blocked overnight at 4 °C in Blotto [5% (w/v) low-fat dried milk/80 mM Na_2HPO_4 /20 mM NaH_2PO_4 /100 mM NaCl (pH 7.5)] and then incubated for 1 h at 37 °C with a monoclonal antibody against β -COP (M3A5). Blots were incubated sequentially with rabbit anti-mouse and horseradish peroxidase-conjugated goat anti-rabbit immunoglobulins. β' -COP was detected with a specific affinity-purified antibody. Detection was per-

formed with the enhanced chemiluminescence Western blotting system (Amersham Corp.) in accordance with the supplier's recommendations.

Phagosome recycling assay

The recycling pathway between phagosomes and the cell surface was assayed as described by Pitt et al. [3]. In brief, J774-E clone macrophages were incubated for 30 min at 4 °C with formaldehyde-fixed *S. aureus* coated sequentially with rabbit anti-mouse antibody and then with mouse ¹²⁵I-labelled anti-N₂ph IgG. To optimize contact between cells and the phagocytic probe, the samples were centrifuged for 10 min at 1800 rev./min (500 g) in a benchtop centrifuge. Uptake was initiated by incubation at 37 °C for 5 min in a water bath and stopped by addition of cold PBS. After extensive washing at 4 °C to remove free particles, cells were incubated at 37 °C for different periods in HBSA. At the appropriate times, the culture medium was removed and assayed for trichloroacetic-acid-precipitable radioactivity (recycled material). The cell-associated radioactivity was assayed after cell solubilization with 0.5% (v/v) Triton X-100 in HBSA. This fraction was centrifuged for 5 min at 12000 g; *S. aureus*-associated radioactivity was measured in the pellet.

Electron microscopy

Purified phagosomes resuspended in HBE buffer were fixed with 1% (w/v) glutaraldehyde (prepared in HBE buffer). After fixation for 10 min, the samples were centrifuged for 15 min at 12000 g and the pellets were processed for transmission electron microscopy.

RESULTS

β-COP binds to highly purified phagosomes

To study the interaction of phagosomes with COP proteins, phagosomes of high purity were prepared with a discontinuous

Table 1 Analysis of the distribution of phagosomal, Golgi and plasma membrane markers in subcellular fractions obtained during the phagosome isolation protocols

The phagosomal marker was ¹²⁵I-labelled *S. aureus*, the Golgi marker was galactosyltransferase activity, and the plasma membrane marker was N₂ph-β-glucuronidase activity. Results are means ± S.D. for six experiments.

Cell fraction	Phagosomal marker (% of total)	Golgi marker (% of total)	Plasma membrane marker (% of total)
PNS	100	100	100
Purified phagosomes	31.5 ± 13.2	0.3 ± 0.5	0.3 ± 0.2

sucrose-density gradient as described in the Experimental section. Electron microscopic analysis of the highly purified phagosomes showed essentially no contamination by other organelles (Figure 1). The efficiency of phagosome isolation was assessed biochemically by measuring different subcellular markers. As shown in Table 1, 31% of the antibody-coated ¹²⁵I-labelled *S. aureus* originally located in the PNS was recovered in the pellet obtained after centrifugation of the PNS through a sucrose cushion. Plasma membrane and other endomembranes of similar density (e.g. endosomes and Golgi vesicles) were present only in trace amounts in the purified phagosomal fraction. Only 0.3% of a plasma membrane marker (N₂ph-β-glucuronidase-anti-N₂ph IgG complex bound to the cell surface) and 0.3% of a Golgi membrane marker (galactosyltransferase activity) of the totals present in the PNS were found in the phagosome fraction (Table 1).

It has been shown that GTP-binding proteins are important in the interaction between COPs and Golgi membranes [28–30]. Therefore, to study the recruitment of COP proteins to phagosomal membranes, we incubated purified phagosomes with cytosol (COPs source) in the presence of guanosine 5'-[γ-




Figure 1 Electron micrographs of purified phagosomes

Electron micrographs showing purified phagosome preparations at low (A) and high (B) magnifications. Scale bars, 0.67 μm (A) and 0.21 μm (B).




Figure 2 Effect of nucleotides and AIF₄ on β -COP binding to phagosomes

Binding of β -COP was assessed (see the Experimental section) after the incubation of purified phagosomes with cytosol for 20 min at 37 °C in the absence or presence of 25 μ M GTP[S] (GTP γ S), 200 μ M [S]ppG (GDP β S), 1 mM GTP, 100 μ M AINH₄(SO₄)₂ plus 10 mM KF or 100 μ M AINH₄(SO₄)₂ plus 10 mM KF plus 1 mM GTP.




Figure 3 Inhibition of GTP[S]-enhanced binding of β -COP to phagosomes by $G\beta\gamma$ subunits

Phagosomes and cytosol were preincubated with GTP[S] (GTP γ S) (20 μ M) or $G\beta\gamma$ subunits (500 nM) for 6 min at 37 °C. Subsequently, either $G\beta\gamma$ subunits or GTP[S] was added to the samples and incubated for a further 12 min. The samples were then processed as indicated in the Experimental section.

thio]triphosphate (GTP[S]), a non-hydrolysable GTP analogue. The binding of β -COP to phagosomes was analysed by SDS/PAGE and Western blot with a specific monoclonal antibody. As shown in Figure 2, cytosolic β -COP associated strongly with purified phagosomes in the presence of GTP[S] (lane 2). GTP also stimulated binding, although to a smaller extent than GTP[S] (Figure 2, lane 4). Lane 3 shows that guanosine 5'-[γ -thio]diphosphate ([S]ppG), a GDP analogue that locks GTP-binding proteins into the GDP-bound form, was a strong inhibitor of β -COP recruitment to phagosomes. As with β -COP, binding of β -COP to phagosome membranes was also observed in the presence of GTP[S] (results not shown).

AIF₄, an agent widely used to study heterotrimeric G-proteins [31], was also tested. As shown in Figure 2, AIF₄ stimulated the association of β -COP with phagosomes (lane 5) and the binding was increased markedly by the simultaneous addition of both GTP and AIF₄ (lane 6). These results suggest that one or more GTP-binding proteins regulate the interaction between COP proteins and phagosomes.

Because AIF₄ is an activator of trimeric G-proteins, the effect of AIF₄ suggests that this family of GTP-binding proteins is involved in the interaction of β -COP with phagosomes. $G\beta\gamma$ subunits of heterotrimeric G-proteins inhibit the function of these GTP-binding-proteins [32,33]. Therefore, if phagosomes and cytosol are preincubated with $G\beta\gamma$ subunits before adding GTP[S], the β -COP binding should not be stimulated by this nucleotide. When this sequential incubation was performed, a significant inhibition of β -COP binding to phagosomes was observed (Figure 3, lane 3). In contrast, when the membranes were first incubated with GTP[S] before the addition of $G\beta\gamma$




Figure 4 Pretreatments of phagosomes with different agents: effect on the binding of β -COP to phagosomes

The phagosomes were treated for 1 h at 4 °C with HBE buffer (control buffer), 0.8 M NaCl, 100 mM Na₂CO₃ or 50 mM EDTA in HBE buffer. For treatment with trypsin, phagosomes were incubated for 1 h at 4 °C with 10 μ g/ml trypsin followed by the addition of protease inhibitors (20 μ g/ml soybean trypsin inhibitor, 1 μ M leupeptin and 10 μ M PMSF) or were incubated with a mixture of trypsin and protease inhibitors. After treatment, the samples were washed with HBE buffer and the phagosomes were recovered by centrifugation. The phagosomes were resuspended with cytosol, incubated for 20 min at 37 °C with GTP[S] (20 μ M) and processed as indicated in the Experimental section.

subunits, the binding of β -COP to phagosomes was not altered (Figure 3, lane 2).

These results confirm the role of heterotrimeric G-proteins on the recruitment of β -COP to phagosomal membranes.

β -COP binding to phagosomal membrane requires membrane-associated proteins

To determine whether a membrane component was required for β -COP binding, phagosomes were treated with different agents. Membranes were stripped of peripherally associated proteins by incubation with 0.8 M NaCl or 100 mM Na₂CO₃. As shown in Figure 4, both treatments moderately decreased β -COP binding (lanes 2 and 3), although binding was not affected by washing the membranes in a buffer containing 50 mM EDTA (lane 4), suggesting that Ca²⁺ was not required for binding. In contrast, trypsin completely abolished the ability of the membranes to support β -COP binding [compare trypsin treatment (Figure 4, lane 5) with mock trypsin treatment (lane 6)].

Taken together, the results suggest that a membrane-associated factor is required for the interaction of β -COP with the phagosomal membranes.

BFA blocks the binding of β -COP to phagosomal membranes: role of ARF1

It has been shown that the binding of coatomer to the Golgi depends on the interaction of the small GTP-binding protein ARF with the membrane [29], a process that is blocked by BFA [10,11]. We therefore tested the effect of BFA on the binding of β -COP to phagosomes. As shown in Figure 5(A), BFA blocked the stimulatory effect of GTP[S] on the binding. The specificity of the effect of BFA is shown by the fact that the inactive analogues BFA-17 and BFA-31 did not inhibit the binding (Figure 5B).

The effect of BFA suggests that the binding of β -COP to phagosomes might be regulated by ARF. We therefore incubated phagosomes and cytosol in the presence of recombinant wild-type ARF1 (WT) or the GTPase-defective mutant (Q71L). As shown in Figure 6, WT bound to and stimulated the binding of β -COP to phagosomes when the protein was activated by adding GTP[S] to the assay (lane 2). As expected, the addition of [S]ppG




Figure 5 Effect of BFA on the binding of β -COP

(A) Phagosomes ($6 \mu\text{g}$ of membrane proteins) were preincubated for 6 min at 37°C in the presence of cytosol with $20 \mu\text{M}$ GTP[S] ($\text{GTP}\gamma\text{S}$) or $200 \mu\text{M}$ BFA, and subsequently for a further 12 min with BFA or GTP[S] respectively. As a control, all samples contained 1% ethanol (EtOH) in the BFA solution. (B) Phagosomes ($6 \mu\text{g}$ of membrane proteins) or Golgi membranes ($6 \mu\text{g}$ of membrane proteins) were preincubated for 6 min at 37°C in the presence of cytosol with either methanol (MeOH, 1%), BFA or one of the BFA analogues B17 or B31 (each at $300 \mu\text{M}$), and subsequently incubated for a further 12 min with GTP[S] ($25 \mu\text{M}$).




Figure 6 Effect of WT and mutant ARF1 on the binding of β -COP to phagosomes

Phagosomes and cytosol were incubated with WT (ARF1wt) or the ARF1 mutant ARF1Q71L (each at $100 \mu\text{g}/\text{ml}$) in the absence or presence of GTP[S] ($\text{GTP}\gamma\text{S}$) ($20 \mu\text{M}$), [S]ppG ($\text{GDP}\beta\text{S}$) ($200 \mu\text{M}$) or GTP ($200 \mu\text{M}$). After 20 min at 37°C , the samples were processed as indicated in the Experimental section.

blocked the recruitment of both ARF1 and β -COP to phagosomal membranes (Figure 6, lane 3). GTP did not support ARF1 and β -COP binding. In contrast, Q71L, the ARF mutant defective in GTP hydrolysis, bound more efficiently than WT and supported the binding of β -COP to phagosomes independently of added nucleotide (Figure 6, lanes 5–8), confirming the participation of active ARF1 in the process.




Figure 7 Effect of BFA on phagosomal recycling

J774-E clone macrophages treated for 30 min at 37°C with either 0.03% methanol (control, ○) or $10 \mu\text{M}$ BFA (●), were incubated for 30 min at 4°C with formaldehyde-fixed *S. aureus* coated sequentially with rabbit anti-mouse antibody followed by mouse ^{125}I -labelled anti- N_2ph IgG. Uptake was initiated by incubation at 37°C for 5 min and stopped by the addition of cold PBS. After being washed at 4°C , cells were incubated at 37°C for different durations in the presence of methanol (control) or BFA. At the indicated times, the culture medium was removed and assayed for trichloroacetic-acid-precipitable radioactivity (recycled material). Recycling was expressed as a percentage of the trichloroacetic-acid-precipitable radioactivity recovered in the medium at 30 min of chase. Results are means \pm S.E.M. for two independent experiments performed in duplicate. *At later time points, the difference between control and BFA-treated cells was statistically significant ($P < 0.05$, Student's *t* test).

BFA inhibits the recycling of protein from phagosomes to the cell surface

To study the recycling pathway between phagosomes and the cell surface, we measured the time-dependent appearance of phagocytosed ^{125}I -labelled anti- N_2ph IgG in the cell culture medium. J774-E clone macrophages internalized for 5 min at 37°C prebound phagocytic particles consisting of formaldehyde-fixed *S. aureus*, coated sequentially with rabbit anti-mouse antibody and mouse ^{125}I -labelled anti- N_2ph IgG. Uptake was stopped by the addition of cold PBS and cells were washed extensively at 4°C to remove free particles. Cells were incubated at 37°C for different durations (Figure 7). At the indicated times, culture media were collected and assayed for trichloroacetic-acid-precipitable radioactivity, which represented the recycled protein. We proved that this radioactivity did not come from detached proteins from the *S. aureus* during internalization because particles did not lose radioactivity after incubation for 30 min in culture medium (results not shown). In contrast, we had shown previously that the radioactive protein released to the medium was not re-internalized by endocytosis [3].

To test whether COP proteins might have a role in the phagosome-recycling pathway, we treated the cells for 30 min at 37°C with BFA before adding the phagocytic particles. BFA was also present during the uptake and the chase periods. The amount of radioactivity recovered in the culture medium was diminished in BFA-treated cells (Figure 7). The differences cannot be due to an effect of the BFA on the release of the radioactive protein from the particles because the *S. aureus*-associated protein was similar in control and treated cells during the incubation (results not shown). This shows that the recycling of proteins from phagosomes to the cell surface was partly inhibited by

BFA, suggesting that a BFA-sensitive factor was involved, at least in part, in the process.

DISCUSSION

Here we present evidence that β -COP, one of the components of the protein complex coatomer [34], is recruited to highly purified phagosomes. We have observed that both GTP[S] and AIF₄ promote the association of β -COP with phagosomal membranes, suggesting that one or more GTP-binding proteins participate in the binding of β -COP to the membranes. This is consistent with our previous observation that several heterotrimeric G-proteins are present on highly purified phagosomal membranes [35]. The results obtained with AIF₄ and G $\beta\gamma$ suggest that heterotrimeric G-proteins are likely to be involved in the recruitment of cytoplasmic proteins (e.g. β -COP) to make a particular type of vesicle, possibly by sending signals across the phagosomal membrane. Indeed, we have evidence that AIF₄ inhibits the recycling of proteins from the phagosomal compartment (M. T. Damiani, A. Savina and M. I. Colombo, unpublished work), suggesting that the formation of recycling vesicles might be regulated by heterotrimeric G-proteins.

We have observed that GTP increased the β -COP binding stimulated by AIF₄. Because AIF₄ activates the trimeric G-proteins only in the GDP-bound form [36], GTP might activate other GTP-binding proteins. As with the results observed with Golgi membranes [10], GTP[S] promoted the association of β -COP to phagosomal membranes and prevented the inhibitory effect of BFA. It has been shown that BFA prevents the assembly of coatomer on the Golgi membrane by inhibiting the factor that exchanges GTP bound to ARF [10–12,28]. Several BFA-sensitive guanine nucleotide exchange factors for ARFs have been identified [37–39]. Our results therefore suggest that a BFA-sensitive member of the ARF family is involved in the binding of β -COP to phagosomes. Indeed, our results indicate that ARF1 participates in the binding of β -COP to phagosomes.

ARF family members have been implicated in several events of intracellular membrane trafficking [40]. Published evidence indicates that ARF is involved in the endocytic pathway. It has been shown that an ARF protein regulates endosome fusion reconstituted in a system *in vitro* [21,22]. Moreover, in intact cells the overexpression of an ARF6 mutant defective in GTP binding altered transferrin recycling and caused the accumulation of non-clathrin-coated structures on peripheral tubes [23,24]. Apparently, in HeLa cells ARF6 is related to a plasma membrane–juxtannuclear compartment recycling pathway that is different from the conventional one characterized by the presence of the transferrin receptors [41]. Nevertheless, in CHO cells, a partial co-localization of ARF6 with transferrin receptor suggests that the ARF6-containing compartment might partly overlap with a recycling endosomal compartment [42]. It has been shown that ARF6 also has a role in Fc receptor-mediated phagocytosis through the regulation of actin-cytoskeleton changes that underlie phagocytosis [43]. However, it has recently been shown that ARF1 and not ARF6 is required for COP recruitment to endosomes and for the formation of endosomal carrier vesicles/multivesicular bodies from endosomal membrane *in vitro* [25].

A function for coatomer in the endocytic pathway has been proposed [17–20]. Microinjection of anti- β -COP antibodies inhibited the entry of viruses via the endocytic pathway [17]. These results, and the observation that ϵ -COP-deficient CHO cells exhibit defective endocytosis [44,45], suggest that coatomer or a related protein complex might have a role in the endocytic system. Indeed, a purified fraction of endosomes was shown to recruit *in vitro* a coat containing α -, β -, β' -, ϵ - and ζ -COP but not

γ - or δ -COP [17,18]. Interestingly, recruitment of these sets of proteins on endosomes was stimulated by GTP[S] and was sensitive to BFA [17]. These results are in agreement with our observation that the binding of β -COP to phagosomes is enhanced by GTP[S] and blocked by BFA.

Phagosome maturation is a complex process that involves the influx and efflux of material from phagosomes that gradually acquire lysosomal characteristics. The influx of material is a consequence of the interaction between phagosomes and endocytic compartments [3,46–52]. Protein and membrane recycling from phagosomes represents the efflux of material during the transformation of phagosomes to phagolysosomes. It has been shown that phagosomal proteins recycle rapidly to the plasma membrane [3,53]. Recycling seems to involve the formation of tubules and transport vesicles that selectively include molecules to be recycled back to the plasma membrane [3]. It is therefore possible that β -COP binds to phagosomes to allow the recycling of phagosome components to the cell surface. Indeed, the inhibition of normal recycling of either low-density lipoprotein or transferrin receptors to plasma membrane was observed in a temperature-sensitive ϵ -COP-deficient CHO cell mutant [19,44,45,54], suggesting that this COP protein is involved in endosome recycling.

It has recently been shown that the removal of transferrin receptor from phagosomes is partly inhibited in ϵ -COP-deficient or BFA-treated cells [55]. The authors suggest that transferrin receptor recycling is in part mediated by a COPI-dependent process. These results are in agreement with our observation that BFA significantly inhibited protein recycling from phagosomes to the cell surface.

Our observations suggest that COPI participates in delivering phagosome components towards the cell surface, a process that is essential for phagosome remodelling and antigen presentation.

We thank Lori LaRose, Marilyn Levy and Cheryl Adles for excellent technical assistance. This work was supported by the National Institutes of Health (to P. D. S.) and the National Research Council of Argentina (to W. B.).

REFERENCES

- Oh, Y. K and Swanson, J. A. (1996) Different fate of phagocytosed particles after delivery into macrophage lysosomes. *J. Cell Biol.* **132**, 585–593
- Aderem, A. and Underhill, D. M. (1999) Mechanisms of phagocytosis in macrophages. *Annu. Rev. Immunol.* **17**, 593–623
- Pitt, A., Mayorga, L. S., Schwartz, A. L. and Stahl, P. D. (1992) Transport of phagosomal components to an endosomal compartment. *J. Biol. Chem.* **267**, 126–132
- Smith, S. L. (1997) Clathrin-coated vesicle formation and protein sorting. *Annu. Rev. Biochem.* **66**, 511–548
- Letourneur, F., Gaynor, E. C., Hennecke, S., Demolliere, C., Duden, R., Erm, S. D., Riezman, H. and Cosson, P. (1994) Coatomer is essential for retrieval of dilysine-tagged proteins to the endoplasmic reticulum. *Cell* **79**, 1199–1207
- Barlowe, C., Orci, L., Yeung, T., Hosobuchi, M., Hamamoto, S., Salama, N., Rexach, M. F., Ravazzola, M. and Schekman, R. (1994) COPII: a membrane coat formed by sec proteins that drive vesicle budding from the ER. *Cell* **77**, 895–907
- Lowe, M. and Kreis, T. (1998) Regulation of membrane trafficking in animal cells by COPI. *Biochim. Biophys. Acta* **1404**, 53–66
- Barlowe, C. (1998) COPII and selective export from the endoplasmic reticulum. *Biochim. Biophys. Acta* **1404**, 67–76
- Kreis, T. E., Lowe, M. and Pepperkok, R. (1995) COPs regulating membrane traffic. *Annu. Rev. Cell Biol.* **11**, 677–706
- Donaldson, J. G., Finazzi, D and Klausner, R. D. (1992) Brefeldin A inhibits Golgi membrane-catalysed exchange of guanine nucleotide onto ARF protein. *Nature (London)* **360**, 350–352
- Helms, J. B. and Rothman, J. E. (1992) Inhibition by brefeldin A of a Golgi membrane enzyme that catalyses exchange of guanine nucleotide bound to ARF. *Nature (London)* **360**, 352–354
- Randazzo, P. A., Yang, Y. C., Rulka, C. and Kahn, R. A. (1993) Activation of ADP-ribosylation factor by Golgi membranes. Evidence for a brefeldin A- and protease-sensitive activating factor on Golgi membranes. *J. Biol. Chem.* **268**, 9555–9563

- 13 Chardin, P. and McCormick, F. (1999) Brefeldin A: the advantage of being uncompetitive. *Cell* **97**, 153–155
- 14 Roth, M. G. (1999) Snapshots of arf1: implications for mechanisms of activation and inactivation. *Cell* **97**, 149–152
- 15 Klausner, R. D., Donaldson, J. G. and Lippincott-Schwartz, J. (1992) Brefeldin A: insights into the control of membrane traffic and organelle structure. *J. Cell Biol.* **116**, 1071–1080
- 16 Stoorvogel, W., Oorschot, V. and Geuze, H. J. (1996) A novel class of clathrin-coated vesicles budding from endosomes. *J. Cell Biol.* **132**, 21–33
- 17 Whitney, J. A., Gomez, M., Sheff, D., Kreis, T. E. and Mellman, I. (1995) Cytoplasmic coat proteins involved in endosome function. *Cell* **83**, 703–713
- 18 Aniento, F., Gu, F., Parton, R. G. and Gruenberg, J. (1996) An endosomal β COP is involved in the pH-dependent formation of transport vesicles destined for late endosomes. *J. Cell Biol.* **133**, 29–41
- 19 Daro, E., Sheff, D., Gomez, M., Kreis, T. and Mellman, I. (1997) Inhibition of endosome function in CHO cell bearing a temperature-sensitive defect in the coatamer (COP) component ϵ -COP. *J. Cell Biol.* **29**, 1747–1759
- 20 Gu, F., Aniento, F., Parton, R. G. and Gruenberg, J. (1997) Functional dissection of COP-I subunits in the biogenesis of multivesicular endosomes. *J. Cell Biol.* **139**, 1183–1195
- 21 Lenhard, J. M., Kahn, R. A. and Stahl, P. D. (1992) Evidence for ADP-ribosylation factor (ARF) as a regulator of in vitro endosome–endosome fusion. *J. Biol. Chem.* **267**, 13047–13052
- 22 Lenhard, J. M., Colombo, M. I. and Stahl, P. D. (1994) Heterotrimeric GTP-binding proteins (G proteins) and ADP-ribosylation factor (ARF) regulate priming of endosomal membrane for fusion. *Arch. Biochem. Biophys.* **312**, 474–479
- 23 D'Souza-Schorey, C., Li, G., Colombo, M. I. and Sathl, P. D. (1995) A regulatory role for Arf6 in receptor mediated endocytosis. *Science* **267**, 1175–1178
- 24 Peters, P. J., Hsu, V. W., Ooi, C. E., Finazzi, D., Teal, S. B., Oorschot, V., Donaldson, J. G. and Klausner, R. D. (1995) Overexpression of wild-type and mutant ARF1 and ARF6: distinct perturbations of nonoverlapping membrane compartments. *J. Cell Biol.* **128**, 1003–1017
- 25 Gu, F. and Gruenberg, J. (2000) ARF1 regulates pH-dependent COP functions in the early endocytic pathway. *J. Biol. Chem.* **275**, 8154–8160
- 26 Balch, W. E., Glick, B. S. and Rothman, J. E. (1984) Sequential intermediates in the pathway of intercompartmental transport in a cell-free system. *Cell* **39**, 525–536
- 27 Brew, K., Shaper, J. H., Olsen, K. W., Trayer, I. P. and Hill, R. L. (1975) Cross-linking of the components of lactose synthetase with dimethylpimelimidate. *J. Biol. Chem.* **250**, 1434–1444
- 28 Donaldson, J. G., Kahn, R. A., Lippincott-Schwartz, J. and Klausner, R. D. (1991) Binding of Arf and β -COP to Golgi membranes: possible regulation by a trimeric G protein. *Science* **254**, 1197–1199
- 29 Donaldson, J. G., Cassel, D., Kahn, R. A. and Klausner, R. D. (1992) ADP-ribosylation factor, a small GTP-binding protein, is required for binding of the coatamer protein β -COP to Golgi membranes. *Proc. Natl. Acad. Sci. U.S.A.* **89**, 6408–6412
- 30 Palmer, D. J., Helms, J. B., Beckers, C. J. M., Orci, L. and Rothman, J. E. (1993) Binding of coatamer to Golgi membranes requires ADP-ribosylation factor. *J. Biol. Chem.* **268**, 12083–12089
- 31 Kahn, R. A. (1991) Fluoride is not an activator of the smaller (20–25 kDa) GTP-binding proteins. *J. Biol. Chem.* **266**, 15595–15597
- 32 Northup, J. K., Smigel, M. D., Sternweis, P. C. and Gilman, A. G. (1983) The subunits of the stimulatory regulatory component of adenylate cyclase. Resolution of the activated 45,000-dalton (α) subunit. *J. Biol. Chem.* **258**, 11369–11376
- 33 Okabe, K., Yatani, A., Evans, T., Ho, Y. K., Codina, J., Birnbaumer, L. and Brown, A. M. (1990) $\beta\gamma$ dimers of G proteins inhibit atrial muscarinic K^+ channels. *J. Biol. Chem.* **265**, 12854–12858
- 34 Waters, M. G., Serafini, T. and Rothman, J. E. (1991) Coatamer: a cytosolic protein complex containing subunits of non-clathrin-coated Golgi transport vesicles. *Nature (London)* **349**, 248–251
- 35 Berón, W., Colombo, M. I., Mayorga, L. S. and Stahl, P. D. (1995) In vitro reconstitution of phagosome–endosome fusion: evidence for regulation by heterotrimeric GTPases. *Arch. Biochem. Biophys.* **317**, 337–342
- 36 Bigay, J., Deterre, P., Pfister, C. and Chabre, M. (1985) Fluoroaluminates activate transducin-GDP by mimicking the γ -phosphate of GTP in its binding site. *FEBS Lett.* **191**, 181–185
- 37 Moss, J. and Vaughan, M. (1998) Molecules in the ARF orbit. *J. Biol. Chem.* **273**, 21431–21434
- 38 Togawa, A., Morinaga, N., Ogasawara, M., Moss, J. and Vaughan, M. (1999) Purification and cloning of a brefeldin A-inhibited guanine nucleotide-exchange protein for ADP-ribosylation factors. *J. Biol. Chem.* **274**, 12308–12315
- 39 Yamaji, R., Adamik, R., Takeda, K., Togawa, A., Pacheco-Rodríguez, G., Ferrans, V. J., Moss, J. and Vaughan, M. (2000) Identification and localization of two brefeldin A-inhibited guanine nucleotide-exchange proteins for ADP-ribosylation factors in a macromolecular complex. *Proc. Natl. Acad. Sci. U.S.A.* **97**, 2567–2572
- 40 Moss, J. and Vaughan, M. (1995) Structure and function of ARF proteins: activators of cholera toxin and critical components of intracellular vesicular transport processes. *J. Biol. Chem.* **270**, 12327–12330
- 41 Radhakrishna, H. and Donaldson, J. G. (1997) ADP-ribosylation factor 6 regulates a novel plasma membrane recycling pathway. *J. Cell Biol.* **139**, 49–61
- 42 D'Souza-Schorey, C., van Donselaar, E., Hsu, V. W., Yang, C., Sathl, P. D. and Peters, J. P. (1998) ARF6 targets recycling vesicles to the plasma membrane: insights from an ultrastructural investigation. *J. Cell Biol.* **140**, 603–616
- 43 Zhang, Q., Cox, D., Tseng, C.-C., Donaldson, J. D. and Greenberg, S. (1998) A requirement for ARF6 in Fc γ receptor-mediated phagocytosis in macrophages. *J. Biol. Chem.* **273**, 19977–19981
- 44 Guo, Q., Vasile, E. and Krieger, M. (1994) Disruptions of Golgi structure and membrane traffic in a conditional lethal mammalian cell mutant are corrected by ϵ -COP. *J. Cell Biol.* **125**, 1213–1224
- 45 Hobbie, L., Fisher, A. S., Lee, S., Flint, A. and Krieger, M. (1994) Isolation of three classes of conditional lethal Chinese hamster ovary cell mutants with temperature-dependent defects in low density lipoprotein receptor stability and intracellular membrane transport. *J. Biol. Chem.* **269**, 20958–20970
- 46 Mayorga, L. S., Bertini, F. and Stahl, P. D. (1991) Fusion of newly formed phagosomes with endosomes in intact cells and in a cell-free system. *J. Biol. Chem.* **266**, 6511–6517
- 47 Pitt, A., Mayorga, L. S., Stahl, P. D. and Schwartz, A. L. (1992) Alterations in the protein composition of maturing phagosomes. *J. Clin. Invest.* **90**, 1978–1983
- 48 Rabinowitz, S., Horstmann, H., Gordon, S. and Griffiths, G. (1992) Immunocytochemical characterization of the endocytic and phagolysosomal compartments in peritoneal macrophages. *J. Cell Biol.* **116**, 95–112
- 49 Desjardins, M., Celis, J. E., van Meer, G., Dieplinger, H., Jahraus, A., Griffiths, G. and Huber, L. A. (1994) Molecular characterization of phagosomes. *J. Biol. Chem.* **269**, 32194–32200
- 50 Desjardins, M., Huber, L. A., Parton, R. G. and Griffiths, G. (1994) Biogenesis of phagolysosomes proceeds through a sequential series of interactions with the endocytic apparatus. *J. Cell Biol.* **124**, 677–688
- 51 Berón, W., Alvarez-Dominguez, C., Mayorga, L. S. and Stahl, P. D. (1995) Membrane trafficking along the phagocytic pathway. *Trends Cell Biol.* **5**, 100–104
- 52 Jahraus, A., Tjelle, T. E., Berg, T., Habermann, A., Storrie, O. U. and Griffith, G. (1998) In vitro fusion of phagosomes with different endocytic organelles from J774 macrophages. *J. Biol. Chem.* **273**, 30379–30390
- 53 Muller, W. A., Steinman, R. M. and Cohn, Z. A. (1980) The membrane proteins of the vacuolar system. II. Bidirectional flow between secondary lysosomes and plasma membrane. *J. Cell Biol.* **86**, 304–314
- 54 Guo, Q., Penman, M., Trigatti, B. L. and Krieger, M. (1996) A single point mutation in ϵ -COP results in temperature-sensitive, lethal defects in membrane transport in a Chinese hamster ovary cell mutant. *J. Biol. Chem.* **271**, 11191–11196
- 55 Botelho, R. J., Hackman, D. J., Schreiber, A. D. and Grinstein, S. (2000) Role of COPI in phagosome maturation. *J. Biol. Chem.* **275**, 15717–15727