



Rectangular b-metric space and contraction principles

R. George^{a,*}, S. Radenović^b, K. P. Reshma^c, S. Shukla^d

^aDepartment of Mathematics and Computer Science, St. Thomas College, Bhilai, Chhattisgarh, India.

^bFaculty of Mechanical Engineering, University of Belgrade, Kraljice Marije 16, 11120 Beograd, Serbia.

^cDepartment of Mathematics, Government VYT PG Autonomous College, Durg, Chhattisgarh, India.

^dDepartment of Applied Mathematics, S.V.I.T.S. Indore (M.P.), India.

Abstract

The concept of rectangular b-metric space is introduced as a generalization of metric space, rectangular metric space and b-metric space. An analogue of Banach contraction principle and Kannan's fixed point theorem is proved in this space. Our result generalizes many known results in fixed point theory.

Keywords: Fixed points, b-metric space, rectangular metric space, rectangular b-metric space.

2010 MSC: 47H10.

1. Introduction and Preliminaries

Since the introduction of Banach contraction principle in 1922, because of its wide applications, the study of existence and uniqueness of fixed points of a mapping and common fixed points of two or more mappings has become a subject of great interest. Many authors proved the Banach contraction Principle in various generalized metric spaces. In the sequel Branciari [9] introduced the concept of rectangular metric space (RMS) by replacing the sum on the right hand side of the triangular inequality in the definition of a metric space by a three-term expression and proved an analogue of the Banach Contraction Principle in such space. Since then many fixed point theorems for various contractions on rectangular metric space appeared (see [1],[3],[4],[10],[15],[16],[17],[18],[19],[22],[23],[25],[26]).

On the other hand, in [5] Bakhtin introduced b-metric space as a generalization of metric space and proved analogue of Banach contraction principle in b-metric space. Since then, several papers have dealt

*Corresponding author

Email addresses: renygeorge02@yahoo.com (R. George), radens@beotel.net (S. Radenović), b4reshma@yahoo.com (K. P. Reshma), satishmathematics@yahoo.co.in (S. Shukla)

with fixed point theory or the variational principle for single-valued and multi-valued operators in b-metric spaces (see [2],[6],[7],[8],[11],[12],[13],[14],[20] and the references therein).

In this paper we have introduced the concept of rectangular b-metric space, which is not necessarily Hausdorff and which generalizes the concept of metric space, rectangular metric space and b-metric space. Note that spaces with non Hausdorff topology plays an important role in Tarskian approach to programming language semantics used in computer science (For some details see [24]). Analog of the Banach contraction principle as well as the Kannan type fixed point theorem in rectangular b-metric space are proved. Some examples are included which shows that our generalizations are genuine.

Definition 1.1 ([5]). Let X be a nonempty set and the mapping $d: X \times X \rightarrow [0, \infty)$ satisfies:

(bM1) $d(x, y) = 0$ if and only if $x = y$ for all $x, y \in X$;

(bM2) $d(x, y) = d(y, x)$ for all $x, y \in X$;

(bM3) there exist a real number $s \geq 1$ such that $d(x, y) \leq s[d(x, z) + d(z, y)]$ for all $x, y, z \in X$.

Then d is called a b-metric on X and (X, d) is called a b-metric space (in short bMS) with coefficient s .

Definition 1.2 ([9]). Let X be a nonempty set and the mapping $d: X \times X \rightarrow [0, \infty)$ satisfies:

(RM1) $d(x, y) = 0$ if and only if $x = y$ for all $x, y \in X$;

(RM2) $d(x, y) = d(y, x)$ for all $x, y \in X$;

(RM3) $d(x, y) \leq d(x, u) + d(u, v) + d(v, y)$ for all $x, y \in X$ and all distinct points $u, v \in X \setminus \{x, y\}$.

Then d is called a rectangular metric on X and (X, d) is called a rectangular metric space (in short RMS).

We define a rectangular b-metric space as follows :

Definition 1.3. Let X be a nonempty set and the mapping $d: X \times X \rightarrow [0, \infty)$ satisfies:

(RbM1) $d(x, y) = 0$ if and only if $x = y$;

(RbM2) $d(x, y) = d(y, x)$ for all $x, y \in X$;

(RbM3) there exists a real number $s \geq 1$ such that $d(x, y) \leq s[d(x, u) + d(u, v) + d(v, y)]$ for all $x, y \in X$ and all distinct points $u, v \in X \setminus \{x, y\}$.

Then d is called a rectangular b-metric on X and (X, d) is called a rectangular b-metric space (in short RbMS) with coefficient s .

Note that every metric space is a rectangular metric space and every rectangular metric space is a rectangular b-metric space (with coefficient $s = 1$). However the converse of the above implication is not necessarily true.

Example 1.4. Let $X = \mathbb{N}$, define $d: X \times X \rightarrow X$ by

$$d(x, y) = \begin{cases} 0, & \text{if } x = y; \\ 4\alpha, & \text{if } x, y \in \{1, 2\} \text{ and } x \neq y; \\ \alpha, & \text{if } x \text{ or } y \notin \{1, 2\} \text{ and } x \neq y, \end{cases}$$

where $\alpha > 0$ is a constant. Then (X, d) is a rectangular b-metric space with coefficient $s = \frac{4}{3} > 1$, but (X, d) is not a rectangular metric space, as $d(1, 2) = 4\alpha > 3\alpha = d(1, 3) + d(3, 4) + d(4, 2)$.

Example 1.5. Let $X = \mathbb{N}$, define $d: X \times X \rightarrow X$ such that $d(x, y) = d(y, x)$ for all $x, y \in X$ and

$$d(x, y) = \begin{cases} 0, & \text{if } x = y; \\ 10\alpha, & \text{if } x = 1, y = 2; \\ \alpha, & \text{if } x \in \{1, 2\} \text{ and } y \in \{3\}; \\ 2\alpha, & \text{if } x \in \{1, 2, 3\} \text{ and } y \in \{4\}; \\ 3\alpha, & \text{if } x \text{ or } y \notin \{1, 2, 3, 4\} \text{ and } x \neq y, \end{cases}$$

where $\alpha > 0$ is a constant. Then (X, d) is a rectangular b-metric space with coefficient $s = 2 > 1$, but (X, d) is not a rectangular metric space, as $d(1, 2) = 10\alpha > 5\alpha = d(1, 3) + d(3, 4) + d(4, 2)$.

Note that every b-metric space with coefficient s is a *RbMS* with coefficient s^2 but the converse is not necessarily true. (See Example 1.7 below).

For any $x \in X$ we define the open ball with center x and radius $r > 0$ by

$$B_r(x) = \{y \in X : d(x, y) < r\}$$

The open balls in *RbMS* are not necessarily open (See Example 1.7 below). Let \mathcal{U} be the collection of all subsets \mathcal{A} of X satisfying the condition that for each $x \in \mathcal{A}$ there exist $r > 0$ such that $B_r(x) \subseteq \mathcal{A}$. Then \mathcal{U} defines a topology for the *RbMS* (X, d) , which is not necessarily Hausdorff (See Example 1.7 below).

We define convergence and Cauchy sequence in rectangular b-metric space and completeness of rectangular b-metric space as follows :

Definition 1.6. Let (X, d) be a rectangular b-metric space, $\{x_n\}$ be a sequence in X and $x \in X$. Then

- (a) The sequence $\{x_n\}$ is said to be convergent in (X, d) and converges to x , if for every $\varepsilon > 0$ there exists $n_0 \in \mathbb{N}$ such that $d(x_n, x) < \varepsilon$ for all $n > n_0$ and this fact is represented by $\lim_{n \rightarrow \infty} x_n = x$ or $x_n \rightarrow x$ as $n \rightarrow \infty$.
- (b) The sequence $\{x_n\}$ is said to be Cauchy sequence in (X, d) if for every $\varepsilon > 0$ there exists $n_0 \in \mathbb{N}$ such that $d(x_n, x_{n+p}) < \varepsilon$ for all $n > n_0, p > 0$ or equivalently, if $\lim_{n \rightarrow \infty} d(x_n, x_{n+p}) = 0$ for all $p > 0$.
- (c) (X, d) is said to be a complete rectangular b-metric space if every Cauchy sequence in X converges to some $x \in X$.

Note that, limit of a sequence in a *RbMS* is not necessarily unique and also every convergent sequence in a *RbMS* is not necessarily a Cauchy sequence. The following example illustrates this fact.

Example 1.7. Let $X = A \cup B$, where $A = \{\frac{1}{n} : n \in \mathbb{N}\}$ and B is the set of all positive integers. Define $d: X \times X \rightarrow [0, \infty)$ such that $d(x, y) = d(y, x)$ for all $x, y \in X$ and

$$d(x, y) = \begin{cases} 0, & \text{if } x = y; \\ 2\alpha, & \text{if } x, y \in A; \\ \frac{\alpha}{2n}, & \text{if } x \in A \text{ and } y \in \{2, 3\}; \\ \alpha, & \text{otherwise,} \end{cases}$$

where $\alpha > 0$ is a constant. Then (X, d) is a rectangular b-metric space with coefficient $s = 2 > 1$. However we have the following :

- 1) (X, d) is not a rectangular metric space, as $d(\frac{1}{2}, \frac{1}{3}) = 2\alpha > \frac{17}{12} = d(\frac{1}{2}, 4) + d(4, 3) + d(3, \frac{1}{3})$ and hence not a metric space.
- 2) There does not exist $s > 0$ satisfying $d(x, y) \leq s[d(x, z) + d(z, y)]$ for all $x, y, z \in X$, and so (X, d) is not a b-metric space.
- 3) $B_{\frac{\alpha}{2}}(\frac{1}{2}) = \{2, 3, \frac{1}{2}\}$ and there does not exist any open ball with center 2 and contained in $B_{\frac{\alpha}{2}}(\frac{1}{2})$. So $B_{\frac{\alpha}{2}}(\frac{1}{2})$ is not an open set.
- 4) The sequence $\{\frac{1}{n}\}$ converges to 2 and 3 in *RbMS* and so limit is not unique. Also $d(\frac{1}{n}, \frac{1}{n+p}) = 2\alpha \not\rightarrow 0$ as $n \rightarrow \infty$, therefore $\{\frac{1}{n}\}$ is not a Cauchy sequence in *RbMS*.
- 5) There does not exist any $r_1, r_2 > 0$ such that $B_{r_1}(2) \cap B_{r_2}(3) = \phi$ and so (X, d) is not Hausdorff.

2. Main results

Following theorem is the analogue of Banach contraction principle in rectangular b-metric space.

Theorem 2.1. Let (X, d) be a complete rectangular b-metric space with coefficient $s > 1$ and $T: X \rightarrow X$ be a mapping satisfying:

$$d(Tx, Ty) \leq \lambda d(x, y) \tag{2.1}$$

for all $x, y \in X$, where $\lambda \in [0, \frac{1}{s}]$. Then T has a unique fixed point.

Proof. Let $x_0 \in X$ be arbitrary. Define the sequence $\{x_n\}$ by $x_{n+1} = Tx_n$ for all $n \geq 0$. We shall show that $\{x_n\}$ is Cauchy sequence. If $x_n = x_{n+1}$ then x_n is fixed point of T . So, suppose that $x_n \neq x_{n+1}$ for all $n \geq 0$. Setting $d(x_n, x_{n+1}) = d_n$, it follows from (2.1) that

$$\begin{aligned} d(x_n, x_{n+1}) &= d(Tx_{n-1}, Tx_n) \leq \lambda d(x_{n-1}, x_n) \\ d_n &\leq \lambda d_{n-1}. \end{aligned}$$

Repeating this process we obtain

$$d_n \leq \lambda^n d_0. \tag{2.2}$$

Also, we can assume that x_0 is not a periodic point of T . Indeed, if $x_0 = x_n$ then using (2.2), for any $n \geq 2$, we have

$$\begin{aligned} d(x_0, Tx_0) &= d(x_n, Tx_n) \\ d(x_0, x_1) &= d(x_n, x_{n+1}) \\ d_0 &= d_n \\ d_0 &\leq \lambda^n d_0, \end{aligned}$$

a contradiction. Therefore, we must have $d_0 = 0$, i.e., $x_0 = x_1$, and so x_0 is a fixed point of T . Thus we assume that $x_n \neq x_m$ for all distinct $n, m \in \mathbb{N}$. Again setting $d(x_n, x_{n+2}) = d_n^*$ and using (2.1) for any $n \in \mathbb{N}$, we obtain

$$\begin{aligned} d(x_n, x_{n+2}) &= d(Tx_{n-1}, Tx_{n+1}) \leq \lambda d(x_{n-1}, x_{n+1}) \\ d_n^* &\leq \lambda d_{n-1}^* \end{aligned}$$

Repeating this process we obtain

$$d(x_n, x_{n+2}) \leq \lambda^n d_0^*. \tag{2.3}$$

For the sequence $\{x_n\}$ we consider $d(x_n, x_{n+p})$ in two cases.

If p is odd say $2m + 1$ then using (2.2) we obtain

$$\begin{aligned} d(x_n, x_{n+2m+1}) &\leq s[d(x_n, x_{n+1}) + d(x_{n+1}, x_{n+2}) + d(x_{n+2}, x_{n+2m+1})] \\ &\leq s[d_n + d_{n+1}] + s^2[d(x_{n+2}, x_{n+3}) + d(x_{n+3}, x_{n+4}) \\ &\quad + d(x_{n+4}, x_{n+2m+1})] \\ &\leq s[d_n + d_{n+1}] + s^2[d_{n+2} + d_{n+3}] + s^3[d_{n+4} + d_{n+5}] \\ &\quad + \dots + s^m d_{n+2m} \\ &\leq s[\lambda^n d_0 + \lambda^{n+1} d_0] + s^2[\lambda^{n+2} d_0 + \lambda^{n+3} d_0] + s^3[\lambda^{n+4} d_0 + \lambda^{n+5} d_0] \\ &\quad + \dots + s^m \lambda^{n+2m} d_0 \\ &\leq s\lambda^n [1 + s\lambda^2 + s^2\lambda^4 + \dots] d_0 + s\lambda^{n+1} [1 + s\lambda^2 + s^2\lambda^4 + \dots] d_0 \\ &= \frac{1 + \lambda}{1 - s\lambda^2} s\lambda^n d_0 \quad (\text{note that } s\lambda^2 < 1). \end{aligned}$$

Therefore,

$$d(x_n, x_{n+2m+1}) \leq \frac{1 + \lambda}{1 - s\lambda^2} s\lambda^n d_0. \tag{2.4}$$

If p is even say $2m$ then using (2.2) and (2.3) we obtain

$$\begin{aligned} d(x_n, x_{n+2m}) &\leq s[d(x_n, x_{n+1}) + d(x_{n+1}, x_{n+2}) + d(x_{n+2}, x_{n+2m})] \\ &\leq s[d_n + d_{n+1}] + s^2[d(x_{n+2}, x_{n+3}) + d(x_{n+3}, x_{n+4}) \\ &\quad + d(x_{n+4}, x_{n+2m})] \\ &\leq s[d_n + d_{n+1}] + s^2[d_{n+2} + d_{n+3}] + s^3[d_{n+4} + d_{n+5}] \\ &\quad + \dots + s^{m-1}[d_{2m-4} + d_{2m-3}] + s^{m-1}d(x_{n+2m-2}, x_{n+2m}) \end{aligned}$$

$$\begin{aligned} &\leq s[\lambda^n d_0 + \lambda^{n+1} d_0] + s^2[\lambda^{n+2} d_0 + \lambda^{n+3} d_0] + s^3[\lambda^{n+4} d_0 + \lambda^{n+5} d_0] \\ &\quad + \dots + s^{m-1}[\lambda^{2m-4} d_0 + \lambda^{2m-3} d_0] + s^{m-1} \lambda^{n+2m-2} d_0^* \\ &\leq s\lambda^n [1 + s\lambda^2 + s^2\lambda^4 + \dots] d_0 + s\lambda^{n+1} [1 + s\lambda^2 + s^2\lambda^4 + \dots] d_0 \\ &\quad + s^{m-1} \lambda^{n+2m-2} d_0^*, \end{aligned}$$

i.e.

$$\begin{aligned} d(x_n, x_{n+2m}) &\leq \frac{1 + \lambda}{1 - s\lambda^2} s\lambda^n d_0 + s^{m-1} \lambda^{n+2m-2} d_0^* \\ &< \frac{1 + \lambda}{1 - s\lambda^2} s\lambda^n d_0 + (s\lambda)^{2m} \lambda^{n-2} d_0^* \quad (\text{as } 1 < s) \\ &\leq \frac{1 + \lambda}{1 - s\lambda^2} s\lambda^n d_0 + \lambda^{n-2} d_0^* \quad (\text{as } \lambda \leq \frac{1}{s}). \end{aligned}$$

Therefore

$$d(x_n, x_{n+2m}) \leq \frac{1 + \lambda}{1 - s\lambda^2} s\lambda^n d_0 + \lambda^{n-2} d_0^*. \tag{2.5}$$

It follows from (2.4) and (2.5) that

$$\lim_{n \rightarrow \infty} d(x_n, x_{n+p}) = 0 \quad \text{for all } p > 0. \tag{2.6}$$

Thus $\{x_n\}$ is a Cauchy sequence in X . By completeness of (X, d) there exists $u \in X$ such that

$$\lim_{n \rightarrow \infty} x_n = u. \tag{2.7}$$

We shall show that u is a fixed point of T . Again, for any $n \in \mathbb{N}$ we have

$$\begin{aligned} d(u, Tu) &\leq s[d(u, x_n) + d(x_n, x_{n+1}) + d(x_{n+1}, Tu)] \\ &= s[d(u, x_n) + d_n + d(Tx_n, Tu)] \\ &\leq s[d(u, x_n) + d_n + \lambda d(x_n, u)]. \end{aligned}$$

Using (2.6) and (2.7) it follows from above inequality that $d(u, Tu) = 0$, i.e., $Tu = u$. Thus u is a fixed point of T .

For uniqueness, let v be another fixed point of T . Then it follows from (2.1) that $d(u, v) = d(Tu, Tv) \leq \lambda d(u, v) < d(u, v)$, a contradiction. Therefore, we must have $d(u, v) = 0$, i.e., $u = v$. Thus fixed point is unique. \square

Example 2.2. Let $X = A \cup B$, where $A = \{\frac{1}{n} : n \in \{2, 3, 4, 5\}\}$ and $B = [1, 2]$. Define $d: X \times X \rightarrow [0, \infty)$ such that $d(x, y) = d(y, x)$ for all $x, y \in X$ and

$$\begin{cases} d(\frac{1}{2}, \frac{1}{3}) = d(\frac{1}{4}, \frac{1}{5}) = 0.03 \\ d(\frac{1}{2}, \frac{1}{5}) = d(\frac{1}{3}, \frac{1}{4}) = 0.02 \\ d(\frac{1}{2}, \frac{1}{4}) = d(\frac{1}{5}, \frac{1}{3}) = 0.6 \\ d(x, y) = |x - y|^2 \quad \text{otherwise} \end{cases}$$

Then (X, d) is a rectangular b-metric space with coefficient $s = 4 > 1$. But (X, d) is neither a metric space nor a rectangular metric space. Let $T : X \rightarrow X$ be defined as :

$$Tx = \begin{cases} \frac{1}{4} & \text{if } x \in A \\ \frac{1}{5} & \text{if } x \in B \end{cases}$$

Then T satisfies the condition of Theorem 2.1 and has a unique fixed point $x = \frac{1}{4}$.

Remark 2.3. We say that $T : X \rightarrow X$ has property P if $F(T) = F(T^n)$ (see [21]) where $F(T) = \{x \in X : Tx = x\}$. It is an easy exercise to see that under the assumptions of Theorem 2.1, T has property P.

Following theorem is the analogue of Kannan type contraction in rectangular b-metric space.

Theorem 2.4. *Let (X, d) be a complete rectangular b-metric space with coefficient $s > 1$ and $T : X \rightarrow X$ be a mapping satisfying:*

$$d(Tx, Ty) \leq \lambda[d(x, Tx) + d(y, Ty)] \tag{2.8}$$

for all $x, y \in X$, where $\lambda \in [0, \frac{1}{s+1}]$. Then T has a unique fixed point.

Proof. Let $x_0 \in X$ be arbitrary. We define a sequence $\{x_n\}$ by $x_{n+1} = Tx_n$ for all $n \geq 0$. We shall show that $\{x_n\}$ is Cauchy sequence. If $x_n = x_{n+1}$ then x_n is fixed point of T . So, suppose that $x_n \neq x_{n+1}$ for all $n \geq 0$. Setting $d(x_n, x_{n+1}) = d_n$, it follows from (2.8) that

$$\begin{aligned} d(x_n, x_{n+1}) &= d(Tx_{n-1}, Tx_n) \leq \lambda[d(x_{n-1}, Tx_{n-1}) + d(x_n, Tx_n)] \\ d(x_n, x_{n+1}) &= \lambda[d(x_{n-1}, x_n) + d(x_n, x_{n+1})] \\ d_n &= \lambda[d_{n-1} + d_n] \\ d_n &\leq \frac{\lambda}{1-\lambda}d_{n-1} = \beta d_{n-1}, \end{aligned}$$

where $\beta = \frac{\lambda}{1-\lambda} < \frac{1}{s}$ (as, $\lambda < \frac{1}{s+1}$). Repeating this process we obtain

$$d_n \leq \beta^n d_0. \tag{2.9}$$

Also, we can assume that x_0 is not a periodic point of T . Indeed, if $x_0 = x_n$ then using (2.9), for any $n \geq 2$, we have

$$\begin{aligned} d(x_0, Tx_0) &= d(x_n, Tx_n) \\ d(x_0, x_1) &= d(x_n, x_{n+1}) \\ d_0 &= d_n \\ d_0 &\leq \beta^n d_0, \end{aligned}$$

a contradiction. Therefore, we must have $d_0 = 0$, i.e., $x_0 = x_1$, and so x_0 is a fixed point of T . Thus we assume that $x_n \neq x_m$ for all distinct $n, m \in \mathbb{N}$. Again using (2.8) and (2.9) for any $n \in \mathbb{N}$, we obtain

$$\begin{aligned} d(x_n, x_{n+2}) &= d(Tx_{n-1}, Tx_{n+1}) \leq \lambda[d(x_{n-1}, Tx_{n-1}) + d(x_{n+1}, Tx_{n+1})] \\ &= \lambda[d(x_{n-1}, x_n) + d(x_{n+1}, x_{n+2})] = \lambda[d_{n-1} + d_{n+1}] \\ &\leq \lambda[\beta^{n-1}d_0 + \beta^{n+1}d_0] \\ &= \lambda\beta^{n-1}[1 + \beta^2]d_0 \\ &= \gamma\beta^{n-1}d_0. \end{aligned}$$

Therefore,

$$d(x_n, x_{n+2}) \leq \gamma\beta^{n-1}d_0, \tag{2.10}$$

where $\gamma = \lambda[1 + \beta^2] > 0$.

For the sequence $\{x_n\}$ we consider $d(x_n, x_{n+p})$ in two cases.

If p is odd say $2m + 1$ then using (2.9) we obtain

$$\begin{aligned} d(x_n, x_{n+2m+1}) &\leq s[d(x_n, x_{n+1}) + d(x_{n+1}, x_{n+2}) + d(x_{n+2}, x_{n+2m+1})] \\ &\leq s[d_n + d_{n+1}] + s^2[d(x_{n+2}, x_{n+3}) + d(x_{n+3}, x_{n+4}) \\ &\quad + d(x_{n+4}, x_{n+2m+1})] \\ &\leq s[d_n + d_{n+1}] + s^2[d_{n+2} + d_{n+3}] + s^3[d_{n+4} + d_{n+5}] \\ &\quad + \dots + s^m d_{n+2m} \end{aligned}$$

$$\begin{aligned} &\leq s[\beta^n d_0 + \beta^{n+1} d_0] + s^2[\beta^{n+2} d_0 + \beta^{n+3} d_0] + s^3[\beta^{n+4} d_0 + \beta^{n+5} d_0] \\ &\quad + \dots + s^m \beta^{n+2m} d_0 \\ &\leq s\beta^n [1 + s\beta^2 + s^2\beta^4 + \dots] d_0 + s\beta^{n+1} [1 + s\beta^2 + s^2\beta^4 + \dots] d_0 \\ &= \frac{1 + \beta}{1 - s\beta^2} s\beta^n d_0 \quad (\text{note that } s\beta^2 < 1). \end{aligned}$$

Therefore,

$$d(x_n, x_{n+2m+1}) \leq \frac{1 + \beta}{1 - s\beta^2} s\beta^n d_0. \tag{2.11}$$

If p is even say $2m$ then using (2.9) and (2.10) we obtain

$$\begin{aligned} d(x_n, x_{n+2m}) &\leq s[d(x_n, x_{n+1}) + d(x_{n+1}, x_{n+2}) + d(x_{n+2}, x_{n+2m})] \\ &\leq s[d_n + d_{n+1}] + s^2[d(x_{n+2}, x_{n+3}) + d(x_{n+3}, x_{n+4}) \\ &\quad + d(x_{n+4}, x_{n+2m})] \\ &\leq s[d_n + d_{n+1}] + s^2[d_{n+2} + d_{n+3}] + s^3[d_{n+4} + d_{n+5}] \\ &\quad + \dots + s^{m-1}[d_{2m-4} + d_{2m-3}] + s^{m-1}d(x_{n+2m-2}, x_{n+2m}) \\ &\leq s[\beta^n d_0 + \beta^{n+1} d_0] + s^2[\beta^{n+2} d_0 + \beta^{n+3} d_0] + s^3[\beta^{n+4} d_0 + \beta^{n+5} d_0] \\ &\quad + \dots + s^{m-1}[\beta^{2m-4} d_0 + \beta^{2m-3} d_0] + s^{m-1}\gamma\beta^{n+2m-3} d_0 \\ &\leq s\beta^n [1 + s\beta^2 + s^2\beta^4 + \dots] d_0 + s\beta^{n+1} [1 + s\beta^2 + s^2\beta^4 + \dots] d_0 \\ &\quad + s^{m-1}\gamma\beta^{n+2m-3} d_0, \end{aligned}$$

i.e.

$$\begin{aligned} d(x_n, x_{n+2m}) &\leq \frac{1 + \beta}{1 - s\beta^2} s\beta^n d_0 + s^{m-1}\gamma\beta^{n+2m-3} d_0 \\ &< \frac{1 + \beta}{1 - s\beta^2} s\beta^n d_0 + \gamma(s\beta)^{2m} \beta^{n-3} d_0 \quad (\text{as } 1 < s) \\ &\leq \frac{1 + \beta}{1 - s\beta^2} s\beta^n d_0 + \gamma\beta^{n-3} d_0 \quad (\text{as } \beta \leq \frac{1}{s}). \end{aligned}$$

Therefore

$$d(x_n, x_{n+2m}) \leq \frac{1 + \beta}{1 - s\beta^2} s\beta^n d_0 + \gamma\beta^{n-3} d_0. \tag{2.12}$$

It follows from (2.11) and (2.12) that

$$\lim_{n \rightarrow \infty} d(x_n, x_{n+p}) = 0 \quad \text{for all } p > 0. \tag{2.13}$$

Thus $\{x_n\}$ is a Cauchy sequence in X . By completeness of (X, d) there exists $u \in X$ such that

$$\lim_{n \rightarrow \infty} x_n = u. \tag{2.14}$$

We shall show that u is a fixed point of T . Again, for any $n \in \mathbb{N}$ we have

$$\begin{aligned} d(u, Tu) &\leq s[d(u, x_n) + d(x_n, x_{n+1}) + d(x_{n+1}, Tu)] \\ &= s[d(u, x_n) + d_n + d(Tx_n, Tu)] \\ &\leq s[d(u, x_n) + d_n + \lambda\{d(x_n, Tx_n) + d(u, Tu)\}] \\ &= s[d(u, x_n) + d_n + \lambda\{d(x_n, x_{n+1}) + d(u, Tu)\}] \\ (1 - s\lambda)d(u, Tu) &\leq s[d(u, x_n) + \beta^n d_0 + \lambda d(x_n, x_{n+1})] \end{aligned}$$

Using (2.13) and (2.14) and the fact that $\lambda < \frac{1}{s+1}$, it follows from above inequality that $d(u, Tu) = 0$, i.e., $Tu = u$. Thus u is a fixed point of T .

For uniqueness, let v be another fixed point of T . Then it follows from (2.8) that $d(u, v) = d(Tu, Tv) \leq \lambda[d(u, Tu) + d(v, Tv)] = \lambda[d(u, u) + d(v, v)] = 0$. Therefore, we have $d(u, v) = 0$, i.e., $u = v$. Thus fixed point is unique. \square

Remark 2.5. On the basis of discussion contained in this paper, we have the following:

- 1) The open ball defined in b-metric space, RMS and RbMS are not necessarily open set.
- 2) The collection of open balls in RbMS, RMS and b-metric space do not necessarily form a basis for a topology.
- 3) RbMS, RMS and b-metric space are not necessarily Hausdorff.

Open Problems :

- 1) In Theorem 2.1, can we extend the range of λ to the case $\frac{1}{s} < \lambda < 1$.
- 2) Prove analogue of Chatterjee contraction, Reich contraction, Ciric contraction and Hardy-Rogers contraction in *RbMS*.

Acknowledgements

The authors are thankful to the learned referees for the valuable suggestions provided, which helped them in bringing this paper in its present form.

References

- [1] T. Abdeljawad, D. Turkoglu, *Locally convex valued rectangular metric spaces and Kannan's fixed point theorem*, arXiv, **2011** (2011), 11 pages.1
- [2] H. Aydi, M. F. Bota, E. Karapinar, S. Moradi, *A common fixed point for weak ϕ -contractions on b-metric spaces*, Fixed Point Theory, **13** (2012), 337–346.1
- [3] A. Azam, M. Arshad, *Kannan Fixed Point Theorems on generalised metric spaces*, J. Nonlinear Sci. Appl., **1** (2008), 45–48.1
- [4] A. Azam, M. Arshad, I. Beg, *Banach contraction principle on cone rectangular metric spaces*, Appl. Anal. Discrete Math., **3** (2009), 236–241.1
- [5] I. A. Bakhtin, *The contraction mapping principle in quasimetric spaces*, Funct. Anal., Unianowsk Gos. Ped. Inst., **30** (1989), 26–37.1, 1.1
- [6] M. Boriceanu, *Strict fixed point theorems for multivalued operators in b-metric spaces*, Inter. J. Mod. Math., **4** (2009), 285–301.1
- [7] M. Boriceanu, M. Bota, A. Petrusel, *Multivalued fractals in b-metric spaces*, Cen. Eur. J. Math., **8** (2010), 367–377.1
- [8] M. Bota, A. Molnar, V. Csaba, *On Ekeland's variational principle in b-metric spaces*, Fixed Point Theory, **12** (2011), 21–28.1
- [9] A. Branciari, *A fixed point theorem of Banach-Caccippoli type on a class of generalised metric spaces*, Publ. Math. Debrecen, **57** (2000), 31–37.1, 1.2
- [10] C. N. Chen, *Common fixed point theorem in complete generalized metric spaces*, J. Appl. Math., **2012** (2012), 14 pages.1
- [11] S. Czerwik, *Contraction mappings in b-metric spaces*, Acta. Math. Inform. Univ. Ostraviensis, **1** (1993), 5–11.1
- [12] S. Czerwik, *Nonlinear set-valued contraction mappings in b-metric spaces*, Atti Sem. Mat. Univ. Modena, **46** (1998), 263–276.1
- [13] S. Czerwik, K. Dlutek, S. L. Singh, *Round-off stability of iteration procedures for operators in b-metric spaces*, J. Natur. Phys. Sci., **11** (1997), 87–94.1
- [14] S. Czerwik, K. Dlutek, S. L. Singh, *Round-off stability of iteration procedures for set valued operators in b-metric spaces*, J. Natur. Phys. Sci., **15**(2001), 1–8.1
- [15] P. Das, *A fixed point theorem on a class of generalized metric spaces*, Korean J. Math. Sci., **9** (2002), 29–33.1
- [16] P. Das, *A fixed point theorem in generalized metric spaces*, Soochow J. Math., **33** (2007), 33–39.1
- [17] P. Das, B. K. Lahri, *Fixed point of a Ljubomir Ciric's quasi-contraction mapping in a generalized metric space*, Publ. Math. Debrecen, **61**(2002), 589–594.1
- [18] P. Das, B. K. Lahri, *Fixed Point of contractive mappings in generalised metric space*, Math. Slovaca, **59** (2009), 499–504.1

- [19] I. M. Erhan, E. Karapinar, T. Sekulic, *Fixed Points of (ψ , ϕ) contractions on generalised metric spaces*, Fixed Point Theory Appl., **2012** (2012), 12 pages.1
- [20] R. George, B. Fisher, *Some generalised results of fixed points in cone b-metric spaces*, Math. Moravic., **17** (2013), 39–50.1
- [21] G. S. Jeong, B. E. Rhoades, *Maps for which $F(T) = F(T^n)$* , Fixed Point Theory Appl., **6** (2007), 71–105.2.3
- [22] M. Jleli, B. Samet, *The Kannan's fixed point theorem in cone rectangular metric space*, J. Nonlinear Sci. Appl., **2** (2009), 161–167.1
- [23] H. Lakzian, B. Samet, *Fixed Points for (ψ, ϕ) -weakly contractive mapping in generalised metric spaces*, Appl. Math. Lett., **25** (2012), 902–906.1
- [24] S. G. Mathews, *Partial Metric Topology*, Papers on general topology appl., Ann. New York Acad. Sci., **728** (1994), 183–197.1
- [25] D. Mihet, *On Kannan fixed point result in generalised metric spaces*, J. Nonlinear Sci. Appl., **2** (2009), 92–96.1
- [26] I. R. Sarma, J. M. Rao, S. S. Rao, *Contractions over generalised metric spaces*, J. Nonlinear Sci. Appl., **2** (2009), 180–182.1