

RECURSIVE LABELLING SYSTEMS AND STABILITY OF RECURSIVE STRUCTURES IN HYPERARITHMETICAL DEGREES

C. J. ASH

ABSTRACT. We show that, under certain assumptions of recursiveness in \mathfrak{A} , the recursive structure \mathfrak{A} is Δ_α^0 -stable for $\alpha < \omega_1^{CK}$ if and only if there is an enumeration of \mathfrak{A} using a Σ_α^0 set of recursive Σ_α infinitary formulae and finitely many parameters from \mathfrak{A} . This extends the results of [1].

To do this, we first obtain results concerning Δ_α^0 paths in recursive labelling systems, also extending results of [1]. We show, more generally, that a path and a labelling can simultaneously be defined, when each node of the path is to be obtained by a Δ_α^0 function from the previous node and its label.

Introduction. We say that a recursive structure \mathfrak{A} is Δ_α^0 -stable if, for every recursive structure $\mathfrak{B} \cong \mathfrak{A}$, every isomorphism from \mathfrak{B} to \mathfrak{A} is Δ_α^0 in Kleene's hyperarithmetical hierarchy. We shall show that, under certain assumptions, \mathfrak{A} is Δ_α^0 -stable iff there exists a "formally Δ_α^0 enumeration of \mathfrak{A} ."

The basic outline of our argument is as in [1], where such a result was obtained for finite α . However, certain technicalities prevent the generalization to infinite α from being as straightforward as might be expected. The argument seems most easily described in terms of recursive labelling systems, which were introduced in [1] for a similar purpose.

The two basic results for α -systems are obtained in §1 and applied in the succeeding sections to the question of Δ_α^0 -stability. These basic results are also used in the related topic of Δ_α^0 -categoricity [2].

In §1 we define the notion of a recursively α -guided recursive labelling system, or α -system, and in Proposition 1 state the desired result similar to that of [1] involving the existence of r.e. points of $2^{\mathbb{N}}$ and of labellings of Δ_α^0 paths in an α -system. To obtain this result for $\alpha \geq \omega$, we need to deal with limit ordinals, and we frame an analogous result [Proposition 2] for $\langle \gamma_n \rangle$ -systems where $\langle \gamma_n \rangle$ is an increasing sequence of ordinals. Propositions 1 and 2 are then established by the Main Lemma.

In §2 we show how the results of §1 apply to showing that structures are not Δ_α^0 -stable, which is the "difficult direction" of our result. We show in Theorem 1 that under suitable conditions on \mathfrak{A} there is a recursive structure \mathfrak{B} and an isomorphism from \mathfrak{B} to \mathfrak{A} which is not Δ_α^0 . In the case where α is a limit ordinal, we also give in Theorem 2 conditions under which there exists such a \mathfrak{B} and an isomorphism which is not Δ_β^0 for any $\beta < \alpha$.

Received by the editors June 18, 1984 and, in revised form September 19, 1985.
1980 *Mathematics Subject Classification* (1985 Revision). Primary 03D30.

©1986 American Mathematical Society
0002-9947/86 \$1.00 + \$.25 per page

The method of proof for Theorems 1 and 2 is a little simpler than that given in [1]. We comment that the method of [1] allows the assumptions of Theorems 1 and 2 to be slightly weakened.

In §3 we relate the results of §2 to recursive infinitary formulae. Theorem 4 shows that, under the conditions of Theorem 1, a recursive structure \mathfrak{A} is Δ_α^0 -stable iff \mathfrak{A} has a formally Δ_α^0 enumeration. Theorem 5 gives a similar result for the notion of $\tilde{\Delta}_\alpha^0$ -stability introduced for Theorem 2.

Theorem 4 can also be expressed in terms of the Scott rank of a structure. For a suitable notion of rank, we observe in Theorem 6 that a recursive structure \mathfrak{A} satisfying the assumptions of Theorem 4 is Δ_α^0 -stable iff \mathfrak{A} has $< 2^{\aleph_0}$ automorphisms and, for some finite \bar{p} , (\mathfrak{A}, \bar{p}) has rank at most α . Lastly in §3, we define \mathfrak{A} to be *Hyp-stable* if for each recursive $\mathfrak{B} \cong \mathfrak{A}$, each isomorphism is hyperarithmetical. We note in Theorem 7 that we do not need methods such as those of §1 to deal with this notion because a recursive structure \mathfrak{A} will be Hyp-stable iff it has fewer than 2^{\aleph_0} automorphisms. It follows that, without further assumptions, a recursive structure \mathfrak{A} is Hyp-stable iff it is Δ_α^0 -stable for some $\alpha < \omega_1^{CK}$.

Our main interest in this paper is the theoretical problem of finding general conditions under which a structure is not Δ_α^0 -stable. The examples of conventional algebraic structures which we have considered do not provide examples of Δ_α^0 -stability for $\alpha \geq \omega$. However, the notions of §§2 and 3 may be illustrated by considering structures based on the ordinal numbers, such as $(\alpha, <)$ and $(\alpha, <, +)$. In §4 we summarize some of the results concerning the stability of such structures.

More details about recursion theory are to be found in [6] and about recursive structures in [1]. We shall use Kleene’s system \mathcal{O} of ordinal notations and the related notions $<_0, +_0$, and $| \cdot |_0$ [or just $| \cdot |$] which are described in §11.7 of [6]. The Δ_α^0 functions are as defined in §16.8 of [6].

We use the phrase “recursive transfinite induction” to indicate Kleene’s method for defining a partial recursive function on \mathcal{O} . This consists of taking a partial recursive function f and using the Recursion Theorem to obtain an e for which $\phi_e(a) = f(e, a)$. The definition of $f(n, a)$ normally involves the restriction $\phi_{k(n,a)}$ of ϕ_n to $\{b: b <_0 a\}$. If this definition guarantees that $f(n, a)$ is defined whenever $\phi_{k(n,a)}(b)$ is defined for all $b <_0 a$, then it follows by classical transfinite induction on $|a|$ that $\phi_e(a)$ is defined for every $a \in \mathcal{O}$.

1. Recursive labelling systems. For applications in this paper, we are interested only in the metric space $2^{\mathbb{N}}$. But without much trouble we can obtain the results of this section for any suitably recursive complete metric space and so we shall do so.

DEFINITION. A *recursive metric space* consists of a metric space X , a family $B(X)$ of nonempty open subsets of X , forming a basis for X , and an enumeration of $B(X)$ with respect to which the relations $\sigma < \tau$ [the closure of σ is a subset of τ] and $\delta(\sigma) < 1/n$ [the diameter of σ is less than $1/n$] are r.e. relations on $B(X) \times B(X)$ and $B(X) \times \mathbb{N}$, respectively.

DEFINITION. An r.e. point of a recursive metric space X is an element x of X such that $\{\sigma \in B(X): x \in \sigma\}$ is r.e.

An easily verified equivalent condition is that $\{x\} = \bigcap \sigma_n$ for some recursive sequence $\langle \sigma_n \rangle$ from $B(X)$ such that $\delta(\sigma_n) \rightarrow 0$ as $n \rightarrow \infty$.

We note that, for the space $2^{\mathbf{N}}$ with its usual basis, the r.e. points are the recursive subsets of \mathbf{N} , not the r.e. subsets. This will be the case whenever $B(X)$ consists of clopen sets and the relation $\sigma \cap \tau = \emptyset$ on $B(X)$ is r.e., because then $x \notin \sigma$ iff, for some $\tau \in B(X)$, $x \in \tau$ and $\sigma \cap \tau = \emptyset$.

By a complete recursive metric space, we mean a recursive metric space which is also complete in the classical sense, without further recursive assumptions. We need to use only the fact that if $\langle \sigma_n \rangle$ is a recursive sequence from $B(X)$, $\delta(\sigma_n) \rightarrow 0$ and for each n , $\sigma_{n+1} < \sigma_n$, then $\bigcap \sigma_n = \{x\}$ where x is an r.e. point. This follows since $x \in \sigma$ iff there exists n for which $\sigma_n < \sigma$.

When discussing trees, we need to refer frequently to the levels of the nodes, so we give our definitions in terms of levels and of the predecessor function.

DEFINITION. A tree T consists of disjoint sets $\{a\}, T_0, T_1, \dots$ and a predecessor function p for which $p: T_{n+1} \rightarrow T_n$, $p: T_0 \rightarrow \{a\}$, and, say, $p(a) = a$. If $p(x) = y$ then y is the predecessor of x and x is a successor of y . The set of nodes of T is the set $\{a\} \cup (\bigcup_n T_n)$. The apex of T is the node a , and the nodes of T of level n are the elements of T_n .

[Thus no level is assigned to the element a . In fact, with our present conventions, the element a plays no essential part in any construction and could be omitted.]

A recursive tree is one for which the set of nodes forms an r.e. set of natural numbers and the function p is partial recursive. In this case, each set T_n will be r.e., uniformly in n .

Since the function p completely specifies T , we may define an index for a recursive tree to be an index for the partial recursive function p .

A path in T is a sequence u_0, u_1, \dots [finite or infinite] for which $u_n \in T_n$ and $p(u_{n+1}) = u_n$.

A recursive labelling system on a recursive metric space X is a quintuple (T, L, S, N, F) for which T is a recursive tree, L is an r.e. set, S is an r.e. subset of $T \times L$, N is an r.e. subset of $T \times L \times T \times L$, and F is a function from L to subsets of X for which the relation $F(l) \cap \sigma \neq \emptyset$ on $L \times B(X)$ is r.e.

A labelling of the path u_0, u_1, \dots in T is a sequence l_0, l_1, \dots from L of the same length such that, for each n , $S(u_n, l_n)$ and $N(u_n, l_n, u_{n+1}, l_{n+1})$. An adherent point of a labelling $\langle l_n \rangle$ of an infinite path $\langle u_n \rangle$ is a point $x \in X$ such that, for every open set U for which $x \in U$, there exists N such that for all $n > N$, $F(l_n) \cap U \neq \emptyset$.

[Usually, for a correct labelling, we shall have $F(l_0) \supseteq F(l_1) \supseteq \dots$, in which case $x \in X$ is adherent iff $x \in \bigcap_n \overline{F(l_n)}$. This will be the case for the α -systems about to be considered whenever $\alpha > 1$. The above definition of an adherent point removes, to some extent, the need to treat the case where $\alpha = 1$ as a special case.]

We proceed to find sufficient conditions under which a nonrecursive path in T nevertheless has a [nonrecursive] labelling with an r.e. adherent point. For some of our applications, however, such a result, generalizing that of [1], is not sufficient. We need also to consider situations where the desired path is determined at each level, not just by the previous node, but also by its label. We therefore make the following definitions.

DEFINITIONS. A path-generating function [or instruction] in a recursive labelling system is a function from $T \times L$ to T mapping each (u, l) for which $S(u, l)$ holds to some successor of u in T .

A labelling of an instruction, p , is an infinite path u_0, u_1, \dots in T together with a labelling l_0, l_1, \dots of this path such that, for each n , $p(u_n, l_n) = u_{n+1}$.

We shall show that the existence of suitable relations on L provides a sufficient condition for every Δ_α^0 instruction to have a Δ_α^0 labelling with an r.e. adherent point.

DEFINITION. Let $\alpha < \omega_1^{CK}$ and let $\alpha = \beta + 1$. A recursively α -guided recursive labelling system [or α -system] is a recursive labelling system (T, L, S, N, F) together with a notation $a \in \mathcal{O}$ for α and a family $\{\triangleleft_\gamma : 1 \leq \gamma < \alpha\}$ of uniformly r.e. binary relations on L , indexed by $\{c : c <_0 a\}$, for which conditions (1) to (7) below are satisfied. [Here $<_0$ denotes the partial ordering on \mathcal{O} , so that $\{c : c <_0 a\}$ is an r.e. linearly ordered set of order type α . Or course, we do not use in this instance the notation for 0.]

(1) For each $u \in T_0$ and each $\sigma \in B(X)$ there exists $l \in L$ such that $S(u, l)$ and $F(l) \cap \sigma \neq \emptyset$.

(2) If $S(u, l)$ and $N(u, l, v, m)$, then $S(v, m)$.

(3) Each \triangleleft_γ is reflexive and transitive.

(4) If $N(u, l, v, m)$, then $l \triangleleft_\beta m$.

(5) If $1 \leq \gamma_1 < \gamma_2 < \alpha$ and $l \triangleleft_{\gamma_2} m$, then $l \triangleleft_{\gamma_1} m$.

(6) If $l \triangleleft_1 m$, then $F(l) \supseteq F(m)$.

(7) Suppose that $S(u, l)$, $F(l) \cap \sigma \neq \emptyset$, and v is a successor of u . [Suppose also that $\alpha > \alpha_k > \dots > \alpha_1 > \alpha_0 \geq 1$ and that $l = l_k \triangleleft_{\alpha_k} \dots \triangleleft_{\alpha_2} l_1 \triangleleft_{\alpha_1} l_0$ where $F(l_0) \cap \sigma \neq \emptyset$.]* Then there exists m for which $N(u, l, v, m)$ and $F(m) \cap \sigma \neq \emptyset$ [and $l_i \triangleleft_{\alpha_i} m$ for $i = 0, 1, \dots, k$]*.

*Notes. The two parts of condition (7) in parentheses are to be omitted in the cases where $\alpha = 1$ or $\alpha = 2$. In the case where $\alpha = 1$, the sequence of \triangleleft_γ 's is to be regarded as the empty sequence and only conditions (1), (2), and (7) are relevant. In the case where $\alpha = 2$, we have $\beta = 1$ and we need only the single relation \triangleleft_1 , so condition (5) is irrelevant.

The following result will be established by the Main Lemma.

PROPOSITION 1. If $1 \leq \alpha < \omega_1^{CK}$, α is a successor ordinal and \mathcal{T} is an α -system over a complete recursive metric space, then for every Δ_α^0 instruction in \mathcal{T} there is a Δ_α^0 labelling having an r.e. adherent point.

In order to prove this, we introduce similar notions for the case where α is a limit ordinal. In this case, we restrict our attention to certain particular classes of Δ_α^0 instructions.

DEFINITION. A special sequence for a limit ordinal α is an increasing sequence $\langle \gamma_n \rangle$ of successor ordinals whose limit is α and for which there is a recursive sequence $\langle c_n \rangle$ of notations from \mathcal{O} such that $c_0 <_0 c_1 <_0 c_2 <_0 \dots$.

In such a case, there will be a notation $a \in \mathcal{O}$ for α such that each $c_n <_0 a$, namely $a = 3.5^e$ where e is an index for the recursive sequence $\langle c_n \rangle$. Conversely, if $\alpha < \omega_1^{CK}$ is any limit ordinal, then one may always obtain a special sequence $\langle \gamma_n \rangle$ for α , for example by taking any notation 3.5^e for α and letting $\gamma_n = |\phi_e(n) +_0 1|$.

DEFINITIONS. Let α be a limit ordinal and let $\langle \gamma_n \rangle$ be a special sequence for α .

A special Δ_α^0 -instruction w.r.t. $\langle \gamma_n \rangle$ [or $\langle \gamma_n \rangle$ -instruction] in a recursive labelling system is an instruction p such that, for some recursive sequence $\langle e_n \rangle$, each e_n is a $\Delta_{\gamma_n}^0$ index for the restriction of p to $\{(u, l) : u \in T_n\}$.

A special α -system w.r.t. $\langle \gamma_n \rangle$ [or $\langle \gamma_n \rangle$ -system] is a recursive labelling system together with, first, a notation $a \in \mathcal{O}$ for α , second a recursive sequence $\langle c_n \rangle$ of notations for the $\langle \gamma_n \rangle$, where $c_0 <_0 c_1 <_0 \dots <_0 a$, and third, a family $\{\triangleleft_\gamma : 1 \leq \gamma < \alpha\}$ of uniformly r.e. binary relations on L indexed by $\{c : c <_0 a\}$ which satisfy conditions (1), (2), (3), (5) and (6) above and conditions (4)' and (7)' below.

(4)' If $N(u, l, v, m)$ where $u \in T_n$ and if $\gamma_n = \beta_n + 1$ where $\beta_n > 0$, then $l \triangleleft_{\beta_n} m$.

(7)' Suppose that $S(u, l), \sigma \in B(X)$ and that v is a successor of u where $u \in T_n$. Suppose also that $\gamma_n > \alpha_k > \dots > \alpha_1 > \alpha_0 \geq 1$ and that $l = l_k \triangleleft_{\alpha_k} \dots \triangleleft_{\alpha_2} l_1 \triangleleft_{\alpha_1} l_0$ where $F(l_0) \cap \sigma \neq \emptyset$. Then there exists m for which $N(u, l, v, m), F(m) \cap \sigma \neq \emptyset$ and $l_i \triangleleft_{\alpha_i} m$ for $i = 0, 1, \dots, (k - 1)$.

For a $\langle \gamma_n \rangle$ -instruction in a $\langle \gamma_n \rangle$ -system, we may well expect a corresponding special sort of Δ_α^0 labelling.

DEFINITION. A $\langle \gamma_n \rangle$ -labelling of an instruction p in a recursive labelling system is a labelling $\langle u_n \rangle, \langle l_n \rangle$ of p such that, for some recursive sequence $\langle e_n \rangle$, each e_n is a $\Delta_{\gamma_n}^0$ index for the pair (u_{n+1}, l_{n+1}) .

The following result will also be established by the Main Lemma.

PROPOSITION 2. Let $\langle \gamma_n \rangle$ be a special sequence for a limit ordinal α , where $\alpha < \omega_1^{CK}$, and let \mathcal{T} be a $\langle \gamma_n \rangle$ -system over a complete recursive metric space. Then for every $\langle \gamma_n \rangle$ -instruction in \mathcal{T} there is a $\langle \gamma_n \rangle$ -labelling having an r.e. adherent point.

COMMENTS ON THE MAIN LEMMA. Before proceeding to the statement and proof of the Main Lemma, we comment that a more intuitive line of argument to establish Proposition 1 [and similarly Proposition 2] would be as follows.

Given an α -system \mathcal{T}_α , we would construct a uniformly recursive sequence $\langle \mathcal{T}_\gamma : 1 \leq \gamma \leq \alpha \rangle$ for which each \mathcal{T}_γ is a γ -system in such a way that for each Δ_α^0 instruction p_α in \mathcal{T}_α we could find a sequence $\langle p_\gamma : 1 \leq \gamma \leq \alpha \rangle$ where each p_γ is a Δ_γ^0 instruction in \mathcal{T}_γ . These instructions would be arranged so that each Δ_γ^0 labelling of p_γ would determine a $\Delta_{\gamma+1}^0$ labelling of $p_{\gamma+1}$ having the same adherent points.

The \mathcal{T}_γ 's for which γ is a limit ordinal would be special $\langle \gamma_n \rangle$ -systems w.r.t. some special sequence $\langle \gamma_n \rangle$ for γ . Then p_γ would be a $\langle \gamma_n \rangle$ -instruction and a labelling of p_γ would be obtained by referring, at the $(n + 1)$ th level of \mathcal{T}_γ , to the labelling of p_{γ_n} in \mathcal{T}_{γ_n} .

The path p_1 in \mathcal{T}_1 would first be labelled to have an r.e. adherent point and then, proceeding by recursive transfinite induction, a Δ_γ^0 labelling with the same adherent point would be obtained for each p_γ , in particular for p_α .

A complication which arises in following precisely this outline, when α is infinite, is that each \mathcal{T}_γ must be defined in terms of later \mathcal{T}_γ 's and eventually in terms of \mathcal{T}_α . Thus, the definition of the \mathcal{T}_γ and similarly the p_γ is not by recursive transfinite induction on γ . Rather, the systems \mathcal{T}_γ need not be defined simultaneously one level at a time. Thus the nodes of \mathcal{T}_γ of level n and associated parts of \mathcal{T}_γ must be defined by recursive transfinite induction on some ordinal $\rho(\gamma, n)$. The function ρ depends on some choice of a suitably nested family of special sequences for the limit ordinals $\gamma \leq \alpha$.

So it seems preferable, for the sake of examining the details carefully, to consider a slight generalization which can be proved more straightforwardly by recursive

transfinite induction on α . We no longer need to construct the sequence of systems \mathcal{T}_β explicitly, although this remains the intuitive guide and could be constructed by the same induction.

The generalization which we prove is that for each α -system \mathcal{T}_α and each $\beta < \alpha$ one can obtain a β -system \mathcal{T}_β for which the problem of labelling Δ_α^0 instructions in \mathcal{T}_α is reducible to that of labelling Δ_β^0 instructions in \mathcal{T}_β . We make this precise:

DEFINITION. Let \mathcal{T}_α be an α -system and let $\beta < \alpha$. A β -precursor of \mathcal{T}_α consists of:

- (A) a β -system \mathcal{T}_β ,
- (B) a partial recursive function which assigns to each index for a Δ_α^0 -instruction p_α in \mathcal{T}_α an index for a Δ_β^0 -instruction p_β ,
- (C) a partial recursive function which for each p_β obtained from p_α as in (B) assigns to each index a Δ_β^0 labelling of p_β and index for a Δ_α^0 labelling of p_α having at least the same adherent points.

Note. In the case where α [or similarly β] is a limit ordinal, it is understood here that by a Δ_α^0 -instruction or Δ_α^0 -labelling we mean a $\langle \gamma_n \rangle$ -instruction or $\langle \gamma_n \rangle$ -labelling, where $\langle \gamma_n \rangle$ is the associated special sequence.

PROOF OF PROPOSITIONS 1 AND 2. If \mathcal{T}_1 is a 1-precursor of an α -system \mathcal{T}_α and p_α is a Δ_α^0 instruction in \mathcal{T}_α , then the corresponding recursive instruction p_1 in \mathcal{T}_1 may easily be recursively labelled, as in Lemma 17 of [1], to have an r.e. adherent point. Thus, Propositions 1 and 2 follow from the existence of precursors, to be proved in our Main Lemma.

MAIN LEMMA. If \mathcal{T}_α is an α -system, $\beta < \alpha$ and $\beta \leq \gamma_0$, in the case where α is a limit ordinal with special sequence $\langle \gamma_n \rangle$, then \mathcal{T}_α has a β -precursor.

PROOF.

1. Method of proof. We define, by recursive transfinite induction on $a \in \mathcal{O}$, a recursive function which we call the precursor operator which assigns to each index for an α -system using the notation a for α and to each $b \in \mathcal{O}$ for which $b <_0 a$ [and $b \leq_0 c_0$ in the case where α is a limit ordinal and $\langle c_n \rangle$ is the sequence of notations for the special sequence] an index for a β -precursor, where $\beta = |b|$.

Thus it is sufficient to describe how to obtain a β -precursor \mathcal{T}_β of an α -system \mathcal{T}_α in terms of given precursors of γ -systems \mathcal{T}_γ for $\gamma < \alpha$.

We need to carry various properties of our precursors through the transfinite induction. The labels of the precursor of an α -system and the associated definitions will be obtained from the corresponding notions in the α -system. More generally

DEFINITIONS. An α -set is a system $\mathcal{L} = (L, F, \triangleleft_\gamma)_{\gamma < \alpha}$ satisfying conditions (3), (5), and (6) of the definition of an α -system.

A determining function for a β -system $\mathcal{T}_\beta = (T_\beta, L_\beta, S_\beta, N_\beta, F_\beta, \triangleleft_\gamma^\beta)_{\gamma < \beta}$ is a function $g: L_\beta \rightarrow L$ for some α -set $\mathcal{L} = (L, F, \triangleleft_\gamma)_{\gamma < \alpha}$, where $\alpha \geq \beta$ such that, for all $l \in L_\beta$, $F_\beta(l) = F(g(l))$ and, for all $l, m \in L_\beta$, $l \triangleleft_\gamma^\beta m$ iff $g(l) \triangleleft_\gamma g(m)$.

We now add to the desired ingredients of a precursor, the inductive assumption that our precursor operation also yields:

- (D) a determining function $g_{\alpha\beta}$ from \mathcal{T}_β to the α -set of \mathcal{T}_α .

We also need to use the fact that our construction of a precursor of \mathcal{T}_α is independent of those labels of \mathcal{T}_α which are suitable for no nodes of \mathcal{T}_α .

DEFINITION. Let g be a determining function from an α -system \mathcal{T}_α to an α -set \mathcal{L} , with the notation as above. We say that \mathcal{T}_α is an α -system w.r.t. g if it satisfies the following strengthening of condition (7).

(7)* Suppose that $S_\alpha(u, l)$, $F_\alpha(l) \cap \sigma = \emptyset$, and v is a successor of u . [Suppose also that $\alpha > \alpha_k > \dots > \alpha_0 \geq 1$, that $g(l) = l_k \triangleleft_{\alpha_k} \dots \triangleleft_{\alpha_1} l_0$, where $F(l_0) \cap \sigma \neq \emptyset$ and each $l_i \in L$ rather than L_α]. Then there exists $m \in L_\alpha$ for which $N_\alpha(u, l, v, m)$ and $F_\alpha(m) \cap \sigma \neq \emptyset$ [and $l_i \triangleleft_{\alpha_i} g(m)$ for $i = 0, 1, \dots, k$].

In the case where α is a limit ordinal, condition (7)' is modified in the same way.

We add to our inductive assumptions:

(E) If g is any determining function for \mathcal{T}_α such that \mathcal{T}_α is an α -system w.r.t. g , then \mathcal{T}_β is a β -system w.r.t. the composite $g \circ g_{\alpha\beta}$.

We need to use the fact that our precursors and the associated operations depend, at each level of the tree, only on this level and preceding levels of the corresponding features of the original α -system.

Notation. For each γ -system \mathcal{T} , let \mathcal{T}^k denote the system consisting of the nodes of \mathcal{T} of levels $\leq k$, the labels suitable for these nodes, and the restrictions to these nodes of S, N and $\langle \triangleleft_\xi \rangle_{\xi < \gamma}$.

We add to our inductive assumptions:

(F) For α -systems \mathcal{T}_α and their β -precursors \mathcal{T}_β , each \mathcal{T}_β^k depends only on \mathcal{T}_α^k .

(G) Similarly, the restriction of the function $g_{\alpha\beta}$, in (D), to \mathcal{T}_β^k depends only on \mathcal{T}_α^k .

(H) Likewise, the restriction of p_β , in (C), to \mathcal{T}_β^k depends only on the restriction of p_α to \mathcal{T}_α^k .

COMMENT. So, in principle, the precursor operator acts on partial systems \mathcal{T}_α^k to yield, in (A), the partial system \mathcal{T}_β^k , in (D) a determining function from the labels of \mathcal{T}_β^k to those of \mathcal{T}_α^k and, in (B) a partial recursive function acting on restriction of instructions p_α to \mathcal{T}_α^k . However, no such requirement is necessary [or indeed possible] for (C).

We add further to our inductive assumptions two simplifications:

(I) The level 0 nodes and their labels are exactly the same for \mathcal{T}_β as for \mathcal{T}_α . The function $g_{\alpha\beta}$ is the identity function for such labels.

(J) The labelling of p_α given in (C) always has the same level 0 node with the same label as in the labelling of p_β .

Finally, we note that if $\gamma < \beta < \alpha$, \mathcal{T}_β is a β -precursor of \mathcal{T}_α and \mathcal{T}_γ is the γ -precursor of \mathcal{T}_β , then \mathcal{T}_γ is a suitable choice for a γ -precursor of \mathcal{T}_α , the associated operations being obtained by composition. It is therefore sufficient to describe how to obtain a β -precursor of an α -system \mathcal{T}_α first when $\alpha = \beta + 1$ [for which we consider the two cases where β is or is not a limit ordinal] and then when α is a limit ordinal and $b = c_0$, where $\langle c_n \rangle$ is the associated sequence of notations. We now do this in parts 2, 3, and 4 below.

In the case where $\alpha = \beta + 1$ and β is a limit ordinal, in order to obtain a γ -precursor for \mathcal{T}_α where $\gamma < \beta$, we must obtain a γ -precursor for the β -precursor of \mathcal{T}_α . Thus, the special sequence $\langle \gamma_n \rangle$ for β must have $\gamma \leq \gamma_0$. We therefore show in this case that any special sequence for β can be used. [This is why, in our definitions, we did not restrict attention only to the fundamental sequence given by the notation for β .]

2. The case $\alpha = \beta + 1$ where β is a successor ordinal. Let $\mathcal{T}_\alpha = \mathcal{T} = (T, L, S, N, F, \triangleleft_\gamma)_{\gamma < \alpha}$. If $\alpha = \beta + 1$, we let the β -precursor of \mathcal{T} be the system $\mathcal{T}' = (T', L', S', N', F', \triangleleft'_\gamma)_{\gamma < \beta}$ defined as follows.

The tree T' consists of all the sequences (u_0, u_1, \dots, u_n) from T for which each u_k has level k in T and each u_{k+1} is a finite successor of u_0 . The predecessor of (u_0, u_1, \dots, u_n) is $(u_0, u_1, \dots, u_{n-1})$.

If $u = (u_0, u_1, \dots, u_n)$ and the path in T with last node u_n is r_0, r_1, \dots, r_n [so $r_n = u_n$], then we define $S'(u, l)$ to hold when $l = (l_0, l_1, \dots, l_n)$ is a labelling in \mathcal{T} of this path. The set L' consists of all such sequences l . The determining function g for \mathcal{T}' is defined by $g(l) = l_n$, and F' and the \triangleleft'_γ are defined according to the definition of a determining function. When all else is done, we may replace (u_0) and (l_0) by u_0 and l_0 for the sake of assumptions (I) and (J).

For $u, v \in T', l, m \in L'$, we define $N'(u, l, v, m)$ as follows. Let $u = (u_0, u_1, \dots, u_n)$. We require that $v = (u_0, u_1, \dots, u_n, u_{n+1})$ for some u_{n+1} and that $S'(u, l)$ and $S'(v, m)$. Thus $l = (l_0, l_1, \dots, l_n)$ and $m = (m_0, m_1, \dots, m_n, m_{n+1})$ are labellings in \mathcal{T} of paths r_0, \dots, r_n and $r_0, \dots, r_{a-1}, s_1, \dots, s_{n+1}$, where $r_n = u_n$ and either $s_a \neq r_a$ or $a = n + 1$. Then we let $N'(u, l, v, m)$ hold if, additionally, $m_0 = l_0, m_1 = l_1, \dots, m_{a-1} = l_{a-1}$, and $l_n \triangleleft'_\gamma m_{n+1}$ where $\beta = \gamma + 1$. We note that, here, $a \geq 1$, by the definition of T' .

Conditions (1) to (6) for a β -system are straightforward. To see that \mathcal{T}' is a β -system w.r.t. g also satisfying assumption (E), suppose that \mathcal{T} is an α -system w.r.t. some determining function h from \mathcal{T} to an α -set $(L^+, F^+, \triangleleft'_\gamma)_{\gamma < \alpha}$. We must verify condition (7)* for \mathcal{T}' and $h \circ g$.

Suppose, then, that v is a successor of u in $T', l \in L', S'(u, l), \beta > \alpha_k > \dots > \alpha_0 \geq 1$, and that $h(g(l)) = l_k^+ \triangleleft_{\alpha_k}^+ \dots \triangleleft_{\alpha_1}^+ l_0^+$, where each $l_i^+ \in L^+$ and $F^+(l_0^+) \cap \sigma \neq \emptyset$. If $\beta = \gamma + 1$, then we may assume that $\alpha_k = \gamma$, since $l_k^+ \triangleleft_\gamma^+ l_k^+$. Using the notation above, we must find a labelling $m = (l_0, \dots, l_{a-1}, m_a, \dots, m_{n+1})$ of the path $r_0, \dots, r_{a-1}, s_a, \dots, s_{n+1}$ in \mathcal{T} .

Since $l = (l_0, \dots, l_n)$ is a labelling of a path in the α -system \mathcal{T} , we have $l_{a-1} \triangleleft_\beta l_n$ and so $h(l_{a-1}) \triangleleft_\beta^+ h(l_n) = l_k^+ \triangleleft_{\alpha_k}^+ l_{k-1}^+ \dots \triangleleft_{\alpha_1}^+ l_0^+$. Now by condition (7)* for \mathcal{T} and h , we may find $m_a \in L$ for which $N(r_{a-1}, l_{a-1}, s_a, m_a), F(m_a) \cap \sigma \neq \emptyset, h(l_{a-1}) \triangleleft_\beta^+ h(m_a)$, and $l_i^+ \triangleleft_{\alpha_i} h(m_a)$ for $0 \leq i \leq k$. We may then choose the remaining m_i in turn using (7) for \mathcal{T} , so that $F(m_{n+1}) \cap \sigma \neq \emptyset$. Then, since $h(l_n) = l_k^+ \triangleleft_{\alpha_k}^+ h(m_a)$ and $\alpha_k = \gamma$, we have $l_n \triangleleft_\gamma m_a \triangleleft_\beta m_{n+1}$, so $l_n \triangleleft_\gamma m_{n+1}$ and $N'(u, l, v, m)$. For $0 \leq i \leq k$, since $l_i^+ \triangleleft_{\alpha_i}^+ h(m_a)$ and $m_a \triangleleft_\beta m_{n+1}$, we have $h(m_a) \triangleleft_\beta^+ h(m_{n+1})$ and so $l_i^+ \triangleleft_{\alpha_i}^+ h(m_{n+1}) = h(g(m))$, as required.

For condition (B), given any Δ_α^0 instruction p in \mathcal{T} , we obtain a corresponding Δ_β^0 instruction p' in \mathcal{T}' as follows. Since $\alpha = \beta + 1$, we may let $p(u, l) = \lim_s \hat{p}(u, l, s)$ where \hat{p} is a Δ_β^0 function. For $u' \in T', l' \in L'$ such that $S'(u', l')$, let $u' = (u_0, \dots, u_n)$ and $l' = (l_0, \dots, l_n)$. Then l' is a labelling of the path r_0, \dots, r_n in \mathcal{T} , where $r_n = u_n$. We consider the least i for which $i = n$ or $\hat{p}(r_i, l_i, n) \neq r_{i+1}$, and let $p'(u', l')$ be $(u_0, \dots, u_n, u_{n+1})$ where u_{n+1} is any level $(n + 1)$ node of T which is a finite successor of $\hat{p}(r_i, l_i, n)$.

Then, for condition (C), suppose that $\langle u'_n \rangle$ and $\langle l'_n \rangle$ form a labelling of p' in \mathcal{T}' . Let $u'_n = (u_{n0}, u_{n1}, \dots, u_{nn})$, $l'_n = (l_{n0}, l_{n1}, \dots, l_{nn})$, and let $r_{n0}, r_{n1}, \dots, r_{nn}$ be the path in \mathcal{T} for which $r_{nn} = u_{nn}$. We may then see, by induction on n , that

for each n the limits $u_n = \lim_s r_{sn}$ and $l_n = \lim_s l_{sn}$ exist and that $\langle u_n \rangle$ and $\langle l_n \rangle$ constitute a labelling of p in \mathcal{T} . If x is an adherent point of the labelling $\langle l'_n \rangle$ and U is an open set where $x \in U$, there exists N such that $F'(l'_n) \cap U \neq \emptyset$ for all $n > N$. Then for $s \geq n > N$ we have $l_{sn} \triangleleft_\beta l_{ss}$, so $F(l_{sn}) \supseteq F(l_{ss}) = F'(l'_s)$ and so, for $n > N$, we have $F(l_n) \cap U \neq \emptyset$. Thus x is also an adherent point of $\langle l_n \rangle$.

3. *The case $\alpha = \beta + 1$ where β is a limit ordinal.* Let $\langle \gamma_n \rangle$ be any special squence for β . Then we may obtain a $\langle \gamma_n \rangle$ -system as a β -precursor of $\mathcal{T}_\alpha = \mathcal{T}$ in very much the same way as in the previous case. The only modifications needed are the following.

In the definition of $N'(u, l, v, m)$, the stipulation $l_n \triangleleft_\gamma m_{n+1}$ where $\beta = \gamma + 1$ is replaced by $l_n \triangleleft_{\beta_n} m_{n+1}$ where $\gamma_n = \beta_n + 1$. In the verification that \mathcal{T}' is a $\langle \gamma_n \rangle$ -system satisfying condition (E), β and γ are replaced by the appropriate γ_n and β_n , respectively.

To obtain a $\langle \gamma_n \rangle$ -instruction p' for \mathcal{T}' corresponding to the instruction p for \mathcal{T} , we first obtain a recursive sequence $\langle e_n \rangle$ where e_n is an index for a $\Delta^0_{\gamma_n}$ function $p_n(u, l)$ and such that $p(u, l) = \lim_n p_n(u, l)$. The instruction p' is now obtained in the same way as before except that $\hat{p}(u, l, n)$ is replaced by $p_n(u, l)$.

4. *The case where α is a limit ordinal.* Let

$$\mathcal{T}_\alpha = \mathcal{T} = (T, L, S, N, F, \langle \gamma \rangle)_{\gamma < \alpha}$$

be an α -system and let $\langle \gamma_n \rangle$ be the associated special sequence for α . We must construct a γ_0 -precursor of \mathcal{T} . We do this by constructing a sequence $\mathcal{T}_0, \mathcal{T}_1, \dots$, where each \mathcal{T}_n is a γ_n -system. Then \mathcal{T}_0 will be the desired γ_0 -precursor of \mathcal{T} .

The \mathcal{T}_n are defined simultaneously as follows. The level 0 nodes of \mathcal{T}_n are exactly the level n nodes of \mathcal{T} , with the same suitable labels. The level $k + 1$ nodes of \mathcal{T}_n and their suitable labels are those of level k of the γ_n -precursor \mathcal{T}'_{n+1} of \mathcal{T}_{n+1} . By assumption (I), the level 0 nodes of \mathcal{T}'_{n+1} are those of \mathcal{T}_{n+1} and so are level $n + 1$ nodes of \mathcal{T} . Thus we may define the predecessor in \mathcal{T}_n of each level 1 node to be as in \mathcal{T} . For nodes of \mathcal{T}_n of levels greater than 1, the predecessors are as in \mathcal{T}'_{n+1} .

In this way, each \mathcal{T}_n is defined in terms of \mathcal{T} and \mathcal{T}_{n+1} . However, the definition is justified since each \mathcal{T}_n^{k+1} depends only on \mathcal{T}^n and $(\mathcal{T}'_{n+1})^k$ and, by assumption (F), $(\mathcal{T}'_{n+1})^k$ depends only on \mathcal{T}_{n+1}^k . Thus the \mathcal{T}_n^k are well defined by induction on k . [Moreover, each \mathcal{T}_n^k depends ultimately only on \mathcal{T}^{k+n} , so \mathcal{T}_0^k depends only on \mathcal{T}^k , as required by condition (F) for a precursor.] The other ingredients of our construction are defined by the same form of induction.

Each \mathcal{T}_n will have a determining function, g_n , to the α -set of \mathcal{T} . The set, L_n , of labels for \mathcal{T}_n consists of the labels of \mathcal{T} suitable for level n nodes of \mathcal{T} , on which g_n is the identity function, together with the labels of \mathcal{T}'_{n+1} on which g_n is $g_{n+1} \circ h_n$, where h_n is the determining function from \mathcal{T}'_{n+1} to \mathcal{T}_{n+1} .

The relations $S_n(u, l)$ and $N_n(u, l, v, m)$ for \mathcal{T}_n are defined to be as in \mathcal{T} if u has level 0 and as in \mathcal{T}'_{n+1} otherwise. The function F and the relations $\langle \triangleleft_\gamma \rangle_{\gamma < \gamma_n}$ are defined in the unique way which makes g_n a determining function for \mathcal{T}_n .

To show that \mathcal{T}_0 is a γ_0 -system w.r.t. g_0 and satisfies condition (E), we suppose that \mathcal{T} is an α -system w.r.t. a determining function h to an α -set $\mathcal{L}^+ = (L^+, F^+, \triangleleft_\gamma^+)_{\gamma < \alpha}$ and prove simultaneously that each \mathcal{T}_n is a γ_n -system w.r.t. $h \circ g_n$.

The arguments are, in principle, by induction on the largest level of labels or nodes involved in each condition for a γ_n -system. The effect of this is that we

may show that \mathcal{T}_n is a γ_n -system w.r.t. $h \circ g_n$ under the assumption that \mathcal{T}_{n+1} is a γ_{n+1} -system w.r.t. $h \circ g_{n+1}$. Conditions (1) to (6) follow immediately from the definitions.

For condition (7)* for \mathcal{T}_n and $h \circ g_n$, suppose that $S_n(u, l), F_n(l) \cap \sigma \neq \emptyset$, and v is a successor of u in \mathcal{T}_n . Suppose also that $\alpha > \alpha_k > \dots > \alpha_0 \geq 1$ and that $h(g_n(l)) = l_k \triangleleft_{\alpha_k}^+ \dots \triangleleft_{\alpha_1}^+ l_0$ where $F^+(l_0) \cap \sigma \neq \emptyset$ and each $l_i \in L^+$.

First, if u has level 0, then $u, v \in T, g_n(l) = l$, and, by (7)* for \mathcal{T} and h , there exists $m \in L$ such that $N(u, l, v, m), F(m) \cap \sigma \neq \emptyset$ and, for $0 \leq i \leq k, l_i \triangleleft_{\alpha_i}^+ h(m)$. But then, by our definitions, $m \in L_n, N_n(u, l, v, m), F_n(m) \cap \sigma = F(g_n(m)) \cap \sigma = F(m) \cap \sigma \neq \emptyset$ and, for $0 \leq i \leq k, l_i \triangleleft_{\alpha_i}^+ h(m) = h(g_n(m))$.

Now suppose that u has level greater than 0. Then u, v are nodes of $\mathcal{T}'_{n+1} = (T'_{n+1}, L'_{n+1}, S'_{n+1}, N'_{n+1}, F'_{n+1}, \triangleleft_{\gamma_n}^m)_{\gamma < \gamma_n}$. By our assumption, \mathcal{T}_{n+1} is a γ_{n+1} -system w.r.t. $h \circ g_{n+1}$, so by the transfinite induction hypothesis \mathcal{T}'_{n+1} is a γ_n -system w.r.t. $h \circ g_{n+1} \circ h_n$. Thus there exists $m \in L'_{n+1}$ such that $N'_{n+1}(u, l, v, m), F'_{n+1}(m) \cap \sigma \neq \emptyset$ and, for $1 \leq i \leq k, l_i \triangleleft_{\alpha_i}^+ h(g_{n+1}(h_n(m))) = h(g_n(m))$.

For (B), suppose that p is a (γ_n) -instruction for \mathcal{T} . We obtain a corresponding $\Delta_{\gamma_0}^0$ instruction p_0 for \mathcal{T}_0 by simultaneously defining for each n a $\Delta_{\gamma_n}^0$ instruction p_n for \mathcal{T}_n . On level 0 nodes of \mathcal{T}_n, p_n acts as p does on level n nodes of \mathcal{T} . On other nodes of \mathcal{T}_n , we let p_n act as does p'_{n+1} , where p'_{n+1} is the instruction for \mathcal{T}'_{n+1} assigned to the instruction p_{n+1} for \mathcal{T}_{n+1} according to (B) of the inductive hypothesis.

For (C), suppose that λ_0 is a $\Delta_{\gamma_0}^0$ labelling of p_0 in \mathcal{T}_0 . We obtain by straightforward induction on n a $\Delta_{\gamma_n}^0$ labelling λ_n of each p_n having the same adherent points. Having obtained λ_n , we obtain λ_{n+1} as follows. We note that λ_n consists of a level 0 node, u_n , of \mathcal{T}_n and a label, l_n , for this, followed by a $\Delta_{\gamma_n}^0$ labelling λ'_{n+1} of p'_{n+1} in \mathcal{T}'_{n+1} . Part (C) of the inductive hypothesis for \mathcal{T}_{n+1} thus yields a labelling λ_{n+1} of p_{n+1} . The sequence of first nodes, u_n , of the λ_n and their labels, l_n , then form [using assumption (J) for the \mathcal{T}_{n+1}] a labelling λ of p in \mathcal{T} .

If x is an adherent point of λ_0 , then it is so for each λ_n , by (C) of the inductive hypothesis, and so certainly for $n \geq 1$ we have $x \in \bigcap \overline{F_n(l)}$ for each label l occurring in λ_n . In particular, $x \in \bigcap \overline{F_n(l_n)}$ for $n \geq 1$, so x is an adherent point of λ .

This concludes the proof.

2. Conditions for instability. Our constructions depend on the “back-and-forth” relations \leq_β between finite sequences, of the same length, of elements of a structure \mathfrak{A} , where β is an ordinal number.

DEFINITION. For each structure \mathfrak{A} we define \leq_β by transfinite induction on $\beta \geq 1$. We define $\bar{a} \leq_1 \bar{b}$ if each finitary universal formula true for \bar{a} in \mathfrak{A} is true for \bar{b} . We define $\bar{a} \leq_{\beta+1} \bar{b}$ if, for each sequence \bar{d} there exists a sequence \bar{c} such that $\bar{a}, \bar{c} \geq_\beta \bar{b}, \bar{d}$. If δ is a limit ordinal, we define $\bar{a} \leq_\beta \bar{b}$ if $\bar{a} \leq_\beta \bar{b}$ for all $\beta < \delta$.

DEFINITION. For each structure \mathfrak{A} , each finite sequence \bar{p} from \mathfrak{A} and each ordinal $\alpha \geq 2$, we define the subset $cl_\alpha(\bar{p})$ of A as follows. If $\alpha = \beta + 1$, then $x \in cl_\alpha(\bar{p})$ if for some \bar{a} , whenever $\bar{p}, x, \bar{a} \leq_\beta \bar{p}, x', \bar{a}'$, then $x = x'$. If δ is a limit ordinal then $cl_\delta(\bar{p}) = \bigcap_{\beta < \delta} cl_\beta(\bar{p})$.

In all the simple examples where \mathfrak{A} is Δ_α^0 -stable we have $A = cl_\alpha(\bar{p})$ for some \bar{p} . We can prove this in general, provided that we make certain additional assumptions about the recursiveness of \mathfrak{A} .

THEOREM 1. *Let \mathfrak{A} be a recursive structure such that for no \bar{p} is $A = \text{cl}_\alpha(\bar{p})$. Then \mathfrak{A} is not Δ_α^0 -stable whenever the following conditions are satisfied.*

- (1) *The existential diagram of \mathfrak{A} is recursive.*
- (2) *For some notation a for α , the relations \leq_β on \mathfrak{A} for $1 \leq \beta < \alpha$ are uniformly r.e. when indexed by $\{b : b <_0 a\}$.*
- (3) *There is a recursive procedure which chooses, for each \bar{p} , an element $x \in A$ for which $x \notin \text{cl}_\alpha(\bar{p})$.*

PROOF. If the domain, A , of \mathfrak{A} is finite, then clearly $A = \text{cl}_\alpha(\bar{p})$ where $A = \{\bar{p}\}$. So we may suppose that $A = \{a_0, a_1, \dots\}$ and let $B = \{b_0, b_1, \dots\}$ be any recursive set. Let L be the language for \mathfrak{A} and let $\{\theta_0, \theta_1, \dots\}$ be an enumeration of the atomic sentences of $L(B)$. We obtain a recursive structure \mathfrak{B} whose domain is B by enumerating, during our construction, the atomic diagram $D(\mathfrak{B})$ of \mathfrak{B} , or, in the terminology of §1, by obtaining an r.e. point x of the metric space $X = 2^{\mathbb{N}}$ with the usual basis as $B(X)$, which yields $D(\mathfrak{B})$ by

$$\begin{cases} \theta_k \in D(\mathfrak{B}) & \text{if } x(k) = 1, \\ -\theta_k \in D(\mathfrak{B}) & \text{if } x(k) = 0. \end{cases}$$

Thus, the nonempty basic open sets $\sigma \in B(X)$ correspond to consistent finite subsets $\Sigma(\sigma)$ of $\{\theta_k : k \in \mathbb{N}\} \cup \{-\theta_k : k \in \mathbb{N}\}$.

Let P be the set of all finite partial one-one functions from B to A . We define $f \in P$ to be *coherent w.r.t. $\sigma \in B(X)$* if there is a bijection $g \supseteq f$ from B to A such that, for each $\theta(b_{i_1}, \dots, b_{i_k}) \in \Sigma(\sigma)$ we have $\mathfrak{A} \models \theta[g(b_{i_1}), \dots, g(b_{i_k})]$. It follows from assumption (1) that this is a recursive relation between f and σ .

The relation \triangleleft_β needed to apply the results of §1 are obtained from similar relations on P defined as follows.

DEFINITION. For $f, g \in P$ and $\beta \geq 1$, let $f \leq_\beta g$ if $\text{dom}(f) \subseteq \text{dom}(g)$ and $a_1, \dots, a_n \leq_\beta b_1, \dots, b_n$, where $\{a_1, \dots, a_n\} = \text{ran}(f)$ and each $b_i = g(f^{-1}(a_i))$.

We may now prove the following [similarly to Lemmas 3, 4, 5, 6, 7 of [1] and using the definition of cl_α].

LEMMA 1. *If $f \in P$ is coherent w.r.t. $\sigma \in B(X)$ then for each $\bar{a} \in A, \bar{b} \in B$, and for each $n = 1, 2, \dots$ there exists $g \in P$ and $\sigma' \in B(X)$ for which $g \supseteq f, \bar{a} \in \text{ran}(g), \bar{b} \in \text{dom}(g), \delta(\sigma') < 1/n$, and g is coherent w.r.t. σ' .*

LEMMA 2. *If $f \triangleleft_\beta g$ and g is coherent w.r.t. σ , then f is coherent w.r.t. σ .*

LEMMA 3. *If $f \leq_\alpha g, g$ is coherent w.r.t. $\sigma, g(y) = x$, and $x \notin \text{cl}_\alpha(\text{ran}(f))$, then for each $\beta < \alpha$ there exists a coherent $h \supseteq f$ for which $g \triangleleft_\beta h$ and $h(y) \neq x$.*

LEMMA 4. *If $\alpha_k > \dots > \alpha_1 > \alpha_0 \geq 1$ and $f_k \triangleleft_{\alpha_k} f_{k-1} \triangleleft_{\alpha_{k-1}} \dots \triangleleft_{\alpha_2} f_1 \triangleleft_{\alpha_1} f_0$ where f_0 is coherent w.r.t. $\sigma \in B(X)$, then there exists $g \supseteq f_k$, coherent w.r.t. σ , such that $f_i \triangleleft_{\alpha_i} g$ for $i = 1, 2, \dots, (k - 1)$.*

To resume the proof, we define an $(\alpha + 1)$ -system as follows. The nodes of the tree of level n consist of sequences of the form $(\mu_0(m_0) = k_0, \dots, \mu_{n-1}(m_{n-1}) = k_{n-1})$ where the μ_e are symbols corresponding to the partial functions ϕ_e^α for some notation a for α . The predecessor function is the usual one for sequences, that is, the function which deletes the last term of a sequence. In this tree there is, as well as the apex, a single node of level 0, namely the empty sequence. A suitable

label for such a node is a pair (f, x) for which $f \in P$, $x \in A$, and $x \notin \text{cl}_\alpha(\text{ran}(f))$. A correct next label on the successor node $(\mu_0(m_0) = k_0, \dots, \mu_n(m_n) = k_n)$ is a suitable label (g, y) such that $g \subseteq f$, $a_n, x \in \text{ran}(g)$, $b_n \in \text{dom}(g)$ and if $m_n = x$, then $g^{-1}(x) \neq k_n$. For each label (f, x) , $F(f, x)$ is defined to be the set of points of 2^N which correspond to structures \mathfrak{B} for which there is an isomorphism from \mathfrak{B} to \mathfrak{A} extending f . This means that $F(f, x) \cap \sigma \neq \emptyset$ iff f is coherent w.r.t. σ .

To show that this is an $\alpha+1$ -system, we define $(f, x) \triangleleft_\beta (g, y)$ if $f \triangleleft_\beta g$, in the case where $\beta < \alpha$, and $(f, x) \triangleleft_\alpha (g, y)$ if $f \subseteq g$. Clause (7) of the definition then requires that, given $\alpha \geq \alpha_k > \dots > \alpha_1 > \alpha_0 \geq 1$, $f_k \triangleleft_{\alpha_k} \dots \triangleleft_{\alpha_2} f_1 \triangleleft_{\alpha_1} f_0$, $\sigma \in B(X)$, and $x \notin \text{cl}_\alpha(\text{ran}(f_k))$, where f_0 is coherent w.r.t. σ , there exists $g \supseteq f_k$ for which g is coherent w.r.t. σ , $x, a_n \in \text{ran}(g)$, $b_n \in \text{dom}(g)$, $f_i \triangleleft_{\alpha_i} g$ for $i = 0, \dots, (k-1)$ and $g^{-1}(x) \neq k$. By Lemma 4 there exists a coherent h for which $h \supseteq f_k$ and $f_i \triangleleft_{\alpha_i} h$ for $i = 0, \dots, (k-1)$. By Lemma 1 we may choose a coherent $h_1 \supseteq h$ for which $x \in \text{ran}(h)$, say $x = h(y)$. By Lemma 3 there exists a coherent $h_2 \supseteq f_k$ for which $h_1 \triangleleft_{\alpha_{k-1}} h_2$ [since $\alpha_{k-1} < \alpha$] and $h_2(y) \neq x$. By Lemma 1 again, we may choose a coherent $g \supseteq h_2$ for which $a_n, x \in \text{ran}(g)$ and $b_n \in \text{dom}(g)$. So $f_k \subseteq h_2 \subseteq g$ and thus $f_k \subseteq g$, and for $i < k$, $f_i \triangleleft_{\alpha_i} h \subseteq h_1 \triangleleft_{\alpha_{k-1}} h_2 \subseteq g$ and thus $f_i \triangleleft_{\alpha_i} g$.

The desired instruction is that which proceeds from a node of level n with label (f, x) to the successor node $(\dots, \mu_n(x) = k)$ for which

$$k = \begin{cases} \phi_n^\alpha(x) & \text{if this is defined,} \\ 0 & \text{(say) otherwise.} \end{cases}$$

Clearly this is a $\Delta_{\alpha+1}^0$ instruction and so, by Proposition 1, there is a correct labelling $(g, x_0), (f_0, x_1), (f_1, x_2), \dots$ of this instruction having an r.e. adherent point. Let \mathfrak{B} be the structure determined by this point and let $f = \bigcup_n f_n$. Then $f: \mathfrak{B} \cong \mathfrak{A}$ and for each e , if ϕ_e^α is total then $f_e^{-1}(x_e) \neq \phi_e^\alpha(x_e)$. Hence f is not Δ_α^0 .

A logical extension of the notion of stability arises from considering a recursive structure \mathfrak{A} which is Δ_α^0 -stable, where α is a limit ordinal, but for each $\beta < \alpha$ is not Δ_β^0 -stable. Then, for each $\beta < \alpha$ there exists $\mathfrak{B}_\beta \cong \mathfrak{A}$ such that no isomorphism from \mathfrak{B}_β to \mathfrak{A} is Δ_β^0 . We ask whether a single such \mathfrak{B} can be chosen independently of β .

DEFINITION. A recursive structure \mathfrak{A} is $\hat{\Delta}_\alpha^0$ -stable, where α is a limit ordinal, if for each recursive $\mathfrak{B} \cong \mathfrak{A}$ and each isomorphism $f: \mathfrak{B} \cong \mathfrak{A}$ there exists $\beta < \alpha$ for which $f \in \Delta_\beta^0$. [Thus $\hat{\Delta}_\alpha^0$ represents the class $\bigcup_{\beta < \alpha} \Delta_\beta^0$.]

THEOREM 2. Suppose that α is a limit ordinal and that, for each $\beta < \alpha$ and each $\bar{p} \in A$, $\text{cl}_\beta(\bar{p}) \neq A$. Then \mathfrak{A} is not $\hat{\Delta}_\alpha^0$ -stable provided that the following conditions are satisfied.

- (1) The existential diagram of \mathfrak{A} is recursive.
- (2) For some notation a for α , the relations \leq_β for $1 \leq \beta < \alpha$ are uniformly r.e. when indexed by $\{b: b <_0 a\}$.
- (3) For the same notation a for α , there is a recursive procedure which chooses, for each \bar{p} , uniformly in a notation $b <_0 a$ for β , an element $x \in A$ for which $x \notin \text{cl}_\beta(\bar{p})$.

PROOF. We use the same terminology as for the proof of Theorem 1. Let a be the notation for α given in assumptions (2) and (3), and let $\langle d_n \rangle$ be a recursive

sequence of notations for ordinals $\langle \beta_n \rangle$ such that $d_0 <_0 d_1 <_0 \dots <_0 a$ and $\lim_n \beta_n = \alpha$. For simplicity we may assume that $\beta_0 \geq 2$. Let $\gamma_n = \beta_n + 1$.

We define a special α -system w.r.t. $\langle \gamma_n \rangle$ as follows. The nodes of the tree of level n are those sequences of the form

$$\left(\mu_{(0)_0}^{(0)_1}(m_0) = k_0, \dots, \mu_{(n-1)_0}^{(n-1)_1}(m_{n-1}) = k_{n-1} \right)$$

where the μ_e^r are symbols corresponding to the partial functions $\phi_e^{d_r}$ and $n = \langle (n)_0, (n)_1 \rangle$ is a recursive pairing function for which $(n)_1 \leq n$.

A suitable label for such a node is a pair (f, x) for which $f \in P$, $x \in A$ and $x \notin \text{cl}_{\beta_n}(\text{ran}(f))$. A correct next label on the successor node $(\dots, \mu_{(n)_0}^{(n)_1}(m_n) = k_n)$ is a suitable label (g, y) such that $g \supseteq f$, $a_n, x \in \text{ran}(g)$, $b_n \in \text{dom}(f)$, and if $m_n = x$, then $g^{-1}(x) \neq k_n$. The function F is again defined so that $F(f, x) \cap \sigma \neq \emptyset$ if and only if f is coherent w.r.t. σ .

For $\beta < \alpha$, we define $(f, x) \triangleleft_\beta (g, y)$ if $f \triangleleft_\beta g$. The verification that the result is a $\langle \gamma_n \rangle$ -system is much as for the proof of Theorem 1.

The desired instruction is that which proceeds from a node of level n to the successor node $(\dots, \mu_{(n)_0}^{(n)_1}(x) = k)$ for which

$$k = \begin{cases} \phi_{(n)_0}^{d_{(n)_1}}(x) & \text{if this is defined,} \\ 0 & \text{(say) otherwise.} \end{cases}$$

Since $d_{(n)_1} <_0 c_0$, we can find a $\Delta_{c_n}^0$ index for this value of k , so we have defined a $\langle \gamma_n \rangle$ -instruction. By Proposition 2, there is a correct labelling of this having an r.e. adherent point giving, as in the previous proof, a recursive structure \mathfrak{B} and an isomorphism $f: \mathfrak{B} \cong \mathfrak{A}$ which is not $\Delta_{\beta_n}^0$ for any n .

COMMENTS ON THE PROOFS. Theorem 1 could instead be proved as in [1] using "binary patterns" as labels for an α -system. The proof given here seems simpler and, incidentally, shows that we could equally construct a recursive \mathfrak{B} and an isomorphism not in any uniform family of total $\Delta_{\alpha+1}^0$ functions. On the other hand, the method of [1] shows that the conditions of Theorem 1 can be weakened to those given in [1]. Assumption (3) can be replaced by:

(3)' There is a Δ_2^0 function which assigns to each \bar{p} an element of $A - \text{cl}_\alpha(\bar{p})$.

A more complicated argument, following the proof of the Main Lemma, shows that Theorem 2 can similarly be modified, replacing assumption (3) by

(3)' There is a Δ_2^0 function which yields, for each \bar{p} and each notation $b <_0 a$ for β , an element of $A - \text{cl}_\beta(\bar{p})$.

3. Infinitary formulae. We define the Σ_α and Π_α formulae of $L_{\omega_1\omega}$ as follows. The Σ_0 and Π_0 formulae are the quantifier-free formulae of $L_{\omega\omega}$. The $\Sigma_{\alpha+1}$ formulae are those of the form $\bigvee_{n \in S} \exists \bar{y}_n \phi_n$ where each ϕ_n as in Π_α formula and \bar{y}_n is a finite sequence of variables. The $\Pi_{\alpha+1}$ formulae are those of the form $\bigwedge_{n \in S} \forall \bar{y}_n \phi_n$ where each ϕ_n is a Σ_α formula. If δ is a limit ordinal, the Σ_δ formulae are $\bigvee_{n \in S} \phi_n$ where each ϕ_n is Σ_β for some $\beta < \delta$ and the Π_δ formulae are $\bigwedge_{n \in S} \phi_n$ where each ϕ_n is Π_β for some $\beta < \delta$.

For $a \in \mathcal{O}$, we define the (recursive) Σ_a and Π_a formulae and simultaneously, by recursive transfinite induction, their Gödel numbers. If $a = 2^b$ and if, for each $n \in W_e$, $\phi_e(n) = \langle i_n, j_n \rangle$ where i_n is a Gödel number for a Π_b formula θ_n and j_n

is a canonical index for a finite sequence of variables \bar{y}_n , then $\langle a, e, 0 \rangle$ is a Gödel number for the Σ_a formula $\bigvee_{n \in W_e} \exists \bar{y}_n \theta_n$. Similarly, if instead each i_n is a Gödel number of a Σ_b formula θ_n , then $\langle a, e, 1 \rangle$ is a Gödel number for the Π_a formula $\bigwedge_{n \in W_e} \forall \bar{y}_n \theta_n$. If $a = 3.5^k$ where $\phi_k(n) = a_n$ and if, for each n , $\phi_e(n)$ is a Gödel number for a Σ_{a_n} formula θ_n , then $\langle a, e, 0 \rangle$ is a Gödel number for the Σ_a formula $\bigvee_{n \in W_e} \theta_n$. If, instead, each θ_n is Π_{a_n} , then $\langle a, e, 1 \rangle$ is a Gödel number for the Π_a formula $\bigwedge_{n \in W_e} \theta_n$.

We define a recursive Σ_a formula to be one which is Σ_a for some $a \in \mathcal{O}$ such that $|a| = \alpha$. We note that, similarly to §16.8 of [1], if $|b| \leq |a|$ then we can effectively find, from a Gödel number of a Σ_b formula, a Gödel number of a logically equivalent Σ_a formula.

We make the following definition as seeming a natural condition under which \mathfrak{A} will be Δ_α^0 -stable.

DEFINITION. A formally Δ_α^0 enumeration of \mathfrak{A} consists of a finite sequence \bar{p} from A , an ordinal notation $a \in \mathcal{O}$ for α and a Σ_α^0 set S of Gödel numbers for a set of recursive Σ_a formulae such that, if ϕ_n denotes the formula with Gödel number n , then, for each $n \in S$, $\mathfrak{A} \models \phi_n[c, \bar{p}]$ for at most one element c of A and for each element c of A there exists at least one $n \in S$ such that $\mathfrak{A} \models \phi_n[c, \bar{p}]$.

COMMENT. Given a formally Δ_α^0 enumeration of \mathfrak{A} , we may convert it into one for which S is recursive and such that, for each $n \in S$, $\mathfrak{A} \models \phi_n[c, \bar{p}]$ for exactly one $c \in A$.

It is easy to see that the existence of such an enumeration is a sufficient condition for Δ_α^0 -stability.

THEOREM 3. *If a recursive structure \mathfrak{A} has a formally Δ_α^0 enumeration, then \mathfrak{A} is Δ_α^0 -stable.*

PROOF. First we note that, by recursive transfinite induction, from a Gödel number for a recursive Σ_a formula, we can compute an index for the Σ_α^0 relation which it determines on a recursive structure.

Now suppose that \mathfrak{B} is a recursive structure and $f: \mathfrak{B} \cong \mathfrak{A}$. Then for some $\bar{q} \in B$ we have $f: (\mathfrak{B}, \bar{q}) \cong (\mathfrak{A}, \bar{p})$ and $f(c) = d$ iff there exists $n \in S$ for which $\mathfrak{A} \models \phi_n[c, \bar{p}]$ and $\mathfrak{B} \models \phi_n[d, \bar{q}]$. But this relation is Σ_α^0 , and so f is Δ_α^0 .

The results of §2 allow us to prove the converse of Theorem 3 under certain assumptions. First, the following may be proved, substantially as for Lemmas 1 and 2 of [1].

LEMMA 5. *Suppose that the relations \leq_β on \mathfrak{A} for $1 \leq \beta < \alpha$ are uniformly r.e. when indexed by $\{b: b <_0 a\}$ for some $a \in \mathcal{O}$ with $|a| = \alpha$ and that the existential diagram of \mathfrak{A} is recursive. Then, from each $\bar{c} \in A$, we can effectively find a Gödel number for a recursive Π_a formula $\phi_{\bar{c}}^\alpha$ such that, for all $\bar{d} \in A$, $\bar{c} \leq_\alpha \bar{d}$ iff $\mathfrak{A} \models \phi_{\bar{c}}^\alpha[\bar{d}]$.*

LEMMA 6. *Under the same assumptions, if, for some \bar{p} , $\text{cl}_\alpha(\bar{p}) = A$, then \mathfrak{A} has a formally Δ_α^0 enumeration.*

Thus, the results of Theorems 1 and 3 may be combined as follows, using Lemma 6. We incorporate the improvements stated in §2.

THEOREM 4. *Suppose that the existential diagram of \mathfrak{A} is recursive, that the relations \leq_β for $1 \leq \beta < \alpha$ are uniformly r.e. when indexed by $\{b: b <_0 a\}$ for*

some $a \in \mathcal{O}$ where $|a| = \alpha$ and that the relation $x \in \text{cl}_\alpha(\bar{p})$ is Δ_2^0 . Then \mathfrak{A} is Δ_α^0 -stable iff \mathfrak{A} has a formally Δ_α^0 enumeration.

Similarly, Theorems 2 and 3 give

THEOREM 5. *Suppose that the existential diagram of \mathfrak{A} is recursive, that the relations \leq_β for $1 \leq \beta < \alpha$ are uniformly r.e., and that the relations $x \in \text{cl}_\beta(\bar{p})$ for $\beta < \alpha$ are uniformly Δ_2^0 . Then \mathfrak{A} is $\hat{\Delta}_\alpha^0$ -stable iff \mathfrak{A} has a formally Δ_β^0 enumeration for some $\beta < \alpha$.*

This establishes the following result, involving only the notions of §2.

COROLLARY. *Under the assumptions of Theorem 5, \mathfrak{A} is $\hat{\Delta}_\alpha^0$ -stable iff \mathfrak{A} is Δ_β^0 -stable for some $\beta < \alpha$.*

Another re-statement of these results concerns the Scott rank of a structure \mathfrak{A} . For sequences \bar{a} and \bar{b} of the same length from A , define $\bar{a} \sim \bar{b}$ if there is an automorphism of \mathfrak{A} which maps each entry of \bar{a} to the corresponding entry of \bar{b} . From several slight variations, we choose to define the rank of \mathfrak{A} to be the least ordinal α for which, for each $\bar{a} \in A$, there exists $\bar{c} \in A$ and $\beta < \alpha$ such that, for all \bar{b}, \bar{d} of the same lengths as \bar{a}, \bar{c} , if $\bar{a}, \bar{c} \leq_\beta \bar{b}, \bar{d}$, then $\bar{a} \sim \bar{b}$. Equivalently, the rank is the least ordinal α such that, for each $\bar{a} \in A$, there is a Σ_α formula θ such that $\bar{b} \sim \bar{a}$ iff $\mathfrak{A} \models \theta[\bar{b}]$. If the rank of \mathfrak{A} is α , then there exists a $\Pi_{\alpha+1}$ sentence ψ such that for each countable \mathfrak{B} , $\mathfrak{B} \cong \mathfrak{A}$ iff $\mathfrak{B} \models \psi$.

In the presence of our other assumptions a recursive structure \mathfrak{A} has a formally Δ_α^0 enumeration iff, for some \bar{p} , the structure (\mathfrak{A}, \bar{p}) has rank at most α and is rigid, that is, has no automorphism other than the identity function. Thus we have:

THEOREM 6. *Under the assumptions of Theorem 4, \mathfrak{A} is Δ_α^0 -stable iff, for some finite sequence \bar{p} from A , (\mathfrak{A}, \bar{p}) is rigid and has rank at most α .*

Similarly, under the assumptions of Theorem 5, \mathfrak{A} is $\hat{\Delta}_\alpha^0$ -stable iff, for some finite sequence \bar{p} from A , (\mathfrak{A}, \bar{p}) is rigid and has rank strictly below α .

COMMENT. The statement of Theorem 6 could be varied using the facts that if \mathfrak{A} has 2^{\aleph_0} automorphisms then so has each (\mathfrak{A}, \bar{p}) , while if \mathfrak{A} has fewer than 2^{\aleph_0} automorphisms then, for some \bar{p} , (\mathfrak{A}, \bar{p}) is rigid.

HYP-STABILITY. A final notion along the lines of this paper would be to define the recursive structure \mathfrak{A} to be *Hyp-stable* if, for each recursive $\mathfrak{B} \cong \mathfrak{A}$, every isomorphism from \mathfrak{B} to \mathfrak{A} is hyperarithmetical, that is, Δ_α^0 -recursive for some $\alpha < \omega_1^{CK}$. However, this matter is quickly disposed of, using the results from [4] that the rank of a recursive structure is at most ω_1^{CK} and that if its rank is equal to ω_1^{CK} , then the structure has 2^{\aleph_0} automorphisms.

Thus, if \mathfrak{A} has rank ω_1^{CK} , then \mathfrak{A} is not Hyp-stable. Now suppose that the rank of \mathfrak{A} is $\alpha < \omega_1^{CK}$. From the definition of rank, we have $\bar{a} \leq_\alpha \bar{b} \Rightarrow \bar{a} \sim \bar{b}$. Without further assumptions, we do not have the conclusion of Lemma 6. However, we can show instead, by recursive transfinite induction, that, uniformly in $\bar{a} \in A$ and a notation for a limit ordinal δ , one can find a Gödel number for a recursive $\Pi_{\delta+2n}$ formula ϕ such that $\bar{a} \leq_{\delta+n} \bar{b}$ iff $\mathfrak{A} \models \phi[\bar{b}]$. Thus, if (\mathfrak{A}, \bar{p}) is rigid, then \mathfrak{A} has a formally $\Delta_{\delta+2n+1}^0$ enumeration, and so by Theorem 3 is $\Delta_{\delta+2n+1}^0$ -stable. We have shown the following theorem.

THEOREM 7. *A recursive structure \mathfrak{A} is Hyp-stable iff \mathfrak{A} has fewer than 2^{\aleph_0} automorphisms. In this case, if the rank of \mathfrak{A} is $\delta + n$, where δ is a limit ordinal or zero, then \mathfrak{A} is $\Delta_{\delta+2n+1}^0$ -stable.*

4. Examples. For many examples of recursive algebraic structures, such as Boolean algebras and abelian groups, the relations \leq_α can be shown to be uniformly recursive. However, those examples of this kind which we have considered and which have infinite rank also have 2^{\aleph_0} automorphisms and so do not provide examples of Δ_α^0 -stable structures for α infinite. [They do provide examples of Δ_α^0 -categorical structures, but this is not our present topic.] We may, however, illustrate the notions of this paper by considering briefly the structures $(\alpha, <)$ and their expansions. These will always be rigid and have arbitrarily large ranks.

We make the following definition.

DEFINITION. A recursive structure is *strictly Δ_α^0 -stable* if it is Δ_α^0 -stable and not Δ_β^0 -stable for any $\beta < \alpha$.

We comment that this notion depends only on the classical isomorphism type of the structure, and that in view of Theorem 7, every recursive structure with fewer than 2^{\aleph_0} isomorphisms is strictly Δ_α^0 -stable for some unique $\alpha < \omega_1^{CK}$. In all our examples, by Theorem 2 and its consequences, the structures which are strictly Δ_α^0 -stable for some limit ordinal α are also not $\hat{\Delta}_\alpha^0$ -stable.

In what follows, δ will always denote a limit ordinal or zero and n will denote a finite ordinal.

EXAMPLE 1. In the case where $\mathfrak{A} = (\alpha, <)$ for some ordinal number α , we note that, if $\bar{a} \leq_\gamma \bar{b}$ for $\gamma \geq 1$, then the elements of the sequence \bar{a} will be in the same order as those of \bar{b} . We need therefore consider only the case where $\bar{a} = a_1, a_2, \dots, a_n$, $a_1 \leq a_2 \leq \dots \leq a_n$, $\bar{b} = b_1, b_2, \dots, b_n$, and $b_1 \leq b_2 \leq \dots \leq b_n$, and we may easily see that the relation $\bar{a} \leq_\gamma \bar{b}$ depends only on the order types of the intervals $[0, a_1), [a_1, a_2), \dots, [a_n, \infty)$, and those of the corresponding intervals for \bar{b} .

So we introduce the corresponding relations \leq_γ between ordinal numbers, including 0, and so between well-ordered sets. Let $\alpha \leq_1 \beta$ if $\alpha = \beta = 0$ or if α is infinite and $\beta > 0$ or if α is finite and $0 < \beta \leq \alpha$. Define $\alpha \leq_{\gamma+1} \beta$ if, whenever $\beta = \beta_0 + 1 + \beta_1 + 1 + \dots + 1 + \beta_m$, then there exist α_i such that $\alpha = \alpha_0 + 1 + \alpha_1 + 1 + \dots + 1 + \alpha_m$, and $\alpha_i \geq_\gamma \beta_i$ for $0 \leq i \leq m$. Define $\alpha \leq_\delta \beta$ to hold for a limit ordinal δ if $\alpha \leq_\gamma \beta$ for all $\gamma < \delta$.

Now we may show, by transfinite induction on γ , that if $a_1 \leq a_2 \leq \dots \leq a_n$ and $b_1 \leq b_2 \leq \dots \leq b_n$, then $\bar{a} \leq_\gamma \bar{b}$ iff $[0, a_1) \leq_\gamma [0, b_1), [a_n, \infty) \leq_\gamma [b_n, \infty)$ and for $1 \leq i < n$, $[a_i, a_{i+1}) \leq_\gamma [b_i, b_{i+1})$.

It remains to find the conditions for which $\alpha \leq_\gamma \beta$, which may be verified by transfinite induction. We note that each α may be written as a finite decreasing sum $\alpha = \sum_\xi \omega^\xi \cdot m_\xi$ where each $m_\xi < \omega$ and also, for each ξ , we may put $\alpha = \omega^\xi \cdot \alpha_\xi + \rho_\xi$ where $\rho_\xi < \omega^\xi$. [Thus $\omega^\xi \cdot \alpha_\xi = \sum_{\eta \geq \xi} \omega^\eta \cdot m_\eta$ and $\rho_\xi = \sum_{\eta < \xi} \omega^\eta \cdot m_\eta$, with the sums again taken in decreasing order.] Similarly, let $\beta = \sum_\xi \omega^\xi \cdot n_\xi$ and for each ξ let $\beta = \omega^\xi \cdot \beta_\xi + \sigma_\xi$ where each $n_\xi < \omega$ and each $\sigma_\xi < \omega^\xi$.

LEMMA 7. (i) *If $\gamma = \delta + 2n + 1$, then $\alpha \leq_\gamma \beta$ iff one of the following conditions hold.*

- (a) $\alpha = \beta$, (b) $\rho_{\delta+n} = \sigma_{\delta+n}$, $\alpha_{\delta+n+1} \geq 1$, and $\beta_{\delta+n} \neq 0$, (c) $\rho_{\sigma+n} = \sigma_{\delta+n}$, $\alpha_{\delta+n+1} = \beta_{\delta+n+1} = 0$, and $m_{\delta+n} \geq n_{\delta+n} > 0$.
- (ii) If $\gamma = \delta + 2n + 2$, then $\alpha \leq_{\gamma} \beta$ iff either
 - (a) $\alpha = \beta$, or (b) $\rho_{\delta+n} = \sigma_{\delta+n}$, $\alpha_{\delta+n+1} \geq 1$, $\beta_{\delta+n+1} \geq 1$, and $m_{\delta+n} \geq n_{\delta+n}$.
 - (iii) If $\gamma = \delta \neq 0$, then $\alpha \leq_{\gamma} \beta$ iff either
 - (a) $\alpha = \beta$, or (b) $\rho_{\delta} = \sigma_{\delta}$, $\alpha_{\delta} \geq 1$, and $\beta_{\delta} \geq 1$.

To use Theorems 1 and 2, we must also examine the sets $cl_{\gamma}(\bar{a})$. It is sufficient to consider only the cases where α is of the form ω^{ξ} . Then from Lemma 7 we may show the following.

LEMMA 8. (i) In $(\omega^{\delta+n}, <)$ for $n \geq 1$, $\delta + n \geq 2$ we have $cl_{\delta+2n}(\emptyset) = \omega^{\delta+n}$ while, for all \bar{a} , $\max\{a_i\} + \omega^{\delta+n-1} \notin cl_{\delta+2n-1}(\bar{a})$.
 (ii) In $(\omega^{\delta}, <)$ for $\delta \neq 0$ we have $cl_{\delta}(\emptyset) = \omega^{\delta}$ while, for all \bar{a} and all $\gamma < \delta$, we have, for example, $\max\{a_i\} + \omega^{\gamma} \notin cl_{\gamma}(\bar{a})$.

Thus, from (i), if $\delta < \omega_1^{CK}$ then by Lemma 6 $(\omega^{\delta+n}, <)$ has a formally $\Delta_{\delta+2n}^0$ enumeration and so is $\Delta_{\delta+2n}^0$ -stable. But, also from (i), for each \bar{p} , $\omega^{\delta+n} \neq cl_{\delta+2n-1}(\bar{p})$, which will show that $(\omega^{\delta+n}, <)$ is not $\Delta_{\delta+2n-1}^0$ -stable provided that the conditions of Theorem 1 can be satisfied. This presents no difficulty since for each notation $a \in \mathcal{O}$ for $\alpha < \omega_1^{CK}$ we may convert $\{b \in \mathcal{O} : b <_0 a\}$ to a recursive well-ordering of type ω^{α} by the process of exponentiation. That is, we let each sequence b_1, \dots, b_n for which $a >_0 b_1 \geq_0 b_2 \geq_0 \dots \geq_0 b_n$ represent the ordinal $\omega^{\beta_1} + \omega^{\beta_2} + \dots + \omega^{\beta_n}$ where $\beta_i = |b_i|$. For this ordering it is clear from Lemma 7 that each relation \leq_{γ} is recursive, uniformly in the notation $c <_0 a$ for γ . Thus condition (2) and similarly, by Lemma 8(i), condition (3) of Theorem 1 are satisfied. Condition (1) is also satisfied since, in this ordering, the successor relation is recursive.

In the same fashion, from Lemma 7 and Lemma 8(ii), we see that $(\omega^{\delta}, <)$ is Δ_{δ}^0 -stable but not $\hat{\Delta}_{\delta}^0$ -stable. Ordinals not of the form ω^{ξ} are nevertheless finite sums of such ordinals and each summand can be treated separately. So we obtain the following.

THEOREM 8. If $\omega^{\delta+n} \leq \alpha < \omega^{\delta+n+1}$ where $\delta + n \leq 2$, then $(\alpha, <)$ is strictly $\Delta_{\delta+2n}^0$ -stable.

The result also holds for $\delta + n = 1$, using Goncharov's Theorem, described in [1].

EXAMPLE 2. For structures of the form $(\alpha, <, S)$ where S is the successor relation, several shortcuts are possible. For example, by arguing in terms of formulae, we can show that $\bar{a} \leq_{\gamma} \bar{b}$ in $(\alpha, <, S)$ iff $\bar{a} \leq_{1+\gamma} \bar{b}$ in $(\alpha, <)$. So using the methods of Example 1 we have

THEOREM 9. If $\omega^{1+\delta+n} \leq \alpha < \omega^{1+\delta+n+1}$ where $\delta + n \geq 0$, then $(\alpha, <, S)$ is strictly $\Delta_{1+\delta+2n}^0$ -stable.

EXAMPLE 3. Similarly, for structures $(\alpha, <, S, L)$ where L is the set of limit ordinals, we have $\bar{a} \leq_{\gamma} \bar{b}$ in $(\alpha, <, S, L)$ iff $\bar{a} \leq_{2+\gamma} \bar{b}$ in $(\alpha, <)$.

This leads to the following theorem.

THEOREM 10. *If $\omega^{1+\delta+n} \leq \alpha < \omega^{1+\delta+n+1}$ where $\delta + n \geq 1$, then $(\alpha, <, S, L)$ is strictly $\Delta_{\delta+2n}^0$ -stable. [For $\alpha < \omega^2$, $(\alpha, <, S, L)$ is Δ_1^0 -stable.]*

EXAMPLE 4. For structures of the form $(\alpha, <, +)$, it is clearly preferable to proceed indirectly. We may first [by careful economy with quantifiers] show that in such a structure the set P_ξ of ordinals of the form $\omega^{\omega^\xi \cdot \beta}$ is given for each ξ by an infinitary recursive $\Pi_{\delta+2n}^0$ formula, where $\xi = \delta + n$ [except that P_0 is Π_1^0] and that the successor relation S_ξ relative to P_ξ is given by a recursive $\Pi_{\delta+2n+1}$ formula. This shows that the structure $(\omega^{\omega^{\delta+n+1}}, <, +)$ has a formally $\Delta_{\delta+2n+2}^0$ enumeration and so is $\Delta_{\delta+2n+2}^0$ -stable.

To show that this structure is strictly $\Delta_{\delta+2n+2}^0$ -stable, we may use Example 1. Since $\omega^{\delta+n+1}$ is strictly $\Delta_{\delta+2n+2}^0$ -stable, there exist recursive orderings of type $\omega^{\delta+n+1}$ which are not $\Delta_{\delta+2n+1}^0$ isomorphic. As in Example 1, these may be converted by exponentiation to structures of isomorphism type $(\omega^{\omega^{\delta+2n+1}}, <, +)$, and these structures cannot be $\Delta_{\delta+2n+1}^0$ -isomorphic. [Note, however, that this transformation may be used only in this direction, since in a recursive structure of type $(\alpha, <, +)$, the elements corresponding to the ordinals ω^ξ may not be recursive.] By similar arguments we have the following.

THEOREM 11. *If $\omega^{\delta+n} \leq \alpha < \omega^{\delta+n+1}$ where $\delta + n \geq 1$, then $(\omega^\alpha, <, +)$ is strictly $\Delta_{\delta+2n}^0$ -stable. [For $\alpha < \omega$, $(\omega^\alpha, <, +)$ is Δ_1^0 -stable.]*

EXAMPLE 5. For structures of the form $(\alpha, <, +, \cdot)$, the existential diagram will not be recursive, so Theorems 1 and 2 as given will certainly not apply. We may, however, follow Example 4 and show that $(\omega^{\omega^{\delta+n}}, <, +, \cdot)$ has a formally $\Delta_{\delta+2n}^0$ enumeration and then apply the same exponentiation construction again, using the fact that $(\omega^{\omega^{\delta+n}}, <, +)$ is strictly $\Delta_{\delta+2n}^0$ -stable, to show the following.

THEOREM 12. *If $\omega^{\delta+n} \leq \alpha < \omega^{\delta+n+1}$, where $\delta + n \geq 1$, then $(\omega^\alpha, <, +, \cdot)$ is strictly $\Delta_{\delta+2n}^0$ -stable. [For $\alpha < \omega$, $(\omega^\alpha, <, +, \cdot)$ is Δ_1^0 -stable.]*

5. Conclusion. The equivalence of Δ_α^0 -stability to the existence of a formally Δ_α^0 enumeration does seem to require further assumptions. A counter-example along these lines for $\alpha = 1$ was given by Goncharov in [5]. The author does not know at this stage whether the conditions of Theorems 1 and 2 can be significantly weakened. Certainly there are many counter-examples yet to be constructed.

The constructions of §1 can also be applied to the notion of Δ_α^0 -categoricity. These results will be given in [2]. E. Barker has obtained results concerning the topic of “intrinsically Σ_α^0 ” relations, which will be given in [3].

REFERENCES

1. C. J. Ash, *Stability of recursive structures in arithmetical degrees*, Ann. Pure Appl. Logic (to appear).
2. —, *Categoricity of recursive structures in hyperarithmetical degrees* (in preparation).
3. E. Barker, *Intrinsically Σ_α^0 relations* (preprint).
4. J. Barwise, *Admissible sets and structures*, Springer-Verlag, Berlin and New York, 1975.
5. S. S. Goncharov, *The quantity of nonautoequivalent constructivizations*, Algebra and Logic **16** (1977), 169–185.
6. H. Rogers, *Theory of recursive functions and effective computability*, McGraw-Hill, New York, 1967.