

*Noora Huotari*

RECYCLING OF WOOD- AND  
PEAT-ASH – A SUCCESSFUL WAY  
TO ESTABLISH FULL PLANT  
COVER AND DENSE BIRCH STAND  
ON A CUT-AWAY PEATLAND

UNIVERSITY OF OULU,  
FACULTY OF SCIENCE,  
DEPARTMENT OF BIOLOGY





ACTA UNIVERSITATIS OULUENSIS  
A Scientiae Rerum Naturalium 576

*NOORA HUOTARI*

**RECYCLING OF WOOD- AND PEAT-  
ASH – A SUCCESSFUL WAY TO  
ESTABLISH FULL PLANT COVER  
AND DENSE BIRCH STAND ON A  
CUT-AWAY PEATLAND**

Academic dissertation to be presented with the assent of  
the Faculty of Science of the University of Oulu for public  
defence in Kuusamonsali (Auditorium YB210), Linnanmaa,  
on 28 October 2011, at 12 noon

UNIVERSITY OF OULU, OULU 2011

Copyright © 2011  
Acta Univ. Oul. A 576, 2011

Supervised by  
Docent Eero Kubin  
Docent Eila Tillman-Sutela  
Professor Jari Oksanen

Reviewed by  
Professor Juhani Päivänen  
Doctor Florence Renou-Wilson

ISBN 978-951-42-9531-7 (Paperback)  
ISBN 978-951-42-9532-4 (PDF)

ISSN 0355-3191 (Printed)  
ISSN 1796-220X (Online)

Cover Design  
Raimo Ahonen

JUVENES PRINT  
TAMPERE 2011

## **Huotari, Noora, Recycling of wood- and peat-ash – a successful way to establish full plant cover and dense birch stand on a cut-away peatland.**

University of Oulu, Faculty of Science, Department of Biology, P.O. Box 3000, FI-90014

University of Oulu, Finland

*Acta Univ. Oul. A 576, 2011*

Oulu, Finland

### ***Abstract***

Mechanical harvesting of peat changes the original mire ecosystem completely, and without active measures these areas may remain non-vegetated even for decades. Afforestation is one of the most popular after-use options for cut-away peatlands in Finland since it has both economic and aesthetic values. Recycling of wood-ash as a fertilizer has been studied extensively in peatlands drained for forestry. Wood-ash is reported to promote tree growth in these areas without any significant negative impact to the environment and could, therefore, be a suitable option also on cut-away peatlands. However, the environmental effects of ash-fertilization on cut-away areas and on ground vegetation are not fully understood.

The impact of wood- and peat-ash application on the early establishment of ground vegetation and downy birch (*Betula pubescens*) seedlings and on post-fertilization element concentrations in plants and peat substrate were studied in a cut-away peatland. Six treatments of wood-ash, peat-ash, biotite or Forest PK-fertilizer were replicated in three blocks in different mixtures and quantities corresponding to 50 kg ha<sup>-1</sup> of phosphorus.

All the fertilizers accelerated the revegetation of the bare peat surface significantly, whereas the establishment of plants in the unfertilized area was non-existent even several years after the peat harvesting had ceased. The most striking difference between the wood- and peat-ash-fertilizers and the commercial Forest PK-fertilizer was the extensive coverage of fire-loving moss species in all the areas where ash was spread. Wood- and peat-ash application also accelerated the germination and early establishment of downy birch seedlings more efficiently than the PK-fertilizer. Ground vegetation proved to be highly important in increasing the early biomass production and carbon sequestration on ash-fertilized cut-away peatland. In addition, the below-ground biomass was equal to the above-ground biomass, or even greater.

Both wood- and peat-ash fertilization ensured an adequate level of nutrients for the early establishment of ground vegetation and birch seedlings in a cut-away peatland. The mosses and herbaceous plants proved to have a major role in retaining the nutrients and heavy metals that otherwise might have leached away from the ash-fertilized cut-away site during the early stages of the afforestation. Although both wood- and peat-ash proved to be suitable for the initial fertilization of afforested cut-away peatlands, a later application of nutrients may be needed to guarantee the growth in a energy-wood stand of downy birch over its entire rotation.

**Keywords:** afforestation, *Betula pubescens* Ehrh., biomass, biotite, carbon stock, fertilization, fire-loving mosses, Forest PK-fertilizer, heavy metals, nutrients, succession



## **Huotari, Noora, Puu- ja turvetuhka edistää kasvillisuuden muodostumista turvetuotannosta vapautuneilla suopohjilla.**

Oulun yliopisto, Luonnontieteellinen tiedekunta, Biologian laitos, PL 3000, 90014 Oulun yliopisto

*Acta Univ. Oul. A 576, 2011*

Oulu

### ***Tiivistelmä***

Turvetuotannon päätyttyä jäljelle jäävä suopohja on aluksi täysin paljas ja vailla maaperän siemenpankkia. Kasvipeitteen luontainen uudistuminen voi viedä jopa vuosikymmeniä. Ympäristönhoidollisesti onkin tärkeää, että suopohjat otetaan uuteen käyttöön mahdollisimman nopeasti tuotannon päätyttyä. Metsitys on tällä hetkellä suosituin suopohjien jälkikäyttömuoto Suomessa. Suopohjan turpeessa on tyypillisesti runsaasti typpeä, mutta niukasti muita kasvuun tarvittavia ravinteita. Puutuhka on osoittautunut pitkäaikaisissa metsäojitetujen turvemaiden tutkimuksissa kilpailukykyiseksi vaihtoehdoksi kaupallisille lannoitteille. Energiantuotannon sivutuotteena syntyvä puu- ja turvetuhka voisi soveltua hyvin myös suopohjien lannoitteeksi. Tuhkan käytöstä lannoitteena turvetuotannosta vapautuneilla suopohjilla ei kuitenkaan ole riittävästi tutkimustietoa.

Tässä työssä tutkittiin puu- ja turvetuhkan vaikutuksia turvetuotannosta vapautuneen suopohjan kasvittumiseen ja puun taimien alkukehitykseen viiden ensimmäisen kasvukauden ajan. Lisäksi tutkittiin kasvillisuuden ravinne- ja raskasmetallipitoisuuksien muutoksia sekä turpeen ravinteisuutta lannoituksen jälkeen.

Kaikki lannoitteet nopeuttivat merkittävästi kasvillisuuden muodostumista paljaalle suopohjalle, kun taas lannoittamaton alue pysyi kasvittomana. Tuhkalannoitetuille alueille syntyi nopeasti laajoja palopaikoilla viihtyvien pioneerisammalten kasvustoja, jotka peittivät ja samalla sitoivat paljaan ja irtonaisen turvemaan pinnan. Tuhkalannoitus edisti myös koivun taimien alkukehitystä tehokkaammin kuin kaupallinen Metsän PK-lannoite. Sammalista ja ruohovartisista kasveista muodostuva aluskasvillisuus ylitti puuntaimet selvästi biomassan määrässä ja toimi metsityksen alkuvaiheessa puuntaimia merkittävämpänä hiilensitojana. Lisäksi kasvien maanalainen biomassaa oli maanpäällistä biomassaa suurempi.

Sekä puu- että turvetuhka takasivat riittävän määrän ravinteita energiapuumetsikön alkukehitykselle. Aluskasvillisuus osoittautui tärkeäksi tuhkasta liukenevien ravinteiden ja raskasmetallien sitojaksi metsityksen alkuvaiheessa. Vaikka sammalten kadmiumpitoisuudet nousivat tuhkalannoituksen seurauksena, ne olivat kuitenkin alhaisia Suomessa aiemmin mitattuihin sammalten yleisiin pitoisuuksiin suhteutettuna. Tuhkalannoitus ei lisännyt haitallisten raskasmetallien pitoisuuksia koivun taimien ja ruohovartisten kasvien lehdissä ja varsissa. Tutkimuksen tulokset tukevat puu- ja turvetuhkan käyttöä energiapuumetsiköiden alkuvaiheen lannoitteena turvetuotannosta vapautuneilla suopohjilla.

**Asiasanat:** biomassa, biotiitti, hieskoivu, hiili, Metsän PK-lannoite, nuotiosammalet, raskasmetallit, ravinteet, suokasvitus, suopohjien metsittäminen, tuhkalannoitus





## Acknowledgements

I started my PhD process in summer 2003 when I was working as a trainee on the joint project “Metsähiisi” that was carried out by the Finnish Forest Research Institute (Metla) Muhos Research Unit and the Vapo Co. At that time, I did the first vegetation inventories at Hirvineva, but I wasn’t yet aware how interesting that research project would be and how far I would eventually go with this research work.

First and foremost, I want to express my deepest gratitude to my excellent supervisor Dr. Eila Tillman-Sutela. Eila, your door has always been open for me and whenever I have asked for help, you have supported me. I want to thank you also for all the interesting and eye-opening discussions during this process. My other supervisor, Dr. Eero Kubin, deserves all my gratitude for always believing in me and providing me the great opportunities to give talks in several meetings, all of which has been invaluable for me. I am also grateful for your efforts on obtaining funding for my research. I also want to thank my third supervisor, Professor Jari Oksanen, who helped me at the early stages of my research.

I wish to express my sincere gratitude to the staff of the Metla research unit at Muhos for providing me excellent working facilities and the best possible working environment. I owe my warmest thanks to everyone who I worked with on the “Metsähiisi” -project. I want to thank Mr. Jorma Issakainen for organizing the field trials, and Mr. Jorma Pasanen and Mr. Kyösti Markkanen for assistance in the field. I also thank Ms. Irene Murtovaara for fine-tuning the figures in my thesis and Mr. Jouni Karhu for helping with the statistical analysis during the research. Mr. Osmo Murtovaara is acknowledged for constructing all the equipment that I have needed during the field work. The staffs of the Metla laboratories in Muhos and Vantaa are acknowledged for the laboratory work.

The co-author of the first original article, Dr. Anneli Kauppi, is warmly thanked for her advice and help, and for the valuable comments on the thesis. I am also grateful to Professor Satu Huttunen for valuable comments on the fourth manuscript and on the thesis, and for all the help on practical matters during the final stages of my doctoral studies. Also, Mr. Mikko Moilanen, Dr. Juha Piispanen and anonymous referees are acknowledged for the valuable comments on the manuscripts.

I would like to express my warmest thanks to Dr Florence Renou-Wilson and Professor Emeritus Juhani Päivänen for reviewing this work. I want to thank also Mr. Andrew Pattison for revising the English text. The study was financially

supported by Metla, Vapo, National Technology Agency of Finland (Tekes), Oulun Energia, the Emil Aaltonen Foundation and the Thule Institute. All those institutions are greatly acknowledged.

All my friends are acknowledged for being a part of my life and for supporting me whenever I have needed it. All the delightful moments with you have been really priceless and refreshing during my studies and my life in general. My warmest thanks to my mom Riitta for her love and support. Without you I would never have been able to come this far! I also want to thank my twin sister Pia and my little brother Eero for being there. Pia, I also want to thank you for your help with the biomass samples! I want to remember also my late little brother Juha and my late dad Seppo. Haluan kiittää mummua ja puffaa kaikesta tuesta ja avusta, jota olen saanut teiltä aina kun olen tarvinnut. I want to thank my parents-in-law Raili and Reino and all my relatives for their support during the hard times and in joy. I should also thank our dogs, Siiri and Lilli, for keeping me in the move during this process.

Above all, I owe my greatest gratitude to my husband Pasi for making it possible to combine studying, research and two small children. I am grateful for your love and understanding. Finally I want to thank our lovely children Iris and Otso for bringing so much love and joy into my life and for reminding me what are the most important things in life.

Oulu, May 2011

Noora Huotari

## List of original articles

The thesis is based on the following articles, which are referred to in the text by their Roman numerals:

- I Huotari N, Tillman-Sutela E, Kauppi A & Kubin E (2007) Fertilization ensures rapid formation of ground vegetation on cut-away peatlands. *Canadian Journal of Forest Research* 37: 874–883.
- II Huotari N, Tillman-Sutela E, Pasanen J & Kubin E (2008) Ash-fertilization improves germination and early establishment of birch (*Betula pubescens* Ehrh.) seedlings on a cut-away peatland. *Forest Ecology and Management* 255: 2870–2875.
- III Huotari N, Tillman-Sutela E & Kubin E (2009) Ground vegetation exceeds tree seedlings in early biomass production and carbon stock on an ash-fertilized cut-away peatland. *Biomass and Bioenergy* 33: 1108–1115.
- IV Huotari N, Tillman-Sutela E & Kubin E (2011) Ground vegetation has a major role in element dynamics in ash-fertilized cut-away peatland. *Forest Ecology and Management* 261: 2081–2088.

*Author's contribution:* N. Huotari planned and carried out the vegetation inventory (I), biomass sampling (III) and collection of the plant samples for carbon, nutrient and heavy metal analyses (III, IV). She participated in birch seedling inventory (II) and in collecting the peat samples (IV). She was responsible for separating the below-ground plant biomass from the peat samples, and for drying and weighing all the biomass samples (III). She also participated in the preparation of the plant samples for the laboratory analyses (III). N. Huotari was responsible for the statistical analyses and was the main author of articles (I–IV).



# Contents

<b>Abstract</b>	
<b>Acknowledgements</b>	<b>7</b>
<b>List of original articles</b>	<b>9</b>
<b>Contents</b>	<b>11</b>
<b>1 Introduction</b>	<b>13</b>
<b>2 Aims of the research</b>	<b>17</b>
<b>3 Materials and methods</b>	<b>19</b>
3.1 Study site.....	19
3.2 Sampling and measurements.....	21
3.3 Laboratory analyses .....	21
3.3.1 Peat samples .....	21
3.3.2 Plant samples.....	22
3.4 Statistical analyses .....	22
<b>4 Results</b>	<b>23</b>
4.1 Total coverage of vegetation .....	23
4.2 Quantity and growth of tree seedlings.....	25
4.2.1 Birch seedlings .....	25
4.2.2 Willow and Scots pine seedlings .....	25
4.3 Plant biomass and C stock .....	26
4.4 Nutrient concentrations in peat .....	26
4.5 Nutrients and heavy metal concentrations in plants.....	27
4.5.1 Mosses.....	27
4.5.2 Herbaceous plants.....	28
4.5.3 Birch seedlings .....	28
4.6 Nutrient pools in peat and vegetation.....	28
<b>5 Discussion</b>	<b>33</b>
<b>6 Conclusions and future prospects</b>	<b>39</b>
<b>References</b>	<b>41</b>
<b>Original articles</b>	<b>47</b>



# 1 Introduction

It is estimated that 500 000–600 000 tons of wood- and peat-ash are produced annually as a by-product of energy production and forest industry in Finland (Emilsson 2006, Motiva 2007). Partly due to strict legislation (Ministry of Agriculture and Forestry 2007, 2009), the present practice is to transport the ash to landfills at a considerable cost. Massive promotion of biomass fuels in Europe (Renewable Energy Directive) is expected to further increase the amount of ash produced in power plants, which will cause a significant disposal problem for the industry in the near future. Consequently, decisions regarding cost-effective and environmentally acceptable recycling options remain a priority.

Recycling of wood-ash as a forest fertilizer has been studied extensively in recent decades, mainly in Scandinavia (Karlton *et al.* 2008). In Finland, pure wood-ash has been reported to promote tree growth on peatlands drained for forestry and the effect can be long-lasting, up to 50 years or more (Silfverberg & Huikari 1985, Silfverberg 1996, Moilanen *et al.* 2002). Wood-ash contains all the essential plant nutrients except nitrogen (N), which is vaporized during the combustion (Karlton *et al.* 2008). In addition to the valuable mineral nutrients, wood-ash also contains heavy metals. The use of wood-ash as a forest fertilizer has been questioned mainly because of its cadmium (Cd) content (Pasanen *et al.* 2001, Kepanen *et al.* 2005, Reimann *et al.* 2008). Although studies have shown that pure wood-ash can be used as a fertilizer on peatland forests without any significant negative impacts on the environment (Lodenius 2003, Moilanen *et al.* 2006, Solla-Gullon *et al.* 2006, Saarsalmi *et al.* 2006, Omil *et al.* 2007), the recycling and utilization of wood-ash in these areas is still insignificant.

In addition to wood, peat is also an important domestic fuel in Finland. Large-scale industrial peat harvesting for energy production started in 1970s due to the worldwide oil crisis (Vasander *et al.* 2003). In 2009, 5% of the Finland's total energy consumption was produced by peat and 20% by wood fuels (Official Statistics of Finland 2009). In recent years, large investments have been brought about to enable the integrated use of wood and peat in existing peat-fired power plants (Paappanen & Leinonen 2005). As a result, the actual amounts of mixed-ash and peat-ash generated in power plants are about two-fold compared to pure wood-ash (Motiva 2007). The elemental content of this ash may vary significantly, depending on the fuel and combustion technique (Emilsson 2006). In contrast to wood-ash, peat-ash contains lower levels of harmful heavy metals, such as Cd and Pb, but is also poorer in mineral nutrients (Nieminen *et al.* 2005, Mandre *et al.*

2010). However, the main concern for the use of peat-ash as a forest fertilizer is the relatively low potassium (K) content, which retards tree growth (Hytönen 1998, 2003).

Cut-away peatlands are former mire ecosystems that have been extensively disturbed by the mechanical harvesting process of peat. Peat extraction is usually accomplished in 20–50 years (Nyrönen 1996) and the remaining cut-away area is completely devoid of plants and viable soil seed bank (Salonen 1987, Huopalainen *et al.* 1998). Over the next decade, around 30 000 ha of cut-away peatlands will be abandoned by the peat industry in Finland (Leinonen 2010). Natural revegetation of these cut-away peatlands is extremely slow due to nutrient deficiency and instability of the loose peat substrate (Salonen & Laaksonen 1994, Campbell *et al.* 2002). With no actions, they will remain non-vegetated even for decades (Salonen 1992, Lavoie & Rochefort 1996, Bérubé *et al.* 2000, Lavoie *et al.* 2005). These bare cut-away areas are also a source of atmospheric carbon (Tuittila & Komulainen 1995, Tuittila *et al.* 1999, Alm *et al.* 2007). Thus, in order to diminish the negative impacts to the environment, such as wind erosion and leaching of nutrients and solid matter into watercourses, rapid revegetation of these bare cut-away areas is important.

In Finland, the peat industry operates mainly on leased areas and the landowners decide about the post-harvesting use of the area. Cut-away peatlands have been adapted for agriculture, reed canary grass (*Phalaris arundinacea* L.) cultivation, afforestation, bird sanctuaries and re-paludification (Selin 1999, Vasander *et al.* 2003). At present, afforestation is the most common after-use option in Finland (Holmgren *et al.* 2008), since it has economical, environmental and aesthetic values. Recently, demands for bioenergy and interest in the establishment of dense stands for producing energy wood in short rotations have increased. Downy birch is a common pioneer tree species on moist peat soils in Finland (Päivänen 2007). However, since the residual peat layer on cut-away areas is typically rich in organic N but poor in mineral nutrients, especially phosphorus (P) and K (Paavilainen & Päivänen 1995, Wind-Mulder *et al.* 1996), soil enrichment is needed. In consequence, recycling and utilization of ash as a fertilizer in these areas is a potential option.

Earlier studies on the effect of wood- and peat-ash application on cut-away peatlands have focused mainly on the nutrition and growth of tree species of economic value (Lehtonen & Tikkanen 1986, Hytönen 1998, Hytönen & Kaunisto 1999, Hytönen 2003, Mandre *et al.* 2010), whereas little attention has been paid to the establishment and development of ground vegetation. Although



ash-fertilization is reported to be a good ecological solution on peatlands drained for forestry, the impact of ash application on cut-away peatlands may be different, since the residual peat layer is considerably thinner and there is no existing vegetation prior to afforestation. Consequently, before its large-scale processing and recycling as a forest fertilizer can be recommended, an evaluation of the environmental consequences of wood- and peat-ash application is necessary.



## 2 Aims of the research

The main objective of the present work was to study the possibility to recycle wood- and peat-ash, a by-product of energy production and forest industry, as fertilizers for cut-away peatlands reserved for energy-wood production. The aim of the present work was to find an efficient method to establish dense, short-rotation birch stands on cut-away areas without generating negative environmental impacts. For these purposes, research was carried out to

1. study the effect of wood- and peat-ash application on the establishment of plant cover in a bare cut-away peatland (I)
2. examine the impact of ash application on the germination and early establishment of tree seedlings, especially downy birch (*Betula pubescens* Ehrh.), in a cut-away peatland (II)
3. quantify the biomass and carbon stock of ground vegetation and tree seedlings at the early stages of the afforestation in a cut-away peatland (III)
4. determine the post-fertilization nutrient status of peat substrate and the nutrient and heavy metal contents in plants (IV)
5. assess the post-fertilization nutrient pools in a young birch stand in a cut-away peatland (III, IV and the results presented here)



## 3 Materials and methods

### 3.1 Study site

The experimental field was established in 2000 at Hirvineva, in Liminka, Finland (64° 44' N, 25° 16' E, 45 m a.s.l.). The cut-away peatland was abandoned by the peat industry in 1996. The residual peat was medium humified moss-sedge peat (SC-p, H5 von Post) with some wood residues, and the thickness of the peat layer varied from 20 to 50 cm. The soil below the peat was sandy till and the groundwater level was about 50 cm. The average yearly temperature sum in the area is 1040 d.d.

The drainage of the experimental area was improved in August 2000 by cleaning the old ditches at 40 m distances. Three blocks were established on the experimental field, and each block was divided into six 40 m x 40 m trial plots (Fig. 1). The plots were split and 1 kg ha<sup>-1</sup> of downy birch (*Betula pubescens* Ehrh.) seeds was hand-sown on randomized halves on the non-vegetated peat surface in early September 2000. The other halves were left to be afforested by natural seed dispersal.

In May 2001 the following treatments were randomized between the plots and replicated in the three blocks: (1) unfertilized, (2) wood-ash 7.9 t ha<sup>-1</sup>, (3) mixed-ash 6.3 t ha<sup>-1</sup>, (4) peat-ash 4.8 t ha<sup>-1</sup>, (5) peat-ash 4.8 t ha<sup>-1</sup> + biotite 1.5 t ha<sup>-1</sup> and (6) Forest PK-fertilizer 0.5 t ha<sup>-1</sup> (Table 1). Prior to spreading, the nutrient contents of the ashes were analyzed in the laboratory of the Finnish Forest Research Institute (Metla) at Muhos (IV). The doses of the fertilizers per plot were adjusted corresponding to 50 kg ha<sup>-1</sup> of phosphorus, as recommended for peatlands drained for forestry (Paavilainen & Päivänen 1995).




Fig. 1. The experimental field at Hirvineva, Liminka.

Table 1. The amounts ( $\text{kg ha}^{-1}$ ) of the fertilizers used and nutrients applied in the experiment, adjusted to correspond to  $50 \text{ kg ha}^{-1}$  of phosphorus (P).

Fertilization treatments ( $\text{kg ha}^{-1}$ )	Applied nutrient amounts ( $\text{kg ha}^{-1}$ )				
	N	P	K	Ca	Mg
1) Unfertilized	-	-	-	-	-
2) Wood-ash	7874	-	50	161	961
3) Mixed-ash	6334	-	50	88	668
Wood-ash	3937	-	25	80	188
Peat-ash	2397	-	25	8	480
4) Peat-ash	4794	-	50	15	376
5) Peat-ash + Biotite	6294	-	50	90	481
Peat-ash	4794	-	50	15	376
Biotite	1500	-	-	75	105
6) Forest PK-fertilizer	556	-	50	89	122

## **3.2 Sampling and measurements**

Percentage coverage of the tree seedlings and ground vegetation was estimated visually by species in systematically placed square plots (60 cm x 60 cm) in July 2003, 2004 and 2005 (I). The quantity and dominant heights of the established tree seedlings were measured by species in systematically placed circular sample plots yearly 2001–2004 (II). The above-ground biomass of tree seedlings and herbaceous plants was harvested using scissors in systematically placed squares in July–August 2004 (III). The dead plant material (litter) was collected separately. The below-ground biomass of plants was sampled by taking volumetric peat samples from 0–10 cm and 10–20 cm peat layers in the center of the biomass squares. The roots, fine roots and rhizomes were then separated from the peat. Lichens and green parts of mosses were collected from the surface of the peat samples (III). All the plant samples were oven-dried and weighed.

The plant samples for analysis of carbon (III), nutrient and heavy metal (IV) were collected in July 2005. The volumetric peat samples for pH and nutrient analysis were collected from 0–10 cm and 10–20 cm peat layers in systematically placed square plots in August–September 2004 (IV).

## **3.3 Laboratory analyses**

### **3.3.1 Peat samples**

The peat samples were prepared and the nutrient concentrations analyzed at the Metla laboratory in Muhos (IV). The pH was measured by soil-water suspension. The concentration of soluble nitrogen (N) was analyzed from potassium sulphate (0.5 M) extract; the  $\text{NH}_4\text{-N}$  concentration by using a salicylate method and the  $\text{NO}_3\text{-N}$  concentration by reduction with hydrazine- copper sulphate mixture and then measuring the nitrite concentration colorimetrically, following the Finnish standard method (SFS 3029). The concentrations of soluble P, K, Calcium (Ca) and Magnesium (Mg) were analyzed from ammonium acetate (0.5 M) extract; K, Ca, and Mg by using an atomic absorption spectrophotometer (AAS) and  $\text{PO}_4\text{-P}$  by using a spectrophotometer. The determination limits for  $\text{NH}_4\text{-N}$  and  $\text{NO}_3\text{-N}$  were  $0.5 \text{ mg kg}^{-1}$  and for  $\text{PO}_4\text{-P}$   $1.5 \text{ mg kg}^{-1}$ , respectively.

The concentrations of total K, Ca and Mg were determined from the dry-ashed peat samples (dissolved by HCl) by using an atomic absorption spectrophotometer (AAS method). The concentrations of total N were determined

by using the Micro-Kjeldahl method (Kubin & Siira 1979) and that of total P photometrically by using the vanado-molybdate method (Halonen *et al.* 1983).

### **3.3.2 Plant samples**

The plant samples for analysis of carbon, nutrient and heavy metal were dried (105 °C), ground to a homogenous powder and digested in HNO<sub>3</sub>/H<sub>2</sub>O<sub>2</sub> -solution using a microwave oven in the Metla laboratory in Muhos (III, IV). Nutrient and heavy metal (K, P, Ca, Mg, sulphur (S), boron (B), manganese (Mn), zinc (Zn), iron (Fe), copper (Cu), nickel (Ni), Cd, aluminium (Al), lead (Pb)) contents were determined from the solution by ICP-emission spectrometry and C and N from the dry matter using CHN-analyzer in the Metla laboratory in Vantaa. The determination limit for Al was 8.0 mg kg<sup>-1</sup>, for Cd 0.05 mg kg<sup>-1</sup> and for Pb 1.50 mg kg<sup>-1</sup>. The C content 488 g kg<sup>-1</sup> (Jackson *et al.* 1997) was used in below-ground biomass and 450 g kg<sup>-1</sup> (Roxburg *et al.* 2006) in litter to estimate the C stock (III).

### **3.4 Statistical analyses**

Statistical methods used for the data analysis were two-way analysis of variance (ANOVA), one-way analysis of variance (ANOVA) with Tukey's multiple comparison tests and Repeated Measures ANOVA. The data that did not meet the assumptions of ANOVA in spite of transformation were tested using a Kruskal–Wallis H test or nonparametric two-factor analysis (by an extension of the Kruskal–Wallis test) with non-parametric Tukey-type multiple comparisons (Nemenyi) tests (Zar 1984). The computer package SPSS for Windows was used for all these statistical tests, with the exception of the nonparametric two-factor analysis with Tukey-type multiple comparisons tests, which was partly manually calculated. The significance level used in all tests was  $p < 0.05$ .



## 4 Results

### 4.1 Total coverage of vegetation

All the fertilizers significantly accelerated the establishment of vegetation in the bare cut-away peatland, whereas the establishment of plants in the unfertilized area was almost non-existent even several years after the peat harvesting had ceased (I and Figs. 2 and 3). Three growing seasons after the fertilization, the small fire-loving mosses, such as *Funaria hygrometrica* (Hedw.) and *Leptobryum pyriforme* (Hedw.) Wils., covered 50–60% of all the ash-fertilized areas, which was 10 times more than in the PK-fertilized areas. During the following two years, the coverage of pioneering moss species gradually decreased on all the fertilized areas as they were replaced by subsequent successional species (I).

Three growing seasons after the fertilization, the coverage of vascular plants in the fertilized areas was 5–8 times greater than in the unfertilized area (I). *Deschampsia cespitosa* (L.) P. Beauv. was by far the most abundant species in all the fertilized areas. At the same time, the dominant species in the unfertilized areas were *D. cespitosa* and *Eriophorum angustifolium* Honck. The total coverage of vascular plants further increased during the following two years in all treatments, but *B. pubescens* displaced *D. cespitosa* as a dominant species in the areas where wood-ash or mixed-ash had been applied.



**Fig. 2. The establishment of plants on the unfertilized area was non-existent even several years after peat harvesting had ceased.**



**Fig. 3. Dense and vigorous vegetation covered the cut-away area three growing seasons after ash application.**

## **4.2 Quantity and growth of tree seedlings**

### **4.2.1 Birch seedlings**

Both wood- and peat-ash application significantly promoted the germination and early establishment of downy birch seedlings on a cut-away peatland (II). More than 200 000 birch seedlings per hectare were observed in all the ash-fertilized areas already after the first growing season (in 2001), which was at its best 16 times greater in comparison to the unfertilized area and 2 times greater than in the PK-fertilized area. Three years later, more than 80% of the birch seedlings which had germinated in 2001 were still alive in the fertilized areas, and the total amount of seedlings had even increased due to natural dissemination. At the same time, the number of birch seedlings in the unfertilized plots had decreased drastically and only 14% of the seedlings germinated in 2001 were still alive in 2004. No significant differences in the quantity of birch seedlings were observed between the sown and the naturally regenerated areas.

Fertilization also significantly improved the growth of birch seedlings over the four-year observation period (II). After one growing season, the mixture of wood- and peat-ash resulted in dominant heights four times, and other fertilizers 2–3 times, greater as compared to seedling heights in the unfertilized areas. This difference became further accentuated during the three following years.

### **4.2.2 Willow and Scots pine seedlings**

The number of willow seedlings which established in the fertilized areas in 2001 varied from 50 000 to 80 000 plants per hectare, whereas the quantity in the unfertilized area was on an average 4 000 plants per ha (II). In contrast to birch and willow, the greatest quantity of Scots pine seedlings was in the unfertilized area, about 8 000 plants per ha, while in the fertilized areas their quantity varied from 0 to 3 000 plants per ha.

Wood- and peat-ash fertilizers resulted in willow seedling dominant heights 4–5 times greater compared to the heights in the unfertilized area already after the first growing season, the dominant heights for the Forest PK-fertilizer being 3 times greater (II). The differences between the treatments remained almost unchanged during the observation period.

There were no significant differences in the dominant heights of Scots pine seedlings between the fertilized and unfertilized treatments after the first growing

season (II). Three years later, the pine seedlings in the ash-fertilized areas had, however, gained more height than the seedlings in the unfertilized or PK-fertilized areas.

### **4.3 Plant biomass and C stock**

Four growing seasons after the fertilization, the total live, above-ground biomass of ground vegetation, comprising mosses and herbaceous plants, varied among all the ash-fertilized areas from 161 to 231 g m<sup>-2</sup>, whereas the corresponding biomass in the PK-fertilized area was 108 g m<sup>-2</sup> and in the unfertilized area was 24 g m<sup>-2</sup> (III). At the same time, the above-ground biomass of tree seedlings in all the ash-fertilized areas was 73–113 g m<sup>-2</sup>, in the PK-fertilized area 57 g m<sup>-2</sup> and in the unfertilized area 33 g m<sup>-2</sup>. The below-ground biomass of herbaceous plants and tree seedlings in the top 20 cm peat layer varied between 197 and 239 g m<sup>-2</sup> in the fertilized areas, whereas the corresponding biomass in the unfertilized area was 71 g m<sup>-2</sup>. The amount of plant litter deposited on the ground during the four growing seasons was 101–244 g m<sup>-2</sup> in the fertilized areas and 34 g m<sup>-2</sup> in the unfertilized area.

After four growing seasons, the greatest C stock of mosses, on average 55 g m<sup>-2</sup>, was in the various ash-fertilized areas, whereas the C stock in the PK-fertilized area was only 8 g m<sup>-2</sup> and in the unfertilized area it was non-existent (III). The corresponding C stock of the above-ground parts of herbaceous plants ranged from 22 g m<sup>-2</sup> to 51 g m<sup>-2</sup> in the fertilized areas and was 11 g m<sup>-2</sup> in the unfertilized area. At the same time, the C stock of tree seedlings was 16 g m<sup>-2</sup> in the unfertilized areas, on an average 44 g m<sup>-2</sup> among the ash-fertilized areas and 27 g m<sup>-2</sup> in the PK-fertilized areas.

The mean C stock of the below-ground parts of herbaceous plants and tree seedlings was 108 g m<sup>-2</sup> in the fertilized areas and 34 g m<sup>-2</sup> in the unfertilized area. The C stock in the litter deposited on the ground surface was from 46 g m<sup>-2</sup> to 110 g m<sup>-2</sup> in the fertilized areas and 20 g m<sup>-2</sup> in the unfertilized area.

### **4.4 Nutrient concentrations in peat**

The mean pH-H<sub>2</sub>O value of the peat in the unfertilized area was 4.6, while the pH in the fertilized areas was 0.1–0.5 units higher (IV). All the fertilization treatments decreased the concentrations of easily soluble nitrogen, NO<sub>3</sub>-N and

NH<sub>4</sub>-N, in peat in comparison to the unfertilized area but had no significant effect on the total N concentrations.

Ash-fertilization resulted in 2–6 times higher concentrations of easily soluble phosphorus (PO<sub>4</sub>-P) in the 0–10 cm peat layer in comparison to the unfertilized and PK-fertilized areas (IV). The concentrations of PO<sub>4</sub>-P were, however, below the reliable detection limit (<1.5 mg kg<sup>-1</sup>) in a third of the peat samples and, consequently, no statistical tests were applied to the data. Overall, fertilization increased the total P concentrations in the 0–10 cm peat layer compared to the unfertilized area, whereas the concentrations in the 10–20 cm peat layer remained unchanged or even decreased.

All the fertilizers, except peat-ash, doubled the concentrations of easily soluble K in the 0–10 cm peat layer compared to the unfertilized area (IV). At the same time, only the treatment using peat-ash + biotite significantly increased the total K concentrations in peat in comparison to the unfertilized area. Wood-ash, for its part, was the only fertilizer that significantly increased the concentrations of Ca and Mg in peat in comparison to the unfertilized area, and the majority of these nutrients were in easily soluble form.

## **4.5 Nutrients and heavy metal concentrations in plants**

### **4.5.1 Mosses**

The nutrient and heavy metal concentrations in mosses were measured only in the fertilized areas, since there was no moss cover in the unfertilized plots (I, IV). Four years after the fertilization, the P concentration of mosses was significantly higher in the peat-ash fertilized area than in the wood-ash or PK-fertilized areas (IV). At the same time, fertilization had no significant effect on the N and K contents of mosses. Wood-ash increased especially the B contents of mosses in comparison to other fertilizers, and peat-ash especially the Na contents.

The cadmium (Cd) contents of mosses were 2–3 times higher in the ash-fertilized areas than in the PK-fertilized area (IV). At the same time, the concentrations of Al, Cu, Fe and Ni in mosses were higher in the PK-fertilized area than in the ash-fertilized areas. The contents of Pb were below the reliable detection limit (1.5 mg kg<sup>-1</sup>) in all the treatments.

#### **4.5.2 Herbaceous plants**

All the fertilization treatments resulted in significantly decreased N, K, S, Zn and Cu contents in *Deschampsia cespitosa* in comparison to the unfertilized treatment (IV). Fertilization also decreased the concentrations of P, Ca, Fe, Ni and Al, although these differences were not statistically significant. Furthermore, the same phenomenon was observed also concerning the Mg, Mn, Zn and Al contents in other herbs and grasses.

Fertilization either had no significant effect or it decreased the concentrations of heavy metals both in *D. cespitosa* and in other herbaceous plants in comparison to the unfertilized area (IV). The contents of Cd and Pb were below the reliable detection limit in the majority of the herbaceous plant samples.

#### **4.5.3 Birch seedlings**

The foliar N contents in all the fertilized areas were comparable to the unfertilized area (IV). All the fertilizers, with the exception of wood-ash, significantly increased the P contents in birch leaves in comparison with the unfertilized treatment. Only the forest PK-fertilizer resulted in significantly increased foliar K contents in comparison to the unfertilized area. In addition, peat-ash + biotite application significantly increased the foliar K contents in comparison to the pure peat-ash. Treatment using peat-ash resulted in higher foliar Ca and Mg concentrations in comparison to the unfertilized area. Overall, the nutrient content in the leaves of birch seedlings was higher than the nutrient content in the stems. None of the fertilization treatments significantly increased the heavy metal concentrations in birch seedlings in comparison with the unfertilized area.


#### **4.6 Nutrient pools in peat and vegetation**

The mean biomasses and nutrient concentrations presented in articles III and IV were used to estimate the N, P, K, Ca and Mg pools ( $\text{kg ha}^{-1}$ ) in the top 20 cm peat substrate and in the above-ground parts of birch seedlings and ground vegetation.

Total N pool in the top 20 cm peat layer was abundant in all the treatments, ranging between 7314 and 8991  $\text{kg ha}^{-1}$  (Fig. 4a). The P pool of peat varied from 258  $\text{kg ha}^{-1}$  in the wood-ash fertilized area to 345  $\text{kg ha}^{-1}$  in the PK-fertilized area (Fig. 4c). At the same time, the largest K pool of peat, 100  $\text{kg ha}^{-1}$ , was in the treatment which used peat-ash + biotite and the smallest, 53  $\text{kg ha}^{-1}$ , in the pure


peat-ash treatment (Fig. 4e). The total Ca pool in peat ranged from 2053 kg ha<sup>-1</sup> in the PK-fertilized area to 3614 kg ha<sup>-1</sup> in the mixed-ash fertilized area (Fig. 5a). The Mg pool of peat was 298 kg ha<sup>-1</sup> in the PK-fertilized area and varied between 405 and 494 kg ha<sup>-1</sup> among the other treatments (Fig. 5c). The proportions of easily soluble N and P in peat were 1% or less of the total pools (Figs. 4a–d). At the same time the corresponding share for K, depending on the treatment, was 28–68% (Fig. 4e–f), for Ca 67–74% (Fig. 5a–b) and for Mg 72–93% (Figs. 5c–d).

The amounts of N, P, Ca and Mg bound in the above-ground plant biomass made up only 1% or less of the total pools in the peat substrate in all the treatments (Figs. 4a, 4c, 5a, 5c), whereas the corresponding share for K was 5% in the unfertilized area and varied from 18% to 40% in the fertilized areas (Fig. 4e). In comparison with the soluble nutrients in peat, the amounts of N, P and K bound in plant biomass were consistently high in all the fertilized areas (Figs. 4b, 4d, 4f).



**Fig. 4.** The total (a, c, e) and soluble (b, d, f) N, P and K pools (kg ha<sup>-1</sup>) in peat in relation to the amounts bound in above-ground plant biomass in the six fertilization treatments. No=unfertilized, Wa=wood-ash, Ma=mixed-ash, Pa=peat-ash, Pa+B=peat-ash+biotite, PK=Forest PK-fertilizer. The nutrient pools were calculated on the basis of the biomass figures and nutrient concentrations presented in articles III and IV.





**Fig. 5.** The total (a, c) and soluble (b, d) Ca and Mg pools (kg ha<sup>-1</sup>) in peat in relation to the amounts bound in above-ground plant biomass in the six fertilization treatments. No=unfertilized, Wa=wood-ash, Ma=mixed-ash, Pa=peat-ash, Pa+B=peat-ash+biotite, PK=Forest PK-fertilizer. The nutrient pools were calculated on the basis of the biomass figures and nutrient concentrations presented in articles III and IV.



## 5 Discussion

Both wood- and peat-ash fertilization ensured a rapid formation of dense ground vegetation on a cut-away peatland, whereas the establishment of plants in the unfertilized areas was sparse even several years after the cessation of peat harvesting (I). Interestingly, ash-fertilizers in particular increased the coverage of fire-loving moss species, such as *Funaria hygrometrica* (Hedw.) and *Leptobryum pyriforme* (Hedw.) Wils., whereas their cover in the PK-fertilized and unfertilized areas was insignificant. The colonization of moss species on the bare ash-fertilized cut-away area resembled a process after severe natural fire (Rees & Juday 2002, Ryömä & Laaka-Lindberg 2005). Furthermore, the ash itself appeared to have a greater impact on the colonization success of fire-loving mosses than did open space or improved nutritional conditions (I), even if the aerial diaspore dispersal ability of mosses in harvested peatlands is reported to be high (Campbell *et al.* 2003). This result indicates that in addition to nutrients, there are also other components in plant-derived ash, comparable to those found in plant-derived smoke (Dixon *et al.* 2009), that promote the germination of different plant species.

Ash application in particular also improved the germination and early establishment of downy birch (*Betula pubescens* Ehrh.) in a cut-away peatland (II). As a pioneer species, birch in many cases colonizes bare grounds rapidly. However, drought and instability of the loose peat surface on bare windswept cut-away areas do not favor the establishment of plants (Salonen *et al.* 1992, Salonen 1994, Cambell *et al.* 2002). In addition, frost heaving is reported to complicate the rooting and survival of young tree seedlings in open cut-away peatlands (Groeneveld & Rochefort 2002, Lavoie *et al.* 2005). The newly risen moss cover in the ash-fertilized areas apparently formed a favorable germination ground for the birch seeds by stabilizing the loose erodible substrate and the moisture content of the peat surface (I, II). On the other hand, the quantity of Scots pine seedlings was more than three times higher on the unfertilized areas compared to the ash-fertilized areas (II), which confirmed the reported retarding effect of ash on the germination of pine seeds (Rikala & Jozefek 1990, Reyes & Casal 2004). In fact, the germination response of plants to fire, smoke or ash varies between species and may therefore influence the species composition in post-fire succession (González-Rabanal & Casal 1995, Zuloaga-Aguilar *et al.* 2011).

The quantities of birch seedlings which germinated in 2001 remained high and almost unchanged during the four-year observation period in all the fertilized

areas, whereas the number of original birch seedlings in the unfertilized area decreased drastically (II). Fertilization also increased the dominant heights of the birch seedlings in comparison to the unfertilized treatment. Surprisingly, the dense herbaceous vegetation, comprised mainly of *Deschampsia cespitosa*, seemed to facilitate rather than prevent the early development of birch seedlings (I, II). This was probably due to a more stable microclimatic temperature and moisture conditions within the vegetation, which protected the birch seedlings from wind and spring and autumn frosts. The results thus confirm the importance of facilitation in the succession mechanism of plant colonization of harvested peat surfaces (Tuittila *et al.* 2000, Groeneveld & Rochefort 2005, Koyama & Tsuyuzaki 2010).

The results proved that the post-fertilization above-ground biomass of mosses and herbaceous plants was up to two times greater than the above-ground biomass of tree seedlings (III). A rough estimation of the amount of C fixed by the above-ground parts of the herbaceous plants in the ash-fertilized areas was  $35 \text{ g C m}^{-2} \text{ a}^{-1}$ , and by mosses  $14 \text{ g C m}^{-2} \text{ a}^{-1}$ . Since the above-ground parts of herbaceous plants represent one year's growth (Mälkönen 1974), the standing biomass of herbaceous plants measured here was assumed to be equal to their annual above-ground production (III). The annual production of mosses was assumed to be a quarter of the standing biomass that they had built up during the four post-fertilization growing seasons. In the present life cycle analysis (LCA) of peat fuel utilization chains, only C fixed by the growing trees is used when estimating the C balance in afforested cut-away peatlands (Kirkinen *et al.* 2007, Hytönen *et al.* 2008). However, the total amount of C fixed by the above-ground parts of the ground vegetation in this study was at least 15% of the quantities ( $45\text{--}330 \text{ g C m}^{-2} \text{ a}^{-1}$ ), that are used in the actual calculations. Those quantities do not include the C bound in the ground vegetation and deposited litter, due mainly to the absence of exact studies. Since the role of stand age is of great importance when explaining the variation of plant biomass during succession process (Muukkonen & Mäkipää 2006), the biomass figures presented in this study are consistent only with the early stages of the afforestation process. Therefore, empirical data on the entire forest vegetation growth during the rotation period in afforested cut-away areas is needed.

The present results showed also that the below-ground biomass of plants, i.e. roots and rhizomes, was equal to the above-ground biomass, or even greater (III). This is in accordance with earlier findings in boreal forests, where more than 70% of the field layer biomass is reported to be in roots (Palviainen *et al.* 2005). Fine-

roots are reported to produce the majority of the annual biomass in tree stands of varying ages (Helmisaari *et al.* 2002). Thus, the total annual C sequestration of ground vegetation may be considerably larger than estimated above (III). The decomposition rates of different types of plant litter, e.g. root litter and leaf litter vary. Therefore, the ratio of litter input to decomposition output should be measured, when attempting to assess the overall effect of ground vegetation on C balance.

All the fertilizers ensured an adequate level of nutrients in the top 20 cm peat layer for the early establishment and growth of vigorous ground vegetation and tree seedlings in a cut-away peatland (IV). In previous studies, successful tree growth in cut-away areas is reported to be limited mainly by low levels of P and K in the peat substrate (Lumme 1988, Hytönen 1994, Aro *et al.* 1997). The present results proved that four years after the fertilization the P concentrations in birch leaves in all the fertilized areas were even higher than the average values reported for varying aged birches not suffering from P deficiency (Ferm and Markkola 1985, Sarjala & Kaunisto 2002). At the same time, the foliar K concentrations of the vigorous birch seedlings in the areas fertilized with mixed-ash or peat-ash were near the potassium deficiency level reported for older birches in cut-away peatlands (Sarjala & Kaunisto 2002, Kaunisto & Sarjala 2003). This implies that the nutrient demands of trees vary at different ages and that, at later stages, deficiency of K may be a limiting factor for tree growth. On the other hand, the dilution effect due to increased biomass may also have decreased the nutrient concentrations in birch leaves. For the further evaluation of nutrient sufficiency over the planned 30-year rotation, the nutrient pools in peat and plant biomass were calculated.

The total N pool in peat was large in comparison to the amounts bound in the above-ground plant biomass (Fig. 4a), which indicates a low risk for future N deficiency. Even if ash itself contains no N, application of ash is reported to accelerate the mineralization of organically bound N in peat due to a rise in pH and bacterial activity (Weber *et al.* 1985, Hytönen 1998, Moilanen *et al.* 2002, Mahmood *et al.* 2003, Pitman 2006 and references within). Furthermore, ground vegetation proved to have a major role in retention of the available N in peat (Fig. 4b) that otherwise could have leached from the unvegetated cut-away peatland area during the early stages of the afforestation (IV). Also, the total P, Ca and Mg stores in peat were abundant in relation to the demands of plants (Figs. 4c, 5a, 5c). However, since only 1% or less of the total P reserve was in easily soluble form

and thus readily available for plants, the sufficiency of P for tree growth in the long term is largely determined by the rate of mineralization.

At the same time, the total K pools in peat were rather scant in all the treatments, except in that of peat-ash + biotite, which implies a potential risk for future deficiency. On the other hand, the K bound in the plant biomass contributed considerably to the total K store of the site. Since K is characterized by high mobility both in peat soils (Rydin & Jeglum 2006) and within plants (Marschner 1995), its biological cycling is efficient and leaching losses usually insignificant (Laiho *et al.* 1999). Thus, even limited K pool may be sufficient for tree growth for the planned 30-year rotation period. Furthermore, the tree roots may penetrate the sub-peat mineral soil within some years and the K deficiency of trees can thus be minimized.

Interestingly, the K store in the area fertilized with pure peat-ash was almost equal to the K store in the area fertilized with wood-ash, even if the amounts of K applied with wood-ash treatment was ten times larger in comparison with that of the peat-ash. Thus, the post-fertilization concentrations of easily soluble K in the wood-ash fertilized area apparently exceeded the sequestration capacity of the vegetation and resulted in leaching of the excess K to the lower soil layers. The solubility of K in peat-ash is reported to be considerably lower than in wood-ash (Nieminen *et al.* 2005), which probably minimized the leaching losses in the peat-ash fertilized area and equalized the fertilization effects of wood- and peat-ash.

The post-fertilization concentrations of Cd and other harmful heavy metals varied among the studied plant groups (IV). The Cd contents in mosses in the ash-fertilized areas were 2–3 times higher than in the PK-fertilized area, whereas the concentrations of Cd and other harmful heavy metals in the shoots of herbaceous plants and birch seedlings remained at low levels despite the ash application. However, the levels of Cd and other harmful heavy metals were relatively low in comparison with the average concentrations measured in mosses in Finland (Poikolainen *et al.* 2004). The main path for elements' uptake in rootless mosses is through the leaves, and contact with upper soil layer is reported to have a major influence on their chemical composition (Reimann *et al.* 2001). Unlike mosses, herbaceous plants and trees absorb nutrients mainly from the soil via their roots. Consequently, the unchanged levels of Cd and other heavy metals in the tissues of vascular plants observed here may be due to differences in the heavy metal concentrations between the above- and below-ground parts of the plants, as has been reported in tree seedlings in the laboratory conditions (Österås *et al.* 2000).

However, the impact of ash-application on the heavy metal levels in the roots of herbaceous plants or trees is not extensively studied.

The present results indicated that dense pioneer moss cover was important in trapping the heavy metals, such as Cd, present in ash (IV). Thus, due to their great biomass (III) and consistently slow decomposition rate in comparison to the herbaceous plants (Lang *et al.* 2009, Turetsky *et al.* 2010), mosses may be more important than previously estimated in preventing the fast recycling of Cd in the ecosystem. These findings were contrary to earlier studies in which wood-ash was applied onto the existing moss cover in mature forests and no significant change in the heavy metal concentrations of mosses was found, even if the moss cover suffered severely from direct contact with large quantities of ash (Jacobson & Gustafsson 2001, Moilanen *et al.* 2002, Ozolinčius & Varnagirytė 2005, Ozolinčius *et al.* 2007). In this study, the ash was spread on the bare peat surface, which favored the colonization of moss species adapted to post-fire conditions, such as *Ceratodon purpureus*, *Funaria hygrometrica* and *Leptobryum pyriforme* (I).

Low levels of Cd (1–5 ppm) are reported to stimulate the germination and early growth of *F. hygrometrica*, whereas higher levels inhibit the germination process completely (Lepp & Roberts 1977). Also Cu, Pb and Zn are known to inhibit the germination and growth of several moss species even at considerably low levels (Francis & Petersen 1989, Tyler 1990 and ref. within) and the same phenomenon has been reported also in some herbaceous plant species (Lefèvre *et al.* 2009). Consequently, since the Cd tolerance levels of *F. hygrometrica* are known to be quite finite, it could be considered as an indicator species for low levels of soluble Cd in the growing substrate.





## 6 Conclusions and future prospects

Natural regeneration together with wood- and peat-ash application proved to be an efficient way to establish a dense birch stand for energy-wood production in a cut-away peatland. Ash itself had a greater impact on the colonization success of several pioneering species, such as fire-loving mosses and birch, than did open space or artificial fertilization.

The establishment of dense ground vegetation of mosses and herbaceous plants in all the ash-fertilized areas had several environmental advantages. The vegetation stabilizes the loose peat surface and decreases erosion and leaching of nutrients and solid matter into watercourses. The vigorous ground vegetation also facilitated rather than prevented the early growth of birch seedlings in a cut-away peatland and should thus be given consideration when afforesting such areas.

The dense ground vegetation was important also in the biomass production and C sequestration at the early stages of the afforestation. However, the long-term dynamics of ground vegetation and litter accumulation in afforested cut-away areas are not well known. To enable more accurate calculations on the C balance in afforested cut-away peatlands for life cycle analysis and climate impact studies, vegetation inventories and biomass measurements at different succession stages, e.g. at five-year intervals, are needed.

Both wood- and peat-ash ensured an adequate level of nutrients for the early establishment and growth of ground vegetation and birch seedlings in a cut-away peatland. However, a later application of nutrients, especially K, may be needed to guarantee successful tree growth over the entire 30-year rotation. Also, future harvesting of energy-wood may disturb the nutrient balance of the forest stand and retard the establishment of second tree generation. Therefore, the nutrient status of the plant community should be measured at ten-year intervals. Information about the stand's nutrient status could be accumulated not only by the measurement of the nutrients in peat and in the leaves of trees, but also in ground vegetation.

The dense pioneer moss cover was important in trapping the heavy metals present in the ash, whereas the levels of Cd and other harmful heavy metals remained at low levels in the shoots of herbaceous plants and birch seedlings. Due to the dilution effect of vigorous vegetation, the post ash-application Cd levels in herbaceous plants were even below the determination limits and in birch seedlings below the natural levels. However, studies on the concentrations in the

roots of trees and herbaceous plants are needed for a more complete view of the heavy metal levels in plants after ash-application.

The results support the use of wood- and peat-ash as fertilizers for the afforestation of cut-away peatlands. The recycling of ash would provide an environmentally sustainable way for producing energy-wood in these areas. In addition, rapid establishment of dense vegetation increases the C fixation into the ecosystem and may simultaneously reduce atmospheric carbon dioxide emissions from bare cut-away areas. However, grading and stabilization of the nutritional quality of ashes prior to granulation or pelletizing is a pre-requisite for large-scale use of ash as a forest fertilizer.

## References

- Alm J, Shurpali NJ, Minkkinen K, Aro L, Hytönen J, Laurila T, Lohila A, Maljanen M, Martikainen PJ, Mäkirinta P, Penttilä T, Saarnio S, Silvan N, Tuittila ES & Laine J (2007) Emission factors and their uncertainty for the exchange of CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O in Finnish managed peatlands. *Boreal Environmental Research* 12: 191–209.
- Aro L, Kaunisto S & Saarinen M (1997) Suopohjien metsitys. Hankeraportti 1986–1995. Summary: Afforestation of peat cutaway areas. Project report in 1986–1995. The Finnish Forest Research Institute, Research paper 634.
- Bérubé M-E & Lavoie C (2000) The natural revegetation of a vacuum-mined peatland: eight years of monitoring. *Canadian Field-Naturalist* 114(2): 279–286.
- Campbell DR, Lavoie C & Rochefort L (2002) Wind erosion and surface stability in abandoned milled peatlands. *Canadian Journal of Soil Science* 82: 85–95.
- Campbell DR, Rochefort L & Lavoie C (2003) Determining the immigration potential of plants colonizing disturbed environments: the case of milled peatlands in Quebec. *Journal of Applied Ecology* 40: 78–91.
- Dixon KW, Merritt DJ, Flematti GR & Ghisalberti EL (2009) Karrikinolide – a phytoreactive compound derived from smoke with applications in horticulture, ecological restoration and agriculture. *Acta Horticulturae* 813: 155–170.
- Emilsson S (2006) International handbook: from extraction of forest fuels to ash recycling. Skogsstyrelsen, Swedish Forest Agency.
- Ferm A & Markkola A (1985) Hieskoivun lehtien, oksien ja silmujen ravinnepitoisuuksien kasvukautainen vaihtelu. Abstract: Nutritional variation of leaves, twigs and buds in *Betula pubescens* stands during the growing season. *Folia Forestalia* 613.
- Francis PC & Petersen RL (1989) Assessment of toxicity of heavy metal ion combinations on spore germination and protonemal growth of *Polytrichum commune*. *The Bryologist* 92(1): 60–67.
- González-Rabanal F & Casal M (1995) Effect of high temperatures and ash on germination of ten species from gorse shrubland. *Vegetation* 116: 123–131.
- Groeneveld E & Rochefort L (2002) Nursing plants in peatland restoration: on their potential use to alleviate frost heaving problems. *Suo* 53(3–4):73–85.
- Groeneveld E & Rochefort L (2005) *Polytrichum strictum* as a solution to frost heaving in disturbed ecosystems: a case study with milled peatlands. *Restoration Ecology* 13(1): 74–82.
- Halonen O, Tulkki H & Derome J (1983) Nutrient analysis methods. The Finnish Forest Research Institute, Research Papers 121: 1–28.
- Helmisaari HS, Makkonen K, Kellomäki S, Valtonen E & Mälkönen E (2002) Below- and above-ground biomass, production and nitrogen use in Scots pine stands in eastern Finland. *Forest Ecology and Management* 165: 317–326.
- Holmgren K, Kirkinen J & Savolainen I (2008) Climate impact of peat fuel utilization. In: Strack M (ed) *Peatlands and Climate Change*. Jyväskylä, International Peat Society: 123–147.

- Huopalainen M, Tuittila E-S, Laine J & Vasander H (1998) Seed and spore bank in a cut-away peatland twenty years after abandonment. *International Peat Journal* 8. International Peat Society.
- Hytönen J (1994) Effect of fertilizer application rate on nutrient status and biomass production in short-rotation plantations of willows on cut-away peatland areas. *Suo* 45(3): 65–77.
- Hytönen J (1998) Puutuhkasta ja muista jäteteineista valmistetut pelletit rauduskoivun taimien ravinnelähteenä. Summary: Pellets made of wood ash and other wastes as nutrient sources for silver birch seedlings. *Suo* 49(2): 49–63.
- Hytönen J (2003) Effects of wood, peat and coal ash fertilization on Scots pine foliar nutrient concentrations and growth on afforested former agricultural peat soils. *Silva Fennica* 37(2): 219–234.
- Hytönen J & Kaunisto S (1999) Effect of fertilization on the biomass production of coppiced mixed birch and willow stands on a cut-away peatland. *Biomass and Bioenergy* 17: 455–69.
- Hytönen J, Aro L, Maljanen M, Mäkiranta P, Potila H, Laine J, Lohila A, Martikainen P, Minkkinen K, Pihlatie M & Shurpali N (2008) The effect of afforestation of organic croplands and cutaway peatlands on greenhouse gas balance. In: Sarkkola S (ed) *Greenhouse effect of the use of peat and peatlands in Finland*. Publications of Ministry of Agriculture and Forestry 11a/2007: 31–37.
- Jackson RB, Mooney HA & Schulze ED (1997) A global budget for fine root biomass, surface area, and nutrient contents. *Proceedings of the National Academy of Sciences USA* 94: 7362–7366.
- Jacobson S & Gustafsson L (2001) Effects on ground vegetation of the application of wood ash to a Swedish Scots pine stand. *Basic and Applied Ecology* 2: 233–241.
- Karltun E, Saarsalmi A, Ingerslev M, Mandre M, Andersson S, Gaitnieks T, Ozolinčius R, Varnagiryte-Kabasinskiene I (2008) Wood ash recycling – possibilities and risks. In: Röser *et al.* (eds.) *Sustainable use of forest biomass for energy: a synthesis with focus on the Baltic and Nordic region*. *Managing Forest Ecosystems* 12: 79–108.
- Kaunisto S & Sarjala T (2003) Foliar potassium concentrations of bilberry, bog bilberry and downy birch as indicators of potassium nutrition of Scots pine on a drained peatland. *Silva Fennica* 37(3): 325–332.
- Kepanen A, Lodenius M, Tulisalo & Hartikainen H (2005) Effects of different wood ashes on the solubility of cadmium in two boreal forest soils. *Boreal Environmental Research* 10: 135–143.
- Kirkinen J, Minkkinen K, Penttilä T, Kojola S, Sievänen R, Alm J, Saarnio S, Silvan N, Laine J & Savolainen I (2007) Greenhouse impact due to different peat fuel utilisation chains in Finland – a life-cycle approach. *Boreal Environmental Research* 12: 211–223.
- Koyama A & Tsuyuzaki S (2010) Effects of sedge and cottongrass tussocks on plant establishment patterns in a post-mined peatland, northern Japan. *Wetlands Ecology and Management* 18: 135–148.

- Kubin E & Siira J (1979) On the suitability of the phenyl hypochlorite procedure for the determination of total nitrogen from plant and soil samples. *Aquilo, Serie Botanica* 17: 11–17.
- Laiho R, Sallantausta T & Laine J (1999) The effect of forestry drainage on vertical distributions of major plant nutrients in peat soils. *Plant and Soil* 207: 169–181.
- Lang S, Cornelissen S, Klahn T, van Logtestijn R, Broekman R, Schweikert W & Aerts R (2009) An experimental comparison of chemical traits and litter decomposition rates in a diverse range of subarctic bryophyte, lichen and vascular plant species. *Journal of Ecology* 97: 886–900.
- Lavoie C & Rochefort L (1996) The natural revegetation of harvested peatland in southern Québec: a spatial and dendroecological analysis. *Écoscience* 3(1): 101–111.
- Lavoie C, Saint-Louis A & Lachance D (2005) Vegetation dynamics on an abandoned vacuum-mined peatland: 5 years of monitoring. *Wetlands Ecology and Management* 13: 621–633.
- Lefèvre I, Marchal G, Corréal E, Zanuzzi A & Lutts S (2009) Variation in response to heavy metals during vegetative growth in *Dorycnium pentaphyllum* Scop. *Plant Growth Regulation* 59: 1–11.
- Lehtonen EM & Tikkanen E (1986) Turvetuhkan vaikutus maahan sekä vesipajun (*Salix cv. aquatica*) ravinnetalouteen ja kasvuun turpeentuotannosta vapautuneella suolla. Summary: Effect of peat ash on soil properties and growth on willow (*Salix cv. Aquatica*) at an abandoned peat production area. Research Institute of Northern Finland. University of Oulu. C 69: 1–100.
- Leinonen A (2010) Turpeen tuotanto ja käyttö. Yhteenveto selvityksistä. VTT Tiedotteita – Research Notes 2550.
- Lepp NW & Roberts MJ (1977) Some effects of Cadmium on growth of Bryophytes. *The Bryologist* 80(3): 533–536.
- Lodenius M (2003) Cadmium concentrations in a boreal forest ecosystem after application of wood ash. *Bulletin of Environmental Contamination and Toxicology* 71: 776–781.
- Lumme I (1988) Early effects of peat ash on growth and mineral nutrition of the silver birch (*Betula pendula*) on a mined peatland. *Silva Fennica* 22(2): 99–112.
- Mahmood S, Finlay RG, Fransson A-M & Wallander H (2003) Effects of hardened wood ash on microbial activity, plant growth and nutrient uptake by ectomycorrhizal spruce seedlings. *FEMS Microbiology Ecology* 43: 121–131.
- Mälkönen E (1974) Annual primary production and nutrient cycle in some Scots pine stands. *Communications Institutii Forestalis Fenniae* 84.5: 1–87.
- Mandre M, Pärn H, Klõšeiko J, Ingerslev M, Stupak I, Kört M & Paasrand K (2010) Use of biofuel ashes for fertilization of *Betula pendula* seedlings on nutrient-poor peat soil. *Biomass and Bioenergy* 34: 1384–1392.
- Marschner H (1995) Mineral nutrition of higher plants. London, Academic Press.
- Ministry of Agriculture and Forestry (2007) Fertilizer product Decree 12/07. Helsinki.
- Ministry of Agriculture and Forestry (2009) Decree 19/09 for changing the Fertilizer product Decree 12/07. Helsinki.

- Moilanen M, Silfverberg K & Hokkanen TJ (2002) Effects of wood-ash on tree growth, vegetation and substrate quality of a drained mire: a case study. *Forest Ecology and Management* 171: 321–338.
- Moilanen M, Fritze H, Nieminen M, Piirainen S, Issakainen J & Piispanen J (2006) Does wood ash application increase heavy metal accumulation in forest berries and mushrooms? *Forest Ecology and Management* 226: 153–160.
- Motiva (2007) Biopolttoaineiden tuhkaa metsälannoitteeksi. Helsinki. (Booklet in Finnish available at: [www.motiva.fi](http://www.motiva.fi))
- Muukkonen P & Mäkipää R (2006) Empirical biomass models of understorey vegetation in boreal forests according to stand and site attributes. *Boreal Environment Research* 11: 355–369.
- Nieminen M, Piirainen S & Moilanen M (2005) Release of mineral nutrients and heavy metals from wood and peat ash fertilizers: Field studies in Finnish forest soils. *Scandinavian Journal of Forest Research* 20: 146–153.
- Nyrönen T (1996) Peat production. In *Global Peat Resources*. Edited by Lappalainen, E. International Peat Society, Jyväskylä, Finland : 315–318.
- Official Statistics of Finland (2009) Energy consumption [e-publication]. ISSN=1798-6869. Helsinki: Statistics Finland. Available at: [http://www.stat.fi/til/ekul/2009/ekul\\_2009\\_2010-12-10\\_tie\\_001\\_en.html](http://www.stat.fi/til/ekul/2009/ekul_2009_2010-12-10_tie_001_en.html), cited 7.3.2011.
- Official Statistics of Finland (OSF) (2009) Energy consumption [e-publication]. ISSN=1798-6869. Helsinki: Statistics Finland. Available at: [http://www.stat.fi/til/ekul/2009/ekul\\_2009\\_2010-12-10\\_tie\\_001\\_en.html](http://www.stat.fi/til/ekul/2009/ekul_2009_2010-12-10_tie_001_en.html), cited 7.3.2011.
- Omil B, Pineiro V & Merino A (2007) Trace elements in soils and plants in temperate forest plantations subjected to single and multiple applications of mixed wood ash. *Science of the Total Environment* 381: 157–168.
- Österås AH, Ekvall L & Greger M (2000) Sensitivity to, and accumulation of, cadmium in *Betula pendula*, *Picea abies*, and *Pinus sylvestris* seedlings from different regions in Sweden. *Canadian Journal of Botany* 78: 1440–1449.
- Ozolinčius R, Buožytė R & Varnagirytė-Kabašinskienė I (2007) Wood ash and nitrogen influence on ground vegetation cover and chemical composition. *Biomass and Bioenergy* 31: 710–716.
- Paappanen T & Leinonen A (eds) (2005) Fuel Peat Industry in EU – Country Reports, VTT Report PRO2/P2079/05, Jyväskylä, Finland. Available at: <http://www.turbaliit.ee/index.php?picfile=21>, cited 21.9.2010.
- Paavilainen E & Päivänen J (1995) Peatland Forestry, Ecology and Principles. *Ecological Studies*, vol. 111. Springer-Verlag, Berlin/Heidelberg/New York.
- Päivänen J (2007) Suot ja suometsät – järkevän käytön perusteet. Metsäkustannus Oy, Hämeenlinna.
- Palviainen M, Finér L, Mannerkoski H, Piirainen S & Starr M (2005) Changes in the above- and below-ground biomass and nutrient pools of ground vegetation after clear-cutting of a mixed boreal forest. *Plant and Soil* 275: 157–167.

- Pasanen J, Louekari K & Malm J (2001) Cadmium in wood ash used as fertilizer in forestry: risks to the environment and human health. Ministry of Agriculture and Forestry, publications 5/2001. Helsinki.
- Pitman RM (2006) Wood ash use in forestry – a review of the environmental impacts. *Forestry* 79(5): 563–588. doi:10.1093/forestry/cpl041
- Poikolainen J, Kubin E, Piispanen J & Karhu J (2004) Atmospheric heavy metal deposition in Finland during 1985–2000 using mosses as bioindicators. *The Science of the Total Environment* 318: 171–185.
- Rees DC & Juday GP (2002) Plant species diversity on logged versus burned sites in central Alaska. *Forest Ecology and Management* 155: 291–302.
- Reimann C, Koller F, Frengstad B, Kashulina G, Niskavaara H & Englmaier P (2001) Comparison of the element composition in several plant species and their substrate from a 1 500 000-km<sup>2</sup> area in Northern Europe. *The Science of the Total Environment* 278: 87–112.
- Reimann C, Ottesen RT, Andersson M, Arnoldussen A, Koller F & Englmaier P (2008) Element levels in birch and spruce wood ashes – green energy? *Science of the Total Environment* 393: 191–197.
- Reyes O & Casal M (2004) Effects of forest fire ash on germination and early growth of four pinus species. *Plant Ecology* 175: 81–89.
- Rikala R & Jozefek HJ (1990) Effect of dolomite lime and wood ash on peat substrate and development of tree seedlings. *Silva Fennica* 24: 323–334.
- Roxburg SH, Wood SW, Mackey BG, Woldendorp G & Gibbons P (2006) Assessing the carbon sequestration potential of managed forests: a case study from temperate Australia. *Journal of Applied Ecology* 43: 1149–59.
- Rydin H & Jeglum J (2006) *The biology of peatlands*. Oxford University Press Inc., New York.
- Ryömä R & Laaka-Lindberg S (2005) Bryophyte recolonization on burnt soil and logs. *Scandinavian Journal of Forest Research* 20(6): 5–16.
- Saarsalmi A, Kukkola M, Moilanen M & Arola M (2006) Long-term effects of ash and N fertilisation on stand growth, tree nutrient status and soil chemistry in a Scots pine stand. *Forest Ecology and Management* 235: 116.
- Salonen V (1987) Relationship between the seed rain and the establishment of vegetation in two areas abandoned after peat harvesting. *Holarctic Ecology* 10: 171–174.
- Salonen V (1992) Plant colonization of harvested peat surfaces. *Biological research reports from the University of Jyväskylä* 29: 1–29.
- Salonen V (1994) Revegetation of harvested peat surfaces in relation to substrate quality. *Journal of Vegetation Science* 5: 403–408.
- Salonen V & Laaksonen M (1994) Effects of fertilization, liming, watering and tillage on plant colonization of bare peat surfaces. *Annales Botanici Fennici* 31: 29–36.
- Salonen V, Penttinen A & Särkkä A (1992) Plant colonization of a bare peat surface: population changes and spatial patterns. *Journal of Vegetation Sciences* 3(1): 113–118.
- Sarjala T & Kaunisto S (2002) Potassium nutrition and free polyamines of *Betula pendula* Roth and *Betula pubescens* Ehrh. *Plant and Soil* 238: 141–149.

- Selin P (1999) Turvevarojen teollinen käyttö ja suopohjien hyödyntäminen Suomessa. Jyväskylä Studies in Biological and Environmental Science 79.
- Silfverberg K (1996) Nutrient status and development of tree stands and vegetation on ash-fertilized drained peatlands in Finland. The Finnish Forest Research Institute, Research paper 588.
- Silfverberg K & Huikari O (1985) Tuhkalannoitus metsäojitetuilla turvemailla. Summary: Wood-ash fertilization on drained peatlands. *Folia Forestalia* 633: 1–25.
- Solla-Gullon F, Santalla M, Rodriguez-Soalleiro RJ & Merino A (2006) Nutritional status and growth of a young *Pseudotsuga menziesii* plantation in a temperate region after application of wood-bark ash. *Forest Ecology and Management* 237: 312–321.
- Tuittila E-S & Komulainen V-M (1995) Vegetation and CO<sub>2</sub> balance in an abandoned harvested peatland in Aitoneva, southern Finland. *Suo* 46 (3): 69–80.
- Tuittila ES, Komulainen VM, Vasander H & Laine J (1999) Restored cut-away peatland as a sink for atmospheric CO<sub>2</sub>. *Oecologia* 120: 563–574.
- Tuittila ES, Rita H, Vasander H & Laine J (2000) Vegetation patterns around *Eriophorum vaginatum* L. tussocks in a cut-away peatland in southern Finland. *Canadian Journal of Botany* 78: 47–58.
- Turesky MR, Mack MC, Hollingsworth TN & Harden JW (2010) The role of mosses in ecosystem succession and function in Alaska's boreal forest. *Canadian Journal of Forest Research* 40: 1237–1264.
- Tyler G (1990) Bryophytes and heavy metals: a literature review. *Botanical Journal of the Linnean Society* 104: 231–253.
- Vasander H, Tuittila ES, Lode E, Lundin L, Ilomets M, Sallantausta T, Heikkilä R, Pitkänen ML & Laine J (2003) Status and restoration of peatlands in northern Europe. *Wetlands Ecology and Management* 11: 51–63.
- Weber A, Karsisto M, Leppänen R, Sundman V & Skujiņš J (1985) Microbial activities in a histosol: Effects of wood ash and NPK fertilizers. *Soil Biology and Biochemistry* 17(3): 291–296.
- Wind-Mulder H, Rochefort L & Vitt DH (1996) Water and peat chemistry comparisons of natural and post-harvested peatlands across Canada and their relevance to peatland restoration. *Ecological Engineering* 7: 161–181.
- Zar JH (1984) *Biostatistical analysis*. 2nd ed. New Jersey: Prentice-Hall International Inc.
- Zuloaga-Aguilar S, Briones O & Orozco-Segovia A (2011) Seed germination of montane forest in response to ash, smoke and heat shock in Mexico. *Acta Oecologica* 37(3): 256–262.



## Original articles

The thesis is based on the following articles, which are referred to in the text by their Roman numerals:

- I Huotari N, Tillman-Sutela E, Kauppi A & Kubin E (2007) Fertilization ensures rapid formation of ground vegetation on cut-away peatlands. *Canadian Journal of Forest Research* 37: 874–883.
- II Huotari N, Tillman-Sutela E, Pasanen J & Kubin E (2008) Ash-fertilization improves germination and early establishment of birch (*Betula pubescens* Ehrh.) seedlings on a cut-away peatland. *Forest Ecology and Management* 255: 2870–2875.
- III Huotari N, Tillman-Sutela E & Kubin E (2009) Ground vegetation exceeds tree seedlings in early biomass production and carbon stock on an ash-fertilized cut-away peatland. *Biomass and Bioenergy* 33: 1108–1115.
- IV Huotari N, Tillman-Sutela E & Kubin E (2011) Ground vegetation has a major role in element dynamics in ash-fertilized cut-away peatland. *Forest Ecology and Management* 261: 2081–2088.

Reprinted with permission from NRC Research Press (I) and Elsevier (II, III, IV).

Original publications are not included in the electronic version of the dissertation.



560. Siirtola, Antti (2010) Algorithmic multiparameterised verification of safety properties : process algebraic approach
561. Lappi, Anna-Kaisa (2010) Mechanisms of protein disulphide isomerase catalyzed disulphide bond formation
562. Sarala, Marian (2010) Elongation of Scots pine seedlings under blue light depletion
563. Vance, Anthony (2010) Why do employees violate is security policies? : insights from multiple theoretical perspectives
564. Karppinen, Katja (2010) Biosynthesis of hypericins and hyperforins in *Hypericum perforatum* L. (St. John's wort) – precursors and genes involved
565. Louhi, Pauliina (2010) Responses of brown trout and benthic invertebrates to catchment-scale disturbance and in-stream restoration measures in boreal river systems
566. Hekkala, Riitta (2011) The many facets of an inter-organisational information system project as perceived by the actors
567. Niittyvuopio, Anne (2011) Adaptation to northern conditions at flowering time genes in *Arabidopsis lyrata* and *Arabidopsis thaliana*
568. Leppälä, Johanna (2011) The genetic basis of incipient speciation in *Arabidopsis lyrata*
569. Kivelä, Sami, Mikael (2011) Evolution of insect life histories in relation to time constraints in seasonal environments : polymorphism and clinal variation
570. Kaartinen, Salla (2011) Space use and habitat selection of the wolf (*Canis lupus*) in human-altered environment in Finland
571. Hilli, Sari (2011) Carbon fractions and stocks in organic layers in boreal forest soils—impacts of climatic and nutritional conditions
572. Jokipii-Lukkari, Soile (2011) Endogenous haemoglobins and heterologous *Vitreoscilla* haemoglobin in hybrid aspen
573. Vuosku, Jaana (2011) A matter of life and death - polyamine metabolism during zygotic embryogenesis of pine
574. Petsalo, Aleksanteri (2011) Development of LC/MS techniques for plant and drug metabolism studies
575. Leppälä, Mirva (2011) Successional changes in vegetation and carbon dynamics during boreal mire development

Book orders:

Granum: Virtual book store  
<http://granum.uta.fi/granum/>

S E R I E S E D I T O R S

**A**  
**SCIENTIAE RERUM NATURALIUM**

*Senior Assistant Jorma Arhippainen*

**B**  
**HUMANIORA**

*Lecturer Santeri Palviainen*

**C**  
**TECHNICA**

*Professor Hannu Heusala*

**D**  
**MEDICA**

*Professor Olli Vuolteenaho*

**E**  
**SCIENTIAE RERUM SOCIALIUM**

*Senior Researcher Eila Estola*

**F**  
**SCRIPTA ACADEMICA**

*Director Sinikka Eskelinen*

**G**  
**OECONOMICA**

*Professor Jari Juga*

EDITOR IN CHIEF

*Professor Olli Vuolteenaho*

PUBLICATIONS EDITOR

*Publications Editor Kirsti Nurkkala*

ISBN 978-951-42-9531-7 (Paperback)

ISBN 978-951-42-9532-4 (PDF)

ISSN 0355-3191 (Print)

ISSN 1796-220X (Online)

