

Cochrane Database of Systematic Reviews

Reduction in saturated fat intake for cardiovascular disease (Review)

Hooper L, Martin N, Jimoh OF, Kirk C, Foster E, Abdelhamid AS

Hooper L, Martin N, Jimoh OF, Kirk C, Foster E, Abdelhamid AS. Reduction in saturated fat intake for cardiovascular disease. *Cochrane Database of Systematic Reviews* 2020, Issue 5. Art. No.: CD011737. DOI: 10.1002/14651858.CD011737.pub2.

www.cochranelibrary.com



TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
SUMMARY OF FINDINGS	3
BACKGROUND	6
OBJECTIVES	7
METHODS	7
RESULTS	11
Figure 1	12
Figure 2	14
Figure 3	17
Figure 4	18
Figure 5	19
Figure 6	22
DISCUSSION	23
AUTHORS' CONCLUSIONS	27
ACKNOWLEDGEMENTS	27
REFERENCES	29
CHARACTERISTICS OF STUDIES	65
DATA AND ANALYSES	118
Analysis 1.1. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 1: ALL-CAUSE MORTALITY	128
Analysis 1.2. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 2: All-cause mortality, SA low summary risk of bias	128
Analysis 1.3. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 3: All-cause mortality, SA aim to reduce SFA	129
Analysis 1.4. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 4: All-cause mortality, SA statistically significant SFA reduction	129
Analysis 1.5. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 5: All-cause mortality, SA TC reduction	130
Analysis 1.6. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 6: All-cause mortality, SA excluding WHI	130
Analysis 1.7. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 7: All-cause mortality, SA Mantel-Haenszel fixed-effect	131
Analysis 1.8. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 8: All-cause mortality, SA Peto fixed- effect	131
Analysis 1.9. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 9: All-cause mortality, subgroup by any substitution	132
Analysis 1.10. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 10: All-cause mortality, subgroup by main substitution	133
Analysis 1.11. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 11: All-cause mortality, subgroup by duration	135
Analysis 1.12. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 12: All-cause mortality, subgroup by	136
Analysis 1.13. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 13: All-cause mortality, subgroup by SFA change	138
Analysis 1.14. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 14: All-cause mortality, subgroup by	139
Analysis 1.15. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 15: All-cause mortality, subgroup by CVD risk	140
Analysis 1.16. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 16: All-cause mortality, subgroup by	141
Analysis 1.17. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 17: All-cause mortality, subgroup decade	142
Analysis 1.18. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 18: CARDIOVASCULAR MORTALITY	143



Analysis 1.19. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 19: CVD mortality, SA low summary risk of bias	143
Analysis 1.20. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 20: CVD mortality, SA aim to reduce SFA	144
Analysis 1.21. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 21: CVD mortality, SA statistically significant SFA reduction	144
Analysis 1.22. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 22: CVD mortality, SA TC reduction	145
Analysis 1.23. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 23: CVD mortality, SA excluding WHI	145
Analysis 1.24. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 24: CVD mortality, SA Mantel-Haenszel fixed-effect	146
Analysis 1.25. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 25: CVD mortality, SA Peto fixed-effect	146
Analysis 1.26. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 26: CVD mortality, subgroup by any substitution	147
Analysis 1.27. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 27: CVD mortality, subgroup by main substitution	148
Analysis 1.28. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 28: CVD mortality, subgroup by duration	150
Analysis 1.29. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 29: CVD mortality, subgroup by baseline SFA	151
Analysis 1.30. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 30: CVD mortality, subgroup by SFA change	153
Analysis 1.31. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 31: CVD mortality, subgroup by sex	154
Analysis 1.32. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 32: CVD mortality, subgroup by CVD risk	155
Analysis 1.33. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 33: CVD mortality, subgroup by TC reduction	156
Analysis 1.34. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 34: CVD mortality, subgroup decade of publication	157
Analysis 1.35. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 35: COMBINED CARDIOVASCULAR EVENTS	158
Analysis 1.36. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 36: CVD events, SA low summary risk of bias	158
Analysis 1.37. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 37: CVD events, SA aim to reduce SFA	159
Analysis 1.38. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 38: CVD events, SA statistically significant SFA reduction	159
Analysis 1.39. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 39: CVD events, SA TC reduction	160
Analysis 1.40. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 40: CVD events, SA excluding WHI	160
Analysis 1.41. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 41: CVD events, SA Mantel-Haenszel fixed-effect	161
Analysis 1.42. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 42: CVD events, SA Peto fixed-effect	161
Analysis 1.43. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 43: CVD events, SA excluding trials with additional interventions	162
Analysis 1.44. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 44: CVD events, subgroup by any substitution	163
Analysis 1.45. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 45: CVD events, subgroup by main substitution	164
Analysis 1.46. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 46: CVD events, subgroup by duration .	166
Analysis 1.47. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 47: CVD events, subgroup by baseline SFA	167
Analysis 1.48. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 48: CVD events, subgroup by SFA change	169
Analysis 1.49. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 49: CVD events, subgroup by sex	170
Analysis 1.50. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 50: CVD events, subgroup by CVD risk	171
Analysis 1.51. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 51: CVD events, subgroup by TC 2 reduction	172



Analysis 1.52. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 52: CVD events, subgroup decade of publication	173
Analysis 2.1. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 1: MYOCARDIAL INFARCTION	189
Analysis 2.2. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 2: MI, SA by low summary risk of bias	189
Analysis 2.3. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 3: MI, SA aim to reduce SFA	190
Analysis 2.4. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 4: MI, SA statistically significant SFA reduction	190
Analysis 2.5. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 5: MI, SA by TC reduction	191
Analysis 2.6. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 6: MI, SA excluding WHI	191
Analysis 2.7. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 7: MI, SA Mantel-Haenszel fixed- effect	192
Analysis 2.8. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 8: MI, SA Peto fixed-effect	192
Analysis 2.9. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 9: MI, subgroup by any substitution	193
Analysis 2.10. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 10: MI, subgroup by main substitution	194
Analysis 2.11. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 11: MI, subgroup by duration	195
Analysis 2.12. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 12: MI, subgroup by baseline SFA	196
Analysis 2.13. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 13: MI, subgroup by SFA change .	197
Analysis 2.14. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 14: MI, subgroup by sex	198
Analysis 2.15. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 15: MI, subgroup by CVD risk	199
Analysis 2.16. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 16: MI, subgroup by TC reduction	200
Analysis 2.17. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 17: MI, subgroup decade of publication	201
Analysis 2.18. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 18: NON-FATAL MYOCARDIAL INFARCTION	202
Analysis 2.19. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 19: Non-fatal MI, SA by low summary risk of bias	202
Analysis 2.20. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 20: Non-fatal MI, SA aim to reduce SFA	203
Analysis 2.21. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 21: Non-fatal MI, SA statistically significant SFA reduction	203
Analysis 2.22. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 22: Non-fatal MI, SA by TC reduction	204
Analysis 2.23. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 23: Non-fatal MI, SA excluding WHI	204
Analysis 2.24. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 24: Non-fatal MI, SA Mantel- Haenszel fixed-effect	205
Analysis 2.25. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 25: Non-fatal MI, SA Peto fixed- effect	205
Analysis 2.26. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 26: Non-fatal MI, subgroup by any substitution	206
Analysis 2.27. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 27: Non-fatal MI, subgroup by main substitution	207
Analysis 2.28. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 28: Non-fatal MI, subgroup by duration	208
Analysis 2.29. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 29: Non-fatal MI, subgroup by baseline SFA	209
Analysis 2.30. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 30: Non-fatal MI, subgroup by SFA change	210
Analysis 2.31. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 31: Non-fatal MI, subgroup by sex	211
Analysis 2.32. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 32: Non-fatal MI, subgroup by	212
CVD risk	



Analysis 2.33. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 33: Non-fatal MI, subgroup by TC reduction	213
Analysis 2.34. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 34: Non-fatal MI, subgroup decade of publication	214
Analysis 2.35. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 35: STROKE	215
Analysis 2.36. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 36: Stroke, SA by low summary risk of bias	215
Analysis 2.37. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 37: Stroke, SA aim to reduce SFA .	216
Analysis 2.38. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 38: Stroke, SA statistically significant SFA reduction	216
Analysis 2.39. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 39: Stroke, SA by TC reduction	217
Analysis 2.40. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 40: Stroke, SA excluding WHI	217
Analysis 2.41. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 41: Stroke, SA Mantel-Haenszel fixed-effect	218
Analysis 2.42. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 42: Stroke, SA Peto fixed-effect	218
Analysis 2.43. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 43: Stroke, subgroup by any substitution	219
Analysis 2.44. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 44: Stroke, subgroup by main substitution	220
Analysis 2.45. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 45: Stroke, subgroup by duration	221
Analysis 2.46. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 46: Stroke, subgroup by baseline SFA	222
Analysis 2.47. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 47: Stroke, subgroup by SFA change	223
Analysis 2.48. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 48: Stroke, subgroup by sex	224
Analysis 2.49. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 49: Stroke, subgroup by CVD risk	225
Analysis 2.50. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 50: Stroke, subgroup by TC reduction	226
Analysis 2.51. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 51: Stroke, subgroup decade of publication	227
Analysis 2.52. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 52: CORONARY HEART DISEASE MORTALITY	228
Analysis 2.53. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 53: CHD mortality, SA by low summary risk of bias	228
Analysis 2.54. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 54: CHD mortality, SA aim to reduce SFA	229
Analysis 2.55. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 55: CHD mortality, SA statistically significant SFA reduction	229
Analysis 2.56. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 56: CHD mortality, SA by TC reduction	230
Analysis 2.57. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 57: CHD mortality, SA excluding WHI	230
Analysis 2.58. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 58: CHD mortality, SA Mantel- Haenszel fixed-effect	231
Analysis 2.59. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 59: CHD mortality, SA Peto fixed- effect	231
Analysis 2.60. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 60: CHD mortality, subgroup by any substitution	232
Analysis 2.61. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 61: CHD mortality, subgroup by main substitution	233
Analysis 2.62. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 62: CHD mortality, subgroup by duration	234
Analysis 2.63. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 63: CHD mortality, subgroup by baseline SFA	235
Analysis 2.64. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 64: CHD mortality, subgroup by SFA change	236

Reduction in saturated fat intake for cardiovascular disease (Review)

Copyright @ 2020 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.



Analysis 2.65. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 65: CHD mortality, subgroup by sex	237
Analysis 2.66. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 66: CHD mortality, subgroup by CVD risk	238
Analysis 2.67. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 67: CHD mortality, subgroup by TC reduction	239
Analysis 2.68. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 68: CHD mortality, subgroup decade of publication	240
Analysis 2.69. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 69: CORONARY HEART DISEASE EVENTS	241
Analysis 2.70. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 70: CHD events, SA by low summary risk of bias	241
Analysis 2.71. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 71: CHD events, SA excluding WHI	242
Analysis 2.72. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 72: CHD events, SA statistically significant SFA reduction	242
Analysis 2.73. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 73: CHD events, SA by TC reduction	243
Analysis 2.74. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 74: CHD events, SA aim to reduce SFA	243
Analysis 2.75. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 75: CHD events, SA Mantel- Haenszel fixed-effect	244
Analysis 2.76. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 76: CHD events, SA Peto fixed-effect	244
Analysis 2.77. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 77: CHD events, subgroup by any substitution	245
Analysis 2.78. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 78: CHD events, subgroup by main substitution	246
Analysis 2.79. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 79: CHD events, subgroup by duration	247
Analysis 2.80. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 80: CHD events, subgroup by baseline SFA	248
Analysis 2.81. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 81: CHD events, subgroup by SFA change	249
Analysis 2.82. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 82: CHD events, subgroup by sex	250
Analysis 2.83. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 83: CHD events, subgroup by CVD risk	251
Analysis 2.84. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 84: CHD events, subgroup by TC reduction	252
Analysis 2.85. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 85: CHD events, subgroup decade of publication	253
Analysis 2.86. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 86: DIABETES DIAGNOSES	254
Analysis 3.1. Comparison 3: SFA reduction vs usual diet - secondary blood outcomes, Outcome 1: Total cholesterol, mmol/L	259
Analysis 3.2. Comparison 3: SFA reduction vs usual diet - secondary blood outcomes, Outcome 2: TC, mmol/L, subgroup by any replacement	260
Analysis 3.3. Comparison 3: SFA reduction vs usual diet - secondary blood outcomes, Outcome 3: TC, mmol/L, subgroup by main replacement	261
Analysis 3.4. Comparison 3: SFA reduction vs usual diet - secondary blood outcomes, Outcome 4: LDL cholesterol, mmol/L	261
Analysis 3.5. Comparison 3: SFA reduction vs usual diet - secondary blood outcomes, Outcome 5: LDL, mmol/L, subgroup by any replacement	262
Analysis 3.6. Comparison 3: SFA reduction vs usual diet - secondary blood outcomes, Outcome 6: LDL, mmol/L, subgroup by main replacement	263
Analysis 3.7. Comparison 3: SFA reduction vs usual diet - secondary blood outcomes, Outcome 7: HDL cholesterol, mmol/L	263
Analysis 3.8. Comparison 3: SFA reduction vs usual diet - secondary blood outcomes, Outcome 8: HDL, mmol/L, subgroup by	264
any replacement	



Analysis 3.9. Comparison 3: SFA reduction vs usual diet - secondary blood outcomes, Outcome 9: HDL, mmol/L, subgroup by main replacement	265
Analysis 3.10. Comparison 3: SFA reduction vs usual diet - secondary blood outcomes, Outcome 10: Triglycerides, mmol/L	265
Analysis 3.11. Comparison 3: SFA reduction vs usual diet - secondary blood outcomes, Outcome 11: TG, mmol/L, subgroup by any replacement	266
Analysis 3.12. Comparison 3: SFA reduction vs usual diet - secondary blood outcomes, Outcome 12: TG, mmol/L, subgroup by main replacement	267
Analysis 3.13. Comparison 3: SFA reduction vs usual diet - secondary blood outcomes, Outcome 13: total cholesterol /HDL ratio	267
Analysis 3.14. Comparison 3: SFA reduction vs usual diet - secondary blood outcomes, Outcome 14: TC /HDL ratio, subgroup by any replacement	268
Analysis 3.15. Comparison 3: SFA reduction vs usual diet - secondary blood outcomes, Outcome 15: TC /HDL ratio, subgroup by main replacement	269
Analysis 3.16. Comparison 3: SFA reduction vs usual diet - secondary blood outcomes, Outcome 16: LDL /HDL ratio	269
Analysis 3.17. Comparison 3: SFA reduction vs usual diet - secondary blood outcomes, Outcome 17: Lp(a), mmol/L	269
Analysis 3.18. Comparison 3: SFA reduction vs usual diet - secondary blood outcomes, Outcome 18: Lp(a), mmol/L, subgroup by any replacement	270
Analysis 3.19. Comparison 3: SFA reduction vs usual diet - secondary blood outcomes, Outcome 19: Lp(a), mmol/L, subgroup by main replacement	271
Analysis 3.20. Comparison 3: SFA reduction vs usual diet - secondary blood outcomes, Outcome 20: Insulin sensitivity	271
Analysis 4.1. Comparison 4: SFA reduction vs usual diet - secondary outcomes including potential adverse effects, Outcome 1: Cancer diagnoses	272
Analysis 4.2. Comparison 4: SFA reduction vs usual diet - secondary outcomes including potential adverse effects, Outcome 2: Cancer deaths	273
Analysis 4.3. Comparison 4: SFA reduction vs usual diet - secondary outcomes including potential adverse effects, Outcome 3: Weight, kg	273
Analysis 4.4. Comparison 4: SFA reduction vs usual diet - secondary outcomes including potential adverse effects, Outcome 4: BMI, kg/m2	273
Analysis 4.5. Comparison 4: SFA reduction vs usual diet - secondary outcomes including potential adverse effects, Outcome 5: Systolic Blood Pressure, mmHg	274
Analysis 4.6. Comparison 4: SFA reduction vs usual diet - secondary outcomes including potential adverse effects, Outcome 6: Diastolic Blood Pressure, mmHg	274
Analysis 4.7. Comparison 4: SFA reduction vs usual diet - secondary outcomes including potential adverse effects, Outcome 7: Quality of Life	274
ADDITIONAL TABLES	274
APPENDICES	282
WHAT'S NEW	285
HISTORY	285
CONTRIBUTIONS OF AUTHORS	286
DECLARATIONS OF INTEREST	286
SOURCES OF SUPPORT	286
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	286
INDEX TERMS	287



[Intervention Review]

Reduction in saturated fat intake for cardiovascular disease

Lee Hooper¹, Nicole Martin², Oluseyi F Jimoh¹, Christian Kirk¹, Eve Foster¹, Asmaa S Abdelhamid¹

¹Norwich Medical School, University of East Anglia, Norwich, UK. ²Institute of Health Informatics Research, University College London, London, UK

Contact address: Lee Hooper, l.hooper@uea.ac.uk.

Editorial group: Cochrane Heart Group. **Publication status and date:** New search for studies and content updated (no change to conclusions), published in Issue 5, 2020.

Citation: Hooper L, Martin N, Jimoh OF, Kirk C, Foster E, Abdelhamid AS. Reduction in saturated fat intake for cardiovascular disease. *Cochrane Database of Systematic Reviews* 2020, Issue 5. Art. No.: CD011737. DOI: 10.1002/14651858.CD011737.pub2.

Copyright © 2020 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

Reducing saturated fat reduces serum cholesterol, but effects on other intermediate outcomes may be less clear. Additionally, it is unclear whether the energy from saturated fats eliminated from the diet are more helpfully replaced by polyunsaturated fats, monounsaturated fats, carbohydrate or protein.

Objectives

To assess the effect of reducing saturated fat intake and replacing it with carbohydrate (CHO), polyunsaturated (PUFA), monounsaturated fat (MUFA) and/or protein on mortality and cardiovascular morbidity, using all available randomised clinical trials.

Search methods

We updated our searches of the Cochrane Central Register of Controlled Trials (CENTRAL), MEDLINE (Ovid) and Embase (Ovid) on 15 October 2019, and searched Clinicaltrials.gov and WHO International Clinical Trials Registry Platform (ICTRP) on 17 October 2019.

Selection criteria

Included trials fulfilled the following criteria: 1) randomised; 2) intention to reduce saturated fat intake OR intention to alter dietary fats and achieving a reduction in saturated fat; 3) compared with higher saturated fat intake or usual diet; 4) not multifactorial; 5) in adult humans with or without cardiovascular disease (but not acutely ill, pregnant or breastfeeding); 6) intervention duration at least 24 months; 7) mortality or cardiovascular morbidity data available.

Data collection and analysis

Two review authors independently assessed inclusion, extracted study data and assessed risk of bias. We performed random-effects metaanalyses, meta-regression, subgrouping, sensitivity analyses, funnel plots and GRADE assessment.

Main results

We included 15 randomised controlled trials (RCTs) (16 comparisons, ~59,000 participants), that used a variety of interventions from providing all food to advice on reducing saturated fat. The included long-term trials suggested that reducing dietary saturated fat reduced the risk of combined cardiovascular events by 21% (risk ratio (RR) 0.79; 95% confidence interval (CI) 0.66 to 0.93, 11 trials, 53,300 participants of whom 8% had a cardiovascular event, I² = 65%, GRADE moderate-quality evidence). Meta-regression suggested that greater reductions in saturated fat (reflected in greater reductions in serum cholesterol) resulted in greater reductions in risk of CVD events, explaining most heterogeneity between trials. The number needed to treat for an additional beneficial outcome (NNTB) was 56 in primary prevention trials, so 56 people need to reduce their saturated fat intake for ~four years for one person to avoid experiencing a CVD event. In secondary prevention trials, the NNTB was 32. Subgrouping did not suggest significant differences between replacement of saturated fat calories with polyunsaturated fat or carbohydrate, and data on replacement with monounsaturated fat and protein was very limited.



We found little or no effect of reducing saturated fat on all-cause mortality (RR 0.96; 95% CI 0.90 to 1.03; 11 trials, 55,858 participants) or cardiovascular mortality (RR 0.95; 95% CI 0.80 to 1.12, 10 trials, 53,421 participants), both with GRADE moderate-quality evidence.

There was little or no effect of reducing saturated fats on non-fatal myocardial infarction (RR 0.97, 95% CI 0.87 to 1.07) or CHD mortality (RR 0.97, 95% CI 0.82 to 1.16, both low-quality evidence), but effects on total (fatal or non-fatal) myocardial infarction, stroke and CHD events (fatal or non-fatal) were all unclear as the evidence was of very low quality. There was little or no effect on cancer mortality, cancer diagnoses, diabetes diagnosis, HDL cholesterol, serum triglycerides or blood pressure, and small reductions in weight, serum total cholesterol, LDL cholesterol and BMI. There was no evidence of harmful effects of reducing saturated fat intakes.

Authors' conclusions

The findings of this updated review suggest that reducing saturated fat intake for at least two years causes a potentially important reduction in combined cardiovascular events. Replacing the energy from saturated fat with polyunsaturated fat or carbohydrate appear to be useful strategies, while effects of replacement with monounsaturated fat are unclear. The reduction in combined cardiovascular events resulting from reducing saturated fat did not alter by study duration, sex or baseline level of cardiovascular risk, but greater reduction in saturated fat caused greater reductions in cardiovascular events.

PLAIN LANGUAGE SUMMARY

Effect of cutting down on the saturated fat we eat on our risk of heart disease

Review question

We wanted to find out the effects on health of cutting down on saturated fat in our food (replacing animal fats and hard vegetable fats with plant oils, unsaturated spreads or starchy foods).

Background

Health guidance suggests that reducing the amount of saturated fat we eat, by cutting down on animal fats, is good for our health. We wanted to combine all available evidence to see whether following this advice leads to a reduced risk of dying or getting cardiovascular disease (heart disease or stroke).

Study characteristics

We assessed the effect of cutting down the amount of saturated fat we eat for at least two years on health outcomes including dying, heart disease and stroke. We only looked at studies of adults (18 years or older). They included men and women with and without cardiovascular disease. We did not include studies of acutely ill people or pregnant or breastfeeding women.

Key results

We found 15 studies with over 59,000 participants. The evidence is current to October 2019. The review found that cutting down on saturated fat led to a 21% reduction in the risk of cardiovascular disease (including heart disease and strokes), but had little effect on the risk of dying. The review found that health benefits arose from replacing saturated fats with polyunsaturated fat or starchy foods. The greater the decrease in saturated fat, and the more serum total cholesterol is reduced, the greater the protection from cardiovascular events. People who are currently healthy appear to benefit as much as those at increased risk of heart disease or stroke (people with high blood pressure, high serum cholesterol or diabetes, for example), and people who have already had heart disease or stroke. There was no difference in effect between men and women.

This means that, if 56 people without cardiovascular disease, or 32 people who already have cardiovascular disease, reduce their saturated fat for around 4 years, then one person will avoid a cardiovascular event (heart attack or stroke) they would otherwise have experienced.

Quality of the evidence

There is a large body of evidence assessing effects of reducing saturated fat for at least two years. These studies provide moderate-quality evidence that reducing saturated fat reduces our risk of cardiovascular disease.

SUMMARY OF FINDINGS

Summary of findings 1. Effect of reducing saturated fat compared to usual saturated fat on CVD risk in adults (note: for the full set of GRADE tables see additional tables 24 to 28)

Low saturated fat compared with usual saturated fat for CVD risk

Patient or population: people at any baseline risk of CVD

Intervention: lower saturated fat intake

Comparison: higher saturated fat intake

Settings: Any, including community-dwelling and institutions. Included RCTs were conducted in North America, Europe and Australia/New Zealand, no studies were carried out in industrialising or developing countries.

e ef- Anticipated a (95% Cl)	absolute effects	No of Partici- pants - (studies)	Quality of the evidence (GRADE)	Comments
, Risk with higher SFA intake	Risk with lower SFA in- take	()	()	
(0.90 62 per 1000	60 per 1000 (56 to 64)	55,858 (12)	⊕⊕⊕⊝ Moder- ate2,3,4,5,6	Critical importance. Reducing saturated fat intake probably makes little or no difference to all-cause mortality.
				· · · · · · · · · · · · · · · · · · ·
(0.78 19 per 1000	18 per 1000 (15 to 22)	53,421 (11)	⊕⊕⊕⊙ Moder- ate ^{2,3,4,6,7}	Critical importance. Reducing saturated fat intake probably makes little or no difference to cardiovas- cular mortality.
(0.66 84 per 1000	67 per 1000 (56 to 79)	53,300 (12)	⊕⊕⊕⊝ Moder- ate ^{4,8,9,10,11}	Critical importance. Reducing saturated fat intake probably reduces cardiovascular mortality (to a greater extent with greater cholesterol reduction).
(0.80 32 per 1000	29 per 1000 (25 to 32)	53,167 (11)	⊕⊝⊝⊝ Very Low 3,4,5,11,12	Critical importance. The effect of reducing saturated fat intake on risk of myocardial infarction is unclear as the evidence is of very low quality.
	eei- Anticipated a (95% CI) i) Risk with higher SFA intake 5 (0.90 62 per 1000 4 (0.78 19 per 1000 9 (0.66 84 per 1000 9 (0.80 32 per 1000	Product of the second secon	Perform Anticipated absolute effects (95% Cl) No of Partici- pants (studies) No of Partici- pants Partici- pants (studies) Risk with higher SFA intake Risk with lower SFA in- take No of Partici- pants (studies) 5 (0.90 62 per 1000 60 per 1000 (56 to 64) 55,858 (12) 6 (0.78 19 per 1000 18 per 1000 (15 to 22) 53,421 (11) 9 (0.66 84 per 1000 67 per 1000 (56 to 79) 53,300 (12) 9 (0.80 32 per 1000 29 per 1000 (25 to 32) 53,167 (11)	$\frac{(95\% \text{ CI})}{(95\% \text{ CI})} \xrightarrow{\text{Risk with}} \frac{\text{Risk with}}{\text{lower SFA in-take}} \xrightarrow{\text{Risk with}} \frac{\text{Risk with}}{\text{lower SFA in-take}} \xrightarrow{\text{Risk of particle}} \text{Risk of particle$

Cochrane Database of Systematic Reviews

Non-fatal MI follow-up mean duration 55 months ¹	RR 0.97 (0.87 to 1.07)	26 per 1000	25 per 1000 (23 to 28)	52,834 (8)	⊕⊕⊝⊝ Low3,4,5,6,13	Critical importance. Reducing saturated fat may have little or no effect on risk of non-fatal myocar- dial infarction.
Stroke follow-up mean duration 59 months ¹	RR 0.92 (0.68 to 1.25)	22 per 1000	20 per 1000 (15 to 27)	50,952 (7)	⊕⊝⊝⊝ Very Low 3,4,6,13,14	Critical importance. The effect of reducing saturat- ed fat on the risk of stroke is unclear as the evidence was of very low quality.
CHD mortality follow-up mean duration 65 months ¹	RR 0.97 (0.82 to 1.16)	16 per 1000	16 per 1000 (13 to 19)	53,159 (9)	⊕⊕⊝⊝ Low2,3,4,6,14	Critical importance. Reducing saturated fat intake may have little or no effect on CHD mortality.

*The risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

CI: Confidence interval; RR: Risk Ratio; CHD: coronary heart disease.

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate. **Low quality:** Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate. **Very low quality:** We are very uncertain about the estimate.

¹Minimum study duration was 24 months.

² **Risk of bias.** Limiting trials to those at low summary risk of bias also suggested little or no effect. Not downgraded.

³ **Inconsistency**. We found no important heterogeneity; $l^2 \le 30\%$. Not downgraded.

⁴ **Indirectness**. These RCTs directly assessed the effect of lower vs higher saturated fat intake on health outcomes of interest. Participants included men and women with and without CVD at baseline (also some participants with CVD risk factors like diabetes, or at risk of cancers). However, no trials included participants from developing countries. Not downgraded.

⁵ Imprecision. The 95% CI includes both no effect and a benefit. Downgraded once.

⁶ Publication bias. The funnel plot, and comparison of fixed- and random-effects meta-analyses did not suggest major small-study (publication) bias. Not downgraded.

 7 $\mathbf{Imprecision}.$ The 95% CI includes both harm and benefit. Downgraded once.

⁸ Risk of bias. Limiting trials to those at low summary risk of bias also suggested that reducing SFA reduced risk of CVD events. Not downgraded.

⁹ Inconsistency. Although heterogeneity was high, I² = 65%, this was mostly explained by the degree of cholesterol-lowering (a dose effect). Not downgraded.

 10 Imprecision. The 95% CI includes only benefit. Not downgraded.

¹¹ **Publication bias**. The funnel plot, and comparison of fixed- and random-effects meta-analyses suggested some small-study (publication) bias. Downgraded once.

¹² **Risk of bias.** Limiting trials to those at low summary risk of bias moved the RR slightly towards 1.0, suggesting little or no effect on total MI. Downgraded once.

4

Copyright © 2020 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd

Reduction in saturated fat intake for cardiovascular disease (Review)

hrane

¹³ **Risk of bias.** Limiting trials to those at low summary risk of bias moved the RR slightly away from 1.0, suggesting that reducing SFA reduces the risk of non-fatal MI. This was also seen in several other sensitivity analyses. Downgraded once.

¹⁴ Imprecision. The 95% CI includes both important benefits and important harms. Downgraded twice.

¹⁵ **Inconsistency**. Heterogeneity was high, $I^2 = 65\%$. Downgraded once.

....



BACKGROUND

In 1949, Ryle and Russell in Oxford documented a dramatic increase in coronary heart disease (CHD), and the Registrar General's Statistical Tables of 1920 to 1955 showed that there had been a 70fold increase in coronary deaths during this 35-year period (Oliver 2000; Ryle 1949). This sudden surge in coronary heart disease sparked research into its causes. A case-control study published in 1953 of 200 post-myocardial infarction patients and age-matched controls established that those with disease had higher low-density lipoprotein (LDL) cholesterol levels (Oliver 1953).

Meanwhile in 1949 in the USA, Gofman had separated lipids into lipoprotein classes through ultra centrifugation, describing the LDL as 'atherosclerogenic' (Gofman 1949). The following year Keys 1950 proposed that the concentration of plasma cholesterol was proportional to dietary saturated fatty acids (SFA) intake. This relationship was confirmed in work by Hegsted (Hegsted 1965; Hegsted 2000), who published an equation explaining the relationship in 1965 and subsequently in 2000. The equation suggests that dietary saturated fat increases serum cholesterol and so increases cardiovascular (CV) risk, while polyunsaturated fats (PUFA) reduce both. This has since been further refined:

 Δ serum cholesterol (in mg/dL) = 2.16 * Δ dietary saturated fat intake (as percentage of energy) – 1.65 * Δ dietary PUFA intake (as percentage of energy, %E) + 6.77 * Δ dietary cholesterol intake (in units of 100 mg/day) – 0.53

The Seven Countries Study compared CHD mortality in 12,000 men aged 40 to 59 in seven countries, and found positive correlations between CHD mortality and total fat intake in 1970, then in 1986 between CHD mortality and saturated fat intake (Keys 1986; Thorogood 1996). A migrant study of Japanese men living in different cultures confirmed in 1974 that men in California had the diet richest in saturated fat and cholesterol, and the highest CHD rates, those in Hawaii had intermediate saturated fat and CHD rates, and those in Japan had a diet lowest in saturated fat and cholesterol, and the least CHD (Kagan 1974; Robertson 1977). However, systematic reviews of the observational data have not confirmed these early studies. Skeaff 2009 included 28 USA and European cohorts (including 6600 CHD deaths among 280,000 participants) investigating the effects of total, saturated, monounsaturated, trans and omega-3 fats on CHD deaths and events. They found no clear relationship between total, saturated or monounsaturated fat (MUFA) intake and coronary heart disease events or deaths. There was evidence that trans fats increased both coronary heart disease events and deaths, and that total PUFAs and omega-3 fats decreased them. Intervention studies are needed to clarify cause and effect, to ensure that confounding is not hiding true relationships, or suggesting relationships where they do not exist. Trials also directly address the issue of whether altering dietary saturated fat in adults is helpful in reducing the risk of CVD in the general population and in those at high risk. Intervention trials are crucial in forming the basis of evidence-based practice in this area.

Most intervention studies have assessed effects of dietary interventions on risk factors for heart disease, and separate work ties the effect of altering these risk factors to changes in disease incidence and mortality. Systematic reviews in this area follow the same pattern. There are systematic reviews of the effect of dietary fat advice on serum lipid levels (Brunner 1997; Clarke 1997; Denke 1995; Kodama 2009; Malhotra 2014; Mensink 1992; Mensink 2003; Rees 2013; Weggemans 2001; Yu-Poth 1999), suggesting that dietary changes cause changes in serum lipids. There are also systematic reviews on the effect of lipid level alterations on CV morbidity and mortality (Briel 2009; De Caterina 2010; Law 1994; Robinson 2009; Rubins 1995; Walsh 1995), suggesting that changes in lipids do affect CVD risk. Other risk factors dealt with in a similar way are blood pressure (Bucher 1996; Law 1991; Shah 2007), body weight or fatness (Astrup 2000; Hession 2009; SIGN 1996), angiographic measurements (Marchioli 1994), antioxidant intake (Ness 1997), metabolic profile (Kodama 2009) and alcohol intake (Rimm 1996). A problem with this two-level approach is that any single dietary alteration may have effects over a wide range of risk factors for CVD. An example of this is the choice of substitution of saturated fats by carbohydrate, PUFAs, MUFAs or protein in the diet. This choice may alter lipid profile, and may also affect blood pressure, body weight, oxidative state, rate of cholesterol efflux from fibroblasts, insulin resistance, post-prandial triacylglycerol response, blood clotting factors, and platelet aggregation. There may also be further risk factors of which we are not yet aware. Evidence of beneficial effect on one risk factor does not rule out an opposite effect on another unstudied risk factor, and therefore an overall null (or harmful) effect of intervention. While understanding the effects of dietary advice on intermediate risk factors helps to ensure diets are truly altered by advice, and illuminates mechanisms, the best way of combining the effects on all of these risk factors is to not study risk factors, but to study the effects of dietary change on important outcomes, on CV morbidity and mortality, and on total mortality.

Substantial randomised controlled trial data on the effects of dietary fat on mortality and morbidity do exist and have been previously reviewed (Abdelhamid 2020; Abdelhamid 2018; Abdelhamid 2019; Brainard 2020; Brown 2019; Deane 2019; Hanson 2020; Hooper 2018; Hooper 2019; Hooper 2012). A recent very large trial, the Women's Health Initiative, that included over 2000 women with, and over 48,000 women without, CVD at baseline for over eight years (WHI 2006)) has raised many questions about both the effects of fat on health and on how we best conduct research to understand the relationship (Astrup 2011; Michels 2009; Prentice 2007; Stein 2006; Yngve 2006). We incorporated these findings into an update of a Cochrane review on dietary fat and CVD risk with a search in 2010 (Hooper 2012), finding reductions in cardiovascular events in studies that modified dietary fat, and in studies of at least two years' duration, but not in studies of fat reduction or studies with less than two years' follow-up.

Why it is important to do this review

Public health dietary advice on prevention of cardiovascular disease (CVD) has changed over time, with a focus on fat modification during the 1960s and fat reduction during the 1990s following the introduction of USA and UK dietary guidance on fat reduction, limiting saturated fat intake to 10% of energy (Harcombe 2015). In 2006, recommendations by the American Heart Association suggested that, among other dietary measures, Americans should "limit intake of saturated fat to 7% of energy, trans fat to 1% of energy, and cholesterol to 300 mg/day by choosing lean meats and vegetable alternatives, fat-free (skim) or low-fat (1% fat) dairy products and minimise intake of partially hydrogenated fats" (Lichtenstein 2006). Current American Heart Association guidelines suggest that Americans should "Aim for a dietary pattern that achieves 5% to 6% of calories from



saturated fat" and "Reduce percent of calories from saturated fat" (both graded as strong evidence on the basis of effects on serum lipids - trials with cardiovascular outcomes are not referenced or discussed, Eckel 2013). European guidance on the treatment of dyslipidaemia is similarly based on dietary effects on lipids, recommending reduction in saturated fats (ESC/EAS 2011) and referencing Mensink 2003, while the Joint British Societies' guidance on preventing CVD recommends a healthy diet including low saturated fat intake (Mach 2019), referencing a variety of evidence including several recent systematic reviews. This is reflected in UK Scientific Advisory Committee on Nutrition recommendations that "dietary reference value for saturated fats remains unchanged: the [population] average contribution of saturated fatty acids to [total] dietary energy be reduced to no more than about 10%", and that "saturated fats are substituted with unsaturated fats. More evidence is available supporting substitution with PUFA than substitution with MUFA" (SACN 2019).

Recent UK National Institute for Health and Care Excellence(NICE) guidance suggests that for people at high risk of or with CVD that they "eat a diet in which total fat intake is 30% or less of total energy intake, saturated fats are 7% or less of total energy intake, intake of dietary cholesterol is less than 300 mg/day and where possible saturated fats are replaced by monounsaturated and polyunsaturated fats". This statement was based on long-term randomised controlled trials reporting hard outcomes, and NICE separately assessed effects of high polyunsaturated diets, including four of the trials included in this review (NICE 2014).

We were interested in assessing the direct evidence from trials of the effects of reducing saturated fats, and considering what macronutrients the saturated fats were replaced by, updating Hooper 2015a. This update also supports a request from the World Health Organization Nutrition Guidance Expert Advisory Group (WHO NUGAG) to more accurately assess effects of reducing saturated fats on all-cause mortality, CV morbidity and other health outcomes, and to consider the differential effects on health outcomes of replacement of the energy from saturated fat by other fats, carbohydrates or protein.

OBJECTIVES

To assess the effect of reducing saturated fat intake and replacing it with carbohydrate (CHO), polyunsaturated (PUFA) or monounsaturated fat (MUFA) and/or protein on mortality and cardiovascular morbidity, using all available randomised clinical trials.

Additional World Health Organization Nutrition Guidance Expert Advisory Group (WHO NUGAG) specific questions included:

- 1. In adults, what is the effect in the population of reduced percentage of energy (%E) intake from saturated fatty acids (SFA) relative to higher intake for reduction in risk of noncommunicable diseases (NCDs)?
- 2. What is the effect on coronary heart disease mortality and coronary heart disease events?
- 3. What is the effect in the population of replacing SFA with polyunsaturated fats (PUFAs), monounsaturated fats (MUFAs), carbohydrates (CHO) (refined versus unrefined), protein or trans fatty acids (TFAs) relative to no replacement for reduction in risk of NCDs?

- 4. What is the effect in the population of consuming < 10%E as SFA relative to > 10%E as SFA for reduction in risk of NCDs?
- 5. What is the effect in the population of a reduction in %E from SFA from 10% in gradual increments relative to higher intake for reduction in risk of NCDs?

METHODS

Criteria for considering studies for this review

Types of studies

Randomised controlled trials only. We accepted randomisation of individuals, or of larger groups (clusters) where there were at least six of these groups randomised. We excluded studies where allocation was not truly randomised (e.g. divisions based on days of the week or first letter of the family name), or where allocation was not stated as randomised, and no further information was available from the authors.

Types of participants

We included studies of adults (18 years or older, no upper age limit) at any risk of cardiovascular disease, with or without existing cardiovascular disease, using or not using lipid-lowering medication. Participants could be of either gender, but we excluded those who were acutely ill, pregnant or lactating.

Types of interventions

We included randomised controlled trials stating an intention to reduce saturated fat (SFA) intake (by suggesting appropriate nutrient-based or food-based aims) OR which provided a general dietary aim, such as improving heart health or reducing total fat, that also achieved a statistically significant saturated fat reduction (P < 0.05) during the trial in the intervention arm compared with the control arm. The intervention had to be dietary advice, supplementation of fats, oils or modified or low-fat foods, or a provided diet, compared to higher saturated fat intake which could be usual diet, higher saturated fat, placebo or a control diet. Intended duration of the dietary intervention was at least two years (24 months or 104 weeks).

Types of outcome measures

Primary outcomes

- All-cause mortality (deaths from any cause)
- Cardiovascular (CVD) mortality (deaths from myocardial infarction, stroke, and/or sudden death)
- Combined CVD events. These included data available on number of people experiencing any of the following: cardiovascular death, cardiovascular morbidity (non-fatal myocardial infarction, angina, stroke, heart failure, peripheral vascular events, atrial fibrillation) and unplanned cardiovascular interventions (coronary artery bypass surgery or angioplasty).

To meet our inclusion criteria, trials had to report either deaths or CVD events. These could be reported as serious adverse events (SAEs) or via communication with authors.

Secondary outcomes

 Additional health events; the outcomes CHD mortality and CHD events were added at the request of the WHO NUGAG group,

and were not present in the original overarching systematic

- review. For each of these, we assessed number of participants experiencing any of these:
- Myocardial infarction, total (fatal and non-fatal)
- * Myocaridal infarction, non-fatal

cochrane

- Stroke
- CHD mortality, which includes death from myocardial infarction or sudden CVD death
- CHD events, which include any of the following: fatal or nonfatal myocardial infarction, angina or sudden CVD death
- type II diabetes incidence
- Blood measures including serum blood lipids
 - total cholesterol (TC, mmol/L)
 - * low-density lipoprotein (LDL) cholesterol, mmol/L
 - high-density lipoprotein (HDL) cholesterol, mmol/L
 - * triglyceride (TG), mmol/L
 - * TG/HDL ratio
 - * LDL/HDL ratio
 - total/HDL ratio
 - lipoprotein (a) (Lp(a)), mmol/L
 - insulin sensitivity including glucose tolerance (homeostatic model assessment (HOMA), intravenous glucose tolerance test (IV-GTT), clamp, glycosylated haemoglobin (HbA1C))
- · Other outcomes including adverse effects reported by study authors
 - cancer diagnoses *
 - cancer deaths
 - body weight, kg
 - body mass index (BMI, kg/m²)
 - systolic blood pressure (sBP, mmHg)
 - diastolic blood pressure (dBP, mmHg)
 - quality of life (any measure)

As all trials collect data on deaths and cardiovascular events (as serious adverse events if not as planned outcome measures), we only included trials where we knew that at least one primary outcome occurred, by communication with authors if necessary. Where we knew that at least one primary outcome occurred, we included the study even where we were unable to use that data in meta-analysis. We excluded studies where we knew that no primary outcome events occurred (for a study to be excluded in this way the paper needed to be very explicit about the lack of all outcomes or we received confirmation from the authors) and this was noted as the reason for exclusion. Lack of a single primary outcome only occurs in very small studies or in young cohorts, so omitting these studies will make no difference to effect sizes and very little difference to absolute effect sizes (NNTs etc). All other trials were considered unclear and where we could not gain clarification on events from authors, they were classified as "awaiting assessment".

For composite outcomes (like CVD events), we worked to collect data on the number of participants in each arm who experienced any type of CVD event, and did not double-count people (so that a person experiencing a stroke and two heart attacks during a trial was counted as one person experiencing CVD events, not as three CVD events).

We extracted event and continuous outcome data for the latest time point available within the trial, and always at least 24 months from inception. We collected change data (with a measure of variance) for continuous outcomes where these were available, and end data where change data were not provided in usable format.

Search methods for identification of studies

Electronic searches

The updated searches were run on 15 October 2019 on the following databases:

- CENTRAL (Issue 10 of 12, 2019, Cochrane Library)
- MEDLINE (Epub Ahead of Print, In-Process & Other Non-Indexed Citations, MEDLINE Daily and MEDLINE, Ovid, 1946 to October 14, 2019)
- Embase (Ovid, 1980 to 2019 week 41).

For this update, we introduced searches of two trials registers on 17 October 2019; Clinicaltrials.gov (www.clinicaltrials.gov) and WHO International Clinical Trials Registry Platform (ICTRP) (apps.who.int/trialsearch/). The searches are described in Appendix 1. The RCT filter for MEDLINE was the Cochrane sensitivity and precision-maximising RCT filter (Lefebvre 2011), and for Embase, terms as recommended in the Cochrane Handbook were applied (Lefebvre 2011).

As we were updating another Cochrane review relating to dietary fat (Hooper 2015b) at the same time, results of the searches for both reviews were combined and de-duplicated before assessment of titles and abstracts.

The search to 2014 is described in Hooper 2015a, and previous searches in Hooper 2012..

Searching other resources

We searched for recent publications of the included studies, to ensure the best possible data set for each study.

Data collection and analysis

Selection of studies

Search results were loaded into Covidence software. All authors independently assessed titles and abstracts from the search, differences were resolved by discussion and, when the findings were not clear cut, the full text was collected for assessment. We only rejected articles on initial screen if the author could determine from the title and abstract that the article was not a report of a randomised controlled trial; the trial did not address a low or modified fat diet; the trial was exclusively in children less than 18 years old, pregnant women or the critically ill; the trial was of less than 24 months duration; or the intervention was multifactorial. When we could not reject a title/abstract with certainty, we obtained the full text of the article for further evaluation.

Data extraction and management

We used a data extraction form designed for earlier versions of this review. We extracted data concerning participants, interventions and outcomes, trial quality characteristics (Chalmers 1990), data on potential effect modifiers including participants' baseline risk of cardiovascular disease, trial duration, intensity of intervention (dietary advice, diet provided, dietary advice

Reduction in saturated fat intake for cardiovascular disease (Review)

plus supplementation, supplementation alone), medications used (particularly lipid-lowering medication) and smoking status, numbers of events and total participant years in trial. Where provided, we collected data on risk factors for cardiovascular disease including blood pressure, lipids and weight.

We defined baseline risk of cardiovascular disease as follows: high risk are participants with existing vascular disease including a history of myocardial infarction, stroke, peripheral vascular disease, angina, heart failure or previous coronary artery bypass grafting or angioplasty; moderate risk are participants with a familial risk, dyslipidaemia, diabetes mellitus, hypertension, chronic renal failure; low risk are other participants or mixedpopulation groups. Those at low or moderate risk combined are primary prevention trials.

Data were extracted independently in duplicate by AA, FOJ and/or LH, alongside assessment of risk of bias.

Assessment of risk of bias in included studies

We carried out 'Risk of bias' assessment independently in duplicate as part of data extraction. We assessed trial risk of bias using the Cochrane tool for assessment of risk of bias (Higgins 2011). For included RCTs, we also assessed whether each study:

- 1. was free of systematic differences in care,
- 2. aimed to reduce SFA intake,
- 3. achieved SFA reduction, or
- 4. achieved total serum cholesterol reduction.

We used the category 'other bias' to note any further issues of methodological concern. Funding was not formally a part of our assessment of bias in RCTs as it is not a core part of the Cochrane 'Risk of bias' tool, but was reported in the Characteristics of included studies.

Two authors (LH, NM) independently extracted validity data from studies identified by the previous search, and resolved differences by discussion.

Poorly concealed allocation is associated with a 40% greater effect size (Schulz 1995), so randomisation and allocation concealment are core issues for all trials. Lack of blinding is associated with bias, though smaller levels of bias than lack of allocation concealment (Savovic 2012), especially in studies with objectively measured outcomes (Wood 2008).

For this review, we introduced the concept of summary risk of bias for whole trials. We considered dietary advice or all-food-provided type trials to be at low summary risk of bias where we judged randomisation, allocation concealment, and blinding of outcome assessors to be adequate. Summary risk of bias was considered moderate to high in all other included trials.

Measures of treatment effect

The effect measures of choice were risk ratios (RR) for dichotomous data and mean difference (MD) for continuous data.

Unit of analysis issues

We did not include any cluster-randomised trials in this review, as no relevant studies included at least six clusters. Where there was more than one relevant intervention arm but only one control arm, we either pooled the relevant intervention arms to create a single pairwise comparison (where the intervention arms were equivalently appropriate for this review) as described in the *Cochrane Handbook* (Higgins 2011), or we excluded intervention arms that were not appropriate for this review, or less appropriate than another arm. When two arms were appropriate for different subgroups (Rose corn oil 1965; Rose olive 1965), then we used the control group once with each intervention arm, and divided the number of events in the control group, and the number of participants in the control group, evenly between the two study comparisons.

In the previous version of this review, data for WHI 2006 were presented separately for those without baseline CVD, and with baseline CVD, for most outcomes. This has been altered in this version of the review, so that both sets of data are presented as a single trial except when subgrouping by CVD risk. This has the effect of representing this study in the same way others are represented (which is appropriate), and slightly reducing the weight of the WHI 2006 study in random-effects meta-analysis, altering the numbers in the analysis.

When assessing event data, we aimed to assess number of participants experiencing an event (rather than numbers of events), to avoid counting more than one outcome event for any one individual within any one comparison. Where we were unclear (for example, where a paper reported numbers of myocardial infarcts but not by arm), we asked authors for further information.

Dealing with missing data

Where trials satisfied the inclusion criteria of our review but did not report mortality and morbidity, or not by study arm, we tried to contact study authors. This allowed inclusion of studies that would otherwise have had to be excluded. We excluded studies which were otherwise relevant but where we could not establish the presence or absence of primary outcomes, despite multiple attempts at author contact.

It was often unclear whether data on primary or secondary outcome events may still have been missing, and so we did not impute data for this review.

Where included studies used methods to infer missing data (such as carrying the latest measurement forward), then we used these data in analyses. Where this was not done, we used the data as presented.

Assessment of heterogeneity

We examined heterogeneity using the I² test, and considered it important where greater than 50% (Higgins 2003; Higgins 2011). Where we identified important clinical or unexplained statistical heterogeneity, we did not pool but instead summarised the studies in a narrative format. We used the assessment of heterogeneity in our GRADE assessments, so that the quality of evidence was downgraded where heterogeneity was important, and not explained by subgrouping or meta-regression.

Assessment of reporting biases

We used funnel plots to examine the possibility of small study bias, including publication bias (Egger 1997), for the primary outcomes of total mortality and combined cardiovascular events. For this

update, we also compared findings of fixed- and random-effects meta-analysis since the two methods weight small trials differently,

Cochrane

Data synthesis

2019).

We carried out data synthesis in the absence of clinical heterogeneity. We used numbers of events in each study arm, and total number of participants randomised, where extracted, and Mantel-Haenszel random-effects meta-analysis carried out in Review Manager 5 software, to assess risk ratios. We extracted event and continuous outcome data for the latest time point available within the trial, and always at least 24 months from inception.

and different effect sizes suggest potential small study bias (Page

We excluded trials where we knew that there were no events in either group. Where trials ran one control group and more than one included intervention group, we used data from the intervention group providing the comparison that best assessed the effect of altering dietary fat. Where the intervention groups appeared equal in this respect, we merged the intervention groups (simply added for dichotomous data, and using the techniques described in Higgins 2011 for continuous data). We had planned that if we identified trials randomised by cluster we would reduce the participant numbers to an "effective sample size" (as described by Hauck 1991); however, we found none that were both included and had cardiovascular events or deaths.

To assess the WHO NUGAG question on the effect of consuming < 10%E as SFA relative to > 10%E as SFA on the risk of noncommunicable diseases (NCDs) in the population, we combined studies with a control group saturated fat intake of > 10%E and an intervention group saturated fat intake of < 10%E. To assess the effect of a reduction in %E from SFA from 10% in gradual increments relative to higher intake, we repeated this with saturated fat cut-offs between 7%E and 13%E.

Subgroup analysis and investigation of heterogeneity

Prespecified analyses included:

Effects of SFA reduction compared with usual or standard diet on all (primary and secondary) outcomes and potential adverse effects. This main analysis addressed the main objective of the review and the first WHO specific question.

Prespecified subgroups for all outcomes included:

• energy substitution - we intended to subgroup studies according to the main energy replacement for SFA - PUFA, MUFA, CHO (refined or unrefined), protein, trans fats, a mixture of these, or unclear. However, when we presented these data to the WHO NUGAG group, they suggested that this subgrouping be altered. They suggested that we use all studies where SFA was reduced and any of PUFA, MUFA, CHO or protein were statistically significantly increased (P < 0.05) in the intervention compared to the control group to assess the effects of replacement by each, regardless of whether or not it constituted the main replacement for SFA. This meant that some studies appeared in more than one subgroup. As there were almost no data in the studies on trans fats, or on refined and unrefined carbohydrates, we did not include a trans group or distinguish by carbohydrate type. This subgrouping addresses the main objective of the review, and the third WHO specific question.

Further subgroups, run for primary and CVD health-related secondary outcomes only, included:

Prespecified:

- Baseline SFA intake, represented by control group SFA intake (up to 12%E from SFA, > 12 to 15%E, > 15 to 18%E, > 18%E from SFA, or unclear)
- Sex (men, women and mixed populations)
- Baseline CVD risk (low-risk or general populations, moderaterisk populations which were defined by risk factors for CVD such as hypertension or diabetes, high-risk populations with existing CVD at baseline)
- Duration in study (mean duration in trial up to 24 months, > 24 to 48 months, > 48 months, and unclear). Duration was a prespecified subgroup that we used in earlier versions of this review to separate studies with duration of less than two years from those of at least two years. As we have excluded shorter studies from this review, and have access to longer studies, we have explored duration over longer time spans. As some long studies had a high proportion of participants whose time in trial was censored, and we wanted to express mean experience of the trial, we used mean duration of participants in the study, rather than the formal study duration for this subgrouping, so that some two-year intervention trials, because they had some deaths or dropouts, had a mean duration in trial of 21 or 22 months.

WHO NUGAG added subgroups:

- Degree of SFA reduction, represented by the difference between SFA intake in the intervention and control groups during the study (up to 4%E from SFA reduction achieved, > 4 to 8% reduction achieved, > 8% reduction achieved, unclear). We prespecified that we intended to explore the degree of SFA reduction in meta-regression, but its addition as a subgroup was post hoc, and requested by the WHO NUGAG group.
- Serum total cholesterol reduction achieved (reduced by a mean of at least 0.2 mmol/L, reduced by less than 0.2 mmol/L or unclear). We prespecified that we intended to explore the degree of serum total cholesterol reduction in meta-regression.
- Ethnic group. Insufficient information was presented to make this feasible. Hence, we report ethnicity information in the Characteristics of included studies.

We explored the effects of different levels of SFA, PUFAs, MUFAs and total dietary fats, and CHO achieved in trials (all as difference between the intervention and control groups, as %E, and for SFA as a percentage of SFA in the intervention compared with control), baseline SFA intake (as %E), change in total cholesterol (difference between intervention and control groups, in mmol/L), sex, study duration in months, and baseline CVD risk using metaregression on total cardiovascular events. We performed randomeffects meta-regression (Berkley 1995) using the STATA command metareg (Sharp 1998; Sterne 2001; Sterne 2009).

To explore the WHO NUGAG specific question about the effect of the population consuming < 10%E as SFA relative to > 10%E SFA, we assessed effects of all studies where the mean assessed intervention SFA intake was < 10%E and the mean control SFA intake was > 10%E. We explored the effect of reduction of %E from SFA in gradual increments by using cut-offs of 7%E (where

all studies with a mean intervention SFA intake < 7%E and mean control SFA intake > 7%E were pooled), 8%, 9%, 10%, 11%, 12% and 13%. We omitted studies where SFA intakes were not reported from these analyses. For each primary outcome, we plotted the pooled risk ratio of that outcome against the cut-off, %E from SFA.

Referee-added subgroups:

In response to the suggestion of a referee of this systematic review, and to better understand the effect of use of statins since the 1990s, we subgrouped studies by decade of publication.

Sensitivity analysis

We carried out sensitivity analyses for primary outcomes assessing the effect of:

- 1. Excluding studies which did not state an aim to reduce SFA
- 2. Excluding studies which did not report SFA intake during the trial, or did not find a statistically significant reduction in SFA in the intervention compared to the control
- Excluding studies where total cholesterol (TC) was not reduced (statistically significant reduction of TC, or of LDL where TC was not reported (considered reduced where P < 0.05), or where reduction was not at least 0.2 mmol/L in intervention compared to control where variance was not reported)
- 4. Excluding the largest study (WHI 2006)
- 5. Analysis run with Mantel-Haenszel fixed-effect model
- 6. Analysis run with Peto fixed-effect model

For this update we also introduced sensitivity analysis excluding trials not at low summary risk of bias. We used results of these analysis to inform GRADE assessment of risk of bias.

GRADE

All primary outcomes, and secondary additional health events, were represented in the 'Summary of findings' table, and underwent GRADE assessment. The GRADE Working Group has developed a common, sensible and transparent approach to grading quality of evidence and strength of recommendations (www.gradeworkinggroup.org/; GRADE 2004). The evidence within this systematic review was first assessed using the GRADE system by the review authors and then discussed and modified by the WHO NUGAG group.

Outcome data were interpreted as follows:

- Is there an effect? (options were 'increased risk', 'decreased risk', or 'little or no effect'). Our main outcome measure was RR so we decided on existence of an effect using RR. RR > 8% (RR < 0.92 or > 1.08) for the highest quality evidence suggested increased or decreased risk (otherwise little or no effect). The presence or not of an effect was decided on the RR for the main analysis and sensitivity analyses, the highest quality evidence (the main analysis, the sensitivity analyses of trials at low summary risk of bias and at low risk of compliance problems).
- 2. For continuous outcomes, reducing SFA was considered to have little or no effect unless effect sizes represented at least 5% change from baseline (or 2% in the case of cumulative outcomes such as adiposity).
- 3. Quality of evidence was assessed using GRADE assessment (GRADE 2004) for key outcomes. We used the five GRADE considerations (risk of bias, consistency of effect, imprecision, indirectness and publication bias) to assess the quality of the body of evidence as it related to the studies that contributed data to the meta-analyses for the prespecified outcomes. We used methods and recommendations described in Section 8.5 and Chapter 12 of the *Cochrane Handbook for Systematic Reviews of Interventions* (Higgins 2011), plus GRADEpro GDT software (GRADEpro 2015). We justified all decisions to downgrade the quality of studies using footnotes and made comments to aid reader's understanding of the review.
- 4. Where there was a suggested effect, the size of effect was assessed using the number needed to treat for an additional beneficial outcome (NNTB), number needed to treat for an additional harmful outcome (NNTH) or absolute risk reduction (ARR).

RESULTS

Description of studies

Results of the search

Figure 1 displays the flow diagram for inclusion of studies. We assessed the 7991 titles and abstracts from the updated electronic search, as well as assessing the 8930 titles and abstracts from the search for our sister review (Hooper 2015b), which de-duplicated to 15,314 titles and abstracts. Of these, 530 were considered potentially relevant to one or both reviews, so were collected as full text. Ten publications were considered relevant for this systematic review, and these were grouped into:







Figure 1. (Continued)



- three new publications for two already included trials (WHI 2006; WINS 2006),
- two publications for one study awaiting assessment (not enough details to confirm inclusion, ICFAMED), and
- five publications for four ongoing trials (ENAbLE due unclear; NCT02481466 due 2020; NCT02938832 due 2023; NEW Soul Study due 2022).

There were no new included trials, but there were new data for WHI 2006 and WINS 2006, as well as the ongoing studies and the study awaiting assessment. This resulted in an updated review including 15 RCTs (16 comparisons as the Rose trial has two comparisons, Rose corn oil 1965 and Rose olive 1965).

Included studies

We included 15 randomised controlled trials (RCTs) in the review (Included studies), and describe them in Characteristics of included studies. The interventions are compared in Table 1.

The main study papers ranged in publication date from 1965 to 2006, but with supplementary publications included up to 2019. The RCTs were conducted in North America (six), Europe (seven), and Australia/New Zealand (two); no studies were carried out in industrialising or developing countries. Six RCTs included only people at high risk of cardiovascular disease, four at moderate risk, and four at low risk (three with raised cancer risk or cancer diagnosis, one with no specific health risks), while one trial included participants at low and high CVD risk (WHI 2006, Table 1; this trial made assessments in each of these groups). Seven studies included only men, three only women, and five both men and women. However, as the largest trial (WHI 2006) was in women only, women are the largest group represented. Trial duration ranged

from two to more than eight years, with a mean duration of 4.7 years.

The form of interventions varied (Table 1). Interventions were of advice to alter intake in 15 of the 16 intervention arms, and additional supplements such as oil or other foods were provided in three trials (four arms: MRC 1968; Oslo Diet-Heart 1966; Rose corn oil 1965; Rose olive 1965), while all food was provided in a residential facility in one RCT (Veterans Admin 1969). Of the 15 arms with an advice element, most interventions were delivered face-toface, but this was unclear in three arms (Houtsmuller 1979; Rose corn oil 1965; Rose olive 1965). Advice was provided individually in nine intervention arms (followed by later group sessions in two arms), in groups only in two trials (Ley 2004; WHI 2006), and was unclear in three RCTs (Black 1994; Houtsmuller 1979; Rose corn oil 1965; Rose olive 1965). Advice was provided by a dietitian in nine arms, a nutritionist in one, a trained nurse in one and was unclear in four. Frequency of study visits for advice and follow-up varied between three times in the first year and twice annually thereafter up to 18 sessions in the first year and quarterly maintenance visits thereafter.

Of the 15 included studies (16 intervention arms), 11 RCTs (12 comparisons) provided data on all-cause mortality (including 55,858 participants and 3518 deaths), 10 RCTs (11 comparisons) on CV mortality (53,421 participants and 1096 cardiovascular deaths), and 11 RCTs (12 comparisons) on combined cardiovascular CVD events (53,300 participants, of whom 4476 participants experienced at least one CVD event) (Table 2). In two included studies, it was clear that events had occurred, but it was not clear in which arm(s) the events had occurred (Oxford Retinopathy 1978; Simon 1997), so that we could not include the data in the meta-analyses. Secondary health events and other secondary outcomes

were reported in varying number of studies (between 1 and 15 studies reported on any single outcome, see Table 2 and Table 3).

Excluded studies

We excluded 520 full-text publications at this update, having assessed the full texts in duplicate. We describe the reasons for some of these exclusions in Characteristics of excluded studies tables. We excluded 29 studies where data on events were not reported in publications and contact with authors confirmed that there had been no deaths or cardiovascular events, where contact with authors confirmed that data were not available, or where we could not establish contact with authors.

Risk of bias in included studies

We display 'Risk of bias' assessments in the individual included study arms in Figure 2.

Figure 2. Methodological quality summary: review authors' judgements about each methodological quality item for each included study. Please note that while Rose 1965 (Rose corn oil 1965; Rose olive 1965) appears twice in this



summary, it is a single trial. Rose 1965 was a 3-arm trial and we have used the two intervention arms separately in the review.



Allocation

All the included trials were randomised controlled trials, and some detail of the randomisation process was provided for all studies, so all were considered at low risk of bias. We excluded those with detected pseudo-random allocation (for example where participants are randomised according to birth date or alphabetically from their name). We judged allocation concealment to be well done in eight RCTs (eight comparisons, Ley 2004; Oslo Diet-Heart 1966; Oxford Retinopathy 1978; STARS 1992; Sydney Diet-Heart 1978; Veterans Admin 1969; WHI 2006; WINS 2006), and unclear in the remainder.



Blinding

Blinding of participants is not easy in dietary studies, as the participants usually have to follow instructions to attain the specific dietary goals. However, it is feasible in some circumstances, including when food is provided via an institutional setting, or meals provided at a central setting and remaining meals packed to take away. It can also be achieved through use of a trial shop, where very specific food-based dietary advice is provided to all participants, or where the same dietary advice is provided to both groups but a different supplement (e.g. dietary advice to reduce fats, then provision of different oils or fats) is provided. Where participants are not blinded, it is difficult to ensure that study staff, healthcare providers and outcome assessors are blinded. The single RCT that appears to have had adequate participant and study personnel blinding was Veterans Admin 1969, and we judged blinding of participants to be inadequate in the remaining studies.

Blinding of outcome assessment was assessed separately for mortality and CVD outcomes. Blinding is not relevant in assessing all-cause mortality, so all trials were considered at low risk of bias for detection bias for this outcome. For CVD outcomes, nine trials were at low risk of detection bias, one was at high risk and the remainder were unclear.

Incomplete outcome data

Assessing whether incomplete outcome data had been addressed was difficult, as the primary outcomes for this review (mortality and cardiovascular events) were often reported as dropouts and exclusions from the original studies, rather than as the primary outcomes of these trials. When mortality or cardiovascular events or both were noted in any one study, it is still feasible that some participants left that study feeling unwell or because the diet was inconvenient, so were simply lost to follow-up from the perspective of the study, and later died or experienced a cardiovascular event. However, six of the studies checked medical records or death registers to ensure that such events were all collected (Black 1994, DART 1989; Oslo Diet-Heart 1966; Sydney Diet-Heart 1978; Veterans Admin 1969; WINS 2006). Within one study, there was extensive tracking of medical records, with assessment of health status by blinded trained adjudicators (WHI 2006), so few major events were likely to have been missed. In the other eight studies, it is not possible to know whether additional deaths or cardiovascular events occurred, that were not counted or ascertained within this review.

Selective reporting

Assessment of selective reporting is difficult when the outcome of interest was simply considered a cause of dropouts in most included studies. We tried to contact all of the trialists to ask about deaths and outcome events, but it is possible that some trialists did not reply as they felt that their data did not reflect the expected or hoped-for pattern of events. All of the included studies have either reported that the participants did not experience any of our primary outcomes, have published their outcome data, or have provided the data they did possess. For this reason, we have graded all the included studies as at low risk of selective reporting.

Other potential sources of bias

Systematic differences in care. We assessed the studies for risk of bias in relation to systematic differences in care. The three

RCTs (four comparisons) that appeared at low risk of systematic differences in care between the study arms included Rose corn oil 1965; Rose olive 1965; Oxford Retinopathy 1978; Veterans Admin 1969, while 11 RCTs clearly did have differences in care, such as differential time provided for those on the intervention to learn a new diet, and/or differential medical follow-up, and one was unclear (Houtsmuller 1979).

Aim to reduce saturated fat. As several studies did not provide clear aims for their interventions (other than to alter specific dietary components, for example), we assessed whether the study stated an aim to reduce saturated fat. Ten RCTs (11 comparisons) clearly aimed to reduce saturated fat in their intervention arms, either directly or indirectly, for example, by stating food goals (DART 1989; Houtsmuller 1979; Moy 2001; MRC 1968; Oslo Diet-Heart 1966; Rose corn oil 1965; Rose olive 1965; STARS 1992; Sydney Diet-Heart 1978; Veterans Admin 1969; WHI 2006), while the remaining five did not (although they did achieve SFA reduction).

Successful saturated fat reduction. Eleven RCTs (11 comparisons) assessed SFA intake during the study period and showed that SFA intake in the intervention arm was statistically significantly lower than that in the control arm (Black 1994; DART 1989; Ley 2004; Moy 2001; Oxford Retinopathy 1978; Simon 1997; STARS 1992; Sydney Diet-Heart 1978; Veterans Admin 1969; WHI 2006; WINS 2006). The remaining studies did not report SFA intake, so we rated them as unclear.

Successful cholesterol reduction. We would expect saturated fat reduction to be reflected in total or LDL cholesterol reductions, which may be more accurate assessments than self-reported saturated fat intake. Nine RCTs (10 comparisons) provided information on serum total or LDL cholesterol levels in the intervention and control arms during the study, and found a reduction in the intervention arm compared to the control (P < 0.05, or where variances were not provided showed a reduction of at least 0.2 mmol/L in the mean intervention measure compared with control). The studies that successfully reduced serum total cholesterol in lower saturated fat arms compared with higher saturated fat arms were DART 1989; Houtsmuller 1979; Simon 1997; STARS 1992; Sydney Diet-Heart 1978; WHI 2006, while Moy 2001 did not report total cholesterol (TC) but showed statistically significant reductions in LDL, and two studies (MRC 1968; Oslo Diet-Heart 1966) did not report variances but did reduce mean TC in the intervention arm compared with control by at least 0.2 mm0l/L. One study (Black 1994) did not report lipid levels during the study, while five others did report lipid levels but did not suggest clear differences between lower and higher saturated fat arms (Ley 2004; Oxford Retinopathy 1978; Rose corn oil 1965; Rose olive 1965; Veterans Admin 1969; WINS 2006).

Dietary changes other than saturated fat. Some trials were partially confounded by aiming to make dietary changes other than those directly related to dietary fat intakes; for example, some studies encouraged intervention participants to make changes to their fat intake as well as changes to fruit and vegetable or fibre or salt intakes. In these studies, any effect on outcomes could be a result of other dietary changes, not of changes in saturated fat intake. The 11 studies (12 comparisons) that appeared free of such differences included Black 1994; DART 1989; Houtsmuller 1979; Ley 2004; MRC 1968; Oxford Retinopathy 1978; Rose corn oil 1965; Rose olive 1965; Simon 1997; Sydney Diet-Heart 1978; Veterans Admin 1969; WINS 2006. This factor was not considered alongside others

Reduction in saturated fat intake for cardiovascular disease (Review) Copyright © 2020 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.



in the formal risk of bias assessment (Figure 2) so is described here. We did not identify any further methodological issues.

Summary risk of bias. We considered dietary advice or all-foodprovided type trials to be at low summary risk of bias where we judged randomisation, allocation concealment, and blinding of outcome assessors to be adequate. For CVD outcomes, five trials were assessed as at low summary risk of bias: Ley 2004; Sydney Diet-Heart 1978; Veterans Admin 1969; WHI 2006; WINS 2006. For all-cause mortality (and lipid outcomes) where blinding of outcome assessors is not important, a further three trials were also at low summary risk of bias, eight in total: Ley 2004; Oslo Diet-Heart 1966; Oxford Retinopathy 1978; STARS 1992; Sydney Diet-Heart 1978; Veterans Admin 1969; WHI 2006; WINS 2006.

Effects of interventions

See: Summary of findings 1 Effect of reducing saturated fat compared to usual saturated fat on CVD risk in adults (note: for the full set of GRADE tables see additional tables 24 to 28)

Primary outcomes

All-cause mortality

GRADE assessment suggests that reducing saturated fat intake probably makes little or no difference to all-cause mortality (moderate-quality evidence, downgraded once for imprecision).

There was little or no effect of lower saturated fat compared to higher saturated fat intake on mortality (risk ratio (RR) 0.96, 95% confidence interval (CI) 0.90 to 1.03, $I^2 = 2\%$, 55,858 participants, 3518 deaths, 11 RCTs, $P_{effect} = 0.42$, Analysis 1.1). This lack of effect was confirmed in sensitivity analyses including only trials at low summary risk of bias (Analysis 1.2), that aimed to reduce saturated fat (Analysis 1.3), that significantly reduced saturated fat intake (Analysis 1.4), that achieved a reduction in total or LDL cholesterol (Analysis 1.5), excluding the largest trial (WHI 2006, Analysis 1.6), or analysing using Mantel-Haenszel or Peto fixed-effect analysis (Analysis 1.7; Analysis 1.8).

Small study bias was assessed using a funnel plot and comparing the results of fixed- and random-effects meta-analysis. The funnel plot did not suggest any small study bias (Figure 3), and the results of fixed- and random-effects meta-analyses were very similar, suggesting that small study bias was not an issue.

Figure 3. Funnel plot of comparison: fat modification or reduction vs usual diet - total mortality.



There was little or no effect, regardless of what nutrients were used to replace the saturated fat removed, including replacement with PUFA, MUFA, CHO and/or protein (Analysis 1.9). Effects did not differ

by main substitution (Analysis 1.10), study duration (Analysis 1.11), baseline saturated fat intake (Analysis 1.12), degree of difference in saturated fat between arms (Analysis 1.13), participant sex



(Analysis 1.14), by baseline CVD risk (Analysis 1.15), by degree of cholesterol reduction (Analysis 1.16) or by decade of publication (Analysis 1.17, Chi^2 test for differences between subgroups all P > 0.05).

Cardiovascular mortality

GRADE assessment suggests that reducing saturated fat intake probably makes little or no difference to cardiovascular mortality (moderate-quality evidence, downgraded once for imprecision).

There was little or no effect of SFA reduction on cardiovascular mortality (RR 0.95, 95% CI 0.80 to 1.12, $I^2 = 30\%$, 10 RCTs, 53,421 participants, 1096 cardiovascular deaths, Analysis 1.18). This lack

of effect was confirmed in sensitivity analyses limiting to trials at low summary risk of bias (Analysis 1.19), explicitly aiming to reduce saturated fat (Analysis 1.20), achieving statistically significant saturated fat reduction (Analysis 1.21), achieving cholesterol reduction (Analysis 1.22), or running fixed-effect analysis (Analysis 1.24; Analysis 1.25). However, excluding the largest single trial (WHI 2006) suggested that reducing saturated fat intake reduced the risk of CVD mortality (Analysis 1.23).

The funnel plot did not suggest small study bias (Figure 4), and the similarity in effect sizes between fixed- and random-effects analysis suggests that small study bias is not important here.

Figure 4. Funnel plot of comparison: fat modification or reduction vs usual diet - cardiovascular mortality



Subgrouping did not suggest important effects of reduced SFA on cardiovascular mortality, regardless of what was substituted for SFA (Analysis 1.26). When subgrouping by main substitution (Analysis 1.27), duration (Analysis 1.28), baseline SFA intake (Analysis 1.29), by difference in SFA (Analysis 1.30), participant sex (Analysis 1.31), baseline CVD risk (Analysis 1.32), or degree of cholesterol reduction (Analysis 1.33), there were no statistically significant differences between subgroups. There was a marginally significant difference between subgroups when ordered by decade of publication, but no clear pattern of effect, so we assumed the effect was probably spurious (Analysis 1.34). Additionally, effects did not appear to relate to statin use, as there was a reduction in risk of CVD mortality in studies published in the 1960s and a marginal increase in risk

in the one trial published during the 1970s (although the 95% confidence interval did include 1.0), both well before statins were in common use (the 4S trial which first showed that use of statins reduced mortality was published in 1994, 4S 1994).

Cardiovascular events

GRADE assessment suggests that reducing SFA intake probably reduces cardiovascular events, to a greater extent with greater cholesterol reduction (moderate-quality evidence, downgraded once for publication bias).

There was a 21% reduction in cardiovascular events in people who had reduced SFA compared with those on higher SFA (RR 0.79, 95%



CI 0.66 to 0.93, $I^2 = 65\%$, 11 RCTs, 53,300 participants, 4476 people with cardiovascular events, $P_{effect} = 0.006$, Analysis 1.35). This protective effect was confirmed in sensitivity analyses including only trials at low summary risk of bias (Analysis 1.36), that aimed to reduce saturated fat (Analysis 1.37), that significantly reduced saturated fat intake (Analysis 1.38), that achieved a reduction in total or LDL cholesterol (Analysis 1.39), or excluding the largest trial (WHI 2006, Analysis 1.40). Analysing using Mantel-Haenszel or Peto fixed-effect analysis suggested more marginal protection (Analysis 1.41; Analysis 1.42). A funnel plot did not suggest severe small-study bias, but it is possible that a few small studies with more cardiovascular events in the intervention groups may be missing from the review (Figure 5). Fixed-effect analyses suggested smaller effects (Analysis 1.41; Analysis 1.42), again suggesting that smaller studies with more cardiovascular events in the intervention groups may be missing. Adding any such studies back would tend to moderate the protective effect of reducing SFA.

Figure 5. Funnel plot of comparison: fat modification or reduction vs usual diet - combined cardiovascular events.



When we subgrouped according to replacement for SFA, the PUFA replacement group suggested a 27% reduction in cardiovascular events, while there were no clear effects of other replacement groups (Analysis 1.44). However, there were no statistically significant differences between subgroups by main replacement (Analysis 1.45), by study duration (Analysis 1.46), or baseline CVD risk (Analysis 1.50). When subgrouping, there was a suggestion of greater effects when baseline SFA was higher (Analysis 1.47), with greater reduction of SFA (Analysis 1.48), in men rather than women (Analysis 1.49), and with greater cholesterol reduction (Analysis 1.51). There were different effects by decade of publication, but no suggestion of a trend or a change following wider introduction of statins in the mid-1990s (Analysis 1.52).

We further explored the effects of dietary fats on cardiovascular events, by using meta-regression of the difference between the control and intervention of total fat intake, SFA intake, MUFA intake, PUFA intake, CHO intake (all by percentage of energy (%E)), serum total cholesterol (in mmol/L) achieved in trials, as well as baseline SFA intake, sex, study duration in months, and CVD risk of participants at baseline (Table 4). As we included only 13 studies for this outcome, we ran meta-regressions exploring single explanatory factors at once, and as data were limited, with many studies not reporting dietary intake data, these analyses were limited in power to assess outcomes. The data suggested that greater reductions in total serum cholesterol levels reduced CVD events more. Greater baseline SFA intake and greater reduction in SFA were also associated with greater improvement in CVD events with SFA reduction, and increases in PUFA and MUFA intakes were slightly protective of CVD events, but none of these relationships were statistically significant. Overall, the relationship with serum total cholesterol was clearest (P = 0.04, accounting for 99% of between-study variation). Sex, study duration and



baseline cardiovascular risk did not appear to influence effect size. Apparent heterogeneity was accounted for by a dose-effect; where SFA reduction resulted in greater serum cholesterol reduction, the reduction in CVD events was greater.

This 21% reduction in risk of CVD events translated into a number needed to treat for an additional beneficial outcome (NNTB) of 56 in primary prevention trials, so that 56 people need to reduce their saturated fat intake over around four years for one person to avoid experiencing a CVD event. In secondary prevention trials, the NNTB was 32.

Secondary outcomes - health events

Myocardial Infarction (fatal and non-fatal)

GRADE assessment suggested that the effect of reducing saturated fat intake on risk of myocardial infarction is unclear as the evidence was of very low-quality (downgraded once each for risk of bias, imprecision and publication bias).

There was a small protective effect of SFA reduction on myocardial infarction (fatal and non-fatal, RR 0.90, 95% CI 0.80 to 1.01, $I^2 = 10\%$, 10 RCTs (11 comparisons) including 53,167 participants, 1714 people experiencing MI, Analysis 2.1). This protective effect was slightly modified in sensitivity analyses, and confirmed in analyses limited to trials that aimed to reduce saturated fat (Analysis 2.3), that achieved a reduction in total or LDL cholesterol (Analysis 2.5), and excluding the largest trial (WHI 2006, Analysis 2.6). Sensitivity analyses including only trials at low summary risk of bias (RR 0.93, 95% CI 0.81 to 1.08, Analysis 2.2), that significantly reduced saturated fat intake (Analysis 2.4), analysed using Mantel-Haenszel or Peto fixed-effect analysis (Analysis 2.7; Analysis 2.8) suggested little or no effect, though risk ratios were still all < 1.0.

The funnel plot was difficult to interpret, but did not raise major concerns about small-study bias (not shown). While effects of random- and fixed-effect meta-analysis were only slightly different, they fell each side of the line suggesting an effect (Analysis 2.7; Analysis 2.8). There may be a small amount of small study bias.

The protective effect of replacing SFA with PUFA appeared to explain the reduction in MI (Analysis 2.9), but there were no statistically significant differences between subgroups by replacement (Analysis 2.10), duration (Analysis 2.11), baseline SFA intake (Analysis 2.12), change in SFA intake (Analysis 2.13), participant sex (Analysis 2.14), baseline CVD risk (Analysis 2.15), cholesterol reduction (Analysis 2.16) or decade of publication (Analysis 2.17).

Myocardial Infarction (non-fatal only)

GRADE assessment suggests that reducing saturated fat may have little or no effect on risk of non-fatal myocardial infarction (lowquality evidence, downgraded once each for risk of bias and imprecision).

There was no clear effect of SFA reduction compared to usual diet on non-fatal myocardial infarction (RR 0.97, 95% Cl 0.87 to 1.07, l^2 = 0%, 7 RCTs, 52,834 participants, 1385 people with at least one non-fatal MI, Analysis 2.18). This lack of effect was not altered in sensitivity analyses retaining only those that aimed to reduce SFA (Analysis 2.20), those showing a reduction in serum cholesterol (Analysis 2.22), or fixed-effect analysis (Analysis 2.24; Analysis 2.25). However, sensitivity analyses retaining only trials at low summary risk of bias (RR 0.89, 95% CI 0.58 to 1.35, Analysis 2.19), those showing a significant reduction in SFA (Analysis 2.21), and omitting the largest trial (WHI 2006, Analysis 2.23) all suggested a reduction in non-fatal MI with reduced SFA.

The funnel plot did not raise major concerns about small-study bias (not shown), and effects of fixed- and random-effects analyses were very similar, reinforcing the lack of small study bias.

Subgrouping by any replacement for SFA suggested reductions in non-fatal MI when replaced by PUFA, but not other replacements (Analysis 2.26). Subgrouping by main substitution (Analysis 2.27), duration (Analysis 2.28), baseline SFA intake (Analysis 2.29), degree of SFA reduction (Analysis 2.30), sex (Analysis 2.31), baseline CVD risk (Analysis 2.32), degree of cholesterol reduction (Analysis 2.33) and decade of publication (Analysis 2.34) did not suggest significant differences between subgroups.

Stroke (any type, fatal or non-fatal)

GRADE assessment suggests that the effect of reducing SFA intake on stroke is unclear as the evidence is of very low-quality (downgraded twice for imprecision and once for risk of bias).

As data on stroke were sparse, it was not possible to tease out differential effects on ischaemic or haemorrhagic strokes, or whether a stroke was fatal. For this analysis, we combined all stroke data from any study. There was little or no effect of SFA reduction compared to usual diet on stroke of any type with any outcome (RR 0.92, 95% CI 0.68 to 1.25, $I^2 = 9\%$, 7 RCTs, 50,952 participants, 1118 people with stroke, Analysis 2.35). This lack of effect was not altered in sensitivity analyses retaining only those that aimed to reduce SFA (Analysis 2.37), those showing a reduction in serum cholesterol (Analysis 2.39), or fixed-effect analysis (Analysis 2.41; Analysis 2.42). However, for sensitivity analyses retaining only trials at low summary risk of bias (RR 0.76, 95% CI 0.42 to 1.38, Analysis 2.36), those showing a significant reduction in SFA (Analysis 2.38), and omitting the largest trial (WHI 2006, Analysis 2.40) the best estimate of effect always suggested a reduction in stroke with reduced SFA, though they were not statistically significant.

We did not create a funnel plot as the analysis only included data from seven RCTs, however RRs generated using fixed-effect analyses were much closer to 1.0 than the random-effects metaanalysis (suggesting a small amount of publication bias), though both suggested little or no effect.

Subgrouping by any substitution for SFA suggested reduction in risk of stroke whether SFA was replaced by PUFA, CHO or protein (Analysis 2.43). Subgrouping did not suggest significant differences between subgroups by main substitution (Analysis 2.44), duration (Analysis 2.62), baseline SFA (Analysis 2.46), SFA change (Analysis 2.47), sex (Analysis 2.48), CVD risk (Analysis 2.49), cholesterol reduction (Analysis 2.50) or decade of publication (Analysis 2.51).

Coronary heart disease (CHD) mortality

GRADE assessment suggests that reducing saturated fat intake may have little or no effect on CHD mortality (low-quality evidence, downgraded twice for imprecision).

Eight RCTs (9 comparisons) suggest little or no effect of reducing saturated fat on risk of CHD mortality (RR 0.97, 95% CI 0.82 to 1.16, $I^2 = 28\%$, 53,159 participants, 927 people died of coronary heart



LIDrary Better health.

ochrane

disease, Analysis 2.52), and this was not altered in any sensitivity analyses (Analysis 2.53; Analysis 2.54; Analysis 2.55; Analysis 2.56; Analysis 2.57; Analysis 2.58; Analysis 2.59).

We did not create a funnel plot as the analysis only included data from seven RCTs, but the results of fixed- and random-effects analyses were nearly identical, suggesting that small study bias is not an issue here.

There was no suggestion of an effect of reducing SFA on CHD mortality regardless of what replaced the SFA (Analysis 2.60). There were no statistically significant differences between subgrouping in any analysis (Analysis 2.61; Analysis 2.62; Analysis 2.63; Analysis 2.64; Analysis 2.65; Analysis 2.66; Analysis 2.67; Analysis 2.68).

Coronary heart disease events

GRADE assessment suggested that the effect of reducing saturated fat on CHD events is unclear as the evidence is of very lowquality (downgraded once each for imprecision, risk of bias and inconsistency).

There was the suggestion of a 17% reduction in CHD events as a result of saturated fat reduction in the main analysis (RR 0.83, 95% CI 0.68 to 1.01, $I^2 = 62\%$, 53,199 participants, 2261 people had at least one coronary heart disease event in 10 RCTs, Analysis 2.69). This did not differ in sensitivity analyses (Analysis 2.74; Analysis 2.72; Analysis 2.73; Analysis 2.71; Analysis 2.75; Analysis 2.76) except when limiting to trials at low summary risk of bias (RR 0.92, 95% CI 0.77 to 1.10, Analysis 2.70).

The funnel plot did not appear unbalanced, and the results of fixed- and random-effects analyses were different, though both suggested that reducing SFA resulted in lower risk of CHD.

Subgrouping by any replacement for SFA suggested that replacement by PUFA may lead to reduced risk of CHD events (Analysis 2.77). There were no statistically significant differences between any other subgroups (Analysis 2.78; Analysis 2.79; Analysis 2.80; Analysis 2.81; Analysis 2.82; Analysis 2.83; Analysis 2.84) except by decade of publication, though this did not suggest any sequence or step change (Analysis 2.85).

Type 2 diabetes, new diagnoses

Only one RCT reported on diagnosis of diabetes (WHI 2006). There was little or no effect of reducing SFA intakes on diagnosis of diabetes in this study (RR 0.96, 95% CI 0.90 to 1.02, 48,835 participants, 3342 developed diabetes, Analysis 2.86). WHI 2006 was assessed at low summary risk of bias, aimed to reduce SFA, and demonstrated significant SFA and cholesterol reduction. With only one trial, we were not able to assess publication bias or carry out subgrouping.

Secondary outcomes - blood levels

Serum blood lipids

Total cholesterol (TC): There was a reduction in TC in participants with reduced SFA compared to higher SFA (mean difference (MD) -0.24 mmol/L, 95% CI -0.36 to -0.13, $I^2 = 60\%$, 13 RCTs, 7115 participants, Analysis 3.1). We did not conduct sensitivity analyses or most subgroupings on secondary outcomes, but there was no clear differential effect on TC depending on the replacement for SFA (PUFA, MUFA, CHO or a mixture, Analysis 3.2; Analysis 3.3).

The funnel plot did not raise concerns about small-study bias (not shown).

Low-density lipoprotein (LDL): There was a reduction in LDL in participants with reduced SFA compared to higher SFA (MD -0.19 mmol/L, 95% CI -0.33 to -0.05, $I^2 = 37\%$, 5 RCTs, 3291 participants, Analysis 3.4). There was no clear differential effect on LDL depending on the replacement for SFA (PUFA, MUFA, CHO or a mixture, Analysis 3.5; Analysis 3.6). We could not interpret the funnel plot due to sparsity of studies (not shown).

High-density lipoprotein (HDL): There was little or no effect of reducing SFA intakes on HDL (MD -0.01 mmol/L, 95% CI -0.02 to 0.01, $I^2 = 0\%$, 7 RCTs, 5147 participants, Analysis 3.7). There was no clear differential effect on HDL depending on the replacement for SFA (PUFA, MUFA, CHO or a mixture, Analysis 3.8; Analysis 3.9). We could not interpret the funnel plot due to sparsity of studies (not shown).

Triglycerides (TG): There was little or no effect of reducing SFA intakes on TG (MD -0.08 mmol/L, 95% CI -0.21 to 0.04, $I^2 = 51\%$, 7 RCTs, 3845 participants, Analysis 3.10). There was no clear differential effect on TG depending on the replacement for SFA (PUFA, MUFA, CHO or a mixture, Analysis 3.11; Analysis 3.12). We could not interpret the funnel plot due to sparsity of studies (not shown).

TG/HDL ratio: We did not find any studies that reported TG/HDL ratio.

TC/HDL ratio: Only three RCTs reported on TC/HDL ratio. There was little or no effect of reducing SFA intakes on TC/HDL (MD -0.10, 95% CI -0.33 to 0.13, $I^2 = 24\%$, 2985 participants, Analysis 3.13). There were no clear differential effects of replacement on TC/HDL (Analysis 3.14; Analysis 3.15). We could not interpret the funnel plot due to sparsity of studies (not shown).

LDL/HDL ratio: Only one RCT reported on LDL/HDL ratio. There was no clear effect of reducing SFA intakes on LDL/HDL in this study (MD -0.36, 95% CI -0.92 to 0.20, 50 participants, Analysis 3.16). This study replaced SFA with CHO (mainly) and PUFA.

Lipoprotein (a) (Lp(a)): Only two RCTs reported on lipoprotein (a), but these included 28,820 participants. There was little or no effect of reducing SFA intakes on Lp(a) (MD 0.00, 95% CI -0.00 to 0.00, $I^2 = 0\%$, Analysis 3.17). There was no suggestion of differential effects of replacement on Lp(a) (Analysis 3.18; Analysis 3.19). We could not interpret the funnel plot due to sparsity of studies (not shown).

Homeostatic model assessment (HOMA): Only one RCT reported on the effects of reducing SFA on insulin resistance using HOMA. There was little or no effect of reducing SFA intakes compared to usual diet on HOMA in this study (MD -0.00, 95% CI -0.04 to 0.04, 2832 participants, Analysis 3.20).

Glucose at two hours post-glucose tolerance test (GTT): Only three RCTs reported on glucose two hours post-GTT. There was a reduction in glucose after reducing SFA intakes compared to usual diet (MD -1.69 mmol/L, 95% CI -2.55 to -0.82, $I^2 = 45\%$, 249 participants, Analysis 3.20). We could not interpret the funnel plot due to sparsity of studies (not shown).

HbA1c (glycosylated haemoglobin): HbA1c was not measured in any included RCT.



Secondary outcomes - other outcomes and potential harms

There was little or no effect of reducing SFA intakes on **cancer diagnoses** of any type (RR 0.94, 95% CI 0.83 to 1.07, $I^2 = 33\%$, 4 RCTs, 52,294 participants, 5476 cancer diagnoses, Analysis 4.1); **cancer deaths** (RR 1.00, 95% CI 0.61 to 1.64, $I^2 = 49\%$, 5 RCTs, 52,283 participants, 2472 cancer deaths, Analysis 4.2); **systolic blood pressure** (MD -0.19 mmHg, 95% CI -1.36 to 0.97, $I^2 = 0\%$, 5 RCTs, 3812 participants, Analysis 4.5); **diastolic blood pressure** (MD -0.36 mmHg, 95% CI -1.03 to 0.32, $I^2 = 0\%$, 5 RCTs, 3812 participants, Analysis 4.6).

There was evidence that reducing SFA intake resulted in small reductions in **body weight** (MD -1.97 kg, 95% CI -3.67 to -0.27, I^2 = 72%, 6 RCTs, 4541 participants, Analysis 4.3), and **body mass index** (MD -0.50, 95% CI -0.82 to -0.19, I^2 = 55%, 6 RCTs, 5553 participants, Analysis 4.4).

Only one RCT reported assessing **quality of life**. The Women's Health Initiative (WHI 2006) assessed quality of life at baseline using the SF-36 tool). They found that being in the lower SFA arm resulted in a small improvement in Global Quality of Life at trial

close-out (on a scale of 0 worst to 10 best, MD 0.04, 95% CI 0.01 to 0.07, Analysis 4.7). This very small effect (less than 1% change) was statistically significant but unlikely to be relevant to individuals. However, it suggests no reduction in quality of life in those reducing their saturated fat.

Other results

To assess the effect in the population of consuming < 10%E as SFA relative to > 10%E as SFA for reduction in risk of noncommunicable diseases (NCDs), we combined studies with a control group saturated fat intake of > 10%E and an intervention group saturated fat intake of < 10%E for all-cause mortality, cardiovascular and coronary heart disease mortality and events, myocardial infarctions, non-fatal myocardial infarctions, and stroke. To assess the effect in the population of a reduction in %E from SFA from 10% in gradual increments relative to higher intake we repeated this with saturated fat cut-offs between 7%E and 13%E. The data for these cut-offs are shown in Table 5, and were plotted for a visual overview (Figure 6). The figure saturated fat intake was greater than 10%E in control groups, and less than 10%E in intervention groups.





Additional WHO NUGAG specific questions:

In adults what is the effect in the population of reduced percentage of energy (%E) intake from saturated fatty acids (SFA) relative to higher intake for reduction in risk of noncommunicable diseases (NCDs)?

We found that reducing saturated fat for at least two years suggested no clear effects on all-cause or cardiovascular mortality, but a 21% reduction in combined cardiovascular events. Heterogeneity in this result was partially explained by greater reductions in cardiovascular events in studies with greater serum total cholesterol reductions (implying greater reductions in SFA intake). Effects of reducing SFA on other cardiovascular and cancer outcomes were either very small or unclear (as the evidence was of very low quality), but it should be noted that risk ratios were all 1.0 or lower - no harm was indicated. Effects on NCD risk factors were small but positive (serum total cholesterol, LDL cholesterol, systolic and diastolic blood pressure, weight and BMI) or neutral (HDL cholesterol and TGs).

What is the effect of reducing SFA on coronary heart disease mortality and coronary heart disease events?

We found little or no effect of reducing SFA on non-fatal MI and CHD events, but the evidence on MI, stroke and CHD events was of very low quality. However, all risk ratios were less than 1.0.

What is the effect in the population of replacing SFA with PUFAs, MUFAs, CHO (refined versus unrefined), protein or trans fatty acids (TFAs) relative to no replacement for reduction in risk of NCDs?

We found greater reductions in cardiovascular events in studies that replaced saturated fats by PUFAs or CHO than in studies with replacement with MUFAs or protein, where there was little evidence of any effect.

What is the effect of replacing some saturated fat with PUFA on risk of CVD in adults?

There is moderate-quality evidence that replacing saturated fat with PUFA probably reduces the risk of CVD events. Replacing SFA with PUFA also appears to reduce the risk of total MI, non-fatal MI, stroke and CHD events, but has little or no effect on all-cause mortality, CVD mortality and CHD mortality.

What is the effect of replacing some saturated fat with MUFA on risk of CVD in adults?

The evidence for effects of replacing SFA with MUFA was very limited, so assessment of health effects was not possible.

What is the effect of replacing some saturated fat with CHO on risk of CVD in adults?

While studies that replaced SFA with CHO reduced CVD events and stroke, effects on all-cause mortality and other CVD outcomes suggested little or no effect.

What is the effect of replacing some saturated fat with protein on risk of CVD in adults?

There was no evidence suggesting that replacing SFA with protein reduced all-cause mortality or any CVD outcomes, but the evidence was limited.

Reduction in saturated fat intake for cardiovascular disease (Review) Copyright © 2020 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

What is the effect in the population of consuming < 10%E as SFA relative to > 10%E as SFA for reduction in risk of NCDs?

Cut-off data were difficult to interpret, and confidence intervals were wide, but they suggested greater reductions in cardiovascular events in studies where saturated fat intake was greater than 10%E in control groups, and less than 10%E in intervention groups (see Figure 6).

What is the effect in the population of a reduction in %E from SFA from 10% in gradual increments relative to higher intake for reduction in risk of NCDs?

The data from RCTs are too limited to be able to address this question.

DISCUSSION

Summary of main results

This systematic review of long-term randomised controlled trials of SFA reduction suggests that reducing saturated fat for at least two years probably has little or no effect on all-cause or cardiovascular mortality, but probably caused a 21% (95% CI 7 to 34%, $I^2 =$ 65%, moderate-quality evidence) reduction in people experiencing cardiovascular events. The heterogeneity in this relationship was explained by greater reduction in CVD events in trials with greater serum cholesterol lowering. This effect on cardiovascular events was retained in sensitivity analyses. Subgrouping suggested that there was a 27% (95% CI 8 to 42%) reduction in cardiovascular events in studies that replaced saturated fats by PUFAs, and a 16% (95% CI -6 to 33%) reduction in studies replacing with CHO, with little information on the effect of replacing with MUFAs or protein. The difference between subgroups was not statistically significant. We could not explore data on trans fats. Meta-regression and subgrouping suggested that greater reductions in SFA intake, greater reductions in total serum cholesterol levels, higher baseline SFA intake and greater increases in PUFA and MUFA intakes reduced CVD events more, but the strongest factor was the degree of cholesterol lowering. This clearly indicates that the cardiovascular effects of reducing saturated fat rely on changes in atherosclerosis via serum cholesterol. The degree of cholesterol lowering reflects greater reduction in SFA and greater increase in PUFA (Hegsted 2000).

There may be little or no effect of SFA reduction on non-fatal MI or CHD mortality (both low-quality evidence), and effects on fatal and non-fatal MI, stroke and CHD events were unclear (as the evidence was of very low-quality). However, risk ratios were less than 1.00 for all of these. While we found small reductions in body weight and body mass index with advice to reduce saturated fats, there was little or no effect on diabetes diagnoses, cancer diagnoses or cancer deaths, or on systolic or diastolic blood pressure.

Reducing saturated fat caused reductions in serum total and LDL cholesterol, which did not differ according to type of replacement. There was little or no effect of saturated fat reduction on serum HDL cholesterol or triglyceride. Data on lipid ratios, Lp(a) and HOMA were very limited and effects unclear, but SFA reduction appears to reduce glucose two hours after a glucose load.



Overall completeness and applicability of evidence

The review included adult participants at varying levels of risk of cardiovascular disease, men and women, with mean ages from 46 to 66 years at baseline, in free-living and institutional settings, and across the past 50 years. All the studies were conducted in industrialised countries, and no data were available from developing or transitional countries. The effectiveness of SFA reduction has been well assessed, with trials of at least 24 months including more than 50,000 participants for all primary and secondary CVD outcomes. Three thousand five hundred and eighteen participants in the included trials died, 1096 died of a cardiovascular cause, and 4476 experienced at least one cardiovascular event.

The external validity of the review in industrialised countries, men and women, people with low, moderate and high risk of cardiovascular disease was high, but it is not clear how this evidence relates to diets in developing and transitional countries.

Quality of the evidence

All 15 trials (16 comparisons) included were randomised controlled trials, allocation concealment was judged well done in eight RCTs and blinding of outcome assessors adequate in nine trials assessing CVD outcomes (and all trials assessing all-cause mortality). Blinding of participants is difficult and expensive in dietary fat trials, but was adequate in one trial. We judged incomplete outcome data not to be a problem in seven RCTs, and selective reporting was not a problem in any trial. Three trials were free of differences in care between the intervention and control arms, 10 RCTs stated an aim to reduce saturated fat, 11 showed evidence they had reduced SFA intake (all studies did one or the other), and nine studies showed clear reductions in total cholesterol. Five trials were at low summary risk of bias.

The lack of blinding of participants in most dietary trials is unlikely to alter outcome assessment when outcomes include death and cardiovascular events, but lack of blinding in the participants may have led those in the control groups to alter their lifestyle and dietary practices (for example, feeling that they have not been helped to reduce their cardiovascular risk, they may act to reduce their own risk by altering other lifestyle behaviours such as smoking or exercise, leading to a potential lessening of the apparent effect of the dietary intervention). Systematic differences in care between arms may have led to intervention groups receiving additional support in areas like self efficacy and gaining support from new social circles, potentially beneficial to health regardless of dietary fat intake, or gaining additional healthcare professional time, possibly leading to earlier diagnosis and treatment of other risk factors such as raised blood pressure. Additional dietary messages such as those around fruit and vegetable intake, fibre, alcohol and sugars, present in many studies, may have been protective, or may have diluted the effect or attainability or both of the saturated fat goals.

The quality of evidence balances the uncertainty over allocation concealment, lack of blinding and presence of systematic differences in care and additional dietary differences between arms (Figure 2) with the scale and consistency of the evidence across studies and across decades, despite very different designs and design flaws. For this reason, there is moderate-quality evidence

that the interventions that reduce dietary saturated fat intake reduce the risk of cardiovascular events.

Complex interventions

With complex interventions, such as dietary interventions, there are additional questions that need to be asked about included studies. Important issues to consider include defining the intervention, searching for and identifying all relevant studies, selecting studies for inclusion and data synthesis (Lenz 2007; Sheppherd 2009), as well as questions around whether the intended intervention was realised in study participants during the study.

For this review, we have worked to define the interventions clearly (see Characteristics of included studies), providing information on the type of intervention, stating the study aims and methods for each arm and the assessed total and saturated fat intakes attained within the study. However, while we have characterised the interventions, no two studies that reduced SFA had exactly the same dietary goals for the intervention groups. Methods of attaining the dietary goals varied from providing a whole diet over several years (in studies based in institutions) to providing advice on diet alongside supplementary foods such as margarines or oils, to providing dietary advice with or without supplementary support in the way of group sessions, cooking classes, shopping tours, feedback, self-efficacy sessions and/or individual counselling. We aimed to use this variety to support generalisability for the effects of the interventions.

We aimed to identify all the relevant studies through use of a broad search strategy, which was time-consuming. However, we believe that we have included most relevant trials. We also carefully defined acceptable interventions for each arm, to simplify decisions on inclusion, and the two independent assessors often agreed. We augmented data synthesis by subgrouping and meta-regression, to help us understand the effects of individual elements of dietary fat changes.

A study that sets out to assess the effect of a 30% reduction in saturated fat intake may attain this level of reduction in some participants, exceed it in some and not achieve it at all in others. The actual mean change attained in the intervention group may be less dramatic than that aimed for, and the participants in the control group may also reduce their saturated fat intake by a small amount, narrowing the difference in saturated fat between the groups further and so reducing the scale of any outcome. This can be dealt with in the systematic review if we meta-regress the difference in saturated fat intake between the intervention and control group with the scale of the outcome (assuming a linear dose response), still allowing us to understand the effect of altering saturated fat intake. However, it is difficult to measure actual saturated fat intake achieved. Some trials did not report it, either because they did not assess it, or did assess it but didn't report this relatively uninteresting outcome. Other trials did report the results of asking people what they were eating, using a food frequency questionnaire or several 24-hour food recalls. However, there is good evidence to believe that asking people how they are eating may produce somewhat biased information (Kristal 2005; Schatzkin 2003), and this may be a greater problem where the participant has been recently urged to eat in a particular way, as in a dietary trial. Assessment of change in total cholesterol is a way to get over self-reporting of dietary intake as reducing saturated

fat reduces total and LDL cholesterol. This review suggests that the relationship between saturated fat reduction and CVD events is moderated by the degree of cholesterol lowering, which is exactly what would be expected of a true effect.

The interventions used in the studies included in this review were varied, with some participants given all their food over a long period of time in an institutional setting, while most participants were given advice on how to achieve dietary changes, with or without the support of supplements such as oils and foods (Table 1). Advice was provided by a variety of health professionals, and with different levels of intensity. The effect of this was that different degrees of saturated fat reduction were achieved in different studies. The level of compliance with interventions involving long-term behaviour change, such as those used in these studies, can vary widely. This is likely to attenuate the pooled effect and bias it towards the null. Insofar as we were able to understand this issue, subgrouping and meta-regression suggested that greater reductions in saturated fats were associated with greater reductions in the risk of cardiovascular disease events. This suggestion of a dose response strengthens our belief that there is a true effect of reducing saturated fat on CVD events.

Potential biases in the review process

In compiling the included studies, we worked hard to locate randomised studies that altered dietary SFA intake for at least 24 months, even when cardiovascular events were not reported in study publications, or where such events were reported incidentally as reasons for participant dropouts. We attempted to contact all authors of potentially includable studies to verify the presence or absence of our outcomes. In many studies, no outcomes relevant to this review occurred or were recorded, and the numbers of events occurring within single studies varied from none to over 2500 deaths, over 500 cardiovascular deaths, and over 3000 participants experiencing at least one cardiovascular event (all within WHI 2006, the largest single study with almost 50,000 female participants for many years).

The number of cardiovascular deaths across the review was relatively small (1096), so while we can be quite confident in reporting a reduction in participants experiencing cardiovascular events (4476 events) with SFA reduction, and a lack of effect on total mortality (3518 deaths) within the studies' time scales, the effect on cardiovascular mortality is less clear. The risk ratio of 0.94 (95% CI 0.78 to 1.13, Analysis 1.18) may translate into a small protective effect, but this is unclear.

The lack of effect on individual cardiovascular events is harder to explain; there were 1714 people experiencing MIs, 1118 strokes and 1385 non-fatal MIs, 2472 cancer deaths, 3342 diabetes diagnoses and 5476 cancer diagnoses. Lack of clear effects on any of these outcomes is surprising, given the effects on total cardiovascular events, but may be due to the relatively short timescale of the included studies, compared to a usual lifespan during which risks of chronic illnesses develop over decades, and to relatively small reductions in saturated fat (and serum cholesterol) in some trials. There are difficulties in finding data on the number of people experiencing composite end points such as cardiovascular events. This end point represents the number of people experiencing any of the following: cardiovascular death, cardiovascular morbidity (non-fatal myocardial infarction, angina, stroke, heart failure, peripheral vascular events, atrial fibrillation) and unplanned cardiovascular interventions (coronary artery bypass surgery or angioplasty). Adding up the number of events is easy, but a single participant may have experienced a stroke, an MI and atrial fibrillation during a trial - and we need to take care not to count this individual three times. So finding such composite end point data involves using the best published composite end point data and supplementing this with author contact where possible. We have underestimated such composite end points rather than overestimated them where exact data are not available. Added to this complex picture, it needs to be remembered that definitions and diagnoses of some end points have altered over time.

Where the funnel plots and comparison of fixed- and randomeffects meta-analyses suggest small-study bias, we have downgraded the quality of the evidence in GRADE, but effects of any such small study bias appear small.

Some trials were partially confounded by aiming to make dietary changes other than those directly related to dietary fat intakes; for example, some studies encouraged intervention participants to make changes to their fat intake as well as changes to fruit and vegetable, fibre or salt intakes. In these studies, any effect on outcomes could be a result of other dietary changes, not of changes in saturated fat intake. The 11 studies (12 comparisons) that appeared free of such differences included Black 1994; DART 1989; Houtsmuller 1979; Ley 2004; MRC 1968; Oxford Retinopathy 1978; Rose corn oil 1965; Rose olive 1965; Simon 1997; Sydney Diet-Heart 1978; Veterans Admin 1969; WINS 2006. On the basis of reviewer comments, we assessed effects of reducing saturated fat intake on combined CVD events including only these trials free of additional interventions. Omitting trials with additional interventions (Oslo Diet-Heart 1966; STARS 1992; WHI 2006) leaves eight studies (nine arms) randomising 3998 participants of whom 750 experienced a CVD event, suggesting a similar reduction in CVD events (RR 0.80, 95% CI 0.64 to 0.99, I² = 48%, Analysis 1.43) to the main analysis (RR 0.79, 95% CI 0.66 to 0.93, $I^2 = 65\%$, > 53,000 participants randomised, Analysis 1.35). This suggests that effects on combined CVD events are not driven by interventions other than reductions in saturated fats and any energy replacements.

One surprising element of this review is the lack of new trials identified in the update, and small numbers of potential ongoing trials. This is likely to be because well-powered trials on cardiovascular end points will need to be large and carried out over several years, so expensive. As the effects of saturated fats are felt to be established and understood, trialists and funders may feel that the money would be better invested in answering other questions. For most of the ongoing trials, information is limited and these trials may or may not be included when fully published. Perhaps the current evidence set is as definitive as we will achieve during the 'statin era'.

Agreements and disagreements with other studies or reviews

In this review, saturated fat reduction had little or no effect on all-cause or cardiovascular mortality but did appear to reduce the risk of cardiovascular events by 21%, although effects on MI and stroke individually were less clear. This result was rather different from those of Siri-Tarino 2010, who systematically reviewed cohort studies that assessed relationships between saturated fat and cardiovascular events. They included 21 studies and did not find associations between saturated fat intake and cardiovascular



disease (RR 1.0, 95% CI 0.89 to 1.11). However, this meta-analysis has been criticised (Katan 2010; Scarborough 2010; Stamler 2010), as results of half of the studies included in their meta-analysis were adjusted for serum cholesterol concentrations, while there is an established relation between saturated fat intake and cholesterol level. The studies included in the meta-analysis also varied widely in the method used to assess intake, as half of the studies collected one-day intake data. However, as with our review, they found no relationship between saturated fat intake and coronary heart disease (RR 1.07, 95% CI 0.96 to 1.19) or stroke (RR 0.81, 95% CI 0.62 to 1.05, Siri-Tarino 2010).

In this review, we found that replacing saturated fat with PUFAs (a modified-fat diet) appeared more protective of cardiovascular events than replacement with carbohydrates (a low-fat diet, Analysis 1.44; Analysis 1.45). This was similar to results within our closely allied systematic review assessing health effects of total fat reduction, where modified-fat diets were protective and low-fat diets were not (Hooper 2012). Meta-regression did not suggest any relationship between either PUFAs or MUFAs and cardiovascular events in this review, although the analysis was underpowered. Alonso 2006 suggested a protective role for MUFA from olive oil, but not from meat sources (the main source of MUFA in the USA and Northern Europe). Our systematic review was not able to explore this issue as we included only one small study (underpowered to assess health outcomes on its own) that replaced SFA with MUFA, using an olive oil supplement (Rose olive 1965). A review by Mozaffarian 2010, which again included very similar studies to the last version of this review, with the Finnish Mental Hospital study and Women's Health Initiative data added, stated that their findings provided evidence that consuming PUFAs in place of saturated fat would reduce coronary heart disease. However, their evidence for this was limited, as they found that modifying fat reduced the risk of myocardial infarction or coronary heart disease death (combined) by 19% (similar to our result). As the mean increase in PUFAs in these studies was 9.9% of energy, they infer an effect of increasing PUFAs by 5% of energy of 10% reduction in risk of myocardial infarction or coronary heart disease death. They provided no suggestion or evidence of a relationship between degree of PUFAs increase and level of risk reduction. Another review carried out during updating of the Nordic Nutritional Recommendations (Schwab 2014) included observational as well as intervention studies, and concluded that there was convincing evidence that partial replacement of SFA with PUFA decreases risk of CVD while replacement with CHO is associated with increased CVD risk. The review included studies performed solely in white participants or with a clear white majority.

Within the meta-regression, we hoped to combine studies that effectively altered saturated fat by different degrees, so that studies that reduced saturated fat very little and studies that reduced it a great deal would all offer data points for the meta-regression against mortality and morbidity end points, and similarly for total fat, polyunsaturated, monounsaturated and trans fats. Unfortunately many of the included studies did not report data on assessed dietary intake during the trial, reducing the quantity of data available to understand the relationships. Another limitation in understanding effects of individual classes of fatty acids on mortality and morbidity (both in trials and in observational studies) was our ability to correctly assess participants' intake. We could overcome this by using biomarkers such as serum LDL cholesterol (differences between the LDL concentration in the intervention and control arms could be seen as a reasonable and independent approximation of saturated fat intake); however, as many studies were carried out in the 1960s to 1990s, few measured and reported LDL cholesterol. We used meta-regression with serum total cholesterol (although this is a composite marker and so less related to saturated fat intake), but although this was available for more studies than LDL it was not available for all studies. Despite the limited data, there was a clear suggestion from meta-regression that there was greater reduction of risk of cardiovascular events in studies with greater total serum cholesterol reduction, supporting the central role of serum lipids in the link between dietary saturated fats and cardiovascular events.

Participants' level of risk

As the rate of events is higher in high-risk groups (by definition), it should require smaller sample sizes and shorter follow-up to observe an effect of an intervention in a high-risk group of participants (Davey Smith 1993). There have been suggestions that randomised controlled trials are unsuitable for assessing the effectiveness of interventions, with very modest levels of effect in low-risk populations, because of the huge numbers of personyears of observation needed to gain sufficient statistical power to avoid Type II errors (Ebrahim 1997). However, with the publication of the Women's Health Initiative trial (WHI 2006) we now have data on more people experiencing cardiovascular events who were originally at low risk of cardiovascular disease than in people with moderate or high risk. The same is true for cardiovascular deaths and total mortality.

When end points such as total mortality are used, the situation becomes more difficult, as in low-risk groups the proportion of deaths which are unrelated to cardiovascular disease (and perhaps unlikely to be influenced by dietary fat changes) rises, again diluting any differences in the numbers of deaths between intervention and control groups. It is more likely that changes in cardiovascular deaths will be seen than in total mortality. The trend is certainly in this direction, with the pooled risk ratio for total mortality 0.96 (95% CI 0.90 to 1.03, Analysis 1.1), and for cardiovascular mortality RR 0.94 (95% CI 0.78 to 1.13, Analysis 1.18). Our best estimate is that SFA reduction results in a reduction of 6% in deaths due to cardiovascular disease, and a reduction of 4% in total deaths, but these are small effects with wide confidence intervals.

The high-risk participants all showed evidence of cardiovascular disease at baseline. Under current guidelines, most high-risk participants with raised lipid levels should be on lipid-lowering medication (Grundy 2019; NICE 2014; O'Gara 2014). This raises the question of whether there is any additional advantage of adherence to a reduced SFA diet in addition to statin therapy. Little evidence exists at present to answer this question. However, in all parts of the world where drug budgets are restricted and use of lipid-lowering medication remains rationed even for those at high risk, the use of reduced SFA diets would appear to be a cost-effective option leading to considerable reductions in cardiovascular events for populations (and so in health budgets) in only a few years.

Low-risk participants are unlikely to be on lipid-lowering medication under current guidelines. The suggestion of protection of low-risk individuals from cardiovascular events with a reduction of roughly 21% of events in just a few years of intervention, as there is no evidence that effects in the low-CVD-risk group are different from effects in the higher-risk groups, would appear to merit



continued public health action. Recent guidelines recommend saturated fat reduction in general populations (SACN 2019).

A factor that may affect participant risk of cardiovascular disease, and also the effectiveness of reducing saturated fat intake, that has altered over time is the level of use of statins to control serum lipids in people at moderate and high risk of CVD. The 4S 1994 trial, which was the first trial to show that use of statins could reduce mortality in people with coronary heart disease, was published in 1994 and led to an explosion of the use of statins. For most health outcomes, we saw no clear effect of a decade of publication on risk, but for combined CVD events and CHD events, there were differences between subgroups. For combined CVD events, there were reductions in risk with reduced saturated fat intakes in the 1960s, 1970s and 1990s (both trials published early in the decade), but no clear effect of reducing saturated fat in the 1980s (one trial with 283 events) or 2000s (three large trials). It is possible (but not clear) that participants in the trials published in the 2000s were protected by higher levels of statin use (statins were allowed in participants in the largest trial, WHI 2006).

AUTHORS' CONCLUSIONS

Implications for practice

Evidence supports the reduction of saturated fat to reduce risk of combined cardiovascular events in people with and without existing cardiovascular disease, in men and women, over at least two years and in industrialised countries. Little or no effect of saturated fat reduction was seen on all-cause and cardiovascular mortality, at least on this timescale.

Practical ways to achieve reductions in dietary saturated fat include switching to lower fat dairy foods and cutting off meat fats, as well as reducing intake of foods high in saturated fats such as cakes, biscuits, pies and pastries, butter, ghee, lard, palm oil, sausages and cured meats, hard cheese, cream, ice cream, milkshakes and chocolate (for further details see NHS 2020).

Implications for research

To complement this review of long-term RCTs, we need reviews of metabolic studies to clarify the effects of specific replacements for saturated fat in the diet, and systematic reviews of cohort studies to clarify longer-term effects of saturated fat reductions.

The financial implications (costs and savings) of appropriate advice and legislation to modify fat intake in those at various levels of cardiovascular risk should be assessed and reflected in health policy. Whilst interventions to alter dietary fat intake in individuals at high cardiovascular risk have been fairly successful, such health promotion initiatives in the general population have been less successful. Further work is needed to help high- and low-risk individuals to make effective changes to reduce saturated fat and to maintain these changes over their lifetimes. Research into the effects of legislation to alter fat contents of foods, improved labelling, pricing initiatives and improved availability of healthier foods, linking food production and processing into the health agenda, may yield huge advances in this area.

It is not clear whether there is an additional benefit of reducing saturated fat in those at high risk of cardiovascular disease who are on lipid-lowering medication. Further research to examine the need for maintenance of reduced saturated fat whilst on lipid-lowering medication would be useful, but not as useful as understanding specific dietary fat replacements for saturated fat. However, we did not identify any relevant ongoing trials in our searches.

All future trials should be of at least 2 years duration (preferably longer), employ excellent methodology in terms of randomisation and allocation concealment, blinding of outcome assessors, highquality assessment of macronutrients and micronutrients during the trial in both arms, and equivalent attention and health professional time to participants in both arms.

ACKNOWLEDGEMENTS

We gratefully acknowledge the help of the following investigators in providing information about their own and others' trials (whether eventually included or excluded): H Arnesen (Ullevål University Hospital), AV Astrup (University of Copenhagen), K Aziz (NICVD, Karachi), F Azizi (Research Institute for Endocrine Sciences, Tehran), SAA Beresford (University of Washington), HS Black (Baylor College of Medicine), BPM Bloemberg (National Institute of Public Health and Environmental Protection, Netherlands), DJ Bowen (Fred Hutchinson Cancer Research Center, Seattle), NF Boyd (University of Toronto), GA Bray (Pennington Biomedical Research Center, Baton Rouge), ML Burr (University of Wales), G Carruba (University of Palermo), L Castagnetta (University of Palermo), JL Curzio (University of Glasgow), RF DeBusk (Stanford University), C Defoort (Méditerranée University), Z Djuric (Wayne State University), A Due (University of Copenhagen), S Druker (University of Massachusetts), RPF Dullaart (University Hospital, Groningen), GE Eyssen (University of Toronto), TM Hayes (University Hospital of Wales, Cardiff), JA Heady (retired, formerly of MRC Social Medicine Research Unit), J Hebert (University of South Carolina), RJ Heine (Free University Hospital, Amsterdam), M-L Hellenius (Karolinska Institute), RF Heller (University of Newcastle), TDR Hockaday (retired, formerly Radcliffe Infirmary), L-E Holm (Swedish Radiation Protection Institute), DJ Hyman (Baylor College of Medicine, Houston), AFM Kardinaal (University of Wageningen), F Khan (Ninewells Hospital and Medical School, Dundee), RH Knopp (University of Washington), D Lairon (Méditerranée University), MEJ Lean (University of Glasgow), B Leelarthaepin (University of Sydney), P Leren (University of Oslo), A Lindman (University of Oslo), S Mackey (Stanford University), R MacLennan (retired, formerly of Queensland Institute of Medical Research), F Macrae (Royal Melbourne Hospital), JI Mann (University of Otago), J Marniemi (Social Insurance Institute), K McManus (Harvard Medical School), RP Mensink (Maastricht University), PA Metcalf (University of Auckland), A Michalsen (University Duisburg-Essen), TF Moy (Johns Hopkins University), AR Ness (University of Bristol), I Okene (University of Massachusetts), GS Oostenbrug (Maastricht University), J Pierce (University of California, San Diego), SD Poppitt (University of Auckland), RJ Reber (University of Illinois),

JE Reseland (University of Oslo), BM Retzlaff (University of Washington), AA Rivellese (Federico II University, Naples), P Roderick (University of Southampton), DP Rose (American Health Foundation), FM Sacks (Harvard School of Public Health), WHM Saris (University of Maastricht), ES Sarkkinen (University of Kuopio), A Schatzkin (National Cancer Institute), B Seppelt (German Institute of Human Nutrition), MS Simon (Wayne State University), B Smith (University of Kentucky), E Søndergaard (Svendborg Hospital, Svendborg), AS St. Leger (University of Manchester), VJ Stevens (Kaiser Permanente Centre for Health Research), A Stoddard (University of Massachusetts), LP Svetkey (Duke



University Medical Center), LC Tapsell (University of Wollongong), BC Tilley (Medical University of South Carolina), H van den Berg (TNO Nutrition and Food Research Institute), W van Herpen (Unilever), K van het Hof (Unilever, Rotterdam), MW Verheijden (Wageningen University), GF Watts (University Hospital of Perth), AS Wierzbichi (St. Thomas's Hospital, London), PT Williams (Stanford University), RR Wing (University of Pittsburgh), PD Wood (Stanford University), I Zazpe (University of Navarra), PL Zock (Wageningen Centre for Food Studies).

Thank you to the peer reviewers who have contributed to this review update: Dr Dimitrios Koutoukidis of Nuffield Department of Primary Care Health Sciences, Zhou-Qing Kang of Liaoning Provincial Hospital of Geriatrics, Kristina Petersen of Pennsylvania State University, and the additional referee (who preferred to remain anonymous) as well as to the Cochrane editors Sarah Hodgkinson and Karen Rees, and Charlene Bridges who expertly ran our searches.

We also gratefully acknowledge the expertise and help of the following who contributed to previous iterations of this systematic review: S Adams (Royal Free Hospital, London), B Anagnostelis (Royal Free Hospital, London), M Brand (Cochrane Hypertension Group), R Clarke (University of Oxford), D Darrah-Morgan (Russian translation), A Donner (University of Western Ontario), D Fagard (University of East Anglia for duplication of inclusion and data extraction), Shweta Gidwani (University of Manchester), G Gubitz (Cochrane Stroke Group), M Haugh (Cochrane Renal Group), IU Haq (Northern General Hospital, Sheffield), J Hooper (Danish, Swedish and Norwegian translation), BK Hurley (Italian translation), L Jones (Systematic Reviews Training Unit, London), SPH Keen (Cochrane Diabetes Group), S Logan (Systematic Reviews Training Unit, London), LI Mennen (INSERM), T Moore (Cochrane Heart Group), J Muscroft (German and French translation), HL Newmark (Rutgers), E Royle (Cochrane Peripheral Vascular Diseases Group), I Tumur (Pfizer Ltd), AS Truswell (University of Sydney), M Turner (Chinese translation), JM Walsh (University of California), A Wierzbicki (St. Thomas' Hospital, London), WC Willett (Harvard School of Public Health), AF Winder (University of London).

Many thanks to those people who contributed to, and were co-authors of, previous versions of this review (Hooper 2000; Hooper 2001; Hooper 2012; Hooper 2015a): Julian PT Higgins, Shah Ebrahim, George Davey Smith, Rudolph Riemersma, Nigel E Capps, Carolyn Summerbell, Rachel Thompson, Helen Moore, Diredre Sills, Felicia Roberts, Dorotheé Fagard and Gillian Clements.

Finally, many thanks for the members of the WHO NUGAG committee who took time to debate and discuss this review, and ensure that it was addressing the questions that WHO NUGAG were addressing. Members of this NUGAG subcommittee included Nahla C Houalla (American University of Beirut), Jim Mann (University of Otago), Barbara O Schneeman (University of California at Davis), Mary R L'Abbé (University of Toronto), Murray Skeaff (University of Otago), Due Li (Zhejiang University), Russell de Souza (McMaster University), Ibrahim Elmadfa (University of Vienna). Thank you too to all of the NUGAG group members, and members of WHO including Jason Montez, Chizuru Nishida and Emma Kennedy.

REFERENCES

References to studies included in this review

Black 1994 {published and unpublished data}

* Black HS, Herd JA, Goldberg LH, , Thornby JI, Rosen T, et al. Effect of a low-fat diet on the incidence of actinic keratosis. *New England Journal of Medicine* 1994; **330**(18):1272-5.

Black HS, Thornby JI, Goldberg LH, Herd JA, Rosen T, et al. Evidence that a low-fat diet reduces the occurrence of nonmelanoma skin cancer. *International Journal of Cancer* 1995; **62**(2):165-9.

Jaax S, Scott LW, , Thornby JI, Black HS. General guidelines for a low-fat diet effective in the management and prevention of nonmelanoma skin cancer. *Nutrition and Cancer* 1997; **27**(2):150-6.

DART 1989 {published and unpublished data}

* Burr ML, Fehily AM, Gilbert JF, Rogers S, Holliday RM, Sweetnam PM, et al. Effects of changes in fat, fish, and fibre intakes on death and myocardial reinfarction: diet and reinfarction trial (DART). *Lancet* 1989; **2**(8666):757-61.

Burr ML, Fehily AM, Rogers S, Welsby E, King S, Sandham S. Diet and reinfarction trial (DART): design, recruitment, and compliance. *European Heart Journal* 1989; **10**(6):558-67.

Burr ML, Fehily AM. Fish and the heart. Lancet 1989; ii:1450-2.

Burr ML, Holliday RM, Fehily AM, Whitehead PJ. Haematological prognostic indices after myocardial infarction: evidence from the diet and reinfarction trial (DART). *European Heart Journal* 1992; **13**(2):166-70.

Burr ML, Sweetham PM, Fehily AM. Diet and reinfarction [letter]. *European Heart Journal* 1994; **15**(8):1152-3.

Fehily AM, Vaughan-Williams E, Shiels K, Williams AH, Horner M, Bingham G, et al. The effect of dietary advice on nutrient intakes: evidence from the diet and reinfarction trial (DART). *Journal of Human Nutrition & Dietetics* 1989; **2**(4):225-5.

Houtsmuller 1979 {published data only}

Houtsmuller AJ, Van Hal-Ferwerda J, Zahn KJ, Henkes HE. Favourable influences of linoleic acid on the progression of diabetic micro- and macroangiopathy. *Nutrition and Metabolism* 1980; **24**(Suppl 1):105-18.

Houtsmuller AJ, Van Hal-Ferwerda J, Zahn KJ, Henkes HE. Influence of different diets on the progression of diabetic retinopathy. *Progress in Food Nutritition and Science* 1980; **4**(5):41-6.

* Houtsmuller AJ, Zahn KJ, Henkes HE. Unsaturated fats and progression of diabetic retinopathy. *Documenta Ophthalmologia* 1979; **48**(2):363-71.

Ley 2004 {published and unpublished data}

* Ley SJ, Metcalf PA, Scragg RK, Swinburn BA. Long-term effects of a reduced fat diet intervention on cardiovascular disease

risk factors in individuals with glucose intolerance. *Diabetes Research and Clinical Practice* 2004; **63**(2):103-12.

Swinburn BA, Metcalf PA, Ley SJ. Long-term (5-year) effects of a reduced-fat diet intervention in individuals with glucose intolerance. *Diabetes Care* 2001; **24**(4):619-24.

Swinburn BA, Woollard GA, Chang EC, Wilson MR. Effects of reduced-fat diets consumed ad libitum on intake of nutrients particularly antioxidant vitamins. *Journal of the American Dietetic Association* 1999; **99**(11):1400-5.

Moy 2001 {published and unpublished data}

Moy TF, Yanek LR, Raqueno JV, Bezirdjian PJ, Blumenthal RS, Wilder LB, et al. Dietary counseling for high blood cholesterol in families at risk of coronary disease. *Preventive Cardiology* 2001; **4**(4):158-64.

MRC 1968 {published and unpublished data}

Ederer F, Leren P, Turpeinen O, . Cancer among men on cholesterol lowering diets: experience of five clinical trials. *Lancet* 1971; **2**(7717):203-6.

Heady JA. Are PUFA harmful? *British Medical Journal* 1974; **1**(898):115-6.

* . Controlled trial of soya-bean oil in myocardial infarction. *Lancet* 1968; **2**(570):693-9.

Oslo Diet-Heart 1966 {published and unpublished data}

Leren P. Prevention of coronary heart disease, some results from the Oslo secondary and primary intervention studies. *Journal of the American College of Nutrition* 1989; **8**(5):407-10.

Leren P. The Oslo diet-heart study. Eleven year report. *Circulation* 1970; **42**(5):935-42.

Leren P. The effect of a cholesterol lowering diet in male survivors of myocardial infarction. A controlled clinical trial) [Virkningen av cholesterolsenkende diett hos menn som har gjennomgatt hjerteinfarkt. Et kontrollert klinisk fors/ok]. *Nordisk Medicin* 1967; **77**(21):658-61.

* Leren P. The effect of plasma cholesterol lowering diet in male survivors of myocardial infarction. A controlled clinical trial. *Acta Medica Scandinavia* 1966; **466 Suppl**:1-92.

Leren P. The effect of plasma-cholesterol-lowering diet in male survivors of myocardial infarction. A controlled clinical trial. *Bulletin of the New York Academy of Medicine* 1968; **44**(8):1012-20.

Oxford Retinopathy 1978 {published and unpublished data}

Coppack SW, Doll HA, Pim B, Hockaday TD. Intravenous glucose tolerance and mortality in non-insulin-dependant diabetes mellitus. *Quarterly Journal of Medicine* 1990; **75**(277):451-60.

Hillson RM, Hockaday TD, Mann JI, Newton DJ. Hyperinsulinaemia is associated with development of ECG abnormalities in diabetics. *Diabetes Research* 1984; **1**(3):143-9.


* Hockaday TD, Hockaday JM, Mann JI, Turner RC. Prospective comparison of modified fat-high-carbohydrate with standard low-carbohydrate dietary advice in the treatment of diabetes: one year follow-up study. *British Journal of Nutrition* 1978; **39**(2):357-62.

Howard-Williams J, Patel P, Jelfs R, Carter RD, Awdry P, Bron A, et al. Polyunsaturated fatty acids and diabetic retinopathy. *British Journal of Ophthalmology* 1985; **69**(1):15-8.

Lopez-Espinoza I, Howard WJ, Mann JI, Carter RD, Hockaday TD. Fatty acid composition of platelet phospholipids in non-insulindependent diabetics randomized for dietary advice. *British Journal of Nutrition* 1984; **52**(1):41-7.

Rose corn oil 1965 {published data only}

Rose GA, Thomson WB, Williams RT. Corn oil in treatment of ischaemic heart disease. *British Medical Journal* 1965; **1**(5449):1531-3.

Rose olive 1965 {published data only}

Rose GA, Thomson WB, Williams RT. Corn oil in treatment of ischaemic heart disease. *British Medical Journal* 1965; **1**(5449):1531-3.

Simon 1997 {published and unpublished data}

Djuric Z, Heilbrun LK, Reading BA, Boomer A, Valeriote FA, Martino S. Effects of a low fat diet on levels of oxidative damage to DNA in human peripheral nucleated blood cells. *Journal of the National Cancer Institute* 1991; **83**(11):766-9.

Djuric Z, Martino S, Heilbrun LK, Hart RW. Dietary modulation of oxidative DNA damage. *Advances In Experimental Medicine and Biology* 1994; **354**:71-83.

Kasim SE, Martino S, , Khilnani S, Boomer A, Depper J, et al. Dietary and anthropometric determinants of plasma lipoproteins during a long-term low-fat diet in healthy women. *American Journal of Clinical Nutrition* 1993; **57**(2):146-53.

* Simon MS, Heilbrun LK, Boomer A, Kresge C, Depper J, Kim PN, et al. A randomised trial of a low-fat dietary intervention in women at high risk for breast cancer. *Nutrition and Cancer* 1997; **27**(2):136-42.

STARS 1992 {published and unpublished data}

Blann AD, Jackson P, Bath PM, Watts GF. Von Willebrand factor, a possible indicator of endothelial cell damage, decreases during long-term compliance with a lipid-lowering diet. *Journal of Internal Medicine* 1995; **237**(6):557-61.

Watts GF, Brunt JN, Coltart DJ, Lewis B. The St. Thomas Atherosclerosis Regression Study (STARS). *Atherosclerosis* 1992; **97**:231.

Watts GF, Jackson P, Burke V, Lewis B. Dietary fatty acids and progression of coronary artery disease in men. *American Journal of Clinical Nutrition* 1996; **64**(2):202-9.

Watts GF, Jackson P, Mandalia S, Brunt JN, Lewis ES, Coltart DJ, et al. Nutrient intake and progression of coronary artery disease. *American Journal of Cardiology* 1994; **73**(5):328-32.

* Watts GF, Lewis B, Brunt JN, Lewis ES, Coltart DJ, Smith LD, et al. Effects on coronary artery disease of lipid-lowering diet, or diet plus cholestyramine, in the St Thomas' Atherosclerosis Regression Study (STARS). *Lancet* 1992; **339**(8793):563-9.

Watts GF, Lewis B, Brunt JNH, Swan AV. Coronary atheroma regression trials. *Lancet* 1992; **339**(i):1241-3.

Watts GF, Lewis B, Jackson P, Burke V, Lewis ES, Brunt JN, et al. Relationships between nutrient intake and progression/ regression of coronary atherosclerosis as assessed by serial quantitative angiography. *Canadian Journal of Cardiology* 1995; **11**(Suppl G):110G-4G.

Watts GF, Mandalia S, Brunt JN, Slavin BM, Coltart DJ, Lewis B. Independent associations between plasma lipoprotein subfraction levels and the course of coronary artery disease in the St. Thomas' Atherosclerosis Regression Study (STARS). *Metabolism: Clinical and Experimental* 1993; **42**(11):1461-7.

Watts GF, Mandalia S, Slavin BM, Brunt JN, Coltart DJ, Lewis B. Metabolic determinants of the course of coronary artery disease in men. *Clinical Chemistry* 1994; **40**(12):2240-6.

Watts GF. Nutritional, metabolic, and genetic determinants of the progression of coronary heart disease. STARS Group. *Journal of Cardiovascular Pharmacology* 1995; **25**(Suppl 4):S11-9.

Sydney Diet-Heart 1978 {published and unpublished data}

Blacket RB, Leelarthaepin B, McGilchrist C, Palmer AJ, Woodhill JM. The synergistic effect of weight loss and changes in dietary lipids on the serum cholesterol of obese men with hypercholesterolaemia: implications for prevention of coronary heart disease. *Australian and New Zealand Journal of Medicine* 1979; **9**(5):521-9.

Ramsden CE, Zamora D, Leelarthaepin B, Majchrzak-Hong SF, Faurot KR, Suchindran CM, et al. Use of dietary linoleic acid for secondary prevention of coronary heart disease and death: evaluation of recovered data from the Sydney Diet Heart Study and updated meta-analysis. *BMJ* 2013; **346**:e8707. [DOI: 10.1136/bmj.e8707]

* Woodhill JM, Palmer AJ, Leelarthaepin B, McGilchrist C, Blacket RB. Low fat, low cholesterol diet in secondary prevention of coronary heart disease. *Advances in Experimental Medicine and Biology* 1978; **109**:317-30.

Veterans Admin 1969 {published data only}

Dayton S, Hashimoto S, Dixon W, Pearce ML. Composition of lipids in human serum and adipose tissue during prolonged feeding of a diet high in unsaturated fat. *Journal of Lipid Research* 1966; **7**(1):103-11.

Dayton S, Hashimoto S, Pearce ML. Adipose tissue linoleic acid as a criterion of adherence to a modified diet. *Journal of Lipid Research* 1967; **8**(5):508-10.

Dayton S, Hashimoto S, Pearce ML. Influence of a diet high in unsaturated fat upon composition of arterial tissue and atheromata in man. *Circulation* 1965; **32**(6):911-24.



Dayton S, Hashimoto S, Rosenblum D, Pearce M. Vitamin E status of humans during prolonged feeding of unsaturated fats. *Journal of Laboratory and Clinical Medicine* 1965; **65**(5):739-47.

Dayton S, Pearce ML, Goldman H, Harnish A, Plotkin D, Shickman M, et al. Controlled trial of a diet high in unsaturated fat for prevention of atherosclerotic complications. *Lancet* 1968; **2**(577):1060-2.

* Dayton S, Pearce ML, Hashimoto S, Dixon WJ, Tomayasu U. A controlled clinical trial of a diet high in unsaturated fat in preventing complications of atherosclerosis. *Circulation* 1969; **15**(1, Suppl 2):II-1-63.

Dayton S, Pearce ML, Hashimoto S, Fakler LJ, Hiscock E, Dixon WJ. A controlled clinical trial of a diet high in unsaturated fat. *New England Journal of Medicine* 1962; **266**:1017-23.

Dayton S, Pearce ML. Diet and atherosclerosis. *Lancet* 1970; **1**(644):473-4.

Dayton S, Pearce ML. Diet and cardiovascular diseases. *Lancet* 1969; **1**(584):51-2.

Dayton S, Pearce ML. Diet high in unsaturated fat: a controlled clinical trial. *Minnesota Medicine* 1969; **52**(8):1237-42.

Dayton S, Pearce ML. Prevention of coronary heart disease and other complications of atherosclerosis by modified diet. *American Journal of Medicine* 1969; **46**(5):751-62.

Dayton S, Pearce ML. Trial of unsaturated-fat diet. *Lancet* 1968; **2**(581):1296-7.

Hiscock E, Dayton S, Pearce ML, Hashimoto S. A palatable diet high in unsaturated fat. *Journal of the American Dietetic Association* 1962; **40**:427-31.

Pearce ML, Dayton S. Incidence of cancer in men on a diet high in polyunsaturated fat. *Lancet* 1971; **1**(697):464-7.

Sturdevant RA, Pearce ML, Dayton S. Increased prevalence of cholelithiasis in men ingesting a serum-cholesterol-lowering diet. *New England Journal of Medicine* 1973; **288**(1):24-7.

Tompkins MJ, Dayton S, Pearce ML. Effect of long-term feeding of various fats on whole blood clotting times in men. *Journal of Laboratory and Clinical Medicine* 1964; **64**(5):763-72.

WHI 2006 {published data only} ZZZ <accessionId ref="info:x-wiley/clinicalTrialsGov/ NCT00000611"> ZZZNCT00000611

Anderson G, Cummings S, Freedman LS, Furberg C, Henderson M, Johnson SR, et al. Design of the Women's Health

Initiative clinical trial and observational study. *Controlled Clinical Trials* 1998; **19**(1):61-109.

Anderson GL, Manson J, Wallace R, Lund B, Hall D, Davis S, et al. Implementation of the Women's Health Initiative study design. *Annals of Epidemiology* 2003; **13**(9 Suppl):S5-17.

Beresford SA, Johnson KC, Ritenbaugh C, Lasser NL, Snetselaar LG, Black HR, et al. Low-fat dietary pattern and risk of colorectal cancer: the Women's Health Initiative Randomized Controlled Dietary Modification Trial. *JAMA* 2006; **295**(6):643-54. Bowen D, Ehret C, Pedersen M, Snetselaar L, Johnson M, Tinker L, et al. Results of an adjunct dietary intervention program in the Women's Health Initiative. *Journal of the American Dietetic Association* 2002; **102**(11):1631-7.

Carty CL, Kooperberg C, Neuhouser ML, Tinker L, Howard B, Wactawski-Wende J, et al. Low-fat dietary pattern and change in body-composition traits in the Women's Health Initiative Dietary Modification Trial. *American Journal of Clinical Nutrition* 2011; **93**(3):516-24.

Curb JD, McTiernan A, Heckbert SR, Kooperberg C, Stanford J, Nevitt M, et al. Outcomes ascertainment and adjudication methods in the Women's Health Initiative. *Annals of Epidemiology* 2003; **13**(9 Suppl):S122-8.

Hays J, Hunt JR, Hubbell FA, Anderson GL, Limacher M, Allen C, et al. The Women's Health Initiative recruitment methods and results. *Annals of Epidemiology* 2003; **13**(9 Suppl):S18-77.

Hebert JR, Patterson RE, Gorfine M, Ebbeling CB, St Jeor ST, Chlebowski RT, et al. Differences between estimated caloric requirements and self-reported caloric intake in the Women's Health Initiative. *Annals of Epidemiology* 2003; **13**(9):629-37.

Howard BV, Curb JD, Eaton CB, Kooperberg C, Ockene J, Kostis JB, et al. Low-fat dietary pattern and lipoprotein risk factors: the Women's Health Initiative dietary modification trial. *American Journal of Clinical Nutrition* 2010; **91**(4):860-74.

Howard BV, Manson JE, Stefanick ML, Beresford SA, Frank G, Jones B, et al. Low-fat dietary pattern and weight change over 7 years: the Women's Health Initiative Dietary Modification Trial. *JAMA* 2006; **295**(1):39-49.

Howard BV, Van Horn L, Hsia J, Manson JE, Stefanick ML, Wassertheil-Smoller S, et al. Low-fat dietary pattern and risk of cardiovascular disease: the Women's Health Initiative Randomized Controlled Dietary Modification Trial. *JAMA* 2006; **295**(6):655-66.

Howard BV. Dietary fat and cardiovascular disease: putting the Women's Health Initiative in perspective. *Nutrition Metabolism & Cardiovascular Diseases* 2007; **17**(3):171-4.

Neuhouser ML, Tinker L, Shaw PA, Schoeller D, Bingham SA, Horn LV, et al. Use of recovery biomarkers to calibrate nutrient consumption self-reports in the Women's Health Initiative. *American Journal of Epidemiology* 2008; **167**(10):1247-59.

Patterson RE, Kristal A, Rodabough R, Caan B, Lillington L, Mossavar-Rahmani Y, et al. Changes in food sources of dietary fat in response to an intensive low-fat dietary intervention: early results from the Women's Health Initiative. *Journal of the American Dietetic Association* 2003; **103**(4):454-60.

Patterson RE, Kristal AR, Tinker LF, Carter RA, Bolton MP, Gurs-Collins T, et al. Measurement characteristics of the Women's Health Initiative food frequency questionnaire. *Annals of Epidemiology* 1999; **9**(3):178-87.

Prentice RL, Aragaki AK, Howard BV, Chlebowski RT, Thomson CA, Van Horn L, et al. Low-fat dietary pattern among postmenopausal women influences long-term cancer,



cardiovascular disease, and diabetes outcomes. *Journal of Nutrition* 2019; **149**(9):1565-74.

* Prentice RL, Aragaki AK, Van Horn L, Thomson CA, Beresford SAA, Robinson J, et al. Low-fat dietary pattern and cardiovascular disease: results from the Women's Health Initiative randomized controlled trial. *American Journal of Clinical Nutrition* 2017; **106**(1):35-43.

Prentice RL, Caan B, Chlebowski RT, Patterson R, Kuller LH, Ockene JK, et al. Low-fat dietary pattern and risk of invasive breast cancer: the Women's Health Initiative Randomized Controlled Dietary Modification Trial. *JAMA* 2006; **295**(6):629-42.

Prentice RL, Thomson CA, Caan B, Hubbell FA, Anderson GL, Beresford SA, et al. Low-fat dietary pattern and cancer incidence in the Women's Health Initiative Dietary Modification Randomized Controlled Trial. *Journal of the National Cancer Institute* 2007; **99**(20):1534-43.

Ritenbaugh C, Patterson RE, Chlebowski RT, Caan B, Fels-Tinker L, Howard B, et al. The Women's Health Initiative Dietary Modification trial: overview and baseline characteristics of participants. *Annals of Epidemiology* 2003; **13**(9 Suppl):S87-97.

Robinson JG, Wallace R, Safford MM, Pettinger M, Cochrane B, Ko MG, et al. Another treatment gap: restarting secondary prevention medications: the Women's Health Initiative. *Journal* of Clinical Lipidology 2010; **4**(1):36-45.

Rossouw JE, Finnegan LP, Harlan WR, Pinn VW, Clifford C, McGowan JA. The evolution of the Women's Health Initiative: perspectives from the NIH. *Journal of the American Medical Women's Association* 1995; **50**(2):50-5.

The Women's Health Initiative Study Group. Design of the Women's Health Initiative clinical trial and observational study. *Controlled Clinical Trials* 1998; **19**(1):61-109.

Tinker LF, Bonds DE, Margolis KL, Manson JE, Howard BV, Larson J, et al. Low-fat dietary pattern and risk of treated diabetes mellitus in postmenopausal women: the Women's Health Initiative randomized controlled dietary modification trial. *Archives of Internal Medicine* 2008; **168**(14):1500-11.

Tinker LF, Perri MG, Patterson RE, Bowen DJ, McIntosh M, Parker LM, et al. The effects of physical and emotional status on adherence to a low-fat dietary pattern in the Women's Health Initiative. *Journal of the American Dietetic Association* 2002; **102**(6):789-800.

Tinker LF, Rosal MC, Young AF, Perri MG, Patterson RE, Van Horn L, et al. Predictors of dietary change and maintenance in the Women's Health Initiative Dietary Modification Trial. *Journal of the American Dietetic Association* 2007; **107**(7):1155-66.

Women's Health Initiative Study Group. Dietary adherence in the Women's Health Initiative Dietary Modification Trial. *Journal of the American Dietetic Association* 2004; **104**(4):654-8.

WINS 2006 {published and unpublished data}

Chlebowski RT, Blackburn GL, Buzzard IM, Rose DP, Martino S, Khandekar JD, et al. Adherence to a dietary fat intake reduction program in postmenopausal women receiving therapy for early breast cancer. The Women's Intervention Nutrition Study. *Journal of Clinical Oncology* 1993; **11**(11):2072-80.

* Chlebowski RT, Blackburn GL, Thomson CA, Nixon DW, Shapiro A, Hoy MK, et al. Dietary fat reduction and breast cancer outcome: interim efficacy results from the Women's Intervention Nutrition study. *Journal of the National Cancer Institute* 2006; **98**(24):1767-76.

Chlebowski RT, Blackburn GL. Final survival analysis from the randomized Womens Intervention Nutrition Study (WINS) evaluating dietary intervention as adjuvant breast cancer therapy. *Cancer Research* 2015; **75**(9 Supplement):S5-08.

Chlebowski RT, Rose D, Blackburn G, Buzzard M, Khandekar J, York R, et al. Feasibility of using dietary fat intake reduction in adjuvant breast cancer management. *Proceedings of American Society of Clinical Oncology* 1991; **86**:86.

Chlebowski RT, Rose DP, Buzzard IM, Blackburn GL, York M, , et al. Dietary fat reduction in adjuvant breast cancer therapy: current rationale and feasibility issues. *Adjuvant: the Cancer Journal* 1990; **6**:357-63.

Hoy MK, Winters BL, Chlebowski RT, Papoutsakis C, Shapiro A, Lubin MP, et al. Implementing a low-fat eating plan in the Women's Intervention Nutrition Study. *Journal of the American Dietetic Association* 2009; **109**(4):688-96.

Rose DP, Chlebowski RT, Connolly JM, Jones LA, Wynder EL. Effects of tamoxifen adjuvant therapy and a low-fat diet on serum binding proteins and estradiol bioavailability in postmenopausal breast cancer patients. *Cancer Research* 1992; **52**(19):5386-90.

Rose DP, Connolly JM, Chlebowski RT, Buzzard IM, Wynder EL. The effects of a low-fat dietary intervention and tamoxifen adjuvant therapy on the serum estrogen and sex hormonebinding globulin concentrations of postmenopausal breast cancer patients. *Breast Cancer Research & Treatment* 1993; **27**(3):253-62.

Wynder EL, Cohen LA, Winters BL. The challenges of assessing fat intake in cancer research investigations. *Journal of the American Dietetic Association* 1997; **97**(7 Suppl):S5-8.

References to studies excluded from this review

Agewall 2001 {published data only}

Agewall S. Multiple risk intervention trial in high risk hypertensive men: comparison of ultrasound intima-media thickness and clinical outcome during 6 years of follow-up. *Journal of Internal Medicine* 2001; **249**(4):305-14.

Ammerman 2003 {published data only}

Ammerman AS, Keyserling TC, Atwood JR, Hosking JD, Zayed H, Krasny C. A randomized controlled trial of a public health nurse directed treatment program for rural patients with high blood cholesterol. *Preventive Medicine* 2003; **36**(3):340-51.



Anderson 1990 {published and unpublished data}

* Anderson JW, Garrity TF, Smith BM, Whitis SE. Follow-up on a clinical trial comparing the effects of two lipid lowering diets. *Arteriosclerosis* 1990; **10**(5):882a.

Anderson JW, Garrity TF, Wood CL, Whitis SE, Smith BM,

Oeltgen PR. Prospective, randomized, controlled comparison of the effects of low-fat and low-fat plus high-fiber diets on serum lipid concentrations. *American Journal of Clinical Nutrition* 1992; **56**(5):887-94.

Aquilani 2000 {published data only}

Aquilani R, Tramarin R, Pedretti RF, Bertolotti G, Sommaruga M, Mariani P, et al. Can a very-low-fat diet achieve cholesterol goals in CAD? *Cardiology Review* 2000; **17**(10):36-40.

Arntzenius 1985 {published data only}

Arntzenius AC, Kromhout D, Barth JD, Reiber JH, Bruschke AV, Buis B, et al. Diet, lipoprotiens and progression of coronary atherosclerosis: the Leiden Intervention Trial. *New England Journal of Medicine* 1985; **312**(13):805-8.

Aro 1990 {published data only}

Aro A, Ahola I, Jauhiainen M, Mutanen M, Valsta LM. Effects of plasma phospholipid fatty acids of rapeseed oil and sunflower oil diets [Abstract]. *Arteriosclerosis* 1990; **10**:877a.

ASSIST 2001 {published data only}

Moher M, Yudkin P, Wright L, Turner R, Fuller A, Schofield T, et al. Cluster randomised controlled trial to compare three methods of promoting secondary prevention of coronary heart disease in primary care. *BMJ* 2001; **322**(7298):1338.

Australian Polyp Prev 95 {published and unpublished data}

* MacLennan R, Macrae F, Bain C, Battistutta D, Chapuis P, Gratten H, et al. Randomized trial of intake of fat, fiber, and beta carotene to prevent colorectal adenomas. The Australian Polyp Prevention Project. *Journal of the National Cancer Institute* 1995; **87**(23):1760-6.

MacLennan R. Effect of fat, fibre and beta-carotene on colorectal adenomas after 24 months. *Gastroenterology* 1991; **100**:A382.

Macrae FA, Hughes NR, Bhathal PS, Tay D, Selbie L, MacLennan R. Dietary suppression of rectal epthelial cell proliferation. *Gastroenterology* 1991; **100**:A383.

Azadbakht 2007 {published and unpublished data}

Azadbakht L, Mirmiran P, Esmaillzadeh A, Azizi F. Better dietary adherence and weight maintenance achieved by a longterm moderate fat diet. *British Journal of Nutrition* 2007; **97**(2):399-404.

Bakx 1997 {published data only}

Bakx JC, Stafleu A, Van Staveren WA, Van den Hoogen HJ, Van Weel C. Long-term effect of nutritional counseling: a study in family medicine. *American Journal of Clinical Nutrition* 1997; **65**(6 Suppl):1946S-50S.

Ball 1965 {published data only}

Ball KP, Hanington E, McAllen PM, Pilkington TR, Richards JM, Sharland DE, et al. Low-fat diet in myocardial infarction: a controlled trial. *Lancet* 1965; **2**(411):501-4.

Barnard 2009 {published data only}

Barnard ND, Cohen J, Jenkins DJ, Turner-McGrievy G, Gloede L, Green A, et al. A low-fat vegan diet and a conventional diabetes diet in the treatment of type 2 diabetes: a randomized, controlled, 74-wk clinical trial. *American Journal of Clinical Nutrition* 2009; **89**(5):1588S-96S.

Barndt 1977 {published data only}

Barndt R, Blankenhorn CH, Crawford DW, Brooks SH. Regression and progression of early femoral atherosclerosis in treated hyperlipidaemic patients. *Annals of Internal Medicine* 1977; **86**(2):139-46.

Baron 1990 {published data only}

Baron JA, Gleason R, Crowe B, Mann JI. Preliminary trial of the effect of general practice based nutritional advice. *British Journal of General Practice* 1990; **40**(333):137-41.

Barr 1990 {published data only}

Barr SL, Ramakrishnan R, Holleran S. A 30% fat diet high in polyunsaturates and a 30% fat diet high in monounsaturates both lower total and low density lipoprotein cholesterol levels in normal males [Abstract]. *Arteriosclerosis* 1990; **10**:872a.

Barsotti 1991 {published data only}

Barsotti A. Modern trends in the therapy of arteriosclerosis in the light of new physiopathological findings [Moderne tendenze della terapia dell'arteriosclerosi alla luce delle nuove acquisizioni fisiopatologiche]. *Cardiologia* 1991; **36**(12 Suppl 1):33-48.

Baumann 1982 {published data only}

Baumann J, Martschick R. Therapy of hyperlipidemia with xanthinol nicotinate as opposed to low fat diet [Therapie der hyperlipidämie mit xantinolnicotinat gegenüber fettarmer diät]. *Die Medizinische Welt* 1982; **33**(4):139-41.

BDIT Pilot Studies 1996 {published and unpublished data}

Boyd NF, Cousins M, Beaton M, Fishell E, Wright B, Fish E, et al. Clinical trial of low-fat, high-carbohydrate diet in subjects with mammographic dysplasia: report of early outcomes. *Journal of the National Cancer Institute* 1988; **80**(15):1244-8.

Boyd NF, Cousins M, Beaton M, Han L, McGuire V. Methodological issues in clinical trials of dietary fat reduction in patients with breast dysplasia. *Progress in Clinical and Biological Research* 1986; **222**:117-24.

Boyd NF, Cousins M, Beaton M, Kriukov V, Lockwood G, Tritchler D. Quantitative changes in dietary fat intake and serum cholesterol in women: results from a randomized, controlled trial. *American Journal of Clinical Nutrition* 1990; **52**(3):470-6.

Boyd NF, Cousins M, Kriukov V. A randomised controlled trial of dietary fat reduction: the retention of subjects and characteristics of drop outs. *Journal of Clinical Epidemiology* 1992; **45**(1):31-8.

Reduction in saturated fat intake for cardiovascular disease (Review)

Copyright $\ensuremath{\mathbb{G}}$ 2020 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.



Boyd NF, Cousins M, Lockwood G, Tritchler D. Dietary fat and breast cancer risk: the feasibility of a clinical trial of breast cancer prevention. *Lipids* 1992; **27**(10):821-6.

Boyd NF, Cousins M, Lockwood G, Tritchler D. The feasibility of testing experimentally the dietary fat-breast cancer hypothesis. *Progress in Clinical and Biological Research* 1990; **346**:231-41.

* Boyd NF, Martin LJ, Beaton M, Cousins M, Kriukov V. Longterm effects of participation in a randomized trial of a low-fat, high-carbohydrate diet. *Cancer Epidemiology, Biomarkers & Prevention* 1996; **5**(3):217-22.

Lee-Han H, Cousins M, Beaton M, McGuire V, Kriukov V, Chipman M, et al. Compliance in a randomized clinical trial of dietary fat reduction in patients with breast dysplasia. *American Journal of Clinical Nutrition* 1988; **48**(3):575-86.

Beckmann 1995 {published data only}

Beckmann SL, Os I, Kjeldsen SE, Eide IK, Westheim AS, Hjermann I. Effect of dietary counselling on blood pressure and arterial plasma catecholamines in primary hypertension. *American Journal of Hypertension* 1995; **8**(7):704-11.

beFIT 1997 {published and unpublished data}

Retzlaff BM, Walden CE, McNeney WB, Buck BL, McCann BS, Knopp RH. Nutritional intake of women and men on the NCEP Step I and Step II diets. *Journal of the American College of Nutrition* 1997; **16**(1):52-61.

* Walden CE, Retzlaff BM, Buck BL, McCann BS, Knopp RH. Lipoprotein lipid response to the National Cholesterol Education Program Step II diet by hypercholesterolemic and combined hyperlipidemic women and men. *Arteriosclerosis, Thrombosis and Vascular Biology* 1997; **17**(2):375-82.

Walden CE, Retzlaff BM, Buck BL, Wallick S, McCann BS, Knopp RH. Differential effect of National Cholesterol Education Program (NCEP) Step II Diet on HDL cholesterol, its subfractions, and apoprotein A-1 levels in hypercholesterolemic women and men after 1 year: the beFIT study. *Arteriosclerosis, Thrombosis and Vascular Biology* 2000; **20**(6):1580-7.

Beresford 1992 {published data only}

Beresford SA, Farmer EM, Feingold L, Graves KL, Sumner SK, Baker RM. Evaluation of a self-help dietary intervention in a primary care setting. *American Journal of Public Health* 1992; **82**(1):79-84.

Bergstrom 1967 {published data only}

Bergstrom G, Svanborg A. Dietary treatment of acute myocardial infarction. *Acta Medica Scandinavica* 1967; **181**(6):717-21.

Bierenbaum 1963 {published data only}

Bierenbaum ML, Fleischman AI, Raichelson RI, Hayton T, Watson P. Ten year experience of modified fat diets on younger men with coronary heart disease. *Lancet* 1973; **1**(7817):1404-7.

Bierenbaum ML, Green DP, Florin A, Fleischman AI, Caldwell AB. Modified-fat dietary management of the young male with coronary disease. A five-year report. *JAMA* 1967; **202**(13):1119-23. * Bierenbaum ML, Green DP, Gherman C, Florin A, Caldwell AB. The effects of two low fat dietary patterns on the blood cholesterol levels of young male coronary patients. *Journal of Chronic Diseases* 1963; **16**:1073-83.

Bloemberg 1991 {published and unpublished data}

Bloemberg BP, Kromhout D, Goddijn HE, Jansen A, Obermannde Boer GL. The impact for the guidelines for a healthy diet of the Netherlands Nutrition Council on total and high density lipoprotein cholesterol in hypercholesterolemic free living men. *American Journal of Epidemiology* 1991; **134**(1):39-48.

Bloomgarden 1987 {published data only}

Bloomgarden ZT, Karmally W, Metzger MJ, Brothers M, Nechemias C, Bookman J, et al. Randomized, controlled trial of diabetic patient education: improved knowledge without improved metabolic status. *Diabetes Care* 1987; **10**(3):263-72.

Bonk 1975 {published data only}

Bonk S, Hubotter E, Nickel C, Stocksmeier U, Vahey P, Volk I, et al. Myocardial infarct patients with and without intensive nutrition consultation over several years-- comparison of physiological and social variables [Herzinfarktpatienten mit und ohne mehrjährige intensive ernährungsberatung - vergleich von physiologischen und sozialen variablen]. *Infusionstherapie und Klinische Ernährung* 1975; **2**(4):290-6.

Bonnema 1995 {published data only}

Bonnema SJ, Jespersen LT, Marving J, Gregersen G. Supplementation with olive oil rather than fish oil increases small arterial compliance in diabetic patients. *Diabetes, Nutrition and Metabolism Clinical and Experimental* 1995; **8**:81-7.

Bosaeus 1992 {published data only}

Bosaeus I, Belfrage L, Lindgren C, Andersson H. Olive oil instead of butter increases net cholesterol excretion from the small bowel. *European Journal of Clinical Nutrition* 1992; **46**(2):111-5.

Boyd 1988 {published and unpublished data}

Boyd NF, McGuire V, Shannon P, Cousins M, Kriukov V, Mahoney L, et al. Effect of a low-fat high-carbohydrate diet on symptoms of cyclical mastopathy. *Lancet* 1988; **2**(8603):128-32.

BREACPNT {published data only}

. The breast cancer personalized nutrition study (BREACPNT). clinicaltrials.gov/ct2/show/NCT04079270 (first posted 5 September 2019).

Brehm 2009 {published data only (unpublished sought but not used)}

Brehm BJ, Lattin BL, Summer SS, Boback JA, Gilchrist GM, Jandacek RJ, et al. One-year comparison of a highmonounsaturated fat diet with a high-carbohydrate diet in type 2 diabetes. *Diabetes Care* 2009; **32**(2):215-20.

Brensike 1982 {published data only}

Brensike JF, Kelsey SF, Passamani ER, Fisher MR, Richardson JM, Loh IK, et al. National Heart, Lung, and Blood Institute type II Coronary Intervention Study: design, methods, and baseline characteristics. *Controlled Clinical Trials* 1982; **3**(2):91-111.



BRIDGES 2001 {published and unpublished data}

Hebert JR, Ebbeling CB, Olendzki BC, Hurley TG, Ma Y, Saal N, et al. Change in women's diet and body mass following intensive intervention for early-stage breast cancer. *Journal of the American Dietetic Association* 2001; **101**(4):421-31.

Broekmans 2003 {published and unpublished data}

Broekmans WM, Klöpping-Ketelaars IA, Weststrate JA, Tijburg LB, Van Poppel G, Vink AA, et al. Decreased carotenoid concentrations due to dietary sucrose polyesters do not affect possible markers of disease risk in humans. *Journal of Nutrition* 2003; **133**(3):720-6.

Brown 1984 {published data only}

Brown GD, Whyte L, Gee MI, Crockford PM, Grace M, Oberle K, et al. Effects of two "lipid-lowering" diets on plasma lipid levels of patients with peripheral vascular disease. *Journal of the American Dietetic Association* 1984; **84**(5):546-50.

Bruce 1994 {published data only}

Bruce SL, Grove SK. The effect of a coronary artery risk evaluation program on serum lipid values and cardiovascular risk levels. *Applied Nursing Research* 1994; **7**(2):67-74.

Bruno 1983 {published data only}

Bruno R, Arnold C, Jacobson L, Winick M, Wynder E. Randomized controlled trial of a nonpharmacologic cholesterol reduction program at the worksite. *Preventive Medicine* 1983; **12**(4):523-32.

Butcher 1990 {published data only}

Butcher LA, O'Dea K, Sinclair AJ, Parkin JD, Smith IL, Blombery P. The effects of very low fat diets enriched with fish or kangaroo meat on cold-induced vasoconstriction and platelet function. *Prostaglandins, Leukotrienes, and Essential Fatty Acids* 1990; **39**(3):221-6.

Byers 1995 {published data only}

Byers T, Mullis R, Anderson J, Dusenbury L, Gorsky R, Kimber C, et al. The costs and effects of a nutritional education program following work-site cholesterol screening. *American Journal of Public Health* 1995; **85**(5):650-5.

Caggiula 1996 {published data only}

Caggiula AW, Watson JE, Kuller LH, Olson MB, Milas NC, Berry M, et al. Cholesterol-lowering intervention program. Effect of the step I diet in community office practices. *Archives of Internal Medicine* 1996; **156**(11):1205-13.

Canadian DBCP 1997 {published data only (unpublished sought but not used)}

Boyd NF, Greenberg C, Lockwood G, Little L, Martin L, Byng J, et al. Effects at two years of a low-fat, high-carbohydrate diet on radiologic features of the breast: results from a randomized trial. *Journal of the Nationl Cancer Institute* 1997; **89**(7):488-96.

CARMEN 2000 {published and unpublished data}

Poppitt SD, Keogh GF, Prentice AM, Williams DE, Sonnemans HM, Valk EE, et al. Long-term effects of ad libitum low-fat, high-carbohydrate diets on body weight and serum lipids in overweight subjects with metabolic syndrome. *American Journal of Clinical Nutrition* 2002; **75**(1):11-20.

Raben A, Astrup A, Vasilaras TH, Prentice AM, , Formiguera X, et al. The CARMEN study [CARMEN-studiet]. *Ugeskrift für Laeger* 2002; **164**(5):627-31.

* Saris WH, Astrup A, Prentice AM, Zunft HJ, Formiguera X, Verboeket-van de Venne WP, et al. Randomized controlled trial of changes in dietary carbohydrate/fat ratio and simple vs complex carbohydrates on body weight and blood lipids: the CARMEN study. *International Journal of Obesity* 2000; **24**(10):1310-8.

Saris WH, Astrup A, Prentice AM, Zunft HJ, Formiguera X. CARMEN Project: European multicentre study on the impact of dietary fat/CHO ratio and simple/complex CHO changes on long term weight control in overweight subjects. *International Journal of Obesity* 1997; **21**(Suppl 2):S71.

Vasilaras TH, Astrup A, Raben A. Micronutrient intake in overweight subjects is not deficient on an ad libitum fatreduced, high-simple carbohydrate diet. *European Journal of Clinical Nutrition* 2004; **58**(2):326-36.

CARMEN substudy 2002 {published and unpublished data}

Poppitt SD, Keogh GF, Prentice AM, Williams DEM, Sonnemans HMW, Valk EEJ, et al. Long-term effects of ad libitum low-fat, high-carbohydrate diets on body weight and serum lipids in overweight subjects with metabolic syndrome. *American Journal of Clinical Nutrition* 2002; **75**(1):11-20.

Casas-Agustench 2013 {published data only}

Casas-Agustench P, Molina S, Espinosa-Salinas I, Olivares M, Reglero G, Ordovás JM, et al. SELP variant modulates plasma HDL-C responses in subjects with moderate cardiovascular risk after skimmed milk consumption. *FASEB Journal* 2013; **27**(1 suppl):640.21.

Loria-Kohen V, Espinosa-Salinas I, Ramirez de Molina A, Casas-Agustench P, Herranz J, Molina S, et al. A genetic variant of PPARA modulates cardiovascular risk biomarkers after milk consumption. *Nutrition* 2014; **30**(10):1144-50.

Cerin 1993 {*published data only*}

Cerin A, Collins A, Landgren BM, Eneroth P. Hormonal and biochemical profiles of premenstrual syndrome. Treatment with essential fatty acids. *Acta Obstetricia et Gynecologica Scandinavica* 1993; **72**(5):337-43.

Chan 1993 {published data only}

Chan JK, McDonald BE, Gerrard JM, Bruce VM, Weaver BJ, Holub BJ. Effect of dietary alpha-linolenic acid and its ratio to linolenic acid on platelet and plasma fatty acids and thrombogenesis. *Lipids* 1993; **28**(9):811-7.

Chapman 1950 {published data only}

Chapman CB, Gibbons T, Henschel A. The effect of the rice-fruit diet on the composition of the body. *New England Journal of Medicine* 1950; **243**(23):899-905.



Charbonnier 1975 {published data only}

Charbonnier A, Nepveux P, Fluteau G, Fluteau D. Immediate effects of ingestion of olive oil on the principal lipid constituents of the plasma. Comparison with other edible fats. *Médecine & Chirurgie Digestives* 1975; **4 Suppl 2**:73-9.

Cheng 2004 {published data only}

Cheng C, Graziani C, Diamond JJ. Cholesterol-lowering effect of the Food for Heart Nutrition Education Program. *Journal of the American Dietetic Association* 2004; **104**(12):1868-72.

Chiostri 1988 {published data only}

Chiostri JE, Kwiterovich PO. Effect of American Heart Association Phase 2 diet versus eater's choice based diet on hypercholesterolaemia. *Circulation* 1988; **78**(4):II-385.

Choudhury 1984 {published data only}

Choudhury S, Jackson P, Katan MB, Marenah CB, Cortese C, Miller NE, et al. A multifactorial diet in the management of hyperlipidaemia. *Atherosclerosis* 1984; **50**(1):93-103.

Clark 1997 {published data only}

Clark M, Ghandour G, Miller NH, Taylor CB, Bandura A, DeBusk RF. Development and evaluation of a computer-based system for dietary management of hyperlipidemia. *Journal of the American Dietetic Association* 1997; **97**(2):146-50.

Clifton 1992 {published data only}

Clifton PM, Wight MB, Nestel PJ. Is fat restriction needed with HMGCoA reductase inhibitor treatment? *Atherosclerosis* 1992; **93**(1-2):59-70.

Cobb 1991 {published data only}

Cobb MM, Teitelbaum HS, Breslow JL. Lovastatin efficacy in reducing low-density lipoprotein cholesterol levels on high-vs low-fat diets. *JAMA* 1991; **265**(8):997-1001.

Cohen 1991 {published data only}

Cohen MD, D'Amico FJ, Merenstein JH. Weight reduction in obese hypertensive patients. *Family Medicine* 1991; **23**(1):25-8.

Cole 1988 {published data only}

Cole TG, Schmeisser D, Prewitt TE. AHA phase 3 diet reduces cholesterol in moderately hypercholesterolemic premenopausal women [Abstract]. *Circulation* 1988; **78**(4):II-73.

Colquhoun 1990 {published data only}

Colquhoun DM, Moores D, Somerset SM. Comparison of the effects of an avocado enriched and American Heart Association diets on lipid levels [Abstract]. *Arteriosclerosis* 1990; **10**:875a.

Consolazio 1946 {published data only}

Consolazio FC, Forbes WH. The effects of high fat diet in a temperate environment. *Journal of Nutrition* 1946; **32**:195-204.

Cox 1996 {published data only}

Cox RH, Gonzales-Vigilar MC, Novascone MA, Silva-Barbeau I. Impact of a cancer intervention on diet-related cardiovascular disease risks of white and African-American EFNEP clients. *Journal of Nutrition Education* 1996; **28**:209-18.

Croft 1986 {published data only}

Croft PR, Brigg D, Smith S, Harrison CB, Branthwaite A, Collins MF. How useful is weight reduction in the management of hypertension? *Journal of the Royal College of General Practitioners* 1986; **36**(291):445-8.

Curzio 1989 {published and unpublished data}

Curzio JL, Kennedy SS, Elliott HL, Farish E, Barnes JF, Howie CA, et al. Hypercholesterolaemia in treated hypertensives: a controlled trial of intensive dietary advice. *Journal of Hypertension* 1989; **7**(6 Suppl):S254-5.

Dalgard 2001 {published data only}

Dalgard C, Thuroe A, Haastrup B, Haghfelt T, Stender S. Saturated fat intake is reduced in patients with ischemic heart disease 1 year after comprehensive counseling but not after brief counseling. *Journal of the American Dietetic Association* 2001; **101**(12):1420-9.

Da Qing IGT 1997 {published data only}

Pan XR, Li GW, Hu YH, Wang JX, Yang WY, An ZX, et al. Effects of diet and exercise in preventing NIDDM in people with impaired glucose tolerance. The Da Qing IGT and Diabetes Study. *Diabetes Care* 1997; **20**(4):537-44.

DAS 2000 {published data only}

Bovbjerg VE, McCann BS, Brief DJ, Follette WC, Retzlaff BM, Dowdy AA, et al. Spouse support and long-term adherence to lipid-lowering diets. *American Journal of Epidemiology* 1995; **141**(5):451-60.

* Knopp RH, Retzlaff B, Walden C, Fish B, Buck B, McCann B. One-year effects of increasingly fat-restricted, carbohydrateenriched diets on lipoprotein levels in free-living subjects. *Proceedings of the Society for Experimental Biology & Medicine* 2000; **225**(3):191-9.

Knopp RH, Walden CE, McCann BS, Retzlaff B, Dowdy A, Gey G, et al. Serial changes in lipoprotein cholesterol in hypercholesterolemic men treated with alternative diets [abstract]. *Arteriosclerosis* 1989; **9**:745A.

Knopp RH, Walden CE, Retzlaff BM, McCann BS, Dowdy AA, Albers JJ, et al. Long-term cholesterol-lowering effects of 4 fat-restricted diets in hypercholesterolaemic and combined hyperlipidaemic men: the Dietary Alternatives Study. *JAMA* 1997; **278**(18):1509-15.

Walden CE, McCann BS, Retzlaff B, Dowdy A, Hanson M, Fish B, et al. Alternative fat-restricted diets for hypercholesterolemia and combined hyperlipidemia: feasibility, design, subject recruitment, and baseline characteristics of the Dietary Alternatives study. *Journal of the American College of Nutrition* 1991; **10**(5):429-42.

DASH 1997 {published data only}

* Appel LJ, Moore TJ, Obarzanek E, Vollmer WM, Svetkey LP, Sacks FM, et al, DASH Collaborative Research Group. A clinical trial of the effects of dietary patterns on blood pressure. *New England Journal of Medicine* 1997; **336**(16):1117-24.

Blackburn GL. Functional foods in the prevention and treatment of disease: significance of the Dietary Approaches to Stop Hypertension Study. *American Journal of Clinical Nutrition* 1997; **66**(5):1067-71.

Davey Smith 2005 {published data only}

Davey Smith G, Bracha Y, Svendsen KH, Neaton JD, Haffner SM, Kuller LH, et al. Incidence of type 2 diabetes in the randomized multiple risk factor intervention trial. *Annals of Internal Medicine* 2005; **142**(5):313-22.

De Boer 1983 {published data only}

De Boer AC, Turek JV, Pannebakker MA, Den Ottolander GJ. The effect of diets high in polyunsaturated and high in saturated fatty acids on blood lipids and platelet tests in patients with coronary artery disease (CAD) [abstract]. *Thrombosis and Haemostasis* 1983; **50**:96.

De Bont 1981 {published and unpublished data}

De Bont AJ, Baker IA, St Leger AS, Sweetnam PM, Wragg KG, Stephens SM, et al. A randomised controlled trial of the effect of low fat diet advice on dietary response in insulin independent diabetic women. *Diabetologia* 1981; **21**(6):529-33.

DeBusk 1994 {published data only}

DeBusk RF, Miller NH, Superko HR, Dennis CA, Thomas RJ, Lew HT, et al. A case-management system for coronary risk factor modification after acute myocardial infarction [see comments]. *Annals of Internal Medicine* 1994; **120**(9):721-9.

DEER 1998 {published data only (unpublished sought but not used)}

Stefanick ML, Mackey S, Sheehan RD, Ellsworth N, Haskell WL, Wood PD. Effects of diet and exercise in men and postmenopausal women with low levels of HDL cholesterol and high levels of LDL cholesterol. *New England Journal of Medicine* 1998; **339**(1):12-20.

Delahanty 2001 {published data only}

Delahanty LM, Hayden D, Ammerman A, Nathan DM. Medical nutrition therapy for hypercholesterolemia positively affects patient satisfaction and quality of life outcomes. *Annals of Behavioral Medicine* 2002; **24**(4):269-78.

* Delahanty LM, Sonnenberg LM, Hayden D, Nathan DM. Clinical and cost outcomes of medical nutrition therapy for hypercholesterolemia: a controlled trial. *Journal of the American Dietetic Association* 2001; **101**(9):1012-23.

Delius 1969 {*published data only*}

Delius L. Treatment of hypotensive circulatory disorder [Die behandlung der hypotonen kreislaufregulationsstörung]. *Deutsche Medizinische Wochenschrift* 1969; **94**(42):2172-3.

Demark 1990 {published data only}

Demark WW, Bowering J, Cohen PS. Reduced serum cholesterol with dietary change using fat-modified and oat bran supplemented diets. *Journal of the American Dietetic Association* 1990; **90**(2):223-9.

Dengel 1995 {published data only}

Dengel JL, Katzel LI, Goldberg AP. Effect of an American Heart Association diet, with or without weight loss, on lipids in obese middle-aged and older men. *American Journal of Clinical Nutrition* 1995; **62**(4):715-21.

Denke 1994 {published data only}

Denke MA, Grundy SM. Individual responses to a cholesterol lowering diet in 50 men with moderate hypercholesterolaemia. *Archives of Internal Medicine* 1994; **154**(3):17-25.

Diabetes CCT 1995 {published data only}

The Diabetes Control and Complications Trial (DCCT) Research Group. Effect of intensive diabetes management on macrovascular events and risk factors in the Diabetes Control and Complications Trial. *American Journal of Cardiology* 1995; **75**(14):894-903.

Diet & Hormone Study 2003 {published data only (unpublished sought but not used)}

Gann PH, Chatterton RT, Gapstur SM, Liu K, Garside D, Giovanazzi S, et al. The effects of a low-fat/high-fiber diet on sex hormone levels and menstrual cycling in premenopausal women: a 12-month randomized trial (the Diet and Hormone Study). *Cancer* 2003; **98**(9):1870-9.

DIET 1998 {published data only}

Dornelas EA, Wylie-Rosett J, Swencionis C. The DIET study: long term outcomes of a cognitive-behavioural weight control intervention in independent-living elders. *Journal of the American Dietetic Association* 1998; **98**(11):1276-81.

Ding 1992 {published data only}

Ding Q. Clinical study of qianxining in the treatment of 60 cases of yang hyperactivity due to yin deficiency type of hypertension. *Chung Kuo Chung Hsi I Chieh Ho Tsa Chih* 1992; **12**(7):409-11, 388-9.

DIPI 2018 {published data only}

ZZZ <accessionId ref="info:x-wiley/clinicalTrialsGov/ NCT02062424"> ZZZNCT02062424

Arentoft JL, Hoppe C, Andersen EW, Overvad K, Tetens I. Associations between adherence to the Danish Food-Based Dietary Guidelines and cardiometabolic risk factors in a Danish adult population: the DIPI study. *British Journal of Nutrition* 2018; **119**(6):664-73.

DIRECT 2009 {published data only (unpublished sought but not used)}

* Ben-Avraham S, Harman-Boehm I, Schwarzfuchs D, Shai I. Dietary strategies for patients with type 2 diabetes in the era of multi-approaches; review and results from the Dietary Intervention Randomized Controlled Trial (DIRECT). *Diabetes Research and Clinical Practice* 2009; **86**(Suppl 1):S41-8.

Resch KL. Dietary Intervention Randomized Controlled Trial (DIRECT) group: weight loss with a lowcarbohydrate, Mediterranean, or low-fat diet. *Forschende Komplementarmedizin* (2006) 2008; **15**(6):351-2.

Shai I, Gepner Y, Shelef I, Schwarzfuchs D, Zelicha H, Tene L, et al. Precision nutrition. *Obesity Facts* 2019; **12**:1.

Shai I, Schwarzfuchs D, Henkin Y, Shahar DR, Witkow S, Greenberg I, et al. Weight loss with a low-carbohydrate, Mediterranean, or low-fat diet. *New England Journal of Medicine* 2008; **359**(3):229-41.

Shai I, Spence JD, Schwarzfuchs D, Henkin Y, Parraga G, Rudich A, et al. Dietary intervention to reverse carotid atherosclerosis. *Circulation* 2010; **121**(10):1200-8.

Shai I. The effect of low-carb, Mediterranean and low-fat diets on renal function; a 2-year dietary intervention randomized controlled trial (DIRECT). *Obesity Facts* 2012; **5**:19.

Dobs 1991 {published data only}

Dobs AS, Sarma PS, Wilder L. Lipid-lowering diets in patients taking pravastatin, a new HMG-CoA reductase inhibitor: compliance and adequacy. *American Journal of Clinical Nutrition* 1991; **54**(4):696-700.

DO IT 2006 {published and unpublished data}

Berstad P, Seljeflot I, Veierod MB, Hjerkinn EM, Arnesen H, Pedersen JI, et al. Supplementation with fish oil affects the association between very long-chain n-3 polyunsaturated fatty acids in serum non-esterified fatty acids and soluble vascular cell adhesion molecule-1. *Clinical Science* 2003; **105**(1):13-20.

Ellingsen I, Hjerkinn EM, Seljeflot I, Arnesen H, Tonstad S, Ellingsen I, et al. Consumption of fruit and berries is inversely associated with carotid atherosclerosis in elderly men. *British Journal of Nutrition* 2008; **99**(3):674-81 Erratum in: British Journal of Nutrition 2008; 99(3): 697.

Ellingsen I, Seljeflot I, Arnesen H, Tonstad S, , , et al. Vitamin C consumption is associated with less progression in carotid intima media thickness in elderly men: a 3-year intervention study. *Nutrition Metabolism & Cardiovascular Diseases* 2009; **19**(1):8-14.

Furenes EB, Seljeflot I, Solheim S, Hjerkinn EM, Arnesen H. Long-term influence of diet and/or omega-3 fatty acids on matrix metalloproteinase-9 and pregnancy-associated plasma protein-A in men at high risk of coronary heart disease. *Scandinavian Journal of Clinical & Laboratory Investigation* 2008; **68**(3):177-84.

* Hjerkinn EM, Abdelnoor M, Breivik L, Bergengen L, Ellingsen I, Seljeflot I, et al. Effect of diet or very long chain omega-3 fatty acids on progression of atherosclerosis, evaluated by carotid plaques, intima-media thickness and by pulse wave propagation in elderly men with hypercholesterolaemia. *European Journal of Cardiovascular Prevention & Rehabilitation* 2006; **13**(3):325-33.

Hjerkinn EM, Seljeflot I, Ellingsen I, Berstad P, Hjermann I, Sandvik L, et al. Influence of long-term intervention with dietary counselling, long-chain n-3 fatty acid supplements, or both on circulating markers of endothelial activation in men with longstanding hyperlipidemia. *American Journal of Clinical Nutrition* 2005; **81**(3):583-9. Lindman AS, Pedersen JI, Hjerkinn EM, Arnesen H, Veierod MB, Ellingsen I, et al. The effects of long-term diet and omega-3 fatty acid supplementation on coagulation factor VII and serum phospholipids with special emphasis on the R353Q polymorphism of the FVII gene. *Thrombosis & Haemostasis* 2004; **91**(6):1097-104.

Troseid M, Arnesen H, Hjerkinn EM, Seljeflot I. Serum levels of interleukin-18 are reduced by diet and n-3 fatty acid intervention in elderly high-risk men. *Metabolism: Clinical & Experimental* 2009; **58**(11):1543-9.

Troseid M, Seljeflot I, Hjerkinn EM, Arnesen H. Interleukin-18 is a strong predictor of cardiovascular events in elderly men with the metabolic syndrome: synergistic effect of inflammation and hyperglycemia. *Diabetes Care* 2009; **32**(3):486-92.

Due 2008 {published and unpublished data}

Bladbjerg EM, Larsen TM, Due A, Jespersen J, Stender S, Astrup A. Postprandial coagulation activation in overweight individuals after weight loss: acute and long-term effects of a high-monounsaturated fat diet and a low-fat diet. *Thrombosis Research* 2014; **133**(3):327-33.

Due A, Larsen TM, Hermansen K, Stender S, Holst JJ, Toubro S, et al. Comparison of the effects on insulin resistance and glucose tolerance of 6-mo high-monounsaturated-fat, low-fat, and control diets. *American Journal of Clinical Nutrition* 2008; **87**(4):855-62.

* Due A, Larsen TM, Mu H, Hermansen K, Stender S, Astrup A. Comparison of 3 ad libitum diets for weight-loss maintenance, risk of cardiovascular disease, and diabetes: a 6-mo randomized, controlled trial. *American Journal of Clinical Nutrition* 2008; **88**(5):1232-41.

Rasmussen LG, Larsen TM, Mortensen PK, Due A, Astrup A. Effect on 24-h energy expenditure of a moderate-fat diet high in monounsaturated fatty acids compared with that of a low-fat, carbohydrate-rich diet: a 6-mo controlled dietary intervention trial. *American Journal of Clinical Nutrition* 2007; **85**(4):1014-22.

Sloth B, Due A, Larsen TM, Holst JJ, Heding A, Astrup A, et al. The effect of a high-MUFA, low-glycaemic index diet and a lowfat diet on appetite and glucose metabolism during a 6-month weight maintenance period. *British Journal of Nutrition* 2009; **101**(12):1846-58.

Duffield 1982 {published data only}

* Duffield RG, Lewis B, Miller NE, Jamieson CW, Brunt JN, Colchester AC. Treatment of hyperlipidaemia retards progression of symptomatic femoral atherosclerosis. A randomised controlled trial. *Lancet* 1983; **2**(8351):639-42.

Duffield RG, Miller NE, Jamieson CW, Lewis B. A controlled trial of plasma lipid reduction in peripheral atherosclerosis - an interim report. *British Journal of Surgery* 1982; **69**(Suppl):S3-S5.

Dullaart 1992 {published and unpublished data}

Dullaart RP, Beusekamp BJ, Meijer S, Hoogenberg K, Van Doormaal JJ, Sluiter WJ. Long-term effects of linoleic-acidenriched diet on albuminuria and lipid levels in type 1 (insulin-

dependent) diabetic patients with elevated urinary albumin excretion. *Diabetologia* 1992; **35**(2):165-72.

Eating Patterns 1997 {published and unpublished data}

Beresford SA, Curry SJ, Kristal AR, Lazovich D, Feng Z, Wagner EH. A dietary intervention in primary care practice: the Eating Patterns Study. *American Journal of Public Health* 1997; **87**(4):610-6.

Ehnholm 1982 {published data only}

Ehnholm C, Huttunen JK, Pietinen P, Leino U, Mutanen M, Kostiainen E, et al. Effect of diet on serum lipoproteins in a population with a high risk of coronary heart disease. *New England Journal of Medicine* 1982; **307**(14):850-5.

Ehnholm 1984 {published data only}

Ehnholm C, Huttunen JK, Pietinen P, Leino U, Mutanen M, Kostiainen E, et al. Effect of a diet low in saturated fatty acids on plasma lipids, lipoproteins, and HDL subfractions. *Arteriosclerosis* 1984; **4**(3):265-9.

Eisenberg 1990 {published data only}

Eisenberg S. The effect of dietary substitution of monounsaturated fatty acids with carbohydrates on lipoprotein levels, structure, and function in a free-living population [abstract]. *Arteriosclerosis* 1990; **10**:872A.

Elder 2000 {published data only}

Elder JP, Candelaria JI, Woodruff SI, Criqui MH, Talavera GA, Rupp JW. Results of language for health: cardiovascular disease nutrition education for Latino English-as-a-second-language students. *Health Education & Behavior* 2000; **27**(1):50-63.

Ellegard 1991 {published data only}

Ellegard L, Bosaeus I. Sterol and nutrient excretion in ileostomists on prudent diets. *European Journal of Clinical Nutrition* 1991; **45**(9):451-7.

Esposito 2003 {published data only}

Esposito K, Pontillo A, Di Palo C, Giugliano G, Masella M, Marfella R, et al. Effect of weight loss and lifestyle changes on vascular inflammatory markers in obese women: a randomized trial. *JAMA* 2003; **289**(14):1799-804.

Esposito 2004 {published data only}

Esposito K, Marfella R, Ciotola M, Di Palo C, Giugliano F, Giugliano G, et al. Effect of a Mediterranean-style diet on endothelial dysfunction and markers of vascular inflammation in the metabolic syndrome: a randomized trial. *JAMA* 2004; **292**(12):1440-6.

EUROACTION 2008 {published data only}

Wood DA, Kotseva K, Connolly S, Jennings C, Mead A, Jones J, et al. Nurse-coordinated multidisciplinary, familybased cardiovascular disease prevention programme (EUROACTION) for patients with coronary heart disease and asymptomatic individuals at high risk of cardiovascular disease: a paired, cluster-randomised controlled trial. *Lancet* 2008; **371**(9629):1999-2012.

FARIS 1997 {published data only}

Goble A, Jackson B, Phillips P, Race E, Oliver RG, Worcester MC. The Family Atherosclerosis Risk Intervention Study (FARIS): risk factor profiles of patients and their relatives following an acute cardiac event. *Australian and New Zealand Journal of Medicine* 1997; **27**(5):568-77.

Fasting HGS 1997 {published data only}

Dyson PA, Hammersley MS, Morris RJ, Holman RR, Turner RC. The Fasting Hyperglycaemia Study: II. Randomized controlled trial of reinforced healthy-living advice in subjects with increased but not diabetic fasting plasma glucose. *Metabolism* 1997; **46**(12 Suppl 1):50-5.

Ferrara 2000 {published data only}

Ferrara LA, Raimondi AS, D'Episcopo L, Guida L, Dello Russo A, Marotta T. Olive oil and reduced need for antihypertensive medications. *Archives of Internal Medicine* 2000; **160**(6):837-42.

Fielding 1995 {published data only}

Fielding CJ, Havel RJ, Todd KM, Yeo KE, Schloetter MC, Weinberg V, et al. Effects of dietary cholesterol and fat saturation on plasma lipoproteins in an ethnically diverse population of healthy young men. *Journal of Clinical Investigation* 1995; **95**(2):611-8.

Finnish Diabet Prev 2000 {published data only}

Uusitupa M, Louheranta A, Lindstrom J, Valle T, Sundvall J, Eriksson J, et al. The Finnish Diabetes Prevention Study. *British Journal of Nutrition* 2000; **83**(Suppl 1):S137-42.

Finnish Mental Hosp 1972 {published data only}

* Miettinen M, Turpeinen O, Karvonen MJ, Elosuo R, Paavilainen E. Effect of cholesterol-lowering diet on mortality from coronary heart-disease and other causes: a twelve-year clinical trial in men and women. *Lancet* 1972; **2**(782):835-8.

Miettinen M, Turpeinen O, Karvonen MJ, Pekkarinen M, Paavilainen E, Elosuo R. Dietary prevention of coronary heart disease in women: the Finnish Mental Hospital study. *International Journal of Epidemiology* 1983; **12**(1):17-25.

Turpeinen O, Miettinen M, Karvonen M, Roine P, Pekkarinen M, Lehtosuo EJ, et al. Dietary prevention of coronary heart disease: long-term experiment. I. Observations on male subjects. *American Journal of Clinical Nutrition* 1968; **21**(4):255-76.

Fisher 1981 {published data only}

Fisher EA, Breslow JL, Zannis VI, Shen G, Blum CB. Dietary saturated fat, not cholesterol, affects plasma lipids and Apo E. *Arteriosclerosis* 1981; **1**(5):364a.

FIT Heart 2011 {published data only}

Mochari-Greenberger H, Terry MB, Mosca L. Does stage of change modify the effectiveness of an educational intervention to improve diet among family members of hospitalized cardiovascular disease patients? *Journal of the American Dietetic Association* 2010; **110**(7):1027-35.

* Mochari-Greenberger H, Terry MB, Mosca L. Gender, age and race/ethnicity do not modify the effectiveness of a diet intervention among family members of hospitalized

cardiovascular disease patients. *Journal of Nutrition Education and Behavior* 2011; **43**(5):366-73.

Fleming 2002 {published data only}

Fleming RM. The effect of high-, moderate-, and low-fat diets on weight loss and cardiovascular disease risk factors. *Preventive Cardiology* 2002; **5**(3):110-5.

Fortmann 1988 {published data only}

Fortmann SP, Haskell WL, Wood PD. Effects of weight loss on clinic and ambulatory blood pressure in normotensive men. *American Journal of Cardiology* 1988; **62**(1):89-93.

Foster 2003 {published data only}

Foster GD, Wyatt HR, Hill JO, McGuckin BG, Brill C, Mohammed BS, et al. A randomized trial of a low-carbohydrate diet for obesity. *New England Journal of Medicine* 2003; **348**(21):2082-90.

Frenkiel 1986 {published data only}

Frenkiel PG, Lee DW, Cohen H, Gilmore CJ, Resser K, Bonorris GG, et al. The effect of diet on bile acid kinetics and biliary lipid secretion in gallstone patients treated with ursodeoxycholic acid. *American Journal of Clinical Nutrition* 1986; **43**(2):239-50.

FRESH START 2007 {published data only}

Denmark-Wahnefried W, Clipp EC, Lipkus IM, Lobach D, Snyder DC, Sloane R, et al. Main outcomes of the FRESH START trial: a sequentially tailored, diet and exercise mailed print intervention among breast and prostate cancer survivors. *Journal of Clinical Oncology* 2007; **25**(19):2709-18.

Gambera 1995 {published data only}

Gambera PJ, Schneeman BO, Davis PA. Use of the Food Guide Pyramid and US Dietary Guidelines to improve dietary intake and reduce cardiovascular risk in active-duty Air Force members. *Journal of the American Dietetic Association* 1995; **95**(11):1268-73.

Gaullier 2007 {published data only}

, Halse J, Hoivik HO, Hoye K, Syvertsen C, Nurminiemi M, et al. Six months supplementation with conjugated linoleic acid induces regional-specific fat mass decreases in overweight and obese. *British Journal of Nutrition* 2007; **97**:550-60.

Ginsberg 1988 {published data only}

Ginsberg H. Both a high monounsaturated fat diet and the step 1 AHA diet significantly reduce plasma cholesterol levels in healthy males [abstract]. *Circulation* 1988; **78**:II73.

Gjone 1972 {published data only}

Gjone E, Nordoy A, Blomhoff JP, Wiencke I. The effects of unsaturated and saturated dietary fats on plasma cholesterol, phospholipids and lecithin: cholesterol acyltransferase activity. *Acta Medica Scandinavica* 1972; **191**(6):481-4.

Glatzel 1966 {published data only}

Glatzel H. The relationship between postprandial triglyceridemia and the fat content of the basic diet [Die

Cochrane Database of Systematic Reviews

Abhängigkeit der postcenalen triglyceridamie vom fettgehalt der grundkost]. *Klinische Wochenschrift* 1966; **44**(5):283-4.

Goodpaster 1999 {published data only}

Goodpaster BH, Kelley DE, Wing RR, Meier A, Thaete FL. Effects of weight loss on regional fat distribution and insulin sensitivity in obesity. *Diabetes* 1999; **48**(4):839-47.

Grundy 1986 {published data only}

Grundy SM, Nix D, Whelan MF, Franklin L. Comparison of three cholesterol-lowering diets in normolipidaemic men. *JAMA* 1986; **256**(17):2351-5.

Hardcastle 2008 {published data only}

Hardcastle S, Taylor A, Bailey M, Castle R. A randomised controlled trial on the effectiveness of a primary health care based counselling intervention on physical activity, diet and CHD risk factors. *Patient Education & Counseling* 2008; **70**(1):31-9.

Harris 1990 {published data only}

Harris WS, Feldman EB. Intensive dietary intervention in hypercholesterolemic patients. Observed versus predicted changes in cholesterol levels [abstract]. *Arteriosclerosis* 1990; **10**:853A.

Hartman 1993 {published data only}

Hartman T, McCarthy P, Himes J. Use of eating pattern messages to evaluate changes in eating behaviors in a worksite cholesterol education program. *Journal of the American Dietetic Association* 1993; **93**(10):1119-23.

Hartwell 1986 {published data only}

Hartwell SL, Kaplan RM, Wallace JP. Comparison of behavioral interventions for control of type II diabetes mellitus. *Behavior Therapy* 1986; **17**:447-61.

Hashim 1960 {published data only}

Hashim SA, Arteaga A, Van Itallie TB. Effect of saturated medium-chain triglyceride on serum-lipids in man. *Lancet* 1960; **1**(7134):1105-7.

Haufe 2011 {published and unpublished data}

* Haufe S, Engeli S, Kast P, Böhnke J, Utz W, Haas V, et al. Randomized comparison of reduced fat and reduced carbohydrate hypocaloric diets on intrahepatic fat in overweight and obese human subjects. *Hepatology* 2011; **53**(5):1504-14.

Haufe S, Utz W, Engeli S, Kast P, Böhnke J, Pofahl M, et al. Left ventricular mass and function with reduced-fat or reduced-carbohydrate hypocaloric diets in overweight and obese subjects. *Hypertension* 2012; **59**(1):70-5.

Haynes 1984 {published data only}

Haynes RB, Harper AC, Costley SR, Johnston M, Logan AG, Flanagan PT, et al. Failure of weight reduction to reduce mildly elevated blood pressure: a randomized trial. *Journal of Hypertension* 1984; **2**(5):535-9.



Heber 1991 {published data only}

Heber D, Ashley JM, Leaf DA, Barnard JA. Reduction of serum estradiol in postmenopausal women given free access to low-fat high carbohydrate diet. *Nutrition* 1991; **7**(2):137-41.

Heine 1989 {published and unpublished data}

Heine RJ, Mulder C, Popp-Snijders C, Van der Meer J, Van der Veen EA. Linoleic-acid-enriched diet: long-term effects on serum lipoprotein and apolipoprotein concentration and insulin sensitivity in noninsulin-dependent diabetic patients. *American Journal of Clinical Nutrition* 1989; **49**(3):448-56.

Hellenius 1995 {published and unpublished data}

Hellenius ML, Brismar KE, Berglund BH, De Faire U. Effects on glucose tolerance, insulin secretion, insulin-like growth factor 1 and its binding protein, IGFBP-1, in a randomized controlled diet and exercise study in healthy, middle-aged men. *Journal of Internal Medicine* 1995; **238**(2):121-30.

Hellenius ML, Dahlof C, Aberg H, Krakau I, De Faire U. Quality of life is not negatively affected by diet and exercise intervention in healthy men with cardiovascular risk factors. *Quality of Life Research* 1995; **4**(1):13-20.

Hellenius ML, De Faire U, Berglund B, Hamsten A, Krakau I. Diet and exercise are equally effective in reducing risk for cardiovascular disease. Results of a randomized controlled study in men with slightly to moderately raised cardiovascular risk factors. *Atherosclerosis* 1993; **103**(1):81-91.

Hellenius ML, Krakau I, De Faire U. Favourable long-term effects from advice on diet and exercise given to healthy men with raised cardiovascular risks. *Nutrition, Metabolism and Cardiovascular Diseases* 1997; **7**:293-300.

*, Prevention of Cardiovascular Disease: Studies on the Role of Diet and Exercise in the Prevention of Cardiovascular Disease among Middle-Aged Men [PhD thesis]. Huddinge, Sweden: Karolinska Intitute, 1995.

Naslund GK, Fredrikson M, Hellenius ML, De Faire U. Effect of diet and physical exercise intervention programmes on coronary heart disease risk in smoking and non-smoking men in Sweden. *Journal of Epidemiology and Community Health* 1996; **50**(2):131-6.

Heller 1993 {published and unpublished data}

* Heller RF, Knapp JC, Valenti LA, Dobson AJ. Secondary prevention after acute myocardial infarction. *American Journal of Cardiology* 1993; **72**(11):759-62.

Heller RF, Walker RJ, Boyle CA, O'Connell DL, Rusakaniko S, Dobson AJ. A randomised controlled trial of a dietary advice program for relatives of heart attack victims. *Medical Journal of Australia* 1994; **161**(9):529-31.

Hildreth 1951 {published data only}

Hildreth EA, Mellinkoff SM, Blair GW, Hildreth DM. The effect of vegetable fat ingestion on human serum cholesterol concentration. *Circulation* 1951; **3**(5):641-6.

Holm 1990 {published data only (unpublished sought but not used)}

* Holm LE, Nordevang E, Ikkala E, Hallstrom L, Callmer E. Dietary intervention as adjuvant therapy in breast cancer patients - a feasibility study. *Breast Cancer Research and Treatment* 1990; **16**(2):103-9.

Nordevang E, Callmer E, Marmur A, Holm LE. Dietary intervention in breast cancer patients: effects on food choice. *European Journal of Clinical Nutrition* 1992; **46**(6):387-96.

Nordevang E, Ikkala E, Callmer E, Hallstrom L, Holm LE. Dietary intervention in breast cancer patients: effects on dietary habits and nutrient intake. *European Journal of Clinical Nutrition* 1990; **44**(9):681-7.

Horlick 1957 {published data only}

* Horlick L, Craig BM. Effect of long-chain polyunsaturated and saturated fatty acids on the serum-lipids of man. *Lancet* 1957; **2**:566-9.

Horlick 1960 {published data only}

Horlick L, O'Neil JB. Effect of modified egg-yolk fats on bloodcholesterol levels [letter]. *Lancet* 1960; **1**:438.

Howard 1977 {published data only}

Howard AN, Marks J. Hypocholesterolaemic effect of milk [letter]. *Lancet* 1977; **2**(8031):255-6.

Hunninghake 1990 {published data only}

Hunninghake DB, Laskarzewski PM. Gender difference in the response to lovastatin administration with and without a cholesterol lowering diet [abstract]. *Arteriosclerosis* 1990; **10**:786A.

Hutchison 1983 {published data only}

Hutchison K, Oberle K, Crockford P, Grace M, Whyte L, Gee M, et al. Effects of dietary manipulation on vascular status of patients with peripheral vascular disease. *JAMA* 1983; **249**(24):3326-30.

Hyman 1998 {published and unpublished data}

Hyman DJ, Ho KS, Dunn K, Simons-Morton D. Dietary intervention for cholesterol reduction in public clinic patients. *American Journal of Preventive Medicine* 1998; **15**(2):139-45.

lacono 1981 {published data only}

Iacono JM, Judd JT, Marshall MW, Canary JJ, Dougherty RM, Mackin JF, et al. The role of dietary essential fatty acids and prostaglandins in reducing blood pressure. *Progress in Lipid Research* 1981; **20**:349-64.

IMPACT 1995 {*published data only*}

Fielding JE, Mason T, Kinght K, Klesges R, Pelletier KR. A randomized trial of the IMPACT worksite cholesterol reduction program. *American Journal of Preventive Medicine* 1995; **11**(2):120-3.

Iso 1991 {published data only}

Iso H, Konishi M, Terao A, Kiyama M, Tanigaki M, Baba M, et al. A community-based education program for serum cholesterol reduction in urban hypercholesterolemic persons:

comparison of intensive and usual education groups. *Nippon.Koshu.Eisei.Zasshi [Japanese Journal of Public Health]* 1991; **38**(9):751-61.

lves 1993 {published data only}

Ives DG, Kuller LH, Traven ND. Use and outcomes of a cholesterol-lowering intervention for rural elderly subjects. *American Journal of Preventive Medicine* 1993; **9**(5):274-81.

Jalkanen 1991 {published data only}

Jalkanen L. The effect of a weight reduction program on cardiovascular risk factors among overweight hypertensives in primary health care. *Scandinavian Journal of Social Medicine* 1991; **19**(1):66-71.

Jerusalem Nut 1992 {published data only}

Berry EM, Eisenberg S, Friedlander Y, Harats D, Kaufmann NA, Norman Y, et al. Effects of diets rich in monounsaturated fatty acids on plasma lipoproteins: the Jerusalem Nutrition Study. II. Monounsaturated fatty acids vs carbohydrates. *American Journal of Clinical Nutrition* 1992; **56**(2):394-403.

Jula 1990 {published data only}

Jula A, Ronnemaa T, Rastas M, Karvetti RL, Maki J. Longterm nonpharmacological treatment for mild to moderate hypertension. *Journal of Internal Medicine* 1990; **227**(6):413-21.

Junker 2001 {published data only}

Junker R, Pieke B, Schulte H, Nofer R, Neufeld M, Assmann G, et al. Changes in hemostasis during treatment of hypertriglyceridemia with a diet rich in monounsaturated and n-3 polyunsaturated fatty acids in comparison with a low-fat diet. *Thrombosis Research* 2001; **101**(5):355-66.

Karmally 1990 {published data only}

Karmally W, Carpentiri C, Viscardi T, Cheverez V, Holleran S, Ramakrishnan R, et al. Replacing monounsaturated by polyunsaturated fatty acids within an AHA step I diet does not affect the plasma levels or metabolism of low density and high density lipoproteins in normal men [abstract]. *Arteriosclerosis* 1990; **10**:877A.

Karvetti 1992 {published data only}

Karvetti RL, Hakala P. A seven-year follow-up of a weight reduction programme in Finnish primary health care. *European Journal of Clinical Nutrition* 1992; **46**(10):743-52.

Kastarinen 2002 {published data only}

Kastarinen MJ, Puska PM, Korhonen MH, Mustonen JN, Salomaa VV, Sundvall JE, et al. Non-pharmacological treatment of hypertension in primary health care: a 2-year open randomized controlled trial of lifestyle intervention against hypertension in eastern Finland. *Journal of Hypertension* 2002; **20**(12):01.

Kather 1985 {published data only}

Kather H, Wildenberg U, Wieland E. Influence of different dietary conditions in ideal-weight subjects on serum levels of free fatty acids and of glycerol in vivo and on lipid mobilization in vitro [abstract]. *European Journal of Clinical Investigation* 1985; **15**(Suppl 1):.

Katzel 1995 {published data only}

Katzel LI, Bleecker ER, Colman EG, Rogus EM, Sorkin JD, Goldberg AP. Effects of weight loss vs aerobic exercise training on risk factors for coronary disease in healthy, obese, middleaged and older men. A randomized controlled trial [see comments]. *JAMA* 1995; **274**(24):1915-21.

Kawamura 1993 {published data only}

Kawamura M, Akasaka T, Kasatsuki T, Nakajima J, Onodera S, Fujiwara T, et al. Blood pressure is reduced by short-time calorie restriction in overweight hypertensive women with a constant intake of sodium and potassium. *Journal of Hypertension* 1993; **11**(Suppl 5):S320-1.

Keidar 1988 {published data only}

Keidar S, Krul ES, Goldberg AC, Bateman J, Schonfield G. Fatfree diet modulates epitope expression of LDL-apo [abstract]. *Arteriosclerosis* 1988; **8**:565A.

Kempner 1948 {published data only}

Kempner W. Treatment of hypertensive vascular disease with rice diet. *American Journal of Medicine* 1948; **4**(4):545-77.

Keys 1957a {published data only}

Keys A, Anderson JT, Grande F. Serum-cholesterol response to dietary fat [letter]. *Lancet* 1957; **1**:787.

Keys 1957b {published data only}

Keys A, Anderson JT, Grande F. Essential fatty acids, degree of unsaturation, and effect of corn (maize) oil on the serum-cholesterol level in man. *Lancet* 1957; **1**(6959):66-8.

Keys 1957c {published data only}

Keys A. Prediction of serum-cholesterol responses of man to changes in fats in the diet. *Lancet* 1957; **2**:959-66.

Khan 2003 {published and unpublished data}

Khan F, Elherik K, Bolton-Smith C, Barr R, Hill A, Murrie I, et al. The effects of dietary fatty acid supplementation on endothelial function and vascular tone in healthy subjects. *Cardiovascular Research* 2003; **59**(4):955-62.

King 2000 {published data only}

King S, David S, Newton H, Hevey D, Rafferty F, Horgan JH. The effect of dietary modification on the training outcome and body composition in patients undergoing undergoing a cardiac rehabilitation programme. *Coronary Health Care* 2000; **4**(2):76-81.

Kingsbury 1961 {published data only}

Kingsbury KJ, Morgan DM, Aylott C, Emmerson R. Effects of ethyl arachidonate, cod-liver oil, and corn oil on the plasmacholesterol level: a comparison in normal volunteers. *Lancet* 1961; **1**(7180):739-41.

KNOTA {published data only}

ZZZ <accessionId ref="info:x-wiley/clinicalTrialsGov/ NCT00692536"> ZZZNCT00692536

Andersson J, Mellberg C, Otten J, Ryberg M, Rinnstrom D, Larsson C, et al. Left ventricular remodelling changes without



concomitant loss of myocardial fat after long-term dietary intervention. *International Journal of Cardiology* 2016; **216**:92-6.

Blomquist C, Chorell E, Ryberg M, Mellberg C, Larsson C, Lindahl B, et al. Beneficial effects on fatty acid composition and indices of fatty acid desaturase activity with a paleolithic-type diet during a two-year intervention in obese postmenopausal women. *Endocrine Reviews* 2016; **37**(2 Suppl 1):SUN 575.

Blomquist C, Chorell E, Ryberg M, Mellberg C, Worrsjo E, Makoveichuk E, et al. Decreased lipogenesis-promoting factors in adipose tissue in postmenopausal women with overweight on a Paleolithic-type diet. *European Journal of Nutrition* 2018; **57**(8):2877-86.

Franklin KA, Eriksson M, Larsson C, Lindahl B, Mellberg C, Sahlin C, et al. Palaeolithic diet and obstructive sleep apnoea in overweight females: a randomised controlled trial. *European Respiratory Journal* 2016; **48**:PA2376.

* Mellberg C, Sandberg S, Ryberg M, Eriksson M, Brage S, Larsson C, et al. Long-term effects of a Palaeolithic-type diet in obese postmenopausal women: a 2-year randomized trial. *European Journal of Clinical Nutrition* 2014; **68**(3):350-7.

Otten J, Mellberg C, Ryberg M, Sandberg S, Kullberg J, Lindahl B, et al. Strong and persistent effect on liver fat with a Paleolithic diet during a two-year intervention. *International Journal of Obesity* 2016; **40**(5):747-53.

Otten J, Mellberg C, Sandberg S, Hauksson J, Larsson C, Lindahl B, et al. Decrease of visceral adipose tissue and liver fat during diet intervention: a predictor of insulin sensitivity improvement? *Diabetes* 2014; **63**:A16-.

Otten J, Ryberg M, Mellberg C, Andersson T, Chorell E, Lindahl B, et al. Postprandial levels of GLP-1, GIP and glucagon after 2 years of weight loss with a Paleolithic diet: a randomised controlled trial in healthy obese women. *European Journal of Endocrinology* 2019; **180**(6):417-27.

Otten J, Ryberg M, Mellberg C, Lindahl B, Larsson C, Juul Holst J, et al. Weight loss by two different diets increases the postprandial response of GLP-1 but only the Paleolithic diet increases the postprandial response of GIP. *Diabetologia* 2017; **60**(1):S233-.

Otten J, Stomby A, Ryberg M, Svensson M, Hauksson J, Olsson T. Effects of a paleolithic diet with and without supervised exercise on liver fat and insulin sensitivity: a randomised controlled trial in individuals with type 2 diabetes. *Diabetologia* 2016; **59**(1):S10-.

Otten J, Stomby A, Waling M, Isaksson A, Soderstrom I, Ryberg M, et al. A heterogeneous response of liver and skeletal muscle fat to the combination of a Paleolithic diet and exercise in obese individuals with type 2 diabetes: a randomised controlled trial. *Diabetologia* 2018; **61**(7):1548-59.

Otten J, Stomby A, Waling M, Isaksson A, Tellstrom A, Lundin-Olsson L, et al. Benefits of a Paleolithic diet with and without supervised exercise on fat mass, insulin sensitivity, and glycemic control: a randomized controlled trial in individuals with type 2 diabetes. *Diabetes/Metabolism Research and* *Reviews* 2017; **33**(1):e2828. [DOI: https://doi.org/10.1002/ dmrr.2828]

Stomby A, Otten J, Ryberg M, Nyberg L, Olsson T, Boraxbekk CJ. A paleolithic diet with and without combined aerobic and resistance exercise increases functional brain responses and hippocampal volume in subjects with type 2 diabetes. *Frontiers in Aging Neuroscience* 2017; **9**(DEC):391.

Stomby A, Simonyte K, Mellberg C, Ryberg M, Stimson RH, Larsson C, et al. Diet-induced weight loss has chronic tissuespecific effects on glucocorticoid metabolism in overweight postmenopausal women. *International Journal of Obesity* 2015; **39**(5):814-9.

Koopman 1990 {published data only}

Koopman H, Spreeuwenberg C, Westerman RF, Donker AJ.
Dietary treatment of patients with mild to moderate hypertension in a general practice: a pilot intervention study (2). Beyond three months. *Journal of Human Hypertension* 1990;
4(4):372-4.

Koranyi 1963 {published data only}

Koranyi A. Prophylaxis and treatment of the coronary syndrome. *Therapia Hungarica* 1963; **11**:17-20.

Korhonen 2003 {published data only}

Korhonen M, Kastarinen M, Uusitupa M, Puska P, Nissinen A. The effect of intensified diet counselling on the diet of hypertensive subjects in primary health care: a 2-year open randomized controlled trial of lifestyle intervention against hypertension in eastern Finland. *Preventive Medicine* 2003; **36**(1):8-16.

Kriketos 2001 {published data only}

Kriketos AD, Robertson RM, Sharp TA, Drougas H, Reed GW, Storlien LH, et al. Role of weight loss and polyunsaturated fatty acids in improving metabolic fitness in moderately obese, moderately hypertensive subjects. *Journal of Hypertension* 2001; **19**(10):1745-54.

Kris 1994 {published data only}

Kris EP, Mustad VA. Chocolate feeding studies: a novel approach for evaluating the plasma lipid effects of stearic acid. *American Journal of Clinical Nutrition* 1994; **60**(6 Suppl):1029S-36S.

Kristal 1997 {published data only}

Kristal AR, Shattuck AL, Bowen DJ, Sponzo RW, Nixon DW. Feasibility of using volunteer research staff to deliver and evaluate a low-fat dietary intervention: the American Cancer Society Breast Cancer Dietary Intervention Project. *Cancer Epidemiology, Biomarkers and Prevention* 1997; **6**(6):459-67.

Kromhout 1987 {published data only}

Kromhout D, Arntzenius AC, Kempen-Voogd N, Kempen HJ, Barth JD, Van der Voort HA, et al. Long-term effects of linoleic-acid enriched diet, changes in body weight and alcohol consumption on serum total and HDL-cholesterol. *Atherosclerosis* 1987; **66**(1-2):99-105.



Kummel 2008 {published data only}

Kummel MV. Effects of an intervention on health behaviors of older coronary artery bypass (CAB) patients. *Archives of Gerontology and Geriatrics* 2008; **2**(2):227-44.

Laitinen 1993 {published data only}

Laitinen JH, Ahola IE, Sarkkinen ES, Winberg RL, Harmaakorpi IP, Uusitupa MI. Impact of intensified dietary therapy on energy and nutrient intakes and fatty acid composition of serum lipids in patients with recently diagnosed non-insulin-dependent diabetes mellitus. *Journal of the American Dietetic Association* 1993; **93**(3):276-83.

Laitinen 1994 {published data only}

Laitinen J, Uusitupa M, Ahola I, Siitonen O. Metabolic and dietary determinants of serum lipids in obese patients with recently diagnosed non-insulin-dependent diabetes. *Annals of Medicine* 1994; **26**(2):119-24.

Lean 1997 {published and unpublished data}

Han TS, Richmond P, Avenell A, Lean ME. Waist circumference reduction and cardiovascular benefits during weight loss in women. *International Journal of Obesity and Related Metabolic Disorders* 1997; **21**(2):127-34.

* Lean ME, Han TS, Prvan T, Richmond PR, Avenell A. Weight loss with high and low carbohydrate 1200 kcal diets in free living women. *European Journal of Clinical Nutrition* 1997; **51**(4):243-8.

Leduc 1994 {published data only}

Leduc CP, Cherniak D, Faucher J. Effectiveness of a group dietary intervention on hypercholesterolaemia: a randomised controlled clinical trial [poster abstract]. *Atherosclerosis* 1994; **19**:149.

Lewis 1958 {published data only}

Lewis B. Effect of certain dietary oils on bile-acid secretion and serum-cholesterol. *Lancet* 1958; **1**(7030):1090-2.

Lewis 1981 {published data only}

Lewis B, Hammett F, Katan M, Kay RM, Merkx I, Nobels A, et al. Towards an improved lipid-lowering diet: additive effects of changes in nutrient intake. *Lancet* 1981; **2**(8259):1310-3.

Lewis 1985 {published data only}

Lewis B. Randomised controlled trial of the treatment of hyperlipidaemia on progression of atherosclerosis. *Acta Medica Scandinavica* 1985; **701**(Suppl):53-7.

Lichtenstein 2002 {published data only}

Lichtenstein AH, Ausman LM, Jalbert SM, Vilella-Bach M, Jauhiainen M, McGladdery S, et al. Efficacy of a therapeutic lifestyle change/step 2 diet in moderately hypercholesterolemic middle-aged and elderly female and male subjects. *Journal of Lipid Research* 2002; **43**(2):264-73.

Lim 2010 {published data only}

Lim SS, Noakes M, Keogh JB, Clifton PM. Long-term effects of a low carbohydrate, low fat or high unsaturated fat diet

compared to a no-intervention control. *Nutrition, Metabolism, and Cardiovascular Diseases* 2010; **20**(8):599-607.

Linko 1957 {published data only}

Linko E. Vegetable oils and serum cholesterol: short-term experiments with rapeseed and sunflower oils. *Acta Medica Scandinavica* 1957; **159**(6):475-88.

Lipid Res Clinic 1984 {published data only}

Gordon DJ, Salz KM, Roggenkamp KJ. Dietary determinants of plasma cholesterol change in the recruitment phase of the Lipid Research Clinics Coronary Primary Prevention Trial. *Arteriosclerosis* 1982; **2**(6):537-48.

* Lipid Research Clinics. The Lipid Research Clinics Coronary Primary Prevention Trial results. I. Reduction in incidence of coronary heart disease. *JAMA* 1984; **251**(3):351-64.

Lipid Research Clinics. The Lipid Research Clinics Coronary Primary Prevention Trial results. II. The relationship of reduction in incidence of coronary heart disease to cholesterol lowering. *JAMA* 1984; **251**(3):365-74.

Little 1990 {published data only}

Little P, Girling G, Hasler A, Craven A, Trafford A. The effect of a combination low sodium, low fat, high fibre diet on serum lipids in treated hypertensive patients. *European Journal of Clinical Nutrition* 1990; **44**(4):293-300.

Little 2004 {published data only}

Little P, Kelly J, Barnett J, Dorward M, Margetts B, Warm D, et al. Randomised controlled factorial trial of dietary advice for patients with a single high blood pressure reading in primary care. *BMJ* 2004; **328**(7447):1054.

Lottenberg 1996 {published data only}

Lottenberg AM, Nunes VS, Lottenberg SA, Shimabukuro AF, Carrilho AJ, Malagutti S, et al. Plasma cholesteryl ester synthesis, cholesteryl ester transfer protein concentration and activity in hypercholesterolemic women: effects of the degree of saturation of dietary fatty acids in the fasting and postprandial states. *Atherosclerosis* 1996; **126**(2):265-75.

Luszczynska 2007 {published data only}

Luszczynska A, Scholz U, Sutton S. Planning to change diet: a controlled trial of an implementation intentions training intervention to reduce saturated fat intake among patients after myocardial infarction. *Journal of Psychosomatic Research* 2007; **63**(5):491-7.

Lyon Diet Heart 1994 {published data only}

* De Lorgeril M, Renaud S, Mamelle N, Salen P, Martin JL, Monjaud I, et al. Mediterranean alpha-linolenic acid-rich diet in secondary prevention of coronary heart disease. *Lancet* 1994; **343**(8911):1454-9.

De Lorgeril M, Salen P, Caillat-Vallet E, , Barthelemy JC, Mamelle N. Control of bias in dietary trial to prevent coronary recurrences: the Lyon Diet Heart study. *European Journal of Clinical Nutrition* 1997; **51**(2):116-22.



De Lorgeril M, Salen P, , Monjaud I, Delaye J, Mamelle N. Mediterranean diet, traditional risk factors, and the rate of cardiovascular complications after myocardial infarction: final report of the Lyon Diet Heart study. *Circulation* 1999; **99**(6):779-85.

De Lorgeril M, Salen P, Martin JL, Mamelle N, Monjaud I, Touboul P, et al. Effect of a Mediterranean type of diet on the rate of cardiovascular complications in patients with coronary artery disease. insights into the cardioprotective effect of certain nutriments. *Journal of the American College of Cardiology* 1996; **28**(5):1103-8.

De Lorgeril M, Salen P, Martin JL, Monjaud I, Boucher P, Mamelle N. Mediterranean dietary pattern in a randomised trial. *Archives of Internal Medicine* 1998; **158**(11):1181-7.

De Lorgeril M, Salen P. Mediterranean diet in secondary prevention of coronary heart disease. *Australian Journal of Nutrition and Dietetics* 1998; **55**(Suppl):s16-s20.

Renaud S, De Lorgeril M, Delaye J, Guidollet J, Jacquard F, Mamelle N, et al. Cretan Mediterranean diet for prevention of coronary heart disease. *American Journal of Clinical Nutrition* 1995; **61**(6 Suppl):1360S-7S.

Lysikova 2003 {published data only}

Lysikova SL, Pogozheva AV, Akol'zina SE, Vasil'ev AV, Vorob'eva LS. The study of the clinical potency of antiatherogenic diet containing flavonoids in cardiovascular patients. *Voprosy Pitaniia* 2003; **72**(3):8-11.

Macdonald 1972 {published data only}

Macdonald I. Relationship between dietary carbohydrates and fats in their influence on serum lipid concentrations. *Clinical Science* 1972; **43**(2):265-74.

Mansel 1990 {published data only}

Mansel RE, Harrison BJ, Melhuish J, Sheridan W, Pye JK, Pritchard G, et al. A randomized trial of dietary intervention with essential fatty acids in patients with categorized cysts. *Annals of the New York Academy of Sciences* 1990; **586**:288-94.

MARGARIN 2002 {published data only}

Bemelmans WJ, Broer J, Feskens EJ, Smit AJ, Muskiet FA, Lefrandt JD, et al. Effect of an increased intake of alphalinolenic acid and group nutritional education on cardiovascular risk factors: the Mediterranean Alpha-linolenic Enriched Groningen Dietary Intervention (MARGARIN) study. *American Journal of Clinical Nutrition* 2002; **75**(2):221-7.

Marniemi 1990 {published and unpublished data}

Hakala P, Karvetti RL. Weight reduction on lactovegetarian and mixed diets. *European Journal of Clinical Nutrition* 1989; **43**(6):421-30.

* Marniemi J, Seppanen A, Hakala P. Long-term effects on lipid metabolism of weight reduction on lactovegetarian and mixed diet. *International Journal of Obesity* 1990; **14**(2):113-25.

Mattson 1985 {published data only}

Mattson FH, Grundy SM. Comparison of effects of dietary saturated, monounsaturated and polyunsaturated fatty acids on plasma lipids and lipoproteins in man. *Journal of Lipid Research* 1985; **26**(2):194-202.

McAuley 2005 {published and unpublished data}

* McAuley KA, Hopkins CM, Smith KJ, McLay RT, Williams SM, Taylor RW, et al. Comparison of a high-fat and high-protein diets with a high-carbohydrate diet in insulin-resistant obese women. *Diabetologia* 2005; **48**:8-16.

McAuley KA, Smith KJ, Taylor RW, McLay RT, Williams SM, Mann JI. Long-term effects of popular dietary approaches on weight loss and features of insulin resistance. *International Journal of Obesity* 2006; **30**(2):342-9.

McCarron 1997 {published data only}

McCarron DA, Oparil S, Chait A, Haynes RB, Kris EP, Stern JS, et al. Nutritional management of cardiovascular risk factors. A randomized clinical trial. *Archives of Internal Medicine* 1997; **157**(2):169-77.

McCarron 2001 {published data only}

McCarron DA, Reusser ME. Reducing cardiovascular disease risk with diet. *Obesity Research* 2001; **9**(Suppl 4):335S-40S.

McKeown-Eyssen 1994 {published and unpublished data}

McKeown-Eyssen GE, Bright SE, Bruce WR, Jazmaji V. A randomized trial of a low fat high fibre diet in the recurrence of colorectal polyps. *Journal of Clinical Epidemiology* 1994; **47**(5):525-36.

McManus 2001 {published and unpublished data}

McManus K, Antinoro L, Sacks F. Randomized controlled trial of a moderate-fat low-energy diet compared with a low fat, lowenergy diet for weight loss in overweight adults. *International Journal of Obesity and Related Metabolic Disorders* 2001; **25**(10):1503-11.

McNamara 1981 {published data only}

McNamara DJ, Kolb R, Parker T, Batwin H, Brown C, Samuel P, et al. Diet and cholesterol homeostasis in men [abstract]. *Arteriosclerosis* 1981; **1**:369A.

MeDiet 2002 {published and unpublished data}

Carruba G, Granata OM, Pala V, Campisi I, Agostara B, Cusimano R, et al. A traditional Mediterranean diet decreases endogenous estrogens in healthy postmenopausal women. *Nutrition and Cancer* 2006; **56**(2):253-9.

* Castagnetta L, Granata OM, Cusimano R, Ravazzolo B, Liquori M, Polito L, et al. The Mediet Project. *Annals of the New York Academy of Science* 2002; **963**:282-9.

Granata OM, Traina A, Ramirez S, Campisi I, Zarcone M, Amodio R, et al. Dietary enterolactone affects androgen and estrogen levels in healthy postmenopausal women. *Annals of the New York Academy of Science* 2009; **1155**:232-6.

MEDINA {published data only}

Papamiltiadous ES, Roberts SK, Nicoll AJ, Ryan MC, Itsiopoulos C, Salim A, et al. A randomised controlled trial of a Mediterranean Dietary Intervention for Adults with Non Alcoholic Fatty Liver Disease (MEDINA): study protocol. *BMC Gastroenterology* 2016; **16**:14.

Medi-RIVAGE 2004 {published and unpublished data}

Borel P, Moussa M, Reboul E, Lyan B, Defoort C, Vincent-Baudry S, et al. Human fasting plasma concentrations of vitamin E and carotenoids, and their association with genetic variants in apo C-III, cholesteryl ester transfer protein, hepatic lipase, intestinal fatty acid binding protein and microsomal triacylglycerol transfer protein. *British Journal of Nutrition* 2009; **101**(5):680-7.

Borel P, Moussa M, Reboul E, Lyan B, Defoort C, Vincent-Baudry S, et al. Human plasma levels of vitamin E and carotenoids are associated with genetic polymorphisms in genes involved in lipid metabolism. *Journal of Nutrition* 2007; **137**(12):2653-9.

Gastaldi M, Diziere S, Defoort C, Portugal H, Lairon D, Darmon M, et al. Sex-specific association of fatty acid binding protein 2 and microsomal triacylglycerol transfer protein variants with response to dietary lipid changes in the 3-mo Medi-RIVAGE primary intervention study. *American Journal of Clinical Nutrition* 2007; **86**(6):1633-41.

* Vincent S, Gerber M, Bernard MC, Defoort C, Loundou A, Portugal H, et al. The Medi-RIVAGE study (Mediterranean Diet, Cardiovascular Risks and Gene Polymorphisms): rationale, recruitment, design, dietary intervention and baseline characteristics of participants. *Public Health Nutrition* 2004; **7**(4):531-42.

Vincent-Baudry S, Defoort C, Gerber M, Bernard MC, Verger P, Helal O, et al. The Medi-RIVAGE study: reduction of cardiovascular disease risk factors after a 3-mo intervention with a Mediterranean-type diet or a low-fat diet. *American Journal of Clinical Nutrition* 2005; **82**(5):964-71.

Mensink 1987 {published data only}

Mensink RP, Katan MB. Effect of monounsaturated fatty acids versus complex carbohydrates on high-density lipoproteins in healthy men and women. *Lancet* 1987; **1**(8525):122-5.

Mensink 1989 {published data only}

Mensink RP, Katan MB. Effect of a diet enriched with monounsaturated or polyunsaturated fatty acids on levels of low density and high density lipoprotein cholesterol in healthy women and men. *New England Journal of Medicine* 1989; **321**(7):436-41.

Mensink 1990a {published data only}

Mensink RP, Katan MB. Effect of dietary trans fatty acids on high density and low density lipoprotein cholesterol levels in healthy subjects. *New England Journal of Medicine* 1990; **323**(7):439-45.

Mensink 1990b {published and unpublished data}

Mensink RP. Effect of Monounsaturated Fatty Acids on High-Density and Low-Density Lipoprotein Cholesterol Levels and Blood Pressure in Healthy Men and Women. Wageningen, Netherlands: Wageningen University and Research Centre, 1990.

Metroville Health 2003 {published data only (unpublished sought but not used)}

Aziz KU, Dennis B, Davis CE, Sun K, Burke G, Manolio T, et al. Efficacy of CVD risk factor modification in a lower-middle class community in Pakistan: the Metroville Health Study. *Asia Pacific Journal of Public Health* 2003; **15**(1):30-6.

Michalsen 2006 {published and unpublished data}

Michalsen A, Lehmann N, Pithan C, Knoblauch NT, Moebus S, Kannenberg F, et al. Mediterranean diet has no effect on markers of inflammation and metabolic risk factors in patients with coronary artery disease. *European Journal of Clinical Nutrition* 2006; **60**(4):478-85.

Miettinen 1994 {published data only}

Miettinen TA, Vanhanen H. Dietary sitostanol related to absorption, synthesis and serum level of cholesterol in different apolipoprotein E phenotypes. *Atherosclerosis* 1994; **105**(2):217-26.

Millar 1973 {published data only}

Millar JH, Zilkha KJ, Langman MJ, Payling-Wright H, Smith AD, Belin J, et al. Double-blind trial of linoleate supplementation of the diet in multiple sclerosis. *BMJ* 1973; **1**(5856):765-8.

Miller 1998 {published data only}

Miller ER, Appel LJ, Risby TH. Effect of dietary patterns on measures of lipid peroxidation: results from a randomised clinical trial. *Circulation* 1998; **98**(22):2390-5.

Miller 2001 {published and unpublished data}

Miller SL, Reber RJ, Chapman-Novakofski K. Prevalence of CVD risk factors and impact of a two-year education program for premenopausal women. *Women's Health Issues* 2001; **11**(6):486-93.

Milne 1994 {published data only}

Milne RM, Mann JI, Chisholm AW, Williams SM. Long-term comparison of three dietary prescriptions in the treatment of NIDDM. *Diabetes Care* 1994; **17**(1):74-80.

Minnesota Coronary 1989 {published data only}

Brewer ER, Ashman PL, Kuba K. The Minnesota Coronary Survey: composition of diets, adherence and serum lipid response [Abstract]. *Circulation* 1975; **51 and 52**(Suppl II):269.

Dawson EA, Gatewood LC. The Minnesota Coronary Survey: methodology and characteristics of the population [Abstract]. *Circulation* 1975; **51 and 52**(Suppl II):271.

* , Dawson EA, Ashman PL, Gatewood LC, Bartsch GE, Kuba K, et al. Test of effect of lipid lowering by diet on cardiovascular risk. The Minnesota Coronary Survey. *Arteriosclerosis* 1989; **9**(1):129-35.

, Dawson EA, Kuba K, Brewer ER, Gatewood LC, Bartsch GE. The Minnesota Coronary Survey: effect of diet on cardiovascular events and deaths [abstract]. *Circulation* 1975; **51 and 52**(Suppl II):4.



Minnesota HHP 1990 {published data only}

Murray DM, Kurth C, Mullis R, Jeffery RW. Cholesterol reduction through low-intensity interventions: results from the Minnesota Heart Health Program. *Preventive Medicine* 1990; **19**(2):181-9.

Mojonnier 1980 {published data only}

Mojonnier ML, Hall Y, Berkson DM, Robinson E, Wethers B, Pannbacker B, et al. Experience in changing food habits of hyperlipidaemic men and women. *Journal of the American Dietetic Association* 1980; **77**(2):140-8.

Mokuno 1988 {published data only}

Mokuno H, Yamada N, Sugimoto T. Cholesterol free diet in heterozygous familial hypercholesterolaemia: significant lowering effect on plasma cholesterol (abstract). *Arteriosclerosis* 1988; **8**(5):590a.

Mortensen 1983 {published data only}

Mortensen JZ, Schmidt EB, Nielsen AH, Dyerberg J. The effect of N-6 and N-3 polyunsaturated fatty acids on hemostasis, blood lipids and blood pressure. *Thrombosis and Haemostasis* 1983; **50**(2):543-6.

Mottalib 2018 {published data only}

Mottalib A, Mitri J, Salsberg V, Ashrafzadeh S, Elseaidy T, Tomah S, et al. Effect of dairy consumption and Its fat content on glycemic control and cardiovascular risk factors in patients with type 2 diabetes — a randomized controlled study. *Diabetes* 2018; **67**(Suppl 1):760-P.

MRFIT substudy 1986 {published data only}

Daniel GJ, Dolecek TA, Caggiula AW, Van Horn LV, Epley L, Randall BL. Increasing the use of meatless meals: a nutrition intervention substudy in the Multiple Risk Factor Intervention Trial (MRFIT). *Journal of the American Dietetic Association* 1986; **86**(6):778-81.

MSDELTA 1995 {published data only}

Ginsberg HN. New directions in dietary studies and heart disease: the National Heart, Lung and Blood Institute sponsored multicenter study of diet effects on lipoproteins and thrombogenic activity. *Advances In Experimental Medicine and Biology* 1995; **369**:241-7.

MSFAT 1997 {published and unpublished data}

* Van het Hoff K, Weststrate JA, Van den Berg H, Velthuis-te Wierik EJ, De Graaf C, Zimmermanns NJ, et al. A long-term study on the effect of spontaneous consumption of reduced fat products as part of a normal diet on indicators of health. *International Journal of Food Sciences and Nutrition* 1997; **48**(1):19-29.

Velthuis-te Wierik EJ, Van Leeuwen RE, Hendriks HF, Verhagen H, Loft S, Poulsen HE, et al. Short-term moderate energy restriction does not affect indicators of oxidative stress and genotoxicity in humans. *Journal of Nutrition* 1995; **125**(10):2631-9.

Velthuis-te Wierik EJ, Van den Berg H, Weststrate JA, Van het Hoff KH, De Graaf C. Consumption of reduced-fat products: effects on parameters of anti-oxidative capacity. *European Journal of Clinical Nutrition* 1996; **50**(4):214-9.

Weststrate JA, Van het Hof KH, Van den Berg H, Velthuis-te Wierik EJ, De Graaf C, Zimmermanns NJ, et al. A comparison of the effect of free access to reduced fat products or their full fat equivalents on food intake, body weight, blood lipids and fat-soluble antioxidants levels and haemostasis variables. *European Journal of Clinical Nutrition* 1998; **52**(6):389-95.

Mujeres Felices 2003 {published data only}

* Fitzgibbon ML, Gapstur SM, Knight SJ. Mujeres felices por ser saludables: a breast cancer risk reduction program for Latino women. *Preventive Medicine* 2003; **36**(5):536-46.

Fitzgibbon ML, Gapstur SM, Knight SJ. Results of Mujeres Felices por ser Saludables: a dietary/breast health randomized clinical trial for Latino women. *Annals of Behavioral Medicine* 2004; **28**(2):95-104.

Mutanen 1997 {published data only}

Mutanen M. Comparison between dietary monounsaturated and polyunsaturated fatty acids as regards diet-related diseases. *Biomedicine and Pharmacotherapy* 1997; **51**(8):314-7.

Muzio 2007 {published data only}

Muzio F, Mondazzi L, Harris WS, Sommariva D, Branchi A. Effects of moderate variations in the macronutrient content of the diet on cardiovascular disease risk factors in obese patients with the metabolic syndrome. *American Journal of Clinical Nutrition* 2007; **86**(4):946-51.

Naglak 2000 {published data only (unpublished sought but not used)}

Naglak MC, Mitchell DC, Shannon BM, Pearson TA, Harkness WL, Kris-Etherton PM. Nutrient adequacy of diets of adults with hypercholesterolemia after a cholesterol-lowering intervention: long term assessment. *Journal of the American Dietetic Association* 2000; **100**(11):1385-91.

NAS 1987 {published data only}

Chlebowski RT, Nixon DW, Blackburn GL, Jochimsen P, Scanlon EF, Insull W, et al. A breast cancer Nutrition Adjuvant Study (NAS): protocol design and initial patient adherence. *Breast Cancer Research and Treatment* 1987; **10**(1):21-9.

National Diet Heart 1968 {published data only}

Baker BM, , Keys A, Kinsell LW, Page IH, Stamler J, et al. The National Diet-Heart Study: an initial report. *JAMA* 1963; **185**:105-6.

Brown HB. The National Diet Heart Study - implications for dietitians and nutritionists. *Journal of the American Dietetic Association* 1968; **52**(4):279-87.

Page IH, Brown HB. Some observations on the National Diet-Heart Study. *Circulation* 1968; **37**(3):313-5.

* The National Diet-Heart Study Research Group. The National Diet-Heart Study final report. *Circulation* 1968; **37**(II):1-428.

The National Diet-Heart Study Research Group. The National Diet-Heart Study. *Nutrition Reviews* 1968; **26**(5):133-6.



NCEP weight 1991 {published and unpublished data}

Wood PD, Stefanick ML, Williams PT, Haskell WL. The effects on plasma lipoproteins of a prudent weight-reducing diet, with or without exercise, in overweight men and women. *New England Journal of Medicine* 1991; **325**(7):461-6.

NCT01954472 {published data only} ZZZ <accessionId ref="info:x-wiley/clinicalTrialsGov/ NCT01954472"> ZZZNCT01954472

. Enhanced multicenter dietary portfolio study (EDP8). clinicaltrials.gov/ct2/show/NCT01954472 (first posted 1 October 2013).

NCT03068078 {published data only}

. A reduced-carbohydrate diet high in monounsaturated fats in type 2 diabetes (ReDuCtion). clinicaltrials.gov/ct2/show/ NCT03068078 (first posted 1 March 2017).

Neil 1995 {published data only}

Neil HA, Roe L, Godlee RJ, Moore JW, Clark GM, Brown J, et al. Randomised trial of lipid lowering dietary advice in general practice: the effects on serum lipids, lipoproteins, and antioxidants [see comments]. *BMJ* 1995; **310**(6979):569-73.

Neverov 1997 {published data only}

Neverov NI, Kaysen GA, Tareyeva IE. Effect of lipid-lowering therapy on the progression of renal disease in nondiabetic nephrotic patients. *Contributions to Nephrology* 1997; **120**:68-78.

Next Step 1995 {published and unpublished data}

Tilley BC, Vernon SW, Glanz K, Myers R, Sanders K, Lu M, et al. Worksite cancer screening and nutrition intervention for highrisk auto workers: design and baseline findings of the Next Step Trial. *Preventive Medicine* 1997; **26**(2):227-35.

* Tilley BC, Vernon SW, Myers R, Glanz K, Lu M, Sanders K, et al. Planning the next step. A screening promotion and nutrition intervention trial in the work site. *Annals of the New York Academy of Sciences* 1995; **768**:296-9.

Nordoy 1971 {published data only}

Nordoy A, Rodset JM. The influence of dietary fats on platelets in man. *Acta Medica Scandinavica* 1971; **190**(1-2):27-34.

Norway Veg Oil 1968 {published data only}

Natvig H, Borchgrevink CF, Dedichen J, Owren PA, Schiotz EH, Westlund K. A controlled trial of the effect of linolenic acid on incidence of coronary heart disease: the Norwegian Vegetable Oil Experiment of 1965-66. *Scandinavian Journal of Clinical and Laboratory Investigation* 1968; **105**(Suppl):1-20.

Nutri-AGEs 2015 {published data only}

. Effect of low-glycemic index Mediterranean diet on ages (Nutri_AGEs). clinicaltrials.gov/ct2/show/NCT02353416 (first posted 2 February 2015).

Nutrition Breast Health {published and unpublished data}

Djuric Z, Poore KM, Depper JB, Uhley VE, Lababidi S, Covington C, et al. Methods to increase fruit and vegetable intake with and without a decrease in fat intake: compliance and effects on body weight in the Nutrition and Breast Health Study. *Nutrition and Cancer* 2002; **43**(2):141-51.

O'Brien 1976 {published data only}

O'Brien JR, Etherington MD, Jamieson S. Effect of a diet of polyunsaturated fats on some platelet-function tests. *Lancet* 1976; **2**(7993):995-6.

ODES 2006 {published data only}

Anderssen S, Holme I, Urdal P, Hjermann I. Diet and exercise intervention have favourable effects on blood pressure in mild hypertensives: the Oslo Diet and Exercise Study (ODES). *Blood Pressure* 1995; **4**(6):343-9.

Anderssen SA, Hjermann I, Urdal P, Torjesen PA, Holme I. Improved carbohydrate metabolism after physical training and dietary intervention in individuals with the "atherothrombogenic syndrome". Oslo Diet and Exercise Study (ODES). A randomized trial. *Journal of Internal Medicine* 1996; **240**(4):203-9.

* Holme I, Haaheim LL, Tonstad S, Hjermann I, Holme I, Haaheim LL, et al. Effect of dietary and antismoking advice on the incidence of myocardial infarction: a 16-year follow-up of the Oslo Diet and Antismoking Study after its close. *Nutrition Metabolism & Cardiovascular Diseases* 2006; **16**(5):330-8.

ODES Investigators. The Oslo Diet and Exercise Study (ODES): design and objectives. *Controlled Clinical Trials* 1993; **14**(3):229-43.

Rokling-Andersen MH, Reseland JE, Veierod MB, Anderssen SA, , Urdal P, et al. Effects of long-term exercise and diet intervention on plasma adipokine concentrations. *American Journal of Clinical Nutrition* 2007; **86**(5):1293-301.

Torjesen PA, Birkeland KI, Anderssen SA, Hjermann I, Holme I, Urdal P. Lifestyle changes may reverse development of the insulin resistance syndrome. The Oslo Diet and Exercise Study: a randomized trial. *Diabetes Care* 1997; **20**(1):26-31.

Oldroyd 2001 {published data only}

Oldroyd JC, Unwin NC, White M, Mathers JC, Alberti KG. Randomised controlled trial evaluating lifestyle interventions in people with impaired glucose tolerance. *Diabetes Research & Clinical Practice* 2006; **72**(2):117-27.

* Oldroyd JC. Randomised controlled trial evaluating the effectiveness of behavioural interventions to modify cardiovascular risk factors in men and women with impaired glucose tolerance: outcomes at 6 months. *Diabetes Research and Clinical Practice* 2001; **52**(1):29-43.

Ole Study 2002 {published and unpublished data}

* Bray GA, Lovejoy JC, Most-Windhauser M, Smith SR, Volaufova J, Denkins Y, et al. A 9-mo randomized clinical trial comparing fat-substituted and fat-reduced diets in healthy obese men: the Ole Study. *American Journal of Clinical Nutrition* 2002; **76**(5):928-34.

Lovejoy JC, Bray GA, Lefevre M, Smith SR, Most MM, Denkins YM, et al. Consumption of a controlled low-fat diet containing olestra for 9 months improves health risk factors in conjunction



with weight loss in obese men: the Ole Study. *International Journal of Obesity and Related Metabolic Disorders* 2003; **27**(10):1242-9.

OLIVE 1997 {published data only (unpublished sought but not used)}

Colquhoun DM, Somerset S, Glasziou PP, Richards D, Weyers J. Comparison of an olive oil enriched diet to a low fat diet intervention study using vascular endpoints: assessed by repeat quantitative angiography (OLIVE study). *Australian Journal of Nutrition and Dietetics* 1998; **55**(Suppl 4):S24-9.

* Colquhoun DM. Rationale and design of the "OLIVE" study (Comparison of an Olive oil enriched to a low fat diet intervention study using vascular endpoints) [Abstract]. In: 11th International Symposium on Atherosclerosis; 1997 October; Paris. 1997:326.

ORIGIN 2008 {published data only}

Gerstein H, Yusuf S, Riddle MC, Ryden L, Bosch J. Rationale, design, and baseline characteristics for a large international trial of cardiovascular disease prevention in people with dysglycemia: the ORIGIN Trial (Outcome Reduction with an Initial Glargine Intervention). *American Heart Journal* 2008; **155**(1):26-32.

Oslo Study 2004 {published data only}

* Hjerkinn EM, Sandvik L, Hjermann I, Arnesen H. Effect of diet intervention on long-term mortality in healthy middleaged men with combined hyperlipidaemia. *Journal of Internal Medicine* 2004; **255**(1):68-73.

Hjermann I, Leren P, Norman N, Helgeland A, Holme I. Serum insulin response to oral glucose load during a dietary intervention trial in healthy coronary high risk men: the Oslo study. *Scandinavian Journal of Clinical and Laboratory Investigation* 1980; **40**(1):89-94.

Hjermann I, Velve BK, Holme I, Leren P. Effect of diet and smoking intervention on the incidence of coronary heart disease. Report from the Oslo Study Group of a randomised trial in healthy men. *Lancet* 1981; **2**(8259):1303-10.

Hjermann I. Intervention of smoking and eating habits in healthy men carrying high risk for coronary heart disease. The Oslo Study. *Acta Medica Scandinavica* 1981; **651**(Suppl):281-4.

Hjermann I. Smoking and diet intervention in healthy coronary high risk men. Methods and 5-year follow-up of risk factors in a randomized trial. The Oslo study. *Journal of the Oslo City Hospitals* 1980; **30**(1):3-17.

Pascale 1995 {published data only}

Pascale RW, Wing RR, Butler BA, Mullen M, Bononi P. Effects of a behavioral weight loss program stressing calorie restriction versus calorie plus fat restriction in obese individuals with NIDDM or a family history of diabetes. *Diabetes Care* 1995; **18**(9):1241-8.

PEP 2001 {published data only}

Ohrig E, Geib HC, , Schwandt P. The prevention education program (PEP) Nuremberg: design and baseline data of a family

oriented intervention study. *International Journal of Obesity* 2001; **25**(Suppl 1):S89-92.

PHYLLIS 1993 {published data only}

Bond GM, Crepaldi G, Zanchetti A, Avogaro P, Marubini E, Maseri A, et al. Plaque hypertension lipid-lowering Italian study (PHYLLIS): a protocol for non-invasive evaluation of carotid atherosclerosis in hypercholesterolaemic hypertensive subjects. *Journal of Hypertension* 1993; **11**(Suppl 5):S314-5.

Pilkington 1960 {published and unpublished data}

Pilkington TR, Stafford JL, Hankin VS, Simmonds FM, Koerselman HB. Practical diets for lowering serum lipids. *British Medical Journal* 1960; **1**(5165):23-5.

Polyp Prevention 1996 {published and unpublished data}

Lanza E, Schatzkin A, Ballard BR, Clifford DC, Paskett E, Hayes D, et al. The polyp prevention trial II: dietary intervention program and participant baseline dietary characteristics. *Cancer Epidemiology, Biomarkers and Prevention* 1996; **5**(5):385-92.

* Schatzkin A, Lanza E, Freedman LS, Tangrea J, Cooper MR, Marshall JR, et al. The polyp prevention trial I: rationale, design, recruitment, and baseline participant characteristics. *Cancer Epidemiology, Biomarkers and Prevention* 1996; **5**(5):375-83.

POUNDS LOST 2009 {published and unpublished data}

Anton SD, Gallagher J, Carey VJ, Laranjo N, Cheng J, Champagne CM, et al. Diet type and changes in food cravings following weight loss: findings from the POUNDS LOST Trial. *Eating & Weight Disorders* 2012; **17**(2):e101-8.

Anton SD, LeBlanc E, Allen HR, Karabetian C, Sacks F, Bray G, et al. Use of a computerized tracking system to monitor and provide feedback on dietary goals for calorie-restricted diets: the POUNDS LOST study. *Journal of Diabetes Science & Technology* 2012; **6**(5):1216-25.

Bray GA, Smith SR, DeJonge L, De Souza R, Rood J, Champagne CM, et al. Effect of diet composition on energy expenditure during weight loss: the POUNDS LOST Study. *International Journal of Obesity* 2012; **36**(3):448-55.

De Jonge L, Bray GA, Smith SR, Ryan DH, De Souza RJ, Loria CM, et al. Effect of diet composition and weight loss on resting energy expenditure in the POUNDS LOST study. *Obesity* 2012; **20**(12):2384-9.

De Souza RJ, Bray GA, Carey VJ, Hall KD, LeBoff MS, Loria CM, et al. Effects of 4 weight-loss diets differing in fat, protein, and carbohydrate on fat mass, lean mass, visceral adipose tissue, and hepatic fat: results from the POUNDS LOST trial. *American Journal of Clinical Nutrition* 2012; **95**(3):614-25.

Mattei J, Qi Q, Hu FB, Sacks FM, Qi L, Mattei J, et al. TCF7L2 genetic variants modulate the effect of dietary fat intake on changes in body composition during a weight-loss intervention. *American Journal of Clinical Nutrition* 2012; **96**(5):1129-36.

Mirzaei K, Xu M, Qi Q, De Jonge L, Bray GA, Sacks F, et al. Variants in glucose- and circadian rhythm-related genes affect the response of energy expenditure to weight-loss diets: the



POUNDS LOST Trial. *American Journal of Clinical Nutrition* 2014; **99**(2):392-9.

Nicklas JM, Sacks FM, Smith SR, LeBoff MS, Rood JC, Bray GA, et al. Effect of dietary composition of weight loss diets on highsensitivity c-reactive protein: the randomized POUNDS LOST trial. *Obesity* 2013; **21**(4):681-9.

Qi Q, Bray GA, Hu FB, Sacks FM, Qi L, Qi Q, et al. Weight-loss diets modify glucose-dependent insulinotropic polypeptide receptor rs2287019 genotype effects on changes in body weight, fasting glucose, and insulin resistance: the Preventing Overweight Using Novel Dietary Strategies trial. *American Journal of Clinical Nutrition* 2012; **95**(2):506-13.

Qi Q, Bray GA, Smith SR, Hu FB, Sacks FM, Qi L, et al. Insulin receptor substrate 1 gene variation modifies insulin resistance response to weight-loss diets in a 2-year randomized trial: the Preventing Overweight Using Novel Dietary Strategies (POUNDS LOST) trial. *Circulation* 2011; **124**(5):563-71.

Qi Q, Xu M, Wu H, Liang L, Champagne CM, Bray GA, et al. IRS1 genotype modulates metabolic syndrome reversion in response to 2-year weight-loss diet intervention: the POUNDS LOST trial. *Diabetes Care* 2013; **36**(11):3442-7.

* Sacks FM, Bray GA, Carey VJ, Smith SR, Ryan DH, Anton SD, et al. Comparison of weight-loss diets with different compositions of fat, protein, and carbohydrates. *New England Journal of Medicine* 2009; **360**(9):859-73.

Williamson DA, Anton SD, Han H, Champagne CM, Allen R, LeBlanc E, et al. Adherence is a multi-dimensional construct in the POUNDS LOST trial. *Journal of Behavioral Medicine* 2010; **33**(1):35-46.

Williamson DA, Anton SD, Han H, Champagne CM, Allen R, LeBlanc E, et al. Early behavioral adherence predicts short and long-term weight loss in the POUNDS LOST study. *Journal of Behavioral Medicine* 2010; **33**(4):305-14.

Xu M, Qi Q, Liang J, Bray GA, Hu FB, Sacks FM, et al. Genetic determinant for amino acid metabolites and changes in body weight and insulin resistance in response to weight-loss diets: the Preventing Overweight Using Novel Dietary Strategies (POUNDS LOST) trial. *Circulation* 2013; **127**(12):1283-9.

Zhang X, Qi Q, Bray GA, Hu FB, Sacks FM, Qi L, et al. APOA5 genotype modulates 2-y changes in lipid profile in response to weight-loss diet intervention: the Pounds Lost Trial. *American Journal of Clinical Nutrition* 2012; **96**(4):917-22.

Zhang X, Qi Q, Zhang C, Smith SR, Hu FB, Sacks FM, et al. FTO genotype and 2-year change in body composition and fat distribution in response to weight-loss diets: the POUNDS LOST trial. *Diabetes* 2012; **61**(11):3005-11 Erratum in: Diabetes. 2013 Feb;62(2):662; Note: Smith, SR [added]; Bray, GA [added].

PREDIMED 2008 {published data only (unpublished sought but not used)}

Buil-Cosiales P, Irimia P, Ros E, Riverol M, Gilabert R, Martinez-Vila E, et al. Dietary fibre intake is inversely associated with carotid intima-media thickness: a cross-sectional assessment in the PREDIMED study. *European Journal of Clinical Nutrition* 2009; **63**(10):1213-9.

Castañer O, Corella D, Covas MI, Sorlí JV, Subirana I, Flores-Mateo G, et al. In vivo transcriptomic profile after a Mediterranean diet in high-cardiovascular risk patients: a randomized controlled trial. *American Journal of Clinical Nutrition* 2013; **98**(3):845-53.

Díaz-López A, Bulló M, , Guasch-Ferré M, Ros E, Basora J, et al. Effects of mediterranean diets on kidney function: a report from the PREDIMED trial. *American Journal of Kidney Diseases* 2012; **60**(3):380-9.

Damasceno NRT, Sala-Vila A, Cofán M, Pérez-Heras AM, Fitó M, Ruiz-Gutiérrez V, et al. Mediterranean diet supplemented with nuts reduces waist circumference and shifts lipoprotein subfractions to a less atherogenic pattern in subjects at high cardiovascular risk. *Atherosclerosis* 2013; **230**(2):347-53.

Estruch R, Martínez-González MA, Corella D, Salas-Salvadó J, Ruiz-Gutiérrez V, Covas MI, et al. Effects of a Mediterranean-style diet on cardiovascular risk factors: a randomized trial. *Annals of Internal Medicine* 2006; **145**(1):1-11.

* Estruch R, Ros E, Salas-Salvadó J, , Corella D, Arós F, et al, PREDIMED Study Investigators. Primary prevention of cardiovascular disease with a Mediterranean diet. *New England Journal of Medicine* 2013; **368**(14):1279-90. [DOI: 10.1056/ NEJMoa1200303]

Fernández-Real JM, Bulló M, Moreno-Navarrete JM, Ricart W, Ros E, Estruch R, et al. A Mediterranean diet enriched with olive oil is associated with higher serum total osteocalcin levels in elderly men at high cardiovascular risk. *Journal of Clinical Endocrinology & Metabolism* 2012; **97**(10):3792-8.

Guasch-Ferré M, Bulló M, Babio N, Martínez-González MA, Estruch R, Covas MI, et al. Mediterranean diet and risk of hyperuricemia in elderly participants at high cardiovascular risk. *Journals of Gerontology Series A - Biological Sciences & Medical Sciences* 2013; **68**(10):1263-70.

Hu EA, Toledo E, Diez-Espino J, Estruch R, Corella D, Salas-Salvado J, et al. Lifestyles and risk factors associated with adherence to the Mediterranean diet: a baseline assessment of the PREDIMED trial. *PIOS One* 2013; **8**(4):e60166.

Martínez-Lapiscina EH, Clavero P, Toledo E, Estruch R, Salas-Salvadó J, San Julián B, et al. Mediterranean diet improves cognition: the PREDIMED-NAVARRA randomised trial. *Journal of Neurology, Neurosurgery, and Psychiatry* 2013; **84**(12):1318-25.

Martínez-Lapiscina EH, Clavero P, Toledo E, San Julián B, Sanchez-Tainta A, Corella D, et al. Virgin olive oil supplementation and long-term cognition: the Predimed-Navarra randomized, trial. *Journal of Nutrition, Health and Aging* 2013; **17**(6):544-52.

Mitjavila MT, Fandos M, Salas-Salvadó J, Covas MI, Borrego S, Estruch R, et al. The Mediterranean diet improves the systemic lipid and DNA oxidative damage in metabolic syndrome individuals. A randomized, controlled, trial. *Clinical Nutrition* 2013; **32**(2):172-8.



Nordmann AJ, Suter-Zimmermann K, Bucher HC, Shai I, Tuttle KR, Estruch R, et al. Meta-analysis comparing Mediterranean to low-fat diets for modification of cardiovascular risk factors. *American Journal of Medicine* 2011; **124**(9):841-51.

Prieto RM, Fiol M, Perello J, Estruch R, Ros E, Sanchis P, et al. Effects of Mediterranean diets with low and high proportions of phytate-rich foods on the urinary phytate excretion. *European Journal of Nutrition* 2010; **49**(6):321-6.

Razquin C, Martinez JA, Martinez-Gonzalez MA, Bes-Rastrollo M, Fernández-Crehuet J, Marti A. A 3-year intervention with a Mediterranean diet modified the association between the rs9939609 gene variant in FTO and body weight changes. *International Journal of Obesity* 2010; **34**(2):266-72.

Razquin C, Martinez JA, Martinez-Gonzalez MA, Fernández-Crehuet J, Santos JM, Marti A. A Mediterranean diet rich in virgin olive oil may reverse the effects of the -174G/C IL6 gene variant on 3-year body weight change. *Molecular Nutrition & Food Research* 2010; **54 Suppl 1**:S75-82.

Razquin C, Martinez JA, Martinez-Gonzalez MA, Mitjavila MT, Estruch R, Marti A, et al. A 3 years follow-up of a Mediterranean diet rich in virgin olive oil is associated with high plasma antioxidant capacity and reduced body weight gain. *European Journal of Clinical Nutrition* 2009; **63**(12):1387-93.

Razquin C, Martinez JA, Martinez-Gonzalez MA, Salas-Salvado J, Estruch R, Marti A. A 3-year Mediterranean-style dietary intervention may modulate the association between adiponectin gene variants and body weight change. *European Journal of Nutrition* 2010; **49**:311-9.

Ruiz-Canela M, Estruch R, Corella D, Salas-Salvadó J, Martínez-González MA. Association of Mediterranean diet with peripheral artery disease: the PREDIMED randomized trial. *JAMA* 2014; **311**(4):415-7.

Sánchez-Taínta A, Estruch R, Bulló M, Corella D, Gómez-Gracia E, Fiol M, et al. Adherence to a Mediterranean-type diet and reduced prevalence of clustered cardiovascular risk factors in a cohort of 3,204 high-risk patients. *European Journal of Cardiovascular Prevention & Rehabilitation* 2008; **15**(5):589-93.

Sánchez-Villegas A, Galbete C, Martínez-González MA, Martinez JA, Razquin C, Salas-Salvadó J, et al. The effect of the Mediterranean diet on plasma brain-derived neurotrophic factor (BDNF) levels: the PREDIMED-NAVARRA randomized trial. *Nutritional Neuroscience* 2011; **14**(5):195-201.

Sánchez-Villegas A, Martínez-González MA, Estruch R, Salas-Salvadó J, Corella D, Covas MI, et al. Mediterranean dietary pattern and depression: the PREDIMED randomized trial. *BMC Medicine* 2013; **11**:208.

Sala-Vila A, Romero-Mamani ES, Gilabert R, Núñez I, De la Torre R, Corella D, et al. Changes in ultrasound-assessed carotid intima-media thickness and plaque with a mediterranean diet: a substudy of the PREDIMED trial. *Arteriosclerosis, Thrombosis, and Vascular Biology* 2014; **34**(2):439-45. Salas-Salvadó J, Bulló M, Babio N, , Ibarrola-Jurado N, Basora J, et al. Reduction in the incidence of type 2 diabetes with the Mediterranean diet: results of the PREDIMED-Reus nutrition intervention randomized trial. *Diabetes Care* 2011; **34**:14-9.

Salas-Salvadó J, Bulló M, Estruch R, Ros E, Covas MI, Ibarrola-Jurado N, et al. Prevention of diabetes with Mediterranean diets: a subgroup analysis of a randomized trial. *Annals of Internal Medicine* 2014; **160**(1):1-10.

Salas-Salvadó J, Fernandez-Ballart J, Ros E, Martínez-González MA, Fito M, Estruch R, et al. Effect of a Mediterranean diet supplemented with nuts on metabolic syndrome status: one-year results of the PREDIMED randomized trial. *Archives of Internal Medicine* 2008; **168**(22):2449-58.

Salas-Salvado J, Garcia-Arellano A, Estruch R, Marquez-Sandoval F, Corella D, Fiol M, et al. Components of the Mediterranean-type food pattern and serum inflammatory markers among patients at high risk for cardiovascular disease. *European Journal of Clinical Nutrition* 2008; **62**(5):651-9.

Schröder H, De la Torre R, Estruch R, Corella D, Martínez-González MA, Salas-Salvadó J, et al. Alcohol consumption is associated with high concentrations of urinary hydroxytyrosol. *American Journal of Clinical Nutrition* 2009; **90**(5):1329-35.

Schröder H, Fitó M, Estruch R, Martínez-González MA, Corella D, Salas-Salvadó J, et al. A short screener is valid for assessing Mediterranean diet adherence among older Spanish men and women. *Journal of Nutrition* 2011; **141**(6):1140-5.

Solá R, Fitó M, Estruch R, Salas-Salvadó J, Corella D, De La Torre R, et al. Effect of a traditional Mediterranean diet on apolipoproteins B, A-I, and their ratio: a randomized, controlled trial. *Atherosclerosis* 2011; **218**(1):174-80.

Toledo E, Delgado-Rodríguez M, Estruch R, Salas-Salvadó J, Corella D, Gomez-Gracia E, et al. Low-fat dairy products and blood pressure: follow-up of 2290 older persons at high cardiovascular risk participating in the PREDIMED study. *British Journal of Nutrition* 2009; **101**(1):59-67.

Toledo E, Hu FB, Estruch R, Buil-Cosiales P, Corella D, Salas-Salvadó J, et al. Effect of the Mediterranean diet on blood pressure in the PREDIMED trial: results from a randomized controlled trial. *BMC Medicine* 2013; **11**:207.

Urpi-Sarda M, Casas R, Chiva-Blanch G, Romero-Mamani ES, Valderas-Martínez P, Arranz S, et al. Virgin olive oil and nuts as key foods of the Mediterranean diet effects on inflammatory biomakers related to atherosclerosis. *Pharmacological Research* 2012; **65**(6):577-83.

Waterhouse AL. Resveratrol metabolites in urine as biomarker of wine intake in free-living subjects: the PREDIMED Study. *Free Radical Biology & Medicine* 2009; **46**(12):1561.

Zamora-Ros R, Urpi-Sarda M, Lamuela-Raventos RM, Estruch R, Martínez-González MA, Bulló M, et al. Resveratrol metabolites in urine as a biomarker of wine intake in free-living subjects: the PREDIMED Study. *Free Radical Biology & Medicine* 2009; **46**(12):1562-6.

Zazpe I, Estruch R, Toledo E, Sánchez-Taínta A, Corella D, Bulló M, et al. Predictors of adherence to a Mediterranean-type diet in the PREDIMED trial. *European Journal of Nutrition* 2010; **49**(2):91-9.

Zazpe I, Sánchez-Taínta A, Estruch R, Lamuela-Raventos RM, Schröder H, Salas-Salvadó J, et al. A large randomized individual and group intervention conducted by registered dietitians increased adherence to Mediterranean-type diets: the PREDIMED study. *Journal of the American Dietetic Association* 2008; **108**(7):1134-44.

PREMIER 2003 {published and unpublished data}

* Appel LJ, Champagne CM, Harsha DW, Cooper LS, Obarzanek E, Elmer PJ, et al. Effects of comprehensive lifestyle modification on blood pressure control: main results of the PREMIER clinical trial. *JAMA* 2003; **289**(16):2083-93.

Elmer PJ, Obarzanek E, Vollmer WM, Simons-Morton D, Stevens VJ, Young DR, et al. Effects of comprehensive lifestyle modification on diet, weight, physical fitness, and blood pressure control: 18-month results of a randomized trial. *Annals of Internal Medicine* 2006; **144**(7):485-95.

Ledikwe JH, Rolls BJ, Smiciklas-Wright H, Mitchell DC, Ard JD, Champagne C, et al. Reductions in dietary energy density are associated with weight loss in overweight and obese participants in the PREMIER trial. *American Journal of Clinical Nutrition* 2007; **85**(5):1212-21.

Lien LF, Brown AJ, Ard JD, Loria C, Erlinger TP, Feldstein AC, et al. Effects of PREMIER lifestyle modifications on participants with and without the metabolic syndrome. *Hypertension* 2007; **50**(4):609-16.

Lin PH, Appel LJ, Funk K, Craddick S, Chen C, Elmer P, et al. The PREMIER intervention helps participants follow the Dietary Approaches to Stop Hypertension dietary pattern and the current Dietary Reference Intakes recommendations. *Journal of the American Dietetic Association* 2007; **107**(9):1541-51.

Lin PH, Miwa S, Li YJ, Wang Y, Levy E, Lastor K, et al. Factors influencing dietary protein sources in the PREMIER trial population. *Journal of the American Dietetic Association* 2010; **110**(2):291-5.

Maruthur NM, Wang NY, Appel LJ. Lifestyle interventions reduce coronary heart disease risk: results from the PREMIER Trial. *Circulation* 2009; **119**(15):2026-31.

McGuire HL, Svetkey LP, Harsha DW, Elmer PJ, Appel LJ, Ard JD. Comprehensive lifestyle modification and blood pressure control: a review of the PREMIER trial. *Journal of Clinical Hypertension* 2004; **6**(7):383-90.

Obarzanek E, Vollmer WM, Lin PH, Cooper LS, Young DR, Ard JD, et al. Effects of individual components of multiple behavior changes: the PREMIER trial. *American Journal of Health Behavior* 2007; **31**(5):545-60.

Svetkey LP, Erlinger TP, Vollmer WM, Feldstein A, Cooper LS, Appel LJ, et al. Effect of lifestyle modifications on blood pressure by race, sex, hypertension status, and age. *Journal of Human Hypertension* 2005; **19**(1):21-31. Svetkey LP, Harsha DW, Vollmer WM, Stevens VJ, Obarzanek E, Elmer PJ, et al. Premier: a clinical trial of comprehensive lifestyle modification for blood pressure control: rationale, design and baseline characteristics. *Annals of Epidemiology* 2003; **13**(6):462-71.

Young DR, Coughlin J, Jerome GJ, Myers V, Chae SE, Brantley PJ. Effects of the PREMIER interventions on health-related quality of life. *Annals of Behavioral Medicine* 2010; **40**(3):302-12.

Pritchard 2002 {published data only}

Pritchard JE, Nowson CA, Billington T, Wark JD. Benefits of a year-long workplace weight loss program on cardiovascular risk factors. *Nutrition and Dietetics* 2002; **59**(2):87-96.

Puget Sound EP 2000 {published and unpublished data}

Kristal AR, Curry SJ, Shattuck AL, Feng Z, Li S. A randomized trial of a tailored, self-help dietary intervention: the Puget Sound Eating Patterns Study. *Preventive Medicine* 2000; **31**(4):380-9.

Rabast 1979 {published data only}

Rabast U, Schonborn J, Kasper H. Dietetic treatment of obesity with low and high-carbohydrate diets: comparative studies and clinical results. *International Journal of Obesity* 1979; **3**(3):201-11.

Rabkin 1981 {published data only}

Rabkin SW, Boyko E, Streja DA. Relationship of weight loss and cigarette smoking to changes in high-density lipoprotein cholesterol. *American Journal of Clinical Nutrition* 1981; **34**(9):1764-8.

Radack 1990 {published data only}

Radack K, Deck C, Huster G. The comparative effects of n-3 and n-6 polyunsaturated fatty acids on plasma fibrinogen levels: a controlled clinical trial in hypertriglyceridemic subjects. *Journal of the American College of Nutrition* 1990; **9**(4):352-7.

Rasmussen 1995 {published data only}

Rasmussen OW, Thomsen CH, Hansen KW, Vesterlund M, Winther E, Hermansen K. Favourable effect of olive oil in patients with non-insulin-dependent diabetes. The effect on blood pressure, blood glucose and lipid levels of a high-fat diet rich in monounsaturated fat compared with a carbohydrate-rich diet [Gunstig virkning af olivenolie hos ikkeinsulinkraevende diabetikere. Virkningen pa blodtryk, blodglukose og lipidniveauer af en dioet med et hojt indhold af monoumoettet fedt sammenlignet med en kulhydratrig dioet]. *Ugeskrift for Laeger* 1995; **157**(8):1028-32.

Reaven 2001 {published data only}

Reaven GM, Abbasi F, Bernhart S, Coulston A, Darnell B, Dashti N, et al. Insulin resistance, dietary cholesterol, and cholesterol concentration in postmenopausal women. *Metabolism: Clinical & Experimental* 2001; **50**(5):594-7.

Reid 2002 {published data only}

Reid R, Fodor G, Lydon-Hassen K, D'Angelo MS, McCrea J, Bowlby M, et al. Dietary counselling for dyslipidemia in primary care: results of a randomized trial. *Canadian Journal of Dietetic Practice & Research* 2002; **63**(4):169-75.



Renaud 1986 {published data only}

Renaud S, Godsey F, Dumont E, Thevenon C, Ortchanian E, Martin JL. Influence of long-term diet modification on platelet function and composition in Moselle farmers. *American Journal of Clinical Nutrition* 1986; **43**(1):136-50.

Rivellese 1994 {published and unpublished data}

Rivellese AA, Auletta P, Marotta G, Saldalamacchia G, Giacco A, Mastrilli V, et al. Long term metabolic effects of two dietary methods of treating hyperlipidaemia. *BMJ* 1994; **308**(6923):227-31.

Rivellese 2003 {published data only}

Rivellese AA, Maffettone A, Vessby B, Uusitupa M, Hermansen K, Berglund L, et al. Effects of dietary saturated, monounsaturated and n-3 fatty acids on fasting lipoproteins, LDL size and post-prandial lipid metabolism in healthy subjects. *Atherosclerosis* 2003; **167**(1):149-58.

Roderick 1997 {published and unpublished data}

Roderick P, Ruddock V, Hunt P, Miller G. A randomized trial to evaluate the effectiveness of dietary advice by practice nurses in lowering diet-related coronary heart disease risk. *British Journal* of General Practice 1997; **47**(414):7-12.

Roman CHD prev 1986 {published data only}

* Research Group of the Rome Project of Coronary Heart Disease Prevention. Eight-year follow-up results from the Rome Project of Coronary Heart Disease Prevention. *Preventive Medicine* 1986; **15**(2):176-91.

Research Group of the Rome Project of Coronary Heart Disease Prevention. The Roman Coronary Disease Prevention Project: effectiveness of intervention and reduction of mortality over a 10-year period [II Progetto Romano di Prevenzione della Cardiopatia Coronarica: efficacia dell'intervento e riduzione della mortalita in 10 anni]. *Giornale Italiano di Cardiologia* 1986; **16**(3):196-202.

Rose 1987 {published data only}

Rose DP, Boyar AP, Cohen C, Strong LE. Effect of a low fat diet on hormone levels in women with cystic breast disease. I. Serum steroids and gonadotropins. *Journal of the National Cancer Institute* 1987; **78**(4):623-6.

Sarkkinen 1995 {published and unpublished data}

Makinen E, Uusitupa MI, Pietinen P, Aro A, Penttila I. Long term effects of three fat modified diets on serum lipids in free living hypercholesterolaemic subjects (abstract). *European Heart Journal* 1991; **12**:162.

 * Sarkkinen E. Long-Term Feasibility and Effects of Three Different Fat-Modified Diets in Free-Living Hypercholesterolemic Subjects [PhD thesis]. Kuopio, Finland: University of Kuopio, 1995.

Sarkkinen ES, Agren JJ, Ahola I, Ovaskainen ML, Uusitupa MI. Fatty acid composition of serum cholesterol esters, and erythrocyte and platelet membranes as indicators of long-term adherence to fat-modified diets. *American Journal of Clinical Nutrition* 1994; **59**(2):364-70. Sarkkinen ES, Uusitupa MI, Nyyssonen K, Parviainen M, Penttila I, Salonen JT. Effects of two low-fat diets, high and low in polyunsaturated fatty acids, on plasma lipid peroxides and serum vitamin E levels in free-living hypercholesterolaemic men. *European Journal of Clinical Nutrition* 1993; **47**(9):623-30.

Sarkkinen ES, Uusitupa MI, Pietinen P, Aro A, Ahola I, Penttila I, et al. Long-term effects of three fat-modified diets in hypercholesterolemic subjects. *Atherosclerosis* 1994; **105**(1):9-23.

Uusitupa MI, Sarkkinen ES, Torpstrom J, Pietinen P, Aro A. Long-term effects of four fat-modified diets on blood pressure. *Journal of Human Hypertension* 1994; **8**(3):209-18.

Schaefer 1995a {published data only}

Schaefer EJ, Lichtenstein AH, Lamon-Fava S, McNamara JR, Schaefer MM, Rasmussen H, et al. Body weight and low density lipoprotein cholesterol changes after consumption of a low fat ad libitum diet. *JAMA* 1995; **274**(18):1450-5.

Schaefer 1995b {published data only}

Schaefer EJ, Lichtenstein AH, Lamon-Fava S, Contois JH, Li Z, Rasmussen H, et al. Efficacy of a National Cholesterol Education Program Step 2 diet in normolipidaemic and hypercholesterolaemic middle-aged and elderly men and women. *Arteriosclerosis, Thrombosis, and Vascular Biology* 1995; **15**(8):1079-85.

Schectman 1996 {published data only}

Schectman G, Wolff N, Byrd JC, Hiatt JG, Hartz A. Physician extenders for cost-effective management of hypercholesterolemia. *Journal of General Internal Medicine* 1996; **11**(5):277-86.

Schlierf 1995 {published data only}

Schlierf G, Schuler G, Hambrecht R, Niebauer J, Hauer K, Vogel G, et al. Treatment of coronary heart disease by diet and exercise. *Journal of Cardiovascular Pharmacology* 1995; **25**(Suppl 4):S32-4.

Seppanen-Laakso 1992 {published data only}

Seppanen-Laakso T, Vanhanen H, Laakso I, Kohtamaki H, Viikari J. Replacement of butter on bread by rapeseed oil and rapeseed oil-containing margarine: effects on plasma fatty acid composition and serum cholesterol. *British Journal of Nutrition* 1992; **68**:639-54.

Seppelt 1996 {published and unpublished data}

Seppelt B, Weststrate JA, Reinert A, Johnson D, Luder W, Zunft HJ. Long-term effects of nutrition with fat-reduced foods on energy consumption and body weight [Langzeiteffekte einer ernährung mit fettreduzierten lebensmitteln auf die energieaufnahme und das körpergewicht]. Zeitschrift für Ernährungswissenschaft 1996; **35**(4):369-77.

Singh 1991 {*published data only*}

Singh RB, Rastogi SS, Sircar AR. Dietary strategies for risk-factor modification to prevent cardiovascular diseases. *Nutrition* 1991; **7**(3):210-4.

Singh 1992 {published data only}

Singh RB, Niaz MA, Agarwal P, Begom R, Rastogi SS. Effect of antioxidant-rich foods on plasma ascorbic acid, cardiac enzyme, and lipid peroxide levels in patients hospitalized with acute myocardial infarction. *Journal of the American Dietetic Association* 1995; **95**(7):775-80.

Singh RB, Niaz MA, Ghosh S. Effect on central obesity and associated disturbances of low-energy, fruit- and vegetableenriched prudent diet in north Indians. *Postgraduate Medical Journal* 1994; **70**(830):895-900.

* Singh RB, Rastogi SS, Verma R, Bolaki L, Singh R. An Indian experiment with nutritional modulation in acute myocardial infarction. *American Journal of Cardiology* 1992; **69**(9):879-85.

Singh RB, Rastogi SS, Verma R, Laxmi B, Singh R, Ghosh S, et al. Randomised controlled trial of cardioprotective diet in patients with recent acute myocardial infarction: results of one year follow up. *BMJ* 1992; **304**(6833):1015-9.

Sirtori 1992 {published data only}

Sirtori CR, Gatti E, Tremoli E, Galli C, Gianfranceschi G, Franceschini G, et al. Olive oil, corn oil, and n-3 fatty acids differently affect lipids, lipoproteins, platelets, and superoxide formation in type II hypercholesterolemia. *American Journal of Clinical Nutrition* 1992; **56**(1):113-22.

SLIM 2008 {published data only}

Roumen C, Corpeleijn E, Feskens EJ, Mensink M, Saris WH, Blaak EE, et al. Impact of 3-year lifestyle intervention on postprandial glucose metabolism: the SLIM study. *Diabetic Medicine* 2008; **25**(5):597-605.

Sopotsinskaia 1992 {published data only}

Sopotsinskaia EB, Balitskii KP, Tarutinov VI, Zhukova VM, Semenchuk DD, Kozlovskaia SG, et al. Experience with the use of a low-calorie diet in breast cancer patients to prevent metastasis [Opyt primeneniia nizkokaloriinoi diety u bol'nykh rakom molochnoi zhelezy s tsel'iu profilaktiki metastazi]. *Voprosy Onkologii* 1992; **38**(5):592-9.

Soul Food Light {published data only}

Anderson-Loftin W, Barnett S, Bunn P, Sullivan P, Hussey J, Tavakoli A. Soul Food Light: culturally competent diabetes education. *Diabetes Educator* 2005; **31**(4):555-63.

Stanford NAP 1997 {published data only}

Howard PB, Winkleby MA, Albright CL, Bruce B, Fortmann SP. The Stanford Nutrition Action Program: a dietary fat intervention for low-literacy adults. *American Journal of Public Health* 1997; **87**(12):1971-6.

Stanford Weight 1994 {published and unpublished data}

Williams PT, Krauss RM, Stefanick ML, Vranizan KM, Wood PD. Effects of low-fat diet, calorie restriction, and running on lipoprotein subfraction concentrations in moderately overweight men. *Metabolism* 1994; **43**(5):655-63.

Starmans 1995 {published data only}

Starmans KM, Lustermans FT, Kragten HA, Struijker BH, Rilla H. Lowering cholesterol in patients with mild

hypercholesterolaemia does not improve functional properties of large arteries [abstract]. *Netherlands Journal Of Medicine* 1995; **46**:A70.

Steinbach 1996 {published data only}

Steinbach M. A Romanian contribution to the epidemiology and prevention of cardiovascular diseases. *Romanian Journal of Internal Medicine* 1996; **34**(1-2):137-48.

Steptoe 2001 {published data only}

Steptoe A, Kerry S, Rink E, Hilton S. The impact of behavioral counselling on stage of change in fat intake, physical activity, and cigarette smoking in adults at increased risk of coronary heart disease. *American Journal of Public Health* 2001; **91**(2):265-9.

Stevens 2002 {published and unpublished data}

Stevens VJ, Glasgow RE, Toobert DJ, Karanja N, Smith KS. Oneyear results from a brief, computer-assisted intervention to decrease consumption of fat and increase consumption of fruits and vegetables. *Preventive Medicine* 2003; **36**(5):594-600.

* Stevens VJ, Glasgow RE, Toobert DJ, Karanja N, Smith KS. Randomized trial of a brief dietary intervention to decrease consumption of fat and increase consumption of fruits and vegetables. *American Journal of Health Promotion* 2002; **16**(3):129-34.

Stevenson 1988 {published data only}

Stevenson DW, Darga LL, Spafford TR, Ahmad N, Lucas CP. Variable effects of weight loss on serum lipids and lipoproteins in obese patients. *International Journal of Obesity* 1988; **12**(6):495-502.

Strychar 2009 {published and unpublished data}

Strychar I, Cohn JS, Renier G, Rivard M, Aris-Jilwan N, Beauregard H, et al. Effects of a diet higher in carbohydrate/ lower in fat versus lower in carbohydrate/higher in monounsaturated fat on postmeal triglyceride concentrations and other cardiovascular risk factors in type 1 diabetes. *Diabetes Care* 2009; **32**(9):1597-9.

Sweeney 2004 {published data only}

Sweeney M. Effects of very low-fat diets on anginal symptoms. *Medical Hypotheses* 2004; **63**(3):553.

Søndergaard 2003 {published and unpublished data}

Søndergaard E, Møller JE, Egstrup K. Effect of dietary intervention and lipid-lowering treatment on brachial vasoreactivity in patients with ischemic heart disease and hypercholesterolemia. *American Heart Journal* 2003; **145**(5):E19.

TAIM 1992 {published data only}

Davis BR, Blaufox MD, Hawkins CM, Langford HG, Oberman A, Swencionis C, et al. Trial of antihypertensive interventions and management. Design, methods, and selected baseline results. *Controlled Clinical Trials* 1989; **10**(1):11-30.

Davis BR, Blaufox MD, Oberman A, Wassertheil SS, Zimbaldi N, Cutler JA, et al. Reduction in long-term antihypertensive medication requirements. Effects of weight reduction by dietary



intervention in overweight persons with mild hypertension. *Archives of Internal Medicine* 1993; **153**(15):1773-82.

Davis BR, Oberman A, Blaufox MD, Wassertheil SS, Hawkins CM, Cutler JA, et al. Effect of antihypertensive therapy on weight loss. The Trial of Antihypertensive Interventions and Management Research Group. *Hypertension* 1992; **19**(4):393-9.

Langford HG, Davis BR, Blaufox D, Oberman A, Wassertheil-Smoller S, Hawkins M, et al. Effect of drug and diet treatment of mild hypertension on diastolic blood pressure. The TAIM Research. *Hypertension* 1991; **17**(2):210-7.

Oberman A, Wassertheil-Smoller S, Langford HG, Blaufox MD, Davis BR, Blaszkowski T, et al. Pharmacologic and nutritional treatment of mild hypertension: changes in cardiovascular risk status. *Annals of Internal Medicine* 1990; **112**(2):89-95.

Wassertheil-Smoller S, Davis BR, Breuer B, Chee JC, Oberman A, Blaufox MD. Differences in precision of dietary estimates among different population subgroups. *Annals of Epidemiology* 1993; **3**(6):619-28.

* Wassertheil-Smoller S, Oberman A, Blaufox MD, Davis B, Langford H. The Trial of Antihypertensive Interventions and Management (TAIM) study. Final results with regard to blood pressure, cardiovascular risk, and quality of life. *American Journal of Hypertension* 1992; **5**(1):37-44.

Wylie-Rosett J, Wassertheil-Smoller S, Blaufox MD, Davis BR, Langford HG, Oberman A, et al. Trial of antihypertensive intervention and management: greater efficacy with weight reduction than with a sodium-potassium intervention. *Journal of the American Dietetic Association* 1993; **93**(4):408-15.

Tapsell 2004 {published data only (unpublished sought but not used)}

Tapsell LC, Hokman A, Sebastiao A, Denmeade S, Martin G, Calvert GD, et al. The impact of usual dietary patterns, selection of significant foods and cuisine choices on changing dietary fat under 'free living' conditions. *Asia Pacific Journal of Clinical Nutrition* 2004; **13**(1):86-91.

THIS DIET 2008 {published data only}

Tuttle KR, Shuler LA, Packard DP, Milton JE, Daratha KB, Bibus DM, et al. Comparison of low-fat versus Mediterraneanstyle dietary intervention after first myocardial infarction (from the Heart Institute of Spokane Diet Intervention and Evaluation Trial). *American Journal of Cardiology* 2008; **101**(11):1523-30.

TOHP I 1992 {published data only}

Kumanyika SK, Hebert PR, Cutler JA, Lasser VI, Sugars CP, Steffen Batey L, et al. Feasibility and efficacy of sodium reduction in the Trials of Hypertension Prevention, phase I. *Hypertension* 1993; **22**(4):502-12.

Satterfield S, Cutler JA, Langford HG, Applegate WB, Borhani NO, Brittain E, et al. Trials of hypertension prevention. *Annals of Epidemiology* 1991; **1**(5):455-71.

Stevens VJ, Corrigan SA, Obarzanek E, Bernauer E, Cook NR, Hebert P, et al, the TOHP Collaborative Research Group. Weight loss intervention in phase I of the trials of hypertension prevention. *Archives of Internal Medicine* 1993; **153**(7):849-58.

* The Trials of Hypertension Prevention Collaborative Research Group. The effects of nonpharmacologic interventions on blood pressure of persons with high normal levels. Results of the Trials of Hypertension Prevention, Phase I. *JAMA* 1992; **267**(9):1213-20.

Whelton PK, Hebert PR, Cutler J, Applegate WB, Eberlein KA, Klag MJ, et al. Baseline characteristics of participants in phase I of the Trials of Hypertension Prevention. *Annals of Epidemiology* 1992; **2**(3):295-310.

Whelton PK, Kumanyika SK, Cook NR, Cutler JA, Borhani NO, Hennekens CH, et al. Efficacy of nonpharmacologic interventions in adults with high-normal blood pressure: results from phase 1 of the Trials of Hypertension Prevention. *American Journal of Clinical Nutrition* 1997; **65**(2 Suppl):652S-60S.

TONE 1997 {published data only}

Whelton PK, Appel LJ, Espeland MA, Applegate WB, Ettinger-WH J, Kostis JB, et al. Sodium reduction and weight loss in the treatment of hypertension in older persons: a randomized controlled trial of nonpharmacologic interventions in the elderly (TONE). *JAMA* 1998; **279**(11):839-46.

* Whelton PK, Babnson J, Appel LJ, Charleston J, Cosgrove N, Espeland MA, et al. Recruitment in the Trial of NOnpharmacologic intervention in the Elderly (TONE). *Journal of the American Geriatrics Society* 1997; **45**(2):185-93.

Toobert 2003 {published data only}

Toobert DJ, Glasgow RE, Strycker LA, , Radcliffe JL, Wander RC, et al. Biologic and quality-of-life outcomes from the Mediterranean Lifestyle Program: a randomized clinical trial. *Diabetes Care* 2003; **26**(8):2288-93.

Towle 1994 {published data only}

Towle LA, Bergman EA, Joseph E. Low-fat bison-hybrid ground meat has no effects on serum lipid levels in a study of 12 men. *Journal of the American Dietetic Association* 1994; **94**(5):546-8.

TRANSFACT 2006 {published data only}

* Chardigny JM, Malpuech-Brugere C, Dionisi F, Bauman DE, German B, Mensink RP, et al. Rationale and design of the TRANSFACT project phase I: a study to assess the effect of the two different dietary sources of trans fatty acids on cardiovascular risk factors in humans. *Contemporary Clinical Trials* 2006; **27**(4):364-73.

Chardigny JM. Do trans fatty acids from industrially produced sources and from natural sources have the same effect on cardiovascular disease risk factors in healthy subjects? Results of the Trans Fatty Acids Collaboration (TRANSFACT) study. *American Journal of Clinical Nutrition* 2008; **108**(3):558-66.

Treatwell 1992 {published and unpublished data}

Sorensen G, Morris DM, Hunt MK, Hebert JR, Harris DR, Stoddard A, et al. Work-site nutrition intervention and employees' dietary habits: the Treatwell program. *American Journal of Public Health* 1992; **82**(6):877-80.



Tromsø Heart 1989 {published data only}

Knutsen SF, Knutsen R. The Tromsø Heart Study: family approach to intervention on CHD. Feasibility of risk factor reduction in high-risk persons--project description. *Scandinavian Journal of Social Medicine* 1989; **17**(1):109-19.

Troyer 2010 {published and unpublished data}

Racine EF, Lyerly J, Troyer JL, Warren-Findlow J, McAuley WJ. The influence of home-delivered dietary approaches to stop hypertension meals on body mass index, energy intake, and percent of energy needs consumed among older adults with hypertension and/or hyperlipidemia. *Journal of the Academy of Nutrition & Dietetics* 2012; **112**(11):1755-62.

* Troyer JL, Racine EF, Ngugi GW, McAuley WJ. The effect of home-delivered Dietary Approach to Stop Hypertension (DASH) meals on the diets of older adults with cardiovascular disease. *American Journal of Clinical Nutrition* 2010; **91**(5):1204-12.

UK PDS 1996 {published data only}

* Turner R, Cull C, Holman R. United Kingdom Prospective Diabetes Study 17: a 9-year update of a randomized, controlled trial on the effect of improved metabolic control on complications in non-insulin-dependent diabetes mellitus. *Annals of Internal Medicine* 1996; **124**(1 Pt 2):136-45.

Turner RC, Holman RR. Lessons from UK prospective diabetes study. *Diabetes Research and Clinical Practice* 1995; **28**(Suppl):S151-7.

Urbach 1952 {published data only}

Urbach R, Hildreth EA, Wackerman MT. The therapeutic uses of low fat, low cholesterol diets: I. Treatment of essential familial xanthomatosis. *Journal of Clinical Nutrition* 1952; **1**(1):52-6.

Uusitupa 1993 {published data only}

Uusitupa M, Laitinen J, Siitonen O, Vanninen E, Pyorala K. The maintenance of improved metabolic control after intensified diet therapy in recent type 2 diabetes. *Diabetes Research and Clinical Practice* 1993; **19**(3):227-38.

VASTKOST 2012 {published data only} ZZZ <accessionId ref="info:x-wiley/clinicalTrialsGov/ NCT01005498"> ZZZNCT01005498

* Guldbrand H, Dizdar B, Bunjaku B, Lindstrom T, Bachrach-Lindstrom M, Fredrikson M, et al. In type 2 diabetes, randomisation to advice to follow a low-carbohydrate diet transiently improves glycaemic control compared with advice to follow a low-fat diet producing a similar weight loss. *Diabetologia* 2012; **55**(8):2118-27.

Guldbrand H, Lindstrom T, Dizdar B, Bunjaku B, Ostgren CJ, Nystrom FH, et al. Randomization to a low-carbohydrate diet advice improves health related quality of life compared with a low-fat diet at similar weight-loss in type 2 diabetes mellitus. *Diabetes Research & Clinical Practice* 2014; **106**(2):221-7.

Jonasson L, Guldbrand H, Lundberg AK, Nystrom FH. Advice to follow a low-carbohydrate diet has a favourable impact on lowgrade inflammation in type 2 diabetes compared with advice to follow a low-fat diet. *Annals of Medicine* 2014; **46**(3):182-7.

Vavrikova 1958 {published data only}

Vavrikova J. Essential fatty acids, lipid metabolism, and atherosclerosis [letter]. *Lancet* 1958; **1**:1337.

Verheiden 2003 {published data only (unpublished sought but not used)}

Verheiden MW, Van der Veen JE, Van Zadelhoff WM, Bakx C, Koelen MA, Van den Hoogen HJ, et al. Nutrition guidance in Dutch family practice: behavioural determinants of reduction in fat consumption. *American Journal of Clinical Nutrition* 2003; **77**(Suppl):1058S-64S.

WAHA 2016 {published and unpublished data} ZZZ <accessionId ref="info:x-wiley/clinicalTrialsGov/ NCT01634841"> ZZZNCT01634841

Bitok E, Jaceldo-Siegl K, Rajaram S, Serra-Mir M, Roth I, Feitas-Simoes T, et al. Favourable nutrient intake and displacement with long-term walnut supplementation among elderly: results of a randomised trial. *British Journal of Nutrition* 2017; **118**(3):201-9. [DOI: 10.1017/S0007114517001957]

Bitok E, Rajaram S, Ros E. Does a daily walnut supplement given for a year result in body weight gain? FASEB Journal 2016; **30**:1157.5.

Huey L, Bitok E, Kazzi N. Dietary compliance of walnut or no walnut intake in a 1-year randomized intervention trial among free-living elderly in the Walnuts and Healthy Aging Study (WAHA). FASEB Journal 2016; **30**:1157.10.

Rajaram S, Valls-Pedret C, Cofan M, Sabate J, Serra-Mir M, Perez-Heras AM, et al. The Walnuts and Healthy Aging Study (WAHA): protocol for a nutritional intervention trial with walnuts on brain aging. *Frontiers in Aging Neuroscience* 2016; **8**:333. [PMID: 28119602]

* Ros E, Rajaram S, Sala-Vila A, Serra-Mir M, Valls-Pedret C, Cofán M, et al. Effect of a 1-year walnut supplementation on blood lipids among older individuals: findings from the Walnuts and Healthy Aging (WAHA) study. FASEB Journal 2016; **30**(1 Suppl):293.4.

Wass 1981 {published data only}

Wass VJ, Jarrett RJ, Meilton V, Start MK, Mattock M, Ogg CS, et al. Effect of a long-term fat-modified diet on serum lipoprotein levels of cholesterol and triglyceride in patients. *Clinical Science* 1981; **60**(1):81-6.

Wassertheil 1985 {published data only}

Wassertheil SS, Blaufox MD, Langford HG, Oberman A, Cutter G, Pressel S. Prediction of response to sodium intervention for blood pressure control. *Journal of Hypertension* 1986; **4**(5 Suppl):S343-6.

* Wassertheil SS, Langford HG, Blaufox MD, Oberman A, Hawkins M, Levine B, et al. Effective dietary intervention in hypertensives: sodium restriction and weight reduction. *Journal of the American Dietetic Association* 1985; **85**(4):423-30.

WATCH 1999 {published and unpublished data}

Ockene IS, Hebert JR, Ockene JK, Saperia GM, Stanek E, Nicolosi R, et al. Effect of a physician-delivered nutrition

counselling training and an office-support program on saturated fat intake, weight, and serum lipid measurements in a hyperlipidemic population: Worcester Area Trial for Counseling in Hyperlipidemia. *Archives of Internal Medicine* 1999; **159**(7):725-31.

Watts 1988 {published data only}

Watts GF, Ahmed W, Quiney J, Houlston R, Jackson P, Iles C, et al. Effective lipid lowering diets including lean meat. *BMJ* (*Clinical Research Ed*) 1988; **296**(6617):235-7.

Weintraub 1992 {published data only}

Weintraub M, Sundaresan PR, Schuster B. Long-term weight control study. VII (weeks 0 to 210). Serum lipid changes. *Clinical Pharmacology and Therapeutics* 1992; **51**(5):634-41.

Westman 2006 {published data only}

Westman EC, , Olsen MK, Dudley T, Guyton JR. Effect of a lowcarbohydrate, ketogenic diet program compared to a low-fat diet on fasting lipoprotein subclasses. *International Journal of Cardiology* 2006; **110**(2):212-6.

Weststrate 1998 {published data only}

Weststrate JA, Meijer GW. Plant sterol enriched margarines and reduction of plasma total-and LDL-cholesterol concentrations in normocholesterolaemic and mildly hypercholesterolaemic subjects. *European Journal of Clinical Nutrition* 1998; **52**(5):334-43.

WHEL 2007 {published data only}

Bardwell WA, Profant J, Casden DR, Dimsdale JE, Ancoli-Israel S, Natarajan L, et al. The relative importance of specific risk factors for insomnia in women treated for early-stage breast cancer. *Psycho-Oncology* 2008; **17**(1):9-18.

Caan BJ, Flatt SW, Rock CL, Ritenbaugh C, Newman V, Pierce JP, et al, Women's Healthy Eating and Living Group. Low-energy reporting in women at risk for breast cancer recurrence. *Cancer Epidemiology, Biomarkers & Prevention* 2000; **9**(10):1091-7.

Gold EB, Flatt SW, Pierce JP, Bardwell WA, Hajek RA, Newman VA, et al. Dietary factors and vasomotor symptoms in breast cancer survivors: the WHEL Study. *Menopause* 2006; **13**(3):423-33.

Gold EB, Pierce JP, Natarajan L, Stefanick ML, Laughlin GA, Caan BJ, et al. Dietary pattern influences breast cancer prognosis in women without hot flashes: the Women's Healthy Eating and Living trial. *Journal of Clinical Oncology* 2009; **27**(3):352-9.

Hernandez-Valero MA, Thomson CA, Hernandez M, Tran T, Detry MA, Theriault RL, et al. Comparison of baseline dietary intake of Hispanic and matched non-Hispanic white breast cancer survivors enrolled in the Women's Healthy Eating and Living study. *Journal of the American Dietetic Association* 2008; **108**(8):1323-9.

Hong S, Bardwell WA, Natarajan L, Flatt SW, Rock CL, Newman VA, et al. Correlates of physical activity level in breast cancer survivors participating in the Women's Healthy Eating and Living (WHEL) study. *Breast Cancer Research & Treatment* 2007; **101**(2):225-32.

Hyder JA, Thomson CA, Natarajan L, Madlensky L, Pu M, Emond J, et al. Adopting a plant-based diet minimally increased food costs in WHEL Study. *American Journal of Health Behavior* 2009; **33**(5):530-9.

Madlensky L, Natarajan L, Flatt SW, Faerber S, Newman VA, Pierce JP, et al. Timing of dietary change in response to a telephone counseling intervention: evidence from the WHEL study. *Health Psychology* 2008; **27**(5):539-47.

Mortimer JE, Flatt SW, Parker BA, Gold EB, Wasserman L, Natarajan L, et al. Tamoxifen, hot flashes and recurrence in breast cancer. *Breast Cancer Research & Treatment* 2008; **108**(3):421-6.

Newman VA, Thomson CA, Rock CL, Flatt SW, Kealey S, Bardwell WA, et al. Achieving substantial changes in eating behavior among women previously treated for breast cancer: an overview of the intervention. *Journal of the American Dietetic Association* 2005; **105**(3):382-91.

Paxton RJ, Jones LA, Chang S, Hernandez M, Hajek RA, Flatt SW, et al. Was race a factor in the outcomes of the Women's Health Eating and Living Study? *Cancer* 2011; **117**(16):3805-13.

Pierce JP, Faerber S, Wright FA, Rock CL, Newman V, Flatt SW, et al. A randomized trial of the effect of a plant-based dietary pattern on additional breast cancer events and survival: the Women's Healthy Eating and Living (WHEL) Study. *Controlled Clinical Trials* 2002; **23**(6):728-56.

Pierce JP, Natarajan L, Caan BJ, Flatt SW, Kealey S, Gold EB, et al. Dietary change and reduced breast cancer events among women without hot flashes after treatment of early-stage breast cancer: subgroup analysis of the Women's Healthy Eating and Living Study. *American Journal of Clinical Nutrition* 2009; **89**(5):1565S-71S.

* Pierce JP, Natarajan L, Caan BJ, Parker BA, Greenberg ER, Flatt SW, et al. Influence of a diet very high in vegetables, fruit, and fiber and low in fat on prognosis following treatment for breast cancer: the Women's Healthy Eating and Living (WHEL) randomized trial. *JAMA* 2007; **298**(3):289-98.

Pierce JP, Natarajan L, Sun S, Al-Delaimy W, Flatt SW, Kealey S, et al. Increases in plasma carotenoid concentrations in response to a major dietary change in the Women's Healthy Eating and Living study. *Cancer Epidemiology, Biomarkers & Prevention* 2006; **15**(10):1886-92.

Pierce JP, Newman VA, Flatt SW, Faerber S, Rock CL, Natarajan L, et al. Telephone counseling intervention increases intakes of micronutrient- and phytochemical-rich vegetables, fruit and fiber in breast cancer survivors. *Journal of Nutrition* 2004; **134**(2):452-8.

Pierce JP. A randomized trial of the effect of a plant-based dietary pattern on additional breast cancer events and survival: the Women's Healthy Eating and Living (WHEL) Study. *Controlled Clinical Trials* 2002; **23**(6):728-56.



Pierce JP. Diet and breast cancer prognosis: making sense of the Women's Healthy Eating and Living and Women's Intervention Nutrition Study trials. *Current Opinion in Obstetrics & Gynecology* 2009; **21**(1):86-91.

Rock CL, Flatt SW, Laughlin GA, Gold EB, Thomson CA, Natarajan L, et al. Reproductive steroid hormones and recurrence-free survival in women with a history of breast cancer. *Cancer Epidemiology, Biomarkers & Prevention* 2008; **17**(3):614-20.

Rock CL, Flatt SW, Newman V, Caan BJ, Haan MN, Stefanick ML, et al. Factors associated with weight gain in women after diagnosis of breast cancer. Women's Healthy Eating and Living Study Group. *Journal of the American Dietetic Association* 1999; **99**(10):1212-21.

Rock CL, Flatt SW, Thomson CA, Stefanick ML, Newman VA, Jones L, et al. Plasma triacylglycerol and HDL cholesterol concentrations confirm self-reported changes in carbohydrate and fat intakes in women in a diet intervention trial. *Journal of Nutrition* 2004; **134**(2):342-7.

Rock CL, Natarajan L, Pu M, Thomson CA, Flatt SW, Caan BJ, et al. Longitudinal biological exposure to carotenoids is associated with breast cancer-free survival in the Women's Healthy Eating and Living Study. *Cancer Epidemiology, Biomarkers & Prevention* 2009; **18**(2):486-94.

Saquib N, Flatt SW, Natarajan L, Thomson CA, Bardwell WA, Caan B, et al. Weight gain and recovery of pre-cancer weight after breast cancer treatments: evidence from the Women's Healthy Eating and Living (WHEL) study. *Breast Cancer Research* & *Treatment* 2007; **105**(2):177-86.

Saxe GA, Madlensky L, Kealey S, Wu DP, Freeman KL, Pierce JP, et al. Disclosure to physicians of CAM use by breast cancer patients: findings from the Women's Healthy Eating and Living Study. *Integrative Cancer Therapies* 2008; **7**(3):122-9.

WHO primary prev 1979 {published data only}

WHO Primary prevention trial. Primary prevention of ischaemic heart disease: WHO coordinated cooperative trial. A summary report. *Bulletin Of The World Health Organization* 1979; **57**(5):801-5.

WHT 1990 {published and unpublished data}

Bowen D, Clifford CK, Coates R, Evans M, Feng Z, Fouad M, et al. The Women's Health Trial feasibility study in minority populations: design and baseline descriptions. *Annals of Epidemiology* 1996; **6**(6):507-19.

Bowen D. The role of participation in the Women's Health trial: feasibility study in minority populations. *Preventive Medicine* 2000; **31**(5):474-80.

Bowen DJ, Kestin M, McTiernan A, Carrell D, Green P. Effects of dietary fat intervention on mental health in women. *Cancer Epidemiology, Biomarkers and Prevention* 1995; **4**(5):555-9.

Gorbach SL, Morrill LA, Woods MN, Dwyer JT, Selles WD, Henderson M, et al. Changes in food patterns during a low-fat dietary intervention in women. *Journal of the American Dietetic Association* 1990; **90**(6):802-9.

Henderson MM, Kushi LH, Thompson DJ, Gorbach SL, Clifford CK, Insull W, et al. Feasibility of a randomized trial of a low-fat diet for the prevention of breast cancer: dietary compliance in the Women's Health Trial Vanguard Study. *Preventive Medicine* 1990; **19**(2):115-33.

* Insull W, Henderson MM, Prentice RL, Thompson DJ, Clifford C, Goldman S, et al. Results of a randomized feasibility study of a low-fat diet. *Archives of Internal Medicine* 1990; **150**(2):421-7.

Kristal AR, White E, Shattuck AL, Curry S, Anderson GL, Fowler A, et al. Long-term maintenance of a low-fat diet: durability of fatrelated dietary habits in the Women's Health Trial. *Journal of the American Dietetic Association* 1992; **92**(5):553-9.

Prentice RL, Kakar F, Hursting S, Sheppard L, Klein R, Kushi LH. Aspects of the rationale for the Women's Health Trial. *Journal of the National Cancer Institute* 1988; **80**(11):802-14.

Self S, Prentice R, Iverson D, Henderson M, Thompson D, Byar D, et al. Statistical design of the Women's Health Trial. *Controlled Clininical Trials* 1988; **9**(2):119-36.

Sheppard L, Kristal AR, Kushi LH. Weight loss in women participating in a randomised trial of low-fat diets. *American Journal of Clinical Nutrition* 1991; **54**(5):821-8.

Urban N, Baker M. The Women's Health Trial as an investment. *Medical Decision Making* 1989; **9**(1):59-64.

White E, Shattuck AL, Kristal AR, Urban N, Prentice RL, Henderson MM, et al. Maintenance of a low-fat diet: follow-up of the Women's Health Trial. *Cancer Epidemiology, Biomarkers and Prevention* 1992; **1**(4):315-23.

WHT Feasibility 2003 {published and unpublished data}

Bowen D, Clifford CK, Coates R, Evans M, Feng Z, Fouad M, et al. The Women's Health Trial Feasibility Study in Minority Populations: design and baseline descriptions. *Annals of Epidemiology* 1996; **6**(6):507-19.

* Hall WD, Feng Z, George VA, Lewis CE, Oberman A, Huber M, et al. Low-fat diet: effect on anthropometrics, blood pressure, glucose, and insulin in older women. *Ethnicity and Disease* 2003; **13**(3):337-43.

Wilke 1974 {*published data only*}

Wilke H, Frahm H. Influence of low-caloric-diet and dtriiodothyronine on serum lipids and body weight (author's trans) [Verhalten der serumlipide und des körpergewichts unter reduktionsdiät und medikamentöser behandlung mit dtrijodthyronin]. *Medizinische Klinik* 1974; **69**(48):1986-9.

Williams 1990 {published data only}

Williams PT, Krauss RM, Vranizan KM, Wood PS. Changes in lipoprotein subfractions during diet-induced and exerciseinduced weight loss in moderately overweight men. *Circulation* 1990; **81**(4):1293-304.



Williams 1992 {published data only}

Williams PT, Krauss RM, Vranizan KM, Albers JJ, Wood PD. Effects of weight-loss by exercise and by diet on apolipoproteins A-I and A-II and the particle-size distribution of high-density lipoproteins in men. *Metabolism: Clinical and Experimental* 1992; **41**(4):441-9.

Williams 1994 {published data only}

Williams PT, Stefanick ML, Vranizan KM, Wood PD. The effects of weight loss by exercise or by dieting on plasma high-density lipoprotein (HDL) levels in men with low, intermediate, and normal-to-high HDL at baseline. *Metabolism* 1994; **43**(7):917-24.

Wilmot 1952 {published data only}

Wilmot VA, Swank RL. The influence of low fat diet on blood lipid levels in health and in multiple sclerosis. *American Journal of the Medical Sciences* 1952; **223**(1):25-34.

Wing 1998 {published data only}

Wing RR, Venditti E, Jakicic JM, Polley BA, Lang W. Lifestyle intervention in overweight individuals with a family history of diabetes. *Diabetes Care* 1998; **21**(3):350-9.

WINS UK 2011 {published and unpublished data}

Parry BM, Milne JM, Yadegarfar G, Rainsbury RM. Dramatic dietary fat reduction is feasible for breast cancer patients: results of the randomised study, WINS (UK) - stage 1. *European Journal of Surgical Oncology* 2011; **37**(10):848-55.

WOMAN 2007 {published data only}

Kuller LH, Kriska AM, Kinzel LS, Simkin-Silverman LR, Sutton-Tyrrell K, Johnson BD, et al. The clinical trial of Women On the Move through Activity and Nutrition (WOMAN) study. *Contemporary Clinical Trials* 2007; **28**(4):370-81.

Wood 1988 {published data only}

Wood PD, Stefanick ML, Dreon DM, Frey HB, Garay SC, Williams PT, et al. Changes in plasma lipids and lipoproteins in overweight men during weight loss through dieting as compared with exercise. *New England Journal of Medicine* 1988; **319**(18):1173-9.

Woollard 2003 {published data only}

Woollard J, Burke V, Beilin LJ, Verheijden M, Bulsara MK. Effects of a general practice-based intervention on diet, body mass index and blood lipids in patients at cardiovascular risk. *Journal* of Cardiovascular Risk 2003; **10**(1):31-40.

Working Well 1996 {published data only}

Sorensen G, Thompson B, Glanz K, Feng Z, Kinne S, DiClemente C, et al. Work site-based cancer prevention: primary results from the Working Well Trial. *American Journal of Public Health* 1996; **86**(7):939-47.

Zock 1995 {published and unpublished data}

Zock PL, Mensink RP, Harryvan J, De Vries JH, Katan MB. Fatty acids in serum cholesteryl esters as quantitative biomarkers of dietary intake in humans. *American Journal of Epidemiology* 1997; **145**(12):1114-22.

* Zock PL. Dietary Fatty Acids and Risk Factors for Coronary Heart Disease: Controlled Studies in Healthy Volunteers [PhD thesis]. Wageningen, Netherlands: Landbouwuniversiteit te Wageningen, 1995.

References to studies awaiting assessment

ICFAMED {published data only}ISRCTN27497769

Lapetra J, Lozano-Rodríguez JM, Caballero-Valderrama MR, Santos-Lozano JM, Ortega-Calvo M, García de la Corte FJ, et al. Effect of a Mediterranean diet on the primary prevention of atrial fibrillation and major cardiovascular events in hypertensive patients with high cardiovascular risk: results of ICFAMED study. In: Obesity and Nutrition in the 21st Century. VIII Symposium Ciber Fisiopatología de la Obesidad y Nutrición; 2017; Madrid. Madrid: Ciber Fisiopatología de la Obesidad y Nutrición, 2017:71.

* Lapetra J, Lozano-Rodriguez JM, Miro-Moriano L, Ortega-Calvo M, Santos-Lozano JM, Garcia-Corte FJ, et al. Effect of a Mediterranean diet on the primary prevention of atrial fibrillation and major cardiovascular events in hypertensive patients with high cardiovascular risk: results of ICFAMED randomized trial. *European Journal of Clinical Investigation* 2018; **48**:182-3.

References to ongoing studies

ENAbLE due unclear {published data only}

English C, Patterson A, MacDonald-Wicks L, Attia J, Callister R, Hillier S, et al. ENAbLE: Secondary prevention of stroke. A physical activity and diet trial protocol. Abstracts Presented at the SMART STROKES 2019 Conference, 8–9 August 2019, Hunter Valley, NSW. *International Journal of Stroke* 2019; **14**(1 Suppl):12. [DOI: 10.1177/1747493019858233]

NCT02481466 due 2020 {published data only}

. The combined portfolio diet and exercise study (PortfolioEx). clinicaltrials.gov/ct2/show/NCT02481466 (first posted 25 June 2015).

NCT02938832 due 2023 {published data only}

. Does the advice to eat a Mediterranean diet with low carbohydrate intake, compared with a low-fat diet, reduce diabetes and cardiovascular disease? clinicaltrials.gov/ct2/ show/NCT02938832 (first posted 19 October 2016).

NEW Soul Study due 2022 {published data only} ZZZ <accessionId ref="info:x-wiley/clinicalTrialsGov/ NCT03354377"> ZZZNCT03354377

. Nutritious Eating With Soul (The NEW Soul Study). https:// clinicaltrials.gov/ct2/show/NCT03354377 (first posted 27 November 2017).

* Turner-McGrievy G, Wilcox S, Frongillo EA, Murphy A, Hutto B, Williams K, et al. The Nutritious Eating with Soul (NEW Soul) Study: Study design and methods of a two-year randomized trial comparing culturally adapted soul food vegan vs. omnivorous diets among African American adults at risk for heart disease. *Contemporary Clinical Trials* 2020; **88**:105897.



Additional references

4S 1994

Scandinavian Simvastatin Survival Study Group. Randomised trial of cholesterol lowering in 4444 patients with coronary heart disease: the Scandinavian Simvastatin Survival Study (4S). *Lancet* 1994; **344**(8934):1383-9.

Abdelhamid 2018

Abdelhamid AS, Martin N, Bridges C, Brainard JS, Wang X, Brown TJ, et al. Polyunsaturated fatty acids for the primary and secondary prevention of cardiovascular disease. *Cochrane Database of Systematic Reviews* 2018, Issue 11. [DOI: 10.1002/14651858.CD012345.pub2] [CD012345]

Abdelhamid 2019

Abdelhamid AS, Hooper L, Sivakaran R, Hayhoe R, Welch AA, PUFAH Group. The relationship between omega-3, omega-6 and total polyunsaturated fat and musculoskeletal health and functional status in adults: a systematic review and meta-analysis of RCTs. *Calcified Tissue International* 2019; **105**(4):353-72. [DOI: 10.1007/s00223-019-00584-3]

Abdelhamid 2020

Abdelhamid AS, Brown TJ, Brainard JS, Biswas P, Thorpe GC, Moore HJ, Deane KHO, Summerbell CD, Worthington HV, Song F, Hooper L. Omega-3 fatty acids for the primary and secondary prevention of cardiovascular disease. *Cochrane Database of Systematic Reviews* 2020, Issue 3. [DOI: 10.1002/14651858.CD003177.pub5] [CD003177]

Alonso 2006

, Ruiz-Gutierrez V, . Monounsaturated fatty acids, olive oil and blood pressure: epidemiological, clinical and experimental evidence. *Public Health Nutrition* 2006; **9**(2):251-7.

Astrup 2000

Astrup A, Ryan L, Grunwald GK, Storgaard M, Saris W, Melanson E, et al. The role of dietary fat in body fatness: evidence from a preliminary meta-analysis of ad libitum lowfat dietary intervention studies. *British Journal of Nutrition* 2000; **83**(Suppl 1):S25-32.

Astrup 2011

Astrup A, Dyerberg J, Elwood P, Hermansen K, Hu FB, Jakobsen MU, et al. The role of reducing intakes of saturated fat in the prevention of cardiovascular disease: where does the evidence stand in 2010? *American Journal of Clinical Nutrition* 2011; **93**(4):684–8.

Berkley 1995

Berkley CS, Hoaglin DC, Mosteller F, Colditz GA. A randomeffects regression model for meta-analysis. *Statistics in Medicine* 1995; **14**(4):395-411.

Blann 1995

Blann AD, Jackson P, Bath PM, Watts GF. Von Willebrand factor, a possible indicator of endothelial cell damage, decreases during long-term compliance with a lipid-lowering diet. *Journal of Internal Medicine* 1995; **237**(6):557-61.

Brainard 2020

Brainard J, Jimoh OF, Deane K, Biswas P, Donaldson D, Maas K, et al. Omega-3, omega-6, and polyunsaturated fat for cognition: systematic review and meta-analysis of randomized trials. Journal of the American Medical Directors Association 2020; **15 April 2020 [Epub ahead of print]**. [DOI: 10.1016/ j.jamda.2020.02.022]

Briel 2009

Briel M, Ferreira-Gonzalez I, You JJ, Karanicolas PJ, Akl EA, Wu P, et al. Association between change in high density lipoprotein cholesterol and cardiovascular disease morbidity and mortality: systematic review and meta-regression analysis. *BMJ* 2009; **338**:b92. [DOI: 10.1136/bmj.b92]

Brown 2019

Brown TJ, Brainard J, Song F, Wang X, Abdelhamid A, Hooper L. Omega-3, omega-6, and total dietary polyunsaturated fat for prevention and treatment of type 2 diabetes mellitus: systematic review and meta-analysis of randomised controlled trials. *BMJ* 2019; **366**:14697.

Brunner 1997

Brunner E, White I, Thorogood M, Bristow A, Curle D, Marmot M. Can dietary interventions change diet and cardiovascular risk factors? A meta-analysis of randomized controlled trials. *American Journal of Public Health* 1997; **87**(9):1415-22.

Bucher 1996

Bucher HC, Cook RJ, Guyatt GH, Lang JD, Cook DJ, Hatala R, et al. Effects of dietary calcium supplementation on blood pressure: a meta-analysis of randomised controlled trials. *JAMA* 1996; **275**(13):1016-22.

Chalmers 1990

Chalmers I, Adams M, Dickersin K, Hetherington J, Tarnow-Mordi W, Meinert C, et al. A cohort study of summary reports of controlled trials. *JAMA* 1990; **263**(10):1401-5.

Clarke 1997

Clarke R, Frost C, Collins R, Appleby P, Peto R. Dietary lipids and blood cholesterol: quantitative meta-analysis of metabolic ward studies. *BMJ* 1997; **314**(7074):112-7.

Davey Smith 1993

Davey Smith G, Song F, Sheldon TA. Cholesterol lowering and mortality: the importance of considering initial level of risk. *BMJ* 1993; **306**(6889):1367-73.

Dayton 1965

Dayton S, Hashimoto S, Rosenblum D, Pearce M. Vitamin E status of humans during prolonged feeding of unsaturated fats. *Journal of Laboratory and Clinical Medicine* 1965; **65**(5):739-47.

De Caterina 2010

De Caterina R, Scarano M, Marfisi RM, Lucisano G, Palma F, Tatasciore A, et al. Cholesterol-lowering interventions and stroke. *Journal of the American College of Cardiology* 2010; **55**(3):198-211.



Deane 2019

Deane KHO, Jimoh OF, Biswas P, O'Brien A, Hanson S, Abdelhamid AS, et al. Omega-3 and polyunsaturated fat for prevention of depression and anxiety symptoms: systematic review and meta-analysis of randomised trials. British Journal of Psychiatry 2019 Oct 24 [E-pub ahead of print]. [DOI: 10.1192/ bjp.2019.234]

Denke 1995

Denke MA. Cholesterol-lowering diets. A review of the evidence. *Archives of Internal Medicine* 1995; **155**(1):17-26.

Ebrahim 1997

Ebrahim S, Davey Smith G. Systematic review of randomised controlled trials of multiple risk factor interventions for preventing coronary heart disease. *BMJ* 1997; **314**(7095):1666.

Eckel 2013

Eckel RH, Jakicic JM, Ard JD, De Jesus JM, Houston Miller N, Hubbard VS, et al. 2013 AHA/ACC guideline on lifestyle management to reduce cardiovascular risk: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Circulation* 2013; **129**(25 suppl 2):S76-99. [DOI: 10.1161/01.cir.0000437740.48606.d1]

Egger 1997

Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple graphical test. *BMJ* 1997; **315**(7109):629-34.

ESC/EAS 2011

Task Force for the management of dyslipidaemias of the European Society of Cardiology (ESC) and the European Atherosclerosis Society (EAS). ESC/EAS Guidelines for the management of dyslipidaemias. *European Heart Journal* 2011; **32**(14):1769-1818. [DOI: 10.1093/eurheartj/ehr158]

Gofman 1949

Gofman JW, Lindgren FT, Elliott HA. Ultracentrifugal studies of lipoproteins of human serum. *Journal of Biological Chemistry* 1949; **179**(2):973-9.

GRADE 2004

Atkins D, Best D, Briss PA, Eccles M, Falck-Ytter Y, Flottorp S, et al. Grading quality of evidence and strength of recommendations. *BMJ* 2004; **328**(7454):1490.

GRADEpro 2015 [Computer program]

McMaster University and Evidence Prime Inc GRADEpro GDT. https://gradepro.org/: McMaster University and Evidence Prime Inc, 2015.

Grundy 2019

Grundy SM, Stone NJ, Bailey AL, Beam C, Birtcher KK, Blumenthal RS, et al. 2018 AHA/ACC/AACVPR/AAPA/ABC/ ACPM/ADA/AGS/APhA/ASPC/NLA/PCNA Guideline on the Management of Blood Cholesterol. *Journal of the American College of Cardiology* 2019; **73**(24):e285-350.

Hanson 2020

Hanson S, Thorpe G, Winstanley L, Abdelhamid AS, Hooper L, PUFAH Group. Omega-3, omega-6 and total dietary polyunsaturated fat on cancer incidence: systematic review and meta-analysis of randomised trials. British Journal of Cancer 2020; **122**:1260-70. [DOI: 10.1038/s41416-020-0761-6]

Harcombe 2015

Harcombe Z, Baker JS, Cooper SM, Davies B, Sculthorpe N, DiNicolantonio JJ, et al. Evidence from randomised controlled trials did not support the introduction of dietary fat guidelines in 1977and 1983: a systematic review and meta-analysis. *Open Heart* 2015; **2**(1):e000196. [DOI: 10.1136/openhrt-2014-000196]

Hauck 1991

Hauck WW, Gilliss CL, Donner A, Gortner S. Randomisation by cluster. *Nursing Research* 1991; **40**(6):356-8.

Hegsted 1965

Hegsted DM, McGandy RB, Myers ML, Stare FJ. Quantitative effects of dietary fat on serum cholesterol in man. *American Journal of Clinical Nutrition* 1965; **17**(5):281-95.

Hegsted 2000

Hegsted DM. From chick nutrition to nutrition policy. *Annual Review of Nutrition* 2000; **20**:1-19.

Hession 2009

Hession M, Rolland C, Kulkarni U, Wise A, Broom J. Systematic review of randomized controlled trials of low-carbohydrate vs. low-fat/low-calorie diets in the management of obesity and its comorbidities. *Obesity Reviews* 2009; **10**(1):36-50.

Higgins 2003

Higgins JPT, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ* 2003; **327**(7414):557-60.

Higgins 2011

Higgins JPT, . Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. Available from handbook.cochrane.org.

Hooper 2015b

Hooper L, Abdelhamid A, Bunn D, Brown T, Summerbell CD, Skeaff CM. Effects of total fat intake on body weight. *Cochrane Database of Systematic Reviews* 2015, Issue 8. [DOI: 10.1002/14651858.CD011834]

Hooper 2018

Hooper L, Al-Khudairy L, Abdelhamid AS, Rees K, Brainard JS, Brown TJ, et al. Omega-6 fats for the primary and secondary prevention of cardiovascular disease. *Cochrane Database of Systematic Reviews* 2018, Issue 11. [DOI: 10.1002/14651858.CD011094.pub3] [CD011094]

Hooper 2019

Hooper L, Abdelhamid A, Brainard J, Deane KHO, Song F. Creation of a database to assess effects of omega-3, omega-6 and total polyunsaturated fats on health: database and methodology for a set of reviews. *BMJ Open* 2019; **9**(5):e029554. [DOI: 10.1136/bmjopen-2019-029554]



Howard 2010

Howard BV, Curb JD, Eaton CB, Kooperberg C, Ockene J, Kostis JB, et al. Low-fat dietary pattern and lipoprotein risk factors: the Women's Health Initiative dietary modification trial. *American Journal of Clinical Nutrition* 2010; **91**(4):860-74.

Kagan 1974

Kagan A, Harris BR, , Johnson KG, Kato H, Syme SL, et al. Epidemiologic studies of coronary heart disease and stroke in Japanese men living in Japan, Hawaii and California: demographic, physical, dietary and biochemical characteristics. *Journal of Chronic Disease* 1974; **27**(7-8):345-64.

Kasim 1993

Kasim SE, Martino S, , Khilnani S, Boomer A, Depper J, et al. Dietary and anthropometric determinants of plasma lipoproteins during a long-term low-fat diet in healthy women. *American Journal of Clinical Nutrition* 1993; **57**(2):146-53.

Katan 2010

Katan MB, Brouwer IA, Clarke R, Geleijnse JM, Mensink RP. Saturated fat and heart disease. *American Journal of Clinical Nutrition* 2010; **92**(2):459-60; author reply 460-1.

Keys 1950

Keys A, Mickelsen O, Miller EV, Carleton B. The relation in man between cholesterol levels in the diet and in the blood. *Science* 1950; **112**(2899):79-81.

Keys 1986

Keys A, Menotti A, Karvonen MJ, Aravanis C, Blackburn H, Buzina RM, et al. The diet and 15-year death rate in the seven countries study. *American Journal of Epidemiology* 1986; **124**(6):903-15.

Kodama 2009

Kodama S, Saito K, Tanaka S, Maki M, Yachi Y, Sato M, et al. Influence of fat and carbohydrate proportions on the metabolic profile in patients with type 2 diabetes: a meta-analysis. *Diabetes Care* 2009; **32**(5):959-65.

Kristal 2005

Kristal AR, Peters U, Potter JD. Is it time to abandon the food frequency questionnaire? *Cancer Epidemiology, Biomarkers and Prevention* 2005; **14**(12):2826-8.

Law 1991

Law MR, Frost CD, Wald NJ. By how much does dietary salt restriction lower blood pressure? *BMJ* 1991; **302**(6780):811-24.

Law 1994

Law MR, Wald NJ, Thompson SG. By how much and how quickly does reduction in serum cholesterol concentration lower risk of ischaemic heart disease? *BMJ* 1994; **308**(6925):367-73.

Lefebvre 2011

Lefebvre C, Manheimer E, Glanville J. Chapter 6: Searching for studies. In: Higgins JPT, Green S (editors). Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. Available from handbook.cochrane.org.

Reduction in saturated fat intake for cardiovascular disease (Review) Copyright © 2020 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

Lenz 2007

Lenz M, Steckelberg A, Richter B, Mühlhauser I. Meta-analysis does not allow appraisal of complex interventions in diabetes and hypertension self-management: a methodological review. *Diabetologia* 2007; **50**(7):1375–83.

Lichtenstein 2006

Lichtenstein AH, Appel LJ, Brands M, Carnethon M, Daniels S, Franch HA, et al. Diet and lifestyle recommendations revision 2006: a scientific statement from the American Heart Association Nutrition Committee. *Circulation* 2006; **114**(1):82-96.

Lopez-Espinoza 1984

Lopez-Espinoza I, Howard WJ, Mann JI, Carter RD, Hockaday TD. Fatty acid composition of platelet phospholipids in non-insulindependent diabetics randomized for dietary advice. *British Journal of Nutrition* 1984; **52**(1):41-7.

Mach 2019

Mach F, Baigent C, Catapano AL, Koskinas KC, Casula M, Badimon L, et al. 2019 ESC/EAS guidelines for the management of dyslipidaemias: lipid modification to reduce cardiovascular risk. *Atherosclerosis* 2019; **290**:140-205.

Malhotra 2014

Malhotra A, Shafiq N, Arora A, Singh M, Kumar R, Malhotra S. Dietary interventions (plant sterols, stanols, omega-3 fatty acids, soy protein and dietary fibers) for familial hypercholesterolaemia. *Cochrane Database of Systematic Reviews* 2014, Issue 6. [DOI: 10.1002/14651858.CD001918.pub3]

Marchioli 1994

Marchioli R, Prieto JC, Tognoni G. Surrogate end-points: the case of trials on coronary atherosclerotic plaque regression. *Clinical Trials and Meta-Analyses* 1994; **29**(2-3):139-76.

Mensink 1992

Mensink RP, Katan MB. Effect of dietary fatty acids on serum lipids and lipoproteins: a meta-analysis of 27 trials. *Arteriosclerosis and Thrombosis* 1992; **12**(8):911-9.

Mensink 2003

Mensink RP, Zock PL, Kester ADM, Katan MB. Effects of dietary fatty acids and carbohydrates on the ratio of serum total to HDL cholesterol and on serum lipids and apolipoproteins: a metaanalysis of 60 controlled trials. *American Journal of Clinical Nutrition* 2003; **77**(5):1146-55.

Michels 2009

Michels KB, Willett WC. The Women's Health Initiative randomized controlled dietary modification trial: a post-mortem. *Breast Cancer Research and Treatment* 2009; **114**(1):1-6.

Mozaffarian 2010

Mozaffarian D, Micha R, Wallace S. Effects of coronary heart disease of increasing polyunsaturated fat in place of saturated fat: a systematic review and meta-analysis of randomized controlled trials. *PLOS Medicine* 2010; **7**(3):e1000252.



Ness 1997

Ness AR, Powles JW. Fruit and vegetables, and cardiovascular disease: a review. *International Journal of Epidemiology* 1997; **26**(1):1-13.

NHS 2020

NHS. How to eat less saturated fat. www.nhs.uk/live-well/eat-well/eat-less-saturated-fat/ (accessed 11 Feb 2020).

NICE 2014

National Institute of Health and Care Excellence. Lipid modification: cardiovascular risk assessment and the modification of blood lipids for the primary and secondary prevention of cardiovascular disease. www.nice.org.uk/ guidance/cg181/resources/guidance-lipid-modificationcardiovascular-risk-assessment-and-the-modification-ofblood-lipids-for-the-primary-and-secondary-prevention-ofcardiovascular-disease-pdf (accessed 29th Dec 2019).

O'Gara 2014

O'Gara PT, Kushner FG, Ascheim DD, , Chung MK, De Lemos JA, et al. 2013 ACCF/AHA guideline for the management of STelevation myocardial infarction: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *Circulation* 2013; **127**(4):e362-e425. [DOI: 10.1161/CIR.0b013e3182742cf6]

Oliver 1953

Oliver MF, Boyd GS. The plasma lipids in coronary artery disease. *British Heart Journal* 1953; **15**(4):387-90.

Oliver 2000

Oliver MF. Pioneer research in Britain into atherosclerosis and coronary heart disease - an historical review. *Atherosclerosis* 2000; **150**(1):1-12.

Page 2019

Page MJ, Higgins JPT, Sterne JAC. Chapter 13: Assessing risk of bias due to missing results in a synthesis. In: Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, Welch VA (editors). Cochrane Handbook for Systematic Reviews of Interventions Version 6.0 (updated July 2019). Cochrane 2019. Available from www.training.cochrane.org/handbook.

Patterson 2003

Patterson RE, Kristal A, Rodabough R, Caan B, Lillington L, Mossavar-Rahmani Y, et al. Changes in food sources of dietary fat in response to an intensive low-fat dietary intervention: early results from the Women's Health Initiative. *Journal of the American Dietetic Association* 2003; **103**(4):454-60.

Prentice 2007

Prentice RL. Observational studies, clinical trials, and the Women's Health Initiative. *Lifetime Data Analysis* 2007; **13**(4):449-62.

Rees 2013

Rees K, Dyakova M, Wilson N, Ward K, Thorogood M, Brunner E. Dietary advice for reducing cardiovascular risk. *Cochrane Database of Systematic Reviews* 2013, Issue 12. [DOI: 10.1002/14651858.CD002128.pub5]

Reduction in saturated fat intake for cardiovascular disease (Review) Copyright © 2020 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

Rimm 1996

Rimm EB, Klatsky A, Grobbee D, Stampfer MJ. Review of moderate alcohol consumption and reduced risk of coronary heart disease: is the effect due to beer, wine or spirits? *BMJ* 1996; **312**(7033):731-6.

Robertson 1977

Robertson TL, Kato H, Rhoads GG, Kagan A, Marmot M, Syme SL, et al. Epidemiologic studies of coronary heart disease and stroke in Japanese men living in Japan, Hawaii and California. Incidence of myocardial infarction and death from coronary heart disease. *American Journal of Cardiology* 1977; **39**(2):239-43.

Robinson 2009

Robinson JG, Wang S, Smith BJ, Jacobson TA. Meta-analysis of the relationship between non-high-density lipoprotein cholesterol reduction and coronary heart disease risk. *Journal of the American College of Cardiology* 2009; **53**(4):316-22.

Rubins 1995

Rubins HB. Cholesterol in patients with coronary heart disease: how low should we go? *Journal of General Internal Medicine* 1995; **10**(8):464-71.

Ryle 1949

Ryle JA, Russell WT. The natural history of coronary disease. A clinical and epidemiological study. *British Heart Journal* 1949; **11**(4):370.

SACN 2019

Scientific Advisory Committee on Nutrition. Saturated fat and health. www.gov.uk/government/publications/saturated-fats-and-health-sacn-report (accessed prior to 30 April 2020).

Savovic 2012

Savovic J, Jones H, Altman D, Harris R, Jüni P, Pildal J, et al. Influence of reported study design characteristics on intervention effect estimates from randomised controlled trials: combined analysis of meta-epidemiological studies. *Health Technology Assessment* 2012; **16**:1-82.

Scarborough 2010

Scarborough P, Rayner M, Van Dis I, Norum K. Meta-analysis of effect of saturated fat intake on cardiovascular disease: overadjustment obscures true associations. *American Journal of Clinical Nutrition* 2010; **92**(2):458-9; author reply 459.

Schatzkin 2003

Schatzkin A, Kipnis V, Carroll RJ, Midthune D, Subar AF, Bingham S, et al. A comparison of a food frequency questionnaire with a 24-hour recall for use in an epidemiological cohort study: results from the biomarkerbased Observing Protein and Energy Nutrition (OPEN) Study. *International Journal of Epidemiology* 2003; **32**(6):1054-62.

Schulz 1995

Schulz KF, Chalmers I, Hayes RJ, Altman DG. Empirical evidence of bias. Dimensions of methodological quality associated with estimates of treatment effects in controlled trials. *JAMA* 1995; **273**(5):408-12.



Schwab 2014

Schwab U, Lauritzen L, Tholstrup T, Haldorssoni T, Riserus U, Uusitupa M, et al. Effect of the amount and type of dietary fat on cardiometabolic risk factors and risk of developing type-2 diabetes, cardiovascular disease, and cancer: a systematic review. Food & Nutrition Research 2014; **58**:25145. [DOI: 10.3402/fnr.v58.25145]

Shah 2007

Shah M, Adams-Huet B, Garg A. Effect of high-carbohydrate or high-cis-monounsaturated fat diets on blood pressure: a meta-analysis of intervention trials. *American Journal of Clinical Nutrition* 2007; **85**(5):1251-6.

Sharp 1998

Sharp S. Meta-analysis regression. *Stata Technical Bulletin* 1998; **42**:16-22.

Sheppherd 2009

Shepperd S, Lewin S, Straus S, Clarke M, Eccles MP, Fitzpatrick R, et al. Can we systematically review studies that evaluate complex interventions? *PLOS Medicine* 2009; **6**(8):e1000086.

SIGN 1996

Scottish Intercollegiate Guidelines Network. Management of obesity (national guideline). https://www.sign.ac.uk/sign-115-management-of-obesity (accessed prior to 30 April 2020);**115**.

Siri-Tarino 2010

Siri-Tarino PW, Sun Q, Hu FB, Krauss RM. Meta-analysis of prospective cohort studies evaluating the association of saturated fat with cardiovascular disease. *American Journal of Clinical Nutrition* 2010; **91**(3):535-46.

Skeaff 2009

Skeaff CM, Miller J. Dietary fat and coronary heart disease: summary of evidence from prospective cohort and randomised controlled trials. *Annals of Nutrition and Metabolism* 2009; **55**(1-3):173-201.

Stamler 2010

Stamler J. Diet-heart: a problematic revisit. *American Journal of Clinical Nutrition* 2010; **91**(3):497-9.

Stein 2006

Stein K. After the media feeding frenzy: whither the Women's Health Initiative Dietary Modification Trial? *Journal of the American Dietetic Association* 2006; **106**(6):794-800.

Sterne 2001

Sterne JA, Bradburn MJ, Egger M. Meta-analysis in STATA. In: Egger M, Davey Smith G, Altman DG, editors(s). Systematic Reviews in Health Care: Meta-analysis in Context. London: BMJ Books, 2001.

Sterne 2009

Sterne JA. Meta-Analysis in Stata: an Updated Collection from the Stata Journal. Texas, USA: STATA Press, 2009.

Thorogood M. Nutrition. In: Lawrence M, Neil A, Mant D, Fowler G, editors(s). Prevention of Cardiovascular Disease: an Evidence-Based Approach. Kings Lynn: Oxford University Press, 1996:54-66.

Walsh 1995

Walsh JME, Grady D. Treatment of hyperlipidaemia in women. *JAMA* 1995; **274**(14):1152-8.

Weggemans 2001

Weggemans RM, Zock PL, Katan MB. Dietary cholesterol from eggs increases the ratio of total cholesterol to high-density lipoprotein cholesterol in humans: a meta-analysis. *American Journal of Clinical Nutrition* 2001; **73**(5):885-91.

Wood 2008

Wood L, Egger M, Gluud LL, Schulz K, Jüni P, Altman DG, et al. Empirical evidence of bias in treatment effect estimates in controlled trials with different interventions and outcomes: meta-epidemiological study. *British Medical Journal* 2008; **336**:601-5.

Yngve 2006

Yngve A, Hambraeus L, Lissner L, Serra Majem L, Vaz de Almeida MD, Berg C, et al. Invited Commentary. The Women's Health Initiative. What is on trial: nutrition and chronic disease? Or misinterpreted science, media havoc and the sound of silence from peers. *Public Health Nutrition* 2006; **9**(2):269-72.

Yu-Poth 1999

Yu-Poth S, Zhao G, Etherton T, Naglak M, Jonnalagadda S, Kris-Etherton PM. Effects of the National Cholesterol Education Program's Step I and Step II dietary intervention programs on cardiovascular disease risk factors: a meta-analysis. *American Journal of Clinical Nutrition* 1999; **69**(4):632-46.

References to other published versions of this review

Hooper 2000

Hooper L, Summerbell CD, Higgins JPT, Thompson RL, Clements G, Capps N, et al. Reduced or modified dietary fat for preventing of cardiovascular disease. *Cochrane Database of Systematic Reviews* 2000, Issue 2. [DOI: 10.1002/14651858.CD002137]

Hooper 2001

Hooper L, Summerbell CD, Higgins JPT, Thompson RL, Capps N, Davey Smith G, et al. Dietary fat intake and prevention of cardiovascular disease: systematic review. *BMJ* 2001; **322**(7289):757-63.

Hooper 2011

Hooper L, Summerbell CD, Thompson R, Sills D, Roberts FG, Moore HJ, et al. Reduced or modified dietary fat for preventing cardiovascular disease. *Cochrane Database of Systematic Reviews* 2011, Issue 7. [DOI: 10.1002/14651858.CD002137.pub2]



Hooper 2012

Hooper L, Summerbell CD, Thompson R, Sills D, Roberts FG, Moore HJ, et al. Reduced or modified dietary fat for preventing cardiovascular disease. *Cochrane Database of Systematic Reviews* 2012, Issue 5. [DOI: 10.1002/14651858.CD002137.pub3]

Hooper 2015a

Hooper L, Martin N, Jimoh OF, Abdelhamid AS. Reduction in saturated fat intake for cardiovascular disease. *Cochrane Database of Systematic Reviews* 2015, Issue 6. [DOI: 10.1002/14651858.CD011737]

* Indicates the major publication for the study

CHARACTERISTICS OF STUDIES

Characteristics of included studies [ordered by study ID]

Black 1994

Study characteristics	
Methods	RCT
	Summary risk of bias: moderate to high
Participants	People with non-melanoma skin cancer (USA) CVD risk: low Control: randomised 67, analysed 58 Intervention: randomised 66, analysed 57 Mean years in trial: 1.9 % male: control 67%, intervention 54% Age: mean control 52.3 (SD 13.2), intervention 50.6 (SD 9.7)
	Ethnicity: white 100% (excluded from study if of Asian, Black, Hispanic or American Indian ancestry)
	Statins use allowed: Unclear
	% taking statins: Not reported
Interventions	Reduced fat vs usual diet
	Control aims: no dietary advice Intervention aims: total fat 20%E, protein 15%E, CHO 65%E
	Control methods: no dietary change, 4-month intervals clinic examination by dermatologist
	Intervention methods: 8 x weekly classes plus monthly follow-up sessions, with behavioural techniques being taught following individual approach (not clear if in a group or individual). 4-month intervals clin- ic examination by dermatologist
	Intervention delivered face-to-face by a dietitian
	Total fat intake, %E ("during study" months 4 - 24): cont 37.8 (SD 4.1), int 20.7 (SD 5.5) (mean difference -17.10, 95% CI -18.88 to -15.32) significant reduction
	Saturated fat intake, %E ("during study", months 4 - 24): cont 12.8 (SD 2.0), int 6.6 (SD 1.8), (mean difference -6.20, 95% CI -6.90 to -5.50) significant reduction
	PUFA intake, %E ("during study", months 4 - 24): cont 7.8 (SD 1.4), int 4.5 (SD 1.3), (mean difference -3.30, 95% CI -3.79 to -2.81) significant reduction
	PUFA n-3 intake: not reported
	PUFA n-6 intake: Linoleic acid, cont 16.9 (SD 5.6) g, int 8.5 (SD 3.3) g
	MUFA intake, %E ("during study", months 4 - 24): cont 14.4 (SD 1.7), int 7.6 (SD 2.2), (mean difference -6.80, 95% CI -7.52 to -6.08) significant reduction
	Cochrane
----	----------
マノ	Library

Black 1994 (Continued)	
	CHO intake, %E ("during study", months 4 - 24): cont 44.6 (SD 6.9), int 60.3 (SD 6.3), (mean difference 15.70, 95% Cl 13.29 to 18.11) significant increase
	Protein intake, %E ("during study", months 4 - 24): cont 15.7 (SD 2.4), int 17.7 (SD 2.2), (mean difference 2.00, 95% CI 1.16 to 2.84) significant increase
	Trans fat intake: not reported
	Replacement for saturated fat: CHO and protein (by dietary aims and achievements), main is CHO
	Style: diet advice
	Setting: community
Outcomes	Stated trial outcomes: incidence of actinic keratosis and non-melanoma skin cancer Data available on total mortality? yes Cardiovascular mortality? yes Events available for combined cardiovascular events: cardiovascular deaths
	Secondary outcomes: cancer deaths (none)
	Tertiary outcomes: none (weight data provided, but no variance info)
Notes	Study duration 24 months.
	Study aim was to achieve low-fat diet, but the study achieved a statistically significant reduction in sat- urated fat intake in the low-fat group compared to control.
	SFA reduction achieved.
	Total serum cholesterol: not reported
	At 2 years control -1.5 kg n = 50?, intervention -1 kg n = 51?
	Trial dates: Study dates not reported (but still recruiting at first publication in 1994)
	Funding: National Cancer Institute
	Declarations of Interest of primary researchers: none stated, all authors worked for academic or health institutions
Risk of bias	

Bias	Authors' judgement	Support for judgement
Random sequence genera- tion (selection bias)	Low risk	"list of randomly generated numbers"
Allocation concealment (selection bias)	Unclear risk	Allocation method not clearly described
Blinding of participants and personnel (perfor- mance bias) All outcomes	High risk	Dietary advice provided, so participants not blinded
Blinding of outcome as- sessment (detection bias) CVD outcomes	Low risk	"examined by dermatologists unaware of their treatment assignments". Deaths (all-cause and CVD) not considered relevant to the intervention
Blinding of outcome as- sessment (detection bias)	Low risk	Blinding is not relevant in assessment of mortality

Reduction in saturated fat intake for cardiovascular disease (Review)

Copyright @ 2020 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.



Black 1994 (Continued) All-cause mortality

Incomplete outcome data (attrition bias) All outcomes	Low risk	Low risk for all-cause and CVD mortality. Unclear for other outcomes
Selective reporting (re- porting bias)	Low risk	Not relevant for primary and secondary outcomes as all trialists asked for data
Free of systematic differ- ence in care?	High risk	Minor, all have 4-monthly clinic visits, the intervention group had 8 behaviour- al technique classes that the control group did not have
Stated aim to reduce SFA	High risk	Aim to reduce SFA not stated
Achieved SFA reduction	Low risk	Statistically significant SFA reduction achieved
Achieved TC reduction	Unclear risk	Not reported
Other bias	Low risk	None noted

DART 1989

Study characteristics	
Methods	Factorial RCT
	Diet And Reinfarction Trial (DART)
	Summary risk of bias: moderate to high
Participants	 Men recovering from an MI (UK) CVD risk: high Control: randomised 1015, analysed unclear Intervention: randomised 1018, analysed unclear Mean years in trial: control 1.9, randomised 1.9 % male: 100% Age: mean control 56.8, intervention 56.4 (< 70) Ethnicity: not stated Statins use allowed? Unclear, but there do not appear to have been any medication-based exclusion criteria and included participants were taking anti-hypertensives, anti-anginals, anti-coagulants, anti-platelet, digoxin and "other cardiac drugs". % taking statins: Not reported, but only 5.4% were taking "other cardiac drugs" which may have include ed statins.
Interventions	Reduced and modified fat vs usual diet Control aims: no dietary advice on fat, weight reducing advice if BMI > 30 Intervention aims: reduce fat intake to 30%E, increase P/S to 1.0, weight-reducing advice if BMI > 30 Note: This was a factorial trial, and so some in each group were randomised to increased fatty fish and/ or increased cereal fibre. Control methods: dietitians provided 'sensible eating' advice without specific information on fats.

Intervention methods: dietitians provided the participants and their wives with initial individual advice



DART 1989 (Continued)

Trusted evidence. Informed decisions. Better health.

	and a diet information sheet; participants were revisited for further advice, recipes, encouragement at 1, 3, 6, 9, 12, 15, 18 and 21 months.
	Intervention delivered individually face-to-face by a dietitian
	Total fat intake, %E (through study): cont 35 (SD 6), int 31 (SD 7) (mean difference -4.00, 95% CI -4.57 to -3.43) significant reduction
	Saturated fat intake, %E (through study): cont 15 (SD3), int 11 (SD3), (mean difference -4.00, 95% CI -4.26 to -3.74) significant reduction
	PUFA intake (through <code>study</code>) * : cont 7 (SD unclear), int 9 (SD unclear), (mean difference 2.00, 95% CI 1.57 to 2.43 assuming SDs of 5) significant increase
	PUFA n-3 intake: EPA, cont 0.6 (SD 0.7) g/wk, Int 2.4 (SD 1.4) g/wk
	PUFA n-6 intake: not reported
	MUFA intake (through <code>study</code>) ‡ : cont 13 (SD unclear), int 11 (SD unclear) (mean difference -2.00, 95% CI -2.43 to -1.57 assuming SDs of 5) significant reduction
	CHO intake (through study): cont 44 (SD 6), int 46 (SD 7) (mean difference 2.00, 95% CI 1.43 to 2.57) sig- nificant increase
	Protein intake (through study): cont 17 (SD 4), int 18 (SD 4) (mean difference 1.00, 95% CI 0.65 to 1.35) significant increase
	Trans fat intake: not reported
	Replacement for saturated fat: PUFA and CHO (by dietary aims), PUFA, CHO and protein (by di- etary achievements), main PUFA
	Style: diet advice
	Setting: community
	<pre>\$Estimated by subtraction (assuming total fat = SFA + PUFA + MUFA) or using the ratio (assuming P/ S = PUFA/SFA)</pre>
Outcomes	Stated trial outcomes: mortality, reinfarction Data available on total mortality? yes Cardiovascular mortality? yes Events available for combined cardiovascular events: cardiovascular deaths (including stroke deaths) plus non-fatal MI
	Secondary outcomes: cancer deaths, total MI, non-fatal MI, CHD mortality, CHD events (total MI)
	Tertiary outcomes: total and HDL cholesterol
Notes	Study duration: 24 months
	Study aim was to achieve low fat diet with raised P/S ratio and saturated fat intake in the intervention group was significantly lower than in the control group.
	SFA reduction aimed and achieved.
	Total serum cholesterol, difference between intervention and control, mmol/L: -0.26 (95% CI -0.36 to -0.16), statistically significant reduction
	Trial dates: Study dates not reported (published in 1989)
	Funding: Welsh Scheme for the Development of Health and Social Research, Welsh Heart Research Foundation, Flora Project, Health Promotion Research Trust. (Seven Seas Health Care and Duncan

Reduction in saturated fat intake for cardiovascular disease (Review)

Copyright @ 2020 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

DART 1989 (Continued)

Flockhart provided the MaxEPA capsules and Norgene provided 'Fybranta' tablets - but these were not used in the comparison discussed in this systematic review)

Declarations of Interest of primary researchers: none stated, all authors worked for academic or health institutions

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence genera- tion (selection bias)	Low risk	Randomised using sealed envelopes
Allocation concealment (selection bias)	Unclear risk	Unclear if envelopes were opaque
Blinding of participants and personnel (perfor- mance bias) All outcomes	High risk	Very difficult to blind trials where participants need to make their own dietary changes
Blinding of outcome as- sessment (detection bias) CVD outcomes	Low risk	Quote: "outcome assessors were not aware of study allocation" (Prof Burr, per- sonal communication). Method of blinding not stated
Blinding of outcome as- sessment (detection bias) All-cause mortality	Low risk	Blinding is not relevant in assessment of mortality
Incomplete outcome data (attrition bias) All outcomes	Low risk	GPs contacted for information on mortality and morbidity when participants did not attend
Selective reporting (re- porting bias)	Low risk	Not relevant for primary and secondary outcomes as we asked all trialists for data
Free of systematic differ- ence in care?	High risk	Different levels of advice appear to have been provided. See control and inter- vention methods in the Interventions section of the table of Characteristics of included studies
Stated aim to reduce SFA	Low risk	Aim to reduce SFA stated
Achieved SFA reduction	Low risk	SFA reduction achieved
Achieved TC reduction	Low risk	Statistically significant TC fall
Other bias	Low risk	None noted

Houtsmuller 1979

Study characteristics	
Methods	RCT
	Summary risk of bias: moderate to high
Participants	Adults with newly-diagnosed diabetes (the Netherlands)

Reduction in saturated fat intake for cardiovascular disease (Review)

Copyright @ 2020 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.



Houtsmuller 1979 (Continued)	CVD risk: moderate
	Control: 51 randomised, unclear how many analysed (all analysed re deaths) Intervention: 51 randomised, unclear how many analysed (all re deaths)
	Mean years in trial: unclear (max duration 6 years) % male: 56% overall Age: mean unclear
	Baseline total fat intake: unclear
	Baseline saturated fat intake: unclear
	Ethnicity: not stated
	Statins use allowed? Unclear
	% taking statins: Not reported (probably none as too early, pre-1980)
Interventions	Modified fat vs usual diet
	Control aims: SFA 35%E, CHO 50%E, protein 15%E Intervention aims: total fat 40%E, 1/3 linoleic acid, CHO 45%E, protein 15%E
	Control methods: unclear, surveyed by dietitian
	Intervention methods: unclear, surveyed by dietitian
	Intervention appears to be delivered by dietitian but no clear details on format or frequency
	Total fat intake: not reported
	Saturated fat intake: not reported (mean difference unclear)
	PUFA intake: not reported
	PUFA n-3 intake: not reported
	PUFA n-6 intake: not reported
	MUFA intake: not reported
	CHO intake: not reported
	Protein intake: not reported
	Trans fat intake: not reported
	Replacement for saturated fat: mainly PUFA (based on dietary aims)
	Style: diet advice?
	Setting: community
Outcomes	Stated trial outcomes: progression of diabetic retinopathy Data available on total mortality? no Cardiovascular mortality? no Events available for combined cardiovascular events: total MI and angina
	Secondary outcomes: total cholesterol, TGs (data read off graph), CHD mortality (fatal MI), CHD events (MI, angina)
Notes	Study duration 6 years. Study aim was for control group to take 35%E as saturated fat, and the inter- vention group 40%E from fat, of which 33% was from linoleic acid (so saturated fat < 27%E), but satu- rated fat intake during trial not reported

Houtsmuller 1979 (Continued)

SFA reduction aimed (unclear whether achieved).

Total serum cholesterol, difference between intervention and control, mmol/L: -0.47(95% CI -0.76 to -0.18), statistically significant reduction

Trial dates: Study recruitment 1973 to (unclear)

Funding: Dutch Heart Foundation

Declarations of Interest of primary researchers: none stated, all authors worked for academic or health institutions

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence genera- tion (selection bias)	Low risk	Participants matched in pairs then randomised
Allocation concealment (selection bias)	Unclear risk	Allocation method not clearly described
Blinding of participants and personnel (perfor- mance bias) All outcomes	High risk	Unclear, though unlikely as dietary advice provided
Blinding of outcome as- sessment (detection bias) CVD outcomes	Unclear risk	Blinding of outcome assessors not mentioned
Blinding of outcome as- sessment (detection bias) All-cause mortality	Low risk	Blinding is not relevant in assessment of mortality
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Unclear, deaths, cancer and CV events are dropouts, trialists asked for data - unclear if any data missing
Selective reporting (re- porting bias)	Low risk	Not relevant for primary and secondary outcomes as we asked all trialists for data
Free of systematic differ- ence in care?	Unclear risk	Level and type of intervention unclear. See control and intervention methods in the Interventions section of the table of Characteristics of included studies
Stated aim to reduce SFA	Low risk	Aim to reduce SFA stated
Achieved SFA reduction	Unclear risk	SFA intake not reported
Achieved TC reduction	Low risk	Statistically significant TC fall
Other bias	Low risk	None noted

Ley 2004

Study characteristics



Ley 2004 (Continued)

Methods	RCT
	Summary risk of bias: low
Participants	People with impaired glucose intolerance or high normal blood glucose (New Zealand) CVD risk: moderate Control: unclear how many randomised (176 between both groups), unclear how many analysed (112 between both groups at 5 years) Intervention: as above Mean years in trial: 4.1 over whole trial % male: control 80%, intervention 68% Age: mean control 52.0 (SE 0.8), intervention 52.5 (SE 0.8)
	Ethnicity: European 67% int, 77% control, Maori 11% int, 7% control, Pacific islander 20% int, 13% con- trol, Other 3% int, 4% control (outcomes not provided by ethnicity)
	Statins use allowed? Unclear
	% taking statins: Not reported
Interventions	Reduced fat vs usual diet
	Control aims: usual diet Intervention aims: reduced fat diet (no specific goal stated)
	Control methods: usual intake plus general advice on healthy eating consistent with the New Zealand guidelines and standard dietary information for people with nutrition-related problems upon entering the trial
	Intervention methods: monthly small group meetings to follow a 1-year structured programme aimed at reducing fat in the diet, includes education, personal goal-setting, self-monitoring
	Total fat intake, %E (at 1 year): int 26.1 (SD 7.7), cont 33.6 (SD 7.8) (mean difference -7.50, 95% CI -10.37 to -4.63) significant reduction
	Intervention delivered in small face-to-face groups but unclear by whom
	Saturated fat intake, %E (at 1 year): cont 13.4 (SD 4.7), int 10.0 (SD 4.2) (mean difference -3.40, 95% Cl -5.05 to -1.75) significant reduction
	PUFA intake, %E (at 1 year): cont 4.8 (SD 1.6), int 4.0 (SD 1.4) (mean difference -0.80, 95% CI -1.36 to -0.24) significant reduction
	PUFA n-3 intake: not reported
	PUFA n-6 intake: not reported
	MUFA intake, %E (at 1 year): cont 11.8 (SD 3.1), int 8.9 (SD 2.8) (mean difference -2.90, 95% CI -3.99 to -1.81) significant reduction
	CHO intake, %E (at 1 year): cont 45.8 (SD 10.9), int 54.2 (SD 10.5) (mean difference 8.40, 95% CI 4.44 to 12.36) significant increase
	Protein intake, %E (at 1 year): cont 16.6 (SD 3.9), int 18.4 (SD 3.5), (mean difference 1.80, 95% CI 0.43 to 3.17) significant increase
	Trans fat intake: not reported
	Replacement for saturated fat: carbohydrate and protein (based on dietary achievements)
	Style: diet advice
	Setting: community



Ley 2004 (Continued)		
Outcomes	Stated trial outcomes: lipids, glucose, blood pressure Data available on total mortality? yes Cardiovascular mortality? yes Events available for combined cardiovascular events: MI, angina, stroke, heart failure Secondary outcomes: total MI, stroke, cancer diagnoses, cancer deaths, CHD events (MI or angina) Tertiary outcomes: weight, total, LDL and HDL cholesterol, TGs, BP	
Notes	Study duration over 4 years	
	Study aim was to reduce total fat (not saturated fat), but saturated fat intake in the intervention group was significantly lower than in the control group.	
	SFA reduction achieved	
	Total serum cholesterol, difference between intervention and control, mmol/L: -0.05 (95% CI -0.46 to 0.36), NO statistically significant reduction and smaller than 0.20	
	Trial dates: Recruitment 1988 to 1990	
	Funding: National Heart Foundation of New Zealand, Auckland Medical Research Foundation, Lotteries Medical Board and the Health Research Council of New Zealand	
	Declarations of Interest of primary researchers: none stated, all authors worked for academic or health institutions	
Risk of bias		

Bias	Authors' judgement	Support for judgement
Random sequence genera- tion (selection bias)	Low risk	Unmarked opaque envelopes were opened by the person recruiting, unable to alter allocation later (trial author stated in their reply to us that randomisation and preparation of the envelopes was by people not involved in recruitment).
Allocation concealment (selection bias)	Low risk	Unmarked opaque envelopes were opened by the person recruiting, unable to alter allocation later.
Blinding of participants and personnel (perfor- mance bias) All outcomes	High risk	Dietary advice, not blinded
Blinding of outcome as- sessment (detection bias) CVD outcomes	Low risk	Trial authors stated that outcome assessors were blinded.
Blinding of outcome as- sessment (detection bias) All-cause mortality	Low risk	Blinding is not relevant in assessment of mortality.
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Unclear, deaths, cancer and CV events are dropouts, trialists were asked for data - unclear if any data missing
Selective reporting (re- porting bias)	Low risk	Not relevant for primary and secondary outcomes as we asked all trialists for data
Free of systematic differ- ence in care?	High risk	See control and intervention methods in the Interventions section of the table of Characteristics of included studies

Reduction in saturated fat intake for cardiovascular disease (Review)

Copyright @ 2020 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.



Ley 2004 (Continued)

Stated aim to reduce SFA	High risk	Aim to reduce SFA not stated
Achieved SFA reduction	Low risk	SFA reduction achieved
Achieved TC reduction	High risk	TC fall small (0.05 mmol/L only) and not statistically significant
Other bias	Low risk	None noted

Moy 2001

Study characteristics	
Methods	RCT
	Summary risk of bias: moderate to high
Participants	Middle-aged siblings of people with early CHD, with at least 1 CVD risk factor (USA) CVD risk: moderate Control: randomised 132, analysed 118 Intervention: randomised 135, analysed 117 Mean years in trial: 1.9 % male: control 49%, intervention 55% Age: control mean 45.7 (SD 7), intervention 46.2 (SD 7)
	Ethnicity: African-American 18% int, 25% control (remainder of group ethnicity not described, and out- comes not presented by ethnicity)
	Statins use allowed? Unclear (raised LDL cholesterol was a condition of entry, so use of statins probably minimal)
	% taking statins: Not reported
Interventions	Reduced fat intake vs usual diet
	Control aim: usual care
	Intervention aim: total fat 40 g/d or less
	Control methods: usual physician care with risk factor management at 0, 1 and 2 years
	Intervention methods: Individualised counselling by trained nurse, appointments 6 - 8 weekly for 2 years
	Intervention delivered individually, face-to-face by a trained nurse.
	Total fat intake, %E (at 2 years): int 34.1 (SD unclear), cont 38.0 (SD unclear) (mean difference -3.90, 95% CI -6.46 to -1.34 assuming SDs of 10) significant reduction
	Saturated fat intake, %E (at 2 years): int 11.5 (SD unclear), cont 14.4 (SD unclear) (mean differ- ence -2.90, 95% CI -4.18 to -1.62 assuming SDs of 5) significant reduction
	PUFA intake: not reported
	PUFA n-3 intake: not reported
	PUFA n-6 intake: not reported
	MUFA intake: not reported
	CHO intake: not reported

Moy 2001 (Continued)	
	Protein intake: not reported
	Trans fat intake: not reported
	Replacement for saturated fat: unclear
	Style: diet advice
	Setting: community
Outcomes	Stated trial outcomes: dietary intake Data available on total mortality? yes, no deaths Cardiovascular mortality? yes, no deaths Events available for combined cardiovascular events: total MI, stroke, unstable angina, PVD and PTCA
	Secondary outcomes: cancer diagnoses (no events), cancer deaths (none), stroke, total and non-fatal MI, CHD mortality (none), CHD events (MI or angina)
	Tertiary outcomes: BMI, HDL and LDL cholesterol, TG
Notes	Study duration 2 years
	Study aim was to reduce total fat based on ATPII dietary guidelines, and preliminary work established that this intervention reduced saturated fat and dietary cholesterol, and saturated fat intake was significantly lower than in the control group
	SFA reduction aimed and achieved
	Total serum cholesterol not reported, but <u>LDL</u> was, difference between intervention and control, mmol/L: -0.29 (95% CI -0.54 to -0.04), statistically significant reduction
	Trial dates: Study recruitment 1991 to 1994
	Funding: National Institute of Nursing Research, General Clinical Research Center of the National Insti- tutes of Health
	Declarations of Interest of primary researchers: none stated, all authors worked for academic or health institutions
Diele of him	

Bias	Authors' judgement	Support for judgement
Random sequence genera- tion (selection bias)	Low risk	Randomly assigned via computerised schema after all eligible siblings from a family had been screened
Allocation concealment (selection bias)	Unclear risk	Allocation method not clearly described
Blinding of participants and personnel (perfor- mance bias) All outcomes	High risk	Participants clear about their allocation
Blinding of outcome as- sessment (detection bias) CVD outcomes	High risk	Trialists clear about allocation
Blinding of outcome as- sessment (detection bias) All-cause mortality	Low risk	Blinding is not relevant in assessment of mortality.

Moy 2001 (Continued)

Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Unclear, deaths, cancer and CV events are dropouts, trialists were asked for data - unclear if any data missing
Selective reporting (re- porting bias)	Low risk	Not relevant for primary and secondary outcomes as all trialists asked for data
Free of systematic differ- ence in care?	High risk	Differences in frequency of follow-up, but unclear what differences in care oc- curred between the physician and nurse-led care. See control and interven- tion methods in Interventions section of the table of Characteristics of includ- ed studies
Stated aim to reduce SFA	Low risk	Aim to reduce SFA stated
Achieved SFA reduction	Low risk	SFA reduction achieved
Achieved TC reduction	Low risk	Statistically significant LDL fall (though TC not reported)
Other bias	Low risk	None noted

MRC 1968

Study characteristics	
Methods	RCT
	Medical Research Council (MRC)
	Summary risk of bias: moderate to high
Participants	Free-living men who have survived a first MI (UK) CVD risk: high Control: randomised 194, analysed 181 at 2 years Intervention: randomised 199, analysed 172 at 2 years Mean years in trial: control 3.7, intervention 3.8 % male: 100 Age: unclear (all < 60) Ethnicity: not stated Stating use allowed? Unclear (anti-coagulants allowed, but few other medications appear to have been
	used)
	% taking statins: Not reported (probably none as too early, pre-1980)
Interventions	Modified fat vs usual diet
	Control aims: usual diet Intervention aims: reduce dietary fat to 35 g fat per day, add 84 g soya oil per day
	Control methods: usual diet plus reducing diet (reduced CHO) for weight management for overweight men
	Intervention methods: instructed to follow a dietary regimen removing saturated fat from the diet plus daily dose of 85 g soya oil; half of it had to be taken unheated. Reduced CHO diet for weight manage- ment in overweight men
	Intervention appears to be delivered and supervised by trial dietitian but unclear how often.



MRC 1968 (Continued)	Total fat intake, %E (at 3.5 years): int 46 (SD unclear), cont 43 (SD unclear) (mean difference 3.00, 95%
	CI 0.91 to 5.09 assuming SDs of 10) significant increase
	Saturated fat intake: not reported (mean difference unclear)
	PUFA intake: not reported
	PUFA n-3 intake: not reported
	PUFA n-6 intake: not reported
	MUFA intake: not reported
	CHO intake: not reported
	Protein intake: not reported
	Trans fat intake: not reported
	Replacement for saturated fat: mainly PUFA (based on dietary goals)
	Style: diet advice & supplement (soy oil)
	Setting: community
Outcomes	Stated trial outcomes: MI or sudden death Data available on total mortality? yes Cardiovascular mortality? yes Events available for combined cardiovascular events: cardiovascular deaths and fatal or non-fatal MI
	Secondary outcomes: total and non-fatal MI, stroke, cancer deaths, CHD mortality, CHD events (CHD mortality or non-fatal MI)
	Tertiary outcomes: none (data for weight, total cholesterol and BP, but no variance info)
Notes	Study duration over 6 years
	Study aim: for intervention "saturated fats were replaced by polyunsaturated fats", but saturated fat intakes during trial were not reported.
	SFA reduction aimed
	Total serum cholesterol, difference between intervention and control, mmol/L: -0.64 (95% CI un- clear), reduction > 0.20
	For all, data at 4 years, control n = 89, intervention n = 88
	Weight change: control -3 kg, intervention 0 kg
	Total cholesterol change: control -0.47 mmol/L, intervention -1.11 mmol/L
	Systolic BP change: control 0 mmHg, intervention +2 mmHg
	Diastolic BP change: control +3 mmHg, intervention -1 mmHg
	Trial dates: Study recruitment 1960 to 1965, analysed 1967
	Funding: Medical Research Council
	Declarations of Interest of primary researchers: none stated, all authors worked for academic or health institutions.
Risk of bias	
Bias	Authors' judgement Support for judgement

Reduction in saturated fat intake for cardiovascular disease (Review)

2

Copyright @ 2020 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.



MRC 1968 (Continued)

Random sequence genera- tion (selection bias)	Low risk	Quote: "using random numbers, by blocks within hospitals"
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding of participants and personnel (perfor- mance bias) All outcomes	High risk	Big changes to fat intake in intervention group while control group ate their usual diet
Blinding of outcome as- sessment (detection bias) CVD outcomes	Low risk	Quote: "Suspected relapses were assessed at regular intervals by a review committee unaware of the patients diet group".
Blinding of outcome as- sessment (detection bias) All-cause mortality	Low risk	Blinding is not relevant in assessment of mortality.
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Data collection was thorough, but some participants dropped out and contact was lost, so some events may have been missed.
Selective reporting (re- porting bias)	Low risk	Not relevant for primary and secondary outcomes as all trialists were asked for data
Free of systematic differ- ence in care?	High risk	Unlikely as control group continued diet as usual, intervention group were likely to have had additional contact. See control and intervention methods in the Interventions section of the table of Characteristics of included studies
Stated aim to reduce SFA	Low risk	Aim to reduce SFA stated
Achieved SFA reduction	Unclear risk	SFA intake not reported
Achieved TC reduction	Low risk	Although statistical significance was not reported or calculable, TC in the in- tervention group was 0.64 mmol/L lower than in the control group, a large fall (and almost certainly statistically significant).
Other bias	Low risk	None noted

Oslo Diet-Heart 1966

Study characteristics	
Methods	RCT
	Oslo Diet-Heart Trial
	Summary risk of bias: moderate to high for CVD outcomes, low for all-cause mortality
Participants	Men with previous MI (Norway) CVD risk: high Control: randomised 206, analysed 148 (at 5 years) Intervention: randomised 206, analysed 152 (at 5 years) Mean years in trial: control 4.3, intervention 4.3 % male: 100

Oslo Diet-Heart 1966 (Continue	^{2d)} Age: mean control 56.3, intervention 56.2 (all 30 - 67)
	Ethnicity: ethnicity not mentioned
	Statins use allowed? Unclear (medications not mentioned as exclusion criteria, most appeared to be on anti-coagulant medications, statins not mentioned)
	% taking statins: Not reported (probably none as too early, pre-1980)
Interventions	Modified fat diet vs control
	Control aims: no dietary advice but direct questions answered, supplement = 1 vitamin tablet daily Intervention aims: reduce meat and dairy fats, increase fish, vegetables, supplement - 1 vitamin tablet daily, 0.5 L soy bean oil per week (free to 25% of participants), sardines in cod liver oil (free at certain times to encourage compliance)
	Control methods: usual diet
	Intervention methods: continuous instruction and supervision by dietitian, including home visits, let- ters and phone calls
	Total fat intake: unclear (note - intake of total fat, carbohydrate, protein and sugar was assessed in 17 "especially conscientious and positive" as well as intelligent dieters, but this was not reported here as unlikely to be representative, and lacking control group data)
	Saturated fat intake: unclear (mean difference unclear)
	PUFA intake: unclear
	PUFA n-3 intake: not reported
	PUFA n-6 intake: not reported
	MUFA intake: unclear
	CHO intake: unclear
	Protein intake: unclear
	Trans fat intake: unclear
	Replacement for saturated fat: PUFA (based on dietary goals)
	Style: diet advice and supplement (food)
	Setting: community
Outcomes	Stated trial outcomes: coronary heart disease morbidity and mortality Data available on total mortality? yes Cardiovascular mortality? yes Events available for combined cardiovascular events: total MI, sudden death, stroke, angina
	Secondary outcomes: non-fatal and total MI, stroke, CHD mortality (fatal MI and sudden death), CHD events (MI, angina and sudden death)
	Tertiary outcomes: weight, total cholesterol, systolic and diastolic BP (but no variance information was provided)
Notes	Study duration over 4 years
	Study aim was to reduce serum cholesterol by a diet "low in saturated fats and in cholesterol, and rich in highly unsaturated fats", saturated fat intakes during study were not reported
	SFA reduction aimed (reduction unclear as not measured except in a highly compliant subgroup)



Oslo Diet-Heart 1966 (Continued)

Total serum cholesterol, difference between intervention and control, mmol/L: -1.07 (95% CI unclear), reduction > 0.20

Weight change from baseline was -0.5 kg in the control group (n \sim 155), -2.5 kg in the intervention group (n \sim 160) at 51 months

Total cholesterol change from baseline was -0.46 mmol/L in the control group and -1.53 mmol/L in the intervention group at 51 months

Systolic BP at baseline was 153.8 mmHg in control and 159.0 in intervention, and mean sBP through trial was 154.3 mmHg in control and 158.2 mmHg in the intervention group.

Diastolic BP at baseline was 93.5 mmHg in control and 97.1 mmHg in intervention, through trial mean dBP was 95.5 mmHg in control and 98.6 mmHg in intervention participants

Trial dates: Recruitment 1956 to 1958

Funding: Det Norske Råd for Hjerte- og karsyk-dommer, A/S Freia Chokoladefabriks Arbeidsfond for Ernærings-forskning, JL Tiedemanns Tobaksfabrik Joh H Andresens medisinske fond, plus A/S Farmacöytisk Industri provided a multivitamin free of charge, DE-NO-FA and Lillleborg Fabriker provided soy bean oil at reduced prices, the Research Laboratory of the Norwegian Canning Industry, Stavanger Preserving Co and Kommendal Packing Comp provided Norwegian sardines in cod liver oil free to those in the intervention group.

Declarations of Interest of primary researchers: none stated, all authors worked for academic or health institutions.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence genera- tion (selection bias)	Low risk	"table of random numbers used", by Prof Knut Westlund
Allocation concealment (selection bias)	Low risk	Randomisation appears to have occurred before medical examination with- in the study, so was not affected by participant characteristics and was con- cealed.
Blinding of participants and personnel (perfor- mance bias) All outcomes	High risk	Participants were aware of their allocation as was the main trialist.
Blinding of outcome as- sessment (detection bias) CVD outcomes	Unclear risk	Outcomes were categorised by a diagnostic board, but their blinded status was unclear.
Blinding of outcome as- sessment (detection bias) All-cause mortality	Low risk	Blinding is not relevant in assessment of mortality.
Incomplete outcome data (attrition bias) All outcomes	Low risk	The participants who could not be directly followed up for the 5 years were fol- lowed until death or study end through personal interviews, or contact with their physicians or relatives.
Selective reporting (re- porting bias)	Low risk	Not relevant for primary and secondary outcomes as all trialists were asked for data

Oslo Diet-Heart 1966 (Continued)

Free of systematic differ- ence in care?	High risk	Dietetic input level very different, although medical care appeared similar. See control and intervention methods in the Interventions section of the table of Characteristics of included studies
Stated aim to reduce SFA	Low risk	Aim to reduce SFA stated
Achieved SFA reduction	Unclear risk	SFA intake not reported
Achieved TC reduction	Low risk	Although statistical significance was not reported or calculable, TC in the in- tervention group was 1.07 mmol/L lower than in the control group, a large fall (and almost certainly statistically significant).
Other bias	Low risk	None noted

Oxford Retinopathy 1978

Study characteristics			
Methods	RCT		
	Summary risk of bias: moderate to high for CVD outcomes, low for all-cause mortality		
Participants	Newly-diagnosed non-insulin-dependent diabetics (UK) CVD risk: moderate Control: number randomised unclear (249 split between the 2 groups, 125?), number analysed for mor- tality unclear (all but 2 overall at 16 years) Intervention: number randomised unclear (249 split between the 2 groups, 125?), number analysed as above Mean years in trial: overall 9.3? % male: overall 49% Age: mean overall 47.1 (all < 65) Ethnicity: not stated		
	Statins use allowed? Unclear		
	% taking statins: Not reported (probably none as too early, pre-1980)		
Interventions	Reduced and modified dietary fat vs average diet		
	Control aims: total fat 40%E, PUFA 12%E, protein 20%E, CHO 40%E (reducing simple sugars), 1500 kcal/ day Intervention aims: total fat 26%E, PUFA 16%E, protein 20%E, CHO 54%E (reducing simple sugars), 1500 kcal/day		
	Control methods: dietary advice from diabetes dietitian		
	Intervention methods: dietary advice from diabetes dietitian		
	Total fat intake, %E (at 7 - 9 years)§: int 32 (SD unclear), cont 41 (SD unclear) (mean difference -9.00, 95% CI -11.48 to -6.52 assuming SDs of 10) significant reduction		
	Saturated fat intake, %E (at 7 - 9 years)§: int 10.7 (SD unclear), cont 20.4 (SD unclear) (mean dif- ference -9.70, 95% CI -10.94 to -8.46 assuming SD of 5) significant reduction		
	PUFA intake, %E (at 7 - 9 years)§: int 11.8 (SD unclear), cont 2.1 (SD unclear) (mean difference 9.70, 95% CI 8.46 to 10.94 assuming SDs of 5) significant increase		
	PUFA n-3 intake: not reported		

Oxford Retinopathy 1978 (Con	^{ntinued)} PUFA n-6 intake: not re	ported	
	MUFA intake, %E (at 7 - 95% CI -10.34 to 7.86 as	9 years)§: int 9.5 (SD unclear), cont 18.6 (SD unclear) (mean difference -9.10, suming SDs of 5) significant reduction	
	Carbohydrate intake: n	ot reported	
	Protein intake: not repo	orted	
	Trans fat intake: not reported		
	Replacement for saturated fat: PUFA and CHO (based on dietary goals and achievements)		
	Style: diet advice		
	Setting: community (ou	tpatients clinic)	
	§validity of these data i Source: Lopez-Espinoza	s questionable as it represents only 3 intervention and 3 control participants. 1984	
Outcomes	Stated trial outcomes: Data available on total deaths at 10 years) Cardiovascular mortalit Events available for cor	etinopathy mortality? yes, but unable to ascertain from which intervention groups (34 cy? no nbined cardiovascular events: none	
	Secondary outcomes: n	one	
	Tertiary outcomes: BMI	, total cholesterol	
Notes	Study duration over 9 y	ears	
	Study aim was to reduce in the intervention grou	e total fat and increase PUFAs (so reducing saturates), and saturated fat intake p was significantly lower than in the control group	
	SFA reduction achieve	d	
	Total serum cholester to 0.48), NO statistical	ol, difference between intervention and control, mmol/L: 0.07 (95% CI -0.34 ly significant reduction and smaller than 0.20	
	Trial dates: Recruitment 1973 to 1976		
	Funding: Oxford Diabet Inc	es Trust, British Diabetic Association, International Sugar Research Foundation	
	Declarations of Interest institutions.	of primary researchers: none stated, all authors worked for academic or health	
Risk of bias			
Bias	Authors' judgement	Support for judgement	
Random sequence genera- tion (selection bias)	Low risk	"random number sequence, provided and allotted by a separate agency" (Prof Richard Peto)	

Allocation concealment "random number sequence, provided and allotted by a separate agency" (Prof Low risk Richard Peto)

Participants were not blinded.

Blinding of participants High risk and personnel (performance bias) All outcomes

(selection bias)



Oxford Retinopathy 1978 (Continued)

Blinding of outcome as- sessment (detection bias) CVD outcomes	Unclear risk	Unclear whether physicians blinded to allocation
Blinding of outcome as- sessment (detection bias) All-cause mortality	Low risk	Blinding is not relevant in assessment of mortality.
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Unclear, deaths, cancer and CV events are dropouts - unclear if any data miss- ing
Selective reporting (re- porting bias)	Low risk	Not relevant for primary and secondary outcomes as all trialists were asked for data
Free of systematic differ- ence in care?	Low risk	Dietetic advice for both groups. See control and intervention methods in the Interventions section of the table of Characteristics of included studies
Stated aim to reduce SFA	High risk	Aim to reduce SFA not stated
Achieved SFA reduction	Low risk	SFA reduction achieved
Achieved TC reduction	High risk	No statistically significant TC fall, and difference only 0.07 mmol/L
Other bias	Low risk	None noted

Rose corn oil 1965

Study characteristics		
Methods	RCT	
	Summary risk of bias: moderate to high	
Participants	Men (?) with angina or following MI (UK) CVD risk: high	
	Intervention - corn: randomised 26, analysed 13 Mean years in trial: control 1.7, corn 1.5 % male: unclear (100%?) Age: mean control 58.8, corn 52.6 (all < 70)	
	Ethnicity: not stated	
	Statins use allowed? Unclear (anti-coagulants not allowed, but all participants received convention treatments at the discretion of their physicians)	
	% taking statins: Not reported (probably none as too early, pre-1980)	
Interventions	Modified fat vs usual diet	
	Control aims: usual diet	
	Intervention aims - corn: restrict dietary fat, plus 80 g/day corn oil provided	
	Control methods: usual physician care plus follow-up clinic monthly, then every 2 months, no dietary fat advice or oil provided	

Reduction in saturated fat intake for cardiovascular disease (Review)

Copyright @ 2020 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.



Rose corn oil 1965 (Continued)	Intervention methods: usual physician care plus follow-up clinic monthly, then every 2 months, dietary fat advice plus oil provided			
	Unclear how the advice was delivered or by whom			
	Total fat intake, %E (at 18 months): corn 50.5 (SD unclear), cont 32.6 (SD unclear) (mean difference 17.90, 95% CI 10.77 to 25.03 assuming SDs of 10) significant increase			
	Saturated fat intake: unclear (mean difference unclear)			
	PUFA intake: unclear			
	PUFA n-3 intake: not reported			
	PUFA n-6 intake: not reported			
	MUFA intake: unclear			
	CHO intake, %E (at 18 months): corn 36.5 (SD unclear), cont 51.5 (SD unclear) (mean difference -15.00, 95% CI -29.27 to -0.73 assuming SDs of 20) significant reduction			
	Protein intake, %E (at 18 months): corn 11.0 (SD unclear), cont 13.2 (SD unclear) (mean difference -2.20, 95% CI -5.77 to 1.37 assuming SDs of 5) no significant difference			
	Trans fat intake: unclear			
	Replacement for saturated fat: mainly PUFA (based on aims and achievements)			
	Style: diet advice and supplement (oil)			
	Setting: community			
Outcomes	Stated trial outcomes: cardiac events Data available on total mortality? yes Cardiovascular mortality? yes Events available for combined cardiovascular events: cardiovascular deaths, non-fatal MI, angina, stroke			
	Secondary outcomes: stroke (none), non-fatal and total MI, CHD mortality (fatal MI and sudden death), CHD events (all MI and sudden death)			
	Tertiary outcomes: total cholesterol			
Notes	Study duration 2 years			
	Study aim was to reduce total fat (by restricting fatty meat, sausages, pastry, ice cream, cheese, cake, milk, eggs and butter) and prescribe vegetable oil (so reducing saturates), but saturated fat intakes during intervention were not reported.			
	SFA reduction aimed (but unclear whether achieved as SFA intake not reported)			
	Total serum cholesterol, difference between intervention and control, mmol/L: -0.58 (95% CI -1.42 to 0.26), NO statistically significant reduction but > 0.20			
	Trial dates: unclear, published in 1965			
	Funding: probably unfunded (they thank the Paddington General Hospital for clinic facilities, and St Mary's and Paddington General Hospital physicians for referral of patients, but no funding acknowl- edged)			
	Declarations of Interest of primary researchers: none stated, all authors worked for academic or health institutions.			

Risk of bias



Rose corn oil 1965 (Continued)

Bias	Authors' judgement	Support for judgement
Random sequence genera- tion (selection bias)	Low risk	Trial was stated as "randomised" but without further detail, apart from use of a sealed envelope as below.
Allocation concealment (selection bias)	Unclear risk	When a new participant was accepted for the trial a sealed envelope was opened containing the allocation instructions. In the case of participants allocated to an oil group, the instructions referred only to a code number.
Blinding of participants and personnel (perfor- mance bias) All outcomes	High risk	The physicians in charge knew which participants were receiving oil, but they did not know until the end of the trial the kind of oil that they were receiving.
Blinding of outcome as- sessment (detection bias) CVD outcomes	Low risk	The electrocardiograms were assessed without the knowledge of the participant's treatment group.
Blinding of outcome as- sessment (detection bias) All-cause mortality	Low risk	Blinding is not relevant in assessment of mortality.
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Some lost to follow-up by 2 years, so some events may have been missed
Selective reporting (re- porting bias)	Low risk	Not relevant for primary and secondary outcomes as all trialists were asked for data.
Free of systematic differ- ence in care?	Low risk	All received conventional treatments at the discretion of the physicians, all at- tended a special follow-up clinic. See control and intervention methods in the Interventions section of the table of Characteristics of included studies
Stated aim to reduce SFA	Low risk	Aim to reduce SFA stated
Achieved SFA reduction	Unclear risk	SFA intake not reported
Achieved TC reduction	High risk	Although the TC in the intervention group was 0.58 mmol/L lower than in the control group, this was not statistically significant in this small study.
Other bias	Low risk	None noted

Rose olive 1965

Study characteristics	
Methods	RCT
	Summary risk of bias: moderate to high
Participants	Men (?) with angina or following MI (UK) CVD risk: high Control: randomised 26, analysed 18 Intervention - olive: randomised 28, analysed 12 Mean years in trial: control 1.7, olive 1.5



Rose olive 1965 (Continued)	% male: unclear (100%?) Age: mean control 58.8, olive 55.0 (all < 70)			
	Ethnicity: Not stated			
	Statins use allowed? Unclear (anti-coagulants not allowed, but all participants received conventional treatments at the discretion of their physicians)			
	% taking statins: Not reported (probably none as too early, pre-1980)			
Interventions	Modified fat vs usual diet			
	Control aims: usual diet Intervention aims - olive: restrict dietary fat, plus 80 g/day olive oil provided			
	Control methods: usual physician care plus follow-up clinic monthly, then every 2 months, no dietary fat advice or oil provided			
	Intervention methods: usual physician care plus follow-up clinic monthly, then every 2 months, dietary fat advice plus oil provided			
	Unclear how the advice was delivered or by whom			
	Total fat intake, %E (at 18 months): olive 46.2 (SD unclear), cont 32.6 (SD unclear) (mean difference 13.60, 95% CI 6.30 to 20.90 assuming SDs of 10) significant increase			
	Saturated fat intake: unclear (mean difference unclear)			
	PUFA intake: unclear			
	PUFA n-3 intake: not reported			
	PUFA n-6 intake: not reported			
	MUFA intake: unclear			
	CHO intake, %E (at 18 months): olive 42.2 (SD unclear), cont 51.5 (SD unclear) (mean difference -9.30, 95% CI -23.91 to 5.31 assuming SDs of 20) no significant difference			
	Protein intake, %E (at 18 months): olive 9.6 (SD unclear), cont 13.2 (SD unclear) (mean difference -3.60, 95% CI -7.25 to 0.05 assuming SDs of 5) no significant difference			
	Trans fat intake: unclear			
	Replacement for saturated fat: mainly MUFA (based on dietary aims)			
	Style: diet advice and supplement (oil)			
	Setting: community			
Outcomes	Stated trial outcomes: cardiac events Data available on total mortality? yes Cardiovascular mortality? yes Events available for combined cardiovascular events: cardiovascular deaths, non-fatal MI, angina, stroke			
	Secondary outcomes: stroke (none), non-fatal and total MI, CHD mortality (fatal MI and sudden death), CHD events (all MI and sudden death)			
	Tertiary outcomes: total cholesterol			
Notes	Study duration 2 years			



Rose olive 1965 (Continued)

Study aim was to reduce total fat (by restricting fatty meat, sausages, pastry, ice cream, cheese, cake, milk, eggs and butter) and prescribe vegetable oil (so reducing saturates), but saturated fat intakes during intervention were not reported

SFA reduction aimed (but unclear whether achieved as SFA intake not reported)

Total serum cholesterol, difference between intervention and control, mmol/L: 0.30 (95% CI -0.93 to 1.53), NO statistically significant reduction, mean total cholesterol rose

Trial dates: unclear, published in 1965

Funding: probably unfunded (they thank the Paddington General Hospital for clinic facilities, and St Mary's and Paddington General Hospital physicians for referral of patients, but no funding acknowledged)

Declarations of Interest of primary researchers: none stated, all authors worked for academic or health institutions.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence genera- tion (selection bias)	Low risk	Trial was stated as "randomised" but without further detail, apart from use of a sealed envelope as below.
Allocation concealment (selection bias)	Unclear risk	When a new participant was accepted for the trial a sealed envelope was opened containing the allocation instructions. In the case of participants allocated to an oil group, the instructions referred only to a code number.
Blinding of participants and personnel (perfor- mance bias) All outcomes	High risk	The physicians in charge knew which participants were receiving oil, but they did not know until the end of the trial the kind of oil that they were receiving.
Blinding of outcome as- sessment (detection bias) CVD outcomes	Low risk	The electrocardiograms were assessed without the knowledge of the participant's treatment group.
Blinding of outcome as- sessment (detection bias) All-cause mortality	Low risk	Blinding is not relevant in assessment of mortality.
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Some lost to follow-up by 2 years, so some events may have been missed
Selective reporting (re- porting bias)	Low risk	Not relevant for primary and secondary outcomes as all trialists were asked for data.
Free of systematic differ- ence in care?	Low risk	All received conventional treatments at the discretion of the physicians, all at- tended a special follow-up clinic. See control and intervention methods in the Interventions section of the table of Characteristics of included studies
Stated aim to reduce SFA	Low risk	Aim to reduce SFA stated
Achieved SFA reduction	Unclear risk	SFA intake not reported
Achieved TC reduction	High risk	Although the TC in the intervention group was 0.58 mmol/L lower than in the control group, this was not statistically significant in this small study.

Reduction in saturated fat intake for cardiovascular disease (Review)

Copyright \odot 2020 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.



Rose olive 1965 (Continued)

Other bias

Low risk

None noted

Simon 1997

Study characteristics	5		
Methods	RCT		
	Summary risk of bias: moderate to high		
Participants	Women with a high risk of breast cancer (USA) CVD risk: low Control: randomised 96, analysed 75 Intervention: randomised 98, analysed 72 Mean years in trial: control 1.8, intervention 1.7 % male: 0 Age: mean control 46, intervention 46		
	Ethnicity: White 89%, African-American 9%, Hispanic 2%		
	Statins use allowed? No (those on lipid-lowering medications were excluded)		
	% taking statins: 0%		
Interventions	Reduced fat vs usual diet		
	Control aims: usual diet Intervention aims: total fat 15%E		
	Control methods: continued usual diet		
	Intervention methods: Bi-weekly individual dietetic appointments over 3 months followed by month- ly individual or group appointments, including education, goal-setting, evaluation, feedback and self- monitoring		
	Intervention delivered face-to-face by a dietitian		
	Total fat intake, %E (at 12 months)§: int 17.6 (SD 5.8), cont 33.8 (SD 7.4) (mean difference -16.20, 95% CI -18.34 to -14.06) significant reduction		
	Saturated fat intake, %E (at 12 months)§: int 6.0 (SD 3.0), cont 12.1 (SD 5.2) (mean difference -6.10, 95% Cl -7.47 to -4.73) significant reduction		
	PUFA intake, %E (at 12 months)§: int 3.8 (SD 1.7), cont 7.3 (SD 4.1) (mean difference -3.50, 95% CI -4.51 to -2.49) significant reduction		
	PUFA n-3 intake: not reported		
	PUFA n-6 intake: not reported		
	MUFA intake, %E (at 12 months)§: int 6.1 (SD 3.0), cont 12.8 (SD 6.3) (mean difference -6.70, 95% CI -8.29 to -5.11) significant reduction		
	CHO intake: not reported		
	Protein intake: not reported		
	Trans fat intake: not reported		
	Replacement for saturated fat: unclear, either carbohydrate or protein (based on aims and achievements)		

Simon 1997 (Continued)	
	Style: diet advice
	Setting: community
	§Kasim 1993
Outcomes	Stated trial outcomes: intervention feasibility Data available on total mortality? yes (2 deaths, but not clear in which arms) Cardiovascular mortality? no Events available for combined cardiovascular events: none
	Secondary outcomes: cancer diagnosis (8 diagnoses, but not clear in which arms)
	Tertiary outcomes: weight, total, LDL and HDL cholesterol, TGs
Notes	Study duration 2 years
	Study aim was to reduce total fat to 15%E (saturated fat not mentioned), but saturated fat intake in the intervention group was significantly lower than in the control group
	SFA reduction achieved
	Total serum cholesterol, difference between intervention and control, mmol/L: -0.34 (95% CI -0.64 to -0.04), statistically significant reduction
	Trial dates: Recruitment 1987 to 1989
	Funding: Marilyn J Smith Fund, Harper-Grace Hospitals, the Wesley Foundation, National Cancer Insti- tute, Karmanos Cancer Institute Core Grant, the United Foundation of Detroit
	Declarations of Interest of primary researchers: none stated, all authors worked for academic or health institutions except PN Kim who was affiliated with Wesley Health Strategies (now Health Strategies, which offers a "full-service health and fitness centre with an educated fitness staff and spacious work-out areas", see healthstrategiesfitness.com/)

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence genera- tion (selection bias)	Low risk	Randomisation method not clearly described, but participants were stratified by age and randomised (block size 2).
Allocation concealment (selection bias)	Unclear risk	Allocation method not clearly described
Blinding of participants and personnel (perfor- mance bias) All outcomes	High risk	Participants knew their allocation.
Blinding of outcome as- sessment (detection bias) CVD outcomes	Unclear risk	Unclear whether physicians knew allocations
Blinding of outcome as- sessment (detection bias) All-cause mortality	Low risk	Blinding is not relevant in assessment of mortality.
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Unclear, deaths, cancer and CV events are dropouts - unclear if any data miss- ing

Reduction in saturated fat intake for cardiovascular disease (Review)

Copyright @ 2020 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

Simon 1997 (Continued)

Cochrane

Library

Selective reporting (re- porting bias)	Low risk	Not relevant for primary and secondary outcomes as all trialists were asked for data
Free of systematic differ- ence in care?	High risk	Very different contact time with dietitian, but medical appointments same in both groups. See control and intervention methods in the Interventions sec- tion of the table of Characteristics of included studies
Stated aim to reduce SFA	High risk	Aim to reduce SFA not stated
Achieved SFA reduction	Low risk	SFA reduction achieved
Achieved TC reduction	Low risk	Statistically significant TC fall
Other bias	Low risk	None noted

STARS 1992

Study characteristics	
Methods	RCT
	St Thomas' Atherosclerosis Regression Study (STARS)
	Summary risk of bias: moderate to high for CVD outcomes, low for all-cause mortality
Participants	Men with angina referred for angiography (UK) CVD risk: high Control: unclear how many randomised (30?), analysed 24 Intervention: unclear how many randomised (30?), analysed 26 Mean years in trial: control 2.9, intervention 3.0 % male: 100% Age: mean control 53.9, intervention 48.9 (all < 66) Ethnicity: not stated
	Statins use allowed? No (1 arm of the trial, not described here, prescribed cholestyramine)
	% taking statins: 0%
Interventions	Reduced and modified fat diet vs usual diet
	Control aims: no diet intervention but advised to lose weight if BMI > 25 Intervention aims: total fat 27%E, SFA 8 - 10%E, omega-3 and omega-6 PUFA 8%E, increase in plant-de- rived soluble fibre, dietary cholesterol 100 mg/1000 kcal, advised to lose weight if BMI > 25
	Control methods: usual care but no formal dietetic counselling. They were counselled against smoking if appropriate and advised about daily exercise level.
	Intervention methods: Usual care plus dietetic individual assessment of diet and advice. Further dietet- ic counselling and food stuffs were given to participants who did not achieve or maintain certain levels of serum cholesterol reduction
	Initial intervention was delivered individually face-to-face by a dietitian and follow-up by a clinician.
	Total fat intake, %E (through study): int 27 (SD 7), cont 37 (SD 5) (mean difference -10.00, 95% CI -13.35 to -6.65) significant reduction
	Saturated fat intake, %E (through study): int 9 (SD 3), cont 16 (SD 4) (mean difference -7.00, 95% CI -8.97 to -5.03) significant reduction



STARS 1992 (Continued)	PUFA intake, %E (throu significant increase	igh study)§: int 7 (SD 2), cont 5 (SD 2) (mean difference 2.00, 95% CI 0.89 to 3.11)
	PUFA n-3 intake: not re	ported
	PUFA n-6 intake: not re	ported
	MUFA intake, %E (throu -4.48) significant reduc	ugh study)§: int 10 (SD 4), cont 17 (SD 5) (mean difference -7.00, 95% CI -9.52 to tion
	CHO intake, %E (throug 11.88) significant incre	gh study)§: int 49 (SD 7), cont 41 (SD 7) (mean difference 8.00, 95% Cl 4.12 to ase
	Protein intake, %E (thr 2.73) no significant effe	ough study)§: int 19 (SD 4), cont 18 (SD 2) (mean difference 1.00, 95% CI -0.73 to ect
	Trans fat intake: not re	ported
	Replacement for satu	rated fat: CHO and PUFA (based on aims and achievements)
	Style: diet advice	
	Setting: community	
	§Blann 1995	
Outcomes	Stated trial outcomes: Data available on total Cardiovascular mortali Events available for con stroke, CABG, angiopla	angiography mortality? yes ty? yes mbined cardiovascular events: cardiovascular deaths, non-fatal MI, angina, sty, stroke, total MI, CHD events, plus cancer deaths (none)
	Secondary outcomes: t glucose (weight and BF means)	otal, HDL, LDL cholesterol, TGs, total/HDL and LDL/HDL ratios, 2-hour post-load "remained similar" but were not reported, Lp(a) reported but as geometric
Notes	Study duration: 3 years	
	Study aim was to redu was significantly reduc	ce saturated fats (to 8 - 10%E), and saturated fat intake in the intervention group ed
	SFA reduction aimed a	and achieved
	Total serum cholester -1.19 to -0.33), statist	ol, difference between intervention and control, mmol/L: -0.76 (95% CI ically significant reduction
	Trial dates: Study dates	s not reported (published in 1992)
	Funding: Unilever plc, 1	he Chemical Pathology Fund of St Thomas' Hospital, and Bristol-Meyers Ltd
	Declarations of Interes institutions.	t of primary researchers: none stated, all authors worked for academic or health
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence genera- tion (selection bias)	Low risk	"blinded random cards issued centrally by statistician advisor"
Allocation concealment (selection bias)	Low risk	"blinded random cards issued centrally by statistician advisor"



STARS 1992 (Continued)		
Blinding of participants and personnel (perfor- mance bias) All outcomes	High risk	Participant blinding: inadequate
Blinding of outcome as- sessment (detection bias) CVD outcomes	Unclear risk	Physician blinding: unclear
Blinding of outcome as- sessment (detection bias) All-cause mortality	Low risk	Blinding is not relevant in assessment of mortality.
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Unclear, deaths, cancer and CV events are dropouts - unclear if any data miss- ing
Selective reporting (re- porting bias)	Low risk	Not relevant for primary and secondary outcomes as all trialists were asked for data
Free of systematic differ- ence in care?	High risk	Usual care in both groups, dietetic counselling only in the intervention group. See control and intervention methods in the Interventions section of the table of Characteristics of included studies
Stated aim to reduce SFA	Low risk	Aim to reduce SFA stated
Achieved SFA reduction	Low risk	SFA reduction achieved
Achieved TC reduction	Low risk	Statistically significant TC fall
Other bias	Low risk	None noted

Sydney Diet-Heart 1978

Study characteristics	
Methods	RCT
	Sydney Diet-Heart Trial
	Summary risk of bias: moderate to high
Participants	Men with previous MI (Australia) CVD risk: high Control: randomised 237, analysed 221 at 2 years Intervention: randomised 221, analysed 205 at 2 years Mean years in trial: control 4.3, intervention 4.3 % male: 100 Age: mean control 49.1 (SD 6.5), intervention 48.7 (SD 6.8) Ethnicity: not stated Statins use allowed? Unclear (use of medication did not appear to be an exclusion criteria) % taking statins: Not reported (probably none as too early, pre-1980)
Interventions	Modified fat diet vs usual diet

Reduction in saturated fat intake for cardiovascular disease (Review)

Copyright @ 2020 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.



Sydney Diet-Heart 1978 (Continued)

Control aims: reduction in energy if overweight, no other specific dietary advice, allowed to use PUFA margarine instead of butter

Intervention aims: SFA 10%E, PUFA 15%E, reduction in energy if overweight, dietary chol < 300 mg/day

Control methods: no specific dietary instruction (except re weight)

Intervention methods: advised and tutored individually, diet assessed 3 times in 1st year and twice annually thereafter

Intervention was delivered face-to-face individually but unclear by whom

Total fat intake, %E ("during follow-up"): int 38.3 (SD 5.9), cont 38.1 (SD 5.4) (mean difference 0.20, 95% CI -0.88 to 1.28) no significant difference

Saturated fat intake, %E ("during follow-up"): int 9.8 (SD 2.6), cont 13.5 (SD 3.2) (mean difference -3.70, 95% CI -4.25 to -3.15) significant reduction

PUFA intake, %E ("during follow-up"): int 15.1 (SD 4.3), cont 8.9 (SD 3.5) (mean difference 6.20, 95% CI 5.45 to 6.95) significant increase

PUFA n-3 intake: not reported

PUFA n-6 intake: not reported

MUFA intake, %E ("during follow-up"): int 11.5 (SD 2.1), cont 13.8 (SD 2.5) (mean difference -2.30, 95% CI -2.74 to -1.86) significant reduction

CHO intake, %E ("during follow-up"): int 40.9 (SD 7.3), cont 40.3 (SD 7.3) (mean difference 0.60, 95% CI -0.79 to 1.99) no significant difference

Protein intake, %E ("during follow-up"): int 15.2 (SD 2.8), cont 15.7 (SD 3.4) (mean difference -0.50, 95% CI -1.09 to 0.09) no significant difference

Trans fat intake: not reported

Primary replacement for saturated fat: mainly PUFA (based on dietary aims and achievements)

Style: diet advice

Setting: community

Outcomes	Stated trial outcomes: cardiovascular mortality and morbidity Data available on total mortality? yes Cardiovascular mortality? yes (exact events included not stated) Events available for combined cardiovascular events: none Secondary outcomes: CHD deaths (exact events included not stated)
	Tertiary outcomes: total cholesterol, TG, BMI, sBP, dBP
Notes	Study duration 7 years
	Study aim was saturated fat 10%E, and saturated fat intake in the intervention group was less than 80% of that in the control (73%)
	SFA reduction aimed and achieved
	Total serum cholesterol, difference between intervention and control, mmol/L: -0.30 (95% CI -0.51 to -0.09), statistically significant reduction
	Trial dates: Recruitment 1966 to [unclear] and followed for 2 to 7 years
	Funding: Life Insurance Medical Research Fund of Australia and New Zealand



Sydney Diet-Heart 1978 (Continued)

Declarations of Interest of primary researchers: none stated, all authors worked for academic or health institutions.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence genera- tion (selection bias)	Low risk	Quote: "table of random numbers generated by a research assistant and was concealed until after medical evaluations and testing at baseline were completed".
Allocation concealment (selection bias)	Low risk	As above
Blinding of participants and personnel (perfor- mance bias) All outcomes	High risk	Very difficult to blind trials where participants need to make their own dietary changes
Blinding of outcome as- sessment (detection bias) CVD outcomes	Low risk	Initially masked to group assignment (though success of blinding not checked)
Blinding of outcome as- sessment (detection bias) All-cause mortality	Low risk	Blinding is not relevant in assessment of mortality.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Survival analysis used
Selective reporting (re- porting bias)	Low risk	Not relevant for primary and secondary outcomes as all trialists were asked for data
Free of systematic differ- ence in care?	High risk	Advice and follow-up in intervention group, not in control. See control and in- tervention methods in the Interventions section of the table of Characteristics of included studies
Stated aim to reduce SFA	Low risk	Aim to reduce SFA stated
Achieved SFA reduction	Low risk	SFA reduction achieved
Achieved TC reduction	Low risk	Statistically significant TC fall
Other bias	Low risk	None noted

Veterans Admin 1969

Study characteristics	
Methods	RCT
	Summary risk of bias: moderate to high
Participants	Men living at the Veterans Administration Center (USA) CVD risk: low



Veterans Admin 1969	(Continued)
	Control: randomised 422, analysed 422 Intervention: randomised 424, analysed 424 Mean years in trial: control 3.7, intervention 3.7 % male: 100 Age: mean control 65.6, intervention 65.4 (all 54 - 88)
	Ethnicity: White 90%, African-American 7%, Asian 1%, Mexican 1%, other 1%
	Statins use allowed? Unclear (only 4 participants were taking nicotinic acid, 17 diuretics, 56 digitalis, none on heparin)
	% taking statins: Not reported (probably none as too early, pre-1980)
Interventions	Modified fat vs usual diet
	Control aims: provided, total fat 40%E Intervention aims: total fat 40%E, ¾ of SFA replaced by unsaturated fats, dietary chol reduced
	Control methods: whole diet provided
	Intervention methods: whole diet provided
	Total fat intake, %E (during trial): int 38.9 (SD unclear), cont 40 (SD unclear) (mean difference -1.10, 95% CI -2.45 to 0.25 assuming SDs of 10) no significant difference
	Saturated fat intake, %E (during trial): int 8.3 (SD unclear), cont 18.5 (SD unclear) (mean differ- ence -10.20, 95% CI -10.87 to -9.53 assuming SDs of 5) significant reduction
	PUFA intake, %E (during trial)§: int 16.0 (SD ?), cont 4.9 (SD 0.10) (mean difference 11.10, 95% Cl 10.62 to 11.58 assuming missing SD was 5) significant increase
	PUFA n-3 intake: not reported
	PUFA n-6 intake: not reported
	MUFA intake, %E (during trial) *: not reported, approx int 14.0, cont 17.2 (mean difference -3.20, 95% CI -3.87 to -2.53) significant reduction
	CHO intake, %E (during trial) *: not reported, approx int 45.9, cont 44.8 (mean difference 1.10, 95% CI -1.60 to 3.80 assuming SDs of 20) no significant difference
	Protein intake, %E (during trial)§: int 15.2 (SD ?), cont 15.2 (SD ?) (mean difference 0.00, 95% CI -0.67 to 0.67 assuming SDs of 5) no significant difference
	Trans fat intake: not reported
	Replacement for saturated fat: mainly PUFA (based on dietary aims and achievements)
	Style: diet provided
	Setting: residential institution
	§Dayton 1965
	<pre>\$Estimated by subtraction (assuming total fat = SFA + PUFA + MUFA or energy intake = energy from fat + CHO + protein)</pre>
Outcomes	Stated trial outcomes: mortality, heart disease Data available on total mortality? yes Cardiovascular mortality? yes Events available for combined cardiovascular events: sudden death, definite MI, definite stroke, angina, PVD events
	Secondary outcomes: cancer deaths, cancer diagnoses, stroke, non-fatal MI, total MI, CHD deaths (fatal MI and sudden death due to CHD), CHD events (any MI or sudden death due to CHD)

Veterans Admin 1969 (Continued)

Tertiary outcomes: none (some data on total cholesterol, but no variance info)

Notes	Study duration over 8 years
	Study aim was to replace 66% of saturated fat by unsaturated fats, and saturated fat intake in the in- tervention group was significantly lower than in control
	SFA reduction aimed and achieved
	Total serum cholesterol, difference between intervention and control, mmol/L: -0.37 (95% CI -0.77 to 0.03), NO statistically significant reduction but reduction > 0.20
	Trial dates: Recruitment 1959 to 1967
	Funding: Veterans Administration, Arthur Dodd Fuller Foundation, National Heart Institute, Los Angeles County Heart Association, plus gifts of foods from Mazola corn oil and Mazola margarine, the National Soybean Processors Association, Pitman-Moore Company (Emdee margarine) and Hi-Saff Imitation Ice- cream from Frozen Desserts Company. Edgmar Farms donated milk refrigeration equipment.
	Declarations of Interest of primary researchers: none stated, all authors worked for academic or health institutions.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence genera- tion (selection bias)	Low risk	Quote: "table of random numbers used"
Allocation concealment (selection bias)	Low risk	Extensive baseline assessment before randomisation
Blinding of participants and personnel (perfor- mance bias) All outcomes	Low risk	Institution provided diet in a masked fashion.
Blinding of outcome as- sessment (detection bias) CVD outcomes	Low risk	Physician knowledge of allocation was assessed and found similar to random.
Blinding of outcome as- sessment (detection bias) All-cause mortality	Low risk	Blinding is not relevant in assessment of mortality.
Incomplete outcome data (attrition bias) All outcomes	Low risk	All followed up via Veterans Admin system
Selective reporting (re- porting bias)	Low risk	Not relevant for primary and secondary outcomes as all trialists were asked for data
Free of systematic differ- ence in care?	Low risk	All ate centre food as usual. See control and intervention methods in the Inter- ventions section of the table of Characteristics of included studies
Stated aim to reduce SFA	Low risk	Aim to reduce SFA stated
Achieved SFA reduction	Low risk	SFA reduction achieved
Achieved TC reduction	High risk	No statistically significant TC fall, though fall was > 0.20 mmol/L



Veterans Admin 1969 (Continued)

Other bias

Low risk

None noted

Study characteristics	S
Methods	RCT
	Women's Health Initiative (WHI)
	Summary risk of bias: low
Participants	Postmenopausal women aged 50 - 79 with or without CVD at baseline (USA) CVD risk: low in those without CVD at baseline, high in those with CVD Control without CVD at baseline: randomised 29,294, analysed 29,294 Intervention without CVD at baseline: randomised 19,541, analysed 19,541
	Control with CVD at baseline: randomised 1369, analysed 1369 Intervention with CVD at baseline: randomised 908, analysed 908 Mean years in trial: control 8.1, intervention 8.1 % male: 0 Age: mean (both with and without CVD at baseline) int 62.3 (SD 6.9), control 62.3 (SD 6.9)
	Ethnicity (women both with and without CVD at baseline): white 82%, black 11%, Asian or Pacific Islan- der 2%, unknown 1%, American Indian or Alaskan native < 1%. No statistically significant effects of the intervention on CHD events was seen for any ethnic subgroup.
	Statins use allowed? Yes
	% taking statins: 12% of women recruited were on lipid-lowering medication (these were a mixture of participants with and without CVD at baseline).
Interventions	Reduced fat vs usual diet
	Control: diet-related education materials Intervention: low-fat diet (20%E from fat), reduce saturated fat to 7%E with increased fruit and vegeta- bles
	Control methods: given copy of 'Dietary Guidelines for Americans'
	Intervention methods: 18 group sessions with trained and certified nutritionists in the 1st year, quarter- ly maintenance sessions thereafter, focusing on diet and behaviour modification
	Intervention delivered face-to-face in a group by nutritionists
	Intake data all relate to the full WHI cohort (not divided by whether participants have CVD at baseline or not)
	Total fat intake, %E (at 6 years): int 28.8 (SD 8.4), cont 37.0 (SD 7.3) (mean difference -8.20, 95% CI -8.34 to -8.06) significant reduction
	Saturated fat intake, %E (at 6 years): int 9.5 (SD3.2), cont 12.4 (SD3.1) (mean difference -2.90, 95% CI -2.96 to -2.84 for full WHI population) significant reduction
	PUFA intake, %E (at 6 years)§: int 6.3 (SD?), cont 7.6 (SD?) (mean difference -1.30, 95% CI -1.72 to -0.88 assuming missing SDs were 5) significant reduction
	PUFA n-3 intake: not reported
	PUFA n-6 intake: not reported



WHI 2006 (Continued)	MUFA intake, %E (at 6 years)§: int 11.1 (SD?), cont 14.3 (SD?) (mean difference -3.20, 95% CI -3.62 to
	CHO intake, %E (at 6 years)§: int 53.9 (SD?), cont 46.3 (SD?) (mean difference 7.60, 95% Cl 5.91 to 9.29 assuming SDs of 20) significant increase
	Protein intake, %E (at 6 years)§: int 17.7 (SD?), cont 17.0 (SD?) (mean difference 0.70, 95% CI 0.28 to 1.12 assuming SDs of 5) significant increase
	Trans fat intake, %E (at 6 years)§: int 1.8 (SD?), cont 2.4 (SD?) (mean difference unclear, no SDs as- sumed)
	Replacement for saturated fat: mainly carbohydrate, some protein (based on dietary achieve- ment)
	Style: dietary advice
	Setting: community
	§Amongst the 881 intervention and 1373 control participants with blood samples at baseline, with or without CVD at baseline (Howard 2010)
Outcomes	Stated trial outcomes: breast cancer, mortality, other cancers, cardiovascular events, diabetes
	Data available on total mortality? yes* Cardiovascular mortality? yes Events available for combined cardiovascular events: CHD, stroke, heart failure, angina, peripheral vas- cular disease, revascularisation, pulmonary embolism, DVT
	Secondary outcomes: cancer deaths*, cancer diagnoses*, stroke, non-fatal MI, diabetes diagnosis*
	Tertiary outcomes: weight, BMI, total, LDL and HDL cholesterol, TGs, systolic and diastolic BP (Lp(a) and HOMA reported as geometric means)
	* these are only available for the whole cohort, not split between low and high CVD risk groups
Notes	Study duration over 8 years
	Study aim was to reduce total fat to 20%E, reduce saturated fat to 7%E and increase fruit and veg- etable intake (Patterson 2003), and saturated fat intake in the intervention group was significantly low- er than in control
	SFA reduction aimed and achieved
	Total serum cholesterol, difference between intervention and control, mmol/L: -0.09 (95% CI -0.15 to -0.02), statistically significant reduction
	Trial dates: Recruitment was between 1993 and 1998
	Funding: National Heart, Lung and Blood Institute of the National Institutes of Health
	Declarations of Interest of primary researchers: Declarations varied from paper to paper, but this is a typical one from Beresford 2006 "Dr Black has received research grants from Pfizer and AstraZeneca, was on the speakers bureaus for Pfizer, Novartis, Sanofi-Aventis, Bristol-Meyers Squibb, Searle, Pharmacia, and Boehringer and served as a consultant of on an advisory board for Myogen, Merck Sharp and Dohme, Novartis, Mylan-Bertek, Pfizer, Bristol-Meyers Squibb, and Sanofi-Aventis. Dr Howard has served on the advisory boards of Merck, Schering Plough, and the Egg Nutrition Council, has received research support from Merck and Pfizer, and has consulted for General Mills. Dr Assaf is an employee of Pfizer. No other disclosures were reported."
Risk of bias	

Authors' judgement Support for judgement

Reduction in saturated fat intake for cardiovascular disease (Review) Copyright @ 2020 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

Bias

WHI 2006 (Continued)

Cochrane

Library

Trusted evidence.

Informed decisions. Better health.

Random sequence genera- tion (selection bias)	Low risk	Computer generated permuted block algorithm stratified by clinical centre and age
Allocation concealment (selection bias)	Low risk	Allocations developed by the WHI Clinical Coordinating Center
Blinding of participants and personnel (perfor- mance bias) All outcomes	High risk	Participants aware of allocation
Blinding of outcome as- sessment (detection bias) CVD outcomes	Low risk	Trained clinic staff, who were responsible for anthropometric assessments and administration of FFQs, were blinded to treatment assignments to the extent practical. The dietary intervention staff did not conduct clinical assessments, and clinic staff were not permitted to participate in any intervention activities; participants were instructed not to discuss nutrition activities with clinic staff.
Blinding of outcome as- sessment (detection bias) All-cause mortality	Low risk	Blinding not relevant for mortality assessment
Incomplete outcome data (attrition bias) All outcomes	Low risk	ITT analysis
Selective reporting (re- porting bias)	Low risk	Trials register 1999, study completion 2005, but outcomes not stated in trials register. However, outcomes were well published; trialists were asked for data.
Free of systematic differ- ence in care?	High risk	Intervention participants received 18 group sessions with behavioural mod- ification plus quarterly maintenance sessions thereafter; control groups re- ceived a leaflet. See control and intervention methods in the Interventions section of the table of Characteristics of included studies
Stated aim to reduce SFA	Low risk	Aim to reduce SFA stated
Achieved SFA reduction	Low risk	SFA reduction achieved
Achieved TC reduction	Low risk	Statistically significant TC fall
Other bias	Low risk	None noted

WINS 2006

Study characteristics	
Methods	RCT
	Women's Intervention Nutrition Study (WINS)
	Summary risk of bias: low
Participants	Women with localised resected breast cancer (USA) CVD risk: low
	Control: 1462 randomised, 1462 analysed



WINS 2006 (Continued)	Intervention: 975 randomised, 975 analysed
	Mean years in trial: overall 5.0 % men: 0
	Age: control mean 58.5 (95% CI 43.6 to 73.4), intervention mean 58.6 (95% CI 44.4 to 72.8) (all post- menopausal)
	Ethnicity: 85% white, 5% black, 4% Hispanic, 5% Asian or Pacific Islander, < 1% American Indian or un- known (no outcome data based on ethnicity)
	Statins use allowed? Not stated (statins not mentioned in inclusion or exclusion criteria within trial pro- tocol)
	% taking statins: Not reported
Interventions	Reduced fat intake vs usual diet
	Control aims: minimal nutritional counselling focused on nutritional adequacy Intervention aims: total fat 15 - 20%E
	Control methods: 1 baseline dietetic session plus 3-monthly sessions
	Intervention methods: 8 bi-weekly individual dietetic sessions plus 3-monthly contact and optional monthly group sessions, incorporating individual fat gram goals, social cognitive theory, self-monitor- ing, goal-setting, modelling, social support and relapse prevention and management
	Intervention was delivered face-to-face individually by trained dietitian
	Total fat intake, %E (at 1 year): int 20.3 (SD 8.1), cont 29.2 (SD 7.4) (mean difference -8.90, 95% CI -9.53 to -8.27)
	Total fat intake, %E (at 5 years): int 23.2 (SD 8.4) n = 380, cont 31.2 (SD 8.9) n = 648 (mean difference -8.00, 95% CI -9.09 to -6.91) significant reduction
	Saturated fat intake*, %E (at 1 year): int 6.4 (SD 0.14 [4.4]), cont 9.8 (SD 0.15 [5.7]) (mean differ- ence -3.40, 95% CI -3.80 to -3.00 assuming reported SDs were actually SEs) significant reduction
	PUFA intake*, %E (at 1 year): int 4.5 (SD 0.09 (2.8)), cont 6.4 (SD 0.10 (3.8)) (mean difference -1.90, 95% CI -2.16 to -1.64) significant reduction
	PUFA n-3 intake: not reported by study arm
	PUFA n-6 intake: not reported by study arm
	MUFA intake*, %E (at 1 year): int 7.6 (SD 0.14 (4.4)), cont 11.5 (SD 0.16 (6.1)) (mean difference -3.90, 95% CI -4.32 to -3.48) significant reduction
	CHO intake, %E (at 6 months): int 60.8 (SD 19.6), cont 50.5 (SD 14.8) (mean difference 10.30, 95% CI 8.85 to 11.75) significant increase
	Protein intake, %E (at 6 months): int 19.1 (SD 5.2), cont 17.6 (SD 4.1) (mean difference 1.50, 95% CI 1.11 to 1.89) significant increase
	Trans fat intake: not reported
	Replacement for saturated fat: CHO and protein (based on dietary achievement)
	Style: dietary advice
	Setting: community
	*SDs appear incorrect, probably SEs?
Outcomes	Stated trial outcomes: dietary fat intake, total cholesterol, weight and waist measurement



WINS 2006 (Continued)	Data available on total mortality? yes Cardiovascular mortality? no Events available for combined cardiovascular events: none Secondary outcomes: cancer diagnoses Tertiary outcomes: weight, BMI, total cholesterol
Notes	Study duration 5 years
	Study aim was to reduce total fat to 15 - 20%E
	SFA reduction achieved
	Total serum cholesterol, difference between intervention and control, mmol/L: -0.14 (95% CI -0.34 to 0.05), NO statistically significant reduction and reduction < 0.20
	Trial dates: Recruitment 1994 to 2001
	Funding: National Cancer Institute, Breast Cancer Research Foundation, American Institute for Cancer Research
	Declarations of Interest of primary researchers: none stated, all authors worked for academic or health institutions except that Njeri Karanja worked for Kaiser Permanente Center for Health Research, Bette Caan for Kaiser Permanente Medical Group, and Barbara L Winters for Campbell's Soup Company.
Risk of bias	

Bias	Authors' judgement	Support for judgement
Random sequence genera- tion (selection bias)	Low risk	Random stratified permuted block design, carried out at the statistical coordi- nating centre of WINS
Allocation concealment (selection bias)	Low risk	Random stratified permuted block design, carried out at the statistical coordi- nating centre of WINS
Blinding of participants and personnel (perfor- mance bias) All outcomes	High risk	Not for dietary advice and participants
Blinding of outcome as- sessment (detection bias) CVD outcomes	Low risk	All outcomes assessed by the blinded outcome committee
Blinding of outcome as- sessment (detection bias) All-cause mortality	Low risk	Outcome assessors blinded
Incomplete outcome data (attrition bias) All outcomes	Low risk	All assessed
Selective reporting (re- porting bias)	Low risk	Not relevant for primary and secondary outcomes as all trialists were asked for data
Free of systematic differ- ence in care?	High risk	Differences in attention - more time for those in intervention group. See con- trol and intervention methods in the Interventions section of the table of Char- acteristics of included studies


WINS 2006 (Continued)

Stated aim to reduce SFA	High risk	Aim to reduce SFA not stated
Achieved SFA reduction	Low risk	SFA reduction achieved
Achieved TC reduction	High risk	No statistically significant TC fall
Other bias	Low risk	None noted

 $\% \mathsf{E}:$ percent of total energy intake

ATPII: Adult treatment panel II BMI: body mass index (weight in kg/ height in m, squared) BP: blood pressure CABG: coronary artery bypass graft CHD: coronary heart disease CHO: carbohydrate chol: cholesterol CI: confidence interval cont: control group CVD: cardiovascular disease DART: Diet And Reinfarction Trial dBP: diastolic blood pressure DVT: deep vein thrombosis EPA: eicosapentaenoic acid GPs: general practitioners HDL: high density lipoprotein HOMA: homeostatic model assessment int: intervention group ITT: Intention to treat analysis LDL: low density lipoprotein Lp(a): lipoprotein (a) MI: myocardial infarction MRC: Medical Research Council MUFA: monounsaturated fat P/S: polyunsaturated/saturated fat ratio PCTA: percutaneous transluminal coronary angioplasty PUFA: polyunsaturated fat PVD: peripheral vascular disease RCT: randomised controlled trial sBP: systolic blood pressure SD: standard deviation SE: standard error SFA: saturated fats STARS: St Thomas' Atherosclerosis Regression Study TC: total cholesterol TG: triglyceride vs: versus WHI: Women's Health Initiative WINS: Women's Intervention Nutrition Study

Characteristics of excluded studies [ordered by study ID]



Study	Reason for exclusion
Agewall 2001	Multifactorial intervention
Ammerman 2003	No appropriate control group (and not low fat vs modified fat)
Anderson 1990	Follow-up less than 24 months
Aquilani 2000	No appropriate control group (and not low fat vs modified fat)
Arntzenius 1985	No appropriate control group (and not low fat vs modified fat)
Aro 1990	Intervention and randomised follow-up less than 6 months
ASSIST 2001	Intervention was not dietary fat modification or low fat diet.
Australian Polyp Prev 95	Neither mortality nor cardiovascular morbidity data available (only decided after contact with at least 1 author)
Azadbakht 2007	Follow-up less than 24 months
Bakx 1997	Multifactorial intervention
Ball 1965	Study aim was to assess effects of a low-fat diet and methods stated that the "nature of the fat con- sumed was not altered". Saturated fat content of diet was not reported.
Barnard 2009	Weight reduction encouraged in the conventional diet, but not in the vegan diet arm
Barndt 1977	No appropriate control group (and not low fat vs modified fat)
Baron 1990	Multifactorial intervention
Barr 1990	Intervention and randomised follow-up less than 6 months
Barsotti 1991	Complex paper in Italian; unclear whether cardiovascular events occurred; contact with authors not established
Baumann 1982	Intervention and randomised follow-up less than 6 months
BDIT Pilot Studies 1996	Study aim was to reduce total fat intake to 15%E with no specific intervention on saturated fat. Sat- urated fat in intervention group was more than 80% of that in the control group.
Beckmann 1995	Intervention was not dietary fat modification or low-fat diet.
beFIT 1997	Follow-up less than 24 months
Beresford 1992	Intervention and randomised follow-up less than 6 months
Bergstrom 1967	Intervention and randomised follow-up less than 6 months
Bierenbaum 1963	No appropriate control group (and not low fat vs modified fat)
Bloemberg 1991	Neither mortality nor cardiovascular morbidity data available (only decided after contact with at least 1 author)
Bloomgarden 1987	Multifactorial intervention



Study	Reason for exclusion
Bonk 1975	Trial, unclear if randomised; contact could not be established with trialists
Bonnema 1995	No appropriate control group (and not low fat vs modified fat)
Bosaeus 1992	Intervention and randomised follow-up less than 6 months
Boyd 1988	Follow-up less than 24 months
BREACPNT	Individual microbiome-based dietary advice vs Mediterranean diet (no suggestion of saturated fat reduction in either arm)
Brehm 2009	Unclear whether any relevant events occurred, not able to contact trialists
Brensike 1982	No appropriate control group (and not low fat vs modified fat)
BRIDGES 2001	Follow-up less than 24 months
Broekmans 2003	Intervention was not dietary fat modification or low fat diet.
Brown 1984	No appropriate control group (and not low fat vs modified fat)
Bruce 1994	No appropriate control group (and not low fat vs modified fat)
Bruno 1983	Multifactorial intervention
Butcher 1990	Intervention and randomised follow-up less than 6 months
Byers 1995	No appropriate control group (and not low fat vs modified fat)
Caggiula 1996	No appropriate control group (and not low fat vs modified fat)
Canadian DBCP 1997	Unable to establish contact with authors to provide data on numbers of deaths and CV events
CARMEN 2000	Follow-up less than 24 months
CARMEN substudy 2002	Follow-up less than 24 months
Casas-Agustench 2013	Less than 24 months duration
Cerin 1993	Intervention and randomised follow-up less than 6 months
Chan 1993	Intervention and randomised follow-up less than 6 months
Chapman 1950	Intervention and randomised follow-up less than 6 months
Charbonnier 1975	Intervention and randomised follow-up less than 6 months
Cheng 2004	Intervention and randomised follow-up less than 6 months
Chiostri 1988	Intervention and randomised follow-up less than 6 months
Choudhury 1984	Intervention and randomised follow-up less than 6 months
Clark 1997	Multifactorial intervention



Study	Reason for exclusion
Clifton 1992	Intervention and randomised follow-up less than 6 months
Cobb 1991	Intervention and randomised follow-up less than 6 months
Cohen 1991	Intervention was not dietary fat modification or low fat diet.
Cole 1988	Intervention and randomised follow-up less than 6 months
Colquhoun 1990	Intervention and randomised follow-up less than 6 months
Consolazio 1946	Intervention and randomised follow-up less than 6 months
Cox 1996	Multifactorial intervention
Croft 1986	Intervention was not dietary fat modification or low fat diet.
Curzio 1989	Follow-up less than 24 months
Da Qing IGT 1997	Intervention was not dietary fat modification or low-fat diet.
Dalgard 2001	No appropriate control group (and not low fat vs modified fat)
DAS 2000	No appropriate control group (and not low fat vs modified fat)
DASH 1997	Intervention and randomised follow-up less than 6 months
Davey Smith 2005	Multifactorial intervention
De Boer 1983	Intervention and randomised follow-up less than 6 months
De Bont 1981	Neither mortality nor cardiovascular morbidity data available as study data have been lost
DeBusk 1994	Multifactorial intervention
DEER 1998	Duration 1 year only
Delahanty 2001	No appropriate control group (and not low fat vs modified fat)
Delius 1969	Intervention was not dietary fat modification or low fat diet.
Demark 1990	Intervention and randomised follow-up less than 6 months
Dengel 1995	No appropriate control group (and not low fat vs modified fat)
Denke 1994	Intervention and randomised follow-up less than 6 months
Diabetes CCT 1995	Intervention was not dietary fat modification or low fat diet.
Diet & Hormone Study 2003	Duration 1 year only
DIET 1998	Multifactorial intervention
Ding 1992	Intervention and randomised follow-up less than 6 months
DIPI 2018	Less than 24 months duration

Study	Reason for exclusion
DIRECT 2009	Unable to establish contact with authors to establish whether relevant events occurred; multiple publications checked and no relevant outcomes found
DO IT 2006	Intervention aim was for a "mediterranean diet" with total fat 27 - 30%E, protein 15 - 18%E, CHO 50 - 55%E, no specific aim to reduce saturated fat (though polyunsaturated margarine given to inter- vention group), and intervention group saturated fat was more than 80% of that in the control.
Dobs 1991	No appropriate control group (and not low fat vs modified fat)
Due 2008	Follow-up less than 24 months
Duffield 1982	Multifactorial intervention
Dullaart 1992	Study authors confirmed that no deaths or cardiovascular events occurred during the study.
Eating Patterns 1997	Neither mortality nor cardiovascular morbidity data available (only decided after contact with at least 1 author)
Ehnholm 1982	Intervention and randomised follow-up less than 6 months
Ehnholm 1984	Intervention and randomised follow-up less than 6 months
Eisenberg 1990	Intervention and randomised follow-up less than 6 months
Elder 2000	No appropriate control group (and not low fat vs modified fat)
Ellegard 1991	Intervention and randomised follow-up less than 6 months
Esposito 2003	No appropriate control group (and not low fat vs modified fat)
Esposito 2004	Unable to establish contact with authors to assess whether any relevant events occurred
EUROACTION 2008	Multifactorial intervention
FARIS 1997	Multifactorial intervention
Fasting HGS 1997	No appropriate control group (and not low fat vs modified fat)
Ferrara 2000	No appropriate control group (and not low fat vs modified fat)
Fielding 1995	Intervention and randomised follow-up less than 6 months
Finnish Diabet Prev 2000	Multifactorial intervention
Finnish Mental Hosp 1972	Not randomised (cluster-randomised, but < 6 clusters)
Fisher 1981	Intervention and randomised follow-up less than 6 months
FIT Heart 2011	Authors confirmed that differences between intervention and control groups included smoking and physical activity, as well as dietary changes.
Fleming 2002	No appropriate control group (and not low fat vs modified fat)
Fortmann 1988	Intervention was not dietary fat modification or low fat diet.



Study	Reason for exclusion
Foster 2003	Weight reduction in 1 arm but not the other
Frenkiel 1986	Follow-up less than 24 months
FRESH START 2007	Participants were newly diagnosed with cancer.
Gambera 1995	Intervention and randomised follow-up less than 6 months
Gaullier 2007	No appropriate control group (and not low fat vs modified fat)
Ginsberg 1988	Intervention and randomised follow-up less than 6 months
Gjone 1972	Intervention and randomised follow-up less than 6 months
Glatzel 1966	No appropriate control group (and not low fat vs modified fat)
Goodpaster 1999	No appropriate control group (and not low fat vs modified fat)
Grundy 1986	Intervention and randomised follow-up less than 6 months
Hardcastle 2008	Multifactorial intervention
Harris 1990	Intervention and randomised follow-up less than 6 months
Hartman 1993	No appropriate control group (and not low fat vs modified fat)
Hartwell 1986	No appropriate control group (and not low fat vs modified fat)
Hashim 1960	Intervention and randomised follow-up less than 6 months
Haufe 2011	Aim was to reduce total fat or reduce carbohydrate, but no saturated fat aims were stated, and ef- fects of the diets on saturated fat intakes were unclear.
Haynes 1984	Intervention was not dietary fat modification or low fat diet.
Heber 1991	Intervention and randomised follow-up less than 6 months
Heine 1989	Neither mortality nor cardiovascular morbidity data available (only decided after contact with at least 1 author)
Hellenius 1995	The study aimed for weight loss in 1 arm and not in the comparison arm.
Heller 1993	Neither mortality nor cardiovascular morbidity data available (only decided after contact with at least 1 author)
Hildreth 1951	No appropriate control group (and not low fat vs modified fat)
Holm 1990	Neither mortality nor cardiovascular morbidity data available (only decided after contact with at least 1 author)
Horlick 1957	Intervention and randomised follow-up less than 6 months
Horlick 1960	Intervention and randomised follow-up less than 6 months
Howard 1977	Intervention and randomised follow-up less than 6 months

Reduction in saturated fat intake for cardiovascular disease (Review)



Study	Reason for exclusion
Hunninghake 1990	Intervention and randomised follow-up less than 6 months
Hutchison 1983	No appropriate control group (and not low fat vs modified fat)
Hyman 1998	Neither mortality nor cardiovascular morbidity data available (only decided after contact with at least 1 author)
lacono 1981	Not randomised; Intervention and randomised follow-up less than 6 months
IMPACT 1995	Multifactorial intervention
lso 1991	No appropriate control group (and not low fat vs modified fat)
lves 1993	Multifactorial intervention
Jalkanen 1991	Multifactorial intervention
Jerusalem Nut 1992	Intervention and randomised follow-up less than 6 months
Jula 1990	Multifactorial intervention
Junker 2001	Intervention and randomised follow-up less than 6 months
Karmally 1990	Intervention and randomised follow-up less than 6 months
Karvetti 1992	Multifactorial intervention
Kastarinen 2002	Multifactorial intervention
Kather 1985	Intervention and randomised follow-up less than 6 months
Katzel 1995	Intervention was not dietary fat modification or low fat diet.
Kawamura 1993	Intervention and randomised follow-up less than 6 months
Keidar 1988	Intervention and randomised follow-up less than 6 months
Kempner 1948	No appropriate control group (and not low fat vs modified fat)
Keys 1957a	Intervention and randomised follow-up less than 6 months
Keys 1957b	Intervention and randomised follow-up less than 6 months
Keys 1957c	Intervention and randomised follow-up less than 6 months
Khan 2003	Neither mortality nor cardiovascular morbidity data available (only decided after contact with at least 1 author)
King 2000	Intervention and randomised follow-up less than 6 months
Kingsbury 1961	Intervention and randomised follow-up less than 6 months
KNOTA	Numerous publications checked, but no relevant outcome data found. Trialists not contacted.
Koopman 1990	Intervention and randomised follow-up less than 6 months

Reduction in saturated fat intake for cardiovascular disease (Review)



Study	Reason for exclusion
Koranyi 1963	Unclear whether randomised, unable to contact authors to discuss
Korhonen 2003	Multifactorial intervention
Kriketos 2001	Intervention and randomised follow-up less than 6 months
Kris 1994	Intervention and randomised follow-up less than 6 months
Kristal 1997	Multifactorial intervention
Kromhout 1987	No appropriate control group (and not low fat vs modified fat)
Kummel 2008	Intervention was not dietary fat modification or low-fat diet.
Laitinen 1993	Multifactorial intervention
Laitinen 1994	Multifactorial intervention
Lean 1997	Follow-up less than 24 months
Leduc 1994	Multifactorial intervention
Lewis 1958	Intervention and randomised follow-up less than 6 months
Lewis 1981	Intervention and randomised follow-up less than 6 months
Lewis 1985	Multifactorial intervention
Lichtenstein 2002	Intervention and randomised follow-up less than 6 months
Lim 2010	Unable to establish contact with authors to gain access to data on health outcomes (none reported in paper)
Linko 1957	Intervention and randomised follow-up less than 6 months
Lipid Res Clinic 1984	No appropriate control group (and not low fat vs modified fat)
Little 1990	Intervention and randomised follow-up less than 6 months
Little 2004	Intervention was not dietary fat modification or low-fat diet.
Lottenberg 1996	Intervention and randomised follow-up less than 6 months
Luszczynska 2007	No appropriate control group (and not low fat vs modified fat)
Lyon Diet Heart 1994	Intervention was not dietary fat modification or low-fat diet.
Lysikova 2003	Intervention and randomised follow-up less than 6 months
Macdonald 1972	Intervention and randomised follow-up less than 6 months
Mansel 1990	Intervention was not dietary fat modification or low-fat diet
MARGARIN 2002	No appropriate control group (and not low fat vs modified fat)



Study	Reason for exclusion
Marniemi 1990	Both intervention groups aimed to lose weight, while the control group did not.
Mattson 1985	Intervention and randomised follow-up less than 6 months
McAuley 2005	Follow-up less than 24 months
McCarron 1997	Intervention and randomised follow-up less than 6 months
McCarron 2001	Intervention was not dietary fat modification or low-fat diet.
McKeown-Eyssen 1994	Intervention aim was to reduce total fat and increase dietary fibre (saturated fat not mentioned), and no saturated fat intakes during trial reported.
McManus 2001	Neither mortality nor cardiovascular morbidity data available (only decided after contact with at least 1 author)
McNamara 1981	Intervention and randomised follow-up less than 6 months
Medi-RIVAGE 2004	Weight reduction for some low-fat diet participants (those with BMI > 25) but not in Mediterranean group
MeDiet 2002	Follow-up less than 24 months
MEDINA	Less than 24 months duration
Mensink 1987	Intervention and randomised follow-up less than 6 months
Mensink 1989	Intervention and randomised follow-up less than 6 months
Mensink 1990a	Intervention and randomised follow-up less than 6 months
Mensink 1990b	Intervention and randomised follow-up less than 6 months
Metroville Health 2003	Unable to establish contact with authors to assess whether any relevant events occurred
Michalsen 2006	Diet plus stress management vs no intervention
Miettinen 1994	Intervention and randomised follow-up less than 6 months
Millar 1973	No appropriate control group (and not low fat vs modified fat)
Miller 1998	Intervention and randomised follow-up less than 6 months
Miller 2001	Neither mortality nor cardiovascular morbidity data available (only decided after contact with at least 1 author)
Milne 1994	No appropriate control group (and not low fat vs modified fat) - the high CHO diet was neither 'usu- al' or 'low fat' to compare with the modified fat diet.
Minnesota Coronary 1989	Although the study proceeded for over 4 years, participants (patients) came and went and mean follow-up was only 1 year.
Minnesota HHP 1990	No appropriate control group (and not low fat vs modified fat)
Mojonnier 1980	Unable to establish contact with authors to assess whether any relevant events occurred

Reduction in saturated fat intake for cardiovascular disease (Review)



Study	Reason for exclusion
Mokuno 1988	Intervention and randomised follow-up less than 6 months
Mortensen 1983	Intervention and randomised follow-up less than 6 months
Mottalib 2018	Less than 24 months duration
MRFIT substudy 1986	Intervention and randomised follow-up less than 6 months
MSDELTA 1995	Intervention and randomised follow-up less than 6 months
MSFAT 1997	Follow-up less than 24 months
Mujeres Felices 2003	Diet and breast self-examination vs no intervention
Mutanen 1997	Intervention and randomised follow-up less than 6 months
Muzio 2007	Intervention and randomised follow-up less than 6 months
Naglak 2000	Unable to establish contact with authors to assess whether any relevant events occurred
NAS 1987	Intervention and randomised follow-up less than 6 months
National Diet Heart 1968	Follow-up less than 24 months
NCEP weight 1991	Neither mortality nor cardiovascular morbidity data available (only decided after contact with at least 1 author)
NCT01954472	Study withdrawn (not completed)
NCT03068078	Less than 24 months duration
Neil 1995	No appropriate control group (and not low fat vs modified fat)
Neverov 1997	Multifactorial intervention
Next Step 1995	Neither mortality nor cardiovascular morbidity data available (only decided after contact with at least 1 author)
Nordoy 1971	Intervention and randomised follow-up less than 6 months
Norway Veg Oil 1968	No appropriate control group (and not low fat vs modified fat)
Nutri-AGEs 2015	Less than 24 months duration
Nutrition Breast Health	Follow-up less than 24 months
O'Brien 1976	Intervention and randomised follow-up less than 6 months
ODES 2006	The study aimed for weight loss in 1 arm and not in the other arm.
Oldroyd 2001	Multifactorial intervention
Ole Study 2002	Follow-up less than 24 months
OLIVE 1997	Unable to establish contact with authors to assess whether any relevant events occurred



Study	Reason for exclusion
ORIGIN 2008	Intervention was not dietary fat modification or low-fat diet.
Oslo Study 2004	Multifactorial intervention
Pascale 1995	Multifactorial intervention
PEP 2001	Multifactorial intervention
PHYLLIS 1993	No appropriate control group (and not low fat vs modified fat)
Pilkington 1960	Neither mortality nor cardiovascular morbidity data available (only decided after contact with at least 1 author)
Polyp Prevention 1996	Intervention aim was to reduce total fat and increase dietary fibre, fruit and vegetables (saturated fat not mentioned), and no saturated fat intakes during trial reported.
POUNDS LOST 2009	All study arms (low or high total fat) prescribed low saturated fat intake (8%E); no usual fat com- parator.
PREDIMED 2008	Total fat goals in the low-fat arm were unclear and authors confirmed that aims were nonspecific (if aims < 30%E, this study would be included).
PREMIER 2003	Follow-up less than 24 months
Pritchard 2002	The study aimed for weight loss in 1 arm and not in the comparison arm.
Puget Sound EP 2000	Neither mortality nor cardiovascular morbidity data available (only decided after contact with at least 1 author)
Rabast 1979	Intervention and randomised follow-up less than 6 months
Rabkin 1981	Intervention and randomised follow-up less than 6 months
Radack 1990	Intervention and randomised follow-up less than 6 months
Rasmussen 1995	Intervention and randomised follow-up less than 6 months
Reaven 2001	Intervention and randomised follow-up less than 6 months
Reid 2002	No appropriate control group (and not low fat vs modified fat)
Renaud 1986	Not randomised
Rivellese 1994	Follow-up less than 24 months
Rivellese 2003	Intervention and randomised follow-up less than 6 months
Roderick 1997	Neither mortality nor cardiovascular morbidity data available (only decided after contact with at least 1 author)
Roman CHD prev 1986	Multifactorial intervention
Rose 1987	No appropriate control group (and not low fat vs modified fat)
Sarkkinen 1995	Follow-up less than 24 months

Reduction in saturated fat intake for cardiovascular disease (Review)



Study	Reason for exclusion
Schaefer 1995a	Intervention and randomised follow-up less than 6 months
Schaefer 1995b	Intervention and randomised follow-up less than 6 months
Schectman 1996	Multifactorial intervention
Schlierf 1995	Multifactorial intervention
Seppanen-Laakso 1992	Intervention and randomised follow-up less than 6 months
Seppelt 1996	Follow-up less than 24 months
Singh 1991	Multifactorial intervention
Singh 1992	No appropriate control group (and not low fat vs modified fat)
Sirtori 1992	Intervention and randomised follow-up less than 6 months
SLIM 2008	Multifactorial intervention
Sopotsinskaia 1992	The study aimed for weight loss in 1 arm and not in the comparison arm.
Soul Food Light	Less than 24 months duration
Stanford NAP 1997	Intervention and randomised follow-up less than 6 months
Stanford Weight 1994	The study aimed for weight loss in 1 arm and not in the comparison arm.
Starmans 1995	Intervention and randomised follow-up less than 6 months
Steinbach 1996	Multifactorial intervention
Steptoe 2001	No appropriate control group (and not low fat vs modified fat)
Stevens 2002	Diet plus breast self examination vs no intervention
Stevenson 1988	No appropriate control group (and not low fat vs modified fat)
Strychar 2009	Follow-up less than 24 months
Sweeney 2004	Intervention was not dietary fat modification or low fat diet.
Søndergaard 2003	Follow-up less than 24 months
TAIM 1992	Intervention was not dietary fat modification or low fat diet.
Tapsell 2004	Unable to establish contact with authors to assess whether any relevant events occurred
THIS DIET 2008	All study arms prescribed low saturated fat intake, no usual fat comparator
TOHP 1992	Multifactorial intervention
TONE 1997	Intervention was not dietary fat modification or low-fat diet.
Toobert 2003	Multifactorial intervention

Reduction in saturated fat intake for cardiovascular disease (Review)



Study	Reason for exclusion
Towle 1994	Intervention and randomised follow-up less than 6 months
TRANSFACT 2006	Intervention and randomised follow-up less than 6 months
Treatwell 1992	Neither mortality nor cardiovascular morbidity data available (only decided after contact with at least 1 author)
Tromsø Heart 1989	Multifactorial intervention
Troyer 2010	Longest duration only 12 months
UK PDS 1996	No appropriate control group (and not low fat vs modified fat)
Urbach 1952	No appropriate control group (and not low fat vs modified fat)
Uusitupa 1993	Multifactorial intervention
VASTKOST 2012	Publications reported than no participants died or experienced CVD during the trial.
Vavrikova 1958	Intervention and randomised follow-up less than 6 months
Verheiden 2003	Unable to establish contact with authors to assess whether any relevant events occurred
WAHA 2016	15%E from walnuts vs usual diet (neither arm aimed to reduce saturated fat intake)
Wass 1981	Intervention and randomised follow-up less than 6 months
Wassertheil 1985	Intervention was not dietary fat modification or low fat diet.
WATCH 1999	Neither mortality nor cardiovascular morbidity data available (only decided after contact with at least 1 author)
Watts 1988	Intervention and randomised follow-up less than 6 months
Weintraub 1992	No appropriate control group (and not low fat vs modified fat)
Westman 2006	Intervention was not dietary fat modification or low fat diet.
Weststrate 1998	Intervention and randomised follow-up less than 6 months
WHEL 2007	Study aimed to reduce total fat, but saturated fat goals were not mentioned, and saturated fat in- take in the intervention group was more than 80% of that in the control (81%).
WHO primary prev 1979	Multifactorial intervention
WHT 1990	Neither mortality nor cardiovascular morbidity data available as such data were not collected in the study
WHT Feasibility 2003	Neither mortality nor cardiovascular morbidity data available (only decided after contact with at least 1 author)
Wilke 1974	Intervention and randomised follow-up less than 6 months
Williams 1990	Intervention was not dietary fat modification or low-fat diet.



Study	Reason for exclusion					
Williams 1992	Intervention was not dietary fat modification or low-fat diet.					
Williams 1994	Intervention was not dietary fat modification or low-fat diet.					
Wilmot 1952	No appropriate control group (and not low fat vs modified fat)					
Wing 1998	No appropriate control group (and not low fat vs modified fat)					
WINS UK 2011	Stated aim was to reduce total fat by 50%; no saturated fat aims					
WOMAN 2007	Lifestyle intervention included exercise and weight as well as diet.					
Wood 1988	Intervention was not dietary fat modification or low-fat diet.					
Woollard 2003	Multifactorial intervention including smoking, weight, exercise and alcohol components					
Working Well 1996	Multifactorial intervention					
Zock 1995	Intervention and randomised follow-up less than 6 months					

CHO: carbohydrate CV: cardiovascular E: energy vs: versus

Characteristics of studies awaiting classification [ordered by study ID]

ICFAMED

Methods	A Mediterranean diet for preventing heart failure and atrial fibrillation in hypertensive patients (IC- FAMED)				
	RCT, 24 months				
Participants	People with hypertension aged 55 to 75 years at high cardiovascular risk, but without existing CVD				
Interventions	MedDiet: Mediterranean-style diet, dietary advice (individual and group) every three months LFD: Low-fat diet according to American Heart Association guidelines, dietary advice (individual and group) every three months				
Outcomes	Primary: heart failure and/or atrial fibrillation				
	Secondary: echocardiographic variables & BP variables				
	Actual outcomes from abstracts: MedDiet: 5 CVD events (atrial fibrillation (AF) 2; ischaemic heart disease (IHD) 2; stroke 1), LFD: 11 CVD events (AF 6, IHD 2, stroke 3). The crude rate for the occurrence of events per 1000 patient-months of follow-up was 197 (95% CI: 06–46) for MedDiet, 451 (95% CI: 3–8.1) for LFD. The HR for patients with MedDiet compared to LFD was 044 (95% CI: 015–126, P > 005).				
Notes	Trials registration: ISRCTN27497769				
	Enrollment began in 2012; appears to have completed in 2017; abstract and poster publications on- ly to date				
	Awaiting assessment because: Unclear whether one arm was higher in saturated fat than the other, awaiting fuller publication to assess				

Reduction in saturated fat intake for cardiovascular disease (Review)



AF: atrial fibrillation CVD: cardiovascular disease HR: hazard ratio ICFAMED: A Mediterranean diet for preventing heart failure and atrial fibrillation in hypertensive patients IHD: Ischaemic heart disease LFD: low fat diet MedDiet: Mediterranean style diet RCT: randomised controlled trial

Characteristics of ongoing studies [ordered by study ID]

ENAbLE due unclear

Study name	ENAbLE				
Methods	RCT, 2 x 2 diet and physical activity interventions, duration unclear				
Participants	Stroke survivors able to walk independently				
Interventions	AusMed diet, adaptation of the Mediterranean diet to the Australian context, including provision of starter foods, menu plans and regular counselling				
	Comparator unclear				
	Telehealth-delivered physical activity and diet interventions in both arms				
Outcomes	Primary: sBP				
	Secondary: lipid profiles and glycaemic control				
Starting date	Mid 2019, planned completion date unclear				
Contact information	Coralie English, University of Western Australia (first author of abstract)				
Notes	Trial registration not found				
	Unclear whether the intervention was truly lower vs higher saturated fat as saturated fat goals not provided, and duration unclear				

NCT02481466 due 2020

Study name	Combined Portfolio diet and Exercise study (PortfolioEx)				
Methods	RCT, 2 x 2 factorial design with exercise intervention, 36 months				
Participants	Men and postmenopausal women with BMI up to 40 kg/m 2 with measurable arterial thickening				
Interventions	Lower saturated fat: advice on a therapeutic diet appropriate for hypercholesterolemia (ie < 7% of energy from saturated fat, < 200 mg/d cholesterol) PLUS the combination of viscous fibres, soy protein, plant sterols and nuts, 5% extra monounsaturated fat, and selection of low glycemic index foods				
	Higher saturated fat: advice to follow a DASH-like diet of whole grains, and low-fat dairy products with fruits and vegetables				
	Both arms with or without instruction on the Laval exercise program — a standardised physical ac- tivity/exercise component supervised by trained kinesiologists (exercise physiologists)				

Reduction in saturated fat intake for cardiovascular disease (Review)

NCT02481466 due 2020 (Continued)

Outcomes	Primary: maximum vessel wall volume of the carotid arteries				
	Secondary: composite end point of myocardial infarction, revascularization, cardiovascular hospi- talisation, cardiovascular mortality and stroke; atrial fibrillation; BP; and vessel outcomes				
Starting date	Nov 2016, planned completion Dec 2022				
Contact information	PI: David J Jenkins, MD, NutritionProject@smh.ca, Risk Factor Modification Centre, St. Michael's Hospital				
Notes	Trials registration: NCT02481466				
	Unclear whether the intervention was truly lower vs higher saturated fat as saturated fat goals not provided for both arms				

NCT02938832 due 2023	
Study name	Does the advice to eat a mediterranean diet with low carbohydrate intake, compared with a low-fat diet, reduce diabetes and cardiovascular disease (CardioDiet)
Methods	RCT, 36 months
Participants	Adults with ischaemic heart disease followed up at cardiac rehabilitation units
Interventions	Mediterranean diet with an energy content (E%) from carbohydrates between 25-30%
	Traditional low-fat diet with 45-60 E% from carbohydrates
Outcomes	Primary: diabetes incidence
	Secondary: CVD disease, quality of life
Starting date	Oct 2016, planned completion Oct 2023
Contact information	PI: Fredrik H Nystrom, Professor, MD, University Hospital, Linkoeping, fredrik.nystrom@regionos- tergotland.se
Notes	Trials registration NCT02938832
	Unclear whether the intervention was truly lower vs higher saturated fat as saturated fat goals not provided.

NEW Soul Study due 2022	
Study name	Nutritious Eating With Soul (NEW Soul) study
Methods	RCT, 24 months
Participants	African-American adults aged 18-65 years with BMI 25- 49.9 kg/m ²
Interventions	Lower saturated fat: plant-based vegan diet, instructing participants to favour a diet built around whole grains, fruits, vegetables, and legumes, supplemented by the Oldways African Heritage and Health programme, which includes a food pyramid guide. A Taste of African Heritage (ATAH) six-les- son nutrition and cooking programme has an online course for health professionals and cooking



NEW Soul Study due 2022 (Continued)

instructors (all research and restaurant team members will complete this course). Interventions include intervention meetings, physical activity, and podcasts/mailings.

Higher saturated fat: low-fat omnivorous diet, supplemented by the Oldways African Heritage and Health programme, which includes a food pyramid guide. A Taste of African Heritage (ATAH) six-lesson nutrition and cooking programme has an online course for health professionals and cooking instructors (all research and restaurant team members will complete this course). Interventions include intervention meetings, physical activity, and podcasts/mailings.

Outcomes	Primary: CVD events				
	Secondary: CVD risk factors (including LDL & BP), body weight				
Starting date	May 2018, planned completion June 2022				
Contact information	PI: Brie Turner-McGrievy, Associate Professor, University of South Carolina				
	Trial website: https://newsoul.org/				
Notes	Trials registration NCT03354377				
	Unclear whether the intervention was truly lower vs higher saturated fat as saturated fat goals not provided				

ATAH: A Taste of African Heritage AusMed: Australian style Mediterranean diet BMI: body mass index BP: blood pressure CVD: cardiovascular disease DASH: Dietary Approaches to Stop Hypertension E: energy LDL: low density lipoprotein PorffolioEx: Combined Portfolio diet and Exercise study RCT: randomised controlled trial sBP: systolic blood pressure

DATA AND ANALYSES

Comparison 1. SFA reduction vs usual diet - primary outcomes

Outcome or subgroup title	No. of studies	No. of partici- pants	Statistical method	Effect size
1.1 ALL-CAUSE MORTALITY	12	55858	Risk Ratio (M-H, Random, 95% CI)	0.96 [0.90, 1.03]
1.2 All-cause mortality, SA low summary risk of bias	7	53219	Risk Ratio (M-H, Random, 95% Cl)	0.95 [0.84, 1.08]
1.3 All-cause mortality, SA aim to reduce SFA	9	53112	Risk Ratio (M-H, Random, 95% Cl)	0.97 [0.89, 1.06]
1.4 All-cause mortality, SA statistically signif- icant SFA reduction	8	54973	Risk Ratio (M-H, Random, 95% Cl)	0.98 [0.92, 1.04]

Reduction in saturated fat intake for cardiovascular disease (Review)

Copyright $\ensuremath{\mathbb S}$ 2020 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.



Cochrane Database of Systematic Reviews

Outcome or subgroup title	No. of studies	No. of partici- pants	Statistical method	Effect size
1.5 All-cause mortality, SA TC reduction	8	53073	Risk Ratio (M-H, Random, 95% CI)	0.97 [0.88, 1.07]
1.6 All-cause mortality, SA excluding WHI	11	7023	Risk Ratio (M-H, Random, 95% CI)	0.95 [0.83, 1.07]
1.7 All-cause mortality, SA Mantel-Haenszel fixed-effect	12	55858	Risk Ratio (M-H, Fixed, 95% CI)	0.97 [0.91, 1.03]
1.8 All-cause mortality, SA Peto fixed-effect	12	55858	Peto Odds Ratio (Peto, Fixed, 95% CI)	0.96 [0.90, 1.04]
1.9 All-cause mortality, subgroup by any substitution	12		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.9.1 replaced by PUFA	7	4238	Risk Ratio (M-H, Random, 95% CI)	0.96 [0.82, 1.13]
1.9.2 replaced by MUFA	1	52	Risk Ratio (M-H, Random, 95% CI)	3.00 [0.33, 26.99]
1.9.3 replaced by CHO	6	53669	Risk Ratio (M-H, Random, 95% CI)	0.97 [0.90, 1.04]
1.9.4 replaced by protein	5	53614	Risk Ratio (M-H, Random, 95% CI)	0.97 [0.90, 1.04]
1.9.5 replacement unclear	0	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
1.10 All-cause mortality, subgroup by main substitution	12		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.10.1 replaced by PUFA	6	4183	Risk Ratio (M-H, Random, 95% CI)	0.97 [0.82, 1.14]
1.10.2 replaced by MUFA	1	52	Risk Ratio (M-H, Random, 95% CI)	3.00 [0.33, 26.99]
1.10.3 replaced by CHO	5	51636	Risk Ratio (M-H, Random, 95% CI)	0.97 [0.90, 1.04]
1.10.4 replaced by protein	0	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
1.10.5 replacement unclear	0	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
1.11 All-cause mortality, subgroup by dura- tion	12	55858	Risk Ratio (M-H, Random, 95% CI)	0.96 [0.90, 1.03]
1.11.1 up to 24mo	4	2246	Risk Ratio (M-H, Random, 95% CI)	0.99 [0.78, 1.26]

Reduction in saturated fat intake for cardiovascular disease (Review)



Cochrane Database of Systematic Reviews

Outcome or subgroup title	No. of studies	No. of partici- pants	Statistical method	Effect size
1.11.2 >24 to 48mo	3	1294	Risk Ratio (M-H, Random, 95% Cl)	0.96 [0.83, 1.12]
1.11.3 >48mo	4	52142	Risk Ratio (M-H, Random, 95% Cl)	0.96 [0.79, 1.16]
1.11.4 unclear duration	1	176	Risk Ratio (M-H, Random, 95% Cl)	0.33 [0.07, 1.61]
1.12 All-cause mortality, subgroup by base- line SFA	12	55858	Risk Ratio (M-H, Random, 95% Cl)	0.96 [0.90, 1.03]
1.12.1 up to 12%E SFA baseline	1	2437	Risk Ratio (M-H, Random, 95% Cl)	0.90 [0.67, 1.21]
1.12.2 >12 to 15%E SFA baseline	5	51635	Risk Ratio (M-H, Random, 95% Cl)	1.01 [0.86, 1.19]
1.12.3 >15 to 18%E SFA baseline	1	55	Risk Ratio (M-H, Random, 95% Cl)	0.35 [0.04, 3.12]
1.12.4 >18%E SFA baseline	1	846	Risk Ratio (M-H, Random, 95% Cl)	0.98 [0.83, 1.15]
1.12.5 unclear	4	885	Risk Ratio (M-H, Random, 95% Cl)	0.80 [0.62, 1.04]
1.13 All-cause mortality, subgroup by SFA change	12	55858	Risk Ratio (M-H, Random, 95% Cl)	0.96 [0.90, 1.03]
1.13.1 up to 4%E difference	5	53939	Risk Ratio (M-H, Random, 95% Cl)	0.99 [0.86, 1.13]
1.13.2 >4 to 8%E difference	2	188	Risk Ratio (M-H, Random, 95% Cl)	0.41 [0.08, 2.07]
1.13.3 >8%E difference	1	846	Risk Ratio (M-H, Random, 95% Cl)	0.98 [0.83, 1.15]
1.13.4 unclear	4	885	Risk Ratio (M-H, Random, 95% Cl)	0.80 [0.62, 1.04]
1.14 All-cause mortality, subgroup by sex	12	55858	Risk Ratio (M-H, Random, 95% Cl)	0.96 [0.90, 1.03]
1.14.1 Men	9	4410	Risk Ratio (M-H, Random, 95% Cl)	0.96 [0.83, 1.11]
1.14.2 Women	2	51272	Risk Ratio (M-H, Random, 95% Cl)	0.97 [0.90, 1.05]
1.14.3 Mixed, men and women	1	176	Risk Ratio (M-H, Random, 95% Cl)	0.33 [0.07, 1.61]

Reduction in saturated fat intake for cardiovascular disease (Review)



Cochrane Database of Systematic Reviews

Outcome or subgroup title	No. of studies	No. of partici- pants	Statistical method	Effect size
1.15 All-cause mortality, subgroup by CVD risk	12	55858	Risk Ratio (M-H, Random, 95% CI)	0.96 [0.90, 1.03]
1.15.1 Low CVD risk	4	52251	Risk Ratio (M-H, Random, 95% CI)	0.97 [0.91, 1.04]
1.15.2 Moderate CVD risk	1	176	Risk Ratio (M-H, Random, 95% CI)	0.33 [0.07, 1.61]
1.15.3 Existing CVD disease	7	3431	Risk Ratio (M-H, Random, 95% Cl)	0.97 [0.76, 1.24]
1.16 All-cause mortality, subgroup by TC re- duction	12		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.16.1 serum chol reduced by at least 0.2mmol/L	7	4238	Risk Ratio (M-H, Random, 95% Cl)	0.96 [0.81, 1.14]
1.16.2 serum chol reduced by <0.2mmol/L	4	51487	Risk Ratio (M-H, Random, 95% Cl)	0.97 [0.90, 1.04]
1.16.3 serum chol reduction unclear	1	133	Risk Ratio (M-H, Random, 95% Cl)	0.51 [0.05, 5.46]
1.17 All-cause mortality, subgroup decade of publication	12	55858	Risk Ratio (M-H, Random, 95% Cl)	0.96 [0.90, 1.03]
1.17.1 1960s	5	1731	Risk Ratio (M-H, Random, 95% Cl)	0.92 [0.80, 1.07]
1.17.2 1970s	1	458	Risk Ratio (M-H, Random, 95% Cl)	1.49 [0.95, 2.34]
1.17.3 1980s	1	2033	Risk Ratio (M-H, Random, 95% Cl)	0.98 [0.76, 1.25]
1.17.4 1990s	2	188	Risk Ratio (M-H, Random, 95% Cl)	0.41 [0.08, 2.07]
1.17.5 2000s	3	51448	Risk Ratio (M-H, Random, 95% Cl)	0.97 [0.88, 1.05]
1.18 CARDIOVASCULAR MORTALITY	11	53421	Risk Ratio (M-H, Random, 95% Cl)	0.94 [0.78, 1.13]
1.19 CVD mortality, SA low summary risk of bias	4	50315	Risk Ratio (M-H, Random, 95% Cl)	0.96 [0.67, 1.38]
1.20 CVD mortality, SA aim to reduce SFA	9	53112	Risk Ratio (M-H, Random, 95% Cl)	0.95 [0.79, 1.14]
1.21 CVD mortality, SA statistically signifi- cant SFA reduction	7	52536	Risk Ratio (M-H, Random, 95% Cl)	0.95 [0.75, 1.21]

Reduction in saturated fat intake for cardiovascular disease (Review)



Cochrane Database of Systematic Reviews

Outcome or subgroup title	No. of studies	No. of partici- pants	Statistical method	Effect size
1.22 CVD mortality, SA TC reduction	8	53073	Risk Ratio (M-H, Random, 95% CI)	0.95 [0.78, 1.15]
1.23 CVD mortality, SA excluding WHI	10	4586	Risk Ratio (M-H, Random, 95% CI)	0.92 [0.72, 1.18]
1.24 CVD mortality, SA Mantel-Haenszel fixed-effect	11	53421	Risk Ratio (M-H, Fixed, 95% Cl)	0.95 [0.85, 1.07]
1.25 CVD mortality, SA Peto fixed-effect	11	53421	Peto Odds Ratio (Peto, Fixed, 95% CI)	0.95 [0.84, 1.08]
1.26 CVD mortality, subgroup by any substi- tution	11		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.26.1 replaced by PUFA	7	4251	Risk Ratio (M-H, Random, 95% CI)	0.95 [0.73, 1.25]
1.26.2 replaced by MUFA	1	52	Risk Ratio (M-H, Random, 95% CI)	3.00 [0.33, 26.99]
1.26.3 replace by CHO	5	51232	Risk Ratio (M-H, Random, 95% CI)	0.99 [0.85, 1.14]
1.26.4 replaced by protein	4	51177	Risk Ratio (M-H, Random, 95% CI)	0.99 [0.86, 1.14]
1.26.5 replacement unclear	0	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
1.27 CVD mortality, subgroup by main sub- stitution	11		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.27.1 replaced by PUFA	6	4196	Risk Ratio (M-H, Random, 95% CI)	0.97 [0.73, 1.28]
1.27.2 replaced by MUFA	1	52	Risk Ratio (M-H, Random, 95% CI)	3.00 [0.33, 26.99]
1.27.3 replace by CHO	4	49199	Risk Ratio (M-H, Random, 95% CI)	0.78 [0.42, 1.46]
1.27.4 replaced by protein	0	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
1.27.5 replacement unclear	0	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
1.28 CVD mortality, subgroup by duration	11	53447	Risk Ratio (M-H, Random, 95% CI)	0.95 [0.78, 1.16]
1.28.1 up to 24mo	4	2272	Risk Ratio (M-H, Random, 95% CI)	1.26 [0.54, 2.94]

Reduction in saturated fat intake for cardiovascular disease (Review)



Cochrane Database of Systematic Reviews

Outcome or subgroup title	No. of studies	No. of partici- pants	Statistical method	Effect size
1.28.2 >24 to 48mo	3	1294	Risk Ratio (M-H, Random, 95% CI)	0.79 [0.57, 1.08]
1.28.3 >48 mo	3	49705	Risk Ratio (M-H, Random, 95% CI)	1.02 [0.73, 1.43]
1.28.4 unclear duration	1	176	Risk Ratio (M-H, Random, 95% CI)	0.25 [0.03, 2.19]
1.29 CVD mortality, subgroup by baseline SFA	11	53447	Risk Ratio (M-H, Random, 95% CI)	0.95 [0.78, 1.16]
1.29.1 up to 12%E SFA baseline	0	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
1.29.2 >12 to 15%E SFA baseline	5	51635	Risk Ratio (M-H, Random, 95% CI)	1.06 [0.84, 1.32]
1.29.3 >15 to 18%E SFA baseline	1	55	Risk Ratio (M-H, Random, 95% CI)	0.35 [0.04, 3.12]
1.29.4 >18%E SFA baseline	1	846	Risk Ratio (M-H, Random, 95% CI)	0.70 [0.51, 0.96]
1.29.5 unclear	4	911	Risk Ratio (M-H, Random, 95% CI)	1.00 [0.61, 1.66]
1.30 CVD mortality, subgroup by SFA change	11	53447	Risk Ratio (M-H, Random, 95% CI)	0.95 [0.78, 1.16]
1.30.1 up to 4%E difference	4	51502	Risk Ratio (M-H, Random, 95% CI)	1.07 [0.85, 1.33]
1.30.2 >4 to 8%E difference	2	188	Risk Ratio (M-H, Random, 95% CI)	0.29 [0.05, 1.70]
1.30.3 >8%E difference	1	846	Risk Ratio (M-H, Random, 95% CI)	0.70 [0.51, 0.96]
1.30.4 unclear	4	911	Risk Ratio (M-H, Random, 95% CI)	1.00 [0.61, 1.66]
1.31 CVD mortality, subgroup by sex	11	53447	Risk Ratio (M-H, Random, 95% CI)	0.95 [0.78, 1.16]
1.31.1 Men	9	4436	Risk Ratio (M-H, Random, 95% CI)	0.96 [0.73, 1.25]
1.31.2 Women	1	48835	Risk Ratio (M-H, Random, 95% CI)	1.00 [0.84, 1.19]
1.31.3 Mixed, men and women	1	176	Risk Ratio (M-H, Random, 95% CI)	0.25 [0.03, 2.19]

Reduction in saturated fat intake for cardiovascular disease (Review)



Cochrane Database of Systematic Reviews

Outcome or subgroup title	No. of studies	No. of partici- pants	Statistical method	Effect size
1.32 CVD mortality, subgroup by CVD risk	11	53447	Risk Ratio (M-H, Random, 95% CI)	0.96 [0.80, 1.14]
1.32.1 Low CVD risk	3	47537	Risk Ratio (M-H, Random, 95% CI)	0.84 [0.60, 1.16]
1.32.2 Moderate CVD risk	1	176	Risk Ratio (M-H, Random, 95% CI)	0.25 [0.03, 2.19]
1.32.3 Existing CVD disease	8	5734	Risk Ratio (M-H, Random, 95% CI)	1.04 [0.83, 1.31]
1.33 CVD mortality, subgroup by TC reduc- tion	11		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.33.1 serum chol reduced by at least 0.2mmol/L	7	4251	Risk Ratio (M-H, Random, 95% CI)	0.95 [0.73, 1.25]
1.33.2 serum chol reduced by <0.2mmol/L	3	49063	Risk Ratio (M-H, Random, 95% Cl)	0.97 [0.47, 2.01]
1.33.3 serum chol reduction unclear	1	133	Risk Ratio (M-H, Random, 95% Cl)	0.20 [0.01, 4.15]
1.34 CVD mortality, subgroup decade of publication	11	53421	Risk Ratio (M-H, Random, 95% Cl)	0.94 [0.78, 1.13]
1.34.1 1960s	5	1731	Risk Ratio (M-H, Random, 95% Cl)	0.78 [0.63, 0.97]
1.34.2 1970s	1	458	Risk Ratio (M-H, Random, 95% CI)	1.59 [0.99, 2.55]
1.34.3 1980s	1	2033	Risk Ratio (M-H, Random, 95% Cl)	1.01 [0.77, 1.31]
1.34.4 1990s	2	188	Risk Ratio (M-H, Random, 95% Cl)	0.29 [0.05, 1.70]
1.34.5 2000s	2	49011	Risk Ratio (M-H, Random, 95% Cl)	0.78 [0.27, 2.21]
1.35 COMBINED CARDIOVASCULAR EVENTS	12	53300	Risk Ratio (M-H, Random, 95% Cl)	0.79 [0.66, 0.93]
1.36 CVD events, SA low summary risk of bias	3	49857	Risk Ratio (M-H, Random, 95% Cl)	0.89 [0.75, 1.07]
1.37 CVD events, SA aim to reduce SFA	10	52991	Risk Ratio (M-H, Random, 95% Cl)	0.79 [0.66, 0.95]
1.38 CVD events, SA statistically significant SFA reduction	7	52313	Risk Ratio (M-H, Random, 95% Cl)	0.85 [0.71, 1.02]

Reduction in saturated fat intake for cardiovascular disease (Review)



Cochrane Database of Systematic Reviews

Outcome or subgroup title	No. of studies	No. of partici- pants	Statistical method	Effect size
1.39 CVD events, SA TC reduction	9	52952	Risk Ratio (M-H, Random, 95% CI)	0.78 [0.65, 0.94]
1.40 CVD events, SA excluding WHI	11	4465	Risk Ratio (M-H, Random, 95% CI)	0.75 [0.61, 0.91]
1.41 CVD events, SA Mantel-Haenszel fixed- effect	12	53300	Risk Ratio (M-H, Fixed, 95% CI)	0.93 [0.88, 0.99]
1.42 CVD events, SA Peto fixed-effect	12	53300	Peto Odds Ratio (Peto, Fixed, 95% CI)	0.92 [0.87, 0.98]
1.43 CVD events, SA excluding trials with ad- ditional interventions	9	3998	Risk Ratio (M-H, Random, 95% CI)	0.80 [0.64, 0.99]
1.44 CVD events, subgroup by any substitu- tion	12		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.44.1 replaced by PUFA	7	3895	Risk Ratio (M-H, Random, 95% CI)	0.73 [0.58, 0.92]
1.44.2 replaced by MUFA	1	52	Risk Ratio (M-H, Random, 95% CI)	1.00 [0.53, 1.89]
1.44.3 replace by CHO	5	51232	Risk Ratio (M-H, Random, 95% Cl)	0.84 [0.67, 1.06]
1.44.4 replaced by protein	4	51177	Risk Ratio (M-H, Random, 95% CI)	0.97 [0.91, 1.03]
1.44.5 replacement unclear	1	235	Risk Ratio (M-H, Random, 95% CI)	1.68 [0.41, 6.87]
1.45 CVD events, subgroup by main substitu- tion	12		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.45.1 replaced by PUFA	6	3840	Risk Ratio (M-H, Random, 95% CI)	0.78 [0.62, 0.97]
1.45.2 replaced by MUFA	1	52	Risk Ratio (M-H, Random, 95% CI)	1.00 [0.53, 1.89]
1.45.3 replace by CHO	4	49199	Risk Ratio (M-H, Random, 95% CI)	0.67 [0.39, 1.16]
1.45.4 replaced by protein	0	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
1.45.5 replacement unclear	1	235	Risk Ratio (M-H, Random, 95% CI)	1.68 [0.41, 6.87]
1.46 CVD events, subgroup by duration	12	53300	Risk Ratio (M-H, Random, 95% CI)	0.79 [0.66, 0.93]

Reduction in saturated fat intake for cardiovascular disease (Review)



Cochrane Database of Systematic Reviews

Outcome or subgroup title	No. of studies	No. of partici- pants	Statistical method	Effect size	
1.46.1 up to 24mo	5	2481	Risk Ratio (M-H, Random, 95% CI)	0.96 [0.78, 1.16]	
1.46.2 >24 to 48mo	3	1294	Risk Ratio (M-H, Random, 95% CI)	0.73 [0.56, 0.95]	
1.46.3 >48mo	2	49247	Risk Ratio (M-H, Random, 95% Cl)	0.85 [0.63, 1.16]	
1.46.4 unclear duration	2	278	Risk Ratio (M-H, Random, 95% Cl)	0.43 [0.17, 1.08]	
1.47 CVD events, subgroup by baseline SFA	12	53300	Risk Ratio (M-H, Random, 95% CI)	0.79 [0.66, 0.93]	
1.47.1 up to 12%E SFA baseline	0	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable	
1.47.2 >12 to 15%E SFA baseline	5	51412	Risk Ratio (M-H, Random, 95% Cl)	0.97 [0.91, 1.03]	
1.47.3 >15 to 18%E SFA baseline	1	55	Risk Ratio (M-H, Random, 95% CI)	0.41 [0.22, 0.78]	
1.47.4 >18%E SFA baseline	1	846	Risk Ratio (M-H, Random, 95% Cl)	0.79 [0.63, 1.00]	
1.47.5 unclear	5	987	Risk Ratio (M-H, Random, 95% Cl)	0.72 [0.51, 1.03]	
1.48 CVD events, subgroup by SFA change	12	53300	Risk Ratio (M-H, Random, 95% Cl)	0.79 [0.66, 0.93]	
1.48.1 up to 4%E difference	4	51279	Risk Ratio (M-H, Random, 95% Cl)	0.97 [0.91, 1.03]	
1.48.2 >4 to 8%E difference	2	188	Risk Ratio (M-H, Random, 95% Cl)	0.40 [0.22, 0.74]	
1.48.3 >8%E difference	1	846	Risk Ratio (M-H, Random, 95% Cl)	0.79 [0.63, 1.00]	
1.48.4 unclear	5	987	Risk Ratio (M-H, Random, 95% Cl)	0.72 [0.51, 1.03]	
1.49 CVD events, subgroup by sex	12	53300	Risk Ratio (M-H, Random, 95% Cl)	0.79 [0.66, 0.93]	
1.49.1 Men	8	3952	Risk Ratio (M-H, Random, 95% Cl)	0.80 [0.69, 0.93]	
1.49.2 Women	1	48835	Risk Ratio (M-H, Random, 95% Cl)	0.98 [0.92, 1.04]	

Reduction in saturated fat intake for cardiovascular disease (Review)



Cochrane Database of Systematic Reviews

Outcome or subgroup title	No. of studies	No. of partici- pants	Statistical method	Effect size
1.49.3 Mixed, men and women	3	513	Risk Ratio (M-H, Random, 95% CI)	0.59 [0.23, 1.49]
1.50 CVD events, subgroup by CVD risk	12	53300	Risk Ratio (M-H, Random, 95% CI)	0.83 [0.72, 0.96]
1.50.1 Low CVD risk	3	47537	Risk Ratio (M-H, Random, 95% CI)	0.89 [0.75, 1.06]
1.50.2 Moderate CVD risk	3	513	Risk Ratio (M-H, Random, 95% CI)	0.59 [0.23, 1.49]
1.50.3 Existing CVD disease	7	5250	Risk Ratio (M-H, Random, 95% CI)	0.86 [0.71, 1.05]
1.51 CVD events, subgroup by TC reduction	12		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.51.1 serum chol reduced by at least 0.2mmol/L	8	4117	Risk Ratio (M-H, Random, 95% CI)	0.74 [0.59, 0.92]
1.51.2 serum chol reduced by <0.2mmol/L	3	49050	Risk Ratio (M-H, Random, 95% CI)	0.98 [0.91, 1.04]
1.51.3 serum chol reduction unclear	1	133	Risk Ratio (M-H, Random, 95% CI)	0.20 [0.01, 4.15]
1.52 CVD events, subgroup decade of publi- cation	12	53300	Risk Ratio (M-H, Random, 95% CI)	0.79 [0.66, 0.93]
1.52.1 1960s	5	1731	Risk Ratio (M-H, Random, 95% Cl)	0.79 [0.69, 0.91]
1.52.2 1970s	1	102	Risk Ratio (M-H, Random, 95% Cl)	0.27 [0.14, 0.52]
1.52.3 1980s	1	2033	Risk Ratio (M-H, Random, 95% Cl)	0.92 [0.74, 1.15]
1.52.4 1990s	2	188	Risk Ratio (M-H, Random, 95% Cl)	0.40 [0.22, 0.74]
1.52.5 2000s	3	49246	Risk Ratio (M-H, Random, 95% CI)	0.98 [0.91, 1.04]

Analysis 1.1. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 1: ALL-CAUSE MORTALITY

	lower	lower SFA		higher SFA		Risk Ratio	Risk Ratio	
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI	
Black 1994	1	66	2	67	0.1%	0.51 [0.05 , 5.46]	
DART 1989	111	1018	113	1015	7.6%	0.98 [0.76 , 1.25	1 🔶	
Ley 2004	2	88	6	88	0.2%	0.33 [0.07 , 1.61]	
MRC 1968	28	199	31	194	2.1%	0.88 [0.55, 1.41	1 _	
Oslo Diet-Heart 1966	48	206	65	206	4.6%	0.74 [0.54 , 1.02	1 _	
Rose corn oil 1965	5	28	0	13	0.1%	5.31 [0.32, 89.44]	
Rose olive 1965	3	26	1	13	0.1%	1.50 [0.17 , 13.05]	
STARS 1992	1	27	3	28	0.1%	0.35 [0.04 , 3.12]	
Sydney Diet-Heart 1978	39	221	28	237	2.3%	1.49 [0.95 , 2.34]	
Veterans Admin 1969	174	424	177	422	17.4%	0.98 [0.83 , 1.15	1 🔶	
WHI 2006 (1)	989	19541	1520	29294	60.2%	0.98 [0.90 , 1.05] 🙀	
WINS 2006 (2)	64	975	107	1462	5.2%	0.90 [0.67 , 1.21	ı -	
Total (95% CI)		22819		33039	100.0%	0.96 [0.90 , 1.03		
Total events:	1465		2053					
Heterogeneity: $Tau^2 = 0.00$; Chi ² = 11.25	, df = 11 (P = 0.42; I	$^{2} = 2\%$			0.01 0.1 1 10	100
Test for overall effect: Z =	1.01 (P = 0.31)	l)					Favours lower SFA Favours his	gher SFA

Test for subgroup differences: Not applicable

Footnotes

(1) All-cause death during study, Prentice 2017

(2) All-cause mortality during trial, Chlebowski 2015

Analysis 1.2. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 2: All-cause mortality, SA low summary risk of bias

	lower	lower SFA		higher SFA		Risk Ratio	Ris	k Ratio
Study or Subgroup	Events Total		Events	Total	Weight	M-H, Random, 95% CI	M-H, Ran	dom, 95% CI
Ley 2004	2	88	6	88	0.6%	0.33 [0.07 , 1.61]	
Oslo Diet-Heart 1966	48	206	65	206	11.5%	0.74 [0.54 , 1.02].	-
STARS 1992	1	27	3	28	0.3%	0.35 [0.04 , 3.12]	<u> </u>
Sydney Diet-Heart 1978	39	221	28	237	6.6%	1.49 [0.95 , 2.34]	
Veterans Admin 1969	174	424	177	422	27.0%	0.98 [0.83 , 1.15]	•
WHI 2006 (1)	989	19541	1520	29294	41.3%	0.98 [0.90 , 1.05]	
WINS 2006	64	975	107	1462	12.7%	0.90 [0.67 , 1.21]	-
Total (95% CI)		21482		31737	100.0%	0.95 [0.84 , 1.08]	
Total events:	1317		1906					
Heterogeneity: Tau ² = 0.01;	Chi ² = 9.25,	df = 6 (P =	= 0.16); I ² =	: 35%			0.01 0.1	1 10 100
Fest for overall effect: $Z = 0.77 (P = 0.44)$							Favours lower SFA	Favours higher SFA

Test for subgroup differences: Not applicable

Footnotes

(1) All-cause death during study, Prentice 2017



Analysis 1.3. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 3: All-cause mortality, SA aim to reduce SFA

	lower	SFA	higher SFA		Risk Ratio		Risk	Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Rano	lom, 95% CI
DART 1989	111	1018	113	1015	10.8%	0.98 [0.76 , 1.25]]	•
MRC 1968	28	199	31	194	3.2%	0.88 [0.55, 1.41]	_	-
Oslo Diet-Heart 1966	48	206	65	206	6.8%	0.74 [0.54, 1.02]	_	+
Rose corn oil 1965	5	28	0	13	0.1%	5.31 [0.32, 89.44]		
Rose olive 1965	3	26	1	13	0.2%	1.50 [0.17, 13.05]]	
STARS 1992	1	27	3	28	0.2%	0.35 [0.04 , 3.12]		
Sydney Diet-Heart 1978	39	221	28	237	3.5%	1.49 [0.95 , 2.34]]	
Veterans Admin 1969	174	424	177	422	22.2%	0.98 [0.83 , 1.15]]	•
WHI 2006 (1)	989	19541	1520	29294	53.1%	0.98 [0.90 , 1.05]]	•
Total (95% CI)		21690		31422	100.0%	0.97 [0.89 , 1.06]]	
Total events:	1398		1938					
Heterogeneity: Tau ² = 0.00;	Chi ² = 8.94,	df = 8 (P =	= 0.35); I ² =	:11%			0.01 0.1	1 10 100
Test for overall effect: $Z = 0$	0.69 (P = 0.49)]	Favours lower SFA	Favours higher SFA
Test for subgroup difference	es: Not applic	able						

Footnotes

(1) All-cause death during study, Prentice 2017

Analysis 1.4. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 4: All-cause mortality, SA statistically significant SFA reduction

	lower	SFA	higher SFA			Risk Ratio	Risk	Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Rand	om, 95% CI
Black 1994	1	66	2	67	0.1%	0.51 [0.05 , 5.46]	
DART 1989	111	1018	113	1015	6.9%	0.98 [0.76 , 1.25] .	•
Ley 2004	2	88	6	88	0.2%	0.33 [0.07 , 1.61]	L
STARS 1992	1	27	3	28	0.1%	0.35 [0.04 , 3.12]	
Sydney Diet-Heart 1978	39	221	28	237	2.1%	1.49 [0.95 , 2.34]	
Veterans Admin 1969	174	424	177	422	16.5%	0.98 [0.83 , 1.15] .	•
WHI 2006 (1)	989	19541	1520	29294	69.4%	0.98 [0.90 , 1.05]	
WINS 2006	64	975	107	1462	4.7%	0.90 [0.67 , 1.21] _	Ŧ
Total (95% CI)		22360		32613	100.0%	0.98 [0.92 , 1.04	1	
Total events:	1381		1956					
Heterogeneity: $Tau^2 = 0.00$; Chi ² = 6.69,	df = 7 (P =	= 0.46); I ² =	= 0%			0.01 0.1	1 10 100
Test for overall effect: Z =	0.68 (P = 0.50)))					Favours lower SFA	Favours higher SFA
Test fear and second difference	aa. Natannlia	abla						

Test for subgroup differences: Not applicable

Footnotes

(1) All-cause death during study, Prentice 2017



Analysis 1.5. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 5: All-cause mortality, SA TC reduction

	lower SFA		higher SFA		Risk Ratio		Ris	k Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Ran	dom, 95% CI
DART 1989	111	1018	113	1015	12.8%	0.98 [0.76 , 1.25	·]	+
MRC 1968	28	199	31	194	4.1%	0.88 [0.55, 1.41] .	•
Oslo Diet-Heart 1966	48	206	65	206	8.3%	0.74 [0.54 , 1.02	.]	•
Rose corn oil 1965	5	28	0	13	0.1%	5.31 [0.32, 89.44	·]	
STARS 1992	1	27	3	28	0.2%	0.35 [0.04 , 3.12	.]	
Sydney Diet-Heart 1978	39	221	28	237	4.5%	1.49 [0.95 , 2.34	.]	
Veterans Admin 1969	174	424	177	422	23.9%	0.98 [0.83 , 1.15]	•
WHI 2006 (1)	989	19541	1520	29294	46.0%	0.98 [0.90 , 1.05	5]	•
Total (95% CI)		21664		31409	100.0%	0.97 [0.88 , 1.07	']	
Total events:	1395		1937					
Heterogeneity: Tau ² = 0.00;	Chi ² = 8.79,	df = 7 (P =	= 0.27); I ² =	= 20%			0.01 0.1	1 10 100
Test for overall effect: $Z = 0$	0.63 (P = 0.53)	5)					Favours lower SFA	Favours higher SFA

Test for subgroup differences: Not applicable

Footnotes

(1) All-cause death during study, Prentice 2017

Analysis 1.6. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 6: All-cause mortality, SA excluding WHI

	lower	lower SFA		higher SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
Black 1994	1	66	2	67	0.3%	0.51 [0.05 , 5.46]	I
DART 1989	111	1018	113	1015	20.0%	0.98 [0.76 , 1.25]	⊢ ∔
Ley 2004	2	88	6	88	0.6%	0.33 [0.07 , 1.61]	I
MRC 1968	28	199	31	194	6.6%	0.88 [0.55, 1.41]	∣
Oslo Diet-Heart 1966	48	206	65	206	13.2%	0.74 [0.54, 1.02]	I -
Rose corn oil 1965	5	28	0	13	0.2%	5.31 [0.32, 89.44]	I
Rose olive 1965	3	26	1	13	0.3%	1.50 [0.17, 13.05]	I
STARS 1992	1	27	3	28	0.3%	0.35 [0.04 , 3.12]	I
Sydney Diet-Heart 1978	39	221	28	237	7.2%	1.49 [0.95 , 2.34]	I
Veterans Admin 1969	174	424	177	422	36.6%	0.98 [0.83 , 1.15]	
WINS 2006	64	975	107	1462	14.7%	0.90 [0.67 , 1.21]	• •
Total (95% CI)		3278		3745	100.0%	0.95 [0.83 , 1.07]	
Total events:	476		533				
Heterogeneity: $Tau^2 = 0.00$; Chi ² = 11.10	df = 10 (P = 0.35; I	$^{2} = 10\%$			0.01 0.1 1 10 100
Test for overall effect: Z =	0.87 (P = 0.39))				1	Favours lower SFA Favours higher SI
							-

Test for subgroup differences: Not applicable



Analysis 1.7. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 7: All-cause mortality, SA Mantel-Haenszel fixed-effect

	lower	SFA	higher	SFA		Risk Ratio	Ris	k Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Fixed, 95% CI	M-H, Fiz	xed, 95% CI
Black 1994	1	66	2	67	0.1%	0.51 [0.05 , 5.46	5]	
DART 1989	111	1018	113	1015	6.5%	0.98 [0.76 , 1.25	5]	+
Ley 2004	2	88	6	88	0.3%	0.33 [0.07 , 1.61]	<u> </u>
MRC 1968	28	199	31	194	1.8%	0.88 [0.55 , 1.41	.] .	-
Oslo Diet-Heart 1966	48	206	65	206	3.8%	0.74 [0.54 , 1.02	.]	•
Rose corn oil 1965	5	28	0	13	0.0%	5.31 [0.32, 89.44	-)	
Rose olive 1965	3	26	1	13	0.1%	1.50 [0.17 , 13.05	j <u> </u>	
STARS 1992	1	27	3	28	0.2%	0.35 [0.04 , 3.12	2]	<u> </u>
Sydney Diet-Heart 1978	39	221	28	237	1.6%	1.49 [0.95 , 2.34	-]	
Veterans Admin 1969	174	424	177	422	10.3%	0.98 [0.83 , 1.15	5]	•
WHI 2006 (1)	989	19541	1520	29294	70.4%	0.98 [0.90 , 1.05	5]	
WINS 2006	64	975	107	1462	5.0%	0.90 [0.67 , 1.21	.]	Ŧ
Total (95% CI)		22819		33039	100.0%	0.97 [0.91 , 1.03	3]	
Total events:	1465		2053					
Heterogeneity: Chi ² = 11.25	5, df = 11 (P =	= 0.42); I ²	= 2%				0.01 0.1	1 10 100
Test for overall effect: Z =	1.01 (P = 0.31)	l)					Favours lower SFA	Favours higher SFA

Test for subgroup differences: Not applicable

Footnotes

(1) All-cause death during study, Prentice 2017

Analysis 1.8. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 8: All-cause mortality, SA Peto fixed-effect

	lower	lower SFA		higher SFA		Peto Odds Ratio	Peto	Odds Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	Peto, Fixed, 95% CI	Peto, F	ixed, 95% CI
Black 1994	1	66	2	67	0.1%	0.52 [0.05 , 5.05]]	•
DART 1989	111	1018	113	1015	6.5%	0.98 [0.74 , 1.29]]	+
Ley 2004	2	88	6	88	0.3%	0.35 [0.09 , 1.45]	I	
MRC 1968	28	199	31	194	1.6%	0.86 [0.50, 1.50]]	
Oslo Diet-Heart 1966	48	206	65	206	2.7%	0.66 [0.43 , 1.02]] .	•
Rose corn oil 1965	5	28	0	13	0.1%	5.09 [0.70, 37.06]]	
Rose olive 1965	3	26	1	13	0.1%	1.50 [0.17 , 13.11]]	
STARS 1992	1	27	3	28	0.1%	0.36 [0.05 , 2.71]	I	
Sydney Diet-Heart 1978	39	221	28	237	1.9%	1.59 [0.95 , 2.68]]	
Veterans Admin 1969	174	424	177	422	6.7%	0.96 [0.73 , 1.27]]	+
WHI 2006 (1)	989	19541	1520	29294	74.8%	0.97 [0.90 , 1.06]]	
WINS 2006	64	975	107	1462	5.0%	0.89 [0.65 , 1.22]]	Ŧ
Total (95% CI)		22819		33039	100.0%	0.96 [0.90 , 1.04]]	
Total events:	1465		2053					
Heterogeneity: $Chi^2 = 13.01$	1, df = 11 (P =	= 0.29); I ² =	= 15%				0.01 0.1	1 10 100
Test for overall effect: Z =	1.00 (P = 0.32)	2)]	Favours lower SFA	Favours higher SFA

Footnotes

(1) All-cause death during study, Prentice 2017

Test for subgroup differences: Not applicable



Analysis 1.9. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 9: All-cause mortality, subgroup by any substitution

Study or Subgroup Events Total Weight M-H, Random, 95% C1 M-H, Random, 95% C1 19.1 replaced by PUEA DART 1989 111 1018 113 1015 24.4% 0.98 [0.76, 1.25] DART 1989 111 1018 113 1015 24.4% 0.088 [0.35, 1.41] Odo Dice-Hean 1966 48 206 65 206 17.5% 0.074 [0.54, 1.02] Study Dice-Hean 1966 1 227 3 228 0.06% 2.22 (0.0.1, 7.2] Study Dice-Hean 1960 174 424 177 422 37.2% 0.098 [0.83, 1.15] Subtool (95% C1) 201 206 418 Hetrogenetity: Trai=0.01; Chi ²⁺ 8.406, df = 61 = 0.23; F ² = 20% Tasta Tasta Tasta 0.06 [0.82, 1.13] Subtool (95% C1) 2 2 2 1000.9% 3.00 [0.33, 26.99] Subtool (95% C1) 2 88 6 88 0.51 [0.05, 5.46] DART 1989 11 1018 113 1015 8.5% 0.09 [0.67, 1.25] 0.09 [0.67, 1.25] 0.09		lower SFA		higher SFA			Risk Ratio	Risk Ratio
J. J. replaced by PUFA DART 1989 111 1018 113 1015 24.4% 0.98 [0.76, 1.25] MRC 1968 28 199 114 9.5% 0.88 [0.55, 1.41] Odo Die-Heart 1966 48 206 65 206 17.5% 0.74 [0.54, 1.02] Rose cano all 1965 5 28 1 13 0.66% 2.32 [0.30, 17.92] Sydney Die-Heart 1978 39 2.21 28 0.5% 0.35 [0.04, 3.12] Sydney Die-Heart 1976 39 2.21 21 50 (0.06% 0.36 [0.82, 1.13] Subtotal (95% C1) 2.123 2.115 100.0% 0.96 [0.82, 1.13] Noae olive 1965 3 2.6 100.0% 3.00 [0.33, 2.6.99] Subtotal (95% C1) 2.6 2.6 100.0% 3.00 [0.33, 2.6.99] Subtotal (95% C1) 2.6 2.6 100.0% 3.00 [0.33, 2.6.99] Subtotal (95% C1) 2.6 2.6 100.0% 3.00 [0.33, 2.6.99] Subtotal (95% C1) 2.6 2.6 100.0% 3.00 [0.33, 2.6.99] Subtotal (95% C1) 2.6 2	Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
DART 1989 111 1018 113 1015 244% 0.98 [0.76, 1.25] MRC 1968 28 199 31 194 9.5% 0.88 [0.55, 1.41] Oko Duct-Heart 1976 48 206 65 206 17.5% 0.74 [0.54, 1.02] Row corn oil 1965 5 28 1 13 0.6% 2.32 [0.30, 17.92] STARS 1992 1 27 3 28 0.5% 0.35 [0.04, 3.12] Sydney Duct-Heart 1978 39 221 28 237 10.3% 1.49 [0.95, 2.34] Veteraus Admin 1969 174 424 177 422 37.2% 0.98 [0.83, 1.15] Subtal (95% CI) 2123 2115 100.0% 0.96 [0.82, 1.13] Total events: 406 418 Heterogeneity: Tau ² = 0.01; $h2 = 8.06$, $d2 = 6 (P = 0.23)$; $P = 26\%$ Test for overall effect: Z = 0.46 (P = 0.64) 19.2 replaced by MUFA Rose oilve 1965 3 26 1 26 100.0% 3.00 [0.33, 26.99] Subtal (95% CI) 26 26 100.0% 3.00 [0.33, 26.99] Subtal (95% CI) 26 26 100.0% 3.00 [0.33, 26.99] Subtal (95% CI) 27 3 28 0.1% 0.51 [0.05, 5.46] DART 1989 111 1018 113 1015 8.5% 0.08 [0.76, 1.25] Ley 2004 2 88 6 88 0.2% 0.33 [0.07, 1.61] STARS 1992 1 27 3 28 0.1% 0.35 [0.05, 5.46] DART 1989 111 1018 113 1015 8.5% 0.98 [0.76, 1.25] WIN 2006 (1) 989 19541 1520 29294 85.3% 0.98 [0.06, 1.25] WIN 2006 (1) 989 19541 1520 29294 85.3% 0.98 [0.06, 1.21] WH 2006 (1) 989 19541 1520 29294 85.3% 0.98 [0.06, 1.21] WH 2006 (1) 989 19541 1520 29294 85.3% 0.98 [0.05, 5.46] DART 1989 111 1018 113 1015 8.5% 0.98 [0.06, 1.25] How results: 1168 1751 Heterogeneity: Tau ² = 0.03 (P = 0.36) 19.4 replaced by protein Biack 1994 1 66 2 67 0.1% 0.51 [0.05, 5.46] DART 1999 111 1018 113 1015 8.5% 0.98 [0.06, 1.25] H1 2016 (1) 989 19541 1520 29294 85.4% 0.98 [0.06, 1.25] H2 replaced by protein Biack 1994 1 66 2 67 0.1% 0.51 [0.05, 5.46] DART 1999 111 1018 113 1015 8.5% 0.98 [0.76, 1.21] Subtal (95% CI) 2 107 1462 5.8% 0.99 [0.67, 1.21] Subtal (95% CI) 2 108 31926 100.0% 0.97 [0.90, 1.04] H2 205 (1) 94 11 66 2 67 0.1% 0.51 [0.05, 5.46] DART 1999 111 1018 113 1015 8.5% 0.98 [0.76, 1.25] Ley 2004 2 88 6 88 0.2% 0.93 [0.76, 1.25] H1 206 (1) 99 99 19541 1520 29294 85.4% 0.99 [0.67, 1.21] Subtal (95% CI) 2 0 0 0 H1 206 (1) 99 99 194 11 520 29294 85.4% 0.99 [0.67, 1.21] Subtal	1.9.1 replaced by PUFA							
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	DART 1989	111	1018	113	1015	24.4%	0.98 [0.76 , 1.25]	1
Oslo Diet-Heart 1966 48 206 65 206 17.5% 0.74 [0.54, 1.02] Rase corn oil 1965 5 28 1 13 0.6% 2.32 [0.30, 17.92] Syluey Diet-Heart 1978 39 221 28 237 10.3% 1.49 [0.95, 2.34] Vectrama Admin 1969 174 424 177 422 37.2% 0.98 [0.83, 1.15] Subtoal (95% CI) 2123 2115 100.0% 0.98 [0.83, 1.15] Total events: 406 418 Heterogeneity: Tau ² = 0.01 Chi ² = 8.06, df = 6 (P = 0.23); P = 26% Test for overall effect: Z = 0.46 (P = 0.23); P = 26% Test for overall effect: Z = 0.46 (P = 0.23); P = 26% Test for overall effect: Z = 0.98 (P = 0.3) 19.3 replaced by MUFA Rose olive 1965 3 226 1 26 100.0% 3.00 [0.33, 26.99] Subtotal (95% CI) 26 26 100.0% 3.00 [0.33, 26.99] Subtotal (95% CI) 26 26 100.0% 3.00 [0.33, 26.99] Subtotal (95% CI) 27 3 28 0.1% 0.51 [0.05, 5.46] DART 1989 111 1018 113 1015 8.5% 0.98 [0.76, 1.25] Ley 2004 2 88 6 88 0.2% 0.33 [0.07, 1.61] Syltasy 100.05(1) 989 19541 1520 29294 85.3% 0.98 [0.06, 1.25] Ley 2004 2 88 6 88 0.2% 0.33 [0.07, 1.61] Subtotal (95% CI) 21713 31954 100.0% 0.97 [0.90, 1.04] Heterogeneity: Tau ³ = 0.00; Chi ³ = 2.31, 0, df = 5 (P = 0.67); P = 0% Test for overall effect: Z = 0.91 (P = 0.36) 19.4 replaced by protein Black 1994 1 66 2 67 0.1% 0.51 [0.05, 5.46] DART 1989 111 1018 113 1015 8.5% 0.98 [0.07, 1.21] Subtotal (95% CI) 21713 31954 100.0% 0.97 [0.90, 1.04] Total events: 1168 1751 Heterogeneity: Tau ³ = 0.00; Chi ³ = 2.34, df = 4 (P = 0.67); P = 0% Test for overall effect: Z = 0.91 (P = 0.36) 19.4 replaced by protein Black 1994 1 66 2 87 0.1% 0.51 [0.05, 5.46] DART 1989 111 1018 113 1015 8.5% 0.98 [0.07, 1.25] Ley 2004 2 88 6 88 0.2% 0.33 [0.07, 1.61] WH 2006 (1) 989 19541 1520 29294 85.4% 0.90 [0.67, 1.25] Ley 2004 2 168 31926 100.0% 0.97 [0.90, 1.04] Total events: 1167 T48 Heterogeneity: Tau ³ = 0.00; Chi ³ = 2.34, df = 4 (P = 0.67); P = 0% Test for overall effect: X = 0.88 (P = 0.38) 19.5 replacement unclear Subtotal (95% CI) 0 0 0 Heterogeneity: Not applicable Test for overall effect: X = 0.88 (P = 0.38) 19.5 replacement unclear Subtotal	MRC 1968	28	199	31	194	9.5%	0.88 [0.55 , 1.41]	
Rose corn oil 1965 5 28 1 13 0.6% 2.32 [0.30, 17.92] STAKS 1992 1 27 3 28 0.5% 0.55 [0.04, 3.12] Sydney Die-Heart 1978 39 21 28 237 [0.3%, 1.49] [0.05, 2.34] Veteras Admin 1969 174 424 177 422 37.2% 0.98 [0.83, 1.15] Subtoal 695% CD 2123 2115 100.0% 0.96 [0.82, 1.13] 1 Total events: 406 418 0.96 [0.82, 1.13] 1 Total events: 406 126 100.0% 3.00 [0.33, 26.99] Subtoal 695% CD 26 26 100.0% 3.00 [0.33, 26.99] Subtoal (95% CD 26 26 100.0% 3.00 [0.33, 26.99] Subtoal (95% CD 2 8 6 8 0.2% 0.33 [0.07, 1.61] Dack 1994 1 66 2 67 0.1% 0.51 [0.05, 5.46] Dack 1994 1 66 2 67 0.1% 0.53 [0.04, 3.12] WH 2006 (1) 989 19541 120 292.94	Oslo Diet-Heart 1966	48	206	65	206	17.5%	0.74 [0.54 , 1.02]	
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	Rose corn oil 1965	5	28	1	13	0.6%	2 32 [0 30 17 92]	
Sydney Die-Heart 1978 39 221 28 237 10.3% 1.49 [0.05, 2.34] Veternas Admin 1960 174 424 177 422 37.2% 0.08 [0.33, 1.15] Subtoal (95% C) 2123 2115 100.0% 0.98 [0.33, 1.15] Total events: 406 418 Heterogeneity: Tai ² = 0.0; Ch ² = 8.06, df = 6 (P = 0.23); P = 26% Test for overall effect: Z = 0.46 (P = 0.64) 1.9.2 replaced by MUFA Rose dive 1965 3 26 1 26 100.0% 3.00 [0.33, 26.99] Subtoal (95% CI) 26 26 26 100.0% 3.00 [0.33, 26.99] Subtoal (95% CI) 26 26 26 100.0% 3.00 [0.33, 26.99] Subtoal (95% CI) 26 26 26 00.0% 3.00 [0.33, 26.99] Subtoal (95% CI) 26 26 00.0% 3.00 [0.33, 26.99] 1.9.3 replaced by MUFA Test for overall effect: Z = 0.98 (P = 0.33) 1.9.3 replaced by CHO Black 1994 1 66 2 67 0.1% 0.51 [0.05, 5.46] DART 1989 111 1018 113 1015 8.5% 0.98 [0.76, 1.25] Ly 2004 2 88 6 88 0.2% 0.33 [0.07, 1.61] STARS 1902 1 277 3 28 0.1% 0.35 [0.04, 3.12] WH 2006 (1) 989 19541 1520 29294 85.3% 0.098 [0.76, 1.25] WH 2006 (2) 64 975 107 1462 5.8% 0.098 [0.76, 1.21] Subtoal (95% CI) 21715 31954 100.0% 0.97 [0.90, 1.04] Heterogeneity: Tai ² = 0.00; Chi ² = 3.19, df = 5 (P = 0.67); P = 0% Test for overall effect: Z = 0.91 (P = 0.36) 1.9.4 replaced by protein Black 1994 1 66 2 67 0.1% 0.51 [0.05, 5.46] DART 1989 111 1018 113 1015 8.5% 0.98 [0.76, 1.22] 4.9 2004 2 88 6 88 0.2% 0.03 [0.07, 1.61] WH 2006 (1) 989 19541 1520 29294 85.4% 0.98 [0.09, 1.05] WH 2006 (1) 989 19541 1520 29294 85.4% 0.98 [0.09, 1.05] WH 2006 (1) 989 19541 1520 29294 85.4% 0.98 [0.09, 1.05] WH 2006 (1) 989 19541 1520 29294 85.4% 0.98 [0.09, 1.05] WH 2006 (1) 989 19541 1520 29294 85.4% 0.98 [0.09, 1.05] WH 2006 (1) 989 19541 1520 29294 85.4% 0.98 [0.09, 1.05] WH 2006 (1) 989 19541 1520 29294 85.4% 0.98 [0.09, 1.05] WH 2006 (1) 989 19541 4 (P = 0.67); P = 0% Test for overall effect: Z = 0.88 (P = 0.38) 1.95 replacement unclear Subtoal (95% CI) 21688 31922 100.0% 0.97 [0.90, 1.04] Test for overall effect: Z = 0.88 (P = 0.38) 1.95 replacement unclear Subtoal (95% CI) 0 0 0 Net estimable Test for overall effect: Not a	STARS 1992	1	20 27	3	28	0.5%	0.35 [0.04 . 3.12]	
Veterans Admin 1969 174 424 177 422 37.2% $0.98 [0.83, 1.15]$ Subtoal 05% C1) 2123 2115 100.0% $0.96 [0.82, 1.13]$ Total events: 406 418 Heterogeneity: Tat" = 0.01: Chi" = 8.06, df = 6 (P = 0.23); P = 26% Test for overall effect: Z = 0.46 (P = 0.64) 1.9.2 replaced by MUFA Rose olive 1095 3 26 1 26 100.0% 3.00 [0.33, 26.99] Subtoal (95% C1) 26 26 100.0% 3.00 [0.33, 26.99] Subtoal (95% C1) 26 26 100.0% 3.00 [0.33, 26.99] Test for overall effect: Z = 0.98 (P = 0.33) 1.11 1018 113 1015 8.5% 0.98 [0.76, 1.25] Ley 2004 2 88 6 88 0.2% 0.33 [0.01, 3.51 [0.05, 5.46] Wh 2006 (1) 98 19541 1520 292.94 8.5.3% 0.98 [0.07, 1.21] Wh 2006 (1) 21715 31954 100.0% 0.97 [0.90, 1.04] 0.97 [0.90, 1.04] 0.97 [0.90, 1.04] Total events: 1168 1751 0.98 [0.76, 1.25] 0.98 [0.76,	Sydney Diet-Heart 1978	39	221	28	237	10.3%	1.49 [0.95 , 2.34]	
Subtoal (95% CI) 2123 2115 100.0% 0.96 [0.82, 1.13] Total events: 406 418 Heterogeneity: Tar# - 0.01; CP = 806, df = 6 (P = 0.23); P = 26% Test for overall effect: Z = 0.46 (P = 0.64) 1.9.2 replaced by MUFA Rose olive 1965 3 26 100.0% 3.00 [0.33, 26.99] Subtoal (95% CI) 26 26 100.0% 3.00 [0.33, 26.99] Subtoal (95% CI) 26 26 0.00.0% 3.00 [0.33, 26.99] Jobal vents: 3 1 1 Hetrogenetiy: SNot applicable Test for overall effect: Z = 0.98 (P = 0.33) 1 1018 113 1015 8.5% 0.98 [0.76, 1.25] DART 1989 111 1018 113 1015 8.5% 0.98 [0.04, 3.12] 1 VINS 2006 (2) 64 975 107 1462 5.8% 0.98 [0.90, 1.05] 1 Total events: 1168 1751 Heterogeneity: Tar# - 0.00; CP = 0.47); P = 0% 1 66 2 67 0.1% 0.97 [0.90, 1.04] 1 Total events: 1168 1751 Heterogeneity: Tar# - 0.00; CP = 0.67); P = 0.67; P = 0.67; P	Veterans Admin 1969	174	424	177	422	37.2%	0.98 [0.83 , 1.15]	
Total events: $406 418 100 10$	Subtotal (95% CI)		2123		2115	100.0%	0.96 [0.82 , 1.13]	I
The terms of the second secon	Total events:	406		418		10000 /0	000 [0002 ; 1110]	Ţ
The sequency in the intervent of the only is the only is the intervent of the only is there only is the only is there only is the only is the only is th	Heterogeneity: $Tau^2 = 0.01$.	$Chi^2 = 8.06$	df = 6 (P =	= 0 23)· I ² =	= 26%			
1.9.2 replaced by MUFA Rose olive 1965 3 26 1 26 100.0% 3.00 [0.33, 26.99] Subtoal (95% C1) 26 26 100.0% 3.00 [0.33, 26.99] Total events: 3 1 Heterogeneity: Not applicable Test for overall effect: Z = 0.98 (P = 0.33) 1.9.3 replaced by CHO Black 1994 1 66 2 67 0.1% 0.51 [0.05, 5.46] DART 1989 111 1018 113 1015 8.5% 0.98 [0.07, 1.21] STARS 1992 1 27 3 28 0.1% 0.33 [0.07, 1.61] WH1 2006 (1) 989 19541 1520 2924 8.3% 0.98 [0.90, 1.05] WH2 2006 (2) 64 975 107 1462 5.8% 0.99 [0.67, 1.21] Subtoal (95% C1) 21715 31954 100.0% 0.97 [0.90, 1.04] 101 Total events: 1168 1751 107 1462 5.8% 0.98 [0.67, 1.25] Ley 2004 2 88 6 88 0.2% 0.33 [007, 1.61]	Test for overall effect: $Z = 0$	0.46 (P = 0.64)	4)	- 0.25), 1 -	2070			
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $								
Rose of ve 1965 3 26 1 26 100.0% 3.00 [0.33, 26.99] Subtotal (95% CI) 26 26 26 100.0% 3.00 [0.33, 26.99] Total events: 3 1 Heterogeneity: Not applicable Test for overall effect: Z = 0.98 (P = 0.33) 1.3. replaced by CHO Black 1994 1 66 2 67 0.1% 0.51 [0.05, 5.46] DART 1989 111 1018 113 1015 8.5% 0.98 [0.76, 1.25] Ley 2004 2 88 6 88 0.2% 0.33 [0.07, 1.61] STARS 1992 1 27 3 282 0.1% 0.35 [0.04, 3.12] WH1 2006 (1) 989 19541 1520 29294 85.3% 0.98 [0.90, 1.05] WINS 2006 (2) 64 975 107 1462 5.8% 0.90 [0.67, 1.21] Subtotal (95% CI) 21715 31954 100.0% 0.97 [0.90, 1.04] Total events: 1168 1751 Heterogeneity: Tau ² = 0.00; Ch ² = 3.19, df = 5 (P = 0.75); P = 0% Test for overall effect: Z = 0.91 (P = 0.36) 1.9.4 replaced by protein Black 1994 1 66 2 67 0.1% 0.51 [0.05, 5.46] UALL 1018 113 1015 8.5% 0.98 [0.76, 1.25] Ley 2004 2 88 6 88 0.2% 0.33 [0.07, 1.61] WH 2006 (1) 989 19541 1520 29294 85.4% 0.98 [0.76, 1.25] Ley 2004 2 88 6 6 88 0.2% 0.33 [0.07, 1.61] WH 2006 (1) 989 19541 1520 29294 85.4% 0.98 [0.90, 1.05] WINS 2006 (2) 64 975 107 1462 5.8% 0.90 [0.67, 1.21] WH 2006 (1) 989 19541 1520 29294 85.4% 0.98 [0.90, 1.05] WINS 2006 (2) 64 975 107 1462 5.8% 0.90 [0.67, 1.21] Subtotal (95% CI) 21688 31926 100.0% 0.97 [0.90, 1.04] Total events: 1167 1748 Heterogeneity: Tau ² = 0.00; Ch ² = 2.34, df = 4 (P = 0.67); P = 0% Test for overall effect: Z = 0.88 (P = 0.38) 1.9.5 replacement unclear Subtotal (95% CI) 0 0 0 Not estimable Total events: 0 0 0 Heterogeneity: Not applicable Test for overall effect: Not applicable Test for overall effect: Not applicable Test for overall effect: Not applicable	1.9.2 replaced by MUFA		-		-			
Subtotal (95% CI) 26 26 100.0% $3.00 [0.33, 26.99]$ Total events: 3 1 Heterogeneity: Not applicable Test for overall effect: Z = 0.98 (P = 0.33) 1.9.3 replaced by CHO Black 1994 1 66 2 67 0.1% 0.51 [0.05, 5.46] DART 1989 111 1018 113 1015 8.5% 0.98 [0.76, 1.25] Ley 2004 2 88 6 88 0.2% 0.33 [0.04, 3.12] WHI 2006 (1) 989 19541 1520 29294 85.3% 0.98 [0.90, 1.05] WHNS 2006 (2) 64 975 107 1462 5.8% 0.90 [0.67, 1.21] Subtotal (95% CI) 21715 31954 100.0% 0.97 [0.90, 1.04] Total events: 1168 1751 Heterogeneity: Tau ² = 0.0; Ch ² = 3.19, dif = 5 (P = 0.67); P = 0% Test for overall effect: Z = 0.91 (P = 0.36) H1 1018 113 1015 8.5% 0.98 [0.76, 1.25] Ley 2004 2 88 6 88 0.2% 0.33 [0.07, 1.61] John Ch ² = 3.19, dif = 5 (P = 0.67); P = 0% Test for overall effect: Z = 0.91 (P = 0.36) H1 2006 (1) 989 19541 1520 29294 85.4% 0.98 [0.90, 1.05] WH 2006 (2) 64 975 107 1462 5.8% 0.98 [0.76, 1.25] Ley 2004 2 88 6 88 0.2% 0.33 [0.07, 1.61] Subtotal (95% CI) 21688 31926 100.0% 0.97 [0.90, 1.04] Total events: 1167 1748 Heterogeneity: Tau ² = 0.00; Ch ² = 2.34, df = 4 (P = 0.67); P = 0% Test for overall effect: Z = 0.38 (P = 0.38) H.5 replacement unclear Subtotal (95% CI) 0 0 0 Not estimable Total events: 0 0 0 Heterogeneity: Tau ² = 0.00; Ch ² = 2.34, df = 4 (P = 0.67); P = 0% Test for overall effect: Z = 0.88 (P = 0.38) H.5 replacement unclear Subtotal (95% CI) 0 0 0 Heterogeneity: Tau ² = 0.00; Ch ² = 2.34, df = 4 (P = 0.67); P = 0% Test for overall effect: X = 0.38 (P = 0.38) H.5 replacement unclear Subtotal (95% CI) 0 0 Heterogeneity: Not applicable Test for overall effect: Not applicable Test for overall effect: Not applicable Test for overall effect: Not applicable	Rose olive 1965	3	26	1	26	100.0%	3.00 [0.33 , 26.99]	
Total events: 3 1 Heterogeneity: Not applicable Test for overall effect: Z = 0.98 (P = 0.33) 1.9.3 replaced by CHO Black 1994 1 66 2 67 0.1% 0.51 [0.05, 5.46] DART 1989 111 1018 113 1015 8.5% 0.98 [0.76, 1.25] Ley 2004 2 88 6 88 0.2% 0.33 [0.07, 1.61] STARS 1992 1 27 3 28 0.1% 0.35 [0.04, 3.12] WH1 2006 (1) 989 19541 1520 22924 85.3% 0.98 [0.90, 1.05] WINS 2006 (2) 64 975 107 1462 5.8% 0.90 [0.67, 1.21] Subtotal (95% CI) 21715 31954 100.0% 0.97 [0.90, 1.04] Total events: 1168 1751 Heterogeneity: Tau ² = 0.00; Chi ² = 3.19, df = 5 (P = 0.67); P = 0% Test for overall effect: Z = 0.91 (P = 0.36) 1.9.4 replaced by protein Black 1994 1 66 2 67 0.1% 0.51 [0.05, 5.46] DART 1989 111 1018 113 1015 8.5% 0.98 [0.76, 1.25] Ley 2004 2 88 6 88 0.2% 0.33 [0.07, 1.61] WH1 2006 (1) 989 19541 1520 29294 85.4% 0.98 [0.90, 1.05] WINS 2006 (2) 64 975 107 1462 5.8% 0.90 [0.67, 1.21] Subtotal 695% CI) 21688 31926 100.0% 0.97 [0.90, 1.04] Total events: 1167 1748 Heterogeneity: Tau ² = 0.00; Chi ² = 2.34, df = 4 (P = 0.67); P = 0% Test for overall effect: Z = 0.38 (P = 0.38) 1.9.5 replacement unclear Subtotal (95% CI) 0 0 0 Not estimable Total events: 0 0 0 Heterogeneity: Tau ² = 0.00; Chi ² = 2.34, df = 4 (P = 0.67); P = 0% Test for overall effect: Z = 0.38 (P = 0.38) 1.9.5 replacement unclear Subtotal (95% CI) 0 0 0 Not estimable Total events: 0 0 0 Heterogeneity: Not applicable Test for overall effect: Not applicable Test for overall effect: Not applicable	Subtotal (95% CI)		26		26	100.0%	3.00 [0.33 , 26.99]	
Heterogeneity: Not applicable Test for overall effect: $Z = 0.98 (P = 0.33)$ 19.3 replaced by CHO Black 1994 1 66 2 67 0.1% 0.51 [0.05, 5.46] DART 1989 111 1018 113 1015 8.5% 0.98 [0.76, 1.25] Ley 2004 2 88 6 88 0.2% 0.33 [0.07, 1.61] STARS 1992 1 27 3 28 0.1% 0.35 [0.04, 3.12] WHI 2006 (1) 989 19541 1520 29294 85.3% 0.98 [0.90, 1.05] WHIN 2006 (2) 64 975 107 1462 5.8% 0.90 [0.67, 1.21] Subtotal (95% CI) 21715 31954 100.0% 0.97 [0.90, 1.04] Total events: 1168 1751 Heterogeneity: Tau ² = 0.00; Chi ² = 3.19, df = 5 (P = 0.67); P = 0% Test for overall effect: $Z = 0.91 (P = 0.36)$ 1.94 replaced by protein Black 1994 1 66 2 67 0.1% 0.51 [0.05, 5.46] DART 1989 111 1018 113 1015 8.5% 0.98 [0.76, 1.25] Ley 2004 2 88 6 88 0.2% 0.33 [0.07, 1.61] WHI 2006 (1) 989 19541 1520 29294 83.4% 0.98 [0.90, 1.05] WINS 2006 (2) 64 975 107 1462 5.8% 0.90 [0.67, 1.21] Subtotal (95% CI) 21688 31926 100.0% 0.97 [0.90, 1.04] WHI 2006 (2) 64 975 107 1462 5.8% 0.90 [0.67, 1.21] Subtotal (95% CI) 21688 31926 100.0% 0.97 [0.90, 1.04] Total events: 1167 1748 Heterogeneity: Tau ² = 0.00; Chi ² = 2.34, df = 4 (P = 0.67); P = 0% Test for overall effect: $Z = 0.88 (P = 0.38)$ 1.95 replacement unclear Subtotal (95% CI) 0 0 0 Not estimable Total events: 0 0 0 Heterogeneity: Not applicable Test for overall effect: Not applicable Test for overall effect: Not applicable Test or overall effect: Not applicable	Total events:	3		1				
Test for overall effect: $Z = 0.98$ (P = 0.33) 1.9.3 replaced by CHO Black 1994 1 66 2 67 0.1% 0.51 [0.05, 5.46] DART 1989 111 1018 113 1015 8.5% 0.98 [0.76, 1.25] Ley 2004 2 88 6 88 0.2% 0.33 [0.07, 1.61] WH 2006 (1) 989 19541 1520 29294 85.3% 0.98 [0.90, 1.05] WINS 2006 (2) 64 975 107 1462 5.8% 0.90 [0.67, 1.21] Subtotal 05% CD 21715 31954 100.0% 0.97 [0.90, 1.04] Total events: 1168 1751 Heterogeneity: Tau ² = 0.00; Chi ² = 3.19, df = 5 (P = 0.67); P = 0% Test for overall effect: $Z = 0.91$ (P = 0.36) 1.9.4 replaced by protein Black 1994 1 66 2 67 0.1% 0.51 [0.05, 5.46] DART 1989 111 1018 113 1015 8.5% 0.98 [0.76, 1.25] Ley 2004 2 88 6 88 0.2% 0.33 [0.07, 1.61] WH 2006 (1) 989 19541 1520 29294 85.4% 0.98 [0.76, 1.25] Ley 2004 2 88 6 88 0.2% 0.33 [0.07, 1.61] WH 2006 (2) 64 975 107 1462 5.8% 0.90 [0.67, 1.21] Subtotal 05% CD 21688 31926 100.0% 0.97 [0.90, 1.04] Total events: 1167 1748 Heterogeneity: Tau ² = 0.00; Chi ² = 2.34, df = 4 (P = 0.67); P = 0% Test for overall effect: Z = 0.88 (P = 0.38) 1.9.5 replacement unclear Subtotal 05% CD 0 0 0 Heterogeneity: Not applicable Test for overall effect: Not applicable	Heterogeneity: Not applicab	ole						
1.9.3 replaced by CHO Black 1994 1 66 2 67 0.1% $0.51 [0.05, 5.46]$ DART 1989 111 1018 113 1015 8.5% $0.98 [0.76, 1.25]$ Ley 2004 2 88 6 88 0.2% $0.33 [0.07, 1.61]$ STARS 1992 1 27 3 28 0.1% $0.35 [0.04, 3.12]$ WH1 206 (1) 989 19541 1520 29294 85.3% $0.98 [0.90, 1.05]$ VINS 2006 (2) 64 975 107 1462 5.8% $0.90 [0.67, 1.21]$ Subtotal (95% CI) 21715 31954 100.0% $0.97 [0.90, 1.04]$ Total events: 1168 1751 Heterogeneity: Tau ² = 0.00; Chi ² = 3.19, df = 5 (P = 0.67); P = 0% Test for overall effect: Z = 0.91 (P = 0.36) Hotologe 10 UH1 2006 (1) 98 [9 99 19541 1520 29294 85.4% 0.98 [0.90, 1.05] 0 DART 1989 111 1018 113 1015 8.5% 0.99 [0.67, 1.21] 0 0	Test for overall effect: $Z = 0$	0.98 (P = 0.33)	3)					
Black 1994 1 66 2 67 0.1% 0.51 [0.05, 5.46] DART 1989 111 1018 113 1015 8.5% 0.98 [0.76, 1.25] Ley 2004 2 88 6 88 0.2% 0.33 [0.07, 1.61] WH 2006 (1) 989 19541 1520 29294 85.3% 0.98 [0.90, 1.05] WINS 2006 (2) 64 975 107 1462 5.8% 0.90 [0.67, 1.21] Subtotal (95% CI) 21715 31954 100.0% 0.97 [0.90, 1.04] Total events: 1168 1751 Heterogeneity: Tau ² = 0.00; Ch ² = 3.19, df = 5 (P = 0.67); P = 0% Test for overall effect: Z = 0.91 (P = 0.36) HN 2006 (2) 64 975 107 1462 5.8% 0.98 [0.76, 1.25] Ley 2004 2 88 6 88 0.2% 0.98 [0.76, 1.25] Ley 2004 2 88 6 88 0.2% 0.98 [0.76, 1.25] WH 2006 (1) 989 19541 1520 29294 85.4% 0.98 [0.76, 1.25] WH 2006 (1) 989 19541 1520 29294 85.4% 0.98 [0.07, 1.61] WH 2006 (1) 989 19541 1520 29294 85.4% 0.98 [0.07, 1.21] Subtotal (95% CI) 21688 31926 100.0% 0.97 [0.90, 1.04] Total events: 1167 1748 Heterogeneity: Tau ² = 0.00; Ch ² = 2.34, df = 4 (P = 0.67); P = 0% Test for overall effect: Z = 0.38 (P = 0.38) 1.9.5 replacement unclear Subtotal (95% CI) 0 0 0 Not estimable Total events: 0 0 0 Heterogeneity: Not applicable Total events: 0 0 0	1.9.3 replaced by CHO							
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	Black 1994	1	66	2	67	0.1%	0.51 [0.05, 5.46]	
Ley 2004 2 88 6 88 0.2% 0.33 [0.07, 1.61] STARS 1992 1 27 3 28 0.1% 0.35 [0.04, 3.12] WH 2006 (1) 989 19541 1520 29294 85.3% 0.98 [0.90, 1.05] WINS 2006 (2) 64 975 107 1462 5.8% 0.90 [0.67, 1.21] Subtotal (95% CI) 21715 31954 100.0% 0.97 [0.90, 1.04] Total events: 1168 1751 Heterogeneity: Tau ² = 0.00; Chi ² = 3.19, df = 5 (P = 0.67); P = 0% Test for overall effect: Z = 0.91 (P = 0.36) 1.9.4 replaced by protein Black 1994 1 66 2 67 0.1% 0.51 [0.05, 5.46] DART 1989 111 1018 113 1015 8.5% 0.98 [0.76, 1.25] Ley 2004 2 88 6 88 0.2% 0.33 [0.07, 1.61] WH 2006 (1) 989 19541 1520 29294 85.4% 0.98 [0.90, 1.05] WINS 2006 (2) 64 975 107 1462 5.8% 0.90 [0.67, 1.21] Subtotal (95% CI) 21688 31926 100.0% 0.97 [0.90, 1.04] Total events: 1167 1748 Heterogeneity: Tau ² = 0.00; Chi ² = 2.34, df = 4 (P = 0.67); P = 0% Test for overall effect: Z = 0.88 (P = 0.38) 1.9.5 replacement unclear Subtotal (95% CI) 0 0 0 Not estimable Total events: 0 0 0 Heterogeneity: Not applicable Test for overall effect: Not applicable Test for overall effect: Not applicable	DART 1989	111	1018	113	1015	8.5%	0.98 [0.76, 1.25]	
STARS 1992 1 27 3 28 0.1% $0.35[0.04, 3.12]$ WHI 2006 (1) 989 19541 1520 29294 85.3% $0.98[0.90, 1.05]$ WINS 2006 (2) 64 975 107 1462 5.8% $0.90[0.67, 1.21]$ Subtotal (95% CI) 21715 31954 100.0% $0.97[0.90, 1.04]$ Total events: 1168 1751 Hetrogeneity: Tau ² = 0.00; Chi ² = 3.19, df = 5 (P = 0.67); P = 0% Test for overall effect: Z = 0.91 (P = 0.36) L9.4 replaced by protein Black 1994 1 66 2 67 0.1% $0.51 [0.05, 5.46]$ DART 1989 111 1018 113 1015 8.5% $0.98 [0.76, 1.25]$ Ley 2004 2 88 6 88 0.2% $0.33 [0.07, 1.61]$ WH1 2006 (1) 989 19541 1520 29294 85.4% $0.98 [0.90, 1.05]$ WIN 2006 (2) 64 975 107 1462 5.8% $0.90 [0.67, 1.21]$ Subtotal (95% CI) 21688 31926 100.0% $0.97 [0.90, 1$	Ley 2004	2	88	6	88	0.2%	0.33 [0.07, 1.61]	
WHI 2006 (1) 989 19541 1520 29294 85.3% 0.98 [0.90, 1.05] WINS 2006 (2) 64 975 107 1462 5.8% 0.90 [0.67, 1.21] Subtotal (95% CI) 21715 31954 100.0% 0.97 [0.90, 1.04] Total events: 1168 1751 Heterogeneity: Tau ² = 0.00; Chi ² = 3.19, df = 5 (P = 0.67); P = 0% Test for overall effect: Z = 0.91 (P = 0.36) 19.4 replaced by protein Black 1994 1 66 2 67 0.1% 0.51 [0.05, 5.46] DART 1989 111 1018 113 1015 8.5% 0.98 [0.76, 1.25] Ley 2004 2 88 6 88 0.2% 0.33 [0.07, 1.61] WH1 2006 (1) 98 19541 1520 29294 85.4% 0.98 [0.90, 1.05] WH2 2006 (2) 64 975 107 1462 5.8% 0.90 [0.67, 1.21] Subtotal (95% CI) 21688 31926 100.0% 0.97 [0.90, 1.04] 101 Total events: 1167 1748 Heterogeneity: Tau ² = 0.00; Chi ² = 2.34, df = 4 (P = 0.67); P = 0% <td>STARS 1992</td> <td>1</td> <td>27</td> <td>3</td> <td>28</td> <td>0.1%</td> <td>0.35 [0.04, 3.12]</td> <td></td>	STARS 1992	1	27	3	28	0.1%	0.35 [0.04, 3.12]	
WINS 2006 (2) 64 975 107 1462 5.8% 0.90 [0.67, 1.21] Subtotal (95% CI) 21715 31954 100.0% 0.97 [0.90, 1.04] Total events: 1168 1751 Heterogeneity: Tau ² = 0.00; Chi ² = 3.19, df = 5 (P = 0.67); P = 0% Test for overall effect: Z = 0.91 (P = 0.36) 19.4 replaced by protein Black 1994 1 66 2 67 0.1% 0.51 [0.05, 5.46] DART 1989 111 1018 113 1015 8.5% 0.98 [0.76, 1.25] Ley 2004 2 88 6 88 0.2% 0.33 [0.07, 1.61] WH1 2006 (1) 989 19541 1520 29294 85.4% 0.98 [0.90, 1.05] WINS 2006 (2) 64 975 107 1462 5.8% 0.90 [0.67, 1.21] Subtotal (95% CI) 21688 31926 100.0% 0.97 [0.90, 1.04] 101 Total events: 1167 1748 Heterogeneity: Tau ² = 0.00; Chi ² = 2.34, df = 4 (P = 0.67); I ² = 0% Test for overall effect: Z = 0.88 (P = 0.38) Not estimable Total events: 0 0	WHI 2006 (1)	989	19541	1520	29294	85.3%	0.98 [0.90, 1.05]	-
Subtotal (95% CI) 21715 31954 100.0% 0.97 [0.90, 1.04] Total events: 1168 1751 Heterogeneity: Tau ² = 0.00; Chi ² = 3.19, df = 5 (P = 0.67); I ² = 0% Test for overall effect: $Z = 0.91$ (P = 0.36) 19.4 replaced by protein Black 1994 1 66 2 67 0.1% 0.51 [0.05, 5.46] DART 1989 111 1018 113 1015 8.5% 0.98 [0.76, 1.25] Ley 2004 2 88 6 88 0.2% 0.33 [0.07, 1.61] WHI 2006 (1) 989 19541 1520 29294 85.4% 0.98 [0.90, 1.05] WINS 2006 (2) 64 975 107 1462 5.8% 0.90 [0.67, 1.21] Subtotal (95% CI) 21688 31926 100.0% 0.97 [0.90, 1.04] 1 Total events: 1167 1748 1748 1748 1 Heterogeneity: Tau ² = 0.00; Chi ² = 2.34, df = 4 (P = 0.67); I ² = 0% 7 10 0 0 19.5 replacement unclear 10 0 0 Not estimable 1 Total events:<	WINS 2006 (2)	64	975	107	1462	5.8%	0.90 [0.67, 1.21]	
Total events: 1168 1751 Heterogeneity: Tau ² = 0.00; Chi ² = 3.19, df = 5 (P = 0.67); l ² = 0% Test for overall effect: $Z = 0.91$ (P = 0.36) 1.9.4 replaced by protein Black 1994 1 66 2 67 0.1% 0.51 [0.05, 5.46] DART 1989 111 1018 113 1015 8.5% 0.98 [0.76, 1.25] Ley 2004 2 88 6 88 0.2% 0.33 [0.07, 1.61] WH 2006 (1) 989 19541 1520 29294 85.4% 0.98 [0.90, 1.05] WINS 2006 (2) 64 975 107 1462 5.8% 0.90 [0.67, 1.21] Subtotal (95% CI) 21688 31926 100.0% 0.97 [0.90, 1.04] 100.0% Total events: 1167 1748 1748 1748 1742 0.00; Chi ² = 2.34, df = 4 (P = 0.67); l ² = 0% 1.9.5 1.9.5 1.9.5 1.9.5 1.9.6 1.9.6 1.9.6 1.9.6 1.9.6 1.9.6 1.0.0% 1.9.97 [0.90; 1.04] 1.0.1 1.9.5 1.9.6 1.9.6 1.9.6 1.9.6 1.9.6 <td>Subtotal (95% CI)</td> <td></td> <td>21715</td> <td></td> <td>31954</td> <td>100.0%</td> <td>0.97 [0.90 , 1.04]</td> <td></td>	Subtotal (95% CI)		21715		31954	100.0%	0.97 [0.90 , 1.04]	
Heterogeneity: Tau ² = 0.00; Chi ² = 3.19, df = 5 (P = 0.67); I ² = 0% Test for overall effect: Z = 0.91 (P = 0.36) 1.9.4 replaced by protein Black 1994 1 66 2 67 0.1% 0.51 [0.05, 5.46] DART 1989 111 1018 113 1015 8.5% 0.98 [0.76, 1.25] Ley 2004 2 88 6 88 0.2% 0.33 [0.07, 1.61] WHI 2006 (1) 989 19541 1520 29294 85.4% 0.98 [0.90, 1.05] WINS 2006 (2) 64 975 107 1462 5.8% 0.90 [0.67, 1.21] Subtotal (95% CI) 21688 31926 100.0% 0.97 [0.90, 1.04] Total events: 1167 1748 Heterogeneity: Tau ² = 0.00; Chi ² = 2.34, df = 4 (P = 0.67); I ² = 0% Test for overall effect: Z = 0.88 (P = 0.38) 1.9.5 replacement unclear Subtotal (95% CI) 0 0 Not estimable Total events: 0 0 0 Heterogeneity: Not applicable Total events: Not applicable	Total events:	1168		1751			- , -	
Test for overall effect: $Z = 0.91$ (P = 0.36) 1.9.4 replaced by protein Black 1994 1 66 2 67 0.1% 0.51 [0.05, 5.46] DART 1989 111 1018 113 1015 8.5% 0.98 [0.76, 1.25] Ley 2004 2 88 6 88 0.2% 0.33 [0.07, 1.61] WH 2006 (1) 989 19541 1520 29294 85.4% 0.98 [0.90, 1.05] WINS 2006 (2) 64 975 107 1462 5.8% 0.90 [0.67, 1.21] Subtotal (95% CI) 21688 31926 100.0% 0.97 [0.90, 1.04] Total events: 1167 1748 Heterogeneity: Tau ² = 0.00; Chi ² = 2.34, df = 4 (P = 0.67); P = 0% Test for overall effect: Z = 0.88 (P = 0.38) 1.9.5 replacement unclear Subtotal (95% CI) 0 0 Not estimable Total events: 0 0 0 Heterogeneity: Not applicable Test for overall effect: Not applicable	Heterogeneity: $Tau^2 = 0.00$;	$Chi^2 = 3.19,$	df = 5 (P =	= 0.67); I ² =	= 0%			
1.9.4 replaced by protein Black 1994 1 66 2 67 0.1% $0.51 [0.05, 5.46]$ DART 1989 111 1018 113 1015 8.5% $0.98 [0.76, 1.25]$ Ley 2004 2 88 6 88 0.2% $0.33 [0.07, 1.61]$ WH 2006 (1) 989 19541 1520 29294 85.4% $0.98 [0.90, 1.05]$ WINS 2006 (2) 64 975 107 1462 5.8% $0.90 [0.67, 1.21]$ Subtotal (95% CI) 21688 31926 100.0% $0.97 [0.90, 1.04]$ Total events: 1167 1748 Heterogeneity: Tau ² = 0.00; Chi ² = 2.34, df = 4 (P = 0.67); l ² = 0% Test for overall effect: Z = 0.88 (P = 0.38) Not estimable Total events: 0 0 Total events: 0 0 Not estimable Total events: 0 0 Heterogeneity: Not applicable Test for overall effect: Not applicable Test for overall effect: Not applicable Test for overall effect: Not applicable Unit applicable Unit appli	Test for overall effect: $Z = 0$	0.91 (P = 0.36)	5)	,.				
Black 1994 1 66 2 67 0.1% 0.51 [0.05, 5.46] DART 1989 111 1018 113 1015 8.5% 0.98 [0.76, 1.25] Ley 2004 2 88 6 88 0.2% 0.33 [0.07, 1.61] WHI 2006 (1) 989 19541 1520 29294 85.4% 0.98 [0.90, 1.05] WINS 2006 (2) 64 975 107 1462 5.8% 0.90 [0.67, 1.21] Subtoal (95% CI) 21688 31926 100.0% 0.97 [0.90, 1.04] Total events: 1167 1748 Heterogeneity: Tau ² = 0.00; Chi ² = 2.34, df = 4 (P = 0.67); I ² = 0% Test for overall effect: $Z = 0.88$ (P = 0.38) 1.9.5 replacement unclear Subtoal (95% CI) 0 0 0 Not estimable Total events: 0 0 0 Heterogeneity: Not applicable Test for overall effect: Not applicable	1.9.4 replaced by protein							
DART 1989 111 1018 113 1015 8.5% 0.98 [0.76, 1.25] Ley 2004 2 88 6 88 0.2% 0.33 [0.07, 1.61] WHI 2006 (1) 989 19541 1520 29294 85.4% 0.98 [0.90, 1.05] WINS 2006 (2) 64 975 107 1462 5.8% 0.90 [0.67, 1.21] Subtotal (95% CI) 21688 31926 100.0% 0.97 [0.90, 1.04] Total events: 1167 1748 Heterogeneity: Tau ² = 0.00; Chi ² = 2.34, df = 4 (P = 0.67); I ² = 0% Test for overall effect: Z = 0.88 (P = 0.38) 1.9.5 replacement unclear Subtotal (95% CI) 0 0 Total events: 0 0 Heterogeneity: Not applicable 0 0 Test for overall effect: Not applicable Use the second secon	Black 1994	1	66	2	67	0.1%	0.51 [0.05 , 5.46]	
Ley 2004 2 88 6 88 0.2% 0.33 [0.07, 1.61] WHI 2006 (1) 989 19541 1520 29294 85.4% 0.98 [0.90, 1.05] WINS 2006 (2) 64 975 107 1462 5.8% 0.90 [0.67, 1.21] Subtotal (95% CI) 21688 31926 100.0% 0.97 [0.90, 1.04] Total events: 1167 1748 Heterogeneity: Tau ² = 0.00; Chi ² = 2.34, df = 4 (P = 0.67); I ² = 0% Test for overall effect: Z = 0.88 (P = 0.38) 1.9.5 replacement unclear Subtotal (95% CI) 0 0 Total events: 0 0 Heterogeneity: Not applicable 0 0 Test for overall effect: Not applicable 10 0 Test for overall effect: Not applicable 5 0 Test for overall effect: Not applicable 5 5 Test for overall effect: Not applicable 5 6	DART 1989	111	1018	113	1015	8.5%	0.98 [0.76 , 1.25]	
WHI 2006 (1) 989 19541 1520 29294 85.4% 0.98 [0.90, 1.05] WINS 2006 (2) 64 975 107 1462 5.8% 0.90 [0.67, 1.21] Subtotal (95% CI) 21688 31926 100.0% 0.97 [0.90, 1.04] Total events: 1167 1748 Heterogeneity: Tau ² = 0.00; Chi ² = 2.34, df = 4 (P = 0.67); I ² = 0% Test for overall effect: Z = 0.88 (P = 0.38) 1.9.5 replacement unclear Subtotal (95% CI) 0 0 Not estimable Total events: 0 0 Heterogeneity: Not applicable 0 0 Test for overall effect: Not applicable 10 0	Lev 2004	2	88	6	88	0.2%	0.33 [0.07 , 1.61]	<u> </u>
WINS 2006 (2) 64 975 107 1462 5.8% 0.90 [0.67, 1.21] Subtotal (95% CI) 21688 31926 100.0% 0.97 [0.90, 1.04] Total events: 1167 1748 Heterogeneity: Tau ² = 0.00; Chi ² = 2.34, df = 4 (P = 0.67); l ² = 0% Test for overall effect: Z = 0.88 (P = 0.38) 1.9.5 replacement unclear Subtotal (95% CI) 0 0 Not estimable Total events: 0 0 Not estimable Total events: 0 0 Not estimable Total events: 0 0 Not estimable	WHI 2006 (1)	989	19541	1520	29294	85.4%	0.98 [0.90 , 1.05]	
Subtotal (95% CI) 21688 31926 100.0% 0.97 [0.90, 1.04] Total events: 1167 1748 Heterogeneity: Tau ² = 0.00; Chi ² = 2.34, df = 4 (P = 0.67); I ² = 0% Test for overall effect: Z = 0.88 (P = 0.38) 1.9.5 replacement unclear Subtotal (95% CI) 0 0 Not estimable Total events: 0 0 Not estimable Total events: 0 0 Heterogeneity: Not applicable Test for overall effect: Not applicable Test for overall effect: Not applicable Test for overall effect: Not applicable	WINS 2006 (2)	64	975	107	1462	5.8%	0.90 [0.67 1.21]	_
Total events:11671748Heterogeneity: Tau ² = 0.00; Chi ² = 2.34, df = 4 (P = 0.67); I ² = 0%Test for overall effect: Z = 0.88 (P = 0.38) 1.9.5 replacement unclear Subtotal (95% CI)00Total events:000Heterogeneity: Not applicableTest for overall effect: Not applicable	Subtotal (95% CI)		21688		31926	100.0%	0.97 [0.90 , 1.04]	
How we have How we have Heterogeneity: Tau ² = 0.00; Chi ² = 2.34, df = 4 (P = 0.67); I ² = 0% Test for overall effect: Z = 0.88 (P = 0.38) 1.9.5 replacement unclear Subtotal (95% CI)0Not estimableTotal events:00Total events:00Heterogeneity: Not applicableTest for overall effect: Not applicable	Total events:	1167		1748				
Test for overall effect: $Z = 0.88$ (P = 0.38) 1.9.5 replacement unclear Subtotal (95% CI) 0 0 0 Total events: 0 1.9.5 for overall effect: Not applicable	Heterogeneity: $Tau^2 = 0.00$.	$Chi^2 = 2.34$	df = 4 (P -	= 0.67)· I ² -	= 0%			
1.9.5 replacement unclearSubtotal (95% CI)00Not estimableTotal events:00Heterogeneity: Not applicable0Test for overall effect: Not applicable	Test for overall effect: $Z = 0$	0.88 (P = 0.38)	3)	<i>,</i> , -	570			
I.9.5 replacement unclear 0 0 Not estimable Subtotal (95% CI) 0 0 0 Total events: 0 0 0 Heterogeneity: Not applicable - - - Test for overall effect: Not applicable - - -	105							
Subtotal (95% Cl) 0 0 Not estimable Total events: 0 0 Heterogeneity: Not applicable	1.9.5 replacement unclear						xy , , y x x	
I otal events: 0 0 Heterogeneity: Not applicable Test for overall effect: Not applicable	Subtotal (95% CI)	~	0	~	0		Not estimable	
Heterogeneity: Not applicable Test for overall effect: Not applicable	Total events:	0		0				
Test for overall effect: Not applicable	Heterogeneity: Not applicab	ole						
	lest for overall effect: Not a	applicable						
Test for subgroup differences: Chi ² = 1.02, df = 3 (P = 0.80), l ² = 0% $0.05 - 0.2 + 1 + 5 + 20$	Test for subgroup difference	es: Chi ² = 1.0	2, df = 3 (P = 0.80), I	$^{2} = 0\%$		_	
Favours lower SFA Favours higher	Footnotes						Fa	vours lower SFA Favours higher SF

(1) All-cause death during study, Prentice 2017

(2) All-cause mortality during trial, Chlebowski 2015

Analysis 1.10. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 10: All-cause mortality, subgroup by main substitution

	lower SFA		higher SFA		Risk Ratio		Risk Ratio	
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI	
1.10.1 replaced by PUFA								
DART 1989	111	1018	113	1015	24.5%	0.98 [0.76, 1.25]		
MRC 1968	28	199	31	194	9.8%	0.88 [0.55, 1.41]		
Oslo Diet-Heart 1966	48	206	65	206	17.7%	0.74 [0.54, 1.02]		
Rose corn oil 1965	5	28	1	13	0.6%	2.32 [0.30, 17.92]		
Sydney Diet-Heart 1978	39	221	28	237	10.6%	1.49 [0.95, 2.34]		
Veterans Admin 1969	174	424	177	422	36.8%	0.98 [0.83, 1.15]		
Subtotal (95% CI)		2096		2087	100.0%	0.97 [0.82, 1.14]	⊥	
Total events:	405		415			- / -	Ţ	
Heterogeneity: $Tau^2 = 0.01$; ($Chi^2 = 7.23,$	df = 5 (P =	= 0.20); I ² =	31%				
Test for overall effect: $Z = 0$.	.39 ($P = 0.70$))						
1.10.2 replaced by MUFA								
Rose olive 1965	3	26	1	26	100.0%	3.00 [0.33 , 26.99]		
Subtotal (95% CI)		26		26	100.0%	3.00 [0.33 , 26.99]		
Total events:	3		1					
Heterogeneity: Not applicabl	le							
Test for overall effect: $Z = 0$.	.98 ($P = 0.33$)						
1.10.3 replaced by CHO								
Black 1994	1	66	2	67	0.1%	0.51 [0.05 , 5.46]		
Ley 2004	2	88	6	88	0.2%	0.33 [0.07, 1.61]		
STARS 1992	1	27	3	28	0.1%	0.35 [0.04, 3.12]		
WHI 2006 (1)	989	19541	1520	29294	93.2%	0.98 [0.90 , 1.05]	-	
WINS 2006 (2)	64	975	107	1462	6.3%	0.90 [0.67 , 1.21]		
Subtotal (95% CI)		20697		30939	100.0%	0.97 [0.90 , 1.04]		
Total events:	1057		1638					
Heterogeneity: $Tau^2 = 0.00$; ($Chi^2 = 3.17,$	df = 4 (P =	= 0.53); I ² =	:0%				
Test for overall effect: $Z = 0$.	.90 (P = 0.37)						
1.10.4 replaced by protein		-		_				
Subtotal (95% CI)	-	0	~	0		Not estimable		
1 otal events:	0		0					
Heterogeneity: Not applicabl Test for overall effect: Not ap	e pplicable							
1.10.5 replacement unclear								
Subtotal (95% CI)		0		0		Not estimable		
Total events:	0		0					
Heterogeneity: Not applicabl	le							
Test for overall effect: Not a	pplicable							
Test for subgroup differences	s: $Chi^2 = 1.0$	2, df = 2 (P = 0.60), I	$^{2} = 0\%$		E	0.05 0.2 1 5 20	
						Га	rouis lower STA Favours night	

Footnotes

(1) All-cause death during study, Prentice 2017

(2) All-cause mortality during trial, Chlebowski 2015



Analysis 1.11. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 11: All-cause mortality, subgroup by duration

	lower SFA		higher SFA			Risk Ratio	Risk Rati	0
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random,	95% CI
1.11.1 up to 24mo								
Black 1994	1	66	2	67	0.1%	0.51 [0.05 , 5.46]		
DART 1989	111	1018	113	1015	7.6%	0.98 [0.76, 1.25]		
Rose corn oil 1965	5	28	0	13	0.1%	5.31 [0.32, 89.44]		
Rose olive 1965	3	26	1	13	0.1%	1.50 [0.17, 13.05]		
Subtotal (95% CI)		1138		1108	7.8%	0.99 [0.78 , 1.26]	•	
Total events:	120		116				Ť	
Heterogeneity: $Tau^2 = 0.003$; Chi ² = 1.82,	df = 3 (P =	= 0.61); I ² =	:0%				
Test for overall effect: $Z = 0$	0.08 (P = 0.94)	4)						
1.11.2 >24 to 48mo								
MRC 1968	28	199	31	194	2.1%	0.88 [0.55, 1.41]	_	
STARS 1992	1	27	3	28	0.1%	0.35 [0.04 , 3.12]		_
Veterans Admin 1969	174	424	177	422	17.4%	0.98 [0.83 , 1.15]		
Subtotal (95% CI)		650		644	19.6%	0.96 [0.83, 1.12]	4	
Total events:	203		211				The second se	
Heterogeneity: $Tau^2 = 0.003$	$Chi^2 = 1.02,$	df = 2 (P = 2)	= 0.60); I ² =	:0%				
Test for overall effect: $Z = 0$	0.49 (P = 0.63)	3)						
1.11.3 >48mo								
Oslo Diet-Heart 1966	48	206	65	206	4.6%	0.74 [0.54 , 1.02]	-	
Sydney Diet-Heart 1978	39	221	28	237	2.3%	1.49 [0.95 , 2.34]		
WHI 2006 (1)	989	19541	1520	29294	60.2%	0.98 [0.90 , 1.05]		
WINS 2006 (2)	64	975	107	1462	5.2%	0.90 [0.67 , 1.21]	-	
Subtotal (95% CI)		20943		31199	72.4%	0.96 [0.79 , 1.16]	•	
Total events:	1140		1720				1	
Heterogeneity: $Tau^2 = 0.02$	$Chi^2 = 6.63,$	df = 3 (P = 3)	= 0.08); I ² =	: 55%				
Test for overall effect: $Z = 0$	0.45 (P = 0.65)	5)						
1.11.4 unclear duration								
Ley 2004	2	88	6	88	0.2%	0.33 [0.07 , 1.61]		
Subtotal (95% CI)		88		88	0.2%	0.33 [0.07 , 1.61]		
Total events:	2		6					
Heterogeneity: Not applical	ole							
Test for overall effect: Z =	1.37 (P = 0.17)	7)						
Total (95% CI)		22819		33039	100.0%	0.96 [0.90 , 1.03]		
Total events:	1465		2053				1	
Heterogeneity: $Tau^2 = 0.003$; Chi ² = 11.25	, df = 11 (P = 0.42; I	$^{2} = 2\%$			0.05 0.2 1	5 20
Test for overall effect: $Z =$	1.01 (P = 0.3)	l)				Fa	vours lower SFA I	Favours higher SFA
Test for subgroup difference	es: $Chi^2 = 1.8$	0, df = 3 (P = 0.61), I	$^{2} = 0\%$				

Footnotes

(1) All-cause death during study, Prentice 2017

(2) All-cause mortality during trial, Chlebowski 2015

Analysis 1.12. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 12: All-cause mortality, subgroup by baseline SFA

			myner	SFA		KISK KAIIO	KISK KADO
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
1.12.1 up to 12%E SFA base	line						
WINS 2006 (1)	64	975	107	1462	5.2%	0.90 [0.67 , 1.21]	-
Subtotal (95% CI)		975		1462	5.2%	0.90 [0.67 , 1.21]	•
Total events:	64		107				T
Heterogeneity: Not applicable							
Test for overall effect: $Z = 0.7$	1 (P = 0.48)	5)					
1.12.2 >12 to 15% E SFA bas	eline						
Black 1994	1	66	2	67	0.1%	0.51 [0.05 , 5.46]	
DART 1989	111	1018	113	1015	7.6%	0.98 [0.76 , 1.25]	
Lev 2004	2	88	6	88	0.2%	0.33 [0.07 , 1.61]	<u> </u>
Sydney Diet-Heart 1978	39	221	28	237	2.3%	1.49 [0.95 , 2.34]	
WHI 2006 (2)	989	19541	1520	29294	60.2%	0.98 [0.90 , 1.05]	
Subtotal (95% CI)	,0,	20934	1020	30701	70.4%	1.01 [0.86 . 1.19]	T
Total events:	1142	-0701	1669	00701		101 [0000 ; 1125]	▼
Heterogeneity: $Tau^2 = 0.01$; C	$hi^2 = 5.48$	df = 4 (P =	$= 0.24$): $I^2 =$	27%			
Test for overall effect: $Z = 0.1$	1 (P = 0.92)	2)	0.2.), 1	2770			
1.12.3 >15 to 18%E SFA bas	eline						
STARS 1992	1	27	3	28	0.1%	0 35 [0 04 3 12]	
Subtotal (95% CI)	1	27	5	28	0.1%	0.35 [0.04, 3.12]	
Total events:	1		3	20	012 /0	0.00 [0.00] (0.12]	
Heterogeneity: Not applicable	-		5				
Test for overall effect: $Z = 0.9$	5 (P = 0.34)					
1.12.4 >18%E SFA baseline							
Veterans Admin 1969	174	424	177	422	17.4%	0.98 [0.83, 1.15]	1
Subtotal (95% CI)		424		422	17.4%	0.98 [0.83, 1.15]	. In the second s
Total events:	174		177			- / -	Ť
Heterogeneity: Not applicable							
Test for overall effect: $Z = 0.2$	7 (P = 0.79)))					
1.12.5 unclear							
MRC 1968	28	199	31	194	2.1%	0.88 [0.55, 1.41]	
Oslo Diet-Heart 1966	48	206	65	206	4.6%	0.74 [0.54 , 1.02]	-
Rose corn oil 1965	5	28	0	13	0.1%	5.31 [0.32, 89.44]	
Rose olive 1965	3	26	1	13	0.1%	1.50 [0.17 , 13.05]	
Subtotal (95% CI)		459		426	6.9%	0.80 [0.62 , 1.04]	
Total events:	84		97			L ,	
Heterogeneity: $Tau^2 = 0.00$: Cl	$hi^2 = 2.50.$	df = 3 (P =	= 0.47); I ² =	:0%			
Test for overall effect: $Z = 1.6$	7 (P = 0.10)))					
Total (95% CI)		22819		33039	100.0%	0.96 [0.90 , 1.03]	
Total events:	1465		2053				Y
Heterogeneity: $Tau^2 = 0.00$: Cl	hi ² = 11.25	df = 11	P = 0.42): I	$^{2} = 2\%$			
Test for overall effect: $Z = 1.0$	1 (P = 0.31)	.)	,,			Fav	ours lower SFA Favours higher SF
Test for subgroup differences:	$Chi^2 = 3.2$	5, df = 4 (1)	P = 0.52), I	$^{2} = 0\%$			e a

Footnotes

(1) All-cause mortality during trial, Chlebowski 2015

(2) All-cause death during study, Prentice 2017


Analysis 1.13. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 13: All-cause mortality, subgroup by SFA change

	lower	SFA	higher	SFA		Risk Ratio	Risk R	atio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Randor	m, 95% CI
1.13.1 up to 4%E differen	ce							
DART 1989	111	1018	113	1015	7.6%	0.98 [0.76, 1.25]		
Ley 2004	2	88	6	88	0.2%	0.33 [0.07, 1.61]		_
Sydney Diet-Heart 1978	39	221	28	237	2.3%	1.49 [0.95 , 2.34]		-
WHI 2006 (1)	989	19541	1520	29294	60.2%	0.98 [0.90 , 1.05]		
WINS 2006 (2)	64	975	107	1462	5.2%	0.90 [0.67, 1.21]	_	
Subtotal (95% CI)		21843		32096	75.5%	0.99 [0.86 , 1.13]	•	
Total events:	1205		1774				Ť	
Heterogeneity: $Tau^2 = 0.01$; Chi ² = 5.53,	df = 4 (P =	= 0.24); I ² =	= 28%				
Test for overall effect: $Z = 0$	0.17 (P = 0.87)	7)						
1.13.2 >4 to 8%E differen	ce							
Black 1994	1	66	2	67	0.1%	0.51 [0.05 , 5.46]		
STARS 1992	1	27	3	28	0.1%	0.35 [0.04 , 3.12]		
Subtotal (95% CI)		93		95	0.2%	0.41 [0.08 , 2.07]		
Total events:	2		5					
Heterogeneity: $Tau^2 = 0.003$; $Chi^2 = 0.05$,	df = 1 (P = 1)	= 0.82); I ² =	= 0%				
Test for overall effect: Z =	1.07 (P = 0.28)	3)						
1.13.3 >8%E difference								
Veterans Admin 1969	174	424	177	422	17.4%	0.98 [0.83 , 1.15]	+	
Subtotal (95% CI)		424		422	17.4%	0.98 [0.83 , 1.15]	•	
Total events:	174		177				Ĩ	
Heterogeneity: Not applical	ble							
Test for overall effect: $Z = 0$	0.27 (P = 0.79)))						
1.13.4 unclear								
MRC 1968	28	199	31	194	2.1%	0.88 [0.55 , 1.41]		-
Oslo Diet-Heart 1966	48	206	65	206	4.6%	0.74 [0.54 , 1.02]		
Rose corn oil 1965	5	28	0	13	0.1%	5.31 [0.32, 89.44]		
Rose olive 1965	3	26	1	13	0.1%	1.50 [0.17 , 13.05]		•
Subtotal (95% CI)		459		426	6.9%	0.80 [0.62 , 1.04]		
Total events:	84		97				•	
Heterogeneity: $Tau^2 = 0.003$; Chi ² = 2.50,	df = 3 (P = 3)	= 0.47); I ² =	= 0%				
Test for overall effect: Z =	1.67 (P = 0.10)))						
Total (95% CI)		22819		33039	100.0%	0.96 [0.90 , 1.03]		
Total events:	1465		2053					
Heterogeneity: $Tau^2 = 0.003$; Chi ² = 11.25	i, df = 11 (P = 0.42; 1	$2^2 = 2\%$			0.05 0.2 1	5 20
Test for overall effect: $Z =$	1.01 (P = 0.3)	1)				F	avours lower SFA	Favours higher SFA
Test for subgroup difference	es: Chi ² = 3.1	3, $df = 3$ (P = 0.37), I	$^{2} = 4.0\%$				

Footnotes

(1) All-cause death during study, Prentice 2017

(2) All-cause mortality during trial, Chlebowski 2015



Analysis 1.14. Comparison 1: SFA reduction vs usual diet -	primary
outcomes, Outcome 14: All-cause mortality, subgroup by	y sex

	lower	lower SFA		higher SFA		Risk Ratio	Risk Ratio	
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI	
1.14.1 Men								
Black 1994	1	66	2	67	0.1%	0.51 [0.05 , 5.46]	•	
DART 1989	111	1018	113	1015	7.6%	0.98 [0.76, 1.25]	_	
MRC 1968	28	199	31	194	2.1%	0.88 [0.55 , 1.41]		
Oslo Diet-Heart 1966	48	206	65	206	4.6%	0.74 [0.54, 1.02]		
Rose corn oil 1965	5	28	0	13	0.1%	5.31 [0.32, 89.44]		→
Rose olive 1965	3	26	1	13	0.1%	1.50 [0.17, 13.05]		
STARS 1992	1	27	3	28	0.1%	0.35 [0.04, 3.12]		
Sydney Diet-Heart 1978	39	221	28	237	2.3%	1.49 [0.95 , 2.34]		
Veterans Admin 1969	174	424	177	422	17.4%	0.98 [0.83, 1.15]		
Subtotal (95% CI)		2215		2195	34.4%	0.96 [0.83, 1.11]	4	
Total events:	410		420				T	
Heterogeneity: $Tau^2 = 0.01$; Chi ² = 9.20,	df = 8 (P =	= 0.33); I ² =	= 13%				
Test for overall effect: $Z = 0$	0.53 (P = 0.60)))						
1.14.2 Women								
WHI 2006 (1)	989	19541	1520	29294	60.2%	0.98 [0.90, 1.05]	_	
WINS 2006 (2)	64	975	107	1462	5.2%	0.90 [0.67 , 1.21]	Ŧ	
Subtotal (95% CI)		20516		30756	65.4%	0.97 [0.90 , 1.05]		
Total events:	1053		1627					
Heterogeneity: $Tau^2 = 0.00$; Chi ² = 0.28,	df = 1 (P = 1)	= 0.59); I ² =	= 0%				
Test for overall effect: $Z = 0$	0.79 (P = 0.43)	3)						
1.14.3 Mixed, men and wo	omen							
Ley 2004	2	88	6	88	0.2%	0.33 [0.07 , 1.61]		
Subtotal (95% CI)		88		88	0.2%	0.33 [0.07 , 1.61]		
Total events:	2		6					
Heterogeneity: Not applical	ble							
Test for overall effect: Z =	1.37 (P = 0.17)	7)						
Total (95% CI)		22819		33039	100.0%	0.96 [0.90 , 1.03]		
Total events:	1465		2053				1	
Heterogeneity: $Tau^2 = 0.00$; Chi ² = 11.25	df = 11 (P = 0.42; I	$^{2} = 2\%$			0.05 0.2 1 5 20)
Test for overall effect: Z =	1.01 (P = 0.31)	1)				Fa	vours lower SFA Favours high	ner SFA
Test for subgroup difference	es: Chi ² = 1.7	7, $df = 2$ (P = 0.41), I	$^{2} = 0\%$				

Footnotes

(1) All-cause death during study, Prentice 2017

(2) All-cause mortality during trial, Chlebowski 2015

Analysis 1.15. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 15: All-cause mortality, subgroup by CVD risk

	lower	SFA	higher	SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
1.15.1 Low CVD risk							
Black 1994	1	66	2	67	0.1%	0.51 [0.05, 5.46]	_
Veterans Admin 1969	174	424	177	422	17.4%	0.98 [0.83 , 1.15]	.
WHI 2006 (1)	989	19541	1520	29294	60.2%	0.98 [0.90 , 1.05]	.
WINS 2006 (2)	64	975	107	1462	5.2%	0.90 [0.67 , 1.21]	_ T
Subtotal (95% CI)		21006		31245	82.9%	0.97 [0.91 , 1.04]	6
Total events:	1228		1806				
Heterogeneity: $Tau^2 = 0.00$	Chi ² = 0.58,	df = 3 (P =	= 0.90); I ² =	0%			
Test for overall effect: $Z = 0$	0.84 (P = 0.40)))					
1.15.2 Moderate CVD risk	Σ.						
Ley 2004	2	88	6	88	0.2%	0.33 [0.07 , 1.61]	
Subtotal (95% CI)		88		88	0.2%	0.33 [0.07 , 1.61]	
Total events:	2		6				
Heterogeneity: Not applical	ole						
Test for overall effect: Z =	1.37 (P = 0.17)	7)					
1.15.3 Existing CVD disea	se						
DART 1989	111	1018	113	1015	7.6%	0.98 [0.76 , 1.25]	
MRC 1968	28	199	31	194	2.1%	0.88 [0.55, 1.41]	
Oslo Diet-Heart 1966	48	206	65	206	4.6%	0.74 [0.54, 1.02]	
Rose corn oil 1965	5	28	0	13	0.1%	5.31 [0.32, 89.44]	_
Rose olive 1965	3	26	1	13	0.1%	1.50 [0.17, 13.05]	•
STARS 1992	1	27	3	28	0.1%	0.35 [0.04 , 3.12]	
Sydney Diet-Heart 1978	39	221	28	237	2.3%	1.49 [0.95 , 2.34]	
Subtotal (95% CI)		1725		1706	16.9%	0.97 [0.76 , 1.24]	•
Total events:	235		241				Ĭ
Heterogeneity: $Tau^2 = 0.03$	Chi ² = 8.89,	df = 6 (P =	= 0.18); I ² =	33%			
Test for overall effect: $Z = 0$	0.26 (P = 0.79)))					
Total (95% CI)		22819		33039	100.0%	0.96 [0.90 , 1.03]	
Total events:	1465		2053				
Heterogeneity: $Tau^2 = 0.00$	Chi ² = 11.25	, df = 11 (P = 0.42; I	² = 2%			0.05 0.2 1 5 20
Test for overall effect: Z =	1.01 (P = 0.31)	1)				Fa	avours lower SFA Favours higher SFA
Test for subgroup differenc	es: Chi ² = 1.7	7, $df = 2$ (P = 0.41), I	$^{2} = 0\%$			

Footnotes

(1) All-cause death during study, Prentice 2017

(2) All-cause mortality during trial, Chlebowski 2015



Analysis 1.16. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 16: All-cause mortality, subgroup by TC reduction

	lower	SFA	higher	SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
1.16.1 serum chol reduced	by at least 0	.2mmol/I	1				
DART 1989	111	1018	113	1015	24.5%	0.98 [0.76 , 1.25]	+
MRC 1968	28	199	31	194	10.3%	0.88 [0.55, 1.41]	
Oslo Diet-Heart 1966	48	206	65	206	18.1%	0.74 [0.54, 1.02]	
Rose corn oil 1965	5	28	0	13	0.4%	5.31 [0.32, 89.44]	
STARS 1992	1	27	3	28	0.6%	0.35 [0.04 , 3.12]	•
Sydney Diet-Heart 1978	39	221	28	237	11.1%	1.49 [0.95 , 2.34]	
Veterans Admin 1969	174	424	177	422	35.1%	0.98 [0.83 , 1.15]	_
Subtotal (95% CI)		2123		2115	100.0%	0.96 [0.81 , 1.14]	▲
Total events:	406		417				Ĭ
Heterogeneity: $Tau^2 = 0.01$;	, Chi ² = 8.76,	df = 6 (P =	= 0.19); I ² =	= 32%			
Test for overall effect: $Z = 0$	0.42 (P = 0.67)	7)					
1.16.2 serum chol reduced	l by <0.2mm	ol/L					
Ley 2004	2	88	6	88	0.2%	0.33 [0.07 , 1.61]	-
Rose olive 1965	3	26	1	13	0.1%	1.50 [0.17 , 13.05]	
WHI 2006 (1)	989	19541	1520	29294	93.3%	0.98 [0.90 , 1.05]	
WINS 2006 (2)	64	975	107	1462	6.4%	0.90 [0.67 , 1.21]	
Subtotal (95% CI)		20630		30857	100.0%	0.97 [0.90 , 1.04]	
Total events:	1058		1634				
Heterogeneity: $Tau^2 = 0.00$;	Chi ² = 2.21,	df = 3 (P =	= 0.53); I ² =	= 0%			
Test for overall effect: $Z = 0$	0.84 (P = 0.40)))					
1.16.3 serum chol reduction	on unclear						
Black 1994	1	66	2	67	100.0%	0.51 [0.05 , 5.46]	
Subtotal (95% CI)		66		67	100.0%	0.51 [0.05 , 5.46]	
Total events:	1		2				
Heterogeneity: Not applicat	ole						
Test for overall effect: $Z = 0$	0.56 (P = 0.58)	3)					
Test for subgroup difference	es: Chi ² = 0.2	9, df = 2 (P = 0.87), I	² = 0%		Fa	$\begin{array}{c c c c c c c c c c c c c c c c c c c $

Footnotes

(1) All-cause death during study, Prentice 2017

(2) All-cause mortality during trial, Chlebowski 2015



Analysis 1.17. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 17: All-cause mortality, subgroup decade of publication

	lower	SFA	higher	SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
1.17.1 1960s							
MRC 1968	28	199	31	194	2.1%	0.88 [0.55 , 1.41]	_
Oslo Diet-Heart 1966	48	206	65	206	4.6%	0.74 [0.54 , 1.02]	-
Rose corn oil 1965	5	28	0	13	0.1%	5.31 [0.32, 89.44]	
Rose olive 1965	3	26	1	13	0.1%	1.50 [0.17, 13.05]	,
Veterans Admin 1969	174	424	177	422	17.4%	0.98 [0.83 , 1.15]	
Subtotal (95% CI)		883		848	24.3%	0.92 [0.80 , 1.07]	
Total events:	258		274				
Heterogeneity: $Tau^2 = 0.00$; Chi ² = 4.09,	df = 4 (P =	= 0.39); I ² =	2%			
Test for overall effect: Z =	1.09 (P = 0.27)	')					
1.17.2 1970s							
Sydney Diet-Heart 1978	39	221	28	237	2.3%	1.49 [0.95 , 2.34]	↓
Subtotal (95% CI)		221		237	2.3%	1.49 [0.95 , 2.34]	
Total events:	39		28				•
Heterogeneity: Not applical	ble						
Test for overall effect: Z =	1.75 (P = 0.08)	3)					
1.17.3 1980s							
DART 1989	111	1018	113	1015	7.6%	0.98 [0.76 , 1.25]	+
Subtotal (95% CI)		1018		1015	7.6%	0.98 [0.76 , 1.25]	
Total events:	111		113				
Heterogeneity: Not applical	ble						
Test for overall effect: $Z = 0$	0.17 (P = 0.87)	')					
1.17.4 1990s							
Black 1994	1	66	2	67	0.1%	0.51 [0.05 , 5.46]	
STARS 1992	1	27	3	28	0.1%	0.35 [0.04 , 3.12]	
Subtotal (95% CI)		93		95	0.2%	0.41 [0.08 , 2.07]	
Total events:	2		5				-
Heterogeneity: $Tau^2 = 0.00$; $Chi^2 = 0.05$,	df = 1 (P =	= 0.82); I ² =	:0%			
Test for overall effect: Z =	1.07 (P = 0.28)	5)					
1.17.5 2000s							
Ley 2004	2	88	6	88	0.2%	0.33 [0.07 , 1.61]	
WHI 2006 (1)	989	19541	1520	29294	60.2%	0.98 [0.90 , 1.05]	•
WINS 2006 (2)	64	975	107	1462	5.2%	0.90 [0.67 , 1.21]	
Subtotal (95% CI)		20604		30844	65.6%	0.97 [0.88, 1.05]	•
Total events:	1055		1633				
Heterogeneity: $Tau^2 = 0.00$; $Chi^2 = 2.05$,	df = 2 (P =	= 0.36); I ² =	: 3%			
Test for overall effect: $Z = 0$	0.79 (P = 0.43	5)					
Total (95% CI)		22819		33039	100.0%	0.96 [0.90 , 1.03]	
Total events:	1465		2053				
Heterogeneity: $Tau^2 = 0.00$; $Chi^2 = 11.25$, df = 11 (P = 0.42; I	$^{2} = 2\%$			0.05 0.2 1 5 20
Test for overall effect: $Z =$	1.01 (P = 0.31))				Fav	Yours lower SFA Favours higher SFA
Test for subgroup difference	es: $Chi^2 = 5.0$	8, df = 4 (P = 0.28), I	$^{2} = 21.2\%$			

Footnotes

(1) All-cause death during study, Prentice 2017

(2) All-cause mortality during trial, Chlebowski 2015



lower SFA **Risk Ratio** higher SFA **Risk Ratio** Study or Subgroup Events Total Events Total Weight M-H, Random, 95% CI M-H, Random, 95% CI Black 1994 0 66 2 67 0.4% 0.20 [0.01, 4.15] DART 1989 101 19.9% 1.01 [0.77, 1.31] 1018 100 1015 Ley 2004 88 4 88 0.7% 0.25 [0.03, 2.19] 1 MRC 1968 27 199 25 194 1.05 [0.63 , 1.75] 9.5% 38 52 0.73 [0.50 , 1.06] Oslo Diet-Heart 1966 206 206 14.2% Rose corn oil 1965 5 28 0 0.4% 5.31 [0.32, 89.44] 13 Rose olive 1965 3 26 1 13 0.7% 1.50 [0.17 , 13.05] **STARS 1992** 1 27 3 28 0.7%0.35 [0.04 , 3.12] Sydney Diet-Heart 1978 37 221 237 10.5% 1.59 [0.99 , 2.55] 25 57 424 422 17.2% 0.70 [0.51, 0.96] Veterans Admin 1969 81 WHI 2006 (1) 213 19541 320 29294 25.7%1.00 [0.84 , 1.19] 0.94 [0.78 , 1.13] Total (95% CI) 21844 31577 100.0% Total events: 483 613 Heterogeneity: Tau² = 0.03; Chi² = 15.67, df = 10 (P = 0.11); I² = 36% 0.01 0.1 10 100 Test for overall effect: Z = 0.68 (P = 0.49)Favours lower SFA Favours higher SFA

Analysis 1.18. Comparison 1: SFA reduction vs usual diet primary outcomes, Outcome 18: CARDIOVASCULAR MORTALITY

Test for subgroup differences: Not applicable

Footnotes

(1) In participants with and without CVD at baseline

Analysis 1.19. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 19: CVD mortality, SA low summary risk of bias

	lower	lower SFA		higher SFA		Risk Ratio	Ris	Risk Ratio		
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Ran	dom, 95% (CI	
Ley 2004	1	88	4	88	2.6%	0.25 [0.03 , 2.19	·]			
Sydney Diet-Heart 1978	37	221	25	237	25.0%	1.59 [0.99 , 2.55	5]			
Veterans Admin 1969	57	424	81	422	32.9%	0.70 [0.51, 0.96	6] -	-		
WHI 2006 (1)	213	19541	320	29294	39.5%	1.00 [0.84 , 1.19	9]	•		
Total (95% CI)		20274		30041	100.0%	0.96 [0.67 , 1.38	3]			
Total events:	308		430					Ť		
Heterogeneity: Tau ² = 0.08;	Chi ² = 9.96,	df = 3 (P :	= 0.02); I ² =	= 70%			0.01 0.1	1 10	100	
Test for overall effect: $Z = 0$	0.21 (P = 0.83)	3)					Favours lower SFA	Favour	s higher SFA	
Test for subgroup difference	es: Not applic	able								

Footnotes

(1) In participants with and without CVD at baseline



Analysis 1.20. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 20: CVD mortality, SA aim to reduce SFA

	lower	SFA	higher SFA			Risk Ratio	Risk	. Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Rano	lom, 95% CI
DART 1989	101	1018	100	1015	20.2%	1.01 [0.77 , 1.31]]	•
MRC 1968	27	199	25	194	9.5%	1.05 [0.63 , 1.75]] .	-
Oslo Diet-Heart 1966	38	206	52	206	14.3%	0.73 [0.50 , 1.06]	4	•
Rose corn oil 1965	5	28	0	13	0.4%	5.31 [0.32, 89.44]		
Rose olive 1965	3	26	1	13	0.7%	1.50 [0.17, 13.05]		.
STARS 1992	1	27	3	28	0.7%	0.35 [0.04 , 3.12]		<u> </u>
Sydney Diet-Heart 1978	37	221	25	237	10.4%	1.59 [0.99 , 2.55]]	-
Veterans Admin 1969	57	424	81	422	17.3%	0.70 [0.51, 0.96]] -	•
WHI 2006 (1)	213	19541	320	29294	26.5%	1.00 [0.84 , 1.19]]	•
Total (95% CI)		21690		31422	100.0%	0.95 [0.79 , 1.14]]	
Total events:	482		607					
Heterogeneity: Tau ² = 0.03;	Chi ² = 13.22	, df = 8 (P)	= 0.10; I ²	= 39%			0.01 0.1	1 10 100
Test for overall effect: $Z = 0$	0.54 (P = 0.59))]	Favours lower SFA	Favours higher SFA
Test for subgroup difference	es: Not applic	able						

Footnotes

(1) In participants with and without CVD at baseline

Analysis 1.21. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 21: CVD mortality, SA statistically significant SFA reduction

	lower	lower SFA		higher SFA		Risk Ratio	Risk R	atio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Randon	n, 95% CI
Black 1994	0	66	2	67	0.6%	0.20 [0.01 , 4.15]	
DART 1989	101	1018	100	1015	26.3%	1.01 [0.77, 1.31]	1 🖕	
Ley 2004	1	88	4	88	1.2%	0.25 [0.03 , 2.19]]	_
STARS 1992	1	27	3	28	1.1%	0.35 [0.04 , 3.12]]	
Sydney Diet-Heart 1978	37	221	25	237	15.3%	1.59 [0.99 , 2.55]] 4	-
Veterans Admin 1969	57	424	81	422	23.3%	0.70 [0.51, 0.96]] 🗕	
WHI 2006 (1)	213	19541	320	29294	32.2%	1.00 [0.84 , 1.19]	J 🛉	
Total (95% CI)		21385		31151	100.0%	0.95 [0.75 , 1.21]]	
Total events:	410		535				Ţ	
Heterogeneity: $Tau^2 = 0.04$; Chi ² = 11.92	, df = 6 (P	P = 0.06; I ²	= 50%			0.01 0.1 1	10 100
Test for overall effect: Z =	0.40 (P = 0.69)))					Favours lower SFA	Favours higher SFA

Test for subgroup differences: Not applicable

Footnotes

(1) In participants with and without CVD at baseline



Analysis 1.22. Comparison 1: SFA reduction vs usual diet primary outcomes, Outcome 22: CVD mortality, SA TC reduction

	lower	SFA	higher SFA		Risk Ratio		Risk	Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Rand	om, 95% CI
DART 1989	101	1018	100	1015	20.2%	1.01 [0.77 , 1.31	.] .	•
MRC 1968	27	199	25	194	9.9%	1.05 [0.63 , 1.75	j] _	-
Oslo Diet-Heart 1966	38	206	52	206	14.6%	0.73 [0.50 , 1.06	6] 🗕 🗕	
Rose corn oil 1965	5	28	0	13	0.5%	5.31 [0.32, 89.44	·]	
STARS 1992	1	27	3	28	0.7%	0.35 [0.04 , 3.12	2]	
Sydney Diet-Heart 1978	37	221	25	237	10.9%	1.59 [0.99 , 2.55	5]	
Veterans Admin 1969	57	424	81	422	17.5%	0.70 [0.51 , 0.96	6] -	
WHI 2006 (1)	213	19541	320	29294	25.7%	1.00 [0.84 , 1.19)]	
Total (95% CI)		21664		31409	100.0%	0.95 [0.78 , 1.15	5]	
Total events:	479		606					
Heterogeneity: Tau ² = 0.03;	Chi ² = 13.05	, df = 7 (P)	= 0.07); I ²	= 46%			0.01 0.1	1 10 100
Test for overall effect: $Z = 0$	0.54 (P = 0.59))					Favours lower SFA	Favours higher SFA

Test for subgroup differences: Not applicable

Footnotes

(1) In participants with and without CVD at baseline

Analysis 1.23. Comparison 1: SFA reduction vs usual diet primary outcomes, Outcome 23: CVD mortality, SA excluding WHI

	lower	SFA	higher	SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
Black 1994	0	66	2	67	0.7%	0.20 [0.01 , 4.15]	I
DART 1989	101	1018	100	1015	24.3%	1.01 [0.77 , 1.31]	· •
Ley 2004	1	88	4	88	1.3%	0.25 [0.03 , 2.19]	I
MRC 1968	27	199	25	194	14.1%	1.05 [0.63 , 1.75]	I 📥
Oslo Diet-Heart 1966	38	206	52	206	19.2%	0.73 [0.50 , 1.06]	I -
Rose corn oil 1965	5	28	0	13	0.8%	5.31 [0.32, 89.44]	I
Rose olive 1965	3	26	1	13	1.3%	1.50 [0.17, 13.05]	I
STARS 1992	1	27	3	28	1.3%	0.35 [0.04 , 3.12]	I
Sydney Diet-Heart 1978	37	221	25	237	15.2%	1.59 [0.99 , 2.55]	I 4
Veterans Admin 1969	57	424	81	422	21.9%	0.70 [0.51 , 0.96]	•
Total (95% CI)		2303		2283	100.0%	0.92 [0.72 , 1.18]	
Total events:	270		293				Y
Heterogeneity: $Tau^2 = 0.05$; Chi ² = 15.03	, df = 9 (P	P = 0.09; I ²	= 40%			0.01 0.1 1 10 100
Test for overall effect: $Z = 0$	0.65 (P = 0.52)	2)]	Favours lower SFA Favours higher SFA
Test for subgroup differenc	es: Not applic	able					



Analysis 1.24. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 24: CVD mortality, SA Mantel-Haenszel fixed-effect

	lower	lower SFA		higher SFA		Risk Ratio	Ris	sk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Fixed, 95% CI	M-H, Fi	xed, 95% CI
Black 1994	0	66	2	67	0.5%	0.20 [0.01 , 4.15	j]	
DART 1989	101	1018	100	1015	18.2%	1.01 [0.77 , 1.31]	+
Ley 2004	1	88	4	88	0.7%	0.25 [0.03 , 2.19) <u> </u>	
MRC 1968	27	199	25	194	4.6%	1.05 [0.63 , 1.75	5]	—
Oslo Diet-Heart 1966	38	206	52	206	9.4%	0.73 [0.50 , 1.06	5]	-
Rose corn oil 1965	5	28	0	13	0.1%	5.31 [0.32, 89.44	-) –	
Rose olive 1965	3	26	1	13	0.2%	1.50 [0.17 , 13.05	j <u> </u>	
STARS 1992	1	27	3	28	0.5%	0.35 [0.04 , 3.12	2]	
Sydney Diet-Heart 1978	37	221	25	237	4.4%	1.59 [0.99 , 2.55	5]	
Veterans Admin 1969	57	424	81	422	14.8%	0.70 [0.51 , 0.96	5]	
WHI 2006 (1)	213	19541	320	29294	46.5%	1.00 [0.84 , 1.19)]	•
Total (95% CI)		21844		31577	100.0%	0.95 [0.85 , 1.07	']	
Total events:	483		613					
Heterogeneity: $Chi^2 = 15.6^{\circ}$	7, df = 10 (P =	= 0.11); I ²	= 36%				0.01 0.1	1 10 100
Test for overall effect: Z =	0.83 (P = 0.41)	l)					Favours lower SFA	Favours higher SFA
T	Nat1:							-

Test for subgroup differences: Not applicable

Footnotes

(1) In participants with and without CVD at baseline

Analysis 1.25. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 25: CVD mortality, SA Peto fixed-effect

	lower	SFA	higher	SFA		Peto Odds Ratio	Peto O	dds Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	Peto, Fixed, 95% CI	Peto, Fix	ed, 95% CI
Black 1994	0	66	2	67	0.2%	0.14 [0.01 , 2.19)]	
DART 1989	101	1018	100	1015	18.3%	1.01 [0.75 , 1.35	5]	+
Ley 2004	1	88	4	88	0.5%	0.29 [0.05 , 1.73	j	
MRC 1968	27	199	25	194	4.6%	1.06 [0.59 , 1.90)] .	-
Oslo Diet-Heart 1966	38	206	52	206	7.1%	0.67 [0.42 , 1.07]	_ _
Rose corn oil 1965	5	28	0	13	0.4%	5.09 [0.70 , 37.06	j]	
Rose olive 1965	3	26	1	13	0.3%	1.50 [0.17 , 13.11]	<u>-</u>
STARS 1992	1	27	3	28	0.4%	0.36 [0.05 , 2.71]	
Sydney Diet-Heart 1978	37	221	25	237	5.4%	1.70 [0.99 , 2.90)]	
Veterans Admin 1969	57	424	81	422	11.7%	0.66 [0.46 , 0.95	j] _	F
WHI 2006 (1)	213	19541	320	29294	51.1%	1.00 [0.84 , 1.19]	•
Total (95% CI)		21844		31577	100.0%	0.95 [0.84 , 1.08	3]	
Total events:	483		613					
Heterogeneity: Chi ² = 18.54	df = 10 (P =	= 0.05); I ² =	= 46%				0.01 0.1	1 10 100
Test for overall effect: $Z = 0$	0.82 (P = 0.41)	.)					Favours lower SFA	Favours higher SFA

Test for subgroup differences: Not applicable

Footnotes

(1) In participants with and without CVD at baseline



Analysis 1.26. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 26: CVD mortality, subgroup by any substitution

indy or Subgroup Events Total Weight M-H, Random, 95% C1 M-H, Random, 95% C1 2.6.1 replaced by PUF. - <th></th> <th>lower</th> <th>SFA</th> <th>higher</th> <th>SFA</th> <th></th> <th>Risk Ratio</th> <th>Risk Ratio</th>		lower	SFA	higher	SFA		Risk Ratio	Risk Ratio
Cols replaced by PUEA ART 1989 101 1018 100 1015 24.1% 1.01 1.07 1.31 ART 1989 101 1018 100 1015 24.1% 1.01 10.77 1.31 Nob Dist-Heart 1966 38 206 52 206 19.7% 0.73 (150 1.05 System 2019 1 27 3 28 1.5% 0.35 (10.41, 1.5) (1.75) System 2019 1 27 3 28 1.5% 0.35 (10.41, 1.5) (10.77, 1.31) System 2019 1 27 3 28 1.5% 0.35 (10.73, 1.25) System 2016 2123 2123 2128 100.0% 0.96 (0.73, 1.25) System 2016 3 2.6 100.07% 3.00 (0.33, 26.99) (0.100.0% 3.00 (0.33, 26.99) (0.100.0% 3.00 (0.33, 26.99) (0.100.0% (0.99 (0.85, 1.14) (0.99 (0.85, 1.14) (0.99 (0.85, 1.14) (0.99 (0.85, 1.14) (0.99 (0.85, 1.14) <th< th=""><th>Study or Subgroup</th><th>Events</th><th>Total</th><th>Events</th><th>Total</th><th>Weight</th><th>M-H, Random, 95% CI</th><th>M-H, Random, 95% CI</th></th<>	Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	1.26.1 replaced by PUFA							
ARC 1968 27 199 25 194 15.0% 1.05 [0.63, 1.75] blo Dici-Heart 1966 38 206 52 206 19.7% 0.73 [0.50, 1.06] Sca corn oil 1965 5 28 1 26 1.6% 4.64 [0.58, 1.75] Max 1992 1 27 3 28 1.5% 0.55 [0.04, 3.12] Valuey Dici-Heart 1978 37 221 25 237 1.59 [0.99, 2.55] Value Vents: 265 287 Ieterogeneity: Tat# = 0.06; ChP = 1.3.37, df = 6 (P = 0.04); P = 55% sis for ovent effect Z = 0.34 (P = 0.73) Z62 26 100.0% 3.00 [0.33, 26.99] uhotal 05% CD 2 2 26 100.0% 3.00 [0.33, 26.99] uhotal 05% CD 2 26 7 0.2% 0.20 [0.01, 4.15] Staf royenel filect: Z = 0.38 (P = 0.3) 2 2.88 0.4% 0.55 [0.01, 3.12] Z63 replace by CHO 108 k 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] 1.01 [0.77, 1.31] y2 204h 1 84 4.88 0.4%	DART 1989	101	1018	100	1015	24.1%	1.01 [0.77, 1.31]	+
blo Die Heart 1966 38 206 52 206 19.7% 0.73 [0.50, 1.06] tose corn oil 1965 5 28 1 26 1.6% 4.64 [0.58, 37.15] tose corn oil 1965 5 287 1.29 10.99, 2.55] everans Admin 1969 377 424 81 422 22.1% 0.70 [0.51, 0.96] whole all 65% C1) 2123 2128 100.9% 0.95 [0.73, 1.25] otal events: 266 287 test for overall effect: Z = 0.34 (P = 0.04); P = 55% test for overall effect: Z = 0.34 (P = 0.04); P = 55% test for overall effect: Z = 0.98 (P = 0.04); P = 55% test for overall effect: Z = 0.98 (P = 0.33) 2.63 replaced by MUFA tesrogeneity: Teat = 0.06; Chi ² = 1.03.7 2.64 replaced by MUFA tesrogeneity: Teat = 0.98 (P = 0.33) 2.65 replaced by MUFA tesrogeneity: Teat = 0.98 (P = 0.33) 2.64 replaced by MUFA tesrogeneity: Teat = 0.98 (P = 0.48); P = 0.48); P = 0.48 tesrogeneity: Net applicable test for overall effect: Z = 0.98 (P = 0.48); P = 0.48; P = 0.48; P = 0.48 2.64 replaced by protein takk 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] tesrogeneity: Taa' = 0.00; Chi ² = 3.51, df = 4 (P = 0.48); P = 0.48 2.64 replaced by protein takk 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] test for overall effect: Z = 0.19 (P = 0.48); P = 0.48 2.64 replaced by protein takk 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] 2.64 replaced by protein takk 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] 2.64 replaced by protein takk 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] 2.64 replaced by protein takk 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] 2.64 replaced by protein takk 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] 2.64 replaced by protein takk 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] 2.65 replaced by protein takk 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] 2.65 replaced by protein takk 1994 0 60 0 0 test for overall effect: X = 0.12 (P = 0.45); P = 0% test for overall effect: X = 0.12 (P = 0.45); P = 0% test for overall effect: X = 0.12 (P = 0.45); P = 0% test for overall effect: X = 0.12 (P = 0.45); P = 0% test for overall effect: X = 0.12 (P = 0.45); P = 0% test for overall effect: X = 0.12 (P = 0.45); P = 0% test for overall effect: X = 0.12 (P =	MRC 1968	27	199	25	194	15.0%	1.05 [0.63 , 1.75]	
base corn oil 1965 5 28 1 26 1.0% $4.64(0.58, 37.15]$ TARS 1992 1 27 3 28 1.5% $0.35(0.04, 3.12]$ jump Diet-Heart 1978 37 221 25 2.37 16.0% $1.59(0.99, 2.55]$ eterans Admin 1969 57 4.24 81 422 22.1% $0.70(0.51, 0.96]$ biotoal 095% CD 2123 2128 100.0% $0.95(0.73, 1.25]$ of al events: 266 287 learcogneticy: Tarl = 0.06; Ch ² = 1.37, df = 6 (P = 0.04); P = 55% ets for overall effect: Z = 0.34 (P = 0.73) 2.62 replaced by MUFA tose olive 1965 3 26 1 26 100.0% $3.00(0.33, 26.99]$ biotoal 095% CD 226 26 100.0% $3.00(0.33, 26.99]$ biotoal 095% CD 26 26 100.0% $3.00(0.33, 26.99]$ biotoal 095% CD 26 26 100.0% $0.20(0.01, 4.15]$ ets for overall effect: Z = 0.98 (P = 0.33) 2.63 replace by CHO lack 1994 0 66 2 67 0.2% $0.20(0.01, 4.15]$ biotoal 095% CD 2074 30492 100.0% $0.99(0.85, 1.14]$ otical events: 316 4 (P = 0.48); P = 0% ets for overall effect: Z = 0.010; P = 0.551; H = 10% ets for overall effect: Z = 0.010; Ch ² = 3.51, df = 4 (P = 0.48); P = 0% ets for overall effect: Z = 0.010; Ch ² = 3.51, df = 4 (P = 0.48); P = 0% ets for overall effect: Z = 0.010; Ch ² = 3.51, df = 4 (P = 0.48); P = 0% ets for overall effect: Z = 0.010; Ch ² = 3.51, df = 4 (P = 0.48); P = 0% ets for overall effect: Z = 0.010; Ch ² = 3.51, df = 4 (P = 0.48); P = 0% ets for overall effect: Z = 0.010; Ch ² = 3.51, df = 4 (P = 0.48); P = 0% ets for overall effect: Z = 0.010; Ch ² = 3.51, df = 3 (P = 0.45); P = 0% ets for overall effect: Z = 0.012; Q = 0.90; 2.65 replacenent unclear biotoal 05% CD 20713 30464 100.03% 0.99 [0.86, 1.14] otal events: 0 Ch ² = 2.63, df = 3 (P = 0.45); P = 0% ets for overall effect: X = 0.012; Q = 0.90; 2.65 replacenent unclear biotoal 05% CD 0 0 eterogeneity; Narl = 0.00; Ch ² = 2.63, df = 3 (P = 0.45); P = 0% ets for overall effect: X = 0.12; Q = 0.90; 2.65 replacenent unclear biotoal 05% CD 0 0 eterogeneity; Ch and = 0, Di = 0.25; P = 0% ets for overall effect: X = 0.12; Q = 0.90; 2.65 replacenent unclear biotoal 05% CD 0 0 eterogeneity; Narl = 0.05; Ch ² = 1.05; df	Oslo Diet-Heart 1966	38	206	52	206	19.7%	0.73 [0.50 , 1.06]	
TARS 1992 1 27 3 28 1.5% 0.55 [0.04, 3.12] yhey Diet-Heart 1978 37 221 25 237 16.0% 1.59 [0.09, 2.55] eterans Admin 1969 57 424 81 422 22.1% 0.00% 0.05 [0.73, 1.25] bubtoal 05% CD 22123 2128 100.0% 0.95 [0.73, 1.25] vial events: 266 287 teterogeneity: Tur ² = 0.06; Ch ² = 1.3.7, df = 6 (P = 0.04); P = 55% est for overall effect: Z = 0.34 (P = 0.73) 2.62 replaced by MUFA teterogeneity: Not applicable test for overall effect: Z = 0.39 (P = 0.33) 2.63 replaced by MUFA teterogeneity: Not applicable test for overall effect: Z = 0.98 (P = 0.33) 2.63 replaced by MUFA teterogeneity: Not applicable test for overall effect: Z = 0.98 (P = 0.33) 2.63 replaced by MUFA teterogeneity: Not applicable test for overall effect: Z = 0.98 (P = 0.33) 2.63 replace by CHO take 1994 0 66 2 67 0.2% ART 1989 101 1018 100 1015 29.8% 1.01 [0.77, 1.31] 4.02 (0.00, 4, 1.19] 4.03 (0.03, 2.6.99] 1.04 (CM) 2 13 19541 320 29294 69.1% 1.00 [0.84, 1.19] 4.04 (95% CD) 20740 30492 100.0% 1.09 [0.85, 1.14] 4.05 (0.02, 1.14] 4.05 (0.02, 1.14] 4.05 (0.02, 1.14] 4.05 (0.02, 1.14] 4.05 (0.02, 1.14] 4.05 (0.02, 1.14] 4.05 (0.01, 4.15] 2.64 replaced by protein 1.64 (199 (Ch) 2.13 19541 320 29294 69.1% 1.00 [0.77, 1.31] 2.64 replaced by protein 1.64 (199 (Ch) 2.13 19541 320 29294 69.4% 1.00 [0.84, 1.19] 4.05 (0.10, 1.213 19541 320 29294 69.4% 1.00 [0.84, 1.19] 4.05 (0.02, 1.14] 4.05 (0.02, 1.14] 4.05 (0.02, 1.14] 4.05 (0.02, 1.14] 4.05 (0.02, 0.29 [0.00, 4, 1.15] 4.05 (0.02, 0.29 [0.00, 4, 1.15] 4.05 (0.00, 0.74 = 2.63, df = 3 (P = 0.45); P = 0% 1.05 (1.077, 1.31] 4.05 (0.00, 0.74 = 2.63, df = 3 (P = 0.45); P = 0% 1.05 (1.077, 1.31] 4.05 (0.00, 0.74 = 2.63, df = 3 (P = 0.45); P = 0% 1.05 (1.077, 1.31] 4.05 (0.00, 0.74 = 2.63, df = 3 (P = 0.45); P = 0% 1.05 (1.077, 1.31] 4.05 (0.00, 0.74 = 2.63, df = 3 (P = 0.45); P = 0% 1.05 (1.077, 1.31] 4.05 (0.00, 0.74 = 2.63, df = 3 (P = 0.75); P = 0% 1.05 (1.077, 1.31] 4.05 (0.00, 0.74 = 2.63, df = 3 (P = 0.75); P = 0% 1.07 (0.07, 1.14] 4.05 (0.00, 0.74	Rose corn oil 1965	5	28	1	26	1.6%	4.64 [0.58, 37.15]	
ydney Diet-Heart 1978 37 221 25 237 16.0% $1.59 [0.99, 2.55]$ (eterans Admin 1969 57 424 81 422 22.1% $0.70 [0.51, 0.96]$ 0.70 [0.51, 0.96] 0.70 [0.51, 0.16] 0.70 [0.1	STARS 1992	1	27	3	28	1.5%	0.35 [0.04, 3.12]	
The contrast Admin 1969 57 424 81 422 22.1% 0.70 [0.51, 0.96] 1.000% 0.95 [0.73, 1.25] 1.000% 0.95 [0.73, 1.25] 1.000% 0.95 [0.73, 1.25] 1.000% 0.95 [0.73, 1.25] 1.000% 0.95 [0.73, 1.25] 1.000% 0.95 [0.73, 1.25] 1.000% 0.95 [0.73, 1.25] 1.000% 0.95 [0.73, 1.25] 1.000% 0.95 [0.73, 1.25] 1.000\% 0.95 [0.73, 1.14] 1.000\% 0.95 [0.73, 1.14] 1.000\% 0	Sydney Diet-Heart 1978	37	221	25	237	16.0%	1.59 [0.99 , 2.55]	
ubiolal (95% CI) 2123 2128 100.0% 0.95 [0.73, 1.25] trait = 0.05; Ch ² = 1.37; df = 6 (P = 0.04); P = 55% test for overall effect: $Z = 0.34$ (P = 0.73) 2.62 replaced by MUFA tose olive 1965 3 26 1 26 100.0% 3.00 [0.33, 26.99] totolal (95% CI) 26 26 26 100.0% 3.00 [0.33, 26.99] totolal (95% CI) 26 26 100.0% 3.00 [0.33, 26.99] totolar events: 3 1 test for overall effect: $Z = 0.98$ (P = 0.33) 2.63 replace by CHO Stack 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] test for overall effect: $Z = 0.98$ (P = 0.33) 2.63 replace by CHO Stack 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] test for overall effect: $Z = 0.93$ (P = 0.48); P = 0% test for overall effect: $Z = 0.99$ (P = 0.85) 2.64 replaced by protein Stack 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] test for overall effect: $Z = 0.19$ (P = 0.48); P = 0% test for overall effect: $Z = 0.19$ (P = 0.85) 2.64 replaced by protein Stack 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] test for overall effect: $Z = 0.19$ (P = 0.85); P = 0% test for overall effect: $Z = 0.19$ (P = 0.45); P = 0% test for overall effect: $Z = 0.12$ (P = 0.45); P = 0% test for overall effect: $Z = 0.12$ (P = 0.45); P = 0% test for overall effect: $Z = 0.12$ (P = 0.45); P = 0% test for overall effect: $Z = 0.12$ (P = 0.45); P = 0% test for overall effect: $Z = 0.12$ (P = 0.90) 2.25 replacement unclear ubiotal (95% CI) 0 0 0 Not estimable test for overall effect: $Z = 0.12$ (P = 0.90) 2.25 replacement unclear ubiotal (95% CI) 0 0 0 State events: 0 0 0 teroremetry: Not applicable test for overall effect: $Z = 0.12$ (P = 0.79), P = 0% test for overall effect: $Z = 0.12$ (P = 0.79), I = 0% test for overall effect: $Z = 0.12$ (P = 0.79), I = 0% test for overall effect: Not applicable test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I = 0% test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I = 0% test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I = 0% test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I = 0% test for	Veterans Admin 1969	57	424	81	422	22.1%	0.70 [0.51, 0.96]	-
ional events: 266 287 leterogeneity: Tau ² = 0.06; Ch ² = 13.37, df = 6 (P = 0.04); P = 55% iest for overall effect: Z = 0.34 (P = 0.73) 26.2 replaced by MUFA tore cuive 1965 3 26 1 26 100.0% 3.00 [0.33, 26.99] obtoal (95% CI) 26 26 26 100.0% 3.00 [0.33, 26.99] obtoal events: 3 1 1 leterogeneity: Not applicable leterogeneity: Not applicable leterogeneity: Not applicable 101 1018 100 1015 29.8% 1.01 [0.77, 1.31] 27.3 replace by CHO 102 004 1 1 88 4 88 0.4% 0.25 [0.03, 2.19] obtoal (95% CI) 20740 30492 100.0% 0.99 [0.85, 1.14] obtoal (95% CI) 20740 30492 100.0% 0.99 [0.85, 1.14] obtoal (95% CI) 20740 30492 100.0% 0.99 [0.85, 1.14] 26.4 replace by protein 102 0(1.84, 1.19] 104 cvents: 316 429 Heterogeneity: Tau ² = 0.00; Ch ² = 3.51, df = 4 (P = 0.48); P = 0% let for overall effect: Z = 0.19 (P = 0.85) 26.4 replaced by protein 104 cvents: 315 426 leterogeneity: Tau ² = 0.00; Ch ² = 2.63, df = 3 (P = 0.48); P = 0% let for overall effect: Z = 0.12 (P = 0.90) 26.5 replacement unclear ubtoal (95% CI) 0 0 0 Not estimable leterogeneity: Not applicable leterogeneity: Not applicable leteroff or Not estim	Subtotal (95% CI)		2123		2128	100.0%	0.95 [0.73, 1.25]	
The reprogeneity: Tau ² = 0.06; Ch ² = 13.37, df = 6 (P = 0.04); P = 55% test for overall effect: Z = 0.34 (P = 0.73) 2.62 replaced by MUFA tose olive 1965 3 26 1 26 100.0% 3.00 [0.33, 26.99] volate events: 3 1 1 feerogeneity: Not applicable test for overall effect: Z = 0.98 (P = 0.33) 2.63 replace by CHO takek 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] output 1018 100 1015 29.8% 1.01 [0.77, 1.31] output 2004 1 88 4 88 0.4% 0.35 [0.04, 3.12] 1.11 1018 100 1015 29.8% 1.01 [0.77, 1.31] output 2014 1 213 19541 320 29294 69.1% 1.00 [0.84, 1.19] output 2015% CD 20740 30492 100.0% 0.99 [0.85, 1.14] output 2016 (D) 213 19541 320 29294 69.1% 1.00 [0.84, 1.19] output 2005 (Ch) 2 20740 30492 100.0% 0.29 [0.03, 1.14] 1.11 1018 100 1015 29.9% 1.01 [0.77, 1.31] 1.12 101 1018 100 1015 29.9% 1.01 [0.77, 1.31] 1.12 10541 320 29294 69.4% 1.00 [0.84, 1.19] 1.13 10541 320 29294 69.4% 1.00 [0.84, 1.19] 1.14 1018 100 1015 29.9% 1.01 [0.77, 1.31] 1.15 426 thereogeneity: Tau ² = 0.00; Chi ² = 3.51, df = 3 (P = 0.45); P = 0% test for overall effect: Z = 0.19 (P = 0.35) 2.64 replaced by protein takek 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] 1.14 1.19 1.10 [0.77, 1.31] 1.11 1.11	Fotal events:	266		287				Ţ
The set for overall effect: $Z = 0.34$ (P = 0.73) 2.62 replaced by MUFA toose olive 1965 3 26 1 26 100.0% 3.00 [0.33, 26.99] 3.00 [0.33, 26.99] 3.01 [0.77, 1.31] 4.01 [0.77, 1.31] 4.02 [0.01, 4.15] 4.02 [0.01, 4.15] 4.01 [0.77, 1.31] 4.01 [0.77, 1.31] 4.01 [0.77, 1.31] 4.01 [0.77, 1.31] 4.01 [0.08, 1.19] 4.01 [0.18, 1.00] 4.11 [0.10] [0.77, 1.31] 4.11 [0.10] [0.77, 1.31] 4.11 [0.10] [0.77, 1.31] 4.11 [0.10] [0.14, 1.19] 4.11 [0.10] [0.14, 1.19] 4.11 [0.10] [0.14, 1.19] 4.11 [0.10] [0.16] [0.10] [0.16	Heterogeneity: Tau ² = 0.06: ($Chi^2 = 13.37$	df = 6 (P)	= 0.04); I ²	= 55%			
2.6.2 replaced by MUFA toxe olive 1965 3 2 6 1 26 100.0% 3.00 [0.33, 26.99] ubtotal (95% CI) 26 26 26 100.0% 3.00 [0.33, 26.99] 3.00 [0.33, 26.99] violal events: 3 1 leterogeneity: Not applicable 'est for overall effect: Z = 0.98 (P = 0.33) 2.6.3 replace by CHO Mack 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] 4.8 4 88 0.4% 0.25 [0.03, 2.19] ANRT 1989 101 1018 100 1015 29.9% 1.01 [0.77, 1.31] ey 2004 1 88 4 88 0.4% 0.25 [0.03, 2.19] ubtotal (95% CI) 20740 30492 100.0% 0.99 [0.85, 1.14] violal events: 316 429 leterogeneity: Tau ² = 0.01; (P = 3.51, df = 4 (P = 0.48); P = 0% 'est for overall effect: Z = 0.19 (P = 0.85) 2.6.4 replaced by protein Black 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] violal events: 316 429 leterogeneity: Tau ² = 0.01; (P = 3.51, df = 4 (P = 0.48); P = 0% 'est for overall effect: Z = 0.19 (P = 0.85) 2.6.4 replaced by protein Black 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] violal events: 315 426 Violal events: 315 426 Violal events: 315 426 leterogeneity: Tau ² = 0.00; Chi ² = 2.63, df = 3 (P = 0.45); P = 0% 'est for overall effect: Z = 0.12 (P = 0.90) 2.6.5 replacement unclear ubtotal (95% CI) 0 0 0 Not estimable 'est for overall effect: Z = 0.12 (P = 0.90) 2.6.5 replacement unclear ubtotal (95% CI) 0 0 leterogeneity: Not applicable 'est for overall effect: Z = 0.12, (P = 0.90) 2.6.5 replacement unclear ubtotal (95% CI) 0 0 leterogeneity: Not applicable 'est for overall effect: Z = 0.12, df = 3 (P = 0.79), P = 0% 'est for overall effect: S = 0.12, df = 3 (P = 0.79), P = 0% 'est for overall effect: S = 0.12, df = 3 (P = 0.79), P = 0% 'est for overall effect: S = 0.12, df = 3 (P = 0.79), P = 0% 'est for overall effect: S = 0.12, df = 3 (P = 0.97), P = 0% 'est for overall effect: S = 0.12, df = 3 (P = 0.97), P = 0% 'est for overall effect: S = 0.12, df = 3 (P = 0.97), P = 0% 'est for overall effect: S = 0.12, df = 3 (P = 0.97), P = 0% 'est for overall effect: S = 0.12, df = 3 (P = 0.97), P = 0% 'est for overall effect: S = 0.12, df = 3 (P = 0.97), P = 0%	Test for overall effect: $Z = 0$.	34 (P = 0.73)	5)	,, -				
to see olive 1965 3 26 1 26 100.0% 3.00 [0.33, 26.99] ubtool (95% CI) 26 26 100.0% 3.00 [0.33, 26.99] abtool (95% CI) 26 26 7 0.2% 3.00 [0.33, 26.99] 26.3 replace by CHO Hack 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] ey 2004 1 88 4 88 0.4% 0.25 [0.03, 2.19] abtool (1) 213 19541 320 29294 69.1% 1.00 [0.84, 1.19] ubtool (95% CI) 20740 30492 100.0% 0.99 [0.85, 1.14] ical events: 316 429 leterogeneity: Tau ² = 0.01; 0'F = 351, df = 4 (P = 0.48); P = 0% icst for overall effect: Z = 0.19 (P = 0.85) 2.64 replaced by protein Hack 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] abtool (95% CI) 20740 30492 100.0% 0.99 [0.85, 1.14] ical events: 316 429 leterogeneity: Tau ² = 0.01; 0'F = 351, df = 4 (P = 0.48); P = 0% icst for overall effect: Z = 0.19 (P = 0.85) 2.64 replaced by protein Hack 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] abtool (95% CI) 20713 30464 100.0% 0.99 [0.86, 1.14] ical events: 315 426 heterogeneity: Tau ² = 0.00; Chi ² = 2.63, df = 3 (P = 0.45); P = 0% icst for overall effect: Z = 0.12 (P = 0.90) 2.65 replacement unclear ubtool (95% CI) 0 0 0 Not estimable ical events: 0 0 0 Heterogeneity: Not applicable icst for overall effect: Z = 0.12, df = 3 (P = 0.79), P = 0% icst for overall effect: Z = 0.12, df = 3 (P = 0.79), P = 0% icst for overall effect: C = 0.12, df = 3 (P = 0.79), P = 0% icst for overall effect: C = 0.12, df = 3 (P = 0.79), P = 0% icst for overall effect: C = 0.12, df = 3 (P = 0.79), P = 0% icst for overall effect: C = 0.12, df = 3 (P = 0.79), P = 0% icst for overall effect: C = 0.12, df = 3 (P = 0.79), P = 0% icst for overall effect: C = 0.12, df = 3 (P = 0.79), P = 0% icst for overall effect: C = 0.12, df = 3 (P = 0.79), P = 0% icst for overall effect: C = 0.12, df = 3 (P = 0.79), P = 0% icst for overall effect: C = 0.12, df = 3 (P = 0.79), P = 0% icst for overall effect: S = 0.12, df = 3 (P = 0.79), P = 0% icst for overall effect: S = 0.12, df = 3 (P = 0.79), P = 0% icst for overall effect: S = 0.12, df = 3 (P = 0.79), P = 0% icst for overall effect: S = 0.1	1.26.2 replaced by MUFA							
abtotal (95% CI) 26 26 100.0% $3.00 [0.33, 26.99]$ fold events: 3 1 leterogeneity: Not applicable is for overall effect: $Z = 0.98 (P = 0.33)$ 26.3 replace by CHO 101 1018 100 1015 29.8% 1.01 [0.77, 1.51] action overall effect: $Z = 0.98 (P = 0.33)$ 2.6 20 0.01, 4.15] 1.01 [0.77, 1.51] action overall effect: $Z = 0.98 (P = 0.33)$ 1.88 4 88 0.4% 0.25 [0.03, 2.19] Actin 1989 101 1018 100 1015 29.8% 1.01 [0.77, 1.51] abtotal (95% CI) 20740 30492 100.0% 0.99 [0.85, 1.14] abtotal (95% CI) 20740 30492 100.0% 0.99 [0.85, 1.14] abtotal (95% CI) 20713 30464 100.0% 0.29 [0.01, 4.15] abtotal (95% CI) 20713 30464 100.0% 0.99 [0.86, 1.14] abtotal (95% CI) 20713 30464 100.0% 0.99 [0.86, 1.14] abtotal (95% CI) 0 0 0 Not estimable otal events: 0 0 0 <td>Rose olive 1965</td> <td>3</td> <td>26</td> <td>1</td> <td>26</td> <td>100.0%</td> <td>3.00 [0.33 , 26.99]</td> <td></td>	Rose olive 1965	3	26	1	26	100.0%	3.00 [0.33 , 26.99]	
bial events: 3 1 leterogeneity: Not applicable iest for overall effect: $Z = 0.98$ (P = 0.33) 2.6.3 replace by CHO lack 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] ART 1989 101 1018 100 1015 29.8% 101 [0.77, 1.31] eve 2004 1 88 4 88 0.4% 0.25 [0.03, 2.19] TARS 1992 1 27 3 28 0.4% 0.35 [0.04, 3.12] VHI 2006 (1) 213 19541 320 29294 69.1% 1.00 [0.84, 1.19] ubtotal (95% CT) 20740 30492 100.0% 0.99 [0.85, 1.14] Vest for overall effect: $Z = 0.19$ (P = 0.85) 2.6.4 replaced by protein Black 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] ART 1989 101 1018 100 1015 29.9% 1.01 [0.77, 1.31] eve 2004 1 88 4 88 0.4% 0.25 [0.03, 2.19] VHI 2006 (1) 213 19541 320 29294 69.4% 1.00 [0.84, 1.19] ubtotal (95% CT) 20713 30464 100.0% 0.99 [0.86, 1.14] vest for overall effect: $Z = 0.12$ (P = 0.45); P = 0% vest for overall effect: $Z = 0.12$ (P = 0.90) 2.6.5 replacement unclear ubtotal (95% CT) 0 0 0 Not estimable vest for overall effect: $Z = 0.12$ (P = 0.90) 2.6.5 replacement unclear ubtotal (95% CT) 0 0 0 leterogeneity: Not applicable vest for overall effect: Not applicable vest for subgroup differences: Ch ² = 1.05, df = 3 (P = 0.79), P = 0% vest for overall effect: Not applicable	Subtotal (95% CI)	-	26	-	 26	100.0%	3.00 [0.33 . 26.99]	
The terrogeneity: Not applicable Test for overall effect: $Z = 0.98$ (P = 0.33) 2.6.3 replace by CHO Nack 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] (a) 101 1018 100 1015 29.8% 1.01 [0.77, 1.31] (b) 20204 1 88 4 88 0.4% 0.25 [0.03, 21.09] (c) 2004 1 88 4 88 0.4% 0.25 [0.03, 21.09] (c) 2004 1 88 4 88 0.4% 0.25 [0.03, 21.09] (c) 213 19541 320 29294 69.1% 1.00 [0.84, 1.19] (c) 20740 30492 100.0% 0.99 [0.85, 1.14] (c) 20740 1018 100 1015 29.9% 1.01 [0.77, 1.31] (c) 2004 1 88 4 88 0.4% 0.25 [0.03, 2.19] (c) 2004 1 88 4 88 0.4% 0.25 [0.03, 2.19] (c) 2004 1 88 4 88 0.4% 0.25 [0.03, 2.19] (c) 2004 1 88 4 88 0.4% 0.99 [0.86, 1.14] (c) 206 (1) 213 19541 320 29294 69.4% 1.00 [0.84, 1.19] (c) 2004 1 88 4 26 (c) 20713 30464 100.0% 0.99 [0.86, 1.14] (c) 206 (2) 213 19541 320 29294 69.4% 1.00 [0.84, 1.19] (c) 206 (2) 20713 30464 100.0% 0.99 [0.86, 1.14] (c) 206 (2) 217 (2) 20713 30464 100.0% 0.99 [0.86, 1.14] (c) 206 (2) 213 19541 320 29294 69.4% 1.00 [0.84, 1.19] (c) 206 (2) 213 19541 320 29294 69.4% 1.00 [0.84, 1.19] (c) 206 (2) 213 19541 320 29294 69.4% 1.00 [0.84, 1.19] (c) 206 (2) 213 19541 320 29294 69.4% 1.00 [0.84, 1.19] (c) 206 (2) 213 19541 320 29294 69.4% 1.00 [0.84, 1.19] (c) 206 (2) 213 19541 320 29294 69.4% 1.00 [0.84, 1.19] (c) 206 (2) 20713 30464 100.0% 0.99 [0.86, 1.14] (c) 206 (2) 20713 2040 (2) 20713 20713 20713 20713 20713 20713 20713 20713 20713 20713 20713 20713 20713 20713 20713 20713 2071	Fotal events:	3	_0	1	20			
The star overall effect: $Z = 0.98 (P = 0.33)$ 2.6.3 replace by CHO Back 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] OART 1989 101 1018 100 1015 29.8% 1.01 [0.77, 1.31] () 2004 1 88 4 88 0.4% 0.25 [0.03, 2.19] () 213 19541 320 29294 69.1% 1.00 [0.84, 1.19] () 213 19541 320 29294 69.1% 1.00 [0.84, 1.19] () 2014 events: 316 429 leterogeneity: Tau ² = 0.00; Chi ² = 3.51, df = 4 (P = 0.48); P = 0% () 264 replaced by protein Hack 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] () 264 replaced by protein Hack 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] () 264 replaced by protein Hack 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] () 204 1 88 4 88 0.4% 0.25 [0.03, 2.19] () 205 (1) 213 19541 320 29294 69.4% 1.00 [0.84, 1.19] () 206 (1) 213 19541 320 29294 69.4% 1.00 [0.84, 1.19] () 206 (1) 213 19541 320 29294 69.4% 1.00 [0.84, 1.19] () 20713 30464 100.0% 0.99 [0.85, 1.14] () 208 (1) 213 19541 320 29294 69.4% 1.00 [0.84, 1.19] () 208 (1) 213 19541 320 29294 69.4% 1.00 [0.84, 1.19] () 208 (1) 213 19541 320 29294 69.4% 1.00 [0.84, 1.19] () 208 (1) 213 19541 320 29294 69.4% 1.00 [0.86, 1.14] () 214 events: 315 426 leterogeneity: Tau ² = 0.00; Chi ² = 2.63, df = 3 (P = 0.45); P = 0% () 265 replacement unclear () 206 (1) 217 = 2.63, df = 3 (P = 0.45); P = 0% () 265 replacement unclear () 0 0 () 1 0 1 0 1 0 0 () 216 events: 0 0 0 () 265 ct) 0 0 () 265 ct) 0 0 () 26	Heterogeneity: Not applicable	e		1				
2.6.3 replace by CHO Hack 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] PART 1989 101 1018 100 1015 29.8% 1.01 [0.77, 1.31] PART 1989 101 213 19541 320 29294 69.1% 1.00 [0.84, 1.19] TARS 1992 1 27 3 28 0.4% 0.35 [0.04, 3.12] TARS 1992 1 277 40 30492 100.0% 0.99 [0.85, 1.14] VHI 2006 (1) 213 19541 320 29294 69.1% 1.00 [0.84, 1.19] PART 1989 101 1018 100 1015 29.9% 1.01 [0.77, 1.31] PART 1989 101 1018 100 0 1015 29.9% 1.01 [0.77, 1.31] PART 1989 101 1018 100 1015 29.9% 1.01 [0.77, 1.31] PART 1989 101 1018 100 0 1015 29.9% 1.01 [0.77, 1.31] PART 1989 101 1018 100 0 1015 29.9% 1.01 [0.77, 1.31] PART 1980 100 0 $PART 1980$ 100	Test for overall effect: $Z = 0$	- 98 (P = 0.33	5)					
2.6.3 replace by CHO Black 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] y 2004 1 88 4 88 0.4% 0.35 [0.04, 3.12] TARS 1992 1 27 3 28 0.4% 0.35 [0.04, 3.12] TARS 1992 1 27 3 28 0.4% 0.35 [0.04, 3.12] VHI 2006 (1) 213 19541 320 29294 69.1% 1.00 [0.84, 1.19] volat events: 316 429 100.0% 0.99 [0.85, 1.14] 100 volat events: 316 429 100.0% 0.99 [0.85, 1.14] 100 volat events: 316 429 100.0% 0.99 [0.85, 1.14] 100 volat vents: 316 429 0 66 2 67 0.20 [0.01, 4.15] 100 volat vents: 315 426 100.0% 0.25 [0.03, 2.19] 100 100 100 100 100 100 100 100 100 100 100 100 100 100 100 100		(= 0.00	/					
Mack 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] ey 2004 1 88 4 88 0.4% 0.25 [0.03, 2.19] ey 2004 1 27 3 28 0.4% 0.35 [0.04, 3.12] VHI 2006 (1) 213 19541 320 29294 69.1% 1.00 [0.84, 1.19] value vents: 316 429 Iterregeneity: Tau ² = 0.00; Chi ² = 3.51, df = 4 (P = 0.48); l ² = 0% vest for overall effect: Z = 0.19 (P = 0.85) 26.4 replaced by protein Mack 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] vest for overall effect: Z = 0.19 (P = 0.48); l2 = 0% $vest for overall effect: Z = 0.19 (P = 0.48); l2 = 0%vest for overall effect: Z = 0.00; Chi2 = 2.63, df = 3 (P = 0.45); l2 = 0% vest for overall effect: Z = 0.12 (P = 0.90)vest for overall effect: Z = 0.12 (P = 0.90)vest for overall effect: S = 0.12 (P = 0.90)vest for overall effect: C = 1.05, df = 3 (P = 0.79), l2 = 0%vest for overall effect: S tot applicable vest for overall effect: S tot applicablevest for overall effect: S tot applicable vest for overall effect: S tot applicablevest for overall effect: S tot applicable vest for subgroup differences: Chi2 = 1.05, df = 3 (P = 0.79), l2 = 0%$	1.26.3 replace by CHO							
AART 1989 101 1018 100 1015 29.8% 1.01 [0.77, 1.31] exp 2004 1 88 4 88 0.4% 0.25 [0.03, 2.19] TARS 1992 1 27 3 28 0.4% 0.35 [0.04, 3.12] WH 2006 (1) 213 19541 320 29294 69.1% 1.00 [0.84, 1.19] Whotal (95% CI) 20740 30492 100.0% 0.99 [0.85, 1.14] We can be calculated by protein Jack 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] exp 2004 1 88 4 88 0.4% 0.25 [0.03, 2.19] WH 2006 (1) 213 19541 320 29294 69.4% 1.00 [0.84, 1.19] Whotal (95% CI) 20713 30464 100.0% 0.99 [0.86, 1.14] Wh 2006 (1) 213 19541 320 29294 69.4% 1.00 [0.84, 1.19] Wh 2006 (1) 213 19541 320 29294 69.4% 1.00 [0.84, 1.19] Wh 2006 (1) 213 19541 320 29294 69.4% 1.00 [0.84, 1.19] Wh 2006 (1) 213 19541 320 29294 69.4% 1.00 [0.84, 1.19] Wh 2006 (1) 213 19541 320 29294 69.4% 1.00 [0.84, 1.19] Wh 2006 (1) 213 19541 320 29294 69.4% 1.00 [0.84, 1.19] Wh 2006 (1) 213 19541 320 29294 69.4% 1.00 [0.84, 1.19] Wh 2006 (1) 213 19541 320 29294 69.4% 1.00 [0.84, 1.19] Wh 2006 (1) 213 19541 320 29294 69.4% 1.00 [0.84, 1.19] Wh 2006 (1) 213 19541 320 29294 69.4% 1.00 [0.84, 1.19] Wh 2006 (1) 20713 30464 100.0% 0.99 [0.86, 1.14] Wh 2016 (15% CI) 0 0 0 Reterogeneity: Not applicable Test for overall effect: X ot applicable Test for overall effect: Not applicable Test for overall effect: Not applicable Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi	Black 1994	0	66	2	67	0.2%	0.20 [0.01 , 4.15]	• • •
ey 2004 1 88 4 88 0.4% 0.25 [0.03, 2.19] TARS 1992 1 27 3 28 0.4% 0.35 [0.04, 3.12] WH 2006 (1) 213 19541 320 29294 69.1% 1.00 [0.84, 1.19] votal events: 316 429 leterogeneity: Tau ² = 0.00; Chi ² = 3.51, df = 4 (P = 0.48); P = 0% 'est for overall effect: Z = 0.19 (P = 0.85) .26.4 replaced by protein Black 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] votal events: 315 426 leterogeneity: Tau ² = 0.00; Chi ² = 3.51, df = 4 (P = 0.48); P = 0% 'est for overall effect: Z = 0.19 (P = 0.48); P = 0% 'est for overall effect: Z = 0.19 (P = 0.48); P = 0% 'at the tau is the ta	DART 1989	101	1018	100	1015	29.8%	1.01 [0.77, 1.31]	+
TARS 1992 1 27 3 28 0.4% 0.35 [0.04, 3.12] VHI 2006 (1) 213 19541 320 29294 69.1% 1.00 [0.84, 1.19] bubtal (95% CI) 20740 30492 100.0% 0.99 [0.85, 1.14] value vents: 316 429 teterogeneity: Tau ² = 0.00; Chi ² = 3.51, df = 4 (P = 0.48); P = 0% vest for overall effect: Z = 0.19 (P = 0.85) .26.4 replaced by protein Black 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] JART 1989 101 1018 100 1015 29.9% 1.01 [0.77, 1.31] Levy 2004 1 88 4 88 0.4% 0.25 [0.03, 2.19] VHI 2006 (1) 213 19541 320 29294 69.4% 1.00 [0.84, 1.19] Jubtotal (95% CI) 20713 30464 100.0% 0.99 [0.86, 1.14] Votal events: 315 426 teterogeneity: Tau ² = 0.00; Chi ² = 2.63, df = 3 (P = 0.45); P = 0% .26.5 replacement unclear ubtotal (95% CI) 0 0 0 .26.5 replacement unclear ubtotal (95% CI) 0 .26.5 replacement unclear .26.5 replacement unclear .27.5 m .26.5 replacement unclear .26.5 replacement unclear .27.5 m .26.5 replacement unclear .27.5 m .27.5 	Ley 2004	1	88	4	88	0.4%	0.25 [0.03 , 2.19]	-
WHI 2006 (1) 213 19541 320 29294 69.1% 1.00 [0.84, 1.19] jabotal (95% CI) 20740 30492 100.0% 0.99 [0.85, 1.14] i otal events: 316 429 feterogeneity: Tau ² = 0.00; Chi ² = 3.51, df = 4 (P = 0.48); P = 0% leterogeneity: Tau ² = 0.00; Chi ² = 3.51, df = 4 (P = 0.48); P = 0% leterogeneity: Tau ² = 0.00; Chi ² = 3.51, df = 4 (P = 0.48); P = 0% leterogeneity: Tau ² = 0.019 (P = 0.85) .26.4 replaced by protein Black 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] .26.4 replaced by protein Black 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] .26.4 replaced by protein Black 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] .29.204 1 88 4 88 0.4% 0.25 [0.03, 2.19] .40.206 (1) 213 19541 320 29294 69.4% 1.00 [0.84, 1.19] .40.206 (1) 213 19541 320 29294 69.4% 1.00 [0.84, 1.19] .40.206 (1) 213 19541 320 29294 69.4% 1.00 [0.84, 1.19] .40.207 13 30464 100.0% 0.99 [0.86, 1.14] .41.200 (1) 213 19541 320 29294 69.4% 1.00 [0.84, 1.19] .42.5 replacement unclear .44.206 leterogeneity: Tau ² = 0.012 (P = 0.90) .26.5 replacement unclear .44.206 leterogeneity: Not applicable leterogeneity: Not applicable leterogeneit	STARS 1992	1	27	3	28	0.4%	0.35 [0.04 , 3.12]	
Subtotal (95% CI) 20740 30492 100.0% 0.99 [0.85, 1.14] Otal events: 316 429 Heterogeneity: Tau ² = 0.00; Chi ² = 3.51 , df = 4 (P = 0.48); l ² = 0%	WHI 2006 (1)	213	19541	320	29294	69.1%	1.00 [0.84 , 1.19]	
Total events: 316 429 Ideterogeneity: Tau ² = 0.00; Chi ² = 3.51, df = 4 (P = 0.48); I ² = 0% Pest for overall effect: Z = 0.19 (P = 0.85) 26.4 replaced by protein Black 1994 0 66 2 67 0.20 [0.01, 4.15] ART 1989 101 1018 100 1015 29.9% 1.01 [0.77, 1.31] ey 2004 1 88 4 88 0.4% 0.25 [0.03, 2.19] VHI 2006 (1) 213 19541 320 29294 69.4% 1.00 [0.84, 1.19] Subtotal (95% CI) 20713 30464 100.0% 0.99 [0.86, 1.14] Votal events: 315 426 Ieterogeneity: Tau ² = 0.00; Chi ² = 2.63, df = 3 (P = 0.45); I ² = 0% Vettotal (95% CI) 0 0 .26.5 replacement unclear	Subtotal (95% CI)		20740		30492	100.0%	0.99 [0.85 , 1.14]	
Heterogeneity: $Tau^2 = 0.00$; $Chi^2 = 3.51$, $df = 4$ (P = 0.48); $l^2 = 0\%$ Perform the event of the even	Fotal events:	316		429				
The set for overall effect: $Z = 0.19$ (P = 0.85) .26.4 replaced by protein Black 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] DART 1989 101 1018 100 1015 29.9% 1.01 [0.77, 1.31] Lev 2004 1 88 4 88 0.4% 0.25 [0.03, 2.19] WH 2006 (1) 213 19541 320 29294 69.4% 1.00 [0.84, 1.19] Divibitotal (95% CI) 20713 30464 100.0% 0.99 [0.86, 1.14] Total events: 315 426 Teletrogeneity: Tau ² = 0.00; Chi ² = 2.63, df = 3 (P = 0.45); I ² = 0% Test for overall effect: $Z = 0.12$ (P = 0.90) .26.5 replacement unclear Divibitotal (95% CI) 0 0 0 Not estimable Total events: 0 0 0 Teletrogeneity: Not applicable Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ²	Heterogeneity: Tau ² = 0.00; G	Chi ² = 3.51,	df = 4 (P =	= 0.48); I ² =	= 0%			
26.4 replaced by protein Black 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] ART 1989 101 1018 100 1015 29.9% 1.01 [0.77, 1.31] ey 2004 1 88 4 88 0.4% 0.25 [0.03, 2.19] without (95% CI) 20713 30464 100.0% 0.99 [0.86, 1.14] ioutotal (95% CI) 20713 30464 100.0% 0.99 [0.86, 1.14] ioutotal (95% CI) 20.3 df = 3 (P = 0.45); P = 0% 0.55 replacement unclear iubtotal (95% CI) 0 0 Not estimable oftal events: 0 0 0 .26.5 replacement unclear 0 0 0 ubtotal (95% CI) 0 0 0 .26.5 replacement unclear 0 0 0 ubtotal (95% CI) 0 0 0 ieterogeneity: Not applicable	Test for overall effect: $Z = 0$.	19 (P = 0.85	i)					
Black 1994 0 66 2 67 0.2% 0.20 [0.01, 4.15] DART 1989 101 1018 100 1015 29.9% 1.01 [0.77, 1.31] ey 2004 1 88 4 88 0.4% 0.25 [0.03, 2.19] WH 2006 (1) 213 19541 320 29294 69.4% 1.00 [0.84, 1.19] bubtotal (95% CI) 20713 30464 100.0% 0.99 [0.86, 1.14] otal events: 315 426 leterogeneity: Tau ² = 0.00; Chi ² = 2.63, df = 3 (P = 0.45); I ² = 0% Test for overall effect: $Z = 0.12$ (P = 0.90) 26.5 replacement unclear bubtotal (95% CI) 0 0 0 Not estimable Otal events: 0 0 0 leterogeneity: Not applicable Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for su	1.26.4 replaced by protein							
DART 1989 101 1018 100 1015 29.9% 1.01 $[0.77, 1.31]$ Ley 2004 1 88 4 88 0.4% 0.25 $[0.03, 2.19]$ VHI 2006 (1) 213 19541 320 29294 69.4% 1.00 $[0.84, 1.19]$ Subtotal (95% CI) 20713 30464 100.0% 0.99 $[0.86, 1.14]$ Total events: 315 426 Heterogeneity: Tau ² = 0.00; Chi ² = 2.63, df = 3 (P = 0.45); l ² = 0% Test for overall effect: Z = 0.12 (P = 0.90) 26.5 replacement unclear Subtotal (95% CI) 0 0 0 Not estimable Total events: 0 0 0 Heterogeneity: Not applicable Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), l ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), l ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), l ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), l ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), l ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), l ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), l ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), l ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), l ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), l ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), l ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), l ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), l ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), l ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), l ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), l ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), l ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), l ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), l ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), l ² = 0% Test for subgroup differences: Chi ² = 1.05,	3lack 1994	0	66	2	67	0.2%	0.20 [0.01 , 4.15]	←
ey 2004 1 88 4 88 0.4% 0.25 [0.03, 2.19] VHI 2006 (1) 213 19541 320 29294 69.4% 1.00 [0.84, 1.19] Subtotal (95% CI) 20713 30464 100.0% 0.99 [0.86, 1.14] Total events: 315 426 Heterogeneity: Tau ² = 0.00; Chi ² = 2.63, df = 3 (P = 0.45); I ² = 0% Test for overall effect: Z = 0.12 (P = 0.90) -26.5 replacement unclear Subtotal (95% CI) 0 O 0 Total events: 0 O 0 State events: 0 O 0 Total events: 0 Yest for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Yest for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Total events 0.01 0.1 1 Total events 0.01 0.1 1 10 </td <td>DART 1989</td> <td>101</td> <td>1018</td> <td>100</td> <td>1015</td> <td>29.9%</td> <td>1.01 [0.77 , 1.31]</td> <td>+</td>	DART 1989	101	1018	100	1015	29.9%	1.01 [0.77 , 1.31]	+
VHI 2006 (1) 213 19541 320 29294 69.4% 1.00 [0.84, 1.19] Subtotal (95% CI) 20713 30464 100.0% 0.99 [0.86, 1.14] Total events: 315 426 Heterogeneity: Tau ² = 0.00; Chi ² = 2.63, df = 3 (P = 0.45); I ² = 0% Not estimable Post for overall effect: Z = 0.12 (P = 0.90) 0 Not estimable State of events: 0 0 Not estimable Post for overall effect: Not applicable $0 = 0$ Not estimable Post for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Favours lower SFA Favours high	Ley 2004	1	88	4	88	0.4%	0.25 [0.03 , 2.19]	
Subtotal (95% CI) 20713 30464 100.0% 0.99 [0.86, 1.14] For the events: 315 426 Heterogeneity: Tau ² = 0.00; Chi ² = 2.63, df = 3 (P = 0.45); I ² = 0% Performent unclear Performent unclear Subtotal (95% CI) 0 0 Not estimable Not estimable Potate events: 0 0 Protected 0 0 Protected 0.01 0.1 Protectes 0.01 0.1	WHI 2006 (1)	213	19541	320	29294	69.4%	1.00 [0.84 , 1.19]	•
Total events: 315 426 Heterogeneity:Tau ² = 0.00; Chi ² = 2.63, df = 3 (P = 0.45); I ² = 0%Peterogeneity:Tau ² = 0.12 (P = 0.90).26.5 replacement unclearSubtotal (95% CI)000Not estimablePoteorement:0O0Heterogeneity:Not estimablePoteorement:0Poteorement: <td>Subtotal (95% CI)</td> <td></td> <td>20713</td> <td></td> <td>30464</td> <td>100.0%</td> <td>0.99 [0.86 , 1.14]</td> <td>₹</td>	Subtotal (95% CI)		20713		30464	100.0%	0.99 [0.86 , 1.14]	₹
Heterogeneity: Tau ² = 0.00; Chi ² = 2.63, df = 3 (P = 0.45); I ² = 0% 'est for overall effect: Z = 0.12 (P = 0.90) .26.5 replacement unclear bubtotal (95% CI) 0 0 0 'otal events: 0 'est for overall effect: Not applicable 'est for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% 'est for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% 'Potnotes	Fotal events:	315		426				Ĭ
Test for overall effect: $Z = 0.12$ (P = 0.90) .26.5 replacement unclear fubtotal (95% CI) 0 0 Not estimable Total events: 0 0 Heterogeneity: Not applicable Test for overall effect: Not applicable Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0%	Heterogeneity: Tau ² = 0.00; C	Chi ² = 2.63,	df = 3 (P =	= 0.45); I ² =	= 0%			
.26.5 replacement unclear Not estimable Subtotal (95% CI) 0 0 Not estimable Not estimable Total events: 0 0 Iderogeneity: Not applicable 0 0 Test for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% 0 0 Protectes 0 0 0 Protectes 0 0 0	Test for overall effect: $Z = 0$.	12 (P = 0.90))					
Subtotal (95% CI) 0 0 Not estimable 'otal events: 0 0 0 Heterogeneity: Not applicable	1.26.5 replacement unclear							
Yotal events: 0 0 Ideterogeneity: Not applicable Ideterogeneity: Not applicable Yest for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% 0.01 0.1 Yest for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% 0.01 0.1 1 Yest for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% 0.01 0.1 1 10 Favours lower SFA Favours high Favours high 10 10 10	Subtotal (95% CI)		0		0		Not estimable	
Heterogeneity: Not applicable 'est for overall effect: Not applicable 'est for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Contracts	Fotal events:	0		0				
The formation of the f	Heterogeneity: Not applicable	e						
Sest for subgroup differences: Chi ² = 1.05, df = 3 (P = 0.79), I ² = 0% Controles	Test for overall effect: Not ar	oplicable						
Yest for subgroup differences: $Chi^2 = 1.05$, $df = 3$ (P = 0.79), $I^2 = 0\%$ 0.01 0.1 1 10 Favours lower SFAFavours highContractes	· · · · · · · · · · · · · · · ·							
Controtes	Fest for subgroup differences	s: Chi ² = 1.0	5, $df = 3$ (1	P = 0.79), I	$^{2} = 0\%$		E,	0.01 0.1 1 10
	Footnotes						Га	ravours lower SI'A Favours light

(1) Women with and without CVD at baseline



Analysis 1.27. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 27: CVD mortality, subgroup by main substitution

	lower	SFA	higher	SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
1.27.1 replaced by PUFA							
DART 1989	101	1018	100	1015	24.3%	1.01 [0.77 , 1.31]	
MRC 1968	27	199	25	194	15.3%	1.05 [0.63 , 1.75]	-
Oslo Diet-Heart 1966	38	206	52	206	20.0%	0.73 [0.50 , 1.06]	_
Rose corn oil 1965	5	28	1	26	1.7%	4.64 [0.58, 37.15]	
Sydney Diet-Heart 1978	37	221	25	237	16.3%	1.59 [0.99 , 2.55]	
Veterans Admin 1969	57	424	81	422	22.4%	0.70 [0.51, 0.96]	-
Subtotal (95% CI)		2096		2100	100.0%	0.97 [0.73 , 1.28]	
Total events:	265		284			- , -	▼
Heterogeneity: $Tau^2 = 0.06$;	$Chi^2 = 12.62$	df = 5 (P)	= 0.03; I ²	= 60%			
Test for overall effect: $Z = 0$	0.22 (P = 0.82)	2)	,,,				
1.27.2 replaced by MUFA							
Rose olive 1965	3	26	1	26	100.0%	3.00 [0.33 , 26.99]	
Subtotal (95% CI)		26		26	100.0%	3.00 [0.33 , 26.99]	
Total events:	3		1				
Heterogeneity: Not applicab	ole						
Test for overall effect: $Z = 0$	0.98 (P = 0.33)	3)					
1.27.3 replace by CHO							
Black 1994	0	66	2	67	4.1%	0.20 [0.01, 4.15]	←
Ley 2004	1	88	4	88	7.6%	0.25 [0.03 , 2.19]	
STARS 1992	1	27	3	28	7.4%	0.35 [0.04, 3.12]	
WHI 2006 (1)	213	19541	320	29294	81.0%	1.00 [0.84 , 1.19]	
Subtotal (95% CI)		19722		29477	100.0%	0.78 [0.42, 1.46]	
Total events:	215		329			- / -	
Heterogeneity: $Tau^2 = 0.12$;	$Chi^2 = 3.47$,	df = 3 (P =	= 0.32); I ² =	= 14%			
Test for overall effect: $Z = 0$	0.78 (P = 0.43)	3)					
1.27.4 replaced by protein							
Subtotal (95% CI)		0		0		Not estimable	
Total events:	0		0				
Heterogeneity: Not applicab	ole						
Test for overall effect: Not a	applicable						
1.27.5 replacement unclear	r						
Subtotal (95% CI)		0		0		Not estimable	
Total events:	0		0				
Heterogeneity: Not applicab	ole						
Test for overall effect: Not a	applicable						
Test for subgroup difference	es: Chi ² = 1.4	6, df = 2 (1)	P = 0.48), I	² = 0%		E.	0.01 0.1 1 10 100
Footnotes						F	avours lower SFA Favours night SFA

(1) Women with and without CVD at baseline



lower SFA higher SFA **Risk Ratio Risk Ratio** Total Study or Subgroup Events Total **Events** Weight M-H, Random, 95% CI M-H, Random, 95% CI 1.28.1 up to 24mo 2 0 Black 1994 66 67 0.4% 0.20 [0.01, 4.15] DART 1989 1015 101 1018 100 19.4% 1.01 [0.77, 1.31] Rose corn oil 1965 5 28 1 26 0.9% 4.64 [0.58, 37.15] Rose olive 1965 3 26 1 26 0.8%3.00 [0.33, 26.99] 1134 Subtotal (95% CI) 1138 21.5% 1.26 [0.54, 2.94] Total events: 109 104 Heterogeneity: Tau² = 0.26; Chi² = 4.07, df = 3 (P = 0.25); I² = 26% Test for overall effect: Z = 0.54 (P = 0.59) 1.28.2 >24 to 48mo **MRC 1968** 27 199 25 194 10.1% 1.05 [0.63, 1.75] **STARS 1992** 3 1 27 28 0.8%0.35 [0.04, 3.12] Veterans Admin 1969 57 424 81 422 17.1% 0.70 [0.51, 0.96] Subtotal (95% CI) 650 644 28.0% 0.79 [0.57, 1.08] 85 Total events: 109 Heterogeneity: Tau² = 0.01; Chi² = 2.33, df = 2 (P = 0.31); I² = 14% Test for overall effect: Z = 1.49 (P = 0.14)1.28.3 >48 mo Oslo Diet-Heart 1966 38 206 52 206 14.5% 0.73 [0.50, 1.06] Sydney Diet-Heart 1978 37 221 25 237 11.0% 1.59 [0.99, 2.55] WHI 2006 (1) 213 19541 320 29294 24.1% 1.00 [0.84, 1.19] Subtotal (95% CI) 19968 29737 49.6% 1.02 [0.73, 1.43] Total events: 288 397 Heterogeneity: Tau² = 0.06; Chi² = 6.40, df = 2 (P = 0.04); I² = 69% Test for overall effect: Z = 0.10 (P = 0.92) 1.28.4 unclear duration 88 88 0.8% Ley 2004 1 4 0.25 [0.03, 2.19] Subtotal (95% CI) 88 88 0.8% 0.25 [0.03, 2.19] Total events: 1 4 Heterogeneity: Not applicable Test for overall effect: Z = 1.25 (P = 0.21) Total (95% CI) 21844 31603 100.0% 0.95 [0.78, 1.16] Total events: 483 614 Heterogeneity: Tau² = 0.04; Chi² = 17.36, df = 10 (P = 0.07); I² = 42% 0.01 0.110 100 Test for overall effect: Z = 0.53 (P = 0.60) Favours lower SFA Favours higher SFA Test for subgroup differences: $Chi^2 = 3.15$, df = 3 (P = 0.37), $I^2 = 4.9\%$

Analysis 1.28. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 28: CVD mortality, subgroup by duration

Footnotes

(1) Women with and without CVD at baseline



Analysis 1.29. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 29: CVD mortality, subgroup by baseline SFA

	lower	SFA	higher	SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
1.29.1 up to 12%E SFA ba	aseline						
Subtotal (95% CI)		0		0		Not estimable	
Total events:	0		0				
Heterogeneity: Not applicab	ole						
Test for overall effect: Not a	applicable						
1.29.2 >12 to 15% E SFA b	oaseline						
Black 1994	0	66	2	67	0.4%	0.20 [0.01 , 4.15]	←
DART 1989	101	1018	100	1015	19.4%	1.01 [0.77 , 1.31]	· ·
Ley 2004	1	88	4	88	0.8%	0.25 [0.03 , 2.19]	
Sydney Diet-Heart 1978	37	221	25	237	11.0%	1.59 [0.99 , 2.55]	
WHI 2006 (1)	213	19541	320	29294	24.1%	1.00 [0.84, 1.19]	
Subtotal (95% CI)		20934		30701	55.8%	1.06 [0.84, 1.32]	▲
Total events:	352		451				T
Heterogeneity: $Tau^2 = 0.02$;	$Chi^2 = 6.12,$	df = 4 (P =	= 0.19); I ² =	= 35%			
Test for overall effect: $Z = 0$	0.48 (P = 0.63)	3)	,.				
1.29.3 >15 to 18% E SFA b	oaseline						
STARS 1992	1	27	3	28	0.8%	0.35 [0.04, 3.12]	
Subtotal (95% CI)		27		28	0.8%	0.35 [0.04, 3.12]	
Total events:	1		3			- / -	
Heterogeneity: Not applicab	ole						
Test for overall effect: $Z = 0$	0.95 (P = 0.34)	4)					
1.29.4 >18%E SFA baselir	ne						
Veterans Admin 1969	57	424	81	422	17.1%	0.70 [0.51, 0.96]	-
Subtotal (95% CI)		424		422	17.1%	0.70 [0.51 , 0.96]	
Total events:	57		81				•
Heterogeneity: Not applicat	ole						
Test for overall effect: $Z = 2$	2.25 (P = 0.02)	2)					
1.29.5 unclear							
MRC 1968	27	199	25	194	10.1%	1.05 [0.63 , 1.75]	_ _
Oslo Diet-Heart 1966	38	206	52	206	14.5%	0.73 [0.50 , 1.06]	
Rose corn oil 1965	5	28	1	26	0.9%	4.64 [0.58 , 37.15]	
Rose olive 1965	3	26	1	26	0.8%	3.00 [0.33 , 26.99]	
Subtotal (95% CI)		459		452	26.3%	1.00 [0.61 , 1.66]	•
Total events:	73		79				Ť
Heterogeneity: $Tau^2 = 0.10$;	Chi ² = 5.16,	df = 3 (P =	= 0.16); I ² =	= 42%			
Test for overall effect: $Z = 0$	0.01 (P = 0.99)))					
Total (95% CI)		21844		31603	100.0%	0.95 [0.78 , 1.16]	4
Total events:	483		614				1
Heterogeneity: $Tau^2 = 0.04$;	Chi ² = 17.36	df = 10 (P = 0.07; I	$^{2} = 42\%$			0.01 0.1 1 10 100
Test for overall effect: $Z = 0$	0.53 (P = 0.60)))				Fa	avours lower SFA Favours higher SF
Test for subgroup difference	es: Chi ² = 5.2	9, $df = 3$ (1	P = 0.15), I	² = 43.3%			

Footnotes

(1) Women with and without CVD at baseline

Reduction in saturated fat intake for cardiovascular disease (Review)

Copyright @ 2020 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.



Analysis 1.30. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 30: CVD mortality, subgroup by SFA change

	lower	SFA	higher	SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight M-H, Random, 95% CI		M-H, Random, 95% CI
1.30.1 up to 4% E differend	ce						
DART 1989	101	1018	100	1015	19.4%	1.01 [0.77 , 1.31]	-
Ley 2004	1	88	4	88	0.8%	0.25 [0.03 , 2.19]	
Sydney Diet-Heart 1978	37	221	25	237	11.0%	1.59 [0.99 , 2.55]	
WHI 2006 (1)	213	19541	320	29294	24.1%	1.00 [0.84 , 1.19]	
Subtotal (95% CI)		20868		30634	55.4%	1.07 [0.85 , 1.33]	•
Total events:	352		449				
Heterogeneity: $Tau^2 = 0.02$;	$Chi^2 = 5.00,$	df = 3 (P = 3)	= 0.17); I ² =	= 40%			
Test for overall effect: $Z = 0$	0.56 (P = 0.57)	7)					
1.30.2 >4 to 8% E difference	e						
Black 1994	0	66	2	67	0.4%	0.20 [0.01 , 4.15]	← → ↓
STARS 1992	1	27	3	28	0.8%	0.35 [0.04 , 3.12]	
Subtotal (95% CI)		93		95	1.2%	0.29 [0.05 , 1.70]	
Total events:	1		5				
Heterogeneity: $Tau^2 = 0.00$;	$Chi^2 = 0.08,$	df = 1 (P = 1)	= 0.78); I ² =	= 0%			
Test for overall effect: $Z = 1$.37 (P = 0.17)	7)					
1.30.3 >8%E difference							
Veterans Admin 1969	57	424	81	422	17.1%	0.70 [0.51, 0.96]	-
Subtotal (95% CI)		424		422	17.1%	0.70 [0.51 , 0.96]	
Total events:	57		81				•
Heterogeneity: Not applicab	ole						
Test for overall effect: $Z = 2$	P.25 (P = 0.02)	2)					
1.30.4 unclear							
MRC 1968	27	199	25	194	10.1%	1.05 [0.63 , 1.75]	_ _
Oslo Diet-Heart 1966	38	206	52	206	14.5%	0.73 [0.50 , 1.06]	
Rose corn oil 1965	5	28	1	26	0.9%	4.64 [0.58, 37.15]	
Rose olive 1965	3	26	1	26	0.8%	3.00 [0.33 , 26.99]	
Subtotal (95% CI)		459		452	26.3%	1.00 [0.61 , 1.66]	•
Total events:	73		79				T
Heterogeneity: $Tau^2 = 0.10$;	Chi ² = 5.16,	df = 3 (P =	= 0.16); I ² =	= 42%			
Test for overall effect: $Z = 0$	0.01 (P = 0.99)))					
Total (95% CI)		21844		31603	100.0%	0.95 [0.78 , 1.16]	•
Total events:	483		614] .
Heterogeneity: $Tau^2 = 0.04;$	Chi ² = 17.36	df = 10 (P = 0.07; I	$1^2 = 42\%$			0.01 0.1 1 10 100
Test for overall effect: $Z = 0$	0.53 (P = 0.60)))				Fa	avours lower SFA Favours higher SFA
Test for subgroup difference	es: $Chi^2 = 6.3$	9. $df = 3.0$	P = 0.09) I	$^{2} = 53.1\%$			

Footnotes

(1) Women with and without CVD at baseline

	lower	SFA	higher	SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
1.31.1 Men							
Black 1994	0	66	2	67	0.4%	0.20 [0.01, 4.15]	←
DART 1989	101	1018	100	1015	19.4%	1.01 [0.77, 1.31]	`
MRC 1968	27	199	25	194	10.1%	1.05 [0.63 , 1.75]	
Oslo Diet-Heart 1966	38	206	52	206	14.5%	0.73 [0.50, 1.06]	_
Rose corn oil 1965	5	28	1	26	0.9%	4.64 [0.58, 37.15]	
Rose olive 1965	3	26	1	26	0.8%	3.00 [0.33, 26.99]	
STARS 1992	1	27	3	28	0.8%	0.35 [0.04, 3.12]	
Sydney Diet-Heart 1978	37	221	25	237	11.0%	1.59 [0.99, 2.55]	
Veterans Admin 1969	57	424	81	422	17.1%	0.70 [0.51, 0.96]	-
Subtotal (95% CI)		2215		2221	75.1%	0.96 [0.73, 1.25]	A
Total events:	269		290				T
Heterogeneity: $Tau^2 = 0.06$; Chi ² = 15.45	df = 8 (F)	P = 0.05; I ²	= 48%			
Test for overall effect: $Z = 0$	0.32 (P = 0.75)	5)					
1.31.2 Women							
WHI 2006 (1)	213	19541	320	29294	24.1%	1.00 [0.84 , 1.19]	
Subtotal (95% CI)		19541		29294	24.1%	1.00 [0.84 , 1.19]	4
Total events:	213		320				Ĭ
Heterogeneity: Not applical	ole						
Test for overall effect: $Z = 0$	0.02 (P = 0.98)	3)					
1.31.3 Mixed, men and wo	omen						
Ley 2004	1	88	4	88	0.8%	0.25 [0.03 , 2.19]	
Subtotal (95% CI)		88		88	0.8%	0.25 [0.03 , 2.19]	
Total events:	1		4				
Heterogeneity: Not applical	ole						
Test for overall effect: Z =	1.25 (P = 0.21)	l)					
Total (95% CI)		21844		31603	100.0%	0.95 [0.78 , 1.16]	•
Total events:	483		614				Ţ
Heterogeneity: $Tau^2 = 0.04$; Chi ² = 17.36	df = 10 (P = 0.07; I	^{[2} = 42%			
Test for overall effect: $Z = 0$	0.53 (P = 0.60)))				Fa	avours lower SFA Favours higher SFA
Test for subgroup differenc	es: Chi ² = 1.5	9, df = 2 (P = 0.45), I	$^{2} = 0\%$			C C

Analysis 1.31. Comparison 1: SFA reduction vs usual diet primary outcomes, Outcome 31: CVD mortality, subgroup by sex

Footnotes

(1) Women with and without CVD at baseline

Analysis 1.32. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 32: CVD mortality, subgroup by CVD risk

	lower	SFA	higher	SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
1.32.1 Low CVD risk							
Black 1994	0	66	2	67	0.3%	0.20 [0.01 , 4.15]	• • • • • • • • • • • • • • • • • • •
Veterans Admin 1969	57	424	81	422	15.2%	0.70 [0.51, 0.96]	•
WHI 2006 (1)	170	18633	258	27925	21.4%	0.99 [0.81 , 1.20]	.
Subtotal (95% CI)		19123		28414	36.9%	0.84 [0.60 , 1.16]	
Total events:	227		341				•
Heterogeneity: $Tau^2 = 0.04$; Chi ² = 4.33,	df = 2 (P =	= 0.11); I ² =	= 54%			
Test for overall effect: $Z =$	1.06 (P = 0.29)))					
1.32.2 Moderate CVD risk	ζ.						
Ley 2004	1	88	4	88	0.6%	0.25 [0.03 , 2.19]	
Subtotal (95% CI)		88		88	0.6%	0.25 [0.03 , 2.19]	
Total events:	1		4				
Heterogeneity: Not applical	ble						
Test for overall effect: $Z =$	1.25 (P = 0.21)	l)					
1.32.3 Existing CVD disea	ise						
DART 1989	101	1018	100	1015	17.6%	1.01 [0.77 , 1.31]	+
MRC 1968	27	199	25	194	8.5%	1.05 [0.63 , 1.75]	_ _
Oslo Diet-Heart 1966	38	206	52	206	12.7%	0.73 [0.50 , 1.06]	-
Rose corn oil 1965	5	28	1	26	0.7%	4.64 [0.58, 37.15]	
Rose olive 1965	3	26	1	26	0.6%	3.00 [0.33 , 26.99]	
STARS 1992	1	27	3	28	0.6%	0.35 [0.04 , 3.12]	.
Sydney Diet-Heart 1978	37	221	25	237	9.4%	1.59 [0.99 , 2.55]	
WHI 2006 (2)	43	908	62	1369	12.3%	1.05 [0.72 , 1.53]	_
Subtotal (95% CI)		2633		3101	62.4%	1.04 [0.83 , 1.31]	•
Total events:	255		269				ľ
Heterogeneity: $Tau^2 = 0.03$; Chi ² = 10.41	, df = 7 (F)	P = 0.17; I ²	= 33%			
Test for overall effect: $Z = 0$	0.37 (P = 0.71)	l)					
Total (95% CI)		21844		31603	100.0%	0.96 [0.80 , 1.14]	
Total events:	483		614				
Heterogeneity: $Tau^2 = 0.03$; Chi ² = 17.45	, df = 11 (P = 0.10; I	2 = 37%		0.0	1 0.1 1 10 100
Test for overall effect: $Z = 0$	0.50 (P = 0.61)	l)				Favo	ours lower SFA Favours higher SI
Test for subgroup differenc	es: Chi ² = 2.6	6, df = 2 (P = 0.26), I	² = 24.9%			

Footnotes

(1) Women without CVD at baseline

(2) Women with CVD at baseline



Analysis 1.33. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 33: CVD mortality, subgroup by TC reduction

	lower	lower SFA		higher SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
1.33.1 serum chol reduced	by at least 0	.2mmol/L					
DART 1989	101	1018	100	1015	24.1%	1.01 [0.77 , 1.31]	+
MRC 1968	27	199	25	194	15.0%	1.05 [0.63 , 1.75]	_
Oslo Diet-Heart 1966	38	206	52	206	19.7%	0.73 [0.50, 1.06]	-
Rose corn oil 1965	5	28	1	26	1.6%	4.64 [0.58, 37.15]	
STARS 1992	1	27	3	28	1.5%	0.35 [0.04 , 3.12]	
Sydney Diet-Heart 1978	37	221	25	237	16.0%	1.59 [0.99 , 2.55]	
Veterans Admin 1969	57	424	81	422	22.1%	0.70 [0.51, 0.96]	-
Subtotal (95% CI)		2123		2128	100.0%	0.95 [0.73 , 1.25]	▲
Total events:	266		287				Ť
Heterogeneity: $Tau^2 = 0.06$;	; Chi ² = 13.37	df = 6 (P	= 0.04; I ²	= 55%			
Test for overall effect: $Z = 0$	0.34 (P = 0.73)	3)					
1.33.2 serum chol reduced	l by <0.2mm	ol/L					
Ley 2004	1	88	4	88	9.9%	0.25 [0.03 , 2.19]	
Rose olive 1965	3	26	1	26	9.7%	3.00 [0.33 , 26.99]	
WHI 2006 (1)	213	19541	320	29294	80.3%	1.00 [0.84 , 1.19]	•
Subtotal (95% CI)		19655		29408	100.0%	0.97 [0.47 , 2.01]	
Total events:	217		325				Ť
Heterogeneity: $Tau^2 = 0.16$;	, Chi ² = 2.53,	df = 2 (P =	= 0.28); I ² =	= 21%			
Test for overall effect: $Z = 0$	0.09 (P = 0.93)	3)					
1.33.3 serum chol reduction	on unclear						
Black 1994	0	66	2	67	100.0%	0.20 [0.01 , 4.15]	←─── ─ ───
Subtotal (95% CI)		66		67	100.0%	0.20 [0.01 , 4.15]	
Total events:	0		2				
Heterogeneity: Not applicat	ole						
Test for overall effect: $Z = 2$	1.04 (P = 0.30)))					
Test for subgroup difference	es: Chi ² = 1.0	1, df = 2 (P = 0.60), I	² = 0%		F	0.01 0.1 1 10 100 avours lower SFA Favours higher SFA
Footnotes							

(1) Women with and without CVD at baseline

Analysis 1.34. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 34: CVD mortality, subgroup decade of publication

	lower	SFA	higher	SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
1.34.1 1960s							
MRC 1968	27	199	25	194	9.5%	1.05 [0.63, 1.75]	
Oslo Diet-Heart 1966	38	206	52	206	14.2%	0.73 [0.50, 1.06]	_
Rose corn oil 1965	5	28	0	13	0.4%	5.31 [0.32, 89.44]	
Rose olive 1965	3	26	1	13	0.7%	1.50 [0.17, 13.05]	
Veterans Admin 1969	57	424	81	422	17.2%	0.70 [0.51, 0.96]	_
Subtotal (95% CI)		883		848	42.1%	0.78 [0.63 , 0.97]	
Total events:	130		159				
Heterogeneity: $Tau^2 = 0.00$	$Chi^2 = 4.07.$	df = 4 (P =	= 0.40): I ² =	- 2%			
Test for overall effect: $Z = 1$	2.22 (P = 0.03)	3)	,,-				
1.34.2 1970s							
Sydney Diet-Heart 1978	37	221	25	237	10.5%	1.59 [0.99 . 2.55]	
Subtotal (95% CI)	- /	221		237	10.5%	1.59 [0.99 , 2.55]	
Total events:	37		25				
Heterogeneity: Not applical	ble		20				
Test for overall effect: $Z =$	1.91 (P = 0.06)	ຄ					
	1.91 (I = 0.00	,					
1.34.3 1980s							
DART 1989	101	1018	100	1015	19.9%	1.01 [0.77 , 1.31]	+
Subtotal (95% CI)		1018		1015	19.9%	1.01 [0.77 , 1.31]	
Total events:	101		100				Ť
Heterogeneity: Not applical	ble						
Test for overall effect: $Z =$	0.05 (P = 0.96)	5)					
1.34.4 1990s							
Black 1994	0	66	2	67	0.4%	0.20 [0.01, 4.15]	←
STARS 1992	1	27	3	28	0.7%	0.35 [0.04, 3.12]	`
Subtotal (95% CI)		93		95	1.1%	0.29 [0.05, 1.70]	
Total events:	1		5				
Heterogeneity: $Tau^2 = 0.00$; $Chi^2 = 0.08$,	df = 1 (P = 1)	= 0.78); I ² =	:0%			
Test for overall effect: $Z =$	1.37 (P = 0.17)	7)					
1.34.5 2000s							
Ley 2004	1	88	4	88	0.7%	0.25 [0.03 , 2.19]	
WHI 2006 (1)	213	19541	320	29294	25.7%	1.00 [0.84 , 1.19]	↓
Subtotal (95% CI)		19629		29382	26.5%	0.78 [0.27 , 2.21]	
Total events:	214		324				
Heterogeneity: $Tau^2 = 0.34$; Chi ² = 1.55,	df = 1 (P =	= 0.21); I ² =	: 36%			
Test for overall effect: $Z =$	0.48 (P = 0.63)	3)					
Total (95% CI)		21844		31577	100.0%	0.94 [0.78 , 1.13]	
Total events:	483		613				٢
Heterogeneity: $Tau^2 = 0.03$; Chi ² = 15.67	df = 10 (P = 0.11; I	² = 36%			0.01 0.1 1 10 100
Test for overall effect: $Z =$	0.68 (P = 0.49)))				F	avours lower SFA Favours higher SF
Test for subgroup differenc	es: Chi ² = 9.4	8, df = 4 (P = 0.05), I	² = 57.8%			-

Footnotes

(1) Women with and without CVD at baseline



Analysis 1.35. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 35: COMBINED CARDIOVASCULAR EVENTS

	lower	lower SFA		SFA	Risk Ratio		Risk Rat		tio	
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-l	H, Random	, 95% CI	
Black 1994	0	66	2	67	0.3%	0.20 [0.01 , 4.15]		_		_
DART 1989	136	1018	147	1015	14.9%	0.92 [0.74, 1.15]				
Houtsmuller 1979	8	51	30	51	4.9%	0.27 [0.14, 0.52]				
Ley 2004	11	88	16	88	4.5%	0.69 [0.34 , 1.40]				
MRC 1968	62	199	74	194	13.1%	0.82 [0.62, 1.07]		-		
Moy 2001	5	117	3	118	1.4%	1.68 [0.41 , 6.87]				
Oslo Diet-Heart 1966	64	206	90	206	13.6%	0.71 [0.55, 0.92]		-		
Rose corn oil 1965	15	28	6	13	4.8%	1.16 [0.59 , 2.29]				
Rose olive 1965	11	26	5	13	3.6%	1.10 [0.48 , 2.50]			-	
STARS 1992	8	27	20	28	5.4%	0.41 [0.22, 0.78]				
Veterans Admin 1969	97	424	122	422	14.5%	0.79 [0.63 , 1.00]		-		
WHI 2006 (1)	1399	19541	2145	29294	19.0%	0.98 [0.92, 1.04]		•		
Total (95% CI)		21791		31509	100.0%	0.79 [0.66 , 0.93]		•		
Total events:	1816		2660					•		
Heterogeneity: $Tau^2 = 0.0$	04; Chi ² = 31.	72, $df = 1$	1 (P = 0.00)	$(08); I^2 = 63$	5%		0.005 0).1 1	10 200	
Test for overall effect: Z	= 2.72 (P = 0)	.006)				F	avours lower	SFA	Favours higher S	FA

Test for subgroup differences: Not applicable

Footnotes

(1) Total CVD during study period, Prentice 2017

Analysis 1.36. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 36: CVD events, SA low summary risk of bias

	lower	lower SFA		higher SFA		Risk Ratio	Risk Ratio	
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI	
Ley 2004	11	88	16	88	5.9%	0.69 [0.34 , 1.40]		
Veterans Admin 1969	97	424	122	422	32.2%	0.79 [0.63 , 1.00]		
WHI 2006 (1)	1399	19541	2145	29294	61.9%	0.98 [0.92 , 1.04]	•	
Total (95% CI)		20053		29804	100.0%	0.89 [0.75 , 1.07]		
Total events:	1507		2283					
Heterogeneity: $Tau^2 = 0.0$	01; $Chi^2 = 3.8$	8, $df = 2$ (P = 0.14; I	0.00)5 0.1 1 10	200		
Test for overall effect: $Z = 1.21$ (P = 0.23)						Favo	urs lower SFA Favours l	nigher SFA

Test for subgroup differences: Not applicable

Footnotes

(1) Total CVD during study period, Prentice 2017



Analysis 1.37. Comparison 1: SFA reduction vs usual diet primary outcomes, Outcome 37: CVD events, SA aim to reduce SFA

	lower	lower SFA		higher SFA		Risk Ratio	Risk Ratio	
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95	% CI
DART 1989	136	1018	147	1015	15.6%	0.92 [0.74 , 1.15]	-	
Houtsmuller 1979	8	51	30	51	5.2%	0.27 [0.14, 0.52]		
MRC 1968	62	199	74	194	13.7%	0.82 [0.62, 1.07]	-	
Moy 2001	5	117	3	118	1.5%	1.68 [0.41 , 6.87]		
Oslo Diet-Heart 1966	64	206	90	206	14.3%	0.71 [0.55, 0.92]	+	
Rose corn oil 1965	15	28	6	13	5.2%	1.16 [0.59 , 2.29]	_ _	
Rose olive 1965	11	26	5	13	3.9%	1.10 [0.48 , 2.50]	_	
STARS 1992	8	27	20	28	5.8%	0.41 [0.22, 0.78]		
Veterans Admin 1969	97	424	122	422	15.1%	0.79 [0.63 , 1.00]	-	
WHI 2006 (1)	1399	19541	2145	29294	19.7%	0.98 [0.92 , 1.04]	+	
Total (95% CI)		21637		31354	100.0%	0.79 [0.66 , 0.95]	•	
Total events:	1805		2642				*	
Heterogeneity: $Tau^2 = 0.0$	04; Chi ² = 30	.07, $df = 9$	(P = 0.000)	4); $I^2 = 70^6$	%	0.0	05 0.1 1	10 200
Test for overall effect: $Z = 2.51$ (P = 0.01)					Favo	ours lower SFA Fav	ours higher SFA	
Test for subgroup differe	nces: Not app	olicable						

Footnotes

(1) Total CVD during study period, Prentice 2017

Analysis 1.38. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 38: CVD events, SA statistically significant SFA reduction

	lower	SFA	higher	SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
Black 1994	0	66	2	67	0.3%	0.20 [0.01 , 4.15]	
DART 1989	136	1018	147	1015	24.9%	0.92 [0.74 , 1.15]	-
Ley 2004	11	88	16	88	5.4%	0.69 [0.34 , 1.40]	
Moy 2001	5	117	3	118	1.5%	1.68 [0.41 , 6.87]	_ _
STARS 1992	8	27	20	28	6.7%	0.41 [0.22, 0.78]	
Veterans Admin 1969	97	424	122	422	23.7%	0.79 [0.63 , 1.00]	-
WHI 2006 (1)	1399	19541	2145	29294	37.5%	0.98 [0.92, 1.04]	•
Total (95% CI)		21281		31032	100.0%	0.85 [0.71 , 1.02]	
Total events:	1656		2455				•
Heterogeneity: $Tau^2 = 0.02$	2; Chi ² = 12.	43, df = 6	(P = 0.05);	$I^2 = 52\%$		(0.005 0.1 1 10 200
Test for overall effect: Z =	1.77 (P = 0	.08)				Fa	wours lower SFA Favours higher SF
Test for subgroup difference	ces: Not app	olicable					

Footnotes

(1) Total CVD during study period, Prentice 2017



Analysis 1.39. Comparison 1: SFA reduction vs usual diet primary outcomes, Outcome 39: CVD events, SA TC reduction

	lower	SFA	higher	SFA		Risk Ratio	Risk Ratio	
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% (CI
DART 1989	136	1018	147	1015	16.1%	0.92 [0.74 , 1.15]	-	
Houtsmuller 1979	8	51	30	51	5.6%	0.27 [0.14 , 0.52]	_	
MRC 1968	62	199	74	194	14.3%	0.82 [0.62 , 1.07]	-	
Moy 2001	5	117	3	118	1.6%	1.68 [0.41 , 6.87]	_	
Oslo Diet-Heart 1966	64	206	90	206	14.9%	0.71 [0.55, 0.92]	+	
Rose corn oil 1965	15	28	6	13	5.5%	1.16 [0.59 , 2.29]		
STARS 1992	8	27	20	28	6.2%	0.41 [0.22, 0.78]		
Veterans Admin 1969	97	424	122	422	15.7%	0.79 [0.63 , 1.00]	-	
WHI 2006 (1)	1399	19541	2145	29294	20.1%	0.98 [0.92 , 1.04]	+	
Total (95% CI)		21611		31341	100.0%	0.78 [0.65 , 0.94]		
Total events:	1794		2637				•	
Heterogeneity: $Tau^2 = 0.0$	04; Chi ² = 29.	91, df = 8	(P = 0.000)	2); $I^2 = 73^{\circ}$	%	0.00	05 0.1 1 10	200
Test for overall effect: $Z = 2.58$ (P = 0.010)						Favor	irs lower SFA Favour	s higher SFA
Test for subgroup differen	nces: Not app	licable						

Footnotes

(1) Total CVD during study period, Prentice 2017

Analysis 1.40. Comparison 1: SFA reduction vs usual diet primary outcomes, Outcome 40: CVD events, SA excluding WHI

	lower	SFA	higher SFA		Risk Ratio		Risk Ratio	
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Randon	n, 95% CI
Black 1994	0	66	2	67	0.4%	0.20 [0.01 , 4.15]		
DART 1989	136	1018	147	1015	18.1%	0.92 [0.74 , 1.15]	-	
Houtsmuller 1979	8	51	30	51	6.1%	0.27 [0.14, 0.52]		
Ley 2004	11	88	16	88	5.7%	0.69 [0.34 , 1.40]		
MRC 1968	62	199	74	194	16.0%	0.82 [0.62, 1.07]	-	
Moy 2001	5	117	3	118	1.8%	1.68 [0.41 , 6.87]		
Oslo Diet-Heart 1966	64	206	90	206	16.7%	0.71 [0.55, 0.92]	-	
Rose corn oil 1965	15	28	6	13	6.1%	1.16 [0.59 , 2.29]	_	_
Rose olive 1965	11	26	5	13	4.6%	1.10 [0.48 , 2.50]		_
STARS 1992	8	27	20	28	6.9%	0.41 [0.22, 0.78]		
Veterans Admin 1969	97	424	122	422	17.6%	0.79 [0.63 , 1.00]	-	
Total (95% CI)		2250		2215	100.0%	0.75 [0.61 , 0.91]	•	
Total events:	417		515				•	
Heterogeneity: $Tau^2 = 0.0$	04; $Chi^2 = 20.$	51, df = 10	0 (P = 0.02)); $I^2 = 51\%$,	0.	.005 0.1 1	10 200
Test for overall effect: Z	= 2.93 (P = 0)	.003)				Fav	ours lower SFA	Favours higher SFA

Test for subgroup differences: Not applicable



Analysis 1.41. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 41: CVD events, SA Mantel-Haenszel fixed-effect

	lower	SFA	higher	SFA		Risk Ratio	Risk Ratio		
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Fixed, 95% CI	M-H, Fixed, 95% Cl	[
Black 1994	0	66	2	67	0.1%	0.20 [0.01 , 4.15]			
DART 1989	136	1018	147	1015	6.6%	0.92 [0.74 , 1.15]	-		
Houtsmuller 1979	8	51	30	51	1.3%	0.27 [0.14, 0.52]			
Ley 2004	11	88	16	88	0.7%	0.69 [0.34 , 1.40]			
MRC 1968	62	199	74	194	3.4%	0.82 [0.62 , 1.07]			
Moy 2001	5	117	3	118	0.1%	1.68 [0.41 , 6.87]			
Oslo Diet-Heart 1966	64	206	90	206	4.0%	0.71 [0.55, 0.92]	-		
Rose corn oil 1965	15	28	6	13	0.4%	1.16 [0.59 , 2.29]	_ _		
Rose olive 1965	11	26	5	13	0.3%	1.10 [0.48 , 2.50]	_ _		
STARS 1992	8	27	20	28	0.9%	0.41 [0.22, 0.78]			
Veterans Admin 1969	97	424	122	422	5.5%	0.79 [0.63 , 1.00]	-		
WHI 2006 (1)	1399	19541	2145	29294	76.7%	0.98 [0.92 , 1.04]	•		
Total (95% CI)		21791		31509	100.0%	0.93 [0.88 , 0.99]			
Total events:	1816		2660						
Heterogeneity: $Chi^2 = 31$.72, df = 11 (P = 0.0008	$(I); I^2 = 65\%$	2		(005 0.1 1 10	200	
Test for overall effect: Z	= 2.44 (P = 0)	.01)				Fa	vours lower SFA Favour	s higher SFA	
Test for subgroup differe	ences: Not app	olicable							

Footnotes

(1) Total CVD during study period, Prentice 2017

	lower	lower SFA		higher SFA		Peto Odds Ratio	Peto Odd	s Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	Peto, Fixed, 95% CI	Peto, Fixed	95% CI
Black 1994	0	66	2	67	0.1%	0.14 [0.01 , 2.19]	_
DART 1989	136	1018	147	1015	6.4%	0.91 [0.71 , 1.17] 🚽	
Houtsmuller 1979	8	51	30	51	0.6%	0.16 [0.07 , 0.36]	
Ley 2004	11	88	16	88	0.6%	0.65 [0.29, 1.47]]	
MRC 1968	62	199	74	194	2.3%	0.73 [0.49 , 1.11]] _	
Moy 2001	5	117	3	118	0.2%	1.69 [0.41 , 6.90]	
Oslo Diet-Heart 1966	64	206	90	206	2.5%	0.58 [0.39, 0.87]] _	
Rose corn oil 1965	15	28	6	13	0.2%	1.34 [0.36 , 4.90]	
Rose olive 1965	11	26	5	13	0.2%	1.17 [0.31 , 4.44]	
STARS 1992	8	27	20	28	0.4%	0.19 [0.07 , 0.55]	
Veterans Admin 1969	97	424	122	422	4.2%	0.73 [0.54, 0.99]] _	
WHI 2006 (1)	1399	19541	2145	29294	82.3%	0.98 [0.91 , 1.05]	
Total (95% CI)		21791		31509	100.0%	0.92 [0.87 , 0.98]	
Total events:	1816		2660				1	
Heterogeneity: Chi ² = 41	.53, df = 11 (1	P < 0.0001); I ² = 74%				0.005 0.1 1	10 200
Test for overall effect: Z	= 2.44 (P = 0)	.01)					Favours lower SFA	Favours higher SFA

Analysis 1.42. Comparison 1: SFA reduction vs usual diet primary outcomes, Outcome 42: CVD events, SA Peto fixed-effect

Test for subgroup differences: Not applicable

Footnotes

(1) Total CVD during study period, Prentice 2017



Analysis 1.43. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 43: CVD events, SA excluding trials with additional interventions

	lower	SFA	higher	SFA		Risk Ratio	Risk Ratio	
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI	
Black 1994	0	66	2	67	0.5%	0.20 [0.01 , 4.15]		
DART 1989	136	1018	147	1015	23.8%	0.92 [0.74 , 1.15]	-	
Houtsmuller 1979	8	51	30	51	8.0%	0.27 [0.14, 0.52]		
Ley 2004	11	88	16	88	7.5%	0.69 [0.34 , 1.40]		
MRC 1968	62	199	74	194	21.0%	0.82 [0.62, 1.07]	-	
Moy 2001	5	117	3	118	2.3%	1.68 [0.41 , 6.87]	_ _	
Rose corn oil 1965	15	28	6	13	7.9%	1.16 [0.59 , 2.29]		
Rose olive 1965	11	26	5	13	5.9%	1.10 [0.48 , 2.50]	_ _	
Veterans Admin 1969	97	424	122	422	23.1%	0.79 [0.63 , 1.00]	-	
Total (95% CI)		2017		1981	100.0%	0.80 [0.64 , 0.99]		
Total events:	345		405				•	
Heterogeneity: $Tau^2 = 0.0$	$04; Chi^2 = 15.$.39, df = 8	(P = 0.05);	$I^2 = 48\%$		0.0	05 0.1 1 10	200
Test for overall effect: Z	= 2.01 (P = 0)	.04)				Favo	urs lower SFA Favours hig	gher SFA
Test for subgroup different	nces: Not app	olicable						

Analysis 1.44. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 44: CVD events, subgroup by any substitution

	lower	SFA	higher	SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
1.44.1 replaced by PUF.	A						
DART 1989	136	1018	147	1015	19.5%	0.92 [0.74 , 1.15]	-
Houtsmuller 1979	8	51	30	51	7.7%	0.27 [0.14, 0.52]	
MRC 1968	62	199	74	194	17.6%	0.82 [0.62, 1.07]	_
Oslo Diet-Heart 1966	64	206	90	206	18.2%	0.71 [0.55, 0.92]	-
Rose corn oil 1965	15	28	11	26	9.6%	1.27 [0.72, 2.23]	
STARS 1992	8	27	20	28	8.5%	0.41 [0.22, 0.78]	
Veterans Admin 1969	97	424	122	422	19.0%	0.79 [0.63 , 1.00]	-
Subtotal (95% CI)		1953		1942	100.0%	0.73 [0.58, 0.92]	
Total events:	390		494				•
Heterogeneity: $Tau^2 = 0.0$	06; Chi ² = 19.	.30, $df = 6$	(P = 0.004)); $I^2 = 69\%$, 0		
Test for overall effect: Z	= 2.68 (P = 0)	.007)					
1.44.2 replaced by MUF	FA						
Rose olive 1965	11	26	11	26	100.0%	1.00 [0.53 , 1.89]	_
Subtotal (95% CI)	-	26	-	26	100.0%	1.00 [0.53 , 1.89]	
Total events:	11		11				
Heterogeneity: Not appli	cable		-				
Test for overall effect: Z	= 0.00 (P = 1)	.00)					
1.44.3 replace by CHO							
Black 1994	0	66	2	67	0.6%	0.20 [0.01 . 4.15]	
DART 1989	136	1018	147	1015	33.8%	0.92 [0.74 , 1.15]	
Lev 2004	11	88	16	88	8.6%	0.69 [0.34 , 1.40]	I
STARS 1992	8	27	20	28	10.5%	0.41 [0.22, 0.78]	
WHI 2006 (1)	1399	19541	2145	29294	46.5%	0.98 [0.92, 1.04]	
Subtotal (95% CI)	1077	20740	2110	30492	100.0%	0.84 [0.67 , 1.06]	7
Total events:	1554		2330	00.02	10000 /0		
Heterogeneity: $Tau^2 = 0.0$	03: $Chi^2 = 9.2$	20. $df = 4$ (P = 0.06;	$[^2 = 57\%]$			
Test for overall effect: Z	= 1.46 (P = 0)	.14)					
1.44.4 replaced by prote	ein						
Black 1994	0	66	2	67	0.0%	0.20 [0.01, 4.15]	
DART 1989	136	1018	147	1015	8.2%	0.92 [0.74, 1.15]	-
Ley 2004	11	88	16	88	0.8%	0.69 [0.34, 1.40]	
WHI 2006 (1)	1399	19541	2145	29294	91.0%	0.98 [0.92, 1.04]	
Subtotal (95% CI)		20713		30464	100.0%	0.97 [0.91 , 1.03]	.
Total events:	1546		2310				
Heterogeneity: $Tau^2 = 0.0$	00; $Chi^2 = 2.2$	20, df = 3 (P = 0.53;	$l^2 = 0\%$			
Test for overall effect: Z	= 0.97 (P = 0)	.33)					
1.44.5 replacement uncl	ear						
Moy 2001	5	117	3	118	100.0%	1.68 [0.41, 6.87]	
Subtotal (95% CI)		117		118	100.0%	1.68 [0.41, 6.87]	
Total events:	5		3			. / 1	
Heterogeneity: Not appli	cable						
Test for overall effect: Z	= 0.72 (P = 0)	.47)					
Test for subgroup differe	nces: Chi ² = 7	7.10, df =	4 (P = 0.13), I ² = 43.7	%		05 0.1 1 10 200
- 1						Favo	burs lower SFA Favours higher S
Footnotes							

(1) Total CVD during study period, Prentice 2017

Analysis 1.45. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 45: CVD events, subgroup by main substitution

	lower	SFA	higher	SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
1.45.1 replaced by PUFA							
DART 1989	136	1018	147	1015	21.8%	0.92 [0.74, 1.15]	
Houtsmuller 1979	8	51	30	51	7.7%	0.27 [0.14, 0.52]]
MRC 1968	62	199	74	194	19.4%	0.82 [0.62, 1.07]	
Oslo Diet-Heart 1966	64	206	90	206	20.1%	0.71 [0.55, 0.92]	_
Rose corn oil 1965	15	28	11	26	9.8%	1.27 [0.72 , 2.23]	
Veterans Admin 1969	97	424	122	422	21.2%	0.79 [0.63 1.00]	_
Subtotal (95% CI)		1926		1914	100.0%	0.78 [0.62 , 0.97]	
Total events:	382	1720	474	1/11	100.0 /0		
Heterogeneity: $Tau^2 = 0.05$.	$Chi^2 = 15$	17 df = 5	(P = 0.010)). $I^2 = 67\%$	1		
Test for overall effect: $Z = 2$	2.25 (P = 0)	.02)	(1 = 0.010),1 = 07 %	, ,		
1.45.2 replaced by MUFA							
Rose olive 1965	11	26	11	26	100.0%	1.00 [0.53 , 1.89]	-
Subtotal (95% CI)		26		26	100.0%	1.00 [0.53 , 1.89]	
Total events:	11		11				Ť
Heterogeneity: Not applicab	ole						
Test for overall effect: $Z = 0$	0.00 (P = 1)	.00)					
1 45 2 monto as by CUO							
1.45.5 replace by CHO	0		2	(7	2.00	0.20 [0.01 4.15]	
Васк 1994 1. 2004	0	00	2	0/	5.0%	0.20 [0.01, 4.13]	
Ley 2004	11	88	16	88	25.2%	0.69 [0.34 , 1.40]	
STAKS 1992	8	27	20	28	27.8%	0.41 [0.22, 0.78]	
WHI 2006 (1)	1399	19541	2145	29294	44.1%	0.98 [0.92 , 1.04]	
Subtotal (95% CI)		19722		29477	100.0%	0.67 [0.39 , 1.16]	\bullet
Total events:	1418		2183				
Heterogeneity: $Tau^2 = 0.17$; Test for overall effect: $Z = 1$	$Chi^2 = 9.0$.44 (P = 0)4, df = 3 ().15)	P = 0.03;	$l^2 = 67\%$			
		,					
1.45.4 replaced by protein							
Subtotal (95% CI)		0		0		Not estimable	
Total events:	0		0				
Heterogeneity: Not applicab	ole						
Test for overall effect: Not a	applicable						
1.45.5 replacement unclear	r						
Moy 2001	5	117	3	118	100.0%	1.68 [0.41 , 6.87]	
Subtotal (95% CI)		117		118	100.0%	1.68 [0.41 , 6.87]	
Total events:	5		3				
Heterogeneity: Not applicab	ole						
Test for overall effect: $Z = 0$	0.72 (P = 0)	.47)					
Test for subgroup difference	es: Chi ² = 2	2.00, $df = 3$	3 (P = 0.57)), $I^2 = 0\%$		+ 0.0 Faun	05 0.1 1 10 200
Footnotes						Favo	ravours ingher SFA
1 OOHOUS							

(1) Total CVD during study period, Prentice 2017



Analysis 1.46. Comparison 1: SFA reduction vs usual diet primary outcomes, Outcome 46: CVD events, subgroup by duration

	lower	SFA	higher	SFA		Risk Ratio	Risk R	atio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Randor	n, 95% CI
1.46.1 up to 24mo								
Black 1994	0	66	2	67	0.3%	0.20 [0.01 , 4.15]		
DART 1989	136	1018	147	1015	14.9%	0.92 [0.74 , 1.15]	_	
Moy 2001	5	117	3	118	1.4%	1.68 [0.41 , 6.87]		
Rose corn oil 1965	15	28	6	13	4.8%	1.16 [0.59 , 2.29]	_	_
Rose olive 1965	11	26	5	13	3.6%	1.10 [0.48 , 2.50]	_	_
Subtotal (95% CI)		1255		1226	25.0%	0.96 [0.78 , 1.16]		
Total events:	167		163				Y	
Heterogeneity: $Tau^2 = 0.0$	0; Chi ² = 2.1	7, df = 4 (P = 0.70; I	$^{2} = 0\%$				
Test for overall effect: Z =	= 0.45 (P = 0)	.65)						
1.46.2 >24 to 48mo								
MRC 1968	62	199	74	194	13.1%	0.82 [0.62, 1.07]	-	
STARS 1992	8	27	20	28	5.4%	0.41 [0.22, 0.78]		
Veterans Admin 1969	97	424	122	422	14.5%	0.79 [0.63 , 1.00]	-	
Subtotal (95% CI)		650		644	33.0%	0.73 [0.56 , 0.95]		
Total events:	167		216				•	
Heterogeneity: $Tau^2 = 0.0$	3; Chi ² = 3.9	7, $df = 2$ (P = 0.14; I	$^{2} = 50\%$				
Test for overall effect: Z =	= 2.31 (P = 0)	.02)						
1.46.3 >48mo								
Oslo Diet-Heart 1966	64	206	90	206	13.6%	0.71 [0.55, 0.92]	-	
WHI 2006 (1)	1399	19541	2145	29294	19.0%	0.98 [0.92 , 1.04]		
Subtotal (95% CI)		19747		29500	32.6%	0.85 [0.63 , 1.16]	•	
Total events:	1463		2235				•	
Heterogeneity: $Tau^2 = 0.0$	4; Chi ² = 5.6	2, $df = 1$ (P = 0.02; I	² = 82%				
Test for overall effect: Z =	= 0.99 (P = 0)	.32)						
1.46.4 unclear duration								
Houtsmuller 1979	8	51	30	51	4.9%	0.27 [0.14, 0.52]		
Ley 2004	11	88	16	88	4.5%	0.69 [0.34 , 1.40]		
Subtotal (95% CI)		139		139	9.4%	0.43 [0.17 , 1.08]	\bullet	
Total events:	19		46				-	
Heterogeneity: $Tau^2 = 0.3$	2; Chi ² = 3.5	9, df = 1 (P = 0.06; I	$^{2} = 72\%$				
Test for overall effect: Z =	= 1.80 (P = 0)	.07)						
Total (95% CI)		21791		31509	100.0%	0.79 [0.66 , 0.93]	♦	
Total events:	1816		2660				, <u>'</u>	
Heterogeneity: $Tau^2 = 0.0$	4; Chi ² = 31.	72, $df = 1$	1 (P = 0.00)	$(18); I^2 = 6$	5%	0.	005 0.1 1	10 200
Test for overall effect: Z =	= 2.72 (P = 0)	.006)				Fav	ours lower SFA	Favours higher SFA
Test for subgroup different	ices: $Chi^2 = 4$	4.71, df = 1	3 (P = 0.19)	$I^2 = 36.3$	%			

Footnotes

(1) Total CVD during study period, Prentice 2017

Analysis 1.47. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 47: CVD events, subgroup by baseline SFA

	lower	SFA	higher SFA			Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
1.47.1 up to 12%E SFA bas	seline						
Subtotal (95% CI)		0		0		Not estimable	
Total events:	0		0				
Heterogeneity: Not applicable	le						
Test for overall effect: Not a	pplicable						
1.47.2 >12 to 15% E SFA ba	aseline						
Black 1994	0	66	2	67	0.3%	0.20 [0.01 , 4.15]	←
DART 1989	136	1018	147	1015	14.9%	0.92 [0.74 , 1.15]	-
Ley 2004	11	88	16	88	4.5%	0.69 [0.34 , 1.40]	_ _
Moy 2001	5	117	3	118	1.4%	1.68 [0.41 , 6.87]	
WHI 2006 (1)	1399	19541	2145	29294	19.0%	0.98 [0.92 , 1.04]	-
Subtotal (95% CI)		20830		30582	40.1%	0.97 [0.91 , 1.03]	
Total events:	1551		2313				
Heterogeneity: $Tau^2 = 0.00$;	Chi ² = 2.7	9, $df = 4$ (P = 0.59; I	$2^2 = 0\%$			
Test for overall effect: $Z = 0$.94 (P = 0	.35)					
1.47.3 >15 to 18% E SFA ba	aseline						
STARS 1992	8	27	20	28	5.4%	0.41 [0.22, 0.78]	
Subtotal (95% CI)		27		28	5.4%	0.41 [0.22, 0.78]	
Total events:	8		20				
Heterogeneity: Not applicable	le						
Test for overall effect: $Z = 2$.75 ($P = 0$.006)					
1.47.4 >18%E SFA baselin	e						
Veterans Admin 1969	97	424	122	422	14.5%	0.79 [0.63, 1.00]	
Subtotal (95% CI)		424		422	14.5%	0.79 [0.63, 1.00]	
Total events:	97		122			- / -	•
Heterogeneity: Not applicable	le						
Test for overall effect: $Z = 1$.99 (P = 0)	.05)					
1.47.5 unclear							
Houtsmuller 1979	8	51	30	51	4.9%	0.27 [0.14, 0.52]	
MRC 1968	62	199	74	194	13.1%	0.82 [0.62, 1.07]	_ _
Oslo Diet-Heart 1966	64	206	90	206	13.6%	0.71 [0.55, 0.92]	
Rose corn oil 1965	15	28	6	13	4.8%	1.16 [0.59 , 2.29]	
Rose olive 1965	11	26	5	13	3.6%	1.10 [0.48, 2.50]	
Subtotal (95% CI)		510		477	40.0%	0.72 [0.51, 1.03]	
Total events:	160		205				
Heterogeneity: $Tau^2 = 0.09$;	$Chi^2 = 12.$	08, $df = 4$	(P = 0.02);	$I^2 = 67\%$			
Test for overall effect: $Z = 1$.80 (P = 0	.07)	, <u>/</u> ,				
Total (95% CI)		21791		31509	100.0%	0.79 [0.66 , 0.93]	
Total events:	1816		2660			L,	•
Heterogeneity: $Tau^2 = 0.04$:	$Chi^2 = 31$.	72, $df = 1$	1 (P = 0.00)	$(08); I^2 = 6$	5%		
Test for overall effect: $Z = 2$.72 ($P = 0$.006)				Fa	avours lower SFA Favours higher SF.
Test for subgroup difference	s: $Chi^2 = 1$	11.84, df =	= 3 (P = 0.0)	$(08), I^2 = 74$	4.7%		C C

Footnotes

(1) Total CVD during study period, Prentice 2017



Analysis 1.48. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 48: CVD events, subgroup by SFA change

	lower	SFA	higher	SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
1.48.1 up to 4% E differen	nce						
DART 1989	136	1018	147	1015	14.9%	0.92 [0.74, 1.15]	1
Ley 2004	11	88	16	88	4.5%	0.69 [0.34, 1.40]	
Moy 2001	5	117	3	118	1.4%	1.68 [0.41, 6.87]	
WHI 2006 (1)	1399	19541	2145	29294	19.0%	0.98 [0.92, 1.04]	
Subtotal (95% CI)		20764		30515	39.8%	0.97 [0.91 , 1.03]	
Total events:	1551		2311				
Heterogeneity: $Tau^2 = 0.00$); Chi ² = 1.7	6, df = 3 (P = 0.62; I	$^{2} = 0\%$			
Test for overall effect: Z =	0.92 (P = 0	.36)					
1.48.2 >4 to 8%E differen	nce						
Black 1994	0	66	2	67	0.3%	0.20 [0.01, 4.15]	
STARS 1992	8	27	20	28	5.4%	0.41 [0.22, 0.78]	
Subtotal (95% CI)		93		95	5.8%	0.40 [0.22, 0.74]	
Total events:	8		22				\bullet
Heterogeneity: $Tau^2 = 0.00$); $Chi^2 = 0.2$	2, df = 1 (P = 0.64; I	$^{2} = 0\%$			
Test for overall effect: $Z =$	2.90 (P = 0)	.004)					
1.48.3 >8% E difference							
Veterans Admin 1969	97	424	122	422	14.5%	0.79 [0.63, 1.00]	-
Subtotal (95% CI)		424		422	14.5%	0.79 [0.63 , 1.00]	
Total events:	97		122				•
Heterogeneity: Not applica	ble						
Test for overall effect: $Z =$	1.99 (P = 0	.05)					
1.48.4 unclear							
Houtsmuller 1979	8	51	30	51	4.9%	0.27 [0.14, 0.52]	—
MRC 1968	62	199	74	194	13.1%	0.82 [0.62, 1.07]	-
Oslo Diet-Heart 1966	64	206	90	206	13.6%	0.71 [0.55, 0.92]	
Rose corn oil 1965	15	28	6	13	4.8%	1.16 [0.59 , 2.29]	
Rose olive 1965	11	26	5	13	3.6%	1.10 [0.48 , 2.50]	
Subtotal (95% CI)		510		477	40.0%	0.72 [0.51 , 1.03]	
Total events:	160		205				•
Heterogeneity: $Tau^2 = 0.09$; $Chi^2 = 12$.	08, df = 4	(P = 0.02);	$I^2 = 67\%$			
Test for overall effect: Z =	1.80 (P = 0	.07)					
Total (95% CI)		21791		31509	100.0%	0.79 [0.66 , 0.93]	
Total events:	1816		2660			- , -	▼
Heterogeneity: $Tau^2 = 0.04$; Chi ² = 31.	72, df = 1	1 (P = 0.00)	$(08); I^2 = 6$	5%	0.0	005 0.1 1 10 200
Test for overall effect: $Z =$	2.72 (P = 0)	.006)				Fave	burs lower SFA Favours higher SFA
Test for subgroup difference	ces: $Chi^2 = 1$	12.67, df =	= 3 (P = 0.00)	$(5), I^2 = 70$	6.3%		

Footnotes

(1) Total CVD during study period, Prentice 2017

	lower	SFA	higher	SFA		Risk Ratio	Risk Ra	tio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random	, 95% CI
1.49.1 Men								
Black 1994	0	66	2	67	0.3%	0.20 [0.01, 4.15]		
DART 1989	136	1018	147	1015	14.9%	0.92 [0.74, 1.15]	-	
MRC 1968	62	199	74	194	13.1%	0.82 [0.62, 1.07]	-	
Oslo Diet-Heart 1966	64	206	90	206	13.6%	0.71 [0.55, 0.92]	-	
Rose corn oil 1965	15	28	6	13	4.8%	1.16 [0.59 , 2.29]		
Rose olive 1965	11	26	5	13	3.6%	1.10 [0.48 , 2.50]	_	-
STARS 1992	8	27	20	28	5.4%	0.41 [0.22, 0.78]		
Veterans Admin 1969	97	424	122	422	14.5%	0.79 [0.63 , 1.00]	-	
Subtotal (95% CI)		1994		1958	70.2%	0.80 [0.69 , 0.93]	•	
Total events:	393		466				*	
Heterogeneity: $Tau^2 = 0.0$	01; $Chi^2 = 9.2$	23, df = 7 (P = 0.24; I	$^{2} = 24\%$				
Test for overall effect: Z	= 2.97 (P = 0)	0.003)						
1.49.2 Women								
WHI 2006 (1)	1399	19541	2145	29294	19.0%	0.98 [0.92, 1.04]		
Subtotal (95% CI)		19541		29294	19.0%	0.98 [0.92 , 1.04]		
Total events:	1399		2145					
Heterogeneity: Not applie	cable							
Test for overall effect: Z	= 0.68 (P = 0)	0.50)						
1.49.3 Mixed, men and	women							
Houtsmuller 1979	8	51	30	51	4.9%	0.27 [0.14, 0.52]		
Ley 2004	11	88	16	88	4.5%	0.69 [0.34 , 1.40]	_	
Moy 2001	5	117	3	118	1.4%	1.68 [0.41 , 6.87]	.	
Subtotal (95% CI)		256		257	10.8%	0.59 [0.23, 1.49]		
Total events:	24		49					
Heterogeneity: $Tau^2 = 0.4$	46; Chi ² = 6.9	93, df = 2 (P = 0.03; I	2 = 71%				
Test for overall effect: Z	= 1.13 (P = 0)	0.26)						
Total (95% CI)		21791		31509	100.0%	0.79 [0.66 , 0.93]	▲	
Total events:	1816		2660				•	
Heterogeneity: $Tau^2 = 0.0$	$04; Chi^2 = 31$.72, $df = 1$	1 (P = 0.00)	$(08); I^2 = 6$	5%	0.0		10 200
Test for overall effect: Z	= 2.72 (P = 0)	0.006)				Favo	ours lower SFA	Favours higher SF.

Analysis 1.49. Comparison 1: SFA reduction vs usual diet primary outcomes, Outcome 49: CVD events, subgroup by sex

Footnotes

(1) Total CVD during study period, Prentice 2017

Test for subgroup differences: Chi² = 6.95, df = 2 (P = 0.03), I² = 71.2%

Analysis 1.50. Comparison 1: SFA reduction vs usual diet primary outcomes, Outcome 50: CVD events, subgroup by CVD risk

	lower	lower SFA		higher SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
1.50.1 Low CVD risk							
Black 1994	0	66	2	67	0.2%	0.20 [0.01, 4.15]	
Veterans Admin 1969	97	424	122	422	12.5%	0.79 [0.63, 1.00]	-
WHI 2006 (1)	1132	18633	1777	27925	17.4%	0.95 [0.89, 1.03]	
Subtotal (95% CI)		19123		28414	30.2%	0.89 [0.75 , 1.06]	4
Total events:	1229		1901				•
Heterogeneity: $Tau^2 = 0.0$	01; Chi ² = 3.3	2, $df = 2$ (P = 0.19; I	$^{2} = 40\%$			
Test for overall effect: Z	= 1.28 (P = 0)	.20)					
1.50.2 Moderate CVD r	isk						
Houtsmuller 1979	8	51	30	51	3.7%	0.27 [0.14, 0.52]	_ —
Ley 2004	11	88	16	88	3.4%	0.69 [0.34 , 1.40]	
Moy 2001	5	117	3	118	1.0%	1.68 [0.41 , 6.87]	
Subtotal (95% CI)		256		257	8.2%	0.59 [0.23, 1.49]	
Total events:	24		49				-
Heterogeneity: $Tau^2 = 0.4$	46; Chi ² = 6.9	3, $df = 2$ (P = 0.03; I	$^{2} = 71\%$			
Test for overall effect: Z	= 1.13 (P = 0)	.26)					
1.50.3 Existing CVD dis	sease						
DART 1989	136	1018	147	1015	13.0%	0.92 [0.74, 1.15]	-
MRC 1968	62	199	74	194	11.1%	0.82 [0.62, 1.07]	-
Oslo Diet-Heart 1966	64	206	90	206	11.7%	0.71 [0.55, 0.92]	+
Rose corn oil 1965	15	28	6	13	3.7%	1.16 [0.59 , 2.29]	_ _
Rose olive 1965	11	26	5	13	2.7%	1.10 [0.48 , 2.50]	_ _
STARS 1992	8	27	20	28	4.2%	0.41 [0.22, 0.78]	
WHI 2006 (2)	225	908	311	1369	15.3%	1.09 [0.94 , 1.27]	_
Subtotal (95% CI)		2412		2838	61.6%	0.86 [0.71 , 1.05]	
Total events:	521		653				*
Heterogeneity: $Tau^2 = 0.0$	04; Chi ² = 16.	43, df = 6	(P = 0.01);	$I^2=63\%$			
Test for overall effect: Z	= 1.45 (P = 0)	.15)					
Total (95% CI)		21791		31509	100.0%	0.83 [0.72 , 0.96]	
Total events:	1774		2603				▼
Heterogeneity: $Tau^2 = 0.0$	03; Chi ² = 34.	25, $df = 1$	2 (P = 0.00)	$(06); I^2 = 6$	5%	0.0	05 0.1 1 10 200
Test for overall effect: Z	= 2.50 (P = 0)	.01)				Favo	urs lower SFA Favours higher SFA

Test for subgroup differences: Chi² = 0.79, df = 2 (P = 0.67), $I^2 = 0\%$

Footnotes

(1) Women without CVD at baseline

(2) Women with CVD at baseline



Analysis 1.51. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 51: CVD events, subgroup by TC reduction

	lower SFA		higher SFA		Risk Ratio		Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
1.51.1 serum chol reduc	ed by at leas	t 0.2mmo	l/L				
DART 1989	136	1018	147	1015	19.7%	0.92 [0.74, 1.15]	+
Houtsmuller 1979	8	51	30	51	7.4%	0.27 [0.14, 0.52]	_ - _
MRC 1968	62	199	74	194	17.7%	0.82 [0.62, 1.07]	-
Moy 2001	5	117	3	118	2.2%	1.68 [0.41 , 6.87]	_ _
Oslo Diet-Heart 1966	64	206	90	206	18.3%	0.71 [0.55, 0.92]	-
Rose corn oil 1965	15	28	6	13	7.3%	1.16 [0.59 , 2.29]	_ _
STARS 1992	8	27	20	28	8.2%	0.41 [0.22, 0.78]	
Veterans Admin 1969	97	424	122	422	19.2%	0.79 [0.63 , 1.00]	-
Subtotal (95% CI)		2070		2047	100.0%	0.74 [0.59, 0.92]	
Total events:	395		492				•
Heterogeneity: $Tau^2 = 0.0$	05; Chi ² = 18.	95, df = 7	(P = 0.008)); $I^2 = 63\%$,		
Test for overall effect: Z	= 2.73 (P = 0)	.006)					
1.51.2 serum chol reduc	ed by <0.2m	mol/L					
Ley 2004	11	88	16	88	0.8%	0.69 [0.34 , 1.40]	
Rose olive 1965	11	26	5	13	0.6%	1.10 [0.48 , 2.50]	_ _
WHI 2006 (1)	1399	19541	2145	29294	98.6%	0.98 [0.92 , 1.04]	
Subtotal (95% CI)		19655		29395	100.0%	0.98 [0.91 , 1.04]	T
Total events:	1421		2166				
Heterogeneity: $Tau^2 = 0.0$	00; $Chi^2 = 1.0$	2, $df = 2$ (P = 0.60; I	$^{2} = 0\%$			
Test for overall effect: Z	= 0.75 (P = 0)	.45)					
1.51.3 serum chol reduc	tion unclear						
Black 1994	0	66	2	67	100.0%	0.20 [0.01 , 4.15]	
Subtotal (95% CI)		66		67	100.0%	0.20 [0.01 , 4.15]	
Total events:	0		2				
Heterogeneity: Not appli-	cable						
Test for overall effect: Z	= 1.04 (P = 0)	.30)					
Test for subgroup differe	nces: Chi ² = 6	5.80, df =	2 (P = 0.03)), $I^2 = 70.6$	%	0.0 Favo	05 0.1 1 10 200 purs lower SFA Favours higher SFA
Footnotes							0.00

(1) Total CVD during study period, Prentice 2017

Analysis 1.52. Comparison 1: SFA reduction vs usual diet - primary outcomes, Outcome 52: CVD events, subgroup decade of publication

	lower	SFA	higher SFA		Risk Ratio		Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
1.52.1 1960s							
MRC 1968	62	199	74	194	13.1%	0.82 [0.62, 1.07]	_
Oslo Diet-Heart 1966	64	206	90	206	13.6%	0.71 [0.55, 0.92]	-
Rose corn oil 1965	15	28	6	13	4.8%	1.16 [0.59 , 2.29]	
Rose olive 1965	11	26	5	13	3.6%	1.10 [0.48 , 2.50]	
Veterans Admin 1969	97	424	122	422	14.5%	0.79 [0.63 , 1.00]	-
Subtotal (95% CI)		883		848	49.6%	0.79 [0.69 , 0.91]	▲
Total events:	249		297				•
Heterogeneity: $Tau^2 = 0.0$	00; Chi ² = 2.5	66, df = 4(P = 0.63;	$I^2 = 0\%$			
Test for overall effect: Z	= 3.25 (P = 0)	.001)					
1.52.2 1970s							
Houtsmuller 1979	8	51	30	51	4.9%	0.27 [0.14, 0.52]	_ _
Subtotal (95% CI)		51		51	4.9%	0.27 [0.14, 0.52]	
Total events:	8		30				•
Heterogeneity: Not appli	cable						
Test for overall effect: Z	= 3.83 (P = 0)	.0001)					
1.52.3 1980s							
DART 1989	136	1018	147	1015	14.9%	0.92 [0.74, 1.15]	4
Subtotal (95% CI)		1018		1015	14.9%	0.92 [0.74 , 1.15]	4
Total events:	136		147				
Heterogeneity: Not appli	cable						
Test for overall effect: Z	= 0.73 (P = 0)	.46)					
1.52.4 1990s							
Black 1994	0	66	2	67	0.3%	0.20 [0.01 , 4.15]	_
STARS 1992	8	27	20	28	5.4%	0.41 [0.22, 0.78]	
Subtotal (95% CI)		93		95	5.8%	0.40 [0.22, 0.74]	
Total events:	8		22				•
Heterogeneity: $Tau^2 = 0.0$	00; $Chi^2 = 0.2$	22, df = 1 (P = 0.64;	$I^2 = 0\%$			
Test for overall effect: Z	= 2.90 (P = 0)	.004)					
1.52.5 2000s							
Ley 2004	11	88	16	88	4.5%	0.69 [0.34 , 1.40]	
Moy 2001	5	117	3	118	1.4%	1.68 [0.41 , 6.87]	_
WHI 2006 (1)	1399	19541	2145	29294	19.0%	0.98 [0.92, 1.04]	+
Subtotal (95% CI)		19746		29500	24.9%	0.98 [0.91 , 1.04]	
Total events:	1415		2164				
Heterogeneity: $Tau^2 = 0.0$	00; $Chi^2 = 1.5$	51, df = 2 (P = 0.47);	$I^2 = 0\%$			
Test for overall effect: Z	= 0.74 (P = 0)	.46)					
Total (95% CI)		21791		31509	100.0%	0.79 [0.66 , 0.93]	♦
Total events:	1816		2660				
Heterogeneity: $Tau^2 = 0.0$	04; Chi ² = 31.	.72, $df = 1$	1 (P = 0.00)	$(08); I^2 = 65$	5%	0.0	05 0.1 1 10 200
Test for overall effect: Z	= 2.72 (P = 0)	.006)				Favo	ours lower SFA Favours higher SF
Test for subgroup differe	ences: $Chi^2 = 2$	27.36, df =	4 (P < 0.0)	001), $I^2 = 8$	35.4%		

Footnotes

(1) Total CVD during study period, Prentice 2017
Comparison 2. SFA reduction vs usual diet - secondary health events

Outcome or subgroup title	No. of studies	No. of partici- pants	Statistical method	Effect size
2.1 MYOCARDIAL INFARCTION	11	53167	Risk Ratio (M-H, Random, 95% Cl)	0.90 [0.80, 1.01]
2.2 MI, SA by low summary risk of bias	3	49857	Risk Ratio (M-H, Random, 95% CI)	0.93 [0.81, 1.08]
2.3 MI, SA aim to reduce SFA	10	52991	Risk Ratio (M-H, Random, 95% CI)	0.89 [0.78, 1.02]
2.4 MI, SA statistically significant SFA re- duction	6	52180	Risk Ratio (M-H, Random, 95% CI)	0.94 [0.85, 1.04]
2.5 MI, SA by TC reduction	9	52952	Risk Ratio (M-H, Random, 95% CI)	0.88 [0.77, 1.01]
2.6 MI, SA excluding WHI	10	4332	Risk Ratio (M-H, Random, 95% CI)	0.85 [0.73, 0.98]
2.7 MI, SA Mantel-Haenszel fixed-effect	11	53167	Risk Ratio (M-H, Fixed, 95% CI)	0.92 [0.84, 1.01]
2.8 MI, SA Peto fixed-effect	11	53167	Peto Odds Ratio (Peto, Fixed, 95% CI)	0.92 [0.83, 1.01]
2.9 MI, subgroup by any substitution	11		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
2.9.1 replaced by PUFA	7	3895	Risk Ratio (M-H, Random, 95% CI)	0.83 [0.67, 1.02]
2.9.2 replaced by MUFA	1	52	Risk Ratio (M-H, Random, 95% CI)	1.40 [0.51, 3.85]
2.9.3 replace by CHO	4	51099	Risk Ratio (M-H, Random, 95% CI)	0.96 [0.86, 1.06]
2.9.4 replaced by protein	3	51044	Risk Ratio (M-H, Random, 95% CI)	0.96 [0.86, 1.07]
2.9.5 replacement unclear	1	235	Risk Ratio (M-H, Random, 95% CI)	2.02 [0.19, 21.94]
2.10 MI, subgroup by main substitution	11		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
2.10.1 replaced by PUFA	6	3840	Risk Ratio (M-H, Random, 95% CI)	0.83 [0.67, 1.04]
2.10.2 replaced by MUFA	1	52	Risk Ratio (M-H, Random, 95% CI)	1.40 [0.51, 3.85]
2.10.3 replace by CHO	3	49066	Risk Ratio (M-H, Random, 95% CI)	0.97 [0.86, 1.09]

Reduction in saturated fat intake for cardiovascular disease (Review)



Cochrane Database of Systematic Reviews

Outcome or subgroup title	No. of studies	No. of partici- pants	Statistical method	Effect size
2.10.4 replaced by protein	0	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
2.10.5 replacement unclear	1	235	Risk Ratio (M-H, Random, 95% CI)	2.02 [0.19, 21.94]
2.11 MI, subgroup by duration	11	53167	Risk Ratio (M-H, Random, 95% CI)	0.90 [0.80, 1.01]
2.11.1 up to 24mo	4	2348	Risk Ratio (M-H, Random, 95% CI)	0.95 [0.77, 1.17]
2.11.2 >24 to 48mo	3	1294	Risk Ratio (M-H, Random, 95% CI)	0.83 [0.64, 1.06]
2.11.3 >48mo	2	49247	Risk Ratio (M-H, Random, 95% CI)	0.81 [0.54, 1.24]
2.11.4 unclear	2	278	Risk Ratio (M-H, Random, 95% CI)	0.41 [0.02, 7.73]
2.12 MI, subgroup by baseline SFA	11	53167	Risk Ratio (M-H, Random, 95% CI)	0.90 [0.80, 1.01]
2.12.1 up to 12%E SFA baseline	0	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
2.12.2 >12 to 15%E SFA baseline	4	51279	Risk Ratio (M-H, Random, 95% CI)	0.96 [0.87, 1.07]
2.12.3 >15 to 18%E SFA baseline	1	55	Risk Ratio (M-H, Random, 95% CI)	0.52 [0.05, 5.39]
2.12.4 >18%E SFA baseline	1	846	Risk Ratio (M-H, Random, 95% CI)	0.76 [0.55, 1.05]
2.12.5 unclear	5	987	Risk Ratio (M-H, Random, 95% CI)	0.84 [0.54, 1.30]
2.13 MI, subgroup by SFA change	11	53167	Risk Ratio (M-H, Random, 95% CI)	0.90 [0.80, 1.01]
2.13.1 up to 4%E difference	4	51279	Risk Ratio (M-H, Random, 95% CI)	0.96 [0.87, 1.07]
2.13.2 >4 to 8%E difference	1	55	Risk Ratio (M-H, Random, 95% CI)	0.52 [0.05, 5.39]
2.13.3 >8%E difference	1	846	Risk Ratio (M-H, Random, 95% CI)	0.76 [0.55, 1.05]
2.13.4 unclear	5	987	Risk Ratio (M-H, Random, 95% CI)	0.84 [0.54, 1.30]

Reduction in saturated fat intake for cardiovascular disease (Review)



Cochrane Database of Systematic Reviews

Outcome or subgroup title	No. of studies	No. of partici- pants	Statistical method	Effect size
2.14 MI, subgroup by sex	11	53167	Risk Ratio (M-H, Random, 95% CI)	0.90 [0.80, 1.01]
2.14.1 Men	7	3819	Risk Ratio (M-H, Random, 95% CI)	0.85 [0.73, 0.98]
2.14.2 Women	1	48835	Risk Ratio (M-H, Random, 95% CI)	0.97 [0.86, 1.09]
2.14.3 Mixed, men and women	3	513	Risk Ratio (M-H, Random, 95% CI)	0.75 [0.13, 4.47]
2.15 MI, subgroup by CVD risk	11	53167	Risk Ratio (M-H, Random, 95% CI)	0.90 [0.80, 1.01]
2.15.1 Low CVD risk	2	49681	Risk Ratio (M-H, Random, 95% CI)	0.90 [0.72, 1.13]
2.15.2 Moderate CVD risk	3	513	Risk Ratio (M-H, Random, 95% CI)	0.75 [0.13, 4.47]
2.15.3 Existing CVD disease	6	2973	Risk Ratio (M-H, Random, 95% CI)	0.87 [0.74, 1.03]
2.16 MI, subgroup by TC reduction	11		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
2.16.1 serum chol reduced by at least 0.2mmol/L	8	4117	Risk Ratio (M-H, Random, 95% CI)	0.83 [0.70, 0.98]
2.16.2 serum chol reduced by <0.2mmol/L	3	49050	Risk Ratio (M-H, Random, 95% CI)	0.98 [0.87, 1.10]
2.16.3 serum chol reduction unclear	0	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
2.17 MI, subgroup decade of publication	11	53167	Risk Ratio (M-H, Random, 95% Cl)	0.90 [0.80, 1.01]
2.17.1 1960s	5	1731	Risk Ratio (M-H, Random, 95% Cl)	0.80 [0.64, 1.00]
2.17.2 1970s	1	102	Risk Ratio (M-H, Random, 95% Cl)	0.08 [0.00, 1.33]
2.17.3 1980s	1	2033	Risk Ratio (M-H, Random, 95% Cl)	0.91 [0.73, 1.14]
2.17.4 1990s	1	55	Risk Ratio (M-H, Random, 95% CI)	0.52 [0.05, 5.39]
2.17.5 2000s	3	49246	Risk Ratio (M-H, Random, 95% Cl)	0.98 [0.87, 1.10]

Reduction in saturated fat intake for cardiovascular disease (Review)



Cochrane Database of Systematic Reviews

Outcome or subgroup title	No. of studies	No. of partici- pants	Statistical method	Effect size
2.18 NON-FATAL MYOCARDIAL INFARCTION	8	52834	Risk Ratio (M-H, Random, 95% CI)	0.97 [0.87, 1.07]
2.19 Non-fatal MI, SA by low summary risk of bias	2	49681	Risk Ratio (M-H, Random, 95% CI)	0.89 [0.58, 1.35]
2.20 Non-fatal MI, SA aim to reduce SFA	8	52834	Risk Ratio (M-H, Random, 95% CI)	0.97 [0.87, 1.07]
2.21 Non-fatal MI, SA statistically signifi- cant SFA reduction	4	51949	Risk Ratio (M-H, Random, 95% CI)	0.90 [0.72, 1.14]
2.22 Non-fatal MI, SA by TC reduction	7	52795	Risk Ratio (M-H, Random, 95% CI)	0.97 [0.87, 1.07]
2.23 Non-fatal MI, SA excluding WHI	7	3999	Risk Ratio (M-H, Random, 95% CI)	0.81 [0.64, 1.04]
2.24 Non-fatal MI, SA Mantel-Haenszel fixed-effect	8	52834	Risk Ratio (M-H, Fixed, 95% CI)	0.97 [0.87, 1.08]
2.25 Non-fatal MI, SA Peto fixed-effect	8	52834	Peto Odds Ratio (Peto, Fixed, 95% CI)	0.97 [0.87, 1.08]
2.26 Non-fatal MI, subgroup by any substi- tution	8		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
2.26.1 replaced by PUFA	5	3738	Risk Ratio (M-H, Random, 95% CI)	0.80 [0.63, 1.03]
2.26.2 replaced by MUFA	1	52	Risk Ratio (M-H, Random, 95% CI)	1.20 [0.42, 3.45]
2.26.3 replace by CHO	2	50868	Risk Ratio (M-H, Random, 95% CI)	0.93 [0.72, 1.21]
2.26.4 replaced by protein	2	50868	Risk Ratio (M-H, Random, 95% CI)	0.93 [0.72, 1.21]
2.26.5 replacement unclear	1	235	Risk Ratio (M-H, Random, 95% CI)	2.02 [0.19, 21.94]
2.27 Non-fatal MI, subgroup by main sub- stitution	8		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
2.27.1 replaced by PUFA	5	3738	Risk Ratio (M-H, Random, 95% CI)	0.80 [0.63, 1.03]
2.27.2 replaced by MUFA	1	52	Risk Ratio (M-H, Random, 95% CI)	1.20 [0.42, 3.45]
2.27.3 replace by CHO	1	48835	Risk Ratio (M-H, Random, 95% CI)	1.01 [0.90, 1.13]

Reduction in saturated fat intake for cardiovascular disease (Review)



Cochrane Database of Systematic Reviews

Outcome or subgroup title	No. of studies	No. of partici- pants	Statistical method	Effect size
2.27.4 replaced by protein	0	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
2.27.5 replacement unclear	1	235	Risk Ratio (M-H, Random, 95% CI)	2.02 [0.19, 21.94]
2.28 Non-fatal MI, subgroup by duration	8	52834	Risk Ratio (M-H, Random, 95% CI)	0.97 [0.87, 1.07]
2.28.1 up to 24mo	4	2348	Risk Ratio (M-H, Random, 95% CI)	0.83 [0.57, 1.22]
2.28.2 >24 to 48mo	2	1239	Risk Ratio (M-H, Random, 95% CI)	0.82 [0.53, 1.27]
2.28.3 >48mo	2	49247	Risk Ratio (M-H, Random, 95% CI)	0.99 [0.88, 1.12]
2.28.4 unclear	0	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
2.29 Non-fatal MI, subgroup by baseline SFA	8	52834	Risk Ratio (M-H, Random, 95% CI)	0.97 [0.87, 1.07]
2.29.1 up to 12%E SFA baseline	0	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
2.29.2 >12 to 15%E SFA baseline	3	51103	Risk Ratio (M-H, Random, 95% CI)	0.97 [0.83, 1.13]
2.29.3 >15 to 18%E SFA baseline	0	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
2.29.4 >18%E SFA baseline	1	846	Risk Ratio (M-H, Random, 95% CI)	0.62 [0.31, 1.21]
2.29.5 unclear	4	885	Risk Ratio (M-H, Random, 95% CI)	0.91 [0.65, 1.27]
2.30 Non-fatal MI, subgroup by SFA change	8	52834	Risk Ratio (M-H, Random, 95% CI)	0.97 [0.87, 1.07]
2.30.1 up to 4%E difference	3	51103	Risk Ratio (M-H, Random, 95% CI)	0.97 [0.83, 1.13]
2.30.2 >4 to 8%E difference	0	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
2.30.3 >8%E difference	1	846	Risk Ratio (M-H, Random, 95% CI)	0.62 [0.31, 1.21]
2.30.4 unclear	4	885	Risk Ratio (M-H, Random, 95% CI)	0.91 [0.65, 1.27]

Reduction in saturated fat intake for cardiovascular disease (Review)



Cochrane Database of Systematic Reviews

Outcome or subgroup title	No. of studies	No. of partici- pants	Statistical method	Effect size
2.31 Non-fatal MI, subgroup by sex	8	52834	Risk Ratio (M-H, Random, 95% CI)	0.97 [0.87, 1.07]
2.31.1 Men	6	3764	Risk Ratio (M-H, Random, 95% CI)	0.81 [0.63, 1.03]
2.31.2 Women	1	48835	Risk Ratio (M-H, Random, 95% CI)	1.01 [0.90, 1.13]
2.31.3 Mixed, men and women	1	235	Risk Ratio (M-H, Random, 95% CI)	2.02 [0.19, 21.94]
2.32 Non-fatal MI, subgroup by CVD risk	8	52834	Risk Ratio (M-H, Random, 95% CI)	0.95 [0.80, 1.13]
2.32.1 Low CVD risk	2	47404	Risk Ratio (M-H, Random, 95% CI)	0.87 [0.68, 1.12]
2.32.2 Moderate CVD risk	1	235	Risk Ratio (M-H, Random, 95% CI)	2.02 [0.19, 21.94]
2.32.3 Existing CVD disease	6	5195	Risk Ratio (M-H, Random, 95% CI)	1.00 [0.76, 1.31]
2.33 Non-fatal MI, subgroup by TC reduc- tion	8		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
2.33.1 serum chol reduced by at least 0.2mmol/L	6	3960	Risk Ratio (M-H, Random, 95% CI)	0.80 [0.62, 1.03]
2.33.2 serum chol reduced by <0.2mmol/L	2	48874	Risk Ratio (M-H, Random, 95% CI)	1.01 [0.90, 1.13]
2.33.3 serum chol reduction unclear	0	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
2.34 Non-fatal MI, subgroup decade of pub- lication	8	52834	Risk Ratio (M-H, Random, 95% CI)	0.97 [0.87, 1.07]
2.34.1 1960s	5	1731	Risk Ratio (M-H, Random, 95% CI)	0.84 [0.62, 1.13]
2.34.2 1970s	0	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
2.34.3 1980s	1	2033	Risk Ratio (M-H, Random, 95% CI)	0.74 [0.48, 1.14]
2.34.4 1990s	0	0	Risk Ratio (M-H, Random, 95% Cl)	Not estimable
2.34.5 2000s	2	49070	Risk Ratio (M-H, Random, 95% Cl)	1.01 [0.90, 1.13]

Reduction in saturated fat intake for cardiovascular disease (Review)



Cochrane Database of Systematic Reviews

Outcome or subgroup title	No. of studies	No. of partici- pants	Statistical method	Effect size
2.35 STROKE	7	50952	Risk Ratio (M-H, Random, 95% CI)	0.92 [0.68, 1.25]
2.36 Stroke, SA by low summary risk of bias	3	49857	Risk Ratio (M-H, Random, 95% CI)	0.76 [0.42, 1.38]
2.37 Stroke, SA aim to reduce SFA	6	50776	Risk Ratio (M-H, Random, 95% CI)	1.01 [0.90, 1.14]
2.38 Stroke, SA statistically significant SFA reduction	5	50147	Risk Ratio (M-H, Random, 95% CI)	0.83 [0.55, 1.25]
2.39 Stroke, SA by TC reduction	6	50776	Risk Ratio (M-H, Random, 95% CI)	1.01 [0.90, 1.14]
2.40 Stroke, SA excluding WHI	6	2117	Risk Ratio (M-H, Random, 95% CI)	0.63 [0.35, 1.14]
2.41 Stroke, SA Mantel-Haenszel fixed-ef- fect	7	50952	Risk Ratio (M-H, Fixed, 95% CI)	1.01 [0.89, 1.13]
2.42 Stroke, SA Peto fixed-effect	7	50952	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.01 [0.89, 1.14]
2.43 Stroke, subgroup by any substitution	7		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
2.43.1 replaced by PUFA	4	1706	Risk Ratio (M-H, Random, 95% Cl)	0.68 [0.37, 1.27]
2.43.2 replaced by MUFA	0	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
2.43.3 replace by CHO	3	49066	Risk Ratio (M-H, Random, 95% CI)	0.73 [0.29, 1.87]
2.43.4 replaced by protein	2	49011	Risk Ratio (M-H, Random, 95% CI)	0.65 [0.15, 2.75]
2.43.5 replacement unclear	1	235	Risk Ratio (M-H, Random, 95% Cl)	1.01 [0.06, 15.93]
2.44 Stroke, subgroup by main substitution	7		Risk Ratio (M-H, Random, 95% Cl)	Subtotals only
2.44.1 replaced by PUFA	3	1651	Risk Ratio (M-H, Random, 95% Cl)	0.92 [0.31, 2.69]
2.44.2 replaced by MUFA	0	0	Risk Ratio (M-H, Random, 95% Cl)	Not estimable
2.44.3 replace by CHO	3	49066	Risk Ratio (M-H, Random, 95% Cl)	0.73 [0.29, 1.87]

Reduction in saturated fat intake for cardiovascular disease (Review)



Cochrane Database of Systematic Reviews

Outcome or subgroup title	No. of studies	No. of partici- pants	Statistical method	Effect size
2.44.4 replaced by protein	0	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
2.44.5 replacement unclear	1	235	Risk Ratio (M-H, Random, 95% CI)	1.01 [0.06, 15.93]
2.45 Stroke, subgroup by duration	6	50559	Risk Ratio (M-H, Random, 95% CI)	0.91 [0.67, 1.23]
2.45.1 up to 24mo	1	235	Risk Ratio (M-H, Random, 95% CI)	1.01 [0.06, 15.93]
2.45.2 >24 to 48mo	2	901	Risk Ratio (M-H, Random, 95% CI)	0.57 [0.30, 1.11]
2.45.3 >48mo	2	49247	Risk Ratio (M-H, Random, 95% CI)	1.03 [0.91, 1.16]
2.45.4 unclear duration	1	176	Risk Ratio (M-H, Random, 95% CI)	0.20 [0.02, 1.68]
2.46 Stroke, subgroup by baseline SFA	6	50559	Risk Ratio (M-H, Random, 95% CI)	0.91 [0.67, 1.23]
2.46.1 up to 12%E SFA baseline	0	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
2.46.2 >12 to 15%E SFA baseline	3	49246	Risk Ratio (M-H, Random, 95% CI)	0.91 [0.50, 1.66]
2.46.3 >15 to 18%E SFA baseline	1	55	Risk Ratio (M-H, Random, 95% CI)	0.35 [0.01, 8.12]
2.46.4 >18%E SFA baseline	1	846	Risk Ratio (M-H, Random, 95% CI)	0.59 [0.30, 1.15]
2.46.5 unclear	1	412	Risk Ratio (M-H, Random, 95% CI)	2.00 [0.18, 21.89]
2.47 Stroke, subgroup by SFA change	6	50559	Risk Ratio (M-H, Random, 95% Cl)	0.91 [0.67, 1.23]
2.47.1 up to 4%E difference	3	49246	Risk Ratio (M-H, Random, 95% CI)	0.91 [0.50, 1.66]
2.47.2 >4 to 8%E difference	1	55	Risk Ratio (M-H, Random, 95% Cl)	0.35 [0.01, 8.12]
2.47.3 >8%E difference	1	846	Risk Ratio (M-H, Random, 95% CI)	0.59 [0.30, 1.15]
2.47.4 unclear	1	412	Risk Ratio (M-H, Random, 95% CI)	2.00 [0.18, 21.89]

Reduction in saturated fat intake for cardiovascular disease (Review)



Cochrane Database of Systematic Reviews

Outcome or subgroup title	No. of studies	No. of partici- pants	Statistical method	Effect size
2.48 Stroke, subgroup by sex	6	50559	Risk Ratio (M-H, Random, 95% CI)	0.91 [0.67, 1.23]
2.48.1 Men	3	1313	Risk Ratio (M-H, Random, 95% CI)	0.63 [0.33, 1.18]
2.48.2 Women	1	48835	Risk Ratio (M-H, Random, 95% CI)	1.03 [0.91, 1.16]
2.48.3 Mixed, men and women	2	411	Risk Ratio (M-H, Random, 95% CI)	0.37 [0.07, 1.97]
2.49 Stroke, subgroup by CVD risk	6	50559	Risk Ratio (M-H, Random, 95% CI)	1.00 [0.89, 1.11]
2.49.1 Low CVD risk	2	47404	Risk Ratio (M-H, Random, 95% CI)	0.86 [0.52, 1.42]
2.49.2 Moderate CVD risk	2	411	Risk Ratio (M-H, Random, 95% CI)	0.37 [0.07, 1.97]
2.49.3 Existing CVD disease	3	2744	Risk Ratio (M-H, Random, 95% CI)	1.01 [0.86, 1.18]
2.50 Stroke, subgroup by TC reduction	7		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
2.50.1 serum chol reduced by at least 0.2mmol/L	5	1941	Risk Ratio (M-H, Random, 95% CI)	0.70 [0.38, 1.28]
2.50.2 serum chol reduced by <0.2mmol/L	2	49011	Risk Ratio (M-H, Random, 95% CI)	0.65 [0.15, 2.75]
2.50.3 serum chol reduction unclear	0	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
2.51 Stroke, subgroup decade of publica- tion	7	50952	Risk Ratio (M-H, Random, 95% CI)	0.92 [0.68, 1.25]
2.51.1 1960s	3	1651	Risk Ratio (M-H, Random, 95% Cl)	0.92 [0.31, 2.69]
2.51.2 1970s	0	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
2.51.3 1980s	0	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
2.51.4 1990s	1	55	Risk Ratio (M-H, Random, 95% Cl)	0.35 [0.01, 8.12]
2.51.5 2000s	3	49246	Risk Ratio (M-H, Random, 95% Cl)	0.91 [0.50, 1.66]

Reduction in saturated fat intake for cardiovascular disease (Review)



Cochrane Database of Systematic Reviews

Outcome or subgroup title	No. of studies	No. of partici- pants	Statistical method	Effect size
2.52 CORONARY HEART DISEASE MORTALI- TY	9	53159	Risk Ratio (M-H, Random, 95% CI)	0.97 [0.82, 1.16]
2.53 CHD mortality, SA by low summary risk of bias	3	50139	Risk Ratio (M-H, Random, 95% CI)	1.05 [0.77, 1.43]
2.54 CHD mortality, SA aim to reduce SFA	9	53159	Risk Ratio (M-H, Random, 95% CI)	0.97 [0.82, 1.16]
2.55 CHD mortality, SA statistically signifi- cant SFA reduction	4	52172	Risk Ratio (M-H, Random, 95% CI)	1.02 [0.84, 1.24]
2.56 CHD mortality, SA by TC reduction	8	53120	Risk Ratio (M-H, Random, 95% CI)	0.97 [0.81, 1.16]
2.57 CHD mortality, SA excluding WHI	8	4324	Risk Ratio (M-H, Random, 95% CI)	0.97 [0.76, 1.24]
2.58 CHD mortality, SA Mantel-Haenszel fixed-effect	9	53159	Risk Ratio (M-H, Fixed, 95% CI)	0.97 [0.86, 1.10]
2.59 CHD mortality, SA Peto fixed-effect	9	53159	Peto Odds Ratio (Peto, Fixed, 95% CI)	0.97 [0.85, 1.11]
2.60 CHD mortality, subgroup by any sub- stitution	9		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
2.60.1 replaced by PUFA	7	4298	Risk Ratio (M-H, Random, 95% CI)	0.98 [0.74, 1.28]
2.60.2 replaced by MUFA	1	52	Risk Ratio (M-H, Random, 95% CI)	3.00 [0.33, 26.99]
2.60.3 replaced by CHO	2	50868	Risk Ratio (M-H, Random, 95% CI)	0.99 [0.85, 1.16]
2.60.4 replaced by protein	2	50868	Risk Ratio (M-H, Random, 95% CI)	0.99 [0.85, 1.16]
2.60.5 replacement unclear	0	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
2.61 CHD mortality, subgroup by main sub- stitution	9		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
2.61.1 replaced by PUFA	7	4298	Risk Ratio (M-H, Random, 95% Cl)	0.98 [0.74, 1.28]
2.61.2 replaced by MUFA	1	52	Risk Ratio (M-H, Random, 95% Cl)	3.00 [0.33, 26.99]
2.61.3 replaced by CHO	1	48835	Risk Ratio (M-H, Random, 95% CI)	0.99 [0.82, 1.20]

Reduction in saturated fat intake for cardiovascular disease (Review)



Cochrane Database of Systematic Reviews

Outcome or subgroup title	No. of studies	No. of partici- pants	Statistical method	Effect size
2.61.4 replaced by protein	0	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
2.61.5 replacement unclear	0	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
2.62 CHD mortality, subgroup by duration	9		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
2.62.1 up to 24mo	3	2113	Risk Ratio (M-H, Random, 95% CI)	1.02 [0.78, 1.33]
2.62.2 >24 to 48months	2	1239	Risk Ratio (M-H, Random, 95% CI)	0.87 [0.64, 1.19]
2.62.3 >48 months	3	49705	Risk Ratio (M-H, Random, 95% CI)	1.02 [0.72, 1.45]
2.62.4 unclear duration	1	102	Risk Ratio (M-H, Random, 95% CI)	0.09 [0.01, 1.60]
2.63 CHD mortality, subgroup by baseline SFA	9		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
2.63.1 up to 12%E SFA baseline	0	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
2.63.2 >12% to 15%E SFA baseline	3	51326	Risk Ratio (M-H, Random, 95% CI)	1.07 [0.86, 1.34]
2.63.3 >15 to 18%E SFA baseline	0	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
2.63.4 >18%E SFA baseline	1	846	Risk Ratio (M-H, Random, 95% CI)	0.82 [0.55, 1.21]
2.63.5 unclear	5	987	Risk Ratio (M-H, Random, 95% CI)	0.85 [0.56, 1.29]
2.64 CHD mortality, subgroup by SFA change	9		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
2.64.1 up to 4%E difference	3	51326	Risk Ratio (M-H, Random, 95% CI)	1.07 [0.86, 1.34]
2.64.2 >4 to 8%E difference	0	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
2.64.3 >8%E difference	1	846	Risk Ratio (M-H, Random, 95% CI)	0.82 [0.55, 1.21]
2.64.4 unclear	5	987	Risk Ratio (M-H, Random, 95% CI)	0.85 [0.56, 1.29]

Reduction in saturated fat intake for cardiovascular disease (Review)



Cochrane Database of Systematic Reviews

Outcome or subgroup title	No. of studies	No. of partici- pants	Statistical method	Effect size
2.65 CHD mortality, subgroup by sex	9		Risk Ratio (M-H, Random, 95% Cl)	Subtotals only
2.65.1 Men	7	4222	Risk Ratio (M-H, Random, 95% Cl)	0.98 [0.79, 1.23]
2.65.2 Women	1	48835	Risk Ratio (M-H, Random, 95% Cl)	0.99 [0.82, 1.20]
2.65.3 Mixed, men and women	1	102	Risk Ratio (M-H, Random, 95% Cl)	0.09 [0.01, 1.60]
2.66 CHD mortality, subgroup by CVD risk	9		Risk Ratio (M-H, Random, 95% Cl)	Subtotals only
2.66.1 Low CVD risk	2	47404	Risk Ratio (M-H, Random, 95% CI)	0.95 [0.78, 1.16]
2.66.2 Moderate CVD risk	1	102	Risk Ratio (M-H, Random, 95% Cl)	0.09 [0.01, 1.60]
2.66.3 Existing CVD disease	7	5653	Risk Ratio (M-H, Random, 95% Cl)	1.03 [0.83, 1.27]
2.67 CHD mortality, subgroup by TC reduc- tion	9		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
2.67.1 serum chol reduced by at least 0.2mmol/L	7	4285	Risk Ratio (M-H, Random, 95% Cl)	0.96 [0.75, 1.24]
2.67.2 serum chol reduced by <0.2mmol/L	2	48874	Risk Ratio (M-H, Random, 95% CI)	0.99 [0.82, 1.20]
2.67.3 serum chol reduction unclear	0	0	Risk Ratio (M-H, Random, 95% Cl)	Not estimable
2.68 CHD mortality, subgroup decade of publication	9	53159	Risk Ratio (M-H, Random, 95% Cl)	0.97 [0.82, 1.16]
2.68.1 1960s	5	1731	Risk Ratio (M-H, Random, 95% Cl)	0.84 [0.66, 1.06]
2.68.2 1970s	2	560	Risk Ratio (M-H, Random, 95% Cl)	0.54 [0.03, 9.26]
2.68.3 1980s	1	2033	Risk Ratio (M-H, Random, 95% Cl)	1.00 [0.76, 1.30]
2.68.4 1990s	0	0	Risk Ratio (M-H, Random, 95% Cl)	Not estimable
2.68.5 2000s	1	48835	Risk Ratio (M-H, Random, 95% Cl)	0.99 [0.82, 1.20]

Reduction in saturated fat intake for cardiovascular disease (Review)



Cochrane Database of Systematic Reviews

Outcome or subgroup title	No. of studies	No. of partici- pants	Statistical method	Effect size
2.69 CORONARY HEART DISEASE EVENTS	11	53199	Risk Ratio (M-H, Random, 95% CI)	0.83 [0.68, 1.01]
2.70 CHD events, SA by low summary risk of bias	3	49857	Risk Ratio (M-H, Random, 95% CI)	0.92 [0.77, 1.10]
2.71 CHD events, SA excluding WHI	10	4364	Risk Ratio (M-H, Random, 95% CI)	0.80 [0.62, 1.03]
2.72 CHD events, SA statistically significant SFA reduction	6	52212	Risk Ratio (M-H, Random, 95% CI)	0.91 [0.77, 1.06]
2.73 CHD events, SA by TC reduction	9	52984	Risk Ratio (M-H, Random, 95% CI)	0.80 [0.65, 0.99]
2.74 CHD events, SA aim to reduce SFA	10	53023	Risk Ratio (M-H, Random, 95% CI)	0.82 [0.67, 1.00]
2.75 CHD events, SA Mantel-Haenszel fixed- effect	11	53199	Risk Ratio (M-H, Fixed, 95% CI)	0.91 [0.84, 0.99]
2.76 CHD events, SA Peto fixed-effect	11	53199	Peto Odds Ratio (Peto, Fixed, 95% CI)	0.90 [0.83, 0.99]
2.77 CHD events, subgroup by any substi- tution	11		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
2.77.1 replaced by PUFA	7	3895	Risk Ratio (M-H, Random, 95% CI)	0.76 [0.57, 1.00]
2.77.2 replaced by MUFA	1	52	Risk Ratio (M-H, Random, 95% CI)	1.50 [0.62, 3.61]
2.77.3 replaced by CHO	4	51104	Risk Ratio (M-H, Random, 95% CI)	0.93 [0.78, 1.11]
2.77.4 replaced by protein	3	51044	Risk Ratio (M-H, Random, 95% CI)	0.96 [0.88, 1.05]
2.77.5 replacement unclear	1	267	Risk Ratio (M-H, Random, 95% CI)	2.93 [0.31, 27.84]
2.78 CHD events, subgroup by main substi- tution	11		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
2.78.1 replaced by PUFA	6	3840	Risk Ratio (M-H, Random, 95% CI)	0.79 [0.60, 1.04]
2.78.2 replaced by MUFA	1	52	Risk Ratio (M-H, Random, 95% Cl)	1.50 [0.62, 3.61]
2.78.3 replaced by CHO	3	49071	Risk Ratio (M-H, Random, 95% Cl)	0.82 [0.39, 1.72]

Reduction in saturated fat intake for cardiovascular disease (Review)



Cochrane Database of Systematic Reviews

Outcome or subgroup title	No. of studies	No. of partici- pants	Statistical method	Effect size
2.78.4 replaced by protein	0	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
2.78.5 replacement unclear	1	267	Risk Ratio (M-H, Random, 95% CI)	2.93 [0.31, 27.84]
2.79 CHD events, subgroup by duration	11		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
2.79.1 up to 24 months	4	2380	Risk Ratio (M-H, Random, 95% CI)	1.01 [0.76, 1.35]
2.79.2 >24 to 48 months	3	1294	Risk Ratio (M-H, Random, 95% CI)	0.79 [0.55, 1.13]
2.79.3 >48 months	2	49247	Risk Ratio (M-H, Random, 95% CI)	0.85 [0.63, 1.15]
2.79.4 unclear duration	2	278	Risk Ratio (M-H, Random, 95% CI)	0.60 [0.10, 3.58]
2.80 CHD events, subgroup by baseline SFA	11		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
2.80.1 up to 12%E SFA baseline	0	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
2.80.2 >12 to 15%E SFA baseline	4	51311	Risk Ratio (M-H, Random, 95% CI)	0.96 [0.88, 1.06]
2.80.3 >15 to 18%E SFA baseline	1	55	Risk Ratio (M-H, Random, 95% CI)	0.31 [0.10, 1.01]
2.80.4 >18%E SFA baseline	1	846	Risk Ratio (M-H, Random, 95% CI)	0.77 [0.56, 1.04]
2.80.5 unclear	5	987	Risk Ratio (M-H, Random, 95% CI)	0.78 [0.49, 1.26]
2.81 CHD events, subgroup by SFA change	11		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
2.81.1 up to 4%E difference	4	51311	Risk Ratio (M-H, Random, 95% CI)	0.96 [0.88, 1.06]
2.81.2 >4 to 8%E difference	1	55	Risk Ratio (M-H, Random, 95% CI)	0.31 [0.10, 1.01]
2.81.3 >8%E difference	1	846	Risk Ratio (M-H, Random, 95% CI)	0.77 [0.56, 1.04]
2.81.4 unclear	5	987	Risk Ratio (M-H, Random, 95% CI)	0.78 [0.49, 1.26]

Reduction in saturated fat intake for cardiovascular disease (Review)



Cochrane Database of Systematic Reviews

Outcome or subgroup title	No. of studies	No. of partici- pants	Statistical method	Effect size
2.82 CHD events, subgroup by sex	11		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
2.82.1 Men	7	3819	Risk Ratio (M-H, Random, 95% CI)	0.84 [0.70, 1.02]
2.82.2 Women	1	48835	Risk Ratio (M-H, Random, 95% CI)	0.97 [0.87, 1.07]
2.82.3 Mixed, men and women	3	545	Risk Ratio (M-H, Random, 95% CI)	0.88 [0.18, 4.36]
2.83 CHD events, subgroup by CVD risk	11		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
2.83.1 Low CVD risk	2	47404	Risk Ratio (M-H, Random, 95% CI)	0.90 [0.76, 1.05]
2.83.2 Moderate CVD risk	3	545	Risk Ratio (M-H, Random, 95% CI)	0.88 [0.18, 4.36]
2.83.3 Existing CVD disease	7	5250	Risk Ratio (M-H, Random, 95% CI)	0.94 [0.75, 1.16]
2.84 CHD events, subgroup by TC reduction	11		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
2.84.1 serum chol reduced by at least 0.2mmol/L	8	4149	Risk Ratio (M-H, Random, 95% CI)	0.76 [0.58, 0.99]
2.84.2 serum chol reduced by <0.2mmol/L	3	49050	Risk Ratio (M-H, Random, 95% CI)	0.97 [0.88, 1.08]
2.84.3 serum chol reduction unclear	0	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
2.85 CHD events, subgroup decade of pub- lication	11	53201	Risk Ratio (M-H, Random, 95% CI)	0.83 [0.68, 1.01]
2.85.1 1960s	5	1731	Risk Ratio (M-H, Random, 95% CI)	0.84 [0.68, 1.05]
2.85.2 1970s	1	102	Risk Ratio (M-H, Random, 95% Cl)	0.27 [0.14, 0.52]
2.85.3 1980s	1	2033	Risk Ratio (M-H, Random, 95% CI)	0.91 [0.73, 1.14]
2.85.4 1990s	1	57	Risk Ratio (M-H, Random, 95% Cl)	0.33 [0.10, 1.09]
2.85.5 2000s	3	49278	Risk Ratio (M-H, Random, 95% CI)	0.97 [0.88, 1.08]

Reduction in saturated fat intake for cardiovascular disease (Review)



Cochrane Database of Systematic Reviews

Outcome or subgroup title	No. of studies	No. of partici- pants	Statistical method	Effect size
2.86 DIABETES DIAGNOSES	1		Risk Ratio (M-H, Random, 95% CI)	Subtotals only

Analysis 2.1. Comparison 2: SFA reduction vs usual diet secondary health events, Outcome 1: MYOCARDIAL INFARCTION

	Lower	SFA	Higher SFA		Risk Ratio		Risk F	atio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Rando	m, 95% CI
DART 1989	132	1018	144	1015	21.9%	0.91 [0.73 , 1.14]	-	
Houtsmuller 1979	0	51	6	51	0.2%	0.08 [0.00 , 1.33]	←	
Ley 2004	4	88	3	88	0.6%	1.33 [0.31 , 5.78]		
MRC 1968	39	199	40	194	8.2%	0.95 [0.64 , 1.41]	_	-
Moy 2001	2	117	1	118	0.2%	2.02 [0.19, 21.94]		
Oslo Diet-Heart 1966	34	206	54	206	8.6%	0.63 [0.43, 0.92]		
Rose corn oil 1965	9	28	3	13	1.1%	1.39 [0.45 , 4.31]		
Rose olive 1965	7	26	2	13	0.7%	1.75 [0.42 , 7.27]		•
STARS 1992	1	27	2	28	0.3%	0.52 [0.05 , 5.39]		
Veterans Admin 1969	54	424	71	422	11.4%	0.76 [0.55 , 1.05]	-	
WHI 2006	435	19541	671	29294	46.7%	0.97 [0.86 , 1.09]	•	
Total (95% CI)		21725		31442	100.0%	0.90 [0.80 , 1.01]	•	
Total events:	717		997				•	
Heterogeneity: $Tau^2 = 0.0$	0; Chi ² = 11.	07, df = 1	0 (P = 0.35)); $I^2 = 10\%$, D		0 02 0 1 1	10 50
Test for overall effect: Z =	= 1.72 (P = 0)	.09)				F	avours lower SFA	Favours higher SFA
Test for subgroup differer	nces: Not app	licable						

Analysis 2.2. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 2: MI, SA by low summary risk of bias

	Lower	SFA	Higher	Higher SFA		Risk Ratio	Risk R	atio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Randor	n, 95% CI
Ley 2004	4	88	3	88	1.0%	1.33 [0.31 , 5.78]		
Veterans Admin 1969	54	424	71	422	17.5%	0.76 [0.55 , 1.05]		
WHI 2006	435	19541	671	29294	81.6%	0.97 [0.86 , 1.09]	-	
Total (95% CI)		20053		29804	100.0%	0.93 [0.81 , 1.08]		
Total events:	493		745					
Heterogeneity: $Tau^2 = 0.0$	00; Chi ² = 2.1	9, $df = 2$ (P = 0.33; I	$l^2 = 9\%$			0.5 0.7 1	1.5 2
Test for overall effect: $Z = 0.94$ (P = 0.35)						Fav	ours lower SFA	Favours higher SFA
Test for subgroup different	nces: Not app	olicable						



Analysis 2.3. Comparison 2: SFA reduction vs usual diet secondary health events, Outcome 3: MI, SA aim to reduce SFA

	Lower	Lower SFA		Higher SFA		Risk Ratio	Risk R	atio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Randon	n, 95% CI
DART 1989	132	1018	144	1015	22.9%	0.91 [0.73 , 1.14]		
Houtsmuller 1979	0	51	6	51	0.2%	0.08 [0.00 , 1.33]		
MRC 1968	39	199	40	194	9.5%	0.95 [0.64 , 1.41]		
Moy 2001	2	117	1	118	0.3%	2.02 [0.19 , 21.94]		<u> </u>
Oslo Diet-Heart 1966	34	206	54	206	9.9%	0.63 [0.43 , 0.92]		
Rose corn oil 1965	9	28	3	13	1.3%	1.39 [0.45 , 4.31]		
Rose olive 1965	7	26	2	13	0.8%	1.75 [0.42 , 7.27]		<u>. </u>
STARS 1992	1	27	2	28	0.3%	0.52 [0.05 , 5.39]		
Veterans Admin 1969	54	424	71	422	12.9%	0.76 [0.55, 1.05]		
WHI 2006	435	19541	671	29294	41.8%	0.97 [0.86 , 1.09]	•	
Total (95% CI)		21637		31354	100.0%	0.89 [0.78 , 1.02]		
Total events:	713		994				▼	
Heterogeneity: $Tau^2 = 0.0$	01; $Chi^2 = 10$.83, df = 9	(P = 0.29);	$I^2 = 17\%$			0.02 0.1 1	10 50
Test for overall effect: Z	= 1.72 (P = 0)	.09)				F	Favours lower SFA	Favours higher SFA

Test for subgroup differences: Not applicable

Analysis 2.4. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 4: MI, SA statistically significant SFA reduction

	Lower SFA		Higher SFA		Risk Ratio		Risk Ratio	
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI	
DART 1989	132	1018	144	1015	20.5%	0.91 [0.73 , 1.14]	l •	
Ley 2004	4	88	3	88	0.5%	1.33 [0.31 , 5.78]	I	
Moy 2001	2	117	1	118	0.2%	2.02 [0.19, 21.94]	I <u> </u>	
STARS 1992	1	27	2	28	0.2%	0.52 [0.05 , 5.39]	I	
Veterans Admin 1969	54	424	71	422	9.2%	0.76 [0.55 , 1.05]	I -	
WHI 2006	435	19541	671	29294	69.5%	0.97 [0.86 , 1.09]	• •	
Total (95% CI)		21215		30965	100.0%	0.94 [0.85 , 1.04		
Total events:	628		892					
Heterogeneity: $Tau^2 = 0.00$); Chi ² = 2.9	1, df = 5 (P = 0.71; I	$^{2} = 0\%$			0.02 0.1 1 10 50	
Test for overall effect: Z =	1.23 (P = 0	.22)]	Favours lower SFA Favours higher SI	FA



Analysis 2.5. Comparison 2: SFA reduction vs usual diet secondary health events, Outcome 5: MI, SA by TC reduction

	Lower	Lower SFA		Higher SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
DART 1989	132	1018	144	1015	23.3%	0.91 [0.73 , 1.14]	•
Houtsmuller 1979	0	51	6	51	0.2%	0.08 [0.00 , 1.33]	←
MRC 1968	39	199	40	194	9.9%	0.95 [0.64 , 1.41]	
Moy 2001	2	117	1	118	0.3%	2.02 [0.19, 21.94]	
Oslo Diet-Heart 1966	34	206	54	206	10.3%	0.63 [0.43, 0.92]	
Rose corn oil 1965	9	28	3	13	1.4%	1.39 [0.45 , 4.31]	_ _
STARS 1992	1	27	2	28	0.3%	0.52 [0.05 , 5.39]	•
Veterans Admin 1969	54	424	71	422	13.3%	0.76 [0.55 , 1.05]	-
WHI 2006	435	19541	671	29294	40.9%	0.97 [0.86 , 1.09]	•
Total (95% CI)		21611		31341	100.0%	0.88 [0.77 , 1.01]	
Total events:	706		992				*
Heterogeneity: $Tau^2 = 0.0$	1; Chi ² = 10.	04, $df = 8$	(P = 0.26);	$I^2 = 20\%$			0.02 0.1 1 10 50
Test for overall effect: Z =	= 1.79 (P = 0)	.07)		Fa	avours lower SFA Favours higher SFA		
Test for subgroup differen	ices: Not app	olicable					

Analysis 2.6. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 6: MI, SA excluding WHI

	Lower	SFA	Higher SFA		Risk Ratio		Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
DART 1989	132	1018	144	1015	44.9%	0.91 [0.73 , 1.14]	
Houtsmuller 1979	0	51	6	51	0.3%	0.08 [0.00 , 1.33]	←
Ley 2004	4	88	3	88	1.0%	1.33 [0.31 , 5.78]	
MRC 1968	39	199	40	194	14.3%	0.95 [0.64 , 1.41]	
Moy 2001	2	117	1	118	0.4%	2.02 [0.19, 21.94]	
Oslo Diet-Heart 1966	34	206	54	206	15.1%	0.63 [0.43, 0.92]	
Rose corn oil 1965	9	28	3	13	1.8%	1.39 [0.45 , 4.31]	_
Rose olive 1965	7	26	2	13	1.1%	1.75 [0.42 , 7.27]	
STARS 1992	1	27	2	28	0.4%	0.52 [0.05 , 5.39]	
Veterans Admin 1969	54	424	71	422	20.7%	0.76 [0.55 , 1.05]	-
Total (95% CI)		2184		2148	100.0%	0.85 [0.73 , 0.98]	
Total events:	282		326				▼
Heterogeneity: $Tau^2 = 0.0$	00; Chi ² = 9.0	7, df = 9	P = 0.43; I		0.02 0.1 1 10 50		
Test for overall effect: Z	= 2.18 (P = 0)	.03)		Fa	avours lower SFA Favours higher SF		



Analysis 2.7. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 7: MI, SA Mantel-Haenszel fixed-effect

	Lower	SFA	Higher SFA		Risk Ratio		Risk Ra	tio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Fixed, 95% CI	M-H, Fixed, 9	95% CI
DART 1989	132	1018	144	1015	16.7%	0.91 [0.73 , 1.14]		
Houtsmuller 1979	0	51	6	51	0.8%	0.08 [0.00 , 1.33]	←	
Ley 2004	4	88	3	88	0.3%	1.33 [0.31 , 5.78]		
MRC 1968	39	199	40	194	4.7%	0.95 [0.64 , 1.41]	_	
Moy 2001	2	117	1	118	0.1%	2.02 [0.19 , 21.94]		
Oslo Diet-Heart 1966	34	206	54	206	6.2%	0.63 [0.43 , 0.92]		
Rose corn oil 1965	9	28	3	13	0.5%	1.39 [0.45 , 4.31]	_	
Rose olive 1965	7	26	2	13	0.3%	1.75 [0.42 , 7.27]		
STARS 1992	1	27	2	28	0.2%	0.52 [0.05 , 5.39]		
Veterans Admin 1969	54	424	71	422	8.2%	0.76 [0.55 , 1.05]		
WHI 2006	435	19541	671	29294	62.0%	0.97 [0.86 , 1.09]	•	
Total (95% CI)		21725		31442	100.0%	0.92 [0.84 , 1.01]		
Total events:	717		997					
Heterogeneity: $Chi^2 = 11.0$	07, df = 10 (1)	P = 0.35;	$I^2 = 10\%$				0.02 0.1 1	10 50
Test for overall effect: Z =	1.73 (P = 0	.08)				F	Favours lower SFA	Favours higher SFA

Test for subgroup differences: Not applicable

Analysis 2.8. Comparison 2: SFA reduction vs usual diet secondary health events, Outcome 8: MI, SA Peto fixed-effect

	Lower	Lower SFA		Higher SFA		Peto Odds Ratio	Peto Odds	Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	Peto, Fixed, 95% CI	Peto, Fixed,	95% CI
DART 1989	132	1018	144	1015	15.4%	0.90 [0.70 , 1.16]] _	
Houtsmuller 1979	0	51	6	51	0.4%	0.12 [0.02 , 0.63]]	
Ley 2004	4	88	3	88	0.4%	1.34 [0.30 , 6.07]]	
MRC 1968	39	199	40	194	4.1%	0.94 [0.57 , 1.54]] _	
Moy 2001	2	117	1	118	0.2%	1.98 [0.20 , 19.18]	ı <u> </u>	•
Oslo Diet-Heart 1966	34	206	54	206	4.5%	0.56 [0.35 , 0.90]]	
Rose corn oil 1965	9	28	3	13	0.5%	1.53 [0.37 , 6.39]]	
Rose olive 1965	7	26	2	13	0.4%	1.88 [0.40 , 8.96]	ı —	
STARS 1992	1	27	2	28	0.2%	0.52 [0.05 , 5.22]	ı	
Veterans Admin 1969	54	424	71	422	6.9%	0.72 [0.49 , 1.06]] _	
WHI 2006	435	19541	671	29294	67.0%	0.97 [0.86 , 1.10]] 📫	
Total (95% CI)		21725		31442	100.0%	0.92 [0.83 , 1.01]	1	
Total events:	717		997				1	
Heterogeneity: $Chi^2 = 14$.	59, df = $10(1)$	P = 0.15);	$I^2 = 31\%$				0.02 0.1 1	10 50
Test for overall effect: Z =	= 1.73 (P = 0)	.08)]	Favours lower SFA	Favours higher SFA

Analysis 2.9. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 9: MI, subgroup by any substitution

	Lower	SFA	Higher	·SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
2.9.1 replaced by PUFA							
DART 1989	132	1018	144	1015	34.0%	0.91 [0.73 , 1.14]	· _
Houtsmuller 1979	0	51	6	51	0.5%	0.08 [0.00 , 1.33]	
MRC 1968	39	199	40	194	18.3%	0.95 [0.64 , 1.41]	·
Oslo Diet-Heart 1966	34	206	54	206	19.0%	0.63 [0.43, 0.92]	
Rose corn oil 1965	9	28	5	26	4.3%	1.67 [0.64 , 4.34]	
STARS 1992	1	27	2	28	0.8%	0.52 [0.05 , 5.39]	
Veterans Admin 1969	54	424	71	422	23.1%	0.76 [0.55, 1.05]	
Subtotal (95% CI)		1953		1942	100.0%	0.83 [0.67, 1.02]	
Total events:	269		322			, , , , , , , , , , , , , , , , , , , ,	▲
Heterogeneity: $Tau^2 = 0.02$	2: $Chi^2 = 8.4$	0. df = 6	P = 0.21);]	$[^2 = 29\%]$			
Test for overall effect: $Z =$	1.79 (P = 0	.07)					
2.9.2 replaced by MUFA							
Rose olive 1965	7	26	5	26	100.0%	1.40 [0.51 , 3.85]	
Subtotal (95% CI)		26		26	100.0%	1.40 [0.51 , 3.85]	
Total events:	7		5				
Heterogeneity: Not applica	ıble						
Test for overall effect: $Z =$	0.65 (P = 0	.51)					
2.9.3 replace by CHO							
DART 1989	132	1018	144	1015	22.6%	0.91 [0.73 , 1.14]	
Ley 2004	4	88	3	88	0.5%	1.33 [0.31 , 5.78]	
STARS 1992	1	27	2	28	0.2%	0.52 [0.05, 5.39]	
WHI 2006	435	19541	671	29294	76.7%	0.97 [0.86, 1.09]	
Subtotal (95% CI)		20674		30425	100.0%	0.96 [0.86 , 1.06]	□
Total events:	572		820				
Heterogeneity: $Tau^2 = 0.00$); $Chi^2 = 0.6$	9, df = 3 (P = 0.88;	$[^2 = 0\%]$			
Test for overall effect: $Z =$	0.79 (P = 0	.43)					
2.9.4 replaced by protein							
DART 1989	132	1018	144	1015	22.7%	0.91 [0.73 , 1.14]	
Ley 2004	4	88	3	88	0.5%	1.33 [0.31 , 5.78]	· · · · · · · · · · · · · · · · · · ·
WHI 2006	435	19541	671	29294	76.8%	0.97 [0.86 , 1.09]	
Subtotal (95% CI)		20647		30397	100.0%	0.96 [0.86 , 1.07]	⊢
Total events:	571		818				
Heterogeneity: $Tau^2 = 0.00$); $Chi^2 = 0.4$	-3, df = 2 (P = 0.81;	$[^2 = 0\%]$			
Test for overall effect: Z =	0.77 (P = 0	.44)					
2.9.5 replacement unclear	r						
Moy 2001	2	117	1	118	100.0%	2.02 [0.19 , 21.94]	
Subtotal (95% CI)		117		118	100.0%	2.02 [0.19 , 21.94]	
Total events:	2		1				
Heterogeneity: Not applica	ıble						
Test for overall effect: $Z =$	0.58 (P = 0	.56)					
Test for subgroup difference	ces: $Chi^2 = 2$	2.70, df =	4 (P = 0.61), $I^2 = 0\%$		F	0.02 0.1 1 10 50 Favours lower SFA Favours higher SF

Reduction in saturated fat intake for cardiovascular disease (Review)

Lower SFA **Higher SFA Risk Ratio Risk Ratio** Total Total M-H, Random, 95% CI M-H, Random, 95% CI Study or Subgroup **Events Events** Weight 2.10.1 replaced by PUFA **DART 1989** 132 1018 144 1015 32.4% 0.91 [0.73, 1.14] Houtsmuller 1979 0 51 6 51 0.6% 0.08 [0.00, 1.33] **MRC 1968** 39 199 40 194 19.0% 0.95 [0.64, 1.41] 206 Oslo Diet-Heart 1966 34 206 54 19.7% 0.63 [0.43, 0.92] 9 Rose corn oil 1965 5 26 28 4.9% 1.67 [0.64, 4.34] Veterans Admin 1969 54 424 71 422 23.4% 0.76 [0.55, 1.05] Subtotal (95% CI) 1926 1914 100.0% 0.83 [0.67, 1.04] Total events: 268 320 Heterogeneity: Tau² = 0.03; Chi² = 8.23, df = 5 (P = 0.14); I² = 39% Test for overall effect: Z = 1.63 (P = 0.10) 2.10.2 replaced by MUFA 1.40 [0.51, 3.85] Rose olive 1965 7 26 5 100.0% 26 Subtotal (95% CI) 26 26 100.0% 1.40 [0.51, 3.85] 7 Total events: 5 Heterogeneity: Not applicable Test for overall effect: Z = 0.65 (P = 0.51) 2.10.3 replace by CHO Ley 2004 4 88 3 88 0.7% 1.33 [0.31, 5.78] **STARS 1992** 27 2 1 28 0.3% 0.52 [0.05, 5.39] WHI 2006 435 19541 671 29294 99.1% 0.97 [0.86, 1.09] Subtotal (95% CI) 19656 29410 100.0% 0.97 [0.86, 1.09] 440 Total events: 676 Heterogeneity: Tau² = 0.00; Chi² = 0.45, df = 2 (P = 0.80); I² = 0% Test for overall effect: Z = 0.46 (P = 0.64) 2.10.4 replaced by protein Subtotal (95% CI) 0 Not estimable 0 Total events: 0 0 Heterogeneity: Not applicable Test for overall effect: Not applicable 2.10.5 replacement unclear Moy 2001 2 117 1 118 100.0% 2.02 [0.19, 21.94] Subtotal (95% CI) 117 118 100.0% 2.02 [0.19, 21.94] Total events: 2 1 Heterogeneity: Not applicable Test for overall effect: Z = 0.58 (P = 0.56) Test for subgroup differences: $Chi^2 = 2.48$, df = 3 (P = 0.48), $I^2 = 0\%$ 0.02 0.1 10 50 Favours lower SFA Favours higher SFA

Analysis 2.10. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 10: MI, subgroup by main substitution

Reduction in saturated fat intake for cardiovascular disease (Review) Copyright © 2020 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

Analysis 2.11. Comparison 2: SFA reduction vs usual diet secondary health events, Outcome 11: MI, subgroup by duration Lower SFA Higher SFA Risk Ratio Risk Ratio pup Events Total Events Total Weight M-H, Random, 95% CI M-H, Random, 95% C

Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Randor	n, 95% CI
2.11.1 up to 24mo								
DART 1989	132	1018	144	1015	21.9%	0.91 [0.73 , 1.14]	-	
Moy 2001	2	117	1	118	0.2%	2.02 [0.19 , 21.94]		
Rose corn oil 1965	9	28	3	13	1.1%	1.39 [0.45 , 4.31]	,	<u> </u>
Rose olive 1965	7	26	2	13	0.7%	1.75 [0.42 , 7.27]		
Subtotal (95% CI)		1189		1159	23.9%	0.95 [0.77 , 1.17]	•	
Total events:	150		150				Ť	
Heterogeneity: $Tau^2 = 0.0$	00; $Chi^2 = 1.6$	5, $df = 3$ (P = 0.65; I	$[^2 = 0\%]$				
Test for overall effect: Z	= 0.50 (P = 0)	.62)						
2.11.2 >24 to 48mo								
MRC 1968	39	199	40	194	8.2%	0.95 [0.64 , 1.41]		
STARS 1992	1	27	2	28	0.3%	0.52 [0.05, 5.39]		
Veterans Admin 1969	54	424	71	422	11.4%	0.76 [0.55, 1.05]		
Subtotal (95% CI)		650		644	19.9%	0.83 [0.64 , 1.06]		
Total events:	94		113				•	
Heterogeneity: $Tau^2 = 0.0$	00; Chi ² = 0.9	1, $df = 2$ (P = 0.63; I	$[^2 = 0\%]$				
Test for overall effect: Z	= 1.49 (P = 0)	.13)						
2.11.3 >48mo								
Oslo Diet-Heart 1966	34	206	54	206	8.6%	0.63 [0.43, 0.92]		
WHI 2006	435	19541	671	29294	46.7%	0.97 [0.86 , 1.09]		
Subtotal (95% CI)		19747		29500	55.4%	0.81 [0.54 , 1.24]		
Total events:	469		725					
Heterogeneity: $Tau^2 = 0.0$	07; Chi ² = 4.5	0, df = 1 (P = 0.03; I	[² = 78%				
Test for overall effect: Z	= 0.96 (P = 0)	.34)						
2.11.4 unclear								
Houtsmuller 1979	0	51	6	51	0.2%	0.08 [0.00 , 1.33]	←	
Ley 2004	4	88	3	88	0.6%	1.33 [0.31 , 5.78]		
Subtotal (95% CI)		139		139	0.8%	0.41 [0.02, 7.73]		
Total events:	4		9					
Heterogeneity: $Tau^2 = 3.3$	31; Chi ² = 3.4	7, df = 1 (P = 0.06; I	$1^2 = 71\%$				
Test for overall effect: Z	= 0.60 (P = 0)	.55)						
Total (95% CI)		21725		31442	100.0%	0.90 [0.80 , 1.01]		
Total events:	717		997				v	
Heterogeneity: $Tau^2 = 0.0$	00; Chi ² = 11.	07, $df = 1$	0 (P = 0.35)); $I^2 = 10\%$	b		0.02 0.1 1	10 50
Test for overall effect: Z	= 1.72 (P = 0)	.09)				F	Favours lower SFA	Favours higher SF.
Test for subgroup differe	nces: Chi ² =	1.10, df =	3 (P = 0.78)), $I^2 = 0\%$				-

Analysis 2.12. Comparison 2: SFA reduction vs usual diet secondary health events, Outcome 12: MI, subgroup by baseline SFA

	Lower	SFA	Higher	Higher SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
2.12.1 up to 12%E SFA b	oaseline						
Subtotal (95% CI)		0		0		Not estimable	
Total events:	0		0				
Heterogeneity: Not applica	able						
Test for overall effect: Not	t applicable						
2.12.2 >12 to 15%E SFA	baseline						
DART 1989	132	1018	144	1015	21.9%	0.91 [0.73 , 1.14]	-
Ley 2004	4	88	3	88	0.6%	1.33 [0.31 , 5.78]	-
Moy 2001	2	117	1	118	0.2%	2.02 [0.19 , 21.94]	
WHI 2006	435	19541	671	29294	46.7%	0.97 [0.86 , 1.09]	•
Subtotal (95% CI)		20764		30515	69.5%	0.96 [0.87 , 1.07]	•
Total events:	573		819				
Heterogeneity: Tau ² = 0.00 Test for overall effect: Z =	0; $Chi^2 = 0.8$ = 0.74 (P = 0	30, df = 3 (.46)	(P = 0.85);	$1^2 = 0\%$			
2.12.3 >15 to 18% E SFA	baseline						
STARS 1992	1	27	2	28	0.3%	0.52 [0.05 , 5.39]	•
Subtotal (95% CI)		27		28	0.3%	0.52 [0.05 , 5.39]	
Total events:	1		2				
Heterogeneity: Not applica	able						
Test for overall effect: Z =	= 0.55 (P = 0)	.58)					
2.12.4 >18%E SFA basel	ine						
Veterans Admin 1969	54	424	71	422	11.4%	0.76 [0.55, 1.05]	
Subtotal (95% CI)		424		422	11.4%	0.76 [0.55 , 1.05]	
Total events:	54		71				•
Heterogeneity: Not applica	able						
Test for overall effect: Z =	= 1.67 (P = 0)	.10)					
2.12.5 unclear							
Houtsmuller 1979	0	51	6	51	0.2%	0.08 [0.00 , 1.33]	←
MRC 1968	39	199	40	194	8.2%	0.95 [0.64 , 1.41]	-
Oslo Diet-Heart 1966	34	206	54	206	8.6%	0.63 [0.43 , 0.92]	
Rose corn oil 1965	9	28	3	13	1.1%	1.39 [0.45 , 4.31]	-
Rose olive 1965	7	26	2	13	0.7%	1.75 [0.42 , 7.27]	
Subtotal (95% CI)		510		477	18.8%	0.84 [0.54 , 1.30]	
Total events:	89		105				
Heterogeneity: $Tau^2 = 0.09$ Test for overall effect: Z =	9; Chi ² = 6.9 : 0.78 (P = 0	98, df = 4 (.44)	(P = 0.14);	l ² = 43%			
Total (95% CI)		21725		31442	100.0%	0.90 [0.80 , 1.01]	
Total events:	717		997				۲
Heterogeneity: $Tau^2 = 0.00$	0; Chi ² = 11.	.07, $df = 1$	0 (P = 0.35)); $I^2 = 10\%$, 9	0	0.02 0.1 1 10 50
Test for overall effect: Z =	= 1.72 (P = 0	.09)				Fav	ours lower SFA Favours higher SFA
Test for subgroup differen	ces: Chi ² = 2	2.34, df =	3 (P = 0.50)), $I^2 = 0\%$			2

Reduction in saturated fat intake for cardiovascular disease (Review)

Analysis 2.13. Comparison 2: SFA reduction vs usual diet secondary health events, Outcome 13: MI, subgroup by SFA change

	Lower	SFA	Higher	SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
2.13.1 up to 4%E differe	nce						
DART 1989	132	1018	144	1015	21.9%	0.91 [0.73 , 1.14]	1 🚽
Ley 2004	4	88	3	88	0.6%	1.33 [0.31 , 5.78]	
Moy 2001	2	117	1	118	0.2%	2.02 [0.19, 21.94]]
WHI 2006	435	19541	671	29294	46.7%	0.97 [0.86 , 1.09]] 🗖
Subtotal (95% CI)		20764		30515	69.5%	0.96 [0.87 , 1.07]	1 4
Total events:	573		819				
Heterogeneity: $Tau^2 = 0.00$); $Chi^2 = 0.8$	30, df = 3	(P = 0.85); I	$^{2} = 0\%$			
Test for overall effect: Z =	0.74 (P = 0	.46)					
2.13.2 >4 to 8%E differen	nce						
STARS 1992	1	27	2	28	0.3%	0.52 [0.05 , 5.39]]
Subtotal (95% CI)		27		28	0.3%	0.52 [0.05 , 5.39]	
Total events:	1		2				
Heterogeneity: Not application	able						
Test for overall effect: Z =	0.55 (P = 0	.58)					
2.13.3 >8%E difference							
Veterans Admin 1969	54	424	71	422	11.4%	0.76 [0.55 , 1.05]]
Subtotal (95% CI)		424		422	11.4%	0.76 [0.55 , 1.05]	
Total events:	54		71				•
Heterogeneity: Not applica	able						
Test for overall effect: Z =	1.67 (P = 0	.10)					
2.13.4 unclear							
Houtsmuller 1979	0	51	6	51	0.2%	0.08 [0.00 , 1.33]	
MRC 1968	39	199	40	194	8.2%	0.95 [0.64 , 1.41]]
Oslo Diet-Heart 1966	34	206	54	206	8.6%	0.63 [0.43, 0.92]]
Rose corn oil 1965	9	28	3	13	1.1%	1.39 [0.45 , 4.31]]
Rose olive 1965	7	26	2	13	0.7%	1.75 [0.42 , 7.27]]
Subtotal (95% CI)		510		477	18.8%	0.84 [0.54, 1.30]	1 🔶
Total events:	89		105				•
Heterogeneity: $Tau^2 = 0.09$	$\Theta; Chi^2 = 6.9$	98, df = 4	(P = 0.14); I	$^{2} = 43\%$			
Test for overall effect: Z =	0.78 (P = 0	.44)					
Total (95% CI)		21725		31442	100.0%	0.90 [0.80 , 1.01]	1
Total events:	717		997				1
Heterogeneity: $Tau^2 = 0.00$); Chi ² = 11.	.07, $df = 1$	0 (P = 0.35)); $I^2 = 10\%$	b		0.02 0.1 1 10 50
Test for overall effect: Z =	1.72 (P = 0)	.09)				1	Favours lower SFA Favours higher SF
Test for subgroup differen	ces: Chi ² = 2	2.34, df =	3 (P = 0.50)), $I^2 = 0\%$			

Analysis 2.14. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 14: MI, subgroup by sex

	Lower	SFA	Higher	SFA		Risk Ratio	Risk	Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Rando	om, 95% CI
2.14.1 Men								
DART 1989	132	1018	144	1015	21.9%	0.91 [0.73 , 1.14]] 🚽	
MRC 1968	39	199	40	194	8.2%	0.95 [0.64 , 1.41]] _	_
Oslo Diet-Heart 1966	34	206	54	206	8.6%	0.63 [0.43, 0.92]] _	
Rose corn oil 1965	9	28	3	13	1.1%	1.39 [0.45 , 4.31]]	
Rose olive 1965	7	26	2	13	0.7%	1.75 [0.42 , 7.27]]	
STARS 1992	1	27	2	28	0.3%	0.52 [0.05 , 5.39]]	
Veterans Admin 1969	54	424	71	422	11.4%	0.76 [0.55 , 1.05]] _	
Subtotal (95% CI)		1928		1891	52.2%	0.85 [0.73, 0.98]]	
Total events:	276		316				×.	
Heterogeneity: $Tau^2 = 0.0$	00; $Chi^2 = 5.4$	6, df = 6	P = 0.49; I	$^{2} = 0\%$				
Test for overall effect: Z	= 2.20 (P = 0)	.03)						
2.14.2 Women								
WHI 2006	435	19541	671	29294	46.7%	0.97 [0.86 , 1.09]	1	
Subtotal (95% CI)		19541		29294	46.7%	0.97 [0.86 , 1.09	1	
Total events:	435		671			- / .		
Heterogeneity: Not appli	cable							
Test for overall effect: Z	= 0.47 (P = 0)	.64)						
2.14.3 Mixed, men and	women							
Houtsmuller 1979	0	51	6	51	0.2%	0.08 [0.00 , 1.33]	1	_
Ley 2004	4	88	3	88	0.6%	1.33 [0.31 , 5.78]		
Mov 2001	2	117	1	118	0.2%	2.02 [0.19, 21.94]	1	
Subtotal (95% CI)		256		257	1.1%	0.75 [0.13 , 4.47		
Total events:	6		10			- , .		
Heterogeneity: $Tau^2 = 1.2$	27; Chi ² = 4.0	4, df = 2 (P = 0.13; I	$^{2} = 51\%$				
Test for overall effect: Z	= 0.31 (P = 0	.75)						
Total (95% CI)		21725		31442	100.0%	0.90 [0.80 , 1.01	1	
Total events:	717		997			- / .		
Heterogeneity: $Tau^2 = 0.0$	00; Chi ² = 11.	.07, df = 1	0 (P = 0.35)); $I^2 = 10\%$	5		$0 \frac{1}{02} 0 \frac{1}{1}$	10 50
Test for overall effect: Z	= 1.72 (P = 0)	.09)	. ,]	Favours lower SFA	Favours higher S
Test for subgroup differe	ences: $Chi^2 = 2$	2.10, df = 1	2 (P = 0.35)	$I^2 = 4.6\%$	6			c

Analysis 2.15. Comparison 2: SFA reduction vs usual diet secondary health events, Outcome 15: MI, subgroup by CVD risk

	Lower	SFA	Higher	SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
2.15.1 Low CVD risk							
Veterans Admin 1969	54	424	71	422	11.4%	0.76 [0.55 , 1.05]	
WHI 2006	435	19541	671	29294	46.7%	0.97 [0.86 , 1.09]	•
Subtotal (95% CI)		19965		29716	58.2%	0.90 [0.72 , 1.13]	▲
Total events:	489		742				•
Heterogeneity: Tau ² = 0.0	02; Chi ² = 1.9	98, df = 1 (P = 0.16; I	$^{2} = 49\%$			
Test for overall effect: Z	= 0.91 (P = 0)	.36)					
2.15.2 Moderate CVD r	isk						
Houtsmuller 1979	0	51	6	51	0.2%	0.08 [0.00 , 1.33]	←
Ley 2004	4	88	3	88	0.6%	1.33 [0.31 , 5.78]	
Moy 2001	2	117	1	118	0.2%	2.02 [0.19, 21.94]	
Subtotal (95% CI)		256		257	1.1%	0.75 [0.13 , 4.47]	
Total events:	6		10				
Heterogeneity: $Tau^2 = 1.2$	27; $Chi^2 = 4.0$	4, df = 2 (P = 0.13; I	$^{2} = 51\%$			
Test for overall effect: Z	= 0.31 (P = 0)	.75)					
2.15.3 Existing CVD dis	sease						
DART 1989	132	1018	144	1015	21.9%	0.91 [0.73, 1.14]	+
MRC 1968	39	199	40	194	8.2%	0.95 [0.64 , 1.41]	
Oslo Diet-Heart 1966	34	206	54	206	8.6%	0.63 [0.43, 0.92]	
Rose corn oil 1965	9	28	3	13	1.1%	1.39 [0.45 , 4.31]	_
Rose olive 1965	7	26	2	13	0.7%	1.75 [0.42 , 7.27]	
STARS 1992	1	27	2	28	0.3%	0.52 [0.05 , 5.39]	
Subtotal (95% CI)		1504		1469	40.8%	0.87 [0.74 , 1.03]	
Total events:	222		245				ľ
Heterogeneity: Tau ² = 0.0	00; $Chi^2 = 4.9$	0, df = 5 (P = 0.43; I	$^{2} = 0\%$			
Test for overall effect: Z	= 1.62 (P = 0)	.10)					
Total (95% CI)		21725		31442	100.0%	0.90 [0.80 , 1.01]	
Total events:	717		997				Ĭ
Heterogeneity: Tau ² = 0.0	00; $Chi^2 = 11$.07, $df = 1$	0 (P = 0.35)); $I^2 = 10\%$			0.02 0.1 1 10 50
Test for overall effect: Z	= 1.72 (P = 0)	.09)				F	Favours lower SFA Favours higher SF
Test for subgroup differe	nces: Chi ² = 0	0.09, df = 2	2 (P = 0.96)), $I^2 = 0\%$			



Analysis 2.16. Comparison 2: SFA reduction vs usual diet secondary health events, Outcome 16: MI, subgroup by TC reduction

	Lower	SFA	Higher	SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
2.16.1 serum chol reduce	ed by at leas	t 0.2mmo	I/L				
DART 1989	132	1018	144	1015	41.0%	0.91 [0.73 , 1.14]	_
Houtsmuller 1979	0	51	6	51	0.4%	0.08 [0.00 , 1.33]	← ⊷ ↓
MRC 1968	39	199	40	194	16.2%	0.95 [0.64 , 1.41]	-
Moy 2001	2	117	1	118	0.5%	2.02 [0.19, 21.94]	.
Oslo Diet-Heart 1966	34	206	54	206	17.0%	0.63 [0.43, 0.92]	
Rose corn oil 1965	9	28	3	13	2.2%	1.39 [0.45 , 4.31]	
STARS 1992	1	27	2	28	0.5%	0.52 [0.05 , 5.39]	
Veterans Admin 1969	54	424	71	422	22.3%	0.76 [0.55 , 1.05]	
Subtotal (95% CI)		2070		2047	100.0%	0.83 [0.70 , 0.98]	
Total events:	271		321				•
Heterogeneity: $Tau^2 = 0.0$	1; Chi ² = 7.6	68, df = 7 (P = 0.36; I	$^{2} = 9\%$			
Test for overall effect: Z =	= 2.17 (P = 0)	.03)					
2.16.2 serum chol reduce	ed by <0.2m	mol/L					
Ley 2004	4	88	3	88	0.7%	1.33 [0.31 , 5.78]	
Rose olive 1965	7	26	2	13	0.7%	1.75 [0.42, 7.27]	
WHI 2006	435	19541	671	29294	98.7%	0.97 [0.86, 1.09]	•
Subtotal (95% CI)		19655		29395	100.0%	0.98 [0.87 , 1.10]	
Total events:	446		676				
Heterogeneity: $Tau^2 = 0.0$	0; Chi ² = 0.8	2, df = 2 (P = 0.66; I	$^{2} = 0\%$			
Test for overall effect: Z =	= 0.37 (P = 0)	.71)					
2.16.3 serum chol reduct	tion unclear						
Subtotal (95% CI)		0		0		Not estimable	
Total events:	0		0				
Heterogeneity: Not applic	able						
Test for overall effect: No	t applicable						
Test for subgroup differer	nces: $Chi^2 = 2$	2.45, df =	1 (P = 0.12)), I ² = 59.1	%	F	$0.02 0.1 \qquad i \qquad 10 50$ avours lower SFA Favours higher SF.

Analysis 2.17. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 17: MI, subgroup decade of publication

Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
2 17 1 10/0-							
2.17.1 1900s							
MRC 1968	39	199	40	194	8.2%	0.95 [0.64 , 1.41]	_
Oslo Diet-Heart 1966	34	206	54	206	8.6%	0.63 [0.43 , 0.92]	
Rose corn oil 1965	9	28	3	13	1.1%	1.39 [0.45 , 4.31]	
Rose olive 1965	7	26	2	13	0.7%	1.75 [0.42, 7.27]	
Veterans Admin 1969	54	424	71	422	11.4%	0.76 [0.55, 1.05]	
Subtotal (95% CI)		883		848	30.0%	0.80 [0.64 , 1.00]	
Total events:	143		170			- / -	•
Heterogeneity: $Tau^2 = 0.0$	1; Chi ² = 4.4	-3, df = 4 (P = 0.35; I	$^{2} = 10\%$			
Test for overall effect: Z =	= 1.98 (P = 0)	.05)					
2.17.2 1970s							
Houtsmuller 1979	0	51	6	51	0.2%	0.08 [0.00 , 1.33]	
Subtotal (95% CI)		51		51	0.2%	0.08 [0.00 , 1.33]	
Total events:	0		6				
Heterogeneity: Not applica	able						
Test for overall effect: Z =	= 1.76 (P = 0)	.08)					
2.17.3 1980s							
DART 1989	132	1018	144	1015	21.9%	0 91 [0 73 1 14]	
Subtotal (95% CI)	152	1018	111	1015	21.9%	0.91 [0.73 , 1.14]	I
Total events:	132	1010	144	1010			Ţ
Heterogeneity: Not applic:	able		111				
Test for overall effect: Z =	= 0.80 (P = 0)	.42)					
2 17 4 1990s							
STARS 1992	1	27	2	28	0.3%	0.52 [0.05 5.39]	
Subtotal (95% CI)	1	27	2	20	0.3%	0.52 [0.05, 5.39]	
Total events:	1	2,	2	20	0.0 /0	0.02 [0.00 ; 0.07]	
Heterogeneity: Not applic	ahle		2				
Test for overall effect: Z =	= 0.55 (P = 0)	.58)					
2.17.5 2000s							
Lev 2004	4	88	3	88	0.6%	1.33 [0.31 , 5.78]	
Mov 2001	2	117	1	118	0.2%	2.02 [0.19 . 21.94]	
WHI 2006	435	19541	671	29294	46.7%	0.97 [0.86 . 1.09]	
Subtotal (95% CI)		19746		29500	47.6%	0.98 [0.87 , 1.10]	T
Total events:	441		675				Ţ
Heterogeneity: $Tau^2 = 0.00$	0; Chi ² = 0.5	3, df = 2 (P = 0.77: I	$^{2} = 0\%$			
Test for overall effect: Z =	= 0.41 (P = 0)	.68)	,,-				
Total (95% CI)		21725		31442	100.0%	0.90 [0.80 . 1.01]	
Total events:	717		997				•
Heterogeneity: $Tau^2 = 0.00$	0; $Chi^2 = 11$	07. $df = 1$	0 (P = 0.35)); $I^2 = 10\%$,		$\frac{1}{12}$
Test for overall effect: $Z =$	= 1.72 (P = 0)	.09)		., - 1070		U.C Favor	urs lower SFA Favours higher Sl
Test for subgroup differen	ces: $Chi^2 = 4$	5.62. df = 4	4 (P = 0.23)	$I^2 = 28.8$	%	14/01	

Reduction in saturated fat intake for cardiovascular disease (Review)



Analysis 2.18. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 18: NON-FATAL MYOCARDIAL INFARCTION

	Lower	SFA	Higher	SFA		Risk Ratio	Risk Rati	0
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random,	95% CI
DART 1989	35	1018	47	1015	6.0%	0.74 [0.48 , 1.14]	
MRC 1968	25	199	25	194	4.1%	0.97 [0.58 , 1.64	.]	
Moy 2001	2	117	1	118	0.2%	2.02 [0.19 , 21.94	.]	
Oslo Diet-Heart 1966	24	206	31	206	4.5%	0.77 [0.47 , 1.27	1	
Rose corn oil 1965	7	28	3	13	0.8%	1.08 [0.33 , 3.53]	_
Rose olive 1965	6	26	2	13	0.5%	1.50 [0.35 , 6.43]	
Veterans Admin 1969	13	424	21	422	2.4%	0.62 [0.31, 1.21]	
WHI 2006 (1)	459	19541	684	29294	81.4%	1.01 [0.90 , 1.13	1	
Total (95% CI)		21559		31275	100.0%	0.97 [0.87 , 1.07	n 🖡	
Total events:	571		814					
Heterogeneity: $Tau^2 = 0.0$	00; Chi ² = 5.1	1, $df = 7$ (P = 0.65; I	$^{2} = 0\%$			0 02 0 1 1	10 50
Test for overall effect: $Z = 0.62$ (P = 0.54)					Favours lower SFA H	Favours higher SFA		
Test for subgroup differe	nces: Not app	licable						

Footnotes

(1) Non-fatal MI during trial, Prentice 2017

Analysis 2.19. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 19: Non-fatal MI, SA by low summary risk of bias

	Lower SFA		Higher SFA		Risk Ratio		Risk Ra	tio
Study or Subgroup	Events	Total	Events	nts Total	Weight	M-H, Random, 95% CI	M-H, Random	, 95% CI
Veterans Admin 1969	13	424	21	422	25.8%	0.62 [0.31 , 1.21]	
WHI 2006 (1)	459	19541	684	29294	74.2%	1.01 [0.90 , 1.13	5] –	
Total (95% CI)		19965		29716	100.0%	0.89 [0.58 , 1.35	5]	
Total events:	472		705					
Heterogeneity: Tau ² = 0.06	; Chi ² = 1.9	5, $df = 1$ (P = 0.16; I	$^{2} = 49\%$			0.02 0.1 1	10 50
Test for overall effect: Z =	0.56 (P = 0	.57)					Favours lower SFA	Favours higher SFA
Test for subgroup difference	es: Not app	olicable						

Footnotes

(1) Non-fatal MI during trial, Prentice 2017



Analysis 2.20. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 20: Non-fatal MI, SA aim to reduce SFA

	Lower	Lower SFA		Higher SFA		Risk Ratio	Risk R	atio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Rando	m, 95% CI
DART 1989	35	1018	47	1015	6.0%	0.74 [0.48 , 1.14	l]	
MRC 1968	25	199	25	194	4.1%	0.97 [0.58 , 1.64	I) 🚽	_
Moy 2001	2	117	1	118	0.2%	2.02 [0.19 , 21.94	l]	•
Oslo Diet-Heart 1966	24	206	31	206	4.5%	0.77 [0.47 , 1.27	′]	
Rose corn oil 1965	7	28	3	13	0.8%	1.08 [0.33 , 3.53	3]	
Rose olive 1965	6	26	2	13	0.5%	1.50 [0.35, 6.43	3]	.
Veterans Admin 1969	13	424	21	422	2.4%	0.62 [0.31, 1.21]	
WHI 2006 (1)	459	19541	684	29294	81.4%	1.01 [0.90 , 1.13	3]	
Total (95% CI)		21559		31275	100.0%	0.97 [0.87 , 1.07	']	
Total events:	571		814					
Heterogeneity: $Tau^2 = 0.0$	0; Chi ² = 5.1	1, df = 7 (P = 0.65; I	$^{2} = 0\%$			0.02 0.1 1	10 50
Test for overall effect: $Z = 0.62$ (P = 0.54)		.54)					Favours lower SFA	Favours higher SFA

Test for subgroup differences: Not applicable

Footnotes

(1) Non-fatal MI during trial, Prentice 2017

Analysis 2.21. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 21: Non-fatal MI, SA statistically significant SFA reduction

	Lower	SFA	Higher	SFA		Risk Ratio	Risk	Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Rando	om, 95% CI
DART 1989	35	1018	47	1015	21.4%	0.74 [0.48 , 1.14] _	_
Moy 2001	2	117	1	118	0.9%	2.02 [0.19, 21.94]	
Veterans Admin 1969	13	424	21	422	10.1%	0.62 [0.31, 1.21]	_
WHI 2006 (1)	459	19541	684	29294	67.5%	1.01 [0.90 , 1.13]	•
Total (95% CI)		21100		30849	100.0%	0.90 [0.72 , 1.14	1	•
Total events:	509		753					
Heterogeneity: $Tau^2 = 0.0$	2; Chi ² = 3.9	4, df = 3 (P = 0.27; I	$l^2 = 24\%$			0.02 0.1	10 50
Test for overall effect: Z =	= 0.87 (P = 0)	.39)					Favours lower SFA	Favours higher SFA
Test for subgroup differer	nces: Not app	olicable						

Footnotes

(1) Non-fatal MI during trial, Prentice 2017



Analysis 2.22. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 22: Non-fatal MI, SA by TC reduction

	Lower SFA		Higher SFA		Risk Ratio		Risk Ratio	
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random	,95% CI
DART 1989	35	1018	47	1015	6.1%	0.74 [0.48 , 1.14] _	
MRC 1968	25	199	25	194	4.2%	0.97 [0.58, 1.64]] 🗕	
Moy 2001	2	117	1	118	0.2%	2.02 [0.19, 21.94]]	•
Oslo Diet-Heart 1966	24	206	31	206	4.5%	0.77 [0.47 , 1.27]]	
Rose corn oil 1965	7	28	3	13	0.8%	1.08 [0.33, 3.53]]	
Veterans Admin 1969	13	424	21	422	2.4%	0.62 [0.31, 1.21]]	
WHI 2006 (1)	459	19541	684	29294	81.8%	1.01 [0.90 , 1.13]	
Total (95% CI)		21533		31262	100.0%	0.97 [0.87 , 1.07	1	
Total events:	565		812					
Heterogeneity: $Tau^2 = 0.00$); Chi ² = 4.7	6, df = 6 (P = 0.57; I	$^{2} = 0\%$			0.02 0.1 1	10 50
Test for overall effect: Z =	0.66 (P = 0	.51)					Favours lower SFA	Favours higher SFA

Test for subgroup differences: Not applicable

Footnotes

(1) Non-fatal MI during trial, Prentice 2017

Analysis 2.23. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 23: Non-fatal MI, SA excluding WHI

	Lower	Lower SFA		Higher SFA		Risk Ratio	Risk Ratio	
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Randon	n, 95% CI
DART 1989	35	1018	47	1015	32.4%	0.74 [0.48 , 1.14]]	
MRC 1968	25	199	25	194	22.3%	0.97 [0.58 , 1.64]] 🚽	
Moy 2001	2	117	1	118	1.0%	2.02 [0.19, 21.94]]	•
Oslo Diet-Heart 1966	24	206	31	206	24.2%	0.77 [0.47 , 1.27]] _	
Rose corn oil 1965	7	28	3	13	4.3%	1.08 [0.33, 3.53]]	
Rose olive 1965	6	26	2	13	2.8%	1.50 [0.35 , 6.43]]	
Veterans Admin 1969	13	424	21	422	13.0%	0.62 [0.31 , 1.21]]	
Total (95% CI)		2018		1981	100.0%	0.81 [0.64 , 1.04	1	
Total events:	112		130				•	
Heterogeneity: $Tau^2 = 0.0$	00; $Chi^2 = 2.7$	9, df = 6 (P = 0.83; I	$1^2 = 0\%$			0.02 0.1 1	10 50
Test for overall effect: Z	= 1.64 (P = 0)	.10)]	Favours lower SFA	Favours higher SFA



Analysis 2.24. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 24: Non-fatal MI, SA Mantel-Haenszel fixed-effect

	Lower SFA		Higher SFA		Risk Ratio		Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Fixed, 95% CI	M-H, Fixed, 95% CI
DART 1989	35	1018	47	1015	6.9%	0.74 [0.48 , 1.14]	-
MRC 1968	25	199	25	194	3.7%	0.97 [0.58 , 1.64]	_
Moy 2001	2	117	1	118	0.1%	2.02 [0.19 , 21.94]	
Oslo Diet-Heart 1966	24	206	31	206	4.6%	0.77 [0.47 , 1.27]	
Rose corn oil 1965	7	28	3	13	0.6%	1.08 [0.33 , 3.53]	
Rose olive 1965	6	26	2	13	0.4%	1.50 [0.35 , 6.43]	_
Veterans Admin 1969	13	424	21	422	3.1%	0.62 [0.31 , 1.21]	_ _
WHI 2006 (1)	459	19541	684	29294	80.5%	1.01 [0.90 , 1.13]	•
Total (95% CI)		21559		31275	100.0%	0.97 [0.87 , 1.08]	
Total events:	571		814				
Heterogeneity: $Chi^2 = 5.1$	1, df = 7 (P = 7)	: 0.65); I ² :	= 0%				0.02 0.1 1 10 50
Test for overall effect: Z =	= 0.61 (P = 0)	.54)				I	Favours lower SFA Favours higher SFA
Test for subgroup differen	ices: Not app	licable					

Footnotes

(1) Non-fatal MI during trial, Prentice 2017

Analysis 2.25. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 25: Non-fatal MI, SA Peto fixed-effect

	Lower	Lower SFA		Higher SFA		Peto Odds Ratio	Peto Od	lds Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	Peto, Fixed, 95% CI	Peto, Fixe	ed, 95% CI
DART 1989	35	1018	47	1015	6.1%	0.73 [0.47 , 1.14]	
MRC 1968	25	199	25	194	3.4%	0.97 [0.54 , 1.76] _	_
Moy 2001	2	117	1	118	0.2%	1.98 [0.20 , 19.18]	
Oslo Diet-Heart 1966	24	206	31	206	3.7%	0.75 [0.42 , 1.32] _	L
Rose corn oil 1965	7	28	3	13	0.5%	1.11 [0.24 , 5.03]	
Rose olive 1965	6	26	2	13	0.4%	1.58 [0.31 , 8.06]	
Veterans Admin 1969	13	424	21	422	2.5%	0.61 [0.31 , 1.21]	<u> </u>
WHI 2006 (1)	459	19541	684	29294	83.1%	1.01 [0.89 , 1.13]	
Total (95% CI)		21559		31275	100.0%	0.97 [0.87 , 1.08]	
Total events:	571		814					1
Heterogeneity: $Chi^2 = 5.2$	21, df = 7 (P = 7)	= 0.63); I ²	= 0%				0.02 0.1	1 10 50
Test for overall effect: Z	= 0.61 (P = 0)	.54)					Favours lower SFA	Favours higher SF
Test for subgroup differen	nces: Not app	olicable						

Footnotes

(1) Non-fatal MI during trial, Prentice 2017

Analysis 2.26. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 26: Non-fatal MI, subgroup by any substitution

	Lower	SFA	Higher	Higher SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
2.26.1 replaced by PUFA							
DART 1989	35	1018	47	1015	33.2%	0.74 [0.48 , 1.14]	-
MRC 1968	25	199	25	194	22.8%	0.97 [0.58, 1.64]	
Oslo Diet-Heart 1966	24	206	31	206	24.8%	0.77 [0.47 , 1.27]	-
Rose corn oil 1965	7	28	5	26	5.9%	1.30 [0.47 , 3.59]	
Veterans Admin 1969	13	424	21	422	13.3%	0.62 [0.31, 1.21]	_ _
Subtotal (95% CI)		1875		1863	100.0%	0.80 [0.63 , 1.03]	
Total events:	104		129				•
Heterogeneity: $Tau^2 = 0.00$; Chi ² = 2.1	4, df = 4 (P = 0.71; 1	$2^2 = 0\%$			
Test for overall effect: $Z =$	1.72 (P = 0)	.09)					
2.26.2 replaced by MUFA	L						
Rose olive 1965	6	26	5	26	100.0%	1.20 [0.42 , 3.45]	
Subtotal (95% CI)		26		26	100.0%	1.20 [0.42, 3.45]	
Total events:	6		5				
Heterogeneity: Not applica	ble						
Test for overall effect: $Z =$	0.34 (P = 0	.73)					
2.26.3 replace by CHO							
DART 1989	35	1018	47	1015	26.0%	0.74 [0.48, 1.14]	-
WHI 2006 (1)	459	19541	684	29294	74.0%	1.01 [0.90 , 1.13]	-
Subtotal (95% CI)		20559		30309	100.0%	0.93 [0.72 , 1.21]	T
Total events:	494		731			- / -	Y
Heterogeneity: $Tau^2 = 0.02$	2; Chi ² = 1.7	9, $df = 1$ (P = 0.18; I	[² = 44%			
Test for overall effect: $Z =$	0.55 (P = 0	.58)					
2.26.4 replaced by protein	n						
DART 1989	35	1018	47	1015	26.0%	0.74 [0.48 , 1.14]	-
WHI 2006 (1)	459	19541	684	29294	74.0%	1.01 [0.90, 1.13]	-
Subtotal (95% CI)		20559		30309	100.0%	0.93 [0.72 , 1.21]	T
Total events:	494		731				▼
Heterogeneity: $Tau^2 = 0.02$	2; Chi ² = 1.7	9, df = 1 (P = 0.18; I	[² = 44%			
Test for overall effect: $Z =$	0.55 (P = 0	.58)					
2.26.5 replacement unclea	ar						
Moy 2001	2	117	1	118	100.0%	2.02 [0.19 , 21.94]	
Subtotal (95% CI)		117		118	100.0%	2.02 [0.19 , 21.94]	
Total events:	2		1				
Heterogeneity: Not applica	ble						
Test for overall effect: $Z =$	0.58 (P = 0	.56)					
Test for subgroup difference	ces: $Chi^2 = 1$.61, df =	4 (P = 0.81)), $I^2 = 0\%$		_ 0.0	
F						Favo	burs lower SFA Favours higher SF.
rootnotes							

(1) Non-fatal MI during trial, Prentice 2017

Reduction in saturated fat intake for cardiovascular disease (Review) Copyright © 2020 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.



Analysis 2.27. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 27: Non-fatal MI, subgroup by main substitution

	Lower SFA		Higher SFA		Risk Ratio		Risk Ratio	
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI	
2.27.1 replaced by PUFA								
DART 1989	35	1018	47	1015	33.2%	0.74 [0.48, 1.14]		
MRC 1968	25	199	25	194	22.8%	0.97 [0.58 , 1.64]		
Oslo Diet-Heart 1966	24	206	31	206	24.8%	0.77 [0.47 , 1.27]	_ _	
Rose corn oil 1965	7	28	5	26	5.9%	1.30 [0.47 , 3.59]		
Veterans Admin 1969	13	424	21	422	13.3%	0.62 [0.31 , 1.21]	_ _	
Subtotal (95% CI)		1875		1863	100.0%	0.80 [0.63 , 1.03]		
Total events:	104		129				•	
Heterogeneity: $Tau^2 = 0.00$; $Chi^2 = 2.1$	4, df = 4 (P = 0.71; I	$^{2} = 0\%$				
Test for overall effect: $Z =$	1.72 (P = 0.1)	09)						
2.27.2 replaced by MUFA								
Rose olive 1965	6	26	5	26	100.0%	1.20 [0.42 , 3.45]		
Subtotal (95% CI)		26		26	100.0%	1.20 [0.42 , 3.45]		
Total events:	6		5					
Heterogeneity: Not applicat	ble							
Test for overall effect: $Z =$	0.34 (P = 0.34)	73)						
2.27.3 replace by CHO								
WHI 2006 (1)	459	19541	684	29294	100.0%	1.01 [0.90 , 1.13]	•	
Subtotal (95% CI)		19541		29294	100.0%	1.01 [0.90 , 1.13]	•	
Total events:	459		684					
Heterogeneity: Not applical	ble							
Test for overall effect: $Z =$	0.10 (P = 0.10)	92)						
2.27.4 replaced by protein	L							
Subtotal (95% CI)		0		0		Not estimable		
Total events:	0		0					
Heterogeneity: Not applicat	ble							
Test for overall effect: Not	applicable							
2.27.5 replacement unclea	r							
Moy 2001	2	117	1	118	100.0%	2.02 [0.19 , 21.94]		
Subtotal (95% CI)		117		118	100.0%	2.02 [0.19 , 21.94]		
Total events:	2		1					
Heterogeneity: Not applicat	ble							
Test for overall effect: $Z =$	0.58 (P = 0.58)	56)						
Test for subgroup differenc	es: Chi ² = 3	6.08, df = 3	3 (P = 0.38)), $I^2 = 2.5\%$	6	0.0 Favo	01 0.1 1 10 100 urs lower SFA Favours higher SI	

Footnotes

(1) Non-fatal MI during trial, Prentice 2017

Reduction in saturated fat intake for cardiovascular disease (Review) Copyright © 2020 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

	Lower	Lower SFA		Higher SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
2.28.1 up to 24mo							
DART 1989	35	1018	47	1015	6.0%	0.74 [0.48 , 1.14]	_ _
Moy 2001	2	117	1	118	0.2%	2.02 [0.19, 21.94]	_
Rose corn oil 1965	7	28	3	13	0.8%	1.08 [0.33 , 3.53]	
Rose olive 1965	6	26	2	13	0.5%	1.50 [0.35 , 6.43]	
Subtotal (95% CI)		1189		1159	7.6%	0.83 [0.57 , 1.22]	
Total events:	50		53				
Heterogeneity: $Tau^2 = 0.0$	00; Chi ² = 1.6	2, $df = 3$ (P = 0.65; I	$^{2} = 0\%$			
Test for overall effect: Z	= 0.94 (P = 0.00)	.35)					
2.28.2 >24 to 48mo							
MRC 1968	25	199	25	194	4.1%	0.97 [0.58, 1.64]	
Veterans Admin 1969	13	424	21	422	2.4%	0.62 [0.31, 1.21]	
Subtotal (95% CI)		623		616	6.6%	0.82 [0.53, 1.27]	
Total events:	38		46				
Heterogeneity: $Tau^2 = 0.0$	01; Chi ² = 1.1	1, df = 1 (P = 0.29; I	$^{2} = 10\%$			
Test for overall effect: Z	= 0.90 (P = 0.00)	.37)					
2.28.3 >48mo							
Oslo Diet-Heart 1966	24	206	31	206	4.5%	0.77 [0.47 , 1.27]	_ _
WHI 2006 (1)	459	19541	684	29294	81.4%	1.01 [0.90 , 1.13]	•
Subtotal (95% CI)		19747		29500	85.9%	0.99 [0.88, 1.12]	T
Total events:	483		715				Ĭ
Heterogeneity: $Tau^2 = 0.0$	00; Chi ² = 1.0	1, df = 1 (P = 0.31; I	$^{2} = 1\%$			
Test for overall effect: Z	= 0.15 (P = 0.15)	.88)					
2.28.4 unclear							
Subtotal (95% CI)		0		0		Not estimable	
Total events:	0		0				
Heterogeneity: Not appli	cable						
Test for overall effect: N	ot applicable						
Total (95% CI)		21559		31275	100.0%	0.97 [0.87 , 1.07]	
Total events:	571		814				Ţ
Heterogeneity: $Tau^2 = 0.0$	00; $Chi^2 = 5.1$	1, df = 7 (P = 0.65; I	$^{2} = 0\%$			
Test for overall effect: Z	= 0.62 (P = 0.00)	.54)	. ,,			Fa	vours lower SFA Favours higher SFA
Test for subgroup differe	nces: $Chi^2 = 1$.30, df =	2 (P = 0.52)), $I^2 = 0\%$			

Analysis 2.28. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 28: Non-fatal MI, subgroup by duration

Footnotes

(1) Non-fatal MI during trial, Prentice 2017

Analysis 2.29. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 29: Non-fatal MI, subgroup by baseline SFA

	Lower SFA		Higher SFA			Risk Ratio	Risk Ratio	
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI	
2.29.1 up to 12%E SFA b	aseline							
Subtotal (95% CI)		0		0		Not estimable		
Total events:	0		0					
Heterogeneity: Not applica	ıble							
Test for overall effect: Not	applicable							
2.29.2 >12 to 15%E SFA	baseline							
DART 1989	35	1018	47	1015	6.0%	0.74 [0.48 , 1.14]		
Moy 2001	2	117	1	118	0.2%	2.02 [0.19, 21.94]		
WHI 2006 (1)	459	19541	684	29294	81.4%	1.01 [0.90 , 1.13]	· · · ·	
Subtotal (95% CI)		20676		30427	87.6%	0.97 [0.83 , 1.13]	—	
Total events:	496		732				Ť	
Heterogeneity: $Tau^2 = 0.00$); $Chi^2 = 2.1$	4, df = 2 (P = 0.34;	$[^2 = 6\%]$				
Test for overall effect: $Z =$	0.35 (P = 0)	.73)						
2.29.3 >15 to 18%E SFA	baseline							
Subtotal (95% CI)		0		0		Not estimable		
Total events:	0		0					
Heterogeneity: Not applica	ble							
Test for overall effect: Not	applicable							
2.29.4 >18%E SFA baseli	ine							
Veterans Admin 1969	13	424	21	422	2.4%	0.62 [0.31, 1.21]		
Subtotal (95% CI)		424		422	2.4%	0.62 [0.31, 1.21]		
Total events:	13		21					
Heterogeneity: Not applica	ible							
Test for overall effect: $Z =$	1.40 (P = 0)	.16)						
2.29.5 unclear								
MRC 1968	25	199	25	194	4.1%	0.97 [0.58, 1.64]		
Oslo Diet-Heart 1966	24	206	31	206	4.5%	0.77 [0.47, 1.27]		
Rose corn oil 1965	7	28	3	13	0.8%	1.08 [0.33 , 3.53]		
Rose olive 1965	6	26	2	13	0.5%	1.50 [0.35 , 6.43]		
Subtotal (95% CI)		459		426	10.0%	0.91 [0.65 , 1.27]		
Total events:	62		61					
Heterogeneity: $Tau^2 = 0.00$); $Chi^2 = 1.0$	1, $df = 3$ (P = 0.80;	$1^2 = 0\%$				
Test for overall effect: $Z =$	0.58 (P = 0)	.56)	. ,,					
Total (95% CI)		21559		31275	100.0%	0.97 [0.87 , 1.07]		
Total events:	571		814			. / 1	Ţ	
Heterogeneity: $Tau^2 = 0.00$); $Chi^2 = 5.1$	1, $df = 7$ (P = 0.65:	$[^2 = 0\%]$			+ $+$ $+$ $+$ $+$ $+$ $+$ $+$ $+$ $+$	
Test for overall effect: $Z =$	0.62 (P = 0)	.54)				Favo	burs lower SFA Favours higher	
Test for subgroup difference	ces: $Chi^2 = 1$.72. df = 100	2(P = 0.42)). $I^2 = 0\%$		1		

Footnotes

(1) Non-fatal MI during trial, Prentice 2017

Reduction in saturated fat intake for cardiovascular disease (Review) Copyright © 2020 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.
Analysis 2.30. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 30: Non-fatal MI, subgroup by SFA change

	Lower	SFA	Higher	SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
2.30.1 up to 4% E differen	nce						
DART 1989	35	1018	47	1015	6.0%	0.74 [0.48 , 1.14]	
Moy 2001	2	117	1	118	0.2%	2.02 [0.19, 21.94]	
WHI 2006 (1)	459	19541	684	29294	81.4%	1.01 [0.90 , 1.13]	•
Subtotal (95% CI)		20676		30427	87.6%	0.97 [0.83, 1.13]	•
Total events:	496		732				
Heterogeneity: $Tau^2 = 0.00$); Chi ² = 2.1	4, $df = 2$ (P = 0.34; I	$^{2} = 6\%$			
Test for overall effect: $Z =$	0.35 (P = 0	.73)					
2.30.2 >4 to 8%E differen	ice						
Subtotal (95% CI)		0		0		Not estimable	
Total events:	0		0				
Heterogeneity: Not applica	ible						
Test for overall effect: Not	applicable						
2.30.3 >8%E difference							
Veterans Admin 1969	13	424	21	422	2.4%	0.62 [0.31, 1.21]	
Subtotal (95% CI)		424		422	2.4%	0.62 [0.31 , 1.21]	
Total events:	13		21				•
Heterogeneity: Not applica	ıble						
Test for overall effect: $Z =$	1.40 (P = 0	.16)					
2.30.4 unclear							
MRC 1968	25	199	25	194	4.1%	0.97 [0.58, 1.64]	
Oslo Diet-Heart 1966	24	206	31	206	4.5%	0.77 [0.47 , 1.27]	
Rose corn oil 1965	7	28	3	13	0.8%	1.08 [0.33 , 3.53]	
Rose olive 1965	6	26	2	13	0.5%	1.50 [0.35 , 6.43]	_
Subtotal (95% CI)		459		426	10.0%	0.91 [0.65 , 1.27]	•
Total events:	62		61				
Heterogeneity: $Tau^2 = 0.00$); $Chi^2 = 1.0$	1, df = 3 (P = 0.80; I	$^{2} = 0\%$			
Test for overall effect: $Z =$	0.58 (P = 0	.56)					
Total (95% CI)		21559		31275	100.0%	0.97 [0.87 , 1.07]	
Total events:	571		814				
Heterogeneity: $Tau^2 = 0.00$); Chi ² = 5.1	1, df = 7 (P = 0.65; I	$^{2} = 0\%$		(0.01 0.1 1 10 100
Test for overall effect: Z =	0.62 (P = 0	.54)				Fa	vours lower SFA Favours higher SF

Test for subgroup differences: Chi² = 1.72, df = 2 (P = 0.42), I² = 0%

Footnotes

(1) Non-fatal MI during trial, Prentice 2017



Analysis 2.31. Comparison 2: SFA reduction vs usual diet secondary health events, Outcome 31: Non-fatal MI, subgroup by sex

	Lower	SFA	Higher	SFA		Risk Ratio	Ris	k Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Ran	dom, 95% CI
2.31.1 Men								
DART 1989	35	1018	47	1015	6.0%	0.74 [0.48 , 1.14	·] _	•
MRC 1968	25	199	25	194	4.1%	0.97 [0.58 , 1.64	·] .	_
Oslo Diet-Heart 1966	24	206	31	206	4.5%	0.77 [0.47 , 1.27] _	•
Rose corn oil 1965	7	28	3	13	0.8%	1.08 [0.33 , 3.53]	
Rose olive 1965	6	26	2	13	0.5%	1.50 [0.35 , 6.43]	
Veterans Admin 1969	13	424	21	422	2.4%	0.62 [0.31, 1.21] _	_
Subtotal (95% CI)		1901		1863	18.4%	0.81 [0.63 , 1.03]	
Total events:	110		129					•
Heterogeneity: $Tau^2 = 0.00$;	$Chi^2 = 2.2$	d3, df = 5 (P = 0.82; I	$^{2} = 0\%$				
Test for overall effect: $Z = 1$	1.71 ($P = 0$.09)						
2.31.2 Women								
WHI 2006 (1)	459	19541	684	29294	81.4%	1.01 [0.90 , 1.13]	
Subtotal (95% CI)		19541		29294	81.4%	1.01 [0.90 , 1.13]	•
Total events:	459		684					
Heterogeneity: Not applicat	ole							
Test for overall effect: $Z = 0$	0.10 (P = 0)	.92)						
2.31.3 Mixed, men and wo	men							
Moy 2001	2	117	1	118	0.2%	2.02 [0.19 , 21.94	·]	
Subtotal (95% CI)		117		118	0.2%	2.02 [0.19 , 21.94		
Total events:	2		1					
Heterogeneity: Not applicat	ole							
Test for overall effect: $Z = 0$	0.58 (P = 0)	.56)						
Total (95% CI)		21559		31275	100.0%	0.97 [0.87 , 1.07]	•
Total events:	571		814					
Heterogeneity: $Tau^2 = 0.00$;	$Chi^2 = 5.1$	1, df = 7 (P = 0.65; I	$^{2} = 0\%$			0.01 0.1	1 10 100
Test for overall effect: $Z = 0$	0.62 (P = 0)	.54)					Favours lower SFA	Favours higher SFA
Test for subgroup difference	es: Chi ² = 2	2.89, df = 2	2 (P = 0.24)), $I^2 = 30.7$	%			

Footnotes

(1) Non-fatal MI during trial, Prentice 2017

Analysis 2.32. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 32: Non-fatal MI, subgroup by CVD risk

	Lower	SFA	Higher	SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
2.32.1 Low CVD risk							
Veterans Admin 1969	13	424	21	422	5.9%	0.62 [0.31, 1.21]	
WHI 2006	353	18633	581	27925	37.3%	0.91 [0.80 , 1.04]	
Subtotal (95% CI)		19057		28347	43.3%	0.87 [0.68, 1.12]	▲
Total events:	366		602				•
Heterogeneity: $Tau^2 = 0.0$	01; Chi ² = 1.2	23, df = 1 (P = 0.27; I	$^{2} = 19\%$			
Test for overall effect: Z	= 1.11 (P = 0)	.27)					
2.32.2 Moderate CVD r	isk						
Moy 2001	2	117	1	118	0.5%	2.02 [0.19, 21.94]	
Subtotal (95% CI)		117		118	0.5%	2.02 [0.19 , 21.94]	
Total events:	2		1				
Heterogeneity: Not appli	cable						
Test for overall effect: Z	= 0.58 (P = 0)	.56)					
2.32.3 Existing CVD dis	sease						
DART 1989	35	1018	47	1015	12.4%	0.74 [0.48 , 1.14]	
MRC 1968	25	199	25	194	9.3%	0.97 [0.58, 1.64]	_ _
Oslo Diet-Heart 1966	24	206	31	206	10.0%	0.77 [0.47 , 1.27]	
Rose corn oil 1965	7	28	3	13	2.1%	1.08 [0.33 , 3.53]	
Rose olive 1965	6	26	2	13	1.4%	1.50 [0.35 , 6.43]	_
WHI 2006 (1)	82	908	90	1369	20.9%	1.37 [1.03 , 1.83]	-
Subtotal (95% CI)		2385		2810	56.2%	1.00 [0.76 , 1.31]	
Total events:	179		198				Ĭ
Heterogeneity: Tau ² = 0.	04; Chi ² = 7.6	63, df = 5 (P = 0.18; I	$2^2 = 34\%$			
Test for overall effect: Z	= 0.01 (P = 0	.99)					
Total (95% CI)		21559		31275	100.0%	0.95 [0.80 , 1.13]	
Total events:	547		801				Ĭ
Heterogeneity: $Tau^2 = 0.0$	02; $Chi^2 = 11$.03, df = 8	(P = 0.20);	$\mathrm{I}^{2}=27\%$			0.01 0.1 1 10 100
Test for overall effect: Z	= 0.56 (P = 0)	.57)				Fa	avours lower SFA Favours higher SFA
Test for subgroup differe	ences: Chi ² = 0	0.98, df = 2	2 (P = 0.61)), $I^2 = 0\%$			

Footnotes

(1) Women with CVD at baseline

Analysis 2.33. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 33: Non-fatal MI, subgroup by TC reduction

Trusted evidence.

Better health.

Informed decisions.

Cochrane

Library

	Lower	SFA	Higher	SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
2.33.1 serum chol reduce	d by at leas	t 0.2mmo	I/L				
DART 1989	35	1018	47	1015	33.4%	0.74 [0.48 , 1.14]	
MRC 1968	25	199	25	194	22.9%	0.97 [0.58, 1.64]	
Moy 2001	2	117	1	118	1.1%	2.02 [0.19, 21.94]	
Oslo Diet-Heart 1966	24	206	31	206	24.9%	0.77 [0.47 , 1.27]	
Rose corn oil 1965	7	28	3	13	4.4%	1.08 [0.33 , 3.53]	
Veterans Admin 1969	13	424	21	422	13.3%	0.62 [0.31, 1.21]	
Subtotal (95% CI)		1992		1968	100.0%	0.80 [0.62 , 1.03]	
Total events:	106		128				•
Heterogeneity: $Tau^2 = 0.00$	0; $Chi^2 = 2.0$	9, df = 5 (P = 0.84; I	$^{2} = 0\%$			
Test for overall effect: Z =	1.76 (P = 0.1)	.08)					
2.33.2 serum chol reduce	d by <0.2m	mol/L					
Rose olive 1965	6	26	2	13	0.6%	1.50 [0.35 , 6.43]	
WHI 2006 (1)	459	19541	684	29294	99.4%	1.01 [0.90 , 1.13]	
Subtotal (95% CI)		19567		29307	100.0%	1.01 [0.90 , 1.13]	•
Total events:	465		686				
Heterogeneity: $Tau^2 = 0.00$	0; $Chi^2 = 0.2$	9, df = 1 (P = 0.59); I	$^{2} = 0\%$			
Test for overall effect: Z =	0.14 (P = 0.14)	.89)					
2 33 3 serum chol reduct	ion unclear						
Subtotal (95% CI)	ion uncicui	0		0		Not estimable	
Total events:	0	Ū	0	v		i tot estimable	
Heterogeneity: Not applica	able		0				
Test for overall effect: Not	t applicable						
rest for overall effect. No	upplicable						
Test for subgroup differen	ces: Chi ² = 2	2.73, df =	1 (P = 0.10)), $I^2 = 63.4$.%	Fa	0.01 0.1 1 10 100 wours lower SFA Favours higher SFA

Footnotes

(1) Non-fatal MI during trial, Prentice 2017

Analysis 2.34. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 34: Non-fatal MI, subgroup decade of publication

	Lower	SFA	Higher	SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
2.34.1 1960s							
MRC 1968	25	199	25	194	4.1%	0.97 [0.58, 1.64]	
Oslo Diet-Heart 1966	24	206	31	206	4.5%	0.77 [0.47 , 1.27]	
Rose corn oil 1965	7	28	3	13	0.8%	1.08 [0.33 , 3.53]	
Rose olive 1965	6	26	2	13	0.5%	1.50 [0.35, 6.43]	
Veterans Admin 1969	13	424	21	422	2.4%	0.62 [0.31, 1.21]	
Subtotal (95% CI)		883		848	12.4%	0.84 [0.62, 1.13]	▲
Total events:	75		82			- / -	•
Heterogeneity: $Tau^2 = 0.0$	00; Chi ² = 2.0	1, df = 4 (P = 0.73;	$2^2 = 0\%$			
Test for overall effect: Z	= 1.14 (P = 0)	.26)					
2.34.2 1970s							
Subtotal (95% CI)		0		0		Not estimable	
Total events:	0		0				
Heterogeneity: Not applie	cable						
Test for overall effect: No	ot applicable						
2.34.3 1980s							
DART 1989	35	1018	47	1015	6.0%	0.74 [0.48 , 1.14]	
Subtotal (95% CI)		1018		1015	6.0%	0.74 [0.48 , 1.14]	•
Total events:	35		47				•
Heterogeneity: Not applie	cable						
Test for overall effect: Z	= 1.36 (P = 0)	.17)					
2.34.4 1990s							
Subtotal (95% CI)		0		0		Not estimable	
Total events:	0		0				
Heterogeneity: Not applie	cable						
Test for overall effect: No	ot applicable						
2.34.5 2000s							
Moy 2001	2	117	1	118	0.2%	2.02 [0.19 , 21.94]	
WHI 2006 (1)	459	19541	684	29294	81.4%	1.01 [0.90 , 1.13]	
Subtotal (95% CI)		19658		29412	81.6%	1.01 [0.90 , 1.13]	•
Total events:	461		685				
Heterogeneity: $Tau^2 = 0.0$	00; Chi ² = 0.3	3, df = 1 (P = 0.57;	$2^2 = 0\%$			
Test for overall effect: Z	= 0.13 (P = 0.13)	.90)					
Total (95% CI)		21559		31275	100.0%	0.97 [0.87 , 1.07]	
Total events:	571		814				
Heterogeneity: $Tau^2 = 0.0$	00; Chi ² = 5.1	1, df = 7 (P = 0.65;	$1^2 = 0\%$		0.0	01 0.1 1 10 10
Test for overall effect: Z	= 0.62 (P = 0)	.54)				Favo	urs lower SFA Favours higher
Test for subgroup differe	nces: $Chi^2 = 2$	2.78, df = 1	2 (P = 0.25)), $I^2 = 28.0$	%		

Footnotes

(1) Non-fatal MI during trial, Prentice 2017

Cochrane
Library

	Lower	SFA	Higher	SFA		Risk Ratio	Risk R	atio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Randor	m, 95% CI
Ley 2004	1	88	5	88	2.0%	0.20 [0.02 , 1.68]		-
MRC 1968	2	199	0	194	1.0%	4.88 [0.24 , 100.89]		
Moy 2001	1	117	1	118	1.2%	1.01 [0.06 , 15.93]		
Oslo Diet-Heart 1966	2	206	1	206	1.6%	2.00 [0.18 , 21.89]		•
STARS 1992	0	27	1	28	0.9%	0.35 [0.01, 8.12]		
Veterans Admin 1969	13	424	22	422	16.4%	0.59 [0.30 , 1.15]		
WHI 2006 (1)	435	19541	634	29294	76.9%	1.03 [0.91 , 1.16]	•	
Total (95% CI)		20602		30350	100.0%	0.92 [0.68 , 1.25]	•	
Total events:	454		664				Ĩ	
Heterogeneity: $Tau^2 = 0.03$; Chi ² = 6.5	9, df = 6 (P = 0.36; I	$^{2} = 9\%$			0.01 0.1 1	10 100
Test for overall effect: Z =	0.52 (P = 0	.60)				F	avours lower SFA	Favours higher SFA

Analysis 2.35. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 35: STROKE

Test for overall effect: Z = 0.52 (P = 0.60)

Test for subgroup differences: Not applicable

Footnotes

(1) During trial, Prentice 2017

Analysis 2.36. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 36: Stroke, SA by low summary risk of bias

	Lower SFA		Higher SFA			Risk Ratio	Risk R	atio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Randor	n, 95% CI
Ley 2004	1	88	5	88	7.0%	0.20 [0.02 , 1.68]	-
Veterans Admin 1969	13	424	22	422	34.2%	0.59 [0.30 , 1.15		
WHI 2006 (1)	435	19541	634	29294	58.8%	1.03 [0.91 , 1.16) <mark>–</mark>	
Total (95% CI)		20053		29804	100.0%	0.76 [0.42 , 1.38		
Total events:	449		661				•	
Heterogeneity: $Tau^2 = 0.15$	5; Chi ² = 4.8	0, df = 2 (P = 0.09; I	$^{2} = 58\%$			0.01 0.1 1	10 100
Test for overall effect: Z =	0.91 (P = 0	.36)					Favours lower SFA	Favours higher SFA
Test for subgroup different	ces: Not app	licable						

Footnotes

(1) During trial, Prentice 2017



Analysis 2.37. Comparison 2: SFA reduction vs usual diet secondary health events, Outcome 37: Stroke, SA aim to reduce SFA

	Lower	SFA	Higher	SFA		Risk Ratio	Risk R	atio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Randor	n, 95% CI
MRC 1968	2	199	0	194	0.2%	4.88 [0.24 , 100.89]		
Moy 2001	1	117	1	118	0.2%	1.01 [0.06 , 15.93]		
Oslo Diet-Heart 1966	2	206	1	206	0.2%	2.00 [0.18, 21.89]		
STARS 1992	0	27	1	28	0.1%	0.35 [0.01, 8.12]		
Veterans Admin 1969	13	424	22	422	3.1%	0.59 [0.30 , 1.15]		
WHI 2006 (1)	435	19541	634	29294	96.2%	1.03 [0.91 , 1.16]		
Total (95% CI)		20514		30262	100.0%	1.01 [0.90 , 1.14]		
Total events:	453		659					
Heterogeneity: $Tau^2 = 0.0$	00; Chi ² = 4.3	6, df = 5 (P = 0.50; I	$^{2} = 0\%$			0.01 0.1 1	10 100
Test for overall effect: Z	= 0.22 (P = 0)	.83)				F	Favours lower SFA	Favours higher SFA
Test for subgroup differe	nces: Not app	olicable						

Footnotes

(1) During trial, Prentice 2017

Analysis 2.38. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 38: Stroke, SA statistically significant SFA reduction

	Lower SFA		Higher SFA		Risk Ratio		Risk Ra	ntio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random	n, 95% CI
Ley 2004	1	88	5	88	3.6%	0.20 [0.02 , 1.68]		
Moy 2001	1	117	1	118	2.2%	1.01 [0.06 , 15.93]		
STARS 1992	0	27	1	28	1.7%	0.35 [0.01, 8.12]		
Veterans Admin 1969	13	424	22	422	24.9%	0.59 [0.30 , 1.15]		
WHI 2006 (1)	435	19541	634	29294	67.6%	1.03 [0.91 , 1.16]	•	
Total (95% CI)		20197		29950	100.0%	0.83 [0.55 , 1.25]		
Total events:	450		663					
Heterogeneity: $Tau^2 = 0.06$	6; Chi ² = 5.2	4, df = 4 (P = 0.26; I	$^{2} = 24\%$			0.01 0.1 1	10 100
Test for overall effect: Z =	0.89 (P = 0	.37)				F	Favours lower SFA	Favours higher SFA

Test for subgroup differences: Not applicable

Footnotes

(1) During trial, Prentice 2017



Analysis 2.39. Comparison 2: SFA reduction vs usual diet secondary health events, Outcome 39: Stroke, SA by TC reduction

	Lower	SFA	Higher	SFA		Risk Ratio	Risk R	atio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Randon	n, 95% CI
MRC 1968	2	199	0	194	0.2%	4.88 [0.24 , 100.89]		_ .
Moy 2001	1	117	1	118	0.2%	1.01 [0.06 , 15.93]	I	
Oslo Diet-Heart 1966	2	206	1	206	0.2%	2.00 [0.18 , 21.89]		<u> </u>
STARS 1992	0	27	1	28	0.1%	0.35 [0.01, 8.12]		
Veterans Admin 1969	13	424	22	422	3.1%	0.59 [0.30 , 1.15]		
WHI 2006 (1)	435	19541	634	29294	96.2%	1.03 [0.91 , 1.16]	· 📫	
Total (95% CI)		20514		30262	100.0%	1.01 [0.90 , 1.14]		
Total events:	453		659					
Heterogeneity: $Tau^2 = 0.0$	00; $Chi^2 = 4.3$	6, df = 5 (P = 0.50; I	$^{2} = 0\%$			0.01 0.1 1	10 100
Test for overall effect: Z	= 0.22 (P = 0)	.83)				H	Favours lower SFA	Favours higher SFA
Test for subgroup differen	nces: Not app	olicable						

Footnotes

(1) During trial, Prentice 2017

Analysis 2.40. Comparison 2: SFA reduction vs usual diet secondary health events, Outcome 40: Stroke, SA excluding WHI

	Lower	SFA	Higher	SFA		Risk Ratio	Risk F	latio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Rando	m, 95% CI
Ley 2004	1	88	5	88	7.5%	0.20 [0.02 , 1.68]]	_
MRC 1968	2	199	0	194	3.7%	4.88 [0.24, 100.89]]	
Moy 2001	1	117	1	118	4.5%	1.01 [0.06, 15.93]]	
Oslo Diet-Heart 1966	2	206	1	206	5.9%	2.00 [0.18, 21.89]]	_
STARS 1992	0	27	1	28	3.4%	0.35 [0.01 , 8.12]]	
Veterans Admin 1969	13	424	22	422	75.0%	0.59 [0.30 , 1.15]]	
Total (95% CI)		1061		1056	100.0%	0.63 [0.35 , 1.14]	1	
Total events:	19		30				•	
Heterogeneity: $Tau^2 = 0.0$	00; Chi ² = 4.0	6, df = 5 (P = 0.54; I	$[^2 = 0\%]$			$\frac{1}{001}$ 01 1	10 100
Test for overall effect: Z	= 1.53 (P = 0)	.13)]	Favours lower SFA	Favours higher SFA

Test for subgroup differences: Not applicable



Analysis 2.41. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 41: Stroke, SA Mantel-Haenszel fixed-effect

	Lower	SFA	Higher	SFA		Risk Ratio	Risk	Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Fixed, 95% CI	M-H, Fixe	d, 95% CI
Ley 2004	1	88	5	88	0.9%	0.20 [0.02 , 1.68]	_
MRC 1968	2	199	0	194	0.1%	4.88 [0.24, 100.89]]	
Moy 2001	1	117	1	118	0.2%	1.01 [0.06 , 15.93]	
Oslo Diet-Heart 1966	2	206	1	206	0.2%	2.00 [0.18 , 21.89]]	
STARS 1992	0	27	1	28	0.3%	0.35 [0.01, 8.12]]	
Veterans Admin 1969	13	424	22	422	4.1%	0.59 [0.30 , 1.15]	-
WHI 2006 (1)	435	19541	634	29294	94.2%	1.03 [0.91 , 1.16]	
Total (95% CI)		20602		30350	100.0%	1.01 [0.89 , 1.13]	
Total events:	454		664					
Heterogeneity: $Chi^2 = 6.5$	59, df = 6 (P =	= 0.36); I ²	= 9%				0.01 0.1	10 100
Test for overall effect: Z					Favours lower SFA	Favours higher SFA		
								-

Test for subgroup differences: Not applicable

Footnotes

(1) During trial, Prentice 2017

Analysis 2.42. Comparison 2: SFA reduction vs usual diet secondary health events, Outcome 42: Stroke, SA Peto fixed-effect

	Lower	SFA	Higher SFA		Peto Odds Ratio		Peto Odd	s Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	Peto, Fixed, 95% CI	Peto, Fixed	,95% CI
Ley 2004	1	88	5	88	0.6%	0.25 [0.05 , 1.29]		
MRC 1968	2	199	0	194	0.2%	7.24 [0.45 , 116.22]		
Moy 2001	1	117	1	118	0.2%	1.01 [0.06 , 16.22]		
Oslo Diet-Heart 1966	2	206	1	206	0.3%	1.95 [0.20 , 18.89]		
STARS 1992	0	27	1	28	0.1%	0.14 [0.00 , 7.07]	• • •	
Veterans Admin 1969	13	424	22	422	3.2%	0.58 [0.30 , 1.15]		
WHI 2006 (1)	435	19541	634	29294	95.5%	1.03 [0.91 , 1.16]		
Total (95% CI)		20602		30350	100.0%	1.01 [0.89 , 1.14]		
Total events:	454		664					
Heterogeneity: $Chi^2 = 8.65$	5, $df = 6 (P = $	= 0.19); I ² =	= 31%				0.01 0.1 1	10 100
Test for overall effect: $Z = 0.11$ (P = 0.92)						F	Favours lower SFA	Favours higher SFA

Test for subgroup differences: Not applicable

Footnotes

(1) During trial, Prentice 2017

Analysis 2.43. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 43: Stroke, subgroup by any substitution

	Lower	SFA	Higher	SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
2.43.1 replaced by PUFA							
MRC 1968	2	199	0	194	4.2%	4.88 [0.24, 100.89]	
Oslo Diet-Heart 1966	2	206	1	206	6.7%	2.00 [0.18 . 21.89]	
STARS 1992	0	27	1	28	3.9%	0.35 [0.01, 8.12]	
Veterans Admin 1969	13	424	22	422	85.2%	0.59 [0.30, 1.15]	
Subtotal (95% CI)		856		850	100.0%	0.68 [0.37 , 1.27]	
Total events:	17		24				
Heterogeneity: $Tau^2 = 0.00$: $Chi^2 = 2.7$	8. $df = 3$ (P = 0.43):]	$2^2 = 0\%$			
Test for overall effect: $Z =$	1.20 (P = 0)	.23)					
2 42 2 words and her MUEA							
2.45.2 replaced by MUFA	L	^		~		ЪТ-4 4° П.Т	
Subtotal (95% CI)	0	0	0	0		Not estimable	
i otal events:	0		0				
Heterogeneity: Not applica	ble						
lest for overall effect: Not	applicable						
2.43.3 replace by CHO							
Ley 2004	1	88	5	88	15.6%	0.20 [0.02 , 1.68]	
STARS 1992	0	27	1	28	7.9%	0.35 [0.01, 8.12]	
WHI 2006 (1)	435	19541	634	29294	76.5%	1.03 [0.91 , 1.16]	•
Subtotal (95% CI)		19656		29410	100.0%	0.73 [0.29 , 1.87]	—
Total events:	436		640				
Heterogeneity: Tau ² = 0.30	; Chi ² = 2.7	3, $df = 2$ (P = 0.26; I	$^{2} = 27\%$			
Test for overall effect: $Z =$	0.65 (P = 0	.51)					
2.43.4 replaced by proteir	1						
Lev 2004	- 1	88	5	88	28.1%	0.20 [0.02 , 1.68]	
WHI 2006 (1)	435	19541	634	29294	71.9%	1.03 [0.91 , 1.16]	
Subtotal (95% CI)		19629		29382	100.0%	0.65 [0.15 , 2.75]	
Total events:	436		639	_/00	/		
Heterogeneity: Tau ² = 0.75	; Chi ² = 2.2	7, $df = 1$ (P = 0.13): 1	$2^2 = 56\%$			
Test for overall effect: $Z =$	0.59 (P = 0)	.56)					
2 43 5 replacement uncles	ar						
Mov 2001	1	117	1	110	100.00/-	1 01 [0 06 15 02]	
Subtotal (050% CI)	1	11/	1	110	100.0%		
Total events:	1	11/	1	118	100.0%	1.01 [0.00 , 15.95]	
I Utal EVEIIIS. Hataroganaity: Nat appliaa	l bla		1				
Test for overall effect: $Z =$	0.01 (P = 1)	.00)					
	- \	- /					
Test for subgroup difference	$ces: Chi^2 = 0$	0.09, df = 1	3 (P = 0.99)), $I^2 = 0\%$			0.005 0.1 1 10 200
						Fav	ours lower SFA Favours highe
Footnotes							

(1) During trial, Prentice 2017

Analysis 2.44. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 44: Stroke, subgroup by main substitution

	Lower	SFA	Higher	SFA		Risk Ratio	Risk Ratio
tudy or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
.44.1 replaced by PUFA							
/IRC 1968	2	199	0	194	11.2%	4.88 [0.24 , 100.89]	
Slo Diet-Heart 1966	2	206	1	206	16.8%	2.00 [0.18 , 21.89]	
veterans Admin 1969	13	424	22	422	72.0%	0.59 [0.30 , 1.15]	
ubtotal (95% CI)		829		822	100.0%	0.92 [0.31 , 2.69]	
otal events:	17		23				–
Heterogeneity: Tau ² = 0.30;	$Chi^2 = 2.6$	0, df = 2 (P = 0.27; I	² = 23%			
Test for overall effect: $Z = 0$	0.16 (P = 0.16)	.87)					
.44.2 replaced by MUFA							
ubtotal (95% CI)		0		0		Not estimable	
otal events:	0		0				
leterogeneity: Not applicat	ole						
est for overall effect: Not a	applicable						
.44.3 replace by CHO							
ey 2004	1	88	5	88	15.6%	0.20 [0.02 , 1.68]	
TARS 1992	0	27	1	28	7.9%	0.35 [0.01, 8.12]	_
VHI 2006 (1)	435	19541	634	29294	76.5%	1.03 [0.91 , 1.16]	
ubtotal (95% CI)		19656		29410	100.0%	0.73 [0.29 , 1.87]	-
otal events:	436		640				
Ieterogeneity: Tau ² = 0.30;	$Chi^2 = 2.7$	3, df = 2 (P = 0.26; I	² = 27%			
Test for overall effect: $Z = 0$	0.65 (P = 0.65)	.51)					
.44.4 replaced by protein							
ubtotal (95% CI)		0		0		Not estimable	
otal events:	0		0				
leterogeneity: Not applicat	ole						
est for overall effect: Not a	applicable						
.44.5 replacement unclea	r						
loy 2001	1	117	1	118	100.0%	1.01 [0.06 , 15.93]	
ubtotal (95% CI)		117		118	100.0%	1.01 [0.06 , 15.93]	
otal events:	1		1				T
leterogeneity: Not applicat	ole						
Test for overall effect: $Z = 0$	0.01 (P = 1)	.00)					
fest for subgroup difference	es: $Chi^2 = 0$	0.12, df = 2	2 (P = 0.94)	$, I^2 = 0\%$			0.005 0.1 1 10 200
						F	Favours lower SFA Favours higher SF
ootnotes							

1) During trial, Prentice 2017

Analysis 2.45. Comparison 2: SFA reduction vs usual diet secondary health events, Outcome 45: Stroke, subgroup by duration

	Lower	SFA	Higher	SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
2.45.1 up to 24mo							
Moy 2001	1	117	1	118	1.2%	1.01 [0.06, 15.93]	1
Subtotal (95% CI)		117		118	1.2%	1.01 [0.06 , 15.93]	
Total events:	1		1				
Heterogeneity: Not applic	able						
Test for overall effect: Z =	= 0.01 (P = 1)	.00)					
2.45.2 >24 to 48mo							
STARS 1992	0	27	1	28	0.9%	0.35 [0.01 , 8.12]]
Veterans Admin 1969	13	424	22	422	16.3%	0.59 [0.30 , 1.15]]
Subtotal (95% CI)		451		450	17.2%	0.57 [0.30 , 1.11]	
Total events:	13		23				•
Heterogeneity: $Tau^2 = 0.0$	0; Chi ² = 0.1	0, df = 1 (P = 0.75; I	$^{2} = 0\%$			
Test for overall effect: Z =	= 1.65 (P = 0)	.10)					
2.45.3 >48mo							
Oslo Diet-Heart 1966	2	206	1	206	1.6%	2.00 [0.18, 21.89]	
WHI 2006 (1)	435	19541	634	29294	78.1%	1.03 [0.91 , 1.16]]
Subtotal (95% CI)		19747		29500	79.6%	1.03 [0.91 , 1.16]	1 1
Total events:	437		635				
Heterogeneity: $Tau^2 = 0.0$	0; Chi ² = 0.3	0, df = 1 (P = 0.59; I	$^{2} = 0\%$			
Test for overall effect: Z =	= 0.49 (P = 0)	.63)					
2.45.4 unclear duration							
Ley 2004	1	88	5	88	2.0%	0.20 [0.02 , 1.68]	1
Subtotal (95% CI)		88		88	2.0%	0.20 [0.02 , 1.68]	
Total events:	1		5				
Heterogeneity: Not applic	able						
Test for overall effect: Z =	= 1.48 (P = 0)	.14)					
Total (95% CI)		20403		30156	100.0%	0.91 [0.67 , 1.23]	1
Total events:	452		664				
Heterogeneity: $Tau^2 = 0.0$	3; Chi ² = 5.5	6, df = 5 (P = 0.35; I	$^{2} = 10\%$			0.005 0.1 1 10 200
Test for overall effect: Z =	= 0.62 (P = 0)	.54)]	Favours lower SFA Favours higher SFA
Test for subgroup differen	nces: $Chi^2 = 5$	5.15, df =	3 (P = 0.16)), $I^2 = 41.8$	%		

Footnotes

(1) During trial, Prentice 2017



Analysis 2.46. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 46: Stroke, subgroup by baseline SFA

	Lower SFA		Higher SFA		Risk Ratio		Risk	Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Rand	om, 95% CI
2.46.1 up to 12%E SFA I	baseline							
Subtotal (95% CI)		0		0		Not estimable		
Total events:	0		0					
Heterogeneity: Not application	able							
Test for overall effect: No	t applicable							
2.46.2 >12 to 15%E SFA	baseline							
Ley 2004	1	88	5	88	2.0%	0.20 [0.02, 1.68]		
Moy 2001	1	117	1	118	1.2%	1.01 [0.06 , 15.93]		
WHI 2006 (1)	435	19541	634	29294	78.1%	1.03 [0.91, 1.16]		
Subtotal (95% CI)		19746		29500	81.2%	0.91 [0.50 , 1.66]		
Total events:	437		640			- / -		
Heterogeneity: $Tau^2 = 0.10$	0; Chi ² = 2.2	7, df = 2 (P = 0.32; I	$^{2} = 12\%$				
Test for overall effect: Z =	= 0.30 (P = 0)	.76)	,,					
2.46.3 >15 to 18%E SFA	baseline							
STARS 1992	0	27	1	28	0.9%	0.35 [0.01, 8.12]		
Subtotal (95% CI)		27		28	0.9%	0.35 [0.01, 8.12]		
Total events:	0		1					
Heterogeneity: Not application	able							
Test for overall effect: Z =	= 0.66 (P = 0)	.51)						
2.46.4 >18%E SFA basel	line							
Veterans Admin 1969	13	424	22	422	16.3%	0.59 [0.30, 1.15]		-
Subtotal (95% CI)		424		422	16.3%	0.59 [0.30 , 1.15]		
Total events:	13		22				•	
Heterogeneity: Not application	able							
Test for overall effect: Z =	= 1.55 (P = 0)	.12)						
2.46.5 unclear								
Oslo Diet-Heart 1966	2	206	1	206	1.6%	2.00 [0.18 , 21.89]		
Subtotal (95% CI)		206		206	1.6%	2.00 [0.18 , 21.89]		
Total events:	2		1			- / -		
Heterogeneity: Not application	able							
Test for overall effect: Z =	= 0.57 (P = 0)	.57)						
Total (95% CI)		20403		30156	100.0%	0.91 [0.67 , 1.23]		
Total events:	452		664			. –		ſ
Heterogeneity: $Tau^2 = 0.02$	3; Chi ² = 5.5	6, df = 5 (P = 0.35; I	$^{2} = 10\%$			0.02 0.1	1 10 50
Test for overall effect: Z =	= 0.62 (P = 0)	.54)				F	Favours lower SFA	Favours higher S
Test for subgroup differen	ces: $Chi^2 = 1$	1.79, df = 3	3 (P = 0.62)	$J_{12} = 0\%$				-

Footnotes

(1) During trial, Prentice 2017

Analysis 2.47. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 47: Stroke, subgroup by SFA change

	Lower	SFA	Higher	SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
2.47.1 up to 4%E differ	ence						
Ley 2004	1	88	5	88	2.0%	0.20 [0.02, 1.68]	I
Moy 2001	1	117	1	118	1.2%	1.01 [0.06 , 15.93]	I
WHI 2006 (1)	435	19541	634	29294	78.1%	1.03 [0.91 , 1.16]	1 💼
Subtotal (95% CI)		19746		29500	81.2%	0.91 [0.50 , 1.66]	I 👗
Total events:	437		640				T
Heterogeneity: $Tau^2 = 0$.	10; $Chi^2 = 2.2$	7, $df = 2$ (P = 0.32; I	$^{2} = 12\%$			
Test for overall effect: Z	= 0.30 (P = 0.1)	.76)					
2.47.2 >4 to 8%E differ	ence						
STARS 1992	0	27	1	28	0.9%	0.35 [0.01, 8.12]	I
Subtotal (95% CI)		27		28	0.9%	0.35 [0.01 , 8.12]	
Total events:	0		1				
Heterogeneity: Not appli	icable						
Test for overall effect: Z	= 0.66 (P = 0.66)	.51)					
2.47.3 >8%E difference	e						
Veterans Admin 1969	13	424	22	422	16.3%	0.59 [0.30 , 1.15]	I
Subtotal (95% CI)		424		422	16.3%	0.59 [0.30 , 1.15]	
Total events:	13		22				•
Heterogeneity: Not appli	cable						
Test for overall effect: Z	= 1.55 (P = 0.	.12)					
2.47.4 unclear							
Oslo Diet-Heart 1966	2	206	1	206	1.6%	2.00 [0.18, 21.89]	I
Subtotal (95% CI)		206		206	1.6%	2.00 [0.18 , 21.89]	
Total events:	2		1				
Heterogeneity: Not appli	icable						
Test for overall effect: Z	= 0.57 (P = 0.57)	.57)					
Total (95% CI)		20403		30156	100.0%	0.91 [0.67 , 1.23]	. ↓
Total events:	452		664]
Heterogeneity: $Tau^2 = 0$.	03; Chi ² = 5.5	6, df = 5 (P = 0.35; I	$^{2} = 10\%$			0.005 0.1 1 10 200
Test for overall effect: Z	= 0.62 (P = 0.62)	.54)				1	Favours lower SFA Favours higher SFA
Test for subgroup differe	ences: Chi ² = 1	.79, df = 1	3 (P = 0.62)), $I^2 = 0\%$			

Footnotes

(1) During trial, Prentice 2017

Analysis 2.48. Comparison 2: SFA reduction vs usual diet secondary health events, Outcome 48: Stroke, subgroup by sex

	Lower SFA		Higher	Higher SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
2.48.1 Men							
Oslo Diet-Heart 1966	2	206	1	206	1.6%	2.00 [0.18 , 21.89]
STARS 1992	0	27	1	28	0.9%	0.35 [0.01, 8.12]
Veterans Admin 1969	13	424	22	422	16.3%	0.59 [0.30 , 1.15]
Subtotal (95% CI)		657		656	18.8%	0.63 [0.33 , 1.18]
Total events:	15		24				•
Heterogeneity: $Tau^2 = 0.00$); $Chi^2 = 1.0$	8, df = 2 (P = 0.58; I	$^{2} = 0\%$			
Test for overall effect: Z =	1.44 (P = 0	.15)					
2.48.2 Women							
WHI 2006 (1)	435	19541	634	29294	78.1%	1.03 [0.91 , 1.16]
Subtotal (95% CI)		19541		29294	78.1%	1.03 [0.91 , 1.16]
Total events:	435		634				
Heterogeneity: Not applica	ıble						
Test for overall effect: Z =	0.46 (P = 0	.65)					
2.48.3 Mixed, men and w	omen						
Ley 2004	1	88	5	88	2.0%	0.20 [0.02, 1.68]
Moy 2001	1	117	1	118	1.2%	1.01 [0.06 , 15.93]
Subtotal (95% CI)		205		206	3.1%	0.37 [0.07 , 1.97	
Total events:	2		6				
Heterogeneity: $Tau^2 = 0.00$); $Chi^2 = 0.8$	4, df = 1 (P = 0.36; I	$^{2} = 0\%$			
Test for overall effect: Z =	1.17 (P = 0	.24)					
Total (95% CI)		20403		30156	100.0%	0.91 [0.67 , 1.23]
Total events:	452		664				Ĭ
Heterogeneity: $Tau^2 = 0.03$	3; Chi ² = 5.5	6, df = 5 (P = 0.35; I	$^{2} = 10\%$			0.005 0.1 1 10 200
Test for overall effect: Z =	0.62 (P = 0	.54)					Favours lower SFA Favours higher SFA
Test for subgroup differen	ces: $Chi^2 = 3$	3.65, df = 2	2 (P = 0.16)), $I^2 = 45.2$.%		

Footnotes

(1) During trial, Prentice 2017

Analysis 2.49. Comparison 2: SFA reduction vs usual diet secondary health events, Outcome 49: Stroke, subgroup by CVD risk

	Lower SFA		Higher	Higher SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
2.49.1 Low CVD risk							
Veterans Admin 1969	13	424	22	422	2.8%	0.59 [0.30 , 1.15]	1
WHI 2006 (1)	228	18633	334	27925	44.6%	1.02 [0.87 , 1.21]	1
Subtotal (95% CI)		19057		28347	47.4%	0.86 [0.52, 1.42]	1 👗
Total events:	241		356				₹
Heterogeneity: $Tau^2 = 0.09$	9; Chi ² = 2.4	5, df = 1 (P = 0.12; I	² = 59%			
Test for overall effect: $Z =$	0.60 (P = 0	.55)					
2.49.2 Moderate CVD ris	k						
Ley 2004	1	88	5	88	0.3%	0.20 [0.02, 1.68]	
Moy 2001	1	117	1	118	0.2%	1.01 [0.06, 15.93]	
Subtotal (95% CI)		205		206	0.4%	0.37 [0.07 , 1.97	
Total events:	2		6				
Heterogeneity: $Tau^2 = 0.00$); $Chi^2 = 0.8$	4, df = 1 (P = 0.36; I	$^{2} = 0\%$			
Test for overall effect: Z =	1.17 (P = 0)	.24)					
2.49.3 Existing CVD dise	ase						
Oslo Diet-Heart 1966	2	206	1	206	0.2%	2.00 [0.18, 21.89]]
STARS 1992	0	27	1	28	0.1%	0.35 [0.01 , 8.12]]
WHI 2006	206	908	308	1369	51.9%	1.01 [0.86 , 1.18]] 🖌
Subtotal (95% CI)		1141		1603	52.2%	1.01 [0.86 , 1.18]	1 4
Total events:	208		310				
Heterogeneity: $Tau^2 = 0.00$); $Chi^2 = 0.7$	6, df = 2 (P = 0.68; I	$^{2} = 0\%$			
Test for overall effect: Z =	0.11 (P = 0	.91)					
Total (95% CI)		20403		30156	100.0%	1.00 [0.89 , 1.11]	1
Total events:	451		672			- ,	-
Heterogeneity: $Tau^2 = 0.00$); Chi ² = 5.4	4, df = 6 (P = 0.49; I	$^{2} = 0\%$			0.005 0.1 1 10 200
Test for overall effect: Z =	0.08 (P = 0)	.94)]	Favours lower SFA Favours higher SFA
Test for subgroup differen	ces: Chi ² = 1	1.71, df = 2	2 (P = 0.42)), $I^2 = 0\%$			

Footnotes

(1) Women without CVD at baseline



Analysis 2.50. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 50: Stroke, subgroup by TC reduction

	Lower	SFA	Higher	SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
2.50.1 serum chol reduc	ed by at leas	t 0.2mmo	I/L				
MRC 1968	2	199	0	194	4.0%	4.88 [0.24, 100.89]	
Moy 2001	1	117	1	118	4.8%	1.01 [0.06 , 15.93]	
Oslo Diet-Heart 1966	2	206	1	206	6.4%	2.00 [0.18, 21.89]	
STARS 1992	0	27	1	28	3.7%	0.35 [0.01, 8.12]	
Veterans Admin 1969	13	424	22	422	81.1%	0.59 [0.30, 1.15]	-
Subtotal (95% CI)		973		968	100.0%	0.70 [0.38, 1.28]	
Total events:	18		25				•
Heterogeneity: $Tau^2 = 0.0$	00; $Chi^2 = 2.8$	35, df = 4 (P = 0.58; 1	$^{2} = 0\%$			
Test for overall effect: Z	= 1.17 (P = 0)	.24)					
2.50.2 serum chol reduc	ed by <0.2m	mol/L					
Ley 2004	1	88	5	88	28.1%	0.20 [0.02, 1.68]	
WHI 2006 (1)	435	19541	634	29294	71.9%	1.03 [0.91 , 1.16]	•
Subtotal (95% CI)		19629		29382	100.0%	0.65 [0.15 , 2.75]	
Total events:	436		639				
Heterogeneity: $Tau^2 = 0.7$	75; Chi ² = 2.2	27, df = 1 (P = 0.13; I	$^{2} = 56\%$			
Test for overall effect: Z	= 0.59 (P = 0)	.56)					
2.50.3 serum chol reduc	tion unclear						
Subtotal (95% CI)		0		0		Not estimable	
Total events:	0		0				
Heterogeneity: Not appli	cable						
Test for overall effect: N	ot applicable						
Test for subgroup differe	nces: Chi ² = 0	0.01, df =	1 (P = 0.93)), $I^2 = 0\%$		F	0.005 0.1 1 10 200 avours lower SFA Favours higher SF

Footnotes

(1) During trial, Prentice 2017

Analysis 2.51. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 51: Stroke, subgroup decade of publication

	Lower	SFA	Higher	SFA		Risk Ratio	Risk I	Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Rando	om, 95% CI
2.51.1 1960s								
MRC 1968	2	199	0	194	1.0%	4.88 [0.24, 100.89]		
Oslo Diet-Heart 1966	2	206	1	206	1.6%	2.00 [0.18 , 21.89]		•
Veterans Admin 1969	13	424	22	422	16.4%	0.59 [0.30 , 1.15]		
Subtotal (95% CI)		829		822	19.0%	0.92 [0.31 , 2.69]		
Total events:	17		23					
Heterogeneity: $Tau^2 = 0.30$); $Chi^2 = 2.6$	0, $df = 2$ (P = 0.27; I	$^{2} = 23\%$				
Test for overall effect: Z =	0.16 (P = 0.16)	.87)						
2.51.2 1970s								
Subtotal (95% CI)		0		0		Not estimable	2	
Total events:	0		0					
Heterogeneity: Not applica	able							
Test for overall effect: Not	t applicable							
2.51.3 1980s								
Subtotal (95% CI)		0		0		Not estimable	e	
Total events:	0		0					
Heterogeneity: Not applica	able							
Test for overall effect: Not	t applicable							
2.51.4 1990s								
STARS 1992	0	27	1	28	0.9%	0.35 [0.01, 8.12]	I	
Subtotal (95% CI)		27		28	0.9%	0.35 [0.01, 8.12]		
Total events:	0		1					
Heterogeneity: Not applica	able							
Test for overall effect: Z =	0.66 (P = 0)	.51)						
2.51.5 2000s								
Ley 2004	1	88	5	88	2.0%	0.20 [0.02 , 1.68]		_
Moy 2001	1	117	1	118	1.2%	1.01 [0.06 , 15.93]]	
WHI 2006 (1)	435	19541	634	29294	76.9%	1.03 [0.91 , 1.16]		
Subtotal (95% CI)		19746		29500	80.1%	0.91 [0.50 , 1.66]		
Total events:	437		640					
Heterogeneity: $Tau^2 = 0.10$); $Chi^2 = 2.2$	7, $df = 2$ (P = 0.32; I	² = 12%				
Test for overall effect: Z =	0.30 (P = 0.0)	.76)						
Total (95% CI)		20602		30350	100.0%	0.92 [0.68 , 1.25]		,
Total events:	454		664				Ţ	
Heterogeneity: $Tau^2 = 0.03$	3; Chi ² = 6.5	9, df = 6 (P = 0.36; I	$^{2} = 9\%$			0.005 0.1 1	10 200
Test for overall effect: $Z =$	0.52 (P = 0.52)	.60)				I	Favours lower SFA	Favours higher SFA
Test for subgroup differen	ces: Chi ² = 0	0.35, df = 1	2 (P = 0.84)	$I^2 = 0\%$				c

Footnotes

(1) During trial, Prentice 2017



Analysis 2.52. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 52: CORONARY HEART DISEASE MORTALITY

	Lower	SFA	Higher	SFA		Risk Ratio	Risk Ratio		
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random	, 95% CI	
DART 1989	97	1018	97	1015	21.9%	1.00 [0.76 , 1.30]			
Houtsmuller 1979	0	51	5	51	0.4%	0.09 [0.01 , 1.60]	←		
MRC 1968	25	199	25	194	9.1%	0.97 [0.58, 1.64]	· _		
Oslo Diet-Heart 1966	37	206	50	206	14.5%	0.74 [0.51, 1.08]			
Rose corn oil 1965	5	28	1	13	0.7%	2.32 [0.30, 17.92]			
Rose olive 1965	3	26	0	13	0.4%	3.63 [0.20, 65.44]		_ _	
Sydney Diet-Heart 1978	35	221	23	237	9.9%	1.63 [1.00 , 2.67]	_	_	
Veterans Admin 1969	41	424	50	422	13.9%	0.82 [0.55, 1.21]			
WHI 2006 (1)	172	19541	261	29294	29.2%	0.99 [0.82 , 1.20]	+		
Total (95% CI)		21714		31445	100.0%	0.97 [0.82 , 1.16]			
Total events:	415		512				The second se		
Heterogeneity: $Tau^2 = 0.02$, Chi ² = 11.18	, df = 8 (P	= 0.19; I ²	= 28%				10 100	
Test for overall effect: $Z = 0.32$ (P = 0.75)					Fa	avours lower SFA	Favours higher SFA		
Test for subgroup differenc	es: Not applic	able							

Footnotes

(1) CHD death during trial, Prentice 2017

Analysis 2.53. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 53: CHD mortality, SA by low summary risk of bias

	Lower	SFA	Higher	SFA		Risk Ratio	Ris	k Ratio	
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Ran	dom, 95% CI	
Sydney Diet-Heart 1978	35	221	23	237	23.3%	1.63 [1.00 , 2.67]		
Veterans Admin 1969	41	424	50	422	29.9%	0.82 [0.55, 1.21] -	•	
WHI 2006 (1)	172	19541	261	29294	46.7%	0.99 [0.82 , 1.20]	•	
Total (95% CI)		20186		29953	100.0%	1.05 [0.77 , 1.43]		
Total events:	248		334					Ť	
Heterogeneity: $Tau^2 = 0.04$; Chi ² = 4.83,	df = 2 (P = 2)	= 0.09); I ² =	= 59%			0.01 0.1	1 10	100
Test for overall effect: Z =	0.30 (P = 0.76)	5)					Favours lower SFA	Favours h	igher SFA
Test for subgroup differenc	es: Not applie	able							

Footnotes

(1) CHD death during trial, Prentice 2017



Analysis 2.54. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 54: CHD mortality, SA aim to reduce SFA

	Lower	SFA	Higher	SFA		Risk Ratio	Risk 1	Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Rando	om, 95% CI
DART 1989	97	1018	97	1015	21.9%	1.00 [0.76 , 1.30]	-	+
Houtsmuller 1979	0	51	5	51	0.4%	0.09 [0.01 , 1.60]	←	_
MRC 1968	25	199	25	194	9.1%	0.97 [0.58, 1.64]	_	_
Oslo Diet-Heart 1966	37	206	50	206	14.5%	0.74 [0.51, 1.08]	-	
Rose corn oil 1965	5	28	1	13	0.7%	2.32 [0.30 , 17.92]		_ .
Rose olive 1965	3	26	0	13	0.4%	3.63 [0.20, 65.44]		
Sydney Diet-Heart 1978	35	221	23	237	9.9%	1.63 [1.00 , 2.67]		
Veterans Admin 1969	41	424	50	422	13.9%	0.82 [0.55, 1.21]		
WHI 2006 (1)	172	19541	261	29294	29.2%	0.99 [0.82 , 1.20]	•	E
Total (95% CI)		21714		31445	100.0%	0.97 [0.82 , 1.16]		,
Total events:	415		512					
Heterogeneity: $Tau^2 = 0.02$;	Chi ² = 11.18	, df = 8 (P)	= 0.19); I ²	= 28%			0.01 0.1 1	10 100
Test for overall effect: $Z = 0.32$ (P = 0.75)					Fa	wours lower SFA	Favours higher SFA	
Test for subgroup difference	es: Not applic	able						

Footnotes

(1) CHD death during trial, Prentice 2017

Analysis 2.55. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 55: CHD mortality, SA statistically significant SFA reduction

	Lower	SFA	Higher SFA			Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
DART 1989	97	1018	97	1015	29.3%	1.00 [0.76 , 1.30]	•
Sydney Diet-Heart 1978	35	221	23	237	12.5%	1.63 [1.00 , 2.67]	
Veterans Admin 1969	41	424	50	422	18.0%	0.82 [0.55, 1.21]	
WHI 2006 (1)	172	19541	261	29294	40.3%	0.99 [0.82 , 1.20]	•
Total (95% CI)		21204		30968	100.0%	1.02 [0.84 , 1.24]	•
Total events:	345		431				
Heterogeneity: Tau ² = 0.01	; Chi ² = 4.84,	df = 3 (P :	= 0.18); I ² =	= 38%		0	0.01 0.1 1 10 100
Test for overall effect: Z =	0.19 (P = 0.85)	5)	Fav	yours lower SFA Favours higher SFA			

Test for subgroup differences: Not applicable

Footnotes

(1) CHD death during trial, Prentice 2017



Analysis 2.56. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 56: CHD mortality, SA by TC reduction

	Lower	SFA	Higher	SFA		Risk Ratio	Risk Ratio	
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI	
DART 1989	97	1018	97	1015	21.9%	1.00 [0.76 , 1.30]	+	
Houtsmuller 1979	0	51	5	51	0.4%	0.09 [0.01 , 1.60]	←	
MRC 1968	25	199	25	194	9.3%	0.97 [0.58, 1.64]		
Oslo Diet-Heart 1966	37	206	50	206	14.7%	0.74 [0.51, 1.08]		
Rose corn oil 1965	5	28	1	13	0.7%	2.32 [0.30, 17.92]		
Sydney Diet-Heart 1978	35	221	23	237	10.0%	1.63 [1.00 , 2.67]		
Veterans Admin 1969	41	424	50	422	14.1%	0.82 [0.55, 1.21]		
WHI 2006 (1)	172	19541	261	29294	28.9%	0.99 [0.82 , 1.20]	•	
Total (95% CI)		21688		31432	100.0%	0.97 [0.81 , 1.16]	•	
Total events:	412		512				Ť	
Heterogeneity: Tau ² = 0.02;	; Chi ² = 10.38	, df = 7 (P)	P = 0.17; I ²	= 33%			0.01 0.1 1 10	100
Test for overall effect: $Z = 0$	0.36 (P = 0.72)	2)				F	avours lower SFA Favours hi	gher SFA

Test for subgroup differences: Not applicable

Footnotes

(1) CHD death during trial, Prentice 2017

Analysis 2.57. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 57: CHD mortality, SA excluding WHI

	Lower	SFA	Higher	SFA		Risk Ratio	Risk	Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Rand	om, 95% CI
DART 1989	97	1018	97	1015	27.0%	1.00 [0.76 , 1.30]	4	•
Houtsmuller 1979	0	51	5	51	0.7%	0.09 [0.01 , 1.60]	←	L
MRC 1968	25	199	25	194	14.4%	0.97 [0.58, 1.64]	-	-
Oslo Diet-Heart 1966	37	206	50	206	20.5%	0.74 [0.51, 1.08]	-	ļ
Rose corn oil 1965	5	28	1	13	1.4%	2.32 [0.30, 17.92]		
Rose olive 1965	3	26	0	13	0.7%	3.63 [0.20, 65.44]		
Sydney Diet-Heart 1978	35	221	23	237	15.3%	1.63 [1.00 , 2.67]		
Veterans Admin 1969	41	424	50	422	19.9%	0.82 [0.55 , 1.21]	-	+
Total (95% CI)		2173		2151	100.0%	0.97 [0.76 , 1.24]		
Total events:	243		251					
Heterogeneity: $Tau^2 = 0.04$; Chi ² = 11.13	df = 7 (P	= 0.13; I ²	= 37%			0.01 0.1	
Test for overall effect: $Z = 0$	0.22 (P = 0.82)	2)				F	Favours lower SFA	Favours higher SFA

Test for subgroup differences: Not applicable



Analysis 2.58. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 58: CHD mortality, SA Mantel-Haenszel fixed-effect

	Lower	SFA	Higher	SFA		Risk Ratio	Ris	k Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Fixed, 95% CI	M-H, Fi	xed, 95% CI
DART 1989	97	1018	97	1015	21.1%	1.00 [0.76 , 1.30]]	•
Houtsmuller 1979	0	51	5	51	1.2%	0.09 [0.01 , 1.60]] ←	
MRC 1968	25	199	25	194	5.5%	0.97 [0.58 , 1.64]]	_
Oslo Diet-Heart 1966	37	206	50	206	10.8%	0.74 [0.51, 1.08]] -	-
Rose corn oil 1965	5	28	1	13	0.3%	2.32 [0.30 , 17.92]	I	
Rose olive 1965	3	26	0	13	0.1%	3.63 [0.20, 65.44]]	
Sydney Diet-Heart 1978	35	221	23	237	4.8%	1.63 [1.00 , 2.67]]	
Veterans Admin 1969	41	424	50	422	10.9%	0.82 [0.55, 1.21]] .	-
WHI 2006 (1)	172	19541	261	29294	45.3%	0.99 [0.82 , 1.20]]	•
Total (95% CI)		21714		31445	100.0%	0.97 [0.86 , 1.10]]	
Total events:	415		512					
Heterogeneity: Chi ² = 11.18	8, df = 8 (P =	0.19); I ² =	28%				0.01 0.1	1 10 100
Test for overall effect: Z =	5)				I	Favours lower SFA	Favours higher SFA	
Test for subgroup differenc	es: Not applic	able						

Footnotes

(1) CHD death during trial, Prentice 2017

health events, Outcome 59: CHD mortality, SA Peto fixed-effect											
Lower SFA Higher SFA Peto Odds Ratio Peto Odds Ratio											
Study or Subgroup	Events	Total	Events	Total	Weight	Peto, Fixed, 95% CI	Ре	to, Fixed, 95% CI			
DART 1989	97	1018	97	1015	20.8%	1.00 [0.74 , 1.34]	+			
Houtsmuller 1979	0	51	5	51	0.6%	0.12 [0.02, 0.75]				
MRC 1968	25	199	25	194	5.2%	0.97 [0.54 , 1.76]	_			
Oslo Diet-Heart 1966	37	206	50	206	8.2%	0.69 [0.43 , 1.10]				
Rose corn oil 1965	5	28	1	13	0.5%	2.21 [0.35 , 13.90]				
Rose olive 1965	3	26	0	13	0.3%	4.87 [0.41 , 57.37]				
Sydney Diet-Heart 1978	35	221	23	237	6.0%	1.74 [1.00 , 3.02]				
Veterans Admin 1969	41	424	50	422	9.6%	0.80 [0.52 , 1.23]				
WHI 2006 (1)	172	19541	261	29294	48.8%	0.99 [0.81 , 1.20]	•			
Total (95% CI)		21714		31445	100.0%	0.97 [0.85 , 1.11]				
Total events:	415		512								
Heterogeneity: $Chi^2 = 14.72$	2, $df = 8 (P =$	0.06 ; $I^2 =$	46%				0.01 0.1	1 10	100		
Test for overall effect: $Z = 0$	0.44 (P = 0.66)	5)					Favours lower	SFA Favours	higher SFA		

Analysis 2.59. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 59: CHD mortality, SA Peto fixed-effect

Test for subgroup differences: Not applicable

Footnotes

(1) CHD death during trial, Prentice 2017

Analysis 2.60. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 60: CHD mortality, subgroup by any substitution

	Lower	SFA	Higher	SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
2.60.1 replaced by PUFA							
DART 1989	97	1018	97	1015	25.6%	1.00 [0.76 , 1.30]	_
Houtsmuller 1979	0	51	5	51	0.9%	0.09 [0.01, 1.60]	<u> </u>
MRC 1968	25	199	25	194	15.2%	0.97 [0.58 , 1.64]	
Oslo Diet-Heart 1966	37	206	50	206	20.5%	0.74 [0.51, 1.08]	
Rose corn oil 1965	5	28	1	26	1.6%	4.64 [0.58 . 37.15]	
Sydney Diet-Heart 1978	35	221	23	237	16.1%	1.63 [1.00 . 2.67]	
Veterans Admin 1969	41	424	50	422	20.0%	0.82 [0.55 , 1.21]	
Subtotal (95% CI)		2147	20	2151	100.0%	0.98 [0.74 , 1.28]	1
Total events:	240	214/	251	2101	100.0 /0	0.50 [0.74 ; 1.20]	T
Heterogeneity: Tau ² – 0.06: ($Chi^2 = 11.81$	df = 6 (P	$= 0.07) \cdot 12$	- 10%			
Test for overall effect: $Z = 0$.	.18 (P = 0.86)	, ui = 0 (r 5)	= 0.07), I [_]	- 4970			
		,					
2.60.2 replaced by MUFA							
Rose olive 1965	3	26	1	26	100.0%	3.00 [0.33 , 26.99]	
Subtotal (95% CI)		26		26	100.0%	3.00 [0.33 , 26.99]	
Total events:	3		1				
Heterogeneity: Not applicabl	e						
Test for overall effect: $Z = 0$.	.98 ($P = 0.33$	3)					
2.60.2 replaced by CUO							
2.00.3 replaced by CHO	07	1010	07	1015	22.00	1 00 [0 76 1 20]	
DART 1989	97	1018	97	20204	55.9%	1.00 [0.76 , 1.30]	<u>+</u>
WHI 2006 (1)	172	19541	201	29294	00.1%	0.99 [0.82 , 1.20]	
Subtotal (95% CI)	2(0)	20559	250	30309	100.0%	0.99 [0.85 , 1.16]	•
Total events:	269		358				
Heterogeneity: $Tau^2 = 0.00$; ($Chi^2 = 0.00,$	df = 1 (P =	= 0.96); I ² =	= 0%			
Test for overall effect: $Z = 0$.	.11 (P = 0.91)					
2.60.4 replaced by protein							
DART 1989	97	1018	97	1015	33.9%	1.00 [0.76 . 1.30]	1
WHI 2006 (1)	172	19541	261	29294	66.1%	0.99 [0.82 1.20]	I
Subtotal (95% CI)		20559	201	30309	100.0%	0.99 [0.85 , 1.16]	T
Total events:	269	-0000	358	00003	100.0 /0		Y
Heterogeneity: $T_{202} = 0.00$:	$Chi^2 = 0.00$	df = 1 (P -	- 0.06)· 12 -	- 0%			
Test for overall effect: $Z = 0.00$,	.11 (P = 0.91)	u = 1 (1 -	- 0.90), 1 -	- 0 /0			
2.60.5 replacement unclear		Δ		Δ		Not optimable	
Subtotal (95% CI)	0	U	0	U		not estimable	
i otai events:	0		0				
Heterogeneity: Not applicabl	e						
Test for overall effect: Not a	pplicable						
Test for subgroup differences	s: $Chi^2 = 0.9$	9. $df = 3.0$	P = 0.80) I	$^{2} = 0\%$, F	
subgroup unrefered		, – 5 (0.10		Favo	urs lower SFA Favours higher SFA
Footnotes							-

(1) CHD death during trial, Prentice 2017



Analysis 2.61. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 61: CHD mortality, subgroup by main substitution

	Lower	SFA	Higher	SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
2.61.1 replaced by PUFA							
DART 1989	97	1018	97	1015	25.6%	1.00 [0.76 , 1.30]	+
Houtsmuller 1979	0	51	5	51	0.9%	0.09 [0.01 , 1.60]	←
MRC 1968	25	199	25	194	15.2%	0.97 [0.58, 1.64]	
Oslo Diet-Heart 1966	37	206	50	206	20.5%	0.74 [0.51, 1.08]	
Rose corn oil 1965	5	28	1	26	1.6%	4.64 [0.58, 37.15]	
Sydney Diet-Heart 1978	35	221	23	237	16.1%	1.63 [1.00 , 2.67]	
Veterans Admin 1969	41	424	50	422	20.0%	0.82 [0.55 , 1.21]	
Subtotal (95% CI)		2147		2151	100.0%	0.98 [0.74 , 1.28]	•
Total events:	240		251				Ţ
Heterogeneity: $Tau^2 = 0.06$; C	$2hi^2 = 11.81$	df = 6 (P	= 0.07; I ²	= 49%			
Test for overall effect: $Z = 0.1$	18 (P = 0.86))					
2.61.2 replaced by MUFA							
Rose olive 1965	3	26	1	26	100.0%	3.00 [0.33 , 26.99]	
Subtotal (95% CI)		26		26	100.0%	3.00 [0.33 , 26.99]	
Total events:	3		1			- / -	
Heterogeneity: Not applicable	,						
Test for overall effect: $Z = 0.9$	P = 0.33)					
2.61.3 replaced by CHO							
WHI 2006 (1)	172	19541	261	29294	100.0%	0.99 [0.82, 1.20]	•
Subtotal (95% CI)		19541		29294	100.0%	0.99 [0.82, 1.20]	•
Total events:	172		261				Ĭ
Heterogeneity: Not applicable	•						
Test for overall effect: $Z = 0.1$	12 (P = 0.90))					
2.61.4 replaced by protein							
Subtotal (95% CI)		0		0		Not estimable	
Total events:	0		0				
Heterogeneity: Not applicable	•						
Test for overall effect: Not ap	plicable						
2.61.5 replacement unclear							
Subtotal (95% CI)		0		0		Not estimable	
Total events:	0		0				
Heterogeneity: Not applicable							
Test for overall effect: Not ap	plicable						
Test for subgroup differences	: $Chi^2 = 0.99$	9, $df = 2$ ($P = 0.61), I^2$	$^{2} = 0\%$			0.01 0.1 1 10 100
_						Fa	avours lower SFA Favours higher SFA
Footnotes							-

(1) CHD death during trial, Prentice 2017

Analysis 2.62. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 62: CHD mortality, subgroup by duration

	Lower	Lower SFA		Higher SFA		Risk Ratio	Risk Ratio		
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random,	95% CI	
2.62.1 up to 24mo									
DART 1989	97	1018	97	1015	97.5%	1.00 [0.76 , 1.30]			
Rose corn oil 1965	5	28	1	13	1.7%	2.32 [0.30, 17.92]			
Rose olive 1965	3	26	0	13	0.8%	3.63 [0.20, 65.44]		-	
Subtotal (95% CI)		1072		1041	100.0%	1.02 [0.78 , 1.33]	•		
Total events:	105		98				Ť		
Heterogeneity: $Tau^2 = 0.00$; Chi ² = 1.40,	df = 2 (P = 2)	= 0.50); I ² =	= 0%					
Test for overall effect: Z =	0.16 (P = 0.87)	7)							
2.62.2 >24 to 48months									
MRC 1968	25	199	25	194	36.2%	0.97 [0.58, 1.64]			
Veterans Admin 1969	41	424	50	422	63.8%	0.82 [0.55, 1.21]	-		
Subtotal (95% CI)		623		616	100.0%	0.87 [0.64 , 1.19]	→		
Total events:	66		75				•		
Heterogeneity: Tau ² = 0.00	; Chi ² = 0.29,	df = 1 (P = 1)	= 0.59); I ² =	= 0%					
Test for overall effect: Z =	0.87 (P = 0.38	3)							
2.62.3 >48 months									
Oslo Diet-Heart 1966	37	206	50	206	31.5%	0.74 [0.51, 1.08]			
Sydney Diet-Heart 1978	35	221	23	237	25.1%	1.63 [1.00 , 2.67]	-		
WHI 2006 (1)	172	19541	261	29294	43.4%	0.99 [0.82, 1.20]	•		
Subtotal (95% CI)		19968		29737	100.0%	1.02 [0.72 , 1.45]	•		
Total events:	244		334				Ť		
Heterogeneity: $Tau^2 = 0.06$; Chi ² = 6.22,	df = 2 (P = 2)	= 0.04); I ² =	= 68%					
Test for overall effect: Z =	0.13 (P = 0.90))							
2.62.4 unclear duration									
Houtsmuller 1979	0	51	5	51	100.0%	0.09 [0.01 , 1.60]			
Subtotal (95% CI)		51		51	100.0%	0.09 [0.01 , 1.60]			
Total events:	0		5						
Heterogeneity: Not applica	ble								
Test for overall effect: Z =	1.64 (P = 0.10)))							
Test for subgroup difference	ces: Chi ² = 3.3	1, df = 3 (P = 0.35), I	² = 9.4%			0.01 0.1 1	10 100	
						Fa	avours lower SFA	Favours higher SFA	
Footnotes									

(1) CHD death during trial, Prentice 2017



Analysis 2.63. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 63: CHD mortality, subgroup by baseline SFA

	Lower	Lower SFA		Higher SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
2.63.1 up to 12%E SFA ba	seline						
Subtotal (95% CI)		0		0		Not estimable	
Total events:	0		0				
Heterogeneity: Not applicab	ole						
Test for overall effect: Not a	applicable						
2.63.2 >12% to 15% E SFA	baseline						
DART 1989	97	1018	97	1015	35.9%	1.00 [0.76 , 1.30]	_
Sydney Diet-Heart 1978	35	221	23	237	15.9%	1.63 [1.00 , 2.67]	
WHI 2006 (1)	172	19541	261	29294	48.2%	0.99 [0.82, 1.20]	
Subtotal (95% CI)		20780		30546	100.0%	1.07 [0.86 , 1.34]	T
Total events:	304		381			- , -	
Heterogeneity: $Tau^2 = 0.02$;	$Chi^2 = 3.58,$	df = 2 (P =	= 0.17); I ² =	- 44%			
Test for overall effect: $Z = 0$	0.63 (P = 0.53)	3)					
2.63.3 >15 to 18%E SFA b	aseline						
Subtotal (95% CI)		0		0		Not estimable	
Total events:	0		0				
Heterogeneity: Not applicab	ole						
Test for overall effect: Not a	applicable						
2.63.4 >18%E SFA baselir	ie						
Veterans Admin 1969	41	424	50	422	100.0%	0.82 [0.55 , 1.21]	
Subtotal (95% CI)		424		422	100.0%	0.82 [0.55 , 1.21]	
Total events:	41		50				•
Heterogeneity: Not applicab	ole						
Test for overall effect: $Z = 1$.02 (P = 0.31))					
2.63.5 unclear							
Houtsmuller 1979	0	51	5	51	2.0%	0.09 [0.01 , 1.60]	←
MRC 1968	25	199	25	194	38.5%	0.97 [0.58, 1.64]	·
Oslo Diet-Heart 1966	37	206	50	206	53.5%	0.74 [0.51, 1.08]	-
Rose corn oil 1965	5	28	1	13	3.9%	2.32 [0.30, 17.92]	
Rose olive 1965	3	26	0	13	2.0%	3.63 [0.20, 65.44]	
Subtotal (95% CI)		510		477	100.0%	0.85 [0.56 , 1.29]	
Total events:	70		81				T
Heterogeneity: $Tau^2 = 0.05$;	$Chi^2 = 4.98,$	df = 4 (P =	= 0.29); I ² =	= 20%			
Test for overall effect: $Z = 0$	0.76 (P = 0.45)	5)					
Test for subgroup difference	es: Chi ² = 1.9	4, df = 2 (P = 0.38), I	$^{2} = 0\%$		0.	
T						Fav	ours lower SFA Favours higher SFA
Footnotes							

(1) CHD death during trial, Prentice 2017

Analysis 2.64. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 64: CHD mortality, subgroup by SFA change

	Lower	SFA	Higher	SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
2.64.1 up to 4%E difference	;						
DART 1989	97	1018	97	1015	35.9%	1.00 [0.76 , 1.30]	_
Sydney Diet-Heart 1978	35	221	23	237	15.9%	1.63 [1.00, 2.67]	T
WHI 2006 (1)	172	19541	261	29294	48.2%	0.99 [0.82 , 1.20]	
Subtotal (95% CI)		20780		30546	100.0%	1.07 [0.86 , 1.34]	T
Total events:	304		381				
Heterogeneity: $Tau^2 = 0.02$; C	Chi ² = 3.58,	df = 2 (P =	= 0.17); I ² =	44%			
Test for overall effect: $Z = 0.0$	63 (P = 0.53	3)					
2.64.2 >4 to 8%E difference							
Subtotal (95% CI)		0		0		Not estimable	
Total events:	0		0				
Heterogeneity: Not applicable	e						
Test for overall effect: Not ap	plicable						
2.64.3 >8%E difference							
Veterans Admin 1969	41	424	50	422	100.0%	0.82 [0.55, 1.21]	
Subtotal (95% CI)		424		422	100.0%	0.82 [0.55 , 1.21]	
Total events:	41		50				•
Heterogeneity: Not applicable	e						
Test for overall effect: $Z = 1.0$	02 (P = 0.31))					
2.64.4 unclear							
Houtsmuller 1979	0	51	5	51	2.0%	0.09 [0.01 , 1.60]	←
MRC 1968	25	199	25	194	38.5%	0.97 [0.58, 1.64]	-
Oslo Diet-Heart 1966	37	206	50	206	53.5%	0.74 [0.51, 1.08]	-
Rose corn oil 1965	5	28	1	13	3.9%	2.32 [0.30 , 17.92]	
Rose olive 1965	3	26	0	13	2.0%	3.63 [0.20, 65.44]	
Subtotal (95% CI)		510		477	100.0%	0.85 [0.56 , 1.29]	
Total events:	70		81				•
Heterogeneity: Tau ² = 0.05; C	Chi ² = 4.98,	df = 4 (P =	= 0.29); I ² =	20%			
Test for overall effect: $Z = 0.7$	76 ($P = 0.45$	5)					
Test for subgroup differences	: Chi² = 1.9	4, df = 2 (P = 0.38), I	$^{2} = 0\%$		0. Favo	01 0.1 1 10 100 Durs lower SFA Favours higher SFA
Footnotes						Tuv	

(1) CHD death during trial, Prentice 2017



Analysis 2.65. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 65: CHD mortality, subgroup by sex

	Lower	SFA	Higher	SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
2.65.1 Men							
DART 1989	97	1018	97	1015	29.6%	1.00 [0.76 , 1.30]	+
MRC 1968	25	199	25	194	13.6%	0.97 [0.58, 1.64]	
Oslo Diet-Heart 1966	37	206	50	206	20.7%	0.74 [0.51, 1.08]	
Rose corn oil 1965	5	28	1	13	1.2%	2.32 [0.30 , 17.92]	
Rose olive 1965	3	26	0	13	0.6%	3.63 [0.20, 65.44]	
Sydney Diet-Heart 1978	35	221	23	237	14.6%	1.63 [1.00 , 2.67]	
Veterans Admin 1969	41	424	50	422	19.9%	0.82 [0.55, 1.21]	
Subtotal (95% CI)		2122		2100	100.0%	0.98 [0.79 , 1.23]	•
Total events:	243		246				Ĭ
Heterogeneity: Tau ² = 0.02; Cl	hi ² = 8.55,	df = 6 (P =	= 0.20); I ² =	= 30%			
Test for overall effect: $Z = 0.1$	6 (P = 0.87)	7)					
2.65.2 Women							
WHI 2006 (1)	172	19541	261	29294	100.0%	0.99 [0.82, 1.20]	
Subtotal (95% CI)		19541		29294	100.0%	0.99 [0.82 , 1.20]	•
Total events:	172		261				The second secon
Heterogeneity: Not applicable							
Test for overall effect: $Z = 0.1$	2 (P = 0.90))					
2.65.3 Mixed, men and wome	en						
Houtsmuller 1979	0	51	5	51	100.0%	0.09 [0.01 , 1.60]	
Subtotal (95% CI)		51		51	100.0%	0.09 [0.01 , 1.60]	
Total events:	0		5				
Heterogeneity: Not applicable							
Test for overall effect: $Z = 1.6$	4 (P = 0.10)))					
Test for subgroup differences:	Chi ² = 2.6	4, df = 2 (P = 0.27), I	² = 24.4%		Fa	0.01 0.1 1 10 100 avours lower SFA Favours higher SFA

Footnotes

(1) CHD death during trial, Prentice 2017

Analysis 2.66. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 66: CHD mortality, subgroup by CVD risk

	Lower	SFA	Higher	SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
2.66.1 Low CVD risk							
Veterans Admin 1969	41	424	50	422	25.2%	0.82 [0.55, 1.21]	-
WHI 2006 (1)	124	18633	185	27925	74.8%	1.00 [0.80 , 1.26]	•
Subtotal (95% CI)		19057		28347	100.0%	0.95 [0.78 , 1.16]	4
Total events:	165		235				Y
Heterogeneity: $Tau^2 = 0.00$; Chi ² = 0.81,	df = 1 (P = 1)	= 0.37); I ² =	= 0%			
Test for overall effect: $Z =$	0.48 (P = 0.63)	3)					
2.66.2 Moderate CVD risl	ĸ						
Houtsmuller 1979	0	51	5	51	100.0%	0.09 [0.01 , 1.60]	←
Subtotal (95% CI)		51		51	100.0%	0.09 [0.01 , 1.60]	
Total events:	0		5				
Heterogeneity: Not applical	ble						
Test for overall effect: Z =	1.64 (P = 0.10)))					
2.66.3 Existing CVD disea	ise						
DART 1989	97	1018	97	1015	32.2%	1.00 [0.76 , 1.30]	.
MRC 1968	25	199	25	194	13.2%	0.97 [0.58, 1.64]	
Oslo Diet-Heart 1966	37	206	50	206	21.1%	0.74 [0.51, 1.08]	-
Rose corn oil 1965	5	28	1	13	1.0%	2.32 [0.30 , 17.92]	
Rose olive 1965	3	26	0	13	0.5%	3.63 [0.20, 65.44]	
Sydney Diet-Heart 1978	35	221	23	237	14.3%	1.63 [1.00 , 2.67]	- - -
WHI 2006 (2)	34	908	49	1369	17.7%	1.05 [0.68 , 1.61]	+
Subtotal (95% CI)		2606		3047	100.0%	1.03 [0.83 , 1.27]	•
Total events:	236		245				Ĭ
Heterogeneity: $Tau^2 = 0.02$; Chi ² = 7.69,	df = 6 (P =	= 0.26); I ² =	= 22%			
Test for overall effect: Z =	0.24 (P = 0.8)	1)					
Test for subgroup differenc	es: Chi ² = 2.9	0, df = 2 (P = 0.23), I	² = 31.0%		F	0.01 0.1 1 10 100 avours lower SFA Favours higher SFA
Eastwates							

Footnotes

(1) Women without CVD at baseline

(2) Women with CVD at baseline



Analysis 2.67. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 67: CHD mortality, subgroup by TC reduction

	Lower	SFA	Higher	SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
2.67.1 serum chol reduced	l by at least (.2mmol/L					
DART 1989	97	1018	97	1015	27.0%	1.00 [0.76 , 1.30]	+
Houtsmuller 1979	0	51	5	51	0.7%	0.09 [0.01 , 1.60]	←
MRC 1968	25	199	25	194	14.6%	0.97 [0.58, 1.64]	
Oslo Diet-Heart 1966	37	206	50	206	20.6%	0.74 [0.51, 1.08]	
Rose corn oil 1965	5	28	1	13	1.4%	2.32 [0.30, 17.92]	
Sydney Diet-Heart 1978	35	221	23	237	15.5%	1.63 [1.00 , 2.67]	-
Veterans Admin 1969	41	424	50	422	20.0%	0.82 [0.55, 1.21]	
Subtotal (95% CI)		2147		2138	100.0%	0.96 [0.75 , 1.24]	▲
Total events:	240		251				Ţ
Heterogeneity: $Tau^2 = 0.04$; Chi ² = 10.31	, df = 6 (P	= 0.11; I ²	= 42%			
Test for overall effect: $Z = 0$	0.29 (P = 0.77)	7)					
2.67.2 serum chol reduced	l by <0.2mm	ol/L					
Rose olive 1965	3	26	0	13	0.4%	3.63 [0.20, 65.44]	
WHI 2006 (1)	172	19541	261	29294	99.6%	0.99 [0.82, 1.20]	•
Subtotal (95% CI)		19567		29307	100.0%	0.99 [0.82 , 1.20]	T
Total events:	175		261				Ť
Heterogeneity: $Tau^2 = 0.00$; Chi ² = 0.78,	df = 1 (P =	= 0.38); I ² =	= 0%			
Test for overall effect: $Z = 0$	0.07 (P = 0.95)	5)					
2.67.3 serum chol reduction	on unclear						
Subtotal (95% CI)		0		0		Not estimable	
Total events:	0		0				
Heterogeneity: Not applical	ble						
Test for overall effect: Not	applicable						
Test for subgroup differenc	es: Chi ² = 0.0	4, df = 1 (P = 0.85), I	² = 0%		Fe	0.01 0.1 1 10 100 avours lower SEA Eavours higher SEA
Footnotes						17	avours lower SFA F avours llight SFA

(1) CHD death during trial, Prentice 2017

Analysis 2.68. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 68: CHD mortality, subgroup decade of publication

	Lower	SFA	Higher	SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
2.68.1 1960s							
MRC 1968	25	199	25	194	9.1%	0.97 [0.58, 1.64]	
Oslo Diet-Heart 1966	37	206	50	206	14.5%	0.74 [0.51, 1.08]	
Rose corn oil 1965	5	28	1	13	0.7%	2.32 [0.30 , 17.92]	
Rose olive 1965	3	26	0	13	0.4%	3.63 [0.20, 65.44]	
Veterans Admin 1969	41	424	50	422	13.9%	0.82 [0.55, 1.21]	
Subtotal (95% CI)		883		848	38.7%	0.84 [0.66 , 1.06]	
Total events:	111		126				•
Heterogeneity: $Tau^2 = 0.00$; C	$Chi^2 = 2.72,$	df = 4 (P =	= 0.61); I ² =	= 0%			
Test for overall effect: $Z = 1.4$	48 (P = 0.14	4)					
2.68.2 1970s							
Houtsmuller 1979	0	51	5	51	0.4%	0.09 [0.01 , 1.60]	← - - -
Sydney Diet-Heart 1978	35	221	23	237	9.9%	1.63 [1.00 , 2.67]	
Subtotal (95% CI)		272		288	10.2%	0.54 [0.03 , 9.26]	
Total events:	35		28				
Heterogeneity: $Tau^2 = 3.35$; C Test for overall effect: $Z = 0.4$	Chi ² = 4.03, 43 (P = 0.67	df = 1 (P = 7)	= 0.04); I ² =	= 75%			
2.68.3 1980s							
DART 1989	97	1018	97	1015	21.9%	1.00 [0.76 , 1.30]	+
Subtotal (95% CI)		1018		1015	21.9%	1.00 [0.76 , 1.30]	
Total events:	97		97				
Heterogeneity: Not applicable Test for overall effect: $Z = 0.0$	e 02 (P = 0.98	3)					
2.68.4 1990s							
Subtotal (95% CI)		0		0		Not estimable	
Total events:	0		0				
Heterogeneity: Not applicable Test for overall effect: Not ap	e plicable						
2.68.5.2000s							
WHI 2006 (1)	172	19541	261	29294	29.2%	0.99 [0.82 1.20]	1
Subtotal (95% CI)	172	19541	201	29294	29.2%	0.99 [0.82 , 1.20]	I
Total events:	172		261		/0	[0:02 , 1:20]	Ţ
Heterogeneity: Not applicable	, ,		201				
Test for overall effect: $Z = 0$.	12 (P = 0.90))					
Total (95% CI)		21714		31445	100.0%	0.97 [0.82 , 1.16]	
Total events:	415		512				Ţ
Heterogeneity: $Tau^2 = 0.02$; C	Chi ² = 11.18	, df = 8 (P	= 0.19; I ²	= 28%			0.01 0.1 1 10 100
Test for overall effect: $Z = 0.3$	32 (P = 0.75)	5)				Fa	vours lower SFA Favours higher SFA
Test for subgroup differences	: Chi ² = 1.5	4, df = 3 (P = 0.67), I	$^{2} = 0\%$			-

Footnotes

(1) CHD death during trial, Prentice 2017



Analysis 2.69. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 69: CORONARY HEART DISEASE EVENTS

	Lower	Lower SFA		Higher SFA		Risk Ratio	Risk R	atio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Rando	m, 95% CI
DART 1989	132	1018	144	1015	17.5%	0.91 [0.73 , 1.14]	-	
Houtsmuller 1979	8	51	30	51	6.2%	0.27 [0.14, 0.52]		
Ley 2004	5	88	3	88	1.9%	1.67 [0.41 , 6.76]		•——
MRC 1968	50	199	50	194	13.6%	0.97 [0.69 , 1.37]	+	
Moy 2001	3	135	1	132	0.8%	2.93 [0.31 , 27.84]		
Oslo Diet-Heart 1966	64	206	90	206	16.3%	0.71 [0.55, 0.92]	-	
Rose corn oil 1965	12	28	3	13	2.9%	1.86 [0.63 , 5.47]	+	•
Rose olive 1965	9	26	3	13	2.8%	1.50 [0.49 , 4.62]		
STARS 1992	3	27	10	28	2.5%	0.31 [0.10 , 1.01]		
Veterans Admin 1969	60	424	78	422	14.5%	0.77 [0.56 , 1.04]		
WHI 2006 (1)	590	19541	913	29294	21.0%	0.97 [0.87 , 1.07]	+	
Total (95% CI)		21743		31456	100.0%	0.83 [0.68 , 1.01]		
Total events:	936		1325				•	
Heterogeneity: $Tau^2 = 0.0$	$05; Chi^2 = 26.$	44, $df = 10^{-10}$	0 (P = 0.003)	3); $I^2 = 62^6$	%		0.01 0.1 1	10 100
Test for overall effect: Z	= 1.82 (P = 0)	.07)				F	avours lower SFA	Favours higher SFA

Test for subgroup differences: Not applicable

Footnotes

(1) CHD events including CHD death & MI, Prentice 2017

Analysis 2.70. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 70: CHD events, SA by low summary risk of bias

	Lower	Lower SFA		Higher SFA		Risk Ratio	Risk F	Risk Ratio		
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Rando	m, 95% CI		
Ley 2004	5	88	3	88	1.6%	1.67 [0.41 , 6.76]	_	•		
Veterans Admin 1969	60	424	78	422	24.7%	0.77 [0.56 , 1.04]	-			
WHI 2006 (1)	590	19541	913	29294	73.7%	0.97 [0.87 , 1.07]	•	l		
Total (95% CI)		20053		29804	100.0%	0.92 [0.77 , 1.10]				
Total events:	655		994				•			
Heterogeneity: $Tau^2 = 0.0$	1; Chi ² = 2.6	4, df = 2 (P = 0.27; I	$^{2} = 24\%$			0.01 0.1 1	10 100		
Test for overall effect: Z =	= 0.90 (P = 0)	.37)				Fa	vours lower SFA	Favours higher SFA		
Test for subgroup different	ices: Not app	licable								

Footnotes



Analysis 2.71. Comparison 2: SFA reduction vs usual diet secondary health events, Outcome 71: CHD events, SA excluding WHI

	Lower SFA		Higher SFA		Risk Ratio		Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
DART 1989	132	1018	144	1015	20.2%	0.91 [0.73 , 1.14]
Houtsmuller 1979	8	51	30	51	9.0%	0.27 [0.14, 0.52]
Ley 2004	5	88	3	88	3.0%	1.67 [0.41 , 6.76]
MRC 1968	50	199	50	194	16.8%	0.97 [0.69 , 1.37	1 🔺
Moy 2001	3	135	1	132	1.2%	2.93 [0.31 , 27.84]
Oslo Diet-Heart 1966	64	206	90	206	19.2%	0.71 [0.55, 0.92] 🗕
Rose corn oil 1965	12	28	3	13	4.6%	1.86 [0.63 , 5.47]
Rose olive 1965	9	26	3	13	4.3%	1.50 [0.49 , 4.62]
STARS 1992	3	27	10	28	4.0%	0.31 [0.10 , 1.01]
Veterans Admin 1969	60	424	78	422	17.7%	0.77 [0.56 , 1.04]
Total (95% CI)		2202		2162	100.0%	0.80 [0.62 , 1.03	1
Total events:	346		412				•
Heterogeneity: $Tau^2 = 0.0^{\circ}$	7; Chi ² = 22.	03, df = 9	(P = 0.009)); $I^2 = 59\%$	2		0.01 0.1 1 10 100
Test for overall effect: Z =	= 1.72 (P = 0)	.09)					Favours lower SFA Favours higher SFA

Test for subgroup differences: Not applicable

Analysis 2.72. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 72: CHD events, SA statistically significant SFA reduction

	Lower	Lower SFA		Higher SFA		Risk Ratio	Risk Ratio	
Study or Subgroup	Events Total		Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI	
DART 1989	132	1018	144	1015	28.4%	0.91 [0.73 , 1.14]		
Ley 2004	5	88	3	88	1.3%	1.67 [0.41 , 6.76]		
Moy 2001	3	135	1	132	0.5%	2.93 [0.31 , 27.84]		
STARS 1992	3	27	10	28	1.8%	0.31 [0.10, 1.01]		
Veterans Admin 1969	60	424	78	422	18.5%	0.77 [0.56 , 1.04]		
WHI 2006 (1)	590	19541	913	29294	49.6%	0.97 [0.87 , 1.07]	•	
Total (95% CI)		21233		30979	100.0%	0.91 [0.77 , 1.06]		
Total events:	793		1149					
Heterogeneity: $Tau^2 = 0.01$	1; Chi ² = 7.1	2, $df = 5$ (P = 0.21; I	$^{2} = 30\%$			0.01 0.1 1 10 10	0
Test for overall effect: $Z = 1.23$ (P = 0.22)				I	Favours lower SFA Favours higher S	SFA		
Test for subgroup differen	ces: Not app	olicable						

Footnotes



Analysis 2.73. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 73: CHD events, SA by TC reduction

	Lower	Lower SFA		Higher SFA		Risk Ratio	Risk Ra	itio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Randon	a, 95% CI
DART 1989	132	1018	144	1015	18.3%	0.91 [0.73 , 1.14]		
Houtsmuller 1979	8	51	30	51	6.7%	0.27 [0.14, 0.52]		
MRC 1968	50	199	50	194	14.3%	0.97 [0.69 , 1.37]	∣ ∔	
Moy 2001	3	135	1	132	0.8%	2.93 [0.31 , 27.84]		_ _
Oslo Diet-Heart 1966	64	206	90	206	17.0%	0.71 [0.55, 0.92]	-	
Rose corn oil 1965	12	28	3	13	3.2%	1.86 [0.63 , 5.47]		<u> </u>
STARS 1992	3	27	10	28	2.7%	0.31 [0.10 , 1.01]	·	
Veterans Admin 1969	60	424	78	422	15.2%	0.77 [0.56 , 1.04]	-	
WHI 2006 (1)	590	19541	913	29294	21.8%	0.97 [0.87 , 1.07]	•	
Total (95% CI)		21629		31355	100.0%	0.80 [0.65 , 0.99]		
Total events:	922		1319				•	
Heterogeneity: $Tau^2 = 0.05$; $Chi^2 = 24.92$, $df = 8$ (P = 0.002); $I^2 = 68\%$							0.01 0.1 1	10 100
Test for overall effect: $Z = 2.05 (P = 0.04)$			I	Favours lower SFA	Favours higher SFA			
Test for subgroup differen	nces: Not app	olicable						

Footnotes

(1) CHD events including CHD death & MI, Prentice 2017

Analysis 2.74. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 74: CHD events, SA aim to reduce SFA

	Lower	SFA	Higher	Higher SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
DART 1989	132	1018	144	1015	17.8%	0.91 [0.73 , 1.14]	
Houtsmuller 1979	8	51	30	51	6.4%	0.27 [0.14, 0.52]	
MRC 1968	50	199	50	194	13.8%	0.97 [0.69 , 1.37]	↓
Moy 2001	3	135	1	132	0.8%	2.93 [0.31, 27.84]	
Oslo Diet-Heart 1966	64	206	90	206	16.6%	0.71 [0.55, 0.92]	-
Rose corn oil 1965	12	28	3	13	3.0%	1.86 [0.63 , 5.47]	
Rose olive 1965	9	26	3	13	2.8%	1.50 [0.49 , 4.62]	
STARS 1992	3	27	10	28	2.6%	0.31 [0.10 , 1.01]	
Veterans Admin 1969	60	424	78	422	14.8%	0.77 [0.56 , 1.04]	-
WHI 2006 (1)	590	19541	913	29294	21.3%	0.97 [0.87 , 1.07]	•
Total (95% CI)		21655		31368	100.0%	0.82 [0.67 , 1.00]	
Total events:	931		1322				•
Heterogeneity: $Tau^2 = 0.0$	05; Chi ² = 25.	71, df = 9	(P = 0.002)); $I^2 = 65\%$,		0.01 0.1 1 10 100
Test for overall effect: Z	= 1.92 (P = 0)	.05)]	Favours lower SFA Favours higher SFA
T 1 1100							

Test for subgroup differences: Not applicable

Footnotes



Analysis 2.75. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 75: CHD events, SA Mantel-Haenszel fixed-effect

	Lower	Lower SFA		Higher SFA		Risk Ratio	Risk	x Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Fixed, 95% CI	M-H, Fix	ed, 95% CI
DART 1989	132	1018	144	1015	12.6%	0.91 [0.73 , 1.14]	•
Houtsmuller 1979	8	51	30	51	2.6%	0.27 [0.14 , 0.52]	
Ley 2004	5	88	3	88	0.3%	1.67 [0.41 , 6.76]	
MRC 1968	50	199	50	194	4.4%	0.97 [0.69 , 1.37] .	+
Moy 2001	3	135	1	132	0.1%	2.93 [0.31 , 27.84]	
Oslo Diet-Heart 1966	64	206	90	206	7.9%	0.71 [0.55 , 0.92] -	•
Rose corn oil 1965	12	28	3	13	0.4%	1.86 [0.63 , 5.47] .	
Rose olive 1965	9	26	3	13	0.3%	1.50 [0.49 , 4.62] _	
STARS 1992	3	27	10	28	0.9%	0.31 [0.10 , 1.01]	_
Veterans Admin 1969	60	424	78	422	6.8%	0.77 [0.56 , 1.04] –	•
WHI 2006 (1)	590	19541	913	29294	63.8%	0.97 [0.87 , 1.07]	•
Total (95% CI)		21743		31456	100.0%	0.91 [0.84 , 0.99]	
Total events:	936		1325					
Heterogeneity: $Chi^2 = 26$.44, df = 10 (P = 0.003)	; I ² = 62%				0.01 0.1	1 10 100
Test for overall effect: $Z = 2.24$ (P = 0.03)						Favours lower SFA	Favours higher SFA	
Test for subgroup differe	nces: Not app	olicable						

Footnotes

(1) CHD events including CHD death & MI, Prentice 2017

	Lower	SFA	Higher	Higher SFA		Peto Odds Ratio		Peto Odds Ratio		
Study or Subgroup	Events	Total	Events	Total	Weight	Peto, Fixed, 95% CI		Peto, Fixed	,95% CI	
DART 1989	132	1018	144	1015	12.0%	0.90 [0.70 , 1.16]		-		
Houtsmuller 1979	8	51	30	51	1.2%	0.16 [0.07 , 0.36]				
Ley 2004	5	88	3	88	0.4%	1.68 [0.41 , 6.93]				
MRC 1968	50	199	50	194	3.8%	0.97 [0.61 , 1.52]		_	-	
Moy 2001	3	135	1	132	0.2%	2.69 [0.37 , 19.30]				
Oslo Diet-Heart 1966	64	206	90	206	4.9%	0.58 [0.39, 0.87]		-		
Rose corn oil 1965	12	28	3	13	0.4%	2.30 [0.60 , 8.85]		_		
Rose olive 1965	9	26	3	13	0.4%	1.70 [0.41 , 7.04]				
STARS 1992	3	27	10	28	0.5%	0.26 [0.08, 0.90]				
Veterans Admin 1969	60	424	78	422	5.8%	0.73 [0.51 , 1.05]				
WHI 2006 (1)	590	19541	913	29294	70.4%	0.97 [0.87 , 1.07]		•	l	
Total (95% CI)		21743		31456	100.0%	0.90 [0.83 , 0.99]	l			
Total events:	936		1325					•		
Heterogeneity: Chi ² = 33	.99, df = 10 (1	P = 0.0002	2); $I^2 = 71\%$				0.01	0.1 1	10	100
Test for overall effect: Z	= 2.25 (P = 0)	.02)]	Favours 1	ower SFA	Favours 1	higher SFA

Analysis 2.76. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 76: CHD events, SA Peto fixed-effect

Test for subgroup differences: Not applicable

Footnotes

Analysis 2.77. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 77: CHD events, subgroup by any substitution

Study or Subgroup	Lower	SFA	Higher SFA		Risk Ratio		Risk Ratio
	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
2.77.1 replaced by PUFA							
DART 1989	132	1018	144	1015	21.1%	0.91 [0.73, 1.14]	_
Houtsmuller 1979	8	51	30	51	9.9%	0.27 [0.14, 0.52]]
MRC 1968	50	199	50	194	17.9%	0.97 [0.69, 1.37]	
Oslo Diet-Heart 1966	64	206	90	206	20.2%	0.71 [0.55, 0.92]	_
Rose corn oil 1965	12	28	6	26	7.7%	1.86 [0.82, 4.22]	
STARS 1992	3	27	10	28	4.5%	0.31 [0.10 , 1.01]	
Veterans Admin 1969	60	424	78	422	18.7%	0.77 [0.56 . 1.04]	
Subtotal (95% CI)		1953		1942	100.0%	0.76 [0.57 . 1.00]	
Total events:	329	1,000	408		10000 /0		
Heterogeneity: $Tau^2 = 0.08$	22° Chi ² – 20	25 df -6	(P - 0.002)). $I^2 - 70\%$,		
Test for overall effect: $Z =$	1.97 (P = 0)	.05)	(1 = 0.002),1 = 70%			
2.77.2 replaced by MUFA	1						
Rose olive 1965	9	26	6	26	100.0%	1.50 [0.62 , 3.61]	
Subtotal (95% CI)		26		26	100.0%	1.50 [0.62 , 3.61]	
Total events:	9		6				
Heterogeneity: Not applica	ıble						
Test for overall effect: Z =	0.90 (P = 0	.37)					
2.77.3 replaced by CHO							
DART 1989	132	1018	144	1015	35.2%	0.91 [0.73 , 1.14]	
Ley 2004	5	88	3	88	1.6%	1.67 [0.41 , 6.76]	_
STARS 1992	3	30	10	30	2.2%	0.30 [0.09 , 0.98]	
WHI 2006 (1)	590	19541	913	29294	61.0%	0.97 [0.87 , 1.07]	•
Subtotal (95% CI)		20677		30427	100.0%	0.93 [0.78 , 1.11]	↓
Total events:	730		1070				1
Heterogeneity: $Tau^2 = 0.01$; Chi ² = 4.4	-9, df = 3 (P = 0.21;	$I^2 = 33\%$			
Test for overall effect: Z =	0.76 (P = 0	.44)					
2.77.4 replaced by protein	n						
DART 1989	132	1018	144	1015	17.7%	0.91 [0.73 , 1.14]	+
Ley 2004	5	88	3	88	0.4%	1.67 [0.41 , 6.76]	
WHI 2006 (1)	590	19541	913	29294	81.9%	0.97 [0.87 , 1.07]	
Subtotal (95% CI)		20647		30397	100.0%	0.96 [0.88 , 1.05]	Ŧ
Total events:	727		1060				
Heterogeneity: Tau ² = 0.00 Test for overall effect: Z =); $Chi^2 = 0.8$ 0.84 (P = 0	2, df = 2 (.40)	(P = 0.66);	$I^2 = 0\%$			
2.77.5 replacement uncles	ar						
Moy 2001	3	135	1	132	100.0%	2.93 [0.31 , 27.84]	
Subtotal (95% CI)		135		132	100.0%	2.93 [0.31 , 27.84]	
Total events:	3		1				
Heterogeneity: Not applica	ıble						
Test for overall effect: $Z =$	0.94 (P = 0	.35)					
Test for subgroup differences: Chi ² = 4.64, df = 4 (P = 0.33), I ² = 13.8%							01 0.1 1 10 100
Footpotos						Favo	uis lower SFA Favours nigner SF
roothotes							

(1) CHD events including CHD death & MI, Prentice 2017


Analysis 2.78. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 78: CHD events, subgroup by main substitution

	Lower	SFA	Higher	SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
2.78.1 replaced by PUFA							
DART 1989	132	1018	144	1015	22.4%	0.91 [0.73 , 1.14]	
Houtsmuller 1979	8	51	30	51	10.1%	0.27 [0.14, 0.52]	
MRC 1968	50	199	50	194	18.7%	0.97 [0.69 , 1.37]	
Oslo Diet-Heart 1966	64	206	90	206	21.3%	0.71 [0.55, 0.92]	-
Rose corn oil 1965	12	28	6	26	7.8%	1.86 [0.82, 4.22]	
Veterans Admin 1969	60	424	78	422	19.7%	0.77 [0.56, 1.04]	_
Subtotal (95% CI)		1926		1914	100.0%	0.79 [0.60 , 1.04]	Ā
Total events:	326		398				•
Heterogeneity: $Tau^2 = 0.07$: $Chi^2 = 17$.	71. df = 5	(P = 0.003)): $I^2 = 72\%$	2		
Test for overall effect: $Z =$	1.68 (P = 0)	.09)	(,,			
2.78.2 replaced by MUFA	L						
Rose olive 1965	9	26	6	26	100.0%	1.50 [0.62 , 3.61]	_ _
Subtotal (95% CI)		26		26	100.0%	1.50 [0.62 , 3.61]	
Total events:	9		6				
Heterogeneity: Not applica Test for overall effect: Z =	ble $0.90 (P = 0$.37)					
2.78.3 replaced by CHO							
Ley 2004	5	88	3	88	19.0%	1.67 [0.41 , 6.76]	
STARS 1992	3	30	10	30	23.5%	0.30 [0.09 , 0.98]	
WHI 2006 (1)	590	19541	913	29294	57.5%	0.97 [0.87, 1.07]	•
Subtotal (95% CI)		19659		29412	100.0%	0.82 [0.39 , 1.72]	•
Total events:	598		926				
Heterogeneity: Tau ² = 0.25 Test for overall effect: Z =	; $Chi^2 = 4.3$ 0.54 (P = 0	1, df = 2 (.59)	P = 0.12;	[² = 54%			
2.78.4 replaced by protein	1						
Subtotal (95% CI)		0		0		Not estimable	
Total events:	0		0				
Heterogeneity: Not applica	ble						
Test for overall effect: Not	applicable						
2.78.5 replacement unclea	ar						
Moy 2001	3	135	1	132	100.0%	2.93 [0.31 , 27.84]	
Subtotal (95% CI)		135		132	100.0%	2.93 [0.31 , 27.84]	
Total events:	3		1				
Heterogeneity: Not applica Test for overall effect: Z =	ble $0.94 (P = 0$.35)					
Test for subgroup difference	ces: Chi ² = 3	3.05, df = 1	3 (P = 0.38)), I ² = 1.7%	6	0. Fave	01 0.1 1 10 100 Durs lower SFA Favours higher SF
Footnotes						1	

(1) CHD events including CHD death & MI, Prentice 2017

	Lower	Lower SFA		Higher SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
2.79.1 up to 24 months							
DART 1989	132	1018	144	1015	85.1%	0.91 [0.73 , 1.14]	_
Moy 2001	3	135	1	132	1.6%	2.93 [0.31, 27.84]	
Rose corn oil 1965	12	28	3	13	6.9%	1.86 [0.63 , 5.47]	
Rose olive 1965	9	26	3	13	6.4%	1.50 [0.49, 4.62]	
Subtotal (95% CI)		1207		1173	100.0%	1.01 [0.76 , 1.35]	▲
Total events:	156		151				T
Heterogeneity: Tau ² = 0.0	01; Chi ² = 3.1	7, df = 3 (P = 0.37; I	$^{2} = 5\%$			
Test for overall effect: Z	= 0.06 (P = 0)	.95)					
2.79.2 >24 to 48 months							
MRC 1968	50	199	50	194	44.3%	0.97 [0.69, 1.37]	-
STARS 1992	3	27	10	28	8.2%	0.31 [0.10, 1.01]	
Veterans Admin 1969	60	424	78	422	47.5%	0.77 [0.56, 1.04]	-
Subtotal (95% CI)		650		644	100.0%	0.79 [0.55, 1.13]	
Total events:	113		138				•
Heterogeneity: Tau ² = 0.0	04; Chi ² = 3.8	2, $df = 2$ (P = 0.15; I	$^{2} = 48\%$			
Test for overall effect: Z	= 1.28 (P = 0)	.20)					
2.79.3 >48 months							
Oslo Diet-Heart 1966	64	206	90	206	42.6%	0.71 [0.55, 0.92]	-
WHI 2006 (1)	590	19541	913	29294	57.4%	0.97 [0.87, 1.07]	_
Subtotal (95% CI)		19747		29500	100.0%	0.85 [0.63, 1.15]	
Total events:	654		1003				•
Heterogeneity: Tau ² = 0.0	04; Chi ² = 4.9	1, df = 1 (P = 0.03; I	$^{2} = 80\%$			
Test for overall effect: Z	= 1.06 (P = 0)	.29)					
2.79.4 unclear duration							
Houtsmuller 1979	8	51	30	51	55.8%	0.27 [0.14, 0.52]	
Ley 2004	5	88	3	88	44.2%	1.67 [0.41 , 6.76]	
Subtotal (95% CI)		139		139	100.0%	0.60 [0.10 , 3.58]	
Total events:	13		33				
Heterogeneity: Tau ² = 1.	37; Chi ² = 5.3	5, df = 1 (P = 0.02; I	$^{2} = 81\%$			
Test for overall effect: Z	= 0.56 (P = 0)	.57)					
Test for subgroup differe	nces: Chi ² = 1	1.42, df = $\frac{1}{2}$	3 (P = 0.70)), $I^2 = 0\%$		(Fa	D.01 0.1 1 10 100 vours lower SFA Favours higher SF

Analysis 2.79. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 79: CHD events, subgroup by duration

Footnotes

(1) CHD events including CHD death & MI, Prentice 2017



Analysis 2.80. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 80: CHD events, subgroup by baseline SFA

	Lower	SFA	Higher	SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
2.80.1 up to 12% E SFA b	aseline						
Subtotal (95% CI)		0		0		Not estimable	
Total events:	0		0				
Heterogeneity: Not applica	ble						
Test for overall effect: Not	applicable						
2.80.2 >12 to 15%E SFA	baseline						
DART 1989	132	1018	144	1015	17.6%	0.91 [0.73 , 1.14]	_
Ley 2004	5	88	3	88	0.4%	1.67 [0.41 , 6.76]	
Moy 2001	3	135	1	132	0.2%	2.93 [0.31, 27.84]	
WHI 2006 (1)	590	19541	913	29294	81.8%	0.97 [0.87, 1.07]	-
Subtotal (95% CI)		20782		30529	100.0%	0.96 [0.88, 1.06]	T
Total events:	730		1061				
Heterogeneity: $Tau^2 = 0.00$); Chi ² = 1.7	6, df = 3 (P = 0.62; I	$2^2 = 0\%$			
Test for overall effect: $Z =$	0.80 (P = 0)	.42)					
2.80.3 >15 to 18%E SFA	baseline						
STARS 1992	3	27	10	28	100.0%	0.31 [0.10 , 1.01]	
Subtotal (95% CI)		27		28	100.0%	0.31 [0.10 , 1.01]	
Total events:	3		10				
Heterogeneity: Not applica	ıble						
Test for overall effect: $Z =$	1.94 (P = 0	.05)					
2.80.4 >18%E SFA baseli	ine						
Veterans Admin 1969	60	424	78	422	100.0%	0.77 [0.56, 1.04]	-
Subtotal (95% CI)		424		422	100.0%	0.77 [0.56 , 1.04]	
Total events:	60		78				•
Heterogeneity: Not applica	ıble						
Test for overall effect: $Z =$	1.70 (P = 0	.09)					
2.80.5 unclear							
Houtsmuller 1979	8	51	30	51	19.5%	0.27 [0.14, 0.52]	
MRC 1968	50	199	50	194	27.6%	0.97 [0.69 , 1.37]	+
Oslo Diet-Heart 1966	64	206	90	206	29.4%	0.71 [0.55, 0.92]	-
Rose corn oil 1965	12	28	3	13	12.1%	1.86 [0.63 , 5.47]	
Rose olive 1965	9	26	3	13	11.5%	1.50 [0.49 , 4.62]	_
Subtotal (95% CI)		510		477	100.0%	0.78 [0.49 , 1.26]	\bullet
Total events:	143		176				•
Heterogeneity: Tau ² = 0.18 Test for overall effect: Z =	$3; Chi^2 = 15.$ 1.00 (P = 0	59, df = 4 .32)	(P = 0.004); I ² = 74%)		
Test for subgroup difference	ces: Chi ² = 5	5.89, $df = 3$	3 (P = 0.12)), I ² = 49.1	%	0.0 Favc	D1 0.1 1 10 10 Durs lower SFA Favours higher

Footnotes

(1) CHD events including CHD death & MI, Prentice 2017

Analysis 2.81. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 81: CHD events, subgroup by SFA change

	Lower	SFA	Higher	SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
2.81.1 up to 4% E differ	ence						
DART 1989	132	1018	144	1015	17.6%	0.91 [0.73 , 1.14]	-
Ley 2004	5	88	3	88	0.4%	1.67 [0.41 , 6.76]	
Moy 2001	3	135	1	132	0.2%	2.93 [0.31 , 27.84]	
WHI 2006 (1)	590	19541	913	29294	81.8%	0.97 [0.87, 1.07]	-
Subtotal (95% CI)		20782		30529	100.0%	0.96 [0.88 , 1.06]	T
Total events:	730		1061				
Heterogeneity: $Tau^2 = 0.0$	00; Chi ² = 1.7	6, df = 3 (P = 0.62; I	$^{2} = 0\%$			
Test for overall effect: Z	= 0.80 (P = 0)	.42)					
2.81.2 >4 to 8%E differ	ence						
STARS 1992	3	27	10	28	100.0%	0.31 [0.10 , 1.01]	_
Subtotal (95% CI)		27		28	100.0%	0.31 [0.10 , 1.01]	
Total events:	3		10				•
Heterogeneity: Not applie	cable						
Test for overall effect: Z	= 1.94 (P = 0)	.05)					
2.81.3 >8%E difference							
Veterans Admin 1969	60	424	78	422	100.0%	0.77 [0.56, 1.04]	
Subtotal (95% CI)		424		422	100.0%	0.77 [0.56 , 1.04]	
Total events:	60		78				•
Heterogeneity: Not applie	cable						
Test for overall effect: Z	= 1.70 (P = 0)	.09)					
2.81.4 unclear							
Houtsmuller 1979	8	51	30	51	19.5%	0.27 [0.14, 0.52]	
MRC 1968	50	199	50	194	27.6%	0.97 [0.69 , 1.37]	-
Oslo Diet-Heart 1966	64	206	90	206	29.4%	0.71 [0.55, 0.92]	-
Rose corn oil 1965	12	28	3	13	12.1%	1.86 [0.63 , 5.47]	
Rose olive 1965	9	26	3	13	11.5%	1.50 [0.49 , 4.62]	_
Subtotal (95% CI)		510		477	100.0%	0.78 [0.49 , 1.26]	
Total events:	143		176				•
Heterogeneity: $Tau^2 = 0.1$	18; Chi ² = 15.	59, $df = 4$	(P = 0.004)	; $I^2 = 74\%$	2		
Test for overall effect: Z	= 1.00 (P = 0)	.32)					
Test for subgroup different	nces: Chi ² = 5	5.89, df =	3 (P = 0.12)	, I ² = 49.1	%	F	0.01 0.1 1 10 100 avours lower SFA Favours higher SFA

Footnotes

(1) CHD events including CHD death & MI, Prentice 2017



Analysis 2.82. Comparison 2: SFA reduction vs usual diet secondary health events, Outcome 82: CHD events, subgroup by sex

	Lower	SFA	Higher	SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
2.82.1 Men							
DART 1989	132	1018	144	1015	28.5%	0.91 [0.73, 1.14]	_
MRC 1968	50	199	50	194	18.4%	0.97 [0.69, 1.37]	_
Oslo Diet-Heart 1966	64	206	90	206	24.9%	0.71 [0.55, 0.92]	-
Rose corn oil 1965	12	28	3	13	2.8%	1.86 [0.63 , 5.47]	
Rose olive 1965	9	26	3	13	2.6%	1.50 [0.49 , 4.62]	_
STARS 1992	3	27	10	28	2.4%	0.31 [0.10, 1.01]	
Veterans Admin 1969	60	424	78	422	20.5%	0.77 [0.56 , 1.04]	
Subtotal (95% CI)		1928		1891	100.0%	0.84 [0.70 , 1.02]	
Total events:	330		378				•
Heterogeneity: $Tau^2 = 0.0$	02; Chi ² = 9.1	2, $df = 6$ (P = 0.17; I	$^{2} = 34\%$			
Test for overall effect: Z	= 1.79 (P = 0)	.07)					
2.82.2 Women							
WHI 2006 (1)	590	19541	913	29294	100.0%	0.97 [0.87, 1.07]	•
Subtotal (95% CI)		19541		29294	100.0%	0.97 [0.87 , 1.07]	T
Total events:	590		913				
Heterogeneity: Not appli	cable						
Test for overall effect: Z	= 0.61 (P = 0)	.54)					
2.82.3 Mixed, men and	women						
Houtsmuller 1979	8	51	30	51	42.2%	0.27 [0.14, 0.52]	
Ley 2004	5	88	3	88	33.8%	1.67 [0.41 , 6.76]	
Moy 2001	3	135	1	132	24.0%	2.93 [0.31, 27.84]	
Subtotal (95% CI)		274		271	100.0%	0.88 [0.18, 4.36]	
Total events:	16		34				
Heterogeneity: $Tau^2 = 1.4$	46; Chi ² = 8.3	38, df = 2 (P = 0.02; I	² = 76%			
Test for overall effect: Z	= 0.16 (P = 0)	.88)					
Test for subgroup differe	nces: Chi ² =	1.64, df =	2 (P = 0.44)), $I^2 = 0\%$		_ o.d	
Footnotes						Favo	urs lower SFA Favours higher SF.

(1) CHD events including CHD death & MI, Prentice 2017

Favours lower SFA

Favours higher SFA

Analysis 2.83. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 83: CHD events, subgroup by CVD risk

	Lower	SFA	Higher	SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
2.83.1 Low CVD risk							
Veterans Admin 1969	60	424	78	422	21.4%	0.77 [0.56 , 1.04]	
WHI 2006 (1)	806	18633	1292	27925	78.6%	0.93 [0.86 , 1.02]	-
Subtotal (95% CI)		19057		28347	100.0%	0.90 [0.76 , 1.05]	4
Total events:	866		1370				•
Heterogeneity: Tau ² = 0.01	; Chi ² = 1.5	0, df = 1 (P = 0.22; I	$^{2} = 33\%$			
Test for overall effect: Z =	1.34 (P = 0)	.18)					
2.83.2 Moderate CVD risk	k						
Houtsmuller 1979	8	51	30	51	42.2%	0.27 [0.14, 0.52]	
Ley 2004	5	88	3	88	33.8%	1.67 [0.41 , 6.76]	_
Moy 2001	3	135	1	132	24.0%	2.93 [0.31 , 27.84]	
Subtotal (95% CI)		274		271	100.0%	0.88 [0.18, 4.36]	
Total events:	16		34				Ť
Heterogeneity: $Tau^2 = 1.46$; Chi ² = 8.3	8, $df = 2$ (P = 0.02; I	$^{2} = 76\%$			
Test for overall effect: $Z =$	0.16 (P = 0)	.88)					
2.83.3 Existing CVD disea	ise						
DART 1989	132	1018	144	1015	23.7%	0.91 [0.73 , 1.14]	+
MRC 1968	50	199	50	194	17.9%	0.97 [0.69 , 1.37]	_
Oslo Diet-Heart 1966	64	206	90	206	21.9%	0.71 [0.55, 0.92]	-
Rose corn oil 1965	12	28	3	13	3.6%	1.86 [0.63 , 5.47]	_ _
Rose olive 1965	9	26	3	13	3.4%	1.50 [0.49 , 4.62]	_ _
STARS 1992	3	27	10	28	3.1%	0.31 [0.10 , 1.01]	_
WHI 2006 (2)	194	908	257	1369	26.4%	1.14 [0.96 , 1.34]	-
Subtotal (95% CI)		2412		2838	100.0%	0.94 [0.75 , 1.16]	•
Total events:	464		557				ľ
Heterogeneity: $Tau^2 = 0.04$; Chi ² = 15.	05, df = 6	(P = 0.02);	$\mathrm{I^2}=60\%$			
Test for overall effect: $Z =$	0.59 (P = 0	.55)					
Test for subgroup differenc	es: Chi ² = (0.10, df = 2	2 (P = 0.95)), $I^2 = 0\%$. h	

Footnotes

(1) Women without CVD at baseline

(2) Women with CVD at baseline



Analysis 2.84. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 84: CHD events, subgroup by TC reduction

	Lower	SFA	Higher	SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
2.84.1 serum chol reduce	d by at leas	t 0.2mmo	I/L				
DART 1989	132	1018	144	1015	21.7%	0.91 [0.73 , 1.14]	-
Houtsmuller 1979	8	51	30	51	9.8%	0.27 [0.14, 0.52]	
MRC 1968	50	199	50	194	18.1%	0.97 [0.69, 1.37]	_
Moy 2001	3	135	1	132	1.4%	2.93 [0.31 , 27.84]	
Oslo Diet-Heart 1966	64	206	90	206	20.6%	0.71 [0.55, 0.92]	-
Rose corn oil 1965	12	28	3	13	5.0%	1.86 [0.63 , 5.47]	
STARS 1992	3	27	10	28	4.3%	0.31 [0.10, 1.01]	
Veterans Admin 1969	60	424	78	422	19.0%	0.77 [0.56 , 1.04]	
Subtotal (95% CI)		2088		2061	100.0%	0.76 [0.58 , 0.99]	
Total events:	332		406				•
Heterogeneity: $Tau^2 = 0.07$	7; Chi ² = 19.	84, df = 7	(P = 0.006)	; $I^2 = 65\%$	2		
Test for overall effect: Z =	= 2.03 (P = 0)	.04)					
	11 (0.0	1/7					
2.84.2 serum chol reduce	d by <0.2m	mol/L			0.50		
Ley 2004	5	88	3	88	0.5%	1.67 [0.41 , 6.76]	_
Rose olive 1965	9	26	3	13	0.8%	1.50 [0.49 , 4.62]	_ <u>_</u>
WHI 2006 (1)	590	19541	913	29294	98.7%	0.97 [0.87 , 1.07]	
Subtotal (95% CI)		19655		29395	100.0%	0.97 [0.88 , 1.08]	•
Total events:	604		919				
Heterogeneity: $Tau^2 = 0.00$	0; Chi ² = 1.1	4, df = 2 (P = 0.56; I	$^{2} = 0\%$			
Test for overall effect: Z =	= 0.49 (P = 0)	.62)					
2.84.3 serum chol reduct	ion unclear						
Subtotal (95% CI)		0		0		Not estimable	
Total events:	0		0				
Heterogeneity: Not applica	able						
Test for overall effect: Not	t applicable						
Test for subgroup differen	ces: Chi ² = 3	3.00, df =	1 (P = 0.08)	, I² = 66.6	%	0.0	
						Favo	urs lower SFA Favours higher SFA
Footnotes							

(1) CHD events including CHD death & MI, Prentice 2017

Analysis 2.85. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 85: CHD events, subgroup decade of publication

	Lower	SFA	Higher	SFA		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
2.85.1 1960s							
MRC 1968	50	199	50	194	13.5%	0.97 [0.69, 1.37]	_
Oslo Diet-Heart 1966	64	206	90	206	16.3%	0.71 [0.55, 0.92]	_
Rose corn oil 1965	12	28	3	13	2.9%	1.86 [0.63, 5.47]	
Rose olive 1965	9	26	3	13	2.7%	1.50 [0.49, 4.62]	
Veterans Admin 1969	60	424	78	422	14.5%	0.77 [0.56, 1.04]	
Subtotal (95% CI)		883		848	50.0%	0.84 [0.68, 1.05]	
Total events:	195		224				
Heterogeneity: $Tau^2 = 0.0$	$02; Chi^2 = 5.7$	70, $df = 4$ (P = 0.22;	$1^2 = 30\%$			
Test for overall effect: Z	= 1.51 (P = 0)).13)					
2.85.2 1970s							
Houtsmuller 1979	8	51	30	51	6.2%	0.27 [0.14, 0.52]	
Subtotal (95% CI)		51		51	6.2%	0.27 [0.14, 0.52]	
Total events:	8		30			- / -	▼
Heterogeneity: Not applic	cable						
Test for overall effect: Z	= 3.83 (P = 0)	0.0001)					
2.85.3 1980s							
DART 1989	132	1018	144	1015	17.6%	0.91 [0.73 . 1.14]	1
Subtotal (95% CI)		1018		1015	17.6%	0.91 [0.73 , 1.14]	1
Total events:	132		144				T
Heterogeneity: Not applic	cable						
Test for overall effect: Z	= 0.80 (P = 0)	0.42)					
2.85.4 1990s							
STARS 1992	3	27	10	30	2.5%	0.33 [0.10 , 1.09]	
Subtotal (95% CI)		27		30	2.5%	0.33 [0.10 , 1.09]	
Total events:	3		10				
Heterogeneity: Not applic	cable						
Test for overall effect: Z	= 1.82 (P = 0)	0.07)					
2.85.5 2000s							
Ley 2004	5	88	3	88	1.8%	1.67 [0.41, 6.76]	
Moy 2001	3	135	1	132	0.7%	2.93 [0.31, 27.84]	
WHI 2006 (1)	590	19541	913	29294	21.2%	0.97 [0.87 , 1.07]	
Subtotal (95% CI)		19764		29514	23.8%	0.97 [0.88 , 1.08]	I I I I I I I I I I I I I I I I I I I
Total events:	598		917				
Heterogeneity: $Tau^2 = 0.0$	$00; Chi^2 = 1.5$	50, $df = 2$ (P = 0.47;	$[^2 = 0\%]$			
Test for overall effect: Z	= 0.51 (P = 0)).61)					
Total (95% CI)		21743		31458	100.0%	0.83 [0.68 , 1.01]	
Total events:	936		1325				•
Heterogeneity: $Tau^2 = 0.0$	05; Chi ² = 26	.03, df = 1	0 (P = 0.00)	4); $I^2 = 62^6$	%		1 0 1 1 10 100
Test for overall effect: Z	= 1.82 (P = 0)	0.07)				Favo	urs lower SFA Favours higher SFA
Test for subgroup differen	nces: Chi ² =	17.51, df =	= 4 (P = 0.0)	02), $I^2 = 77$	7.2%		÷

Footnotes

(1) CHD events including CHD death & MI, Prentice 2017

Analysis 2.86. Comparison 2: SFA reduction vs usual diet - secondary health events, Outcome 86: DIABETES DIAGNOSES

	Lower	SFA	Higher SFA		Risk Ratio	Risk Ratio		
Study or Subgroup	Events	Total	Events	Total	M-H, Random, 95% CI	M-H, Random, 95% Cl	[
WHI 2006	1303	19541	2039	29294	0.96 [0.90 , 1.02]	-+		
Test for subgroup differe	ences: Not a	pplicable			F	0.85 0.9 1 1.1 avours lower SFA Favours	1.2 higher SFA	

Comparison 3. SFA reduction vs usual diet - secondary blood outcomes

Outcome or subgroup title	No. of studies	No. of partici- pants	Statistical method	Effect size
3.1 Total cholesterol, mmol/L	14	7115	Mean Difference (IV, Random, 95% CI)	-0.24 [-0.36, -0.13]
3.2 TC, mmol/L, subgroup by any replace- ment	14		Mean Difference (IV, Random, 95% CI)	Subtotals only
3.2.1 replaced by PUFA	9	3888	Mean Difference (IV, Random, 95% CI)	-0.33 [-0.47, -0.19]
3.2.2 replace by MUFA	1	24	Mean Difference (IV, Random, 95% CI)	0.30 [-0.93, 1.53]
3.2.3 replace by CHO	6	5094	Mean Difference (IV, Random, 95% CI)	-0.18 [-0.32, -0.04]
3.2.4 replace by protein	4	4986	Mean Difference (IV, Random, 95% CI)	-0.15 [-0.27, -0.04]
3.2.5 replacement unclear	1	72	Mean Difference (IV, Random, 95% CI)	-0.34 [-0.64, -0.04]
3.3 TC, mmol/L, subgroup by main re- placement	14		Mean Difference (IV, Random, 95% CI)	Subtotals only
3.3.1 replaced by PUFA	8	3838	Mean Difference (IV, Random, 95% CI)	-0.28 [-0.37, -0.19]
3.3.2 replace by MUFA	1	24	Mean Difference (IV, Random, 95% CI)	0.30 [-0.93, 1.53]
3.3.3 replace by CHO	4	3181	Mean Difference (IV, Random, 95% CI)	-0.19 [-0.40, 0.01]
3.3.4 replace by protein	0	0	Mean Difference (IV, Random, 95% CI)	Not estimable
3.3.5 replacement unclear	1	72	Mean Difference (IV, Random, 95% CI)	-0.34 [-0.64, -0.04]

Reduction in saturated fat intake for cardiovascular disease (Review)



Cochrane Database of Systematic Reviews

Outcome or subgroup title	No. of studies	No. of partici- pants	Statistical method	Effect size
3.4 LDL cholesterol, mmol/L	5	3291	Mean Difference (IV, Random, 95% CI)	-0.19 [-0.33, -0.05]
3.5 LDL, mmol/L, subgroup by any re- placement	5		Mean Difference (IV, Random, 95% CI)	Subtotals only
3.5.1 replaced by PUFA	1	50	Mean Difference (IV, Random, 95% CI)	-0.48 [-0.90, -0.06]
3.5.2 replace by MUFA	0	0	Mean Difference (IV, Random, 95% CI)	Not estimable
3.5.3 replace by CHO	3	2985	Mean Difference (IV, Random, 95% CI)	-0.16 [-0.35, 0.02]
3.5.4 replace by protein	2	2935	Mean Difference (IV, Random, 95% CI)	-0.09 [-0.15, -0.04]
3.5.5 replacement unclear	2	306	Mean Difference (IV, Random, 95% CI)	-0.29 [-0.51, -0.08]
3.6 LDL, mmol/L, subgroup by main re- placement	5		Mean Difference (IV, Random, 95% CI)	Subtotals only
3.6.1 replaced by PUFA	0	0	Mean Difference (IV, Random, 95% CI)	Not estimable
3.6.2 replace by MUFA	0	0	Mean Difference (IV, Random, 95% CI)	Not estimable
3.6.3 replace by CHO	3	2985	Mean Difference (IV, Random, 95% CI)	-0.16 [-0.35, 0.02]
3.6.4 replace by protein	0	0	Mean Difference (IV, Random, 95% CI)	Not estimable
3.6.5 replacement unclear	2	306	Mean Difference (IV, Random, 95% CI)	-0.29 [-0.51, -0.08]
3.7 HDL cholesterol, mmol/L	6	5147	Mean Difference (IV, Random, 95% CI)	-0.01 [-0.02, 0.01]
3.8 HDL, mmol/L, subgroup by any re- placement	6		Mean Difference (IV, Random, 95% CI)	Subtotals only
3.8.1 replaced by PUFA	2	1905	Mean Difference (IV, Random, 95% CI)	-0.01 [-0.04, 0.01]
3.8.2 replace by MUFA	0	0	Mean Difference (IV, Random, 95% CI)	Not estimable
3.8.3 replace by CHO	4	4840	Mean Difference (IV, Random, 95% CI)	-0.01 [-0.03, 0.00]

Reduction in saturated fat intake for cardiovascular disease (Review)



Cochrane Database of Systematic Reviews

Outcome or subgroup title	No. of studies	No. of partici- pants	Statistical method	Effect size
3.8.4 replace by protein	3	4790	Mean Difference (IV, Random, 95% CI)	-0.01 [-0.03, 0.00]
3.8.5 replacement unclear	2	307	Mean Difference (IV, Random, 95% CI)	0.01 [-0.10, 0.12]
3.9 HDL, mmol/L, subgroup by main re- placement	6		Mean Difference (IV, Random, 95% CI)	Subtotals only
3.9.1 replaced by PUFA	1	1855	Mean Difference (IV, Random, 95% CI)	-0.01 [-0.04, 0.02]
3.9.2 replace by MUFA	0	0	Mean Difference (IV, Random, 95% CI)	Not estimable
3.9.3 replace by CHO	3	2985	Mean Difference (IV, Random, 95% CI)	-0.01 [-0.03, 0.01]
3.9.4 replace by protein	0	0	Mean Difference (IV, Random, 95% CI)	Not estimable
3.9.5 replacement unclear	2	307	Mean Difference (IV, Random, 95% CI)	0.01 [-0.10, 0.12]
3.10 Triglycerides, mmol/L	7	3845	Mean Difference (IV, Random, 95% CI)	-0.08 [-0.21, 0.04]
3.11 TG, mmol/L, subgroup by any re- placement	7		Mean Difference (IV, Random, 95% CI)	Subtotals only
3.11.1 replaced by PUFA	3	604	Mean Difference (IV, Random, 95% CI)	-0.19 [-0.35, -0.02]
3.11.2 replace by MUFA	0	0	Mean Difference (IV, Random, 95% CI)	Not estimable
3.11.3 replace by CHO	3	2985	Mean Difference (IV, Random, 95% CI)	-0.04 [-0.32, 0.25]
3.11.4 replace by protein	2	2935	Mean Difference (IV, Random, 95% CI)	0.01 [-0.08, 0.09]
3.11.5 replacement unclear	2	306	Mean Difference (IV, Random, 95% CI)	-0.09 [-0.52, 0.33]
3.12 TG, mmol/L, subgroup by main re- placement	7		Mean Difference (IV, Random, 95% CI)	Subtotals only
3.12.1 replaced by PUFA	2	554	Mean Difference (IV, Random, 95% CI)	-0.16 [-0.30, -0.01]
3.12.2 replace by MUFA	0	0	Mean Difference (IV, Random, 95% CI)	Not estimable

Reduction in saturated fat intake for cardiovascular disease (Review)



Cochrane Database of Systematic Reviews

Outcome or subgroup title	No. of studies	No. of partici- pants	Statistical method	Effect size
3.12.3 replace by CHO	3	2985	Mean Difference (IV, Random, 95% CI)	-0.04 [-0.32, 0.25]
3.12.4 replace by protein	0	0	Mean Difference (IV, Random, 95% CI)	Not estimable
3.12.5 replacement unclear	2	306	Mean Difference (IV, Random, 95% CI)	-0.09 [-0.52, 0.33]
3.13 total cholesterol /HDL ratio	3	2985	Mean Difference (IV, Random, 95% CI)	-0.10 [-0.33, 0.13]
3.14 TC /HDL ratio, subgroup by any re- placement	3		Mean Difference (IV, Random, 95% CI)	Subtotals only
3.14.1 replaced by PUFA	1	50	Mean Difference (IV, Random, 95% CI)	-0.58 [-1.33, 0.17]
3.14.2 replace by MUFA	0	0	Mean Difference (IV, Random, 95% CI)	Not estimable
3.14.3 replace by CHO	3	2985	Mean Difference (IV, Random, 95% CI)	-0.10 [-0.33, 0.13]
3.14.4 replace by protein	2	2935	Mean Difference (IV, Random, 95% CI)	-0.09 [-0.21, 0.04]
3.14.5 replacement unclear	0	0	Mean Difference (IV, Random, 95% CI)	Not estimable
3.15 TC /HDL ratio, subgroup by main re- placement	3		Mean Difference (IV, Random, 95% CI)	Subtotals only
3.15.1 replaced by PUFA	0	0	Mean Difference (IV, Random, 95% CI)	Not estimable
3.15.2 replace by MUFA	0	0	Mean Difference (IV, Random, 95% CI)	Not estimable
3.15.3 replace by CHO	3	2985	Mean Difference (IV, Random, 95% CI)	-0.10 [-0.33, 0.13]
3.15.4 replace by protein	0	0	Mean Difference (IV, Random, 95% CI)	Not estimable
3.15.5 replacement unclear	0	0	Mean Difference (IV, Random, 95% CI)	Not estimable
3.16 LDL /HDL ratio	1		Mean Difference (IV, Random, 95% CI)	Subtotals only
3.17 Lp(a), mmol/L	2	2882	Mean Difference (IV, Random, 95% CI)	0.00 [-0.00, 0.00]

Reduction in saturated fat intake for cardiovascular disease (Review)



Cochrane Database of Systematic Reviews

Outcome or subgroup title	No. of studies	No. of partici- pants	Statistical method	Effect size
3.18 Lp(a), mmol/L, subgroup by any re- placement	2		Mean Difference (IV, Random, 95% CI)	Subtotals only
3.18.1 replaced by PUFA	1	50	Mean Difference (IV, Random, 95% CI)	0.00 [-1.37, 1.37]
3.18.2 replace by MUFA	0	0	Mean Difference (IV, Random, 95% CI)	Not estimable
3.18.3 replace by CHO	2	2882	Mean Difference (IV, Random, 95% CI)	0.00 [-0.00, 0.00]
3.18.4 replace by protein	1	2832	Mean Difference (IV, Random, 95% CI)	0.00 [-0.00, 0.00]
3.18.5 replacement unclear	0	0	Mean Difference (IV, Random, 95% CI)	Not estimable
3.19 Lp(a), mmol/L, subgroup by main re- placement	2		Mean Difference (IV, Random, 95% CI)	Subtotals only
3.19.1 replaced by PUFA	0	0	Mean Difference (IV, Random, 95% CI)	Not estimable
3.19.2 replace by MUFA	0	0	Mean Difference (IV, Random, 95% CI)	Not estimable
3.19.3 replace by CHO	2	2882	Mean Difference (IV, Random, 95% CI)	0.00 [-0.00, 0.00]
3.19.4 replace by protein	0	0	Mean Difference (IV, Random, 95% CI)	Not estimable
3.19.5 replacement unclear	0	0	Mean Difference (IV, Random, 95% CI)	Not estimable
3.20 Insulin sensitivity	4		Mean Difference (IV, Random, 95% CI)	Subtotals only
3.20.1 HbA1c (glycosylated haemoglo- bin), %	0	0	Mean Difference (IV, Random, 95% CI)	Not estimable
3.20.2 GTT (glucose tolerance test), glu- cose at 2 hours, mmol/L	3	249	Mean Difference (IV, Random, 95% CI)	-1.69 [-2.55, -0.82]
3.20.3 HOMA	1	2832	Mean Difference (IV, Random, 95% CI)	0.00 [-0.04, 0.04]



Cochrane

Library

Analysis 3.1. Comparison 3: SFA reduction vs usual diet secondary blood outcomes, Outcome 1: Total cholesterol, mmol/L

	Le	ower SFA		н	HIgher SFA			Mean Difference	Mean Difference
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% C	I IV, Random, 95% CI
DART 1989	6.31	1.14	924	6.57	1.16	931	17.0%	-0.26 [-0.36 , -0.1	6] 📕
Houtsmuller 1979	6.43	0.65	48	6.9	0.81	48	8.3%	-0.47 [-0.76 , -0.1	8]
Ley 2004	-0.2	0.79	51	-0.15	1.3	52	5.3%	-0.05 [-0.46 , 0.3	6]
MRC 1968	-1.11	0	88	-0.47	0	89		Not estimab	le
Oslo Diet-Heart 1966	6.12	0	168	7.19	0	161		Not estimab	le
Oxford Retinopathy 1978	4.94	0.82	29	4.87	0.79	29	5.3%	0.07 [-0.34 , 0.4	8]
Rose corn oil 1965	-0.78	0.923	13	-0.2	1.031	9	1.6%	-0.58 [-1.42 , 0.2	6]
Rose olive 1965	0.1	2.023	15	-0.2	1.031	9	0.8%	0.30 [-0.93 , 1.5	3]
STARS 1992	6.17	0.459	26	6.93	0.98	24	5.0%	-0.76 [-1.19 , -0.3	3]
Simon 1997	4.87	0.87	34	5.21	0.18	38	8.2%	-0.34 [-0.64 , -0.04	4]
Sydney Diet-Heart 1978	6.5	1.2	221	6.8	1.1	237	11.6%	-0.30 [-0.51 , -0.0	9]
Veterans Admin 1969	4.93	3.72	423	5.3	1.87	420	5.6%	-0.37 [-0.77 , 0.0	3]
WHI 2006	-0.264	0.828	1133	-0.178	0.825	1699	18.9%	-0.09 [-0.15 , -0.0	2]
WINS 2006	-0.0005	0.728	96	0.142	0.675	100	12.3%	-0.14 [-0.34 , 0.0	5] _
Total (95% CI)			3269			3846	100.0%	-0.24 [-0.36 , -0.1	3]
Heterogeneity: $Tau^2 = 0.02$; C	Chi ² = 27.85, d	f = 11 (P = 1)	= 0.003); I	$^{2} = 60\%$					•
Test for overall effect: $Z = 4$.	32 (P < 0.000	l)							-2 -1 0 1 2
Test for subgroup differences	: Not applicat	ole							Favours lower SFA Favours higher SFA

Analysis 3.2. Comparison 3: SFA reduction vs usual diet - secondary blood outcomes, Outcome 2: TC, mmol/L, subgroup by any replacement

	Lower SFA		HIgher SFA					Mean Difference	Mean Difference		
tudy or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI		
2.2.1 replaced by PUFA											
DART 1989	6.31	1.14	924	6.57	1.16	931	33.9%	-0.26 [-0.36, -0.16]			
Houtsmuller 1979	6.43	0.65	48	6.9	0.81	48	14.8%	-0.47 [-0.76, -0.18]			
/IRC 1968	-1.11	0	88	-0.47	0	89		Not estimable			
Slo Diet-Heart 1966	6.12	0	168	7.19	0	161		Not estimable			
0xford Retinopathy 1978	4.94	0.82	29	4.87	0.79	29	9.0%	0.07 [-0.34 , 0.48]			
ose corn oil 1965	-0.78	0.923	13	-0.2	1.031	9	2.6%	-0.58 [-1.42, 0.26]	←		
TARS 1992	6.17	0.459	26	6.93	0.98	24	8.5%	-0.76 [-1.19 , -0.33]	▲		
ydney Diet-Heart 1978	6.5	1.2	221	6.8	1.1	237	21.5%	-0.30 [-0.51 , -0.09]			
eterans Admin 1969	4.93	3.72	423	5.3	1.87	420	9.6%	-0.37 [-0.77 , 0.03]	_		
ubtotal (95% CI)			1940			1948	100.0%	-0.33 [-0.47 , -0.19]			
eterogeneity: Tau ² = 0.01; Ch	i ² = 9.85, df	= 6 (P = 0)	$(.13); I^2 = 3$	39%					•		
est for overall effect: $Z = 4.62$	(P < 0.0000)1)									
.2.2 replace by MUFA											
ose olive 1965	0.1	2.023	15	-0.2	1.031	9	100.0%	0.30 [-0.93 , 1.53]			
ubtotal (95% CI)			15			9	100.0%	0.30 [-0.93 , 1.53]			
leterogeneity: Not applicable											
est for overall effect: $Z = 0.48$	(P = 0.63)										
.2.3 replace by CHO											
ART 1989	6.31	1.14	924	6.57	1.16	931	26.8%	-0.26 [-0.36 , -0.16]			
ey 2004	-0.2	0.79	51	-0.15	1.3	52	8.2%	-0.05 [-0.46, 0.36]			
xford Retinopathy 1978	4.94	0.82	29	4.87	0.79	29	8.2%	0.07 [-0.34 , 0.48]			
TARS 1992	6.17	0.459	26	6.93	0.98	24	7.8%	-0.76 [-1.19 , -0.33]			
/HI 2006	-0.264	0.828	1133	-0.178	0.825	1699	29.7%	-0.09 [-0.15, -0.02]	•		
/INS 2006	-0.0005	0.728	96	0.142	0.675	100	19.3%	-0.14 [-0.34, 0.05]			
ubtotal (95% CI)			2259			2835	100.0%	-0.18 [-0.32 , -0.04]			
eterogeneity: Tau ² = 0.02; Ch	i ² = 17.09, d	f = 5 (P =	0.004); I ²	= 71%					•		
est for overall effect: $Z = 2.55$	(P = 0.01)										
.2.4 replace by protein											
ART 1989	6.31	1.14	924	6.57	1.16	931	33.3%	-0.26 [-0.36 , -0.16]			
ey 2004	-0.2	0.79	51	-0.15	1.3	52	6.4%	-0.05 [-0.46 , 0.36]			
/HI 2006	-0.264	0.828	1133	-0.178	0.825	1699	40.9%	-0.09 [-0.15 , -0.02]	-		
/INS 2006	-0.0005	0.728	96	0.142	0.675	100	19.3%	-0.14 [-0.34 , 0.05]			
ubtotal (95% CI)			2204			2782	100.0%	-0.15 [-0.27 , -0.04]			
eterogeneity: Tau ² = 0.01; Ch	i ² = 8.01, df	= 3 (P = 0)	$(0.05); I^2 = 0$	53%				- / *			
est for overall effect: $Z = 2.64$	(P = 0.008)										
.2.5 replacement unclear											
imon 1997	4.87	0.87	34	5.21	0.18	38	100.0%	-0.34 [-0.64 , -0.04]			
ubtotal (95% CI)			34			38	100.0%	-0.34 [-0.64 , -0.04]			
leterogeneity: Not applicable			-					,			
'est for overall effect: $Z = 2.24$	(P = 0.03)										
est for subgroup differences:	$hi^2 - 5.40$	$df = A (\mathbf{P} -$	-025)12.	- 25 0%							
est for subgroup differences.	JIII – J.40,	ur – + (r =	- 0.23), 123	- 23.370				F	-0.5-0.25 0 0.25 0.5 avours lower SFA Favours hi		



Analysis 3.3. Comparison 3: SFA reduction vs usual diet - secondary blood outcomes, Outcome 3: TC, mmol/L, subgroup by main replacement

	Lo	wer SFA		HIgher SFA				Mean Difference	Mean Difference	
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI	
3.3.1 replaced by PUFA										
DART 1989	6.31	1.14	924	6.57	1.16	931	61.4%	-0.26 [-0.36 , -0.16]		
Houtsmuller 1979	6.43	0.65	48	6.9	0.81	48	9.5%	-0.47 [-0.76 , -0.18]		
MRC 1968	-1.11	0	88	-0.47	0	89		Not estimable		
Oslo Diet-Heart 1966	6.12	0	168	7.19	0	161		Not estimable		
Oxford Retinopathy 1978	4.94	0.82	29	4.87	0.79	29	4.8%	0.07 [-0.34, 0.48]		
Rose corn oil 1965	-0.78	0.923	13	-0.2	1.031	9	1.2%	-0.58 [-1.42, 0.26]	←	
Sydney Diet-Heart 1978	6.5	1.2	221	6.8	1.1	237	17.8%	-0.30 [-0.51 , -0.09]	·	
Veterans Admin 1969	4.93	3.72	423	5.3	1.87	420	5.3%	-0.37 [-0.77, 0.03]		
Subtotal (95% CI)			1914			1924	100.0%	-0.28 [-0.37 , -0.19]		
Heterogeneity: Tau ² = 0.00; Ch	$i^2 = 5.21$, df	= 5 (P = 0)).39); I ² = 4	4%					•	
Test for overall effect: $Z = 5.98$	8 (P < 0.0000	01)								
3.3.2 replace by MUFA										
Rose olive 1965	0.1	2.023	15	-0.2	1.031	9	100.0%	0.30 [-0.93 , 1.53]		
Subtotal (95% CI)			15			9	100.0%	0.30 [-0.93 , 1.53]		
Heterogeneity: Not applicable										
Test for overall effect: $Z = 0.48$	B (P = 0.63)									
3.3.3 replace by CHO										
Ley 2004	-0.2	0.79	51	-0.15	1.3	52	15.1%	-0.05 [-0.46 , 0.36]	_	
STARS 1992	6.17	0.459	26	6.93	0.98	24	14.4%	-0.76 [-1.19 , -0.33]	←	
WHI 2006	-0.264	0.828	1133	-0.178	0.825	1699	40.4%	-0.09 [-0.15 , -0.02]	-	
WINS 2006	-0.0005	0.728	96	0.142	0.675	100	30.0%	-0.14 [-0.34 , 0.05]		
Subtotal (95% CI)			1306			1875	100.0%	-0.19 [-0.40 , 0.01]		
Heterogeneity: $Tau^2 = 0.03$; Ch Test for overall effect: $Z = 1.89$	$i^2 = 9.48$, df P(P = 0.06)	= 3 (P = 0)	$(0.02); I^2 = 0$	58%					•	
3.3.4 replace by protein										
Subtotal (95% CI)			0			0		Not estimable		
Heterogeneity: Not applicable										
Test for overall effect: Not app	licable									
3.3.5 replacement unclear										
Simon 1997	4.87	0.87	34	5.21	0.18	38	100.0%	-0.34 [-0.64 , -0.04]		
Subtotal (95% CI)			34			38	100.0%	-0.34 [-0.64 , -0.04]		
Heterogeneity: Not applicable									-	
Test for overall effect: $Z = 2.24$	(P = 0.03)									
Test for subgroup differences:	Chi ² = 1.63,	df = 3 (P =	= 0.65), I ²	= 0%				F	-0.5-0.25 0 0.25 0.5 avours lower SEA Eavours higher S	

Analysis 3.4. Comparison 3: SFA reduction vs usual diet secondary blood outcomes, Outcome 4: LDL cholesterol, mmol/L

	Lo	wer SFA		HIgher SFA				Mean Difference	Mean Difference	
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI	
Ley 2004	-0.32	0.64	51	-0.16	1.15	52	11.7%	-0.16 [-0.52 , 0.20]		
Moy 2001	-0.69	1.1	117	-0.4	0.8	118	20.0%	-0.29 [-0.54 , -0.04]]	
STARS 1992	4.19	0.51	26	4.67	0.931	24	9.0%	-0.48 [-0.90 , -0.06]	I	
Simon 1997	2.79	0.82	34	3.09	0.99	37	9.0%	-0.30 [-0.72 , 0.12]	I	
WHI 2006	-0.251	0.758	1133	-0.16	0.753	1699	50.3%	-0.09 [-0.15 , -0.03]	1 🖷	
Total (95% CI)			1361			1930	100.0%	-0.19 [-0.33 , -0.05]		
Heterogeneity: Tau ² = 0	$0.01; Chi^2 = 6.$	30, $df = 4$	(P = 0.18)	; I ² = 37%					•	
Test for overall effect: 2	Z = 2.73 (P =	0.006)							-0.5 -0.25 0 0.25 0.5	
Test for subgroup differ	rences: Not ap	1	Favours lower SFA Favours higher S							

Reduction in saturated fat intake for cardiovascular disease (Review)

Analysis 3.5. Comparison 3: SFA reduction vs usual diet - secondary blood outcomes, Outcome 5: LDL, mmol/L, subgroup by any replacement

	Lo	Lower SFA			HIgher SFA			Mean Difference	Mean Difference	
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI	
3.5.1 replaced by PUFA	A									
STARS 1992	4.19	0.51	26	4.67	0.931	24	100.0%	-0.48 [-0.90 , -0.06]		
Subtotal (95% CI)			26			24	100.0%	-0.48 [-0.90 , -0.06]		
Heterogeneity: Not appl	licable									
Test for overall effect: Z	Z = 2.24 (P =	0.03)								
3.5.2 replace by MUFA	1									
Subtotal (95% CI)			0			0		Not estimable		
Heterogeneity: Not appl	licable									
Test for overall effect: N	Not applicable									
3.5.3 replace by CHO										
Ley 2004	-0.32	0.64	51	-0.16	1.15	52	19.7%	-0.16 [-0.52 , 0.20]		
STARS 1992	4.19	0.51	26	4.67	0.931	24	15.5%	-0.48 [-0.90 , -0.06]		
WHI 2006	-0.251	0.758	1133	-0.16	0.753	1699	64.8%	-0.09 [-0.15 , -0.03]		
Subtotal (95% CI)			1210			1775	100.0%	-0.16 [-0.35 , 0.02]		
Heterogeneity: Tau ² = 0	.01; Chi ² = 3.	33, df = 2	(P = 0.19)	; I ² = 40%						
Test for overall effect: Z	Z = 1.72 (P =	0.09)								
3.5.4 replace by protein	n									
Ley 2004	-0.32	0.64	51	-0.16	1.15	52	2.5%	-0.16 [-0.52 , 0.20]		
WHI 2006	-0.251	0.758	1133	-0.16	0.753	1699	97.5%	-0.09 [-0.15 , -0.03]		
Subtotal (95% CI)			1184			1751	100.0%	-0.09 [-0.15 , -0.04]	•	
Heterogeneity: Tau ² = 0	$0.00; Chi^2 = 0.$	14, $df = 1$	(P = 0.71)	; $I^2 = 0\%$					•	
Test for overall effect: Z	Z = 3.24 (P =	0.001)								
3.5.5 replacement uncl	ear									
Moy 2001	-0.69	1.1	117	-0.4	0.8	118	74.6%	-0.29 [-0.54 , -0.04]	_ 	
Simon 1997	2.79	0.82	34	3.09	0.99	37	25.4%	-0.30 [-0.72, 0.12]		
Subtotal (95% CI)			151			155	100.0%	-0.29 [-0.51 , -0.08]		
Heterogeneity: Tau ² = 0	.00; Chi ² = 0.	00, $df = 1$	(P = 0.97)	; $I^2 = 0\%$					•	
Test for overall effect: Z	Z = 2.70 (P =	0.007)								
Test for subgroup differ	ences: Chi ² =	6.44, df =	= 3 (P = 0.0	09), I ² = 53.	4%				-0.5-0.25 0 0.25 0.5	

Favours lower SFA Favours higher SFA

Analysis 3.6. Comparison 3: SFA reduction vs usual diet - secondary blood outcomes, Outcome 6: LDL, mmol/L, subgroup by main replacement

	Lo	wer SFA		н	HIgher SFA			Mean Difference	Mean Difference	
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI	
3.6.1 replaced by PUF.	A									
Subtotal (95% CI)			0			0		Not estimable		
Heterogeneity: Not appl	licable									
Test for overall effect: N	Not applicable	;								
3.6.2 replace by MUFA	4									
Subtotal (95% CI)			0			0		Not estimable		
Heterogeneity: Not appl	licable									
Test for overall effect: N	Not applicable									
3.6.3 replace by CHO										
Ley 2004	-0.32	0.64	51	-0.16	1.15	52	19.7%	-0.16 [-0.52 , 0.20]		
STARS 1992	4.19	0.51	26	4.67	0.931	24	15.5%	-0.48 [-0.90 , -0.06]	_	
WHI 2006	-0.251	0.758	1133	-0.16	0.753	1699	64.8%	-0.09 [-0.15 , -0.03]		
Subtotal (95% CI)			1210			1775	100.0%	-0.16 [-0.35 , 0.02]		
Heterogeneity: Tau ² = 0	0.01; Chi ² = 3.	33, df = 2	(P = 0.19)	; $I^2 = 40\%$					•	
Test for overall effect: 2	Z = 1.72 (P =	0.09)								
3.6.4 replace by protei	n									
Subtotal (95% CI)			0			0		Not estimable		
Heterogeneity: Not appl	licable									
Test for overall effect: N	Not applicable	;								
3.6.5 replacement uncl	lear									
Moy 2001	-0.69	1.1	117	-0.4	0.8	118	74.6%	-0.29 [-0.54 , -0.04]		
Simon 1997	2.79	0.82	34	3.09	0.99	37	25.4%	-0.30 [-0.72 , 0.12]		
Subtotal (95% CI)			151			155	100.0%	-0.29 [-0.51 , -0.08]		
Heterogeneity: Tau ² = 0	0.00; Chi ² = 0.	00, df = 1	(P = 0.97)	; $I^2 = 0\%$					-	
Test for overall effect: 2	Z = 2.70 (P =	0.007)								
Test for subgroup differ	ences: Chi ² =	0.78, df =	1 (P = 0.3	8), I ² = 0%					-0.5-0.25 0 0.25 0.5	
								Fa	avours lower SFA Favours higher	

Analysis 3.7. Comparison 3: SFA reduction vs usual diet secondary blood outcomes, Outcome 7: HDL cholesterol, mmol/L

	Le	ower SFA		н	gher SFA			Mean Difference	Mean Difference
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI
DART 1989	1.04	0.31	924	1.05	0.3	931	28.9%	-0.01 [-0.04 , 0.02]	
Ley 2004	0.01	0.14	51	0.06	0.36	52	2.0%	-0.05 [-0.16 , 0.06]	
Moy 2001	0.044	0.3	117	0.008	0.2	118	5.2%	0.04 [-0.03, 0.10]	+- -
STARS 1992	1.14	0.153	26	1.21	0.294	24	1.3%	-0.07 [-0.20, 0.06]	.
Simon 1997	1.44	0.58	34	1.56	0.55	38	0.3%	-0.12 [-0.38, 0.14]	
WHI 2006	-0.018	0.243	1133	-0.008	0.264	1699	62.2%	-0.01 [-0.03 , 0.01]	•
Total (95% CI)			2285			2862	100.0%	-0.01 [-0.02 , 0.01]	
Heterogeneity: $Tau^2 = 0$	$0.00; Chi^2 = 3$.94, df = 5	(P = 0.56)	; $I^2 = 0\%$					N
Test for overall effect: 2	Z = 1.25 (P =	0.21)							-0.2-0.1 0 0.1 0.2
Test for subgroup differ	ences: Not ap	pplicable						Fa	avours higher SFA Favours lower SFA

Analysis 3.8. Comparison 3: SFA reduction vs usual diet - secondary blood outcomes, Outcome 8: HDL, mmol/L, subgroup by any replacement

	Lo	ower SFA		HI	HIgher SFA			Mean Difference	Mean Difference
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI
3.8.1 replaced by PUF	4								
DART 1989	1.04	0.31	924	1.05	0.3	931	95.7%	-0.01 [-0.04, 0.02]	•
STARS 1992	1.14	0.153	26	1.21	0.294	24	4.3%	-0.07 [-0.20, 0.06]	
Subtotal (95% CI)			950			955	100.0%	-0.01 [-0.04 , 0.01]	▲
Heterogeneity: Tau ² = 0	.00; Chi ² = 0.	.77, $df = 1$	(P = 0.38)	; I ² = 0%					
Test for overall effect: 2	z = 0.91 (P =	0.36)							
3.8.2 replace by MUFA	`								
Subtotal (95% CI)			0			0		Not estimable	
Heterogeneity: Not appl	icable								
Test for overall effect: N	Not applicable	e							
3.8.3 replace by CHO									
DART 1989	1.04	0.31	924	1.05	0.3	931	30.6%	-0.01 [-0.04 , 0.02]	4
Ley 2004	0.01	0.14	51	0.06	0.36	52	2.1%	-0.05 [-0.16, 0.06]	
STARS 1992	1.14	0.153	26	1.21	0.294	24	1.4%	-0.07 [-0.20, 0.06]	
WHI 2006	-0.018	0.243	1133	-0.008	0.264	1699	65.9%	-0.01 [-0.03, 0.01]	
Subtotal (95% CI)			2134			2706	100.0%	-0.01 [-0.03 , 0.00]	
Heterogeneity: Tau ² = 0	.00; Chi ² = 1.	.31, $df = 3$	(P = 0.73)	; I ² = 0%					Y
Test for overall effect: 2	Z = 1.49 (P =	0.14)							
3.8.4 replace by protein	n								
DART 1989	1.04	0.31	924	1.05	0.3	931	31.0%	-0.01 [-0.04, 0.02]	
Ley 2004	0.01	0.14	51	0.06	0.36	52	2.2%	-0.05 [-0.16, 0.06]	
WHI 2006	-0.018	0.243	1133	-0.008	0.264	1699	66.8%	-0.01 [-0.03, 0.01]	
Subtotal (95% CI)			2108			2682	100.0%	-0.01 [-0.03 , 0.00]	
Heterogeneity: Tau ² = 0	.00; Chi ² = 0.	.54, $df = 2$	(P = 0.76)	; I ² = 0%					
Test for overall effect: 2	Z = 1.38 (P =	0.17)							
3.8.5 replacement uncl	ear								
Moy 2001	0.044	0.3	117	0.008	0.2	118	84.4%	0.04 [-0.03, 0.10]	
Simon 1997	1.44	0.58	34	1.56	0.55	38	15.6%	-0.12 [-0.38, 0.14]	_
Subtotal (95% CI)			151			156	100.0%	0.01 [-0.10 , 0.12]	-
Heterogeneity: $Tau^2 = 0$.00; Chi ² = 1.	.28, $df = 1$	(P = 0.26)	; I ² = 22%					T
Test for overall effect: 2	Z = 0.21 (P =	0.84)							
Test for subgroup differ	ences: Chi ² =	0.18, df =	= 3 (P = 0.9	98), I ² = 0%				-	-0.2-0.1 0 0.1 0 2

-0.2-0.1 0 0.1 0.2 Favours higher SFA Favours lower SFA

Analysis 3.9. Comparison 3: SFA reduction vs usual diet - secondary blood outcomes, Outcome 9: HDL, mmol/L, subgroup by main replacement

	Lo	Lower SFA			HIgher SFA			Mean Difference	Mean Difference	
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI	
3.9.1 replaced by PUF	A									
DART 1989	1.04	0.31	924	1.05	0.3	931	100.0%	-0.01 [-0.04 , 0.02]		
Subtotal (95% CI)			924			931	100.0%	-0.01 [-0.04 , 0.02]		
Heterogeneity: Not app	licable								1	
Test for overall effect: 2	Z = 0.71 (P =	0.48)								
3.9.2 replace by MUFA	A									
Subtotal (95% CI)			0			0		Not estimable		
Heterogeneity: Not app	licable									
Test for overall effect: 1	Not applicable	e								
3.9.3 replace by CHO										
Ley 2004	0.01	0.14	51	0.06	0.36	52	3.1%	-0.05 [-0.16 , 0.06]		
STARS 1992	1.14	0.153	26	1.21	0.294	24	2.0%	-0.07 [-0.20 , 0.06]		
WHI 2006	-0.018	0.243	1133	-0.008	0.264	1699	95.0%	-0.01 [-0.03 , 0.01]	-	
Subtotal (95% CI)			1210			1775	100.0%	-0.01 [-0.03 , 0.01]		
Heterogeneity: Tau ² = (0.00; Chi ² = 1.	29, df = 2	P = 0.52	; $I^2 = 0\%$					•	
Test for overall effect: 2	Z = 1.32 (P =	0.19)								
3.9.4 replace by protei	in									
Subtotal (95% CI)			0			0		Not estimable		
Heterogeneity: Not app	licable									
Test for overall effect: 1	Not applicable	e								
3.9.5 replacement unc	lear									
Moy 2001	0.044	0.3	117	0.008	0.2	118	84.4%	0.04 [-0.03 , 0.10]		
Simon 1997	1.44	0.58	34	1.56	0.55	38	15.6%	-0.12 [-0.38 , 0.14]	-	
Subtotal (95% CI)			151			156	100.0%	0.01 [-0.10 , 0.12]		
Heterogeneity: Tau ² = (0.00; Chi ² = 1.	28, df = 1	(P = 0.26)	; I ² = 22%						
Test for overall effect: 2	Z = 0.21 (P =	0.84)								
Test for subgroup differ	rences: Chi ² =	0.19, df =	= 2 (P = 0.9	91), $I^2 = 0\%$					-0.2-0.1 0 0.1 0.2	
								Favo	ours higher SFA Favours low	

Analysis 3.10. Comparison 3: SFA reduction vs usual diet secondary blood outcomes, Outcome 10: Triglycerides, mmol/L

	Lo	wer SFA		HIgher SFA				Mean Difference	Mean Difference
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% Cl	IV, Random, 95% CI
Houtsmuller 1979	0.79	0.6	48	1.05	0.6	48	15.9%	-0.26 [-0.50 , -0.02	2]
Ley 2004	0.37	0.71	51	0.12	1.59	52	6.0%	0.25 [-0.22 , 0.72	2]
Moy 2001	-0.4	2	117	-0.06	1.9	118	5.5%	-0.34 [-0.84 , 0.16	5]
STARS 1992	1.85	1.02	26	2.35	0.98	24	4.6%	-0.50 [-1.05 , 0.05	5]
Simon 1997	1.35	1.05	34	1.25	0.61	37	7.8%	0.10 [-0.30 , 0.50)]
Sydney Diet-Heart 1978	1.6	0.9	221	1.7	0.9	237	22.9%	-0.10 [-0.26 , 0.00	5]
WHI 2006	0.011	0.005	1133	0.011	0.003	1699	37.5%	0.00 [-0.00 , 0.00	D] •
Total (95% CI)			1630			2215	100.0%	-0.08 [-0.21 , 0.04	4]
Heterogeneity: Tau ² = 0.01;	Chi ² = 12.13,	df = 6 (P	= 0.06); I ²	= 51%					•
Test for overall effect: $Z = 1$.	.29 (P = 0.20))							-1 -0.5 0 0.5 1
Test for subgroup differences	s: Not applica	able							Favours lower SFA Favours higher SFA

Analysis 3.11. Comparison 3: SFA reduction vs usual diet - secondary blood outcomes, Outcome 11: TG, mmol/L, subgroup by any replacement

	L	ower SFA		HI	gher SFA			Mean Difference	Mean Difference
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI
3.11.1 replaced by PUFA									
Houtsmuller 1979	0.79	0.6	48	1.05	0.6	48	34.8%	-0.26 [-0.50 , -0.02]
STARS 1992	1.85	1.02	26	2.35	0.98	24	8.3%	-0.50 [-1.05, 0.05	
Sydney Diet-Heart 1978	1.6	0.9	221	1.7	0.9	237	56.9%	-0.10 [-0.26, 0.06	
Subtotal (95% CI)			295			309	100.0%	-0.19 [-0.35 , -0.02	i 📥
Heterogeneity: $Tau^2 = 0.01$;	Chi ² = 2.59,	df = 2 (P =	0.27); I ² =	= 23%					- •
Test for overall effect: $Z = 2$.24 (P = 0.03)	3)							
3.11.2 replace by MUFA									
Subtotal (95% CI)			0			0		Not estimable	e
Heterogeneity: Not applicab	le								
Test for overall effect: Not a	pplicable								
3.11.3 replace by CHO									
Ley 2004	0.37	0.71	51	0.12	1.59	52	22.8%	0.25 [-0.22, 0.72	
STARS 1992	1.85	1.02	26	2.35	0.98	24	18.6%	-0.50 [-1.05, 0.05	
WHI 2006	0.011	0.005	1133	0.011	0.003	1699	58.7%	0.00 [-0.00 , 0.00	
Subtotal (95% CI)			1210			1775	100.0%	-0.04 [-0.32 , 0.25	i 📥
Heterogeneity: $Tau^2 = 0.04$;	Chi ² = 4.19,	df = 2 (P =	0.12); I ² =	= 52%					
Test for overall effect: $Z = 0$.24 (P = 0.81))							
3.11.4 replace by protein									
Ley 2004	0.37	0.71	51	0.12	1.59	52	3.2%	0.25 [-0.22, 0.72	1
WHI 2006	0.011	0.005	1133	0.011	0.003	1699	96.8%	0.00 [-0.00 , 0.00	
Subtotal (95% CI)			1184			1751	100.0%	0.01 [-0.08 , 0.09	1 👗
Heterogeneity: $Tau^2 = 0.00$;	Chi ² = 1.07,	df = 1 (P =	: 0.30); I ² =	= 6%					-
Test for overall effect: $Z = 0$.18 (P = 0.86)	ō)							
3.11.5 replacement unclear									
Moy 2001	-0.4	2	117	-0.06	1.9	118	44.2%	-0.34 [-0.84 , 0.16]
Simon 1997	1.35	1.05	34	1.25	0.61	37	55.8%	0.10 [-0.30 , 0.50]
Subtotal (95% CI)			151			155	100.0%	-0.09 [-0.52 , 0.33	
Heterogeneity: $Tau^2 = 0.04$;	Chi ² = 1.80,	df = 1 (P =	0.18); I ² =	= 45%				. ,	
Test for overall effect: $Z = 0$.43 (P = 0.66	ō)							
Test for subgroup difference	s: Chi² = 4.3	6, df = 3 (F	P = 0.22), I	¹² = 31.2%					-+ -+ -+ -+ -1 -0.5 0 0.5 1 Favours lower SFA Favours higher SFA

Analysis 3.12. Comparison 3: SFA reduction vs usual diet - secondary blood outcomes, Outcome 12: TG, mmol/L, subgroup by main replacement

	Lo	Lower SFA			HIgher SFA			Mean Difference	Mean Difference	
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% Cl	IV, Random, 95% CI	
3.12.1 replaced by PUFA										
Houtsmuller 1979	0.79	0.6	48	1.05	0.6	48	34.5%	-0.26 [-0.50 , -0.02	2]	
Sydney Diet-Heart 1978	1.6	0.9	221	1.7	0.9	237	65.5%	-0.10 [-0.26 , 0.00	5]	
Subtotal (95% CI)			269			285	100.0%	-0.16 [-0.30 , -0.0]	l] 📥	
Heterogeneity: Tau ² = 0.00;	Chi ² = 1.16, d	f = 1 (P = 1)	= 0.28); I ² =	= 14%					•	
Test for overall effect: $Z = 2$	2.04 (P = 0.04))								
3.12.2 replace by MUFA										
Subtotal (95% CI)			0			0		Not estimab	le	
Heterogeneity: Not applicat	ole									
Test for overall effect: Not a	applicable									
3.12.3 replace by CHO										
Ley 2004	0.37	0.71	51	0.12	1.59	52	22.8%	0.25 [-0.22 , 0.72	2]	
STARS 1992	1.85	1.02	26	2.35	0.98	24	18.6%	-0.50 [-1.05 , 0.05	5]	
WHI 2006	0.011	0.005	1133	0.011	0.003	1699	58.7%	0.00 [-0.00 , 0.00)]	
Subtotal (95% CI)			1210			1775	100.0%	-0.04 [-0.32 , 0.25	5] 📥	
Heterogeneity: Tau ² = 0.04;	Chi ² = 4.19, d	f = 2 (P = 2)	= 0.12); I ² =	= 52%						
Test for overall effect: $Z = 0$	0.24 (P = 0.81))								
3.12.4 replace by protein										
Subtotal (95% CI)			0			0		Not estimab	le	
Heterogeneity: Not applicat Test for overall effect: Not a	ole applicable									
3.12.5 replacement unclea	r									
Moy 2001	-0.4	2	117	-0.06	1.9	118	44.2%	-0.34 [-0.84 , 0.10	5]	
Simon 1997	1.35	1.05	34	1.25	0.61	37	55.8%	0.10 [-0.30 , 0.50		
Subtotal (95% CI)			151			155	100.0%	-0.09 [-0.52 , 0.3.	3]	
Heterogeneity: Tau ² = 0.04;	Chi ² = 1.80, d	lf = 1 (P =	= 0.18); I ² =	= 45%				- /		
Test for overall effect: $Z = 0$	0.43 (P = 0.66)									
Test for subgroup difference	es: Chi ² = 0.54	, df = 2 (I	P = 0.76), I	$1^2 = 0\%$					-1 -0.5 0 0.5 1	
									Favours lower SFA Favours higher S	

Analysis 3.13. Comparison 3: SFA reduction vs usual diet secondary blood outcomes, Outcome 13: total cholesterol /HDL ratio

	Lower SFA			HIgher SFA				Mean Difference	Mean Difference	
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI	
Ley 2004	-0.34	1	51	-0.53	1.73	52	14.6%	0.19 [-0.35 , 0.73]		
STARS 1992	5.5	0.918	26	6.08	1.666	24	8.3%	-0.58 [-1.33 , 0.17]	←	
WHI 2006	-0.2	0.8	1133	-0.1	1	1699	77.0%	-0.10 [-0.17 , -0.03]		
Total (95% CI)			1210			1775	100.0%	-0.10 [-0.33 , 0.13]		
Heterogeneity: $Tau^2 = 0$	$0.02; Chi^2 = 2.$.65, df = 2	P = 0.27	; I ² = 24%						
Test for overall effect: 2	Z = 0.83 (P =	0.40)							-0.5-0.25 0 0.25 0.5	
Test for subgroup differ	rences: Not ap	plicable						F	avours lower SFA Favours higher SFA	

Analysis 3.14. Comparison 3: SFA reduction vs usual diet - secondary blood outcomes, Outcome 14: TC /HDL ratio, subgroup by any replacement

	Lower SFA			HIgher SFA				Mean Difference	Mean Difference		
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI		
3.14.1 replaced by PU	FA										
STARS 1992	5.5	0.918	26	6.08	1.666	24	100.0%	-0.58 [-1.33 , 0.17]	← ■ ───────────────────────────────────		
Subtotal (95% CI)			26			24	100.0%	-0.58 [-1.33 , 0.17]			
Heterogeneity: Not app	licable										
Test for overall effect: 2	Z = 1.51 (P =	0.13)									
3.14.2 replace by MUI	FA										
Subtotal (95% CI)			0			0		Not estimable			
Heterogeneity: Not app	licable										
Test for overall effect: 1	Not applicable	e									
3.14.3 replace by CHC)										
Ley 2004	-0.34	1	51	-0.53	1.73	52	14.6%	0.19 [-0.35 , 0.73]			
STARS 1992	5.5	0.918	26	6.08	1.666	24	8.3%	-0.58 [-1.33 , 0.17]			
WHI 2006	-0.2	0.8	1133	-0.1	1	1699	77.0%	-0.10 [-0.17 , -0.03]			
Subtotal (95% CI)			1210			1775	100.0%	-0.10 [-0.33 , 0.13]			
Heterogeneity: Tau ² = ().02; Chi ² = 2.	65, $df = 2$	(P = 0.27)	; I ² = 24%							
Test for overall effect: 2	Z = 0.83 (P =	0.40)									
3.14.4 replace by prote	ein										
Ley 2004	-0.34	1	51	-0.53	1.73	52	4.8%	0.19 [-0.35 , 0.73]			
WHI 2006	-0.2	0.8	1133	-0.1	1	1699	95.2%	-0.10 [-0.17 , -0.03]			
Subtotal (95% CI)			1184			1751	100.0%	-0.09 [-0.21 , 0.04]			
Heterogeneity: Tau ² = (0.00; Chi ² = 1.	07, df = 1	(P = 0.30)	; I ² = 7%					•		
Test for overall effect: 2	Z = 1.39 (P =	0.17)									
3.14.5 replacement un	clear										
Subtotal (95% CI)			0			0		Not estimable			
Heterogeneity: Not app	licable										
Test for overall effect: 1	Not applicable										
Test for subgroup differ	rences: Chi ² =	1.61, df =	= 2 (P = 0.4)	5), $I^2 = 0\%$					-0.5-0.25 0 0.25 0.5		
								Fa	avours lower SFA Favours highe		

Analysis 3.15. Comparison 3: SFA reduction vs usual diet - secondary blood outcomes, Outcome 15: TC /HDL ratio, subgroup by main replacement

	Lo	Lower SFA			HIgher SFA			Mean Difference	Mean Difference
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI
3.15.1 replaced by PUFA	1								
Subtotal (95% CI)			0			0		Not estimable	
Heterogeneity: Not applic	able								
Test for overall effect: No	ot applicable	;							
3.15.2 replace by MUFA									
Subtotal (95% CI)			0			0		Not estimable	
Heterogeneity: Not applic	able								
Test for overall effect: No	ot applicable	;							
3.15.3 replace by CHO									
Ley 2004	-0.34	1	51	-0.53	1.73	52	14.6%	0.19 [-0.35 , 0.73]	
STARS 1992	5.5	0.918	26	6.08	1.666	24	8.3%	-0.58 [-1.33 , 0.17]	←
WHI 2006	-0.2	0.8	1133	-0.1	1	1699	77.0%	-0.10 [-0.17 , -0.03]	
Subtotal (95% CI)			1210			1775	100.0%	-0.10 [-0.33 , 0.13]	
Heterogeneity: Tau ² = 0.0	2; Chi ² = 2.	65, df = 2 (P = 0.27)	; I ² = 24%					-
Test for overall effect: Z =	= 0.83 (P =	0.40)							
3.15.4 replace by protein	1								
Subtotal (95% CI)			0			0		Not estimable	
Heterogeneity: Not applic	able								
Test for overall effect: No	ot applicable								
3.15.5 replacement uncle	ear								
Subtotal (95% CI)			0			0		Not estimable	
Heterogeneity: Not applic	able								
Test for overall effect: No	ot applicable								
Test for subgroup differen	nces: Not ap	plicable							-0.5-0.25 0 0.25 0.5
								F	avours lower SFA Favours highe

Analysis 3.16. Comparison 3: SFA reduction vs usual diet - secondary blood outcomes, Outcome 16: LDL /HDL ratio

	Lower SFA			н	gher SFA		Mean Difference	Mean Difference
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	IV, Random, 95% CI	IV, Random, 95% CI
STARS 1992	3.74	0.71	26	4.1	1.22	24	-0.36 [-0.92 , 0.20]	
Test for subgroup differe	nces: Not ap	oplicable					Fa	-0.5-0.25 0 0.25 0.5 vours lower SFA Favours higher SFA

Analysis 3.17. Comparison 3: SFA reduction vs usual diet - secondary blood outcomes, Outcome 17: Lp(a), mmol/L

	Le	Lower SFA			HIgher SFA			Mean Difference	Mean Difference		
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI		
STARS 1992	0.96	1.89	26	0.96	2.89	24	0.0%	0.00 [-1.37 , 1.37]			
WHI 2006	0.03	0.02	1133	0.03	0.02	1699	100.0%	0.00 [-0.00 , 0.00]			
Total (95% CI)			1159			1723	100.0%	0.00 [-0.00 , 0.00]			
Heterogeneity: Tau ² = 0	$0.00; Chi^2 = 0.00;$.00, $df = 1$	(P = 1.00)	; $I^2 = 0\%$							
Test for overall effect: 2	Z = 0.00 (P =	1.00)							-1 -0.5 0 0.5 1		
Test for subgroup differences: Not applicable								Fa	vours lower SFA Favours higher SF		

Analysis 3.18. Comparison 3: SFA reduction vs usual diet - secondary blood outcomes, Outcome 18: Lp(a), mmol/L, subgroup by any replacement

	Lo	Lower SFA			HIgher SFA			Mean Difference	Mean Difference	
Study or Subgroup	Mean	Mean SD		Total Mean		SD Total		IV, Random, 95% CI	IV, Random, 95% CI	
3.18.1 replaced by PU	FA									
STARS 1992	0.96	1.89	26	0.96	2.89	24	100.0%	0.00 [-1.37 , 1.37]		
Subtotal (95% CI)			26			24	100.0%	0.00 [-1.37 , 1.37]		
Heterogeneity: Not app	olicable									
Test for overall effect: 2	Z = 0.00 (P =	1.00)								
3.18.2 replace by MUI	FA									
Subtotal (95% CI)			0			0		Not estimable		
Heterogeneity: Not app	olicable									
Test for overall effect: 1	Not applicable	e								
3.18.3 replace by CHC)									
STARS 1992	0.96	1.89	26	0.96	2.89	24	0.0%	0.00 [-1.37 , 1.37]		
WHI 2006	0.03	0.02	1133	0.03	0.02	1699	100.0%	0.00 [-0.00 , 0.00]		
Subtotal (95% CI)			1159			1723	100.0%	0.00 [-0.00 , 0.00]	Т	
Heterogeneity: Tau ² = 0	$0.00; Chi^2 = 0.$	00, $df = 1$	(P = 1.00)	; $I^2 = 0\%$						
Test for overall effect: 2	Z = 0.00 (P =	1.00)								
3.18.4 replace by prot	ein									
WHI 2006	0.03	0.02	1133	0.03	0.02	1699	100.0%	0.00 [-0.00 , 0.00]		
Subtotal (95% CI)			1133			1699	100.0%	0.00 [-0.00 , 0.00]	T	
Heterogeneity: Not app	olicable									
Test for overall effect:	Z = 0.00 (P =	1.00)								
3.18.5 replacement un	clear									
Subtotal (95% CI)			0			0		Not estimable		
Heterogeneity: Not app	olicable									
Test for overall effect:	Not applicable	2								
Test for subgroup differ	rences: Chi ² =	0.00, df =	= 2 (P = 1.0	0), $I^2 = 0\%$					-2 -1 0 1 2	
								Fa	avours lower SFA Favours hig	

Analysis 3.19. Comparison 3: SFA reduction vs usual diet - secondary blood outcomes, Outcome 19: Lp(a), mmol/L, subgroup by main replacement

	Lo	Lower SFA			HIgher SFA			Mean Difference	Mean Difference
Study or Subgroup	Mean	SD	Total	Total Mean		SD Total		IV, Random, 95% CI	IV, Random, 95% CI
3.19.1 replaced by PU	FA								
Subtotal (95% CI)			0			0		Not estimable	,
Heterogeneity: Not app	licable								
Test for overall effect:	Not applicable	e							
3.19.2 replace by MUI	FA								
Subtotal (95% CI)			0			0		Not estimable	
Heterogeneity: Not app	olicable								
Test for overall effect: 1	Not applicable	e							
3.19.3 replace by CHC)								
STARS 1992	0.96	1.89	26	0.96	2.89	24	0.0%	0.00 [-1.37 , 1.37]	I
WHI 2006	0.03	0.02	1133	0.03	0.02	1699	100.0%	0.00 [-0.00 , 0.00]	
Subtotal (95% CI)			1159			1723	100.0%	0.00 [-0.00 , 0.00]	
Heterogeneity: Tau ² = 0	$0.00; Chi^2 = 0.$.00, $df = 1$	(P = 1.00)	; $I^2 = 0\%$					
Test for overall effect: 2	Z = 0.00 (P =	1.00)							
3.19.4 replace by prot	ein								
Subtotal (95% CI)			0			0		Not estimable	
Heterogeneity: Not app	olicable								
Test for overall effect: 1	Not applicable	e							
3.19.5 replacement un	clear								
Subtotal (95% CI)			0			0		Not estimable	
Heterogeneity: Not app	olicable								
Test for overall effect: I	Not applicable	e							
Test for subgroup differ	rences: Not ap	plicable							-2 -1 0 1 2
								I	Favours lower SFA Favours highe

Analysis 3.20. Comparison 3: SFA reduction vs usual diet - secondary blood outcomes, Outcome 20: Insulin sensitivity

	L	Lower SFA			HIgher SFA			Mean Difference	Mean Difference		
Study or Subgroup	Mean	Mean SD Total		Mean	SD	SD Total		IV, Random, 95% CI	IV, Random, 95% CI		
3.20.1 HbA1c (glycosy	lated haemo	globin), %	,								
Subtotal (95% CI)			0			0		Not estimable			
Heterogeneity: Not appl	licable										
Test for overall effect: N	Not applicabl	e									
3.20.2 GTT (glucose to	lerance test)), glucose a	at 2 hours	, mmol/L							
Houtsmuller 1979	7.7	2.5	48	10.2	2.7	48	35.8%	-2.50 [-3.54 , -1.46]	_		
Ley 2004	1.02	2.9	51	2.3	3.9	52	27.0%	-1.28 [-2.61 , 0.05]		-	
STARS 1992	4.6	1.7	26	5.8	1.9	24	37.2%	-1.20 [-2.20 , -0.20]			
Subtotal (95% CI)			125			124	100.0%	-1.69 [-2.55 , -0.82]			
Heterogeneity: $Tau^2 = 0$.26; Chi ² = 3	.61, df = 2	(P = 0.16)); I ² = 45%					-		
Test for overall effect: 2	Z = 3.83 (P =	0.0001)									
3.20.3 HOMA											
WHI 2006	1.1	0.5	1133	1.1	0.6	1699	100.0%	0.00 [-0.04 , 0.04]			
Subtotal (95% CI)			1133			1699	100.0%	0.00 [-0.04 , 0.04]		T	
Heterogeneity: Not appl	licable										
Test for overall effect: 2	Z = 0.00 (P =	1.00)									
Test for subgroup differ	ences: Chi ² =	= 14.65, df =	= 1 (P = 0)	0.0001), I ² =	93.2%				-2 -1		
								Fa	wours lower SFA	Favours higher S	

Outcome or subgroup title	No. of studies	No. of partici- pants	Statistical method	Effect size
4.1 Cancer diagnoses	4	52294	Risk Ratio (M-H, Random, 95% CI)	0.94 [0.83, 1.07]
4.2 Cancer deaths	5	52283	Risk Ratio (M-H, Random, 95% CI)	1.00 [0.61, 1.64]
4.3 Weight, kg	6	43062	Mean Difference (IV, Random, 95% CI)	-1.77 [-3.54, -0.01]
4.4 BMI, kg/m2	6	43894	Mean Difference (IV, Random, 95% CI)	-0.42 [-0.72, -0.12]
4.5 Systolic Blood Pressure, mmHg	5	3812	Mean Difference (IV, Random, 95% CI)	-0.19 [-1.36, 0.97]
4.6 Diastolic Blood Pressure, mmHg	5	3812	Mean Difference (IV, Random, 95% CI)	-0.36 [-1.03, 0.32]
4.7 Quality of Life	1	40130	Mean Difference (IV, Random, 95% CI)	0.04 [0.01, 0.07]

Comparison 4. SFA reduction vs usual diet - secondary outcomes including potential adverse effects

Analysis 4.1. Comparison 4: SFA reduction vs usual diet - secondary outcomes including potential adverse effects, Outcome 1: Cancer diagnoses

	Lower	SFA	Higher SFA			Risk Ratio	Risk	Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Rando	om, 95% CI
Ley 2004	3	88	3	88	0.6%	1.00 [0.21 , 4.82]]	
Veterans Admin 1969	70	424	59	422	12.8%	1.18 [0.86 , 1.62]] _	-
WHI 2006	1946	19541	3040	29294	61.7%	0.96 [0.91 , 1.01]]	
WINS 2006	124	975	231	1462	24.9%	0.80 [0.66 , 0.99]] –	
Total (95% CI)		21028		31266	100.0%	0.94 [0.83 , 1.07]	1	,
Total events:	2143		3333				•	
Heterogeneity: $Tau^2 = 0.02$	1; Chi ² = 4.4	7, df = 3 (P = 0.22; I	$^{2} = 33\%$			0.1 0.2 0.5 1	2 5 10
Test for overall effect: Z =	0.91 (P = 0	.37)]	Favours lower SFA	Favours higher SFA
Test for subgroup differen	ces: Not app	licable						



Analysis 4.2. Comparison 4: SFA reduction vs usual diet - secondary outcomes including potential adverse effects, Outcome 2: Cancer deaths

	Lower	SFA	Higher SFA			Risk Ratio	Ris	Risk Ratio					
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Ran	dom, 95% CI					
DART 1989	4	1018	6	1015	11.7%	0.66 [0.19 , 2.35]						
Ley 2004	1	88	2	88	3.9%	0.50 [0.05 , 5.41]						
MRC 1968	1	199	6	194	4.9%	0.16 [0.02 , 1.34	·]	+					
Veterans Admin 1969	31	424	17	422	29.8%	1.81 [1.02 , 3.23]						
WHI 2006	950	19541	1454	29294	49.6%	0.98 [0.90 , 1.06]	•					
Total (95% CI)		21270		31013	100.0%	1.00 [0.61 , 1.64] .						
Total events:	987		1485					Ť					
Heterogeneity: $Tau^2 = 0.13$; $Chi^2 = 7.84$, $df = 4$ (P = 0.10); $I^2 = 49\%$							0.01 0.1	1 10 100					
Test for overall effect: Z =	0.01 (P = 0	.99)					Favours lower SFA	Favours higher SFA					

Test for subgroup differences: Not applicable

Analysis 4.3. Comparison 4: SFA reduction vs usual diet - secondary outcomes including potential adverse effects, Outcome 3: Weight, kg

	Lower SFA			Higher SFA				Mean Difference	Mean Difference
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI
Ley 2004	1.06	4.57	51	1.26	4.9	52	26.9%	-0.20 [-2.03 , 1.63]
MRC 1968	0	0	88	-3	0	89		Not estimable	e
Oslo Diet-Heart 1966	-2.5	0	168	-0.5	0	161		Not estimable	e
Simon 1997	63.4	11.1	34	71.9	11.7	38	8.7%	-8.50 [-13.77 , -3.23] ←=
WHI 2006 (1)	-0.8	10.1	16297	-0.1	10.1	25056	37.7%	-0.70 [-0.90 , -0.50]
WINS 2006	70.6	14.6	380	73.3	14.9	648	26.7%	-2.70 [-4.56 , -0.84	-1 _
Total (95% CI) 17018						26044	100.0%	-1.77 [-3.54 , -0.01]
Heterogeneity: $Tau^2 = 2.1$	4; Chi ² = 13.	06, df = 3	(P = 0.005)	5); $I^2 = 77\%$,				•
Test for overall effect: Z =	= 1.97 (P = 0)	.05)							-10 -5 0 5 10
Test for subgroup differer	nces: Not app	olicable							Favours lower SFA Favours higher SFA

Footnotes

(1) Change from baseline to 7,5 years

Analysis 4.4. Comparison 4: SFA reduction vs usual diet - secondary outcomes including potential adverse effects, Outcome 4: BMI, kg/m2

	Le	Lower SFA			Higher SFA			Mean Difference	Mean Difference
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI
Moy 2001	-0.1	1	117	0.21	2	118	21.4%	-0.31 [-0.71 , 0.09]
Oxford Retinopathy 1978	-1	2.8	29	-0.7	1.8	29	5.3%	-0.30 [-1.51 , 0.91]
Simon 1997	23.8	4.7	34	27.4	4.9	38	1.8%	-3.60 [-5.82 , -1.38]
Sydney Diet-Heart 1978	24.3	1.5	179	24.5	2.8	192	19.5%	-0.20 [-0.65 , 0.25]
WHI 2006 (1)	0.03	3.2	16230	0.3	3.1	24943	34.3%	-0.27 [-0.33 , -0.21] 🗧
WINS 2006	26.8	5.608	755	27.6	5.368	1230	17.8%	-0.80 [-1.30 , -0.30]
Total (95% CI)			17344			26550	100.0%	-0.42 [-0.72 , -0.12]
Heterogeneity: $Tau^2 = 0.07$; C	Chi ² = 13.00, d	f = 5 (P =	0.02); I ² =	: 62%					•
Test for overall effect: $Z = 2.2$	72 (P = 0.006))							-4 -2 0 2 4
Test for subgroup differences	: Not applicab	ole							Favours lower SFA Favours higher SFA

Footnotes

(1) Change to 7.5 years

Cochrane Library

Trusted evidence. Informed decisions. Better health.

Analysis 4.5. Comparison 4: SFA reduction vs usual diet - secondary outcomes including potential adverse effects, Outcome 5: Systolic Blood Pressure, mmHg

	Lo	Lower SFA			Higher SFA			Mean Difference	Mean Difference
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI
Ley 2004	-3.5	17.71	51	1.31	24.37	52	2.0%	-4.81 [-13.03 , 3.41	
MRC 1968	2	0	88	0	0	89		Not estimabl	le
Oslo Diet-Heart 1966	158.2	0	168	154.3	0	161		Not estimabl	le
Sydney Diet-Heart 1978	136.4	17.8	179	136.5	21.6	192	8.4%	-0.10 [-4.12 , 3.92	2]
WHI 2006	-2.2	16.3	1133	-2.1	16.4	1699	89.6%	-0.10 [-1.33 , 1.13	B]
Total (95% CI)			1619			2193	100.0%	-0.19 [-1.36 , 0.97	7]
Heterogeneity: Tau ² = 0.00;	Chi ² = 1.24, d	df = 2 (P =	= 0.54); I ² =	= 0%					T
Test for overall effect: $Z = 0$	0.33 (P = 0.74))							-++++++
Test for subgroup difference							Favours lower SFA Favours higher SFA		

Analysis 4.6. Comparison 4: SFA reduction vs usual diet - secondary outcomes including potential adverse effects, Outcome 6: Diastolic Blood Pressure, mmHg

	Lo	Lower SFA			Higher SFA			Mean Difference	Mean Difference		
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI		
Ley 2004	-7.16	12	51	-4.2	13.85	52	1.8%	-2.96 [-7.96 , 2.04]			
MRC 1968	-1	0	88	3	0	89		Not estimable			
Oslo Diet-Heart 1966	98.6	0	168	95.5	0	161		Not estimable			
Sydney Diet-Heart 1978	87.5	12.3	179	87.9	13.8	192	6.5%	-0.40 [-3.06 , 2.26]			
WHI 2006	-2.6	9.4	1133	-2.3	9.4	1699	91.7%	-0.30 [-1.01 , 0.41]			
Total (95% CI)			1619			2193	100.0%	-0.36 [-1.03 , 0.32]			
Heterogeneity: Tau ² = 0.00;	Chi ² = 1.07, d	f = 2 (P = 2)	= 0.59); I ² =	= 0%					•		
Test for overall effect: $Z = 1$.03 (P = 0.30)							-4 -2 0 2 4		
est for subgroup differences: Not applicable								F	Favours lower SFA Favours higher		

Analysis 4.7. Comparison 4: SFA reduction vs usual diet - secondary outcomes including potential adverse effects, Outcome 7: Quality of Life

	Lower SFA			HIgher SFA				Mean Difference	Mean D	lifference
Study or Subgroup	Mean SD		Total	Mean	SD	Total	Weight	IV, Random, 95% CI	IV, Rando	om, 95% CI
WHI 2006 (1)	0.07	1.41	15788	0.03	1.44	24342	100.0%	0.04 [0.01 , 0.07]		-
Total (95% CI)			15788			24342	100.0%	0.04 [0.01 , 0.07]		•
Heterogeneity: Not appl	icable									
Test for overall effect: Z							-0.2 -0.1	0 0.1 0.2		
Test for subgroup differences: Not applicable								F	Favours lower SFA	Favours higher SFA

Footnotes

(1) Change in Global Quality of Life to trial close-out (0 worst to 10 best), Assaf 2016

ADDITIONAL TABLES

Reference	Population	CVD risk category	Is interven- tion deliv- ered to In- dividual or group?	interven- tion given by?	Face-to- face or oth- er?	Number of visits	Is intervention advice only or other interven- tion?
Black 1994	People with non-melanoma skin cancer	Low	Unclear	Dietitian	Face-to-face	8 x weekly classes then monthly follow-up sessions	Advice (behaviour tech- niques learning)
DART 1989	Men recovering from a MI	High	Individual	Dietitian	Face-to-face	9	Advice (diet advice, recipes and encourage- ment)
Houtsmuller 1979	Adults with newly-diag- nosed diabetes	Moderate	Unclear	Dietitian	Unclear	Unclear	Advice?
Ley 2004	People with impaired glu- cose intolerance or high normal blood glucose	Moderate	Small group	Unclear	Face-to-face	Monthly meetings	Advice (education, per- sonal goal-setting, self- monitoring)
Moy 2001	Middle-aged siblings of peo- ple with early CHD, with at least 1 CVD risk factor	Moderate	Individual	Trained nurse	Face-to-face	6 - 8 weekly for 2 years	Advice (individualised counselling sessions)
MRC 1968	Free-living men who have survived a 1st MI	High	Individual	Dietitian	Face-to-face	Unclear	Advice and supplement (soy oil)
Oslo Di- et-Heart 1966	Men with previous MI	High	Individual	Dietitian	Face-to-face and other	Unclear	Advice and supplement (food)
Oxford Retinopa- thy 1978	Newly-diagnosed non-in- sulin-dependent diabetics	Moderate	Individual	Diabetes di- etitian	Face-to-face	After 1 month then at 3-month intervals	Advice
Rose corn oil 1965	Men (?) with angina or fol- lowing MI	High	Unclear	Unclear	Unclear	Follow-up clinic monthly, then every 2 months	Advice and supplement (oil)
Rose olive 1965	Men (?) with angina or fol- lowing MI	High	Unclear	Unclear	Unclear	Follow-up clinic monthly, then every 2 months	Advice and supplement (oil)
Simon 1997	Women with a high risk of breast cancer	Low	Individual followed by	Dietitian	Face-to-face	Bi-weekly over 3 months fol- lowed by monthly	Advice (individualised of ing plan and counsellin sessions)

Cochrane Library

Trusted evidence. Informed decisions. Better health.

			individual or group				
STARS 1992	Men with angina referred for angiography	High	Individual	Dietitian	Face-to-face	Clinic visits at 3-month inter- vals	Advice
Sydney Di- et-Heart 1978	Men with angina referred for angiography	High	Individual	Unclear	Face-to-face	3 times in 1st year and twice an- nually thereafter	Advice
Veterans Admin 1969	Men living at the Veterans Administration Center	Low	Individual	Unclear (whole diet provided)	N/A	N/A	Diet provided
WHI 2006	Postmenopausal women aged 50 - 79 with or without CVD at baseline	Low and High	Group	Nutritionists	Face-to-face	18 sessions/1st yr and quarterly maintenance sessions after	Advice
WINS 2006	Women with localised re- sected breast cancer	Low	Individual followed by group	Dietitian	Face-to-face	8 bi-weekly sessions, then 3- monthly contact and optional monthly sessions	Advice

MI: myocardial infarction

N/A: not applicable

Table 2. Number of participants and number of outcomes for dichotomous variables (by intervention arm)

Table 1. Comparison of study interventions for included RCTs (Continued)

	Participants	All-cause mortality	CV mor- tality	CVD events	МІ	Non- fatal MI	Stroke	CHD mortal- ity	CHD events	Dia- betes Diag- noses
Black 1994	133	133	133	133	0	0	0	0	0	0
DART 1989	2033	2033	2033	2033	2033	2033	0	2033	2033	0
Houtsmuller 1979	102	0	0	102	102	0	0	102	102	0
Ley 2004	176	176	176	176	176	0	176	0	176	0
Moy 2001	267	0	0	235	235	235	235	0	267	0

•,**4**µ0•

Cochrane Library

Trusted evidence. Informed decisions. Better health.

able 2. Number of participants and number of outcomes for dichotomous variables (by intervention arm) (Continued)										
MRC 1968	393	393	393	393	393	393	393	393	393	0
Oslo Diet-Heart 1966	412	412	412	412	412	412	412	412	412	0
Oxford Retinopathy 1978	249 (data not provid- ed by arm)	0	0	0	0	0	0	0	0	0
Rose corn oil 1965	41	41	41	41	41	41	0	41	41	0
Rose olive 1965	39	39	39	39	39	39	0	39	39	0
Simon 1997	194 (data not provid- ed by arm)	0	0	0	0	0	0	0	0	0
STARS 1992	60	55	55	55	55	0	55	0	55	0
Sydney Diet-Heart 1978	458	458	458	0	0	0	0	458	0	0
Veterans Admin 1969	846	846	846	846	846	846	846	846	846	0
WHI with CVD 2006	2277	0	2277	2277	0	2277	2277	2277	2277	0
WHI 2006	48,835	48,835	46,558	46,558	48,835	46,558	46,558	46,558	46,558	48,835
WINS 2006	2437	2437	0	0	0	0	0	0	0	0
Total Participants	58,509	55,858	53,421	53,300	53,167	52,834	50,952	53,159	53,204	48,835
Percent of participants for this outcome	100%	95%	91%	91%	91%	90%	87%	91%	91%	83%

These numbers are the numbers of participants in each study who were available for assessment of outcomes within meta-analysis (not necessarily the number of participants randomised within the trial).

CHD: coronary heart disease

CV: cardiovascular

CVD: cardiovascular disease

Table 3.	Number of participants and number	er of participants with data for co	ontinuous outcomes (by intervention arm)
----------	-----------------------------------	-------------------------------------	--

Participants	Total cho- lesterol	LDL cho-	HDL cho-	Triglyc- erides	TG/ HDL	Total choles-	LDL/ HDL ratio	LP (a)	Insulin sensi- tivity	
--------------	------------------------	-------------	-------------	--------------------	------------	------------------	----------------------	-----------	-----------------------------	--

Table 3. Number of participants and number of	of participants with data for continu		lous outcomes (by intervent les- les- terol terol			ion arm) (Continued): ra- terol/HDL tio ratio		d) DL		
Black 1994	133	0	0	0	0	0	0	0	0	0
DART 1989	2033	1855	0	1855	0	0	0	0	0	0
Houtsmuller 1979	102	96	0	0	96	0	0	0	0	96
Ley 2004	176	103	103	103	103	0	103	0	0	103
Moy 2001	267	0	235	235	235	0	0	0	0	0
MRC 1968	393	177	0	0	0	0	0	0	0	0
Oslo Diet-Heart 1966	412	329	0	0	0	0	0	0	0	0
Oxford Retinopathy 1978	249	58	0	0	0	0	0	0	0	0
Rose corn oil 1965	41	22	0	0	0	0	0	0	0	0
Rose olive 1965	39	24	0	0	0	0	0	0	0	0
Simon 1997	194	72	71	72	71	0	0	0	0	0
STARS 1992	55	50	50	50	50	0	50	50	50	50
Sydney Diet-Heart 1978	458	458	0	0	458	0	0	0	0	0
Veterans Admin 1969	846	843	0	0	0	0	0	0	0	0
WHI with CVD 2006	2277	0	0	0	0	0	0	0	0	0
WHI 2006	48,835	2832	2832	2832	2832	0	2832	0	2832	2832
WINS 2006	2437	196	0	0	0	0	0	0	0	0
Total Participants	58952	7115	3291	5147	3845	0	2985	50	2882	3081
Percent of participants for this outcome	100%	12%	6%	9%	7%	0%	5%	0.1%	5%	5%

These numbers are the numbers of participants in each study who were available for assessment of outcomes within meta-analysis (not necessarily the number of participants randomised within the trial). HDL: high density lipoprotein

LDL: low density lipoprotein Lp(a): lipoprotein (a)

TG: triglyceride

Regression factor	No. of studies	Constant	Coefficient (95% CI)	P value	Proportion of between study vari- ation ex- plained
Change in SFA as %E	8	0.01	0.05 (-0.03 to 0.13)	0.16	89%
Change in SFA as % of control	8	0.26	0.01 (-0.01 to 0.03)	0.14	89%
Baseline SFA as %E	8	0.68	-0.06 (-0.15 to 0.04)	0.19	81%
Change in TC, mmol/L	12	0.03	0.69 (0.05 to 1.33)	0.04	99%
Change in PUFA as %E	5	-0.01	-0.02 (-0.08 to 0.03)	0.25	100%
Change in MUFA as %E	5	-0.26	-0.03 (-0.14 to 0.09)	0.50	-87%
Change in CHO as %E	7	-0.11	-0.00 (-0.05 to 0.05)	0.92	-273%
Change in total fat intake as %E	9	-0.17	-0.01 (-0.03 to 0.01)	0.28	100%
Gender*	13	-0.17	-0.14 (-0.63 to 0.35)	0.55	-13%
Study duration	13	-0.47	0.00 (-0.01 to 0.02)	0.76	-24.8%
CVD risk at baseline**	13	-0.44	0.03 (-0.48 to 0.55)	0.89	-39%

Table 4. Meta-regression of effects of SFA reduction on cardiovascular events

*Gender was coded as follows: 0 = women, 1 = mixed, 2 = men

**CVD risk at baseline was coded as follows: 1 = Low CVD risk, 2 = Moderate CVD risk, 3 = existing CVD

CHO: carbohydrate

CI: confidence interval

CVD: cardiovascular disease

E: energy

MUFA: monounsaturated fatty acid

PUFA: polyunsaturated fatty fat

SFA: saturated fatty acid

TC: total cholesterol

Cut- off	RR of all- cause mor- tality	RR of CVD mortality	RR of CVD events	RR of MI	RR of non-fatal MI	RR of stroke	RR of CHD mortality	RR of CHD events
7%E	0.89 (0.66 to 1.20)	0.20 (0.01 to 4.15)	0.20 (0.01 to 4.15)	N/A	N/A	N/A	N/A	N/A
8%E	0.89 (0.66 to 1.20)	0.20 (0.01 to 4.15)	0.20 (0.01 to 4.15)	N/A	N/A	N/A	N/A	N/A
9%E	0.96 (0.83 to 1.10)	0.69 (0.51 to 0.94)	0.79 (0.62 to 0.99)	0.76 (0.55 to 1.05)	0.62 (0.31 to 1.21)	0.59 (0.30 to 1.15)	0.82 (0.55 to 1.21)	0.77 (0.56 to 1.04)
10%E	0.99 (0.90 to 1.09)	0.95 (0.67 to 1.35)	0.78 (0.57 to 1.05)	0.93 (0.80 to 1.08)	0.89 (0.58 to 1.35)	0.87 (0.58 to 1.33)	1.05 (0.77 to 1.43)	0.82 (0.60 to 1.13)
11%E	0.99 (0.88 to 1.12)	0.92 (0.65 to 1.31)	0.77 (0.59 to 1.02)	0.94 (0.84 to 1.06)	0.89 (0.58 to 1.35)	0.76 (0.45 to 1.30)	1.02 (0.84 to 1.24)	0.85 (0.63 to 1.15)
12%E	0.98 (0.91 to 1.07)	0.95 (0.75 to 1.21)	0.85 (0.71 to 1.02)	0.94 (0.85 to 1.04)	0.90 (0.72 to 1.14)	0.93 (0.55 to 1.25)	1.02 (0.84 to 1.24)	0.90 (0.77 to 1.06)
13%E	1.02 (0.83 to 1.25)	0.93 (0.63 to 1.38)	0.78 (0.61 to 1.00)	0.87 (0.73 to 1.04)	0.72 (0.50 to 1.03)	0.54 (0.29 to 1.00)	1.06 (0.76 to 1.48)	0.84 (0.63 to 1.12)

CHD: coronary heart disease

Table 5. SFA cut-off data

CVD: cardiovascular disease

E: energy

MI: myocardial infarction

N/A: not applicable (no relevant studies)

RR: risk ratio

SFA: saturated fat, as percentage of energy

Cochrane Library


APPENDICES

Appendix 1. Search strategies 2019 CENTRAL #1 lipid near (low* or reduc* or modifi*) #2 cholesterol* near (low* or modifi* or reduc*) #3 (#1 or #2) #4 MeSH descriptor: [Nutrition Therapy] explode all trees #5 diet* or food* or nutrition* #6 (#4 or #5) #7 (#3 and #6) #8 fat* near (low* or reduc* or modifi* or animal* or saturat* or unsaturat*) #9 MeSH descriptor: [Diet, Atherogenic] explode all trees #10 MeSH descriptor: [Diet Therapy] explode all trees #11 (#7 or #8 or #9 or #10) #12 MeSH descriptor: [Cardiovascular Diseases] this term only #13 MeSH descriptor: [Heart Diseases] explode all trees #14 MeSH descriptor: [Vascular Diseases] explode all trees #15 MeSH descriptor: [Cerebrovascular Disorders] this term only #16 MeSH descriptor: [Brain Ischemia] explode all trees #17 MeSH descriptor: [Carotid Artery Diseases] explode all trees #18 MeSH descriptor: [Dementia, Vascular] explode all trees #19 MeSH descriptor: [Intracranial Arterial Diseases] explode all trees #20 MeSH descriptor: [Intracranial Embolism and Thrombosis] explode all trees #21 MeSH descriptor: [Intracranial Hemorrhages] explode all trees #22 MeSH descriptor: [Stroke] explode all trees #23 coronar* near (bypas* or graft* or disease* or event*) #24 cerebrovasc* or cardiovasc* or mortal* or angina* or stroke or strokes or tia or ischaem* or ischem* #25 myocardi* near (infarct* or revascular* or ischaem* or ischem*) #26 morbid* near (heart* or coronar* or ischaem* or ischem* or myocard*) #27 vascular* near (peripheral* or disease* or complication*) #28 heart* near (disease* or attack* or bypas*) #29 (#12 or #13 or #14 or #15 or #16 or #17 or #18 or #19 or #20 or #21 or #22 or #23 or #24 or #25 or #26 or #27 or #28) #30 (#11 and #29) Date added to CENTRAL trials database: 05/03/2014-15/10/2019

MEDLINE OVID

Reduction in saturated fat intake for cardiovascular disease (Review) Copyright © 2020 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.



- 1. (lipid\$ adj5 (low\$ or reduc\$ or modifi\$)).mp.
- 2. (cholesterol\$ adj5 (low\$ or modific\$ or reduc\$)).mp.

3.1 or 2

- 4. exp Nutrition Therapy/
- 5. (diet\$ or food\$ or nutrition\$).mp.
- 6.4 or 5

7.3 and 6

8. (fat adj5 (low\$ or reduc\$ or modifi\$ or animal\$ or saturat\$ or unsatur\$)).mp.

9. exp Diet, Atherogenic/

10. exp Diet Therapy/

11. 7 or 8 or 9 or 10

12. cardiovascular diseases/ or exp heart diseases/ or exp vascular diseases/

13. cerebrovascular disorders/ or exp brain ischemia/ or exp carotid artery diseases/ or exp dementia, vascular/ or exp intracranial arterial diseases/ or exp "intracranial embolism and thrombosis"/ or exp intracranial hemorrhages/ or exp stroke/

- 14. (coronar\$ adj5 (bypas\$ or graft\$ or disease\$ or event\$)).mp
- 15. (cerebrovasc\$ or cardiovasc\$ or mortal\$ or angina\$ or stroke or strokes).mp.
- 16. (myocardi\$ adj5 (infarct\$ or revascular\$ or ischaemi\$ or ischemi\$)).mp.
- 17. (morbid\$ adj5 (heart\$ or coronar\$ or ischaem\$ or ischem\$ or myocard\$)).mp.
- 18. (vascular\$ adj5 (peripheral\$ or disease\$ or complication\$)).mp.
- 19. (heart\$ adj5 (disease\$ or attack\$ or bypass\$)).mp.

20. 12 or 13 or 14 or 15 or 16 or 17 or 18 or 19

- 21. 11 and 20
- 22. randomized controlled trial.pt.
- 23. controlled clinical trial.pt.
- 24. randomized.ab.
- 25. placebo.ab.
- 26. drug therapy.fs.
- 27. randomly.ab.
- 28. trial.ab.
- 29. groups.ab.
- 30. 22 or 23 or 24 or 25 or 26 or 27 or 28 or 29
- 31. exp animals/ not humans.sh.
- 32. 30 not 31
- 33. 21 and 32
- 34. limit 33 to ed=20140305-20191015



Embase OVID

1. cardiovascular diseases/ or exp heart diseases/ or exp vascular diseases/

2. cerebrovascular disorders/ or exp brain ischemia/ or exp carotid artery diseases/ or exp dementia, vascular/ or exp intracranial arterial diseases/ or exp "intracranial embolism and thrombosis"/ or exp intracranial hemorrhages/ or exp stroke/

- 3. (coronar\$ adj5 (bypas\$ or graft\$ or disease\$ or event\$)).mp.
- 4. (cerebrovasc\$ or cardiovasc\$ or mortal\$ or angina\$ or stroke or strokes).mp.
- 5. (myocardi\$ adj5 (infarct\$ or revascular\$ or ischaemi\$ or ischemi\$)).mp.
- 6. (morbid\$ adj5 (heart\$ or coronar\$ or ischaem\$ or ischem\$ or myocard\$)).mp.
- 7. (vascular\$ adj5 (peripheral\$ or disease\$ or complication\$)).mp.
- 8. (heart\$ adj5 (disease\$ or attack\$ or bypass\$)).mp.
- 9. or/1-8
- 10. (lipid\$ adj5 (low\$ or reduc\$ or modifi\$)).mp.
- 11. (cholesterol\$ adj5 (low\$ or modific\$ or reduc\$)).mp.
- 12. 10 or 11
- 13. (diet\$ or food\$ or eat\$ or nutrition\$).mp.
- 14. exp nutrition/
- 15. 13 or 14
- 16. 12 and 15
- 17. (fat adj5 (low\$ or reduc\$ or modifi\$ or animal\$ or saturat\$ or unsatur\$)).mp.
- 18. exp lipid diet/ or exp fat intake/ or exp low fat diet/
- 19. 16 or 17 or 18
- 20. 9 and 19
- 21. random\$.tw.
- 22. factorial\$.tw.
- 23. crossover\$.tw.
- 24. cross over\$.tw.
- 25. cross-over\$.tw.
- 26. placebo\$.tw.
- 27. (doubl\$ adj blind\$).tw.
- 28. (singl\$ adj blind\$).tw.
- 29. assign\$.tw.
- 30. allocat\$.tw.
- 31. volunteer\$.tw.
- 32. crossover procedure/
- 33. double blind procedure/



Trusted evidence. Informed decisions. Better health.

34. randomized controlled trial/

- 35. single blind procedure/
- 36. 21 or 22 or 23 or 24 or 25 or 26 or 27 or 28 or 29 or 30 or 31 or 32 or 33 or 34 or 35
- 37. (animal/ or nonhuman/) not human/
- 38. 36 not 37
- 39. 20 and 38
- 40. limit 39 to dd=20140305-20191015

Clinicaltrials.gov

Condition or disease: Cardiovascular Diseases OR CVD OR "heart disease"

Intervention/treatment: Dietary Fats OR saturated OR unsaturated OR fat

Study type: Interventional Studies (Clinical Trials)

ICTRP

Condition: Cardiovascular Diseases OR CVD OR heart disease

Intervention: Dietary Fats OR saturated OR unsaturated OR fat

WHAT'S NEW

Date	Event	Description
9 January 2020	New citation required but conclusions have not changed	No new trials included, but four ongoing trials and one study awaiting assessment, and we found new data for two of the al- ready included trials (WHI 2006; WINS 2006). We updated assess- ment of risk of bias, including assessment of summary risk of bias for each trial, and carrying out sensitivity analyses omitting trials not at low summary risk of bias. We updated assessment of small study bias by comparing results of fixed- and random-ef- fects meta-analyses. Data, results, GRADE assessment and con- clusions updated.
29 December 2019	New search has been performed	Searches updated to October 2019, searches of trials registers added.

HISTORY

Protocol first published: Issue 2, 1999 Review first published: Issue 6, 2015

Date	Event	Description
27 March 2015	New citation required and conclusions have changed	We split a previously published review (Reduced or modi- fied dietary fat for preventing cardiovascular disease, DOI: 10.1002/14651858.CD002137.pub3) into six smaller review up- dates. The conclusions are therefore now focused on reduction in saturated fat intake instead of reducing or modifying fat intake overall on its effect on cardiovascular disease risk.

Reduction in saturated fat intake for cardiovascular disease (Review) Copyright © 2020 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.



Trusted evidence. Informed decisions. Better health.

Date	Event	Description
		This split review update includes 15 randomised controlled tri- als.
5 March 2014	New search has been performed	The search has been updated to 5 March 2014.

CONTRIBUTIONS OF AUTHORS

All authors were active in the design of the review and in providing critical revisions of the manuscript, all authors took part in assessment of the results of the updated search, and assessment of inclusion of potentially relevant studies. All authors edited, proof-read and agreed the final version of the review.

LH was the principal author of earlier versions (Hooper 2000; Hooper 2001; Hooper 2012; Hooper 2015a), originated and was primarily responsible for planning and carrying out this systematic review, liaising with WHO NUGAG, carrying out the statistical analyses, and writing the first draft of this review.

AA, OFJ, CK, EF and LH were responsible for data extraction and assessment of validity.

DECLARATIONS OF INTEREST

Lee Hooper: LH is a member of the World Health Organization Nutrition Guidance Expert Advisory Group (NUGAG). WHO paid for her travel, accommodation and expenses to attend NUGAG meetings in Geneva, China and South Korea where the evidence of effects of dietary fats on health was discussed and guidance developed. LH's institution was given grant funding from WHO to carry out the 2019 update of this systematic review, to update a systematic review on the relationship between total fat intake and body weight and a series of systematic reviews on the health effects of polyunsaturated fatty acids.

Nicole Martin: None known

Asmaa Abdelhamid: None known

Oluseyi Florence Jimoh: This review was funded by a grant from the World Health Organization.

Eve Foster: None known

Christian Kirk: None known

SOURCES OF SUPPORT

Internal sources

- University of East Anglia, UK
- Help with acquiring papers for previous versions of this review, and allowing time for Lee Hooper to work on the review
- University of Manchester, UK

Support with collection of papers for the first version of this review

External sources

- Studentship, Systematic Reviews Training Unit, Institute of Child Health, University of London, UK
- Funding to support Lee Hooper to carry out the first version of the systematic review
- World Health Organization, Other

WHO funded the most recent update of this review

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

This review is the result of updating the searches for Hooper 2015a. The objective and outcomes have been widened since the protocol to address queries by WHO NUGAG and the inclusion criteria have changed to focus on saturated fat and long-term trials (24 months instead of six months).



INDEX TERMS

Medical Subject Headings (MeSH)

Cardiovascular Diseases [mortality] [*prevention & control]; Cause of Death; Dietary Carbohydrates [administration & dosage]; Dietary Fats [*administration & dosage]; Dietary Fats, Unsaturated [administration & dosage]; Dietary Proteins [administration & dosage]; Energy Intake; Fatty Acids [*administration & dosage]; Myocardial Infarction [mortality] [prevention & control]; Randomized Controlled Trials as Topic; Stroke [prevention & control]

MeSH check words

Adult; Humans