

Published in final edited form as:

J Org Chem. 2011 October 7; 76(19): 7842–7848. doi:10.1021/jo201192c.

Reduction of Amine *N*-Oxides by Diboron Reagents

Hari Prasad Kokatla, Paul F. Thomson, Suyeal Bae, Venkata Ramana Doddi, and Mahesh K. Lakshman*

Department of Chemistry, The City College and The City University of New York, 160 Convent Avenue, New York, NY 10031-9198

Abstract



Facile reduction of alkylamino-, anilino-, and pyridyl-*N*-oxides can be achieved via the use of diboron reagents, predominantly bis(pinacolato)- and in some cases bis(catecholato)diboron [(pinB)₂ and (catB)₂, respectively]. Reductions occur upon simply mixing the amine *N*-oxide and the diboron reagent in a suitable solvent, at a suitable temperature. Extremely fast reductions of alkylamino- and anilino-*N*-oxides occur, whereas pyridyl-*N*-oxides undergo slower reduction. The reaction is tolerant of a variety of functionalities such as hydroxyl, thiol, and cyano groups, as well as halogens. Notably, a sensitive nucleoside *N*-oxide has also been reduced efficiently. The different rates with which alkylamino- and pyridyl-*N*-oxides are reduced has been used to perform stepwise reduction of the *N,N'*-dioxide of (*S*)-(-)-nicotine. Because it was observed that (pinB)₂ was unaffected by the water of hydration in amine oxides, the feasibility of using water as solvent was evaluated. These reactions also proceeded exceptionally well, giving high product yields. In contrast to the reactions with (pinB)₂, triethylborane reduced alkylamino-*N*-oxides, but pyridine *N*-oxide did not undergo efficient reduction even at elevated temperature. Finally, the mechanism of the reductive process by (pinB)₂ has been probed by ¹H and ¹¹B NMR.

INTRODUCTION

Amine *N*-oxides are frequently encountered in organic synthesis,¹ and often the chemical methodology calls for a reduction of the *N*-oxide to the amine. A variety of methods involving metals have been developed for the *N*-oxide to amine conversion.^{2–21} There are fewer methods that do not rely on the use of metals, such as processes involving sulfurous acid,²² SO₂,²³ sulfur monoxide,²⁴ trimethyl(ethyl)amine–SO₂ complex,²⁵ PCl₃,²⁶ PPh₃ at high temperatures,²⁷ di-*n*-propyl sulfoxylate,²⁸ CS₂,²⁹ bakers' yeast,³⁰ and alcohols and a base.³¹

Many of these methods for *N*-oxide to amine conversion are encumbered with functional group incompatibility problems, need for high temperatures and/or sealed tube techniques, reagents that produce undesired side reactions, and difficult access to reagents and/or catalysts. Thus, a mild and simple method for the selective conversion of amine *N*-oxides to the corresponding amines continues to be synthetically desirable.

*Tel (212) 650-7835; fax (212) 650-6107; lakshman@sci.cuny.cuny.edu.

SUPPORTING INFORMATION

¹H NMR spectra of all *N*-oxides and their reduction products, as well as the ¹H–¹H COSY spectrum for **1**. This information is available free of charge via the Internet at <http://pubs.acs.org>.

In the course of recent work, we had observed an unusual deoxygenation of *O*⁶-(benzotriazolyl)inosine and 2'-deoxyinosine derivatives (a class of purine-benzotriazole ethers) upon exposure to bis(pinacolato)diboron [(pinB)₂] and Cs₂CO₃ (Scheme 1), and had proposed a plausible mechanism for the conversion.³² The products from the deoxygenation reactions are C-6 benzotriazolyl purine nucleoside derivatives, which were isolated in good yields.³² In this reaction, it appears that the oxygen atom attached to the benzotriazolyl moiety is transferred to (pinB)₂, which undergoes oxidation to (pinB)₂O.

(PinB)₂ and the corresponding catechol derivative [(catB)₂] have been briefly investigated for their abilities to reduce *N*-oxides of pyridine, 4-phenylpyridine, and 4-methylmorpholine. Although pyridine *N*-oxide was stated to be reduced in 90% yield by (catB)₂, no preparative use of these reagents has been reported to date.³³ Nevertheless, deoxygenation by diboron reagents is anticipated to be an exceptionally mild and selective process, with broad functional group compatibility. Both (catB)₂ and (pinB)₂ are commercially available. The latter is a comparatively less-expensive, stable compound, which requires no special handling protocols. These considerations led us to explore the use of diboron reagents for the reduction of amine *N*-oxides.

RESULTS AND DISCUSSION

The generally oxophilic nature of boron and the relatively stable B–O bond (~120–130 kcal/mol^{34,35}) in comparison to the B–B bond (~68 kcal/mol³³), and the large favorable enthalpy associated with the formation of two B–O bonds (ΔH ca. 180 kcal/mol³³) all bode well for the use of diboron reagents as reducing agents. Furthermore, trimethylamine *N*-oxide has been known for some time to be an effective oxidant for converting organoboranes to alcohols.^{36–39} With these data, we conducted an NMR tube experiment, where 4-methylmorpholine *N*-oxide was exposed to a slight stoichiometric excess of (pinB)₂, in CDCl₃ at room temperature. No additional precautions were taken, such as the use of inert atmosphere or anhydrous conditions. On the basis of ¹H NMR evaluation, this reaction was extremely rapid (Figure 1) and *complete reduction took no more than 10 minutes at room temperature*. In C₆D₆ this reaction is reported to be complete within 1 h at 25 °C, leading to a 74% isolated yield of a morpholine•(pinB)₂O adduct.³³ Contact of 4-methylmorpholine *N*-oxide with (pinB)₂ at room temperature, in the absence of solvent, led to an immediate, violent reaction with evolution of fumes.

We then queried whether the hybridization of the nitrogen atom made any difference on the course of the reaction. We chose to conduct the reduction with 4-methoxypyridine *N*-oxide hydrate since this compound was expected to be reasonably reactive based on the anticipated reaction mechanism (vide infra). Again, an experiment was conducted in an NMR tube using CD₃CN as solvent (based upon solubility of reagents and the fact that the reaction mixture could be heated, if necessary), and upon mixing 4-methoxypyridine *N*-oxide hydrate and (pinB)₂, a reaction was clearly evident. However, this was much slower than that of 4-methylmorpholine *N*-oxide, and reached completion within 20 h at room temperature. In comparison to 4-methylmorpholine *N*-oxide, a similar slower reduction of pyridine *N*-oxide by (pinB)₂ has been noted.³³ The exotherm associated with the deoxygenation of 4-methylmorpholine-*N*-oxide and 4-methoxypyridine *N*-oxide was determined to be about 0.6 kcal/g by calorimetry.

With the information gleaned from NMR studies, we turned our attention to the broader synthetic generality of the reductive process. We were aware that alkylamine *N*-oxides underwent rapid reactions, whereas pyridine *N*-oxides were slower to react. One other issue remained to be addressed: the formation of amine complexes with the boron-based byproduct, (pinB)₂O,³³ which we also observed. The formation of amine•(pinB)₂O adducts

has been attributed to the increased Lewis acidity at the boron center after oxidation of the B–B bond.³³ We reasoned that addition of a strongly coordinating alkyldiamine should liberate the desired amine from the complex, and we chose water-soluble ethylenediamine for the workup protocol. The results of the reduction procedure are presented in Table 1.

Some notable points emerge from the results in Table 1. *N*-Oxides or their hydrates can be reduced and water does not seem to hamper this reaction by degradation of (pinB)₂ (entries 3 and 5). Reactions of some amine *N*-oxides were slow at room temperature, and heating accelerated the reduction. There does not appear to be an appreciable effect of solvent, and solubility of reactants as well as reaction temperature may be the only considerations for solvent selection (entries 4 and 11). Thiol and hydroxyl groups did not need any specific caution or protection (entries 8 and 9). Halogens are well tolerated, and steric crowding by two chlorine atoms does not deter the reduction (entries 10 and 11). Electron-deficient *p*-cyanopyridine *N*-oxide also underwent clean reduction (entry 12). The substituents in examples 10–12 could be incompatible with some other reduction methods. Oxides of *N,N*-diethylaniline, *N*-benzylpiperidine, and *N*-benzylmorpholine were all reduced rapidly (entries 13–15).

In the cases of pyridine, 2-picoline, 3-picoline, and 2,6-lutidine *N*-oxides, although reductions with (pinB)₂ proceeded smoothly, product volatility made their separation from pinacol-based byproducts somewhat more cumbersome. Use of bis(catecholato)diboron [(catB)₂] ameliorated the purification problems (entries 16–19). In our hands, reactions with (catB)₂ appeared slower as compared to (pinB)₂, and (catB)₂ seems to be moisture-sensitive. As an example, in CH₃CN, reduction of pyridine *N*-oxide with stoichiometric (pinB)₂ was complete within 10 h at 70 °C. On the other hand, even with 1.5 mol equiv of (catB)₂, a comparable reaction was incomplete at 24 h and a temperature of 120 °C was necessary. Nevertheless, reductions can be conducted with (catB)₂, and the steric bulk in 2,6-lutidine *N*-oxide was not an impediment. In prior work, a difference has been noted in the reactivities of (pinB)₂ and (catB)₂, but reduction of pyridine *N*-oxide with (catB)₂ was complete within 3 h at 25 °C, whereas reaction with (pinB)₂ required 6 h at 70 °C.³³

Next we wanted to test the deoxygenation of more complex substrates. Given our interest in nucleoside functionalization, we synthesized 2',3',5'-tri-*O*-(*tert*-butyldimethylsilyl)adenosine *N*1-oxide (**1** in Scheme 2) using a known procedure.⁴⁰ This compound is significantly more fragile as compared to the substrates in Table 1. Exposure of a 0.08 M solution of trisilyl nucleoside **1** to (pinB)₂ in either CH₃CN at 70 °C, or in diglyme at 120 °C led to clean reduction, and trisilyl adenosine **2** was isolated in 71% and 87% yield, respectively (Scheme 2).

(*S*)-(-)-Nicotine presents an interesting case study because it contains alkylamino and pyridyl nitrogen atoms. Both the *N*-oxide **4** and the *N,N'*-dioxide **5** were synthesized using known procedures.^{41,42} These compounds gave us excellent models to test selective as well as comprehensive *N*-oxide reduction (Scheme 3). Reaction of nicotine *N*-oxide **4** with 1 mol equiv of (pinB)₂ gave nicotine (**3**) in 95% yield within 10 minutes, at room temperature. On the other hand, reduction of the *N,N'*-dioxide **5** with 1 mol equiv of (pinB)₂ led exclusively to *N*-oxide **6**^{30,42} in 96% yield, within 10 min at room temperature. By contrast, reduction of **5** to **6** has previously been conducted using SO₂/EtOH, in an overnight reaction.⁴³ The present method provides straightforward, scalable access to the expensive *N*-oxide **6** (commercially *ca.* \$1–3/mg) from readily prepared *N,N'*-dioxide **5**.

Finally, reaction of *N,N'*-dioxide **5** with 2 mol equiv of (pinB)₂ showed formation of *N*-oxide **6** within 10 min at room temperature, and complete reduction to (*S*)-(-)-nicotine (**3**) was then achieved within 1 h, by heating the reaction mixture to 70 °C (93% yield). In

contrast to the workup procedure with the simpler amines, which required addition of ethylenediamine to completely remove boron-containing byproducts, reactions of **4** and **5** did not require this step. The chiral center adjacent to the pyrrolidine nitrogen atom remains unaffected in the reduction, as seen from the optical rotation of the total reduction product **3** in comparison to the (*S*)-(-)-nicotine sample used for this work.

Use of Water as Reaction Medium

Because the water of hydration in the amine *N*-oxides did not seem to affect the reductions by (pinB)₂ (entries 3 and 5 in Table 1; also see the NMR studies described below), we queried whether water could be used as a reaction medium. Results from the reactions of five amine *N*-oxides in water are shown in Table 2. Not only did these reactions prove comparable to those accomplished in organic solvents, but also product isolation was just as readily achieved.

Comparison of Reductions by Triethyl Borane and (pinB)₂

Although pinB–H has been used for preparing (pinB)₂O by reduction of trimethylamine-*N*-oxide,⁴⁴ its utility as a reducing reagent is limited by functional group compatibility issues, and by moisture sensitivity considerations. Further, pinB–OH initially formed in the reaction of pinB–H and the amine *N*-oxide, consumes an equivalent of pinB–H, releasing hydrogen.⁴⁴ Formation of hydrogen by reaction of B–H containing organoboranes and amine *N*-oxides is known.^{36,37} Thus, minimally 2 mol equiv of pinB–H will be required for reducing amine oxides. In principle, trialkylboranes can be used to reduce amine *N*-oxides, as has been demonstrated with trimethylamine *N*-oxide.^{36–39} Although 3 mol of a *N*-oxide can be reduced per 1 mol of a trialkylborane, it has been shown that the rate for each sequential B–C oxidation step is slower than the previous one.³⁷ For example, reaction of 2 mol equiv of trimethylamine *N*-oxide with 1 mol equiv of *n*-Bu₃B is rapid.⁴⁵ However, reaction with the third molar equivalent of the *N*-oxide required 24 h at reflux.⁴⁵ On the basis of these results, we conducted several experiments. First, we assessed selectivity of the reduction by exposing a solution of nicotine *N,N'*-dioxide **5** in CH₃CN to 0.3 mol equiv of commercially available Et₃B, at room temperature. This reaction yielded exclusively *N*-oxide **6** in 53% yield. Because only two alkylamine *N*-oxides can be expected to undergo reduction per Et₃B, at room temperature,⁴⁵ in theory the maximal yield should be 60%. Second, we conducted a reaction of *N*-benzylpiperidine *N*-oxide with 1 mol equiv of Et₃B, also in CH₃CN. This reaction was complete within 10 min affording *N*-benzylpiperidine in 83% yield. Finally, with 1 mol equiv of Et₃B, practically no reduction of pyridine *N*-oxide was observed at room temperature in CH₃CN, and just a trace of pyridine was observed by TLC, after heating at 80 °C for 20 h. It has been reported that in boiling xylene, only a single B–C bond of an alkylborane is oxidized by 4-methylpyridine *N*-oxide.³⁷ Thus, reactions of alkylboranes with *N*-oxides of alkylamines are facile in comparison to reactions with pyridine-*N*-oxides. On the basis of these collective data, (pinB)₂ appears to be a reagent endowed with broad *N*-oxide reducing ability as well as selectivity.

Evaluation of the Reaction Mechanism

To gain insight into the mechanism of this reaction, we conducted some ¹H and ¹¹B NMR experiments. A 0.28 M solution of 4-methoxypyridine *N*-oxide in CD₃CN was exposed to 1.05 mol equiv of (pinB)₂ in an NMR tube. Figure 2, panel A shows that within 10 min at room temperature, a fast reaction ensued. Continued monitoring of this reaction showed that substantial reduction was complete within 125 min, and finally no trace of the *N*-oxide was detectable at 20 h. The disappearance of the OCH₃ resonance in 4-methoxypyridine *N*-oxide (δ = 3.90 ppm) and formation of the corresponding resonance in 4-methoxypyridine (δ = 3.89 ppm) was also observed (not shown in Figure 2A, panel A). The resonances of

(pinB)₂ and its oxidation product appear at $\delta = 1.25$ – 1.26 ppm but are not particularly informative.

¹¹B NMR (Figure 2, panel B) showed disappearance of the resonance at $\delta = 30.6$ ppm corresponding to (pinB)₂ and formation of a single new entity with a resonance at $\delta = 22.5$ ppm. From the ¹¹B NMR chemical shift it is plausible that the oxidation product formed is (pinB)₂O because the chemical shift is consistent for a boron bonded to an oxygen atom.⁴⁶ For comparison, authentic (pinB)₂O was synthesized as reported,⁴⁴ and its ¹¹B resonance was recorded in several solvents.⁴⁷ The chemical shift of the boron-containing product from the deoxygenation reaction is consistent with (pinB)₂O. Because it is not of particular relevance to the chemical methodology, we have not determined whether this product is or contains hydrolyzed pinB–OH. The literature indicates very similar chemical shifts for (pinB)₂O and pinB–OH,^{44,48,49} and amine can form a (pinB)₂O adduct.³³

To investigate by NMR whether water in the *N*-oxide hydrate had any influence on the reaction course, the reduction of 4-phenylpyridine *N*-oxide (which was not a hydrate) was also monitored by ¹H and ¹¹B NMR. The results from the ¹¹B NMR experiment in this case were comparable to those from the reduction of 4-methoxypyridine *N*-oxide hydrate.

On the basis of the ¹¹B NMR experiments, where only two boron resonances ascribable to (pinB)₂ and a final product were discernible, a plausible mechanism is shown in Scheme 4 that is consistent with the one proposed previously.³³ An N–O–B bond is formed by reaction of the *N*-oxide at a boron center, and in a subsequent step (pinB)₂O is produced with liberation of the amine that can remain bound to the boron. The ¹H NMR results, where only two sets of resonances were observed during the course of the reduction, also support this proposal. Consistent with the mechanism proposed, the highly electron-depleted *p*-nitropyridine *N*-oxide was a recalcitrant substrate, whereas the electron-rich *p*-methoxypyridine *N*-oxide underwent reaction at room temperature.

CONCLUSIONS

We have described a facile reduction method for alkylamino-, anilino-, and pyridyl-*N*-oxides. The reactions rely on a simple exposure of the *N*-oxide to (pinB)₂, in a suitable solvent, and at a suitable temperature to attain a reasonable reaction rate. The reductions are generally mild, require no specialized handling, are compatible with a wide range of functionalities, and can be conducted with stoichiometric (pinB)₂. Product isolation is also straightforward. In addition to the broad range of amine oxides that can be reduced, (pinB)₂ offers selectivity in reduction. Thus, an alkylamino-*N*-oxide can be reduced in the presence of a pyridyl-*N*-oxide, as demonstrated with the *N,N'*-dioxide of (*S*)-(–)-nicotine. Water of hydration does not impact the reductions, and notably reductions of amine *N*-oxides can be conducted in water as reaction medium. In recent years, heterocyclic *N*-oxides have proven to be excellent substrates for metal-catalyzed C–H bond activation processes, wherein the *N*-oxide moiety is finally reduced using Pd/HCO₂NH₄, Zn/NH₄Cl, or PCl₃.^{50,51} In this context, the present approach provides an excellent, mild, and functional group-compatible method for the deoxygenation. Also, deoxygenative functionalization of pyridines has been recently reported.⁵² Thus, various natural products containing both pyridyl and alkylamino groups can be functionalized at the pyridine ring via the selectivity offered by this route. All these factors render this method attractive for wide ranging applications in organic synthesis, and our efforts are focused on tapping its utility for more complex functionalizations.

EXPERIMENTAL SECTION

General Experimental Considerations

Thin layer chromatography was performed on 250 μm silica plates, and column chromatographic purifications were performed on 200–300 mesh silica gel. MeCN was distilled over CaH_2 , 1,4-dioxane was distilled over LiAlH_4 and then over Na. All other reagents were obtained from commercial sources. *N*-Oxides in entries 1–6, 8, 9, 11, 12, 16–19 were obtained from commercial suppliers and were used without further purification, whereas *N*-oxides in entries 7, 10, 13, 14, 15 were synthesized. Trisilyl adenosine *N*1-oxide **1** was prepared via a known route (reference 40). Nicotine *N*-oxide **4** and the *N,N'*-dioxide **5** were synthesized by reported routes (reference 41). ^1H NMR spectra were recorded at 500 MHz in CDCl_3 or $\text{DMSO-}d_6$ and are referenced to the residual protonated solvent resonance. All products are known compounds and, except for *N*-benzylmorpholine, are commercially available. Characterization data of *N*-benzylpiperidine and *N*-benzylmorpholine are reported in the literature.^{53,54}

General Procedure for Synthesis of Amine *N*-Oxides

To a stirred solution of amine (5.5–6.7 mmol) in CHCl_3 (2 mL) was added 70% *m*-CPBA (1 mol equiv), portionwise at 0 °C. The resulting mixture was stirred at room temperature for 12 h, at which time complete consumption of starting material was observed by TLC. The reaction mixture was diluted with CHCl_3 , and solid K_2CO_3 (4 mol equiv) was added. The resulting mixture was stirred for an additional 10 min. The solid was separated by filtration, and the filtrate was dried over Na_2SO_4 and concentrated under reduced pressure to afford the amine *N*-oxide in yields of 86–90%.

General Procedure for the Reduction of Amine *N*-Oxides (reactions were conducted on the 0.5–1.0 mmol scale). Caution: these reductions, particularly those of alkyl amine *N*-oxides can be quite vigorous and exothermic. Hence appropriate precautions should be taken while conducting them.

For high boiling amines—A 1 M solution of amine *N*-oxide in the appropriate solvent (see Table 1) was stirred in an oven-dried reaction vial. $(\text{PinB})_2$ (1 mol equiv) was added, the vial was flushed with nitrogen gas, and the mixture was stirred at the appropriate temperature (see Table 1 for details). When TLC indicated the reaction to be complete, ethylenediamine (20 mol equiv) was added to the mixture, and the stirring was continued for 1 h at room temperature. Then the reaction mixture was diluted with water (10 mL) and extracted with Et_2O (3×10 mL). The organic layer was dried over Na_2SO_4 and concentrated under reduced pressure. The crude material was then purified via a silica gel column using EtOAc /hexanes as eluting solvent.

For low boiling amines—A 1 M solution of amine *N*-oxide in MeCN was stirred in an oven-dried reaction vial. $(\text{CatB})_2$ (1.5 mol equiv) was added, the vial was flushed with nitrogen gas, and the mixture was stirred at 120 °C. When TLC indicated the reaction to be complete, ethylenediamine (20 mol equiv) was added to the mixture, and the stirring was continued for 1 h at room temperature. Then the reaction mixture was diluted with water (10 mL) and extracted with Et_2O (3×10 mL). The organic layer was extracted with 1 N HCl. The aqueous layer was separated, and carefully neutralized with 1 N NaOH. The free amine was extracted into Et_2O . The organic layer was dried over Na_2SO_4 and concentrated under reduced pressure. The crude material was then purified via a silica gel column using EtOAc /hexanes as eluting solvent.

General Procedure for the Reduction of Amine *N*-Oxides in Water

To a 1 M solution of amine *N*-oxide in water was added (pinB)₂ (1 mol equiv) with stirring. The mixture was stirred at the appropriate temperature (see Table 2 for details). When TLC indicated the reaction to be complete, ethylenediamine (20 mol equiv) was added to the mixture, and the stirring was continued for 1 h at room temperature. Then the reaction mixture was diluted with water (10 mL) and extracted with Et₂O (3 × 10 mL). The organic layer was dried over Na₂SO₄ and concentrated under reduced pressure. The crude material was then purified via a silica gel column using EtOAc/hexanes as eluting solvent.

2',3',5'-Tri-*O*-(*tert*-butyldimethylsilyl)adenosine *N*1-oxide (**1**)⁴⁰

In an oven-dried 10 mL round-bottom flask, equipped with a stirring bar, was placed a solution of 2',3',5'-tri-*O*-(*t*-butyldimethylsilyl)adenosine (**2**, 200 mg, 0.32 mmol) in MeOH (5 mL). To this well-stirred mixture was added 70% *m*-CPBA (79 mg, 1.4 mol equiv), the vial was flushed with nitrogen gas, and the mixture was stirred at room temperature. After 12 h, TLC indicated the reaction to be complete. The mixture was diluted with CHCl₃ (30 mL) and washed with aqueous NaHCO₃ (3 × 20 mL). The organic layer was dried over Na₂SO₄, filtered, and evaporated under reduced pressure. The crude material was purified via a silica gel column using 10% MeOH in EtOAc to give **1** as a brown solid (179 mg, 87% yield). ¹H NMR (500 MHz, CDCl₃): δ 8.68 (s, 1H, Ar-H), 8.30 (s, 1H, Ar-H), 5.98 (d, 1H, H-1', *J* = 5.4 Hz), 4.57 (t, 1H, H-2', *J* = 4.4 Hz), 4.28 (t, 1H, H-3', *J* = 3.9 Hz), 4.27 (q, 1H, H-4', *J* = 3.1 Hz), 4.00 (dd, 1H, H-5, *J* = 3.9, 11.5 Hz), 3.78 (dd, 1H, H-5', *J* = 2.4, 11.5 Hz), 0.96, 0.92, and 0.80 (3s, 27H, *t*-Bu), 0.15, 0.13, 0.10, 0.09, -0.03, and -0.23 (6s, 18H, SiCH₃). HRMS (ESI) calcd for C₂₈H₅₆N₅O₅Si₃ [M + H]⁺ 626.3584, found 626.3608.

Reduction of 2',3',5'-tri-*O*-(*t*-butyldimethylsilyl)adenosine *N*1-oxide (**1**)

In an oven-dried reaction vial equipped with a stirring bar was placed the adenosine *N*-oxide **1** (50.0 mg, 0.08 mmol) in diglyme (1 mL). (PinB)₂ (20.0 mg, 0.08 mmol) was added, the vial was flushed with nitrogen gas, and the mixture was allowed to stir at 120 °C. After 2 h, TLC indicated the reaction to be complete. Ethylenediamine (20 mol equiv) was added to the mixture, and the stirring was continued for 1 h at room temperature. The mixture was diluted with water (10 mL) and extracted with CH₂Cl₂ (3 × 10 mL). The organic layer was dried over Na₂SO₄ and concentrated under reduced pressure. The crude material was then purified via a silica gel column using 30% EtOAc in hexanes as eluting solvent to give 42 mg (87% yield) of **2** as a white solid.

Nicotine *N*-Oxide **4**.⁴¹

Synthesized by the procedure reported in reference 41. ¹H NMR (500 MHz, CDCl₃): δ 8.63 (dd, 1H, Ar-H, *J* = 1.4, 4.8 Hz), 8.58 (d, 1H, Ar-H, *J* = 1.4 Hz), 8.29 (d, 1H, Ar-H, *J* = 7.8 Hz), 7.36 (dd, 1H, Ar-H, *J* = 4.8, 7.8 Hz), 4.24 (dd, 1H, *J* = 7.3, 11.7 Hz), 3.78 (t, 1H, *J* = 8.7 Hz), 3.63 (app q, 1H, *J*_{app} ~ 9.9 Hz), 2.98 (s, 3H, CH₃), 2.78–2.69 (m, 1H), 2.63–2.54 (m, 1H), 2.31–2.25 (m, 1H), 2.10–2.03 (m, 1H). [α]_D²⁵ = -73.5 (*c* = 2 mg/mL, CHCl₃).

Nicotine *N,N*-Dioxide **5**.⁴¹

Synthesized by the procedure reported in reference 41. ¹H NMR (500 MHz, CDCl₃): δ 8.35 (br s, 1H, Ar-H), 8.22 (d, 1H, Ar-H, *J* = 6.3 Hz), 7.84 (d, 1H, Ar-H, *J* = 7.8 Hz), 7.30 (t, 1H, Ar-H, *J* = 7.0 Hz), 4.14 (dd, 1H, *J* = 7.6, 11.2 Hz), 3.85 (t, 1H, *J* = 8.7 Hz), 3.62 (app q, 1H, *J*_{app} ~ 9.8 Hz), 3.06 (s, 3H, CH₃), 2.70–2.52 (m, 2H), 2.35–2.29 (m, 1H), 2.12–2.03 (m, 1H). [α]_D²⁵ = +1093. (*c* = 5 mg/mL, CHCl₃).

Reduction of Nicotine *N*-Oxide **4** to (*S*)-(-)-Nicotine (**3**)

In an oven-dried reaction vial equipped with a stirring bar was placed the nicotine *N*-oxide **4** (50.0 mg, 0.280 mmol) in CH₃CN (0.25 mL). (PinB)₂ (71.0 mg, 0.280 mmol) was added, the vial was flushed with nitrogen gas, and the mixture was stirred at the room temperature. After 10 min, TLC indicated the reaction to be complete. The reaction mixture was diluted with water (10 mL) and extracted with Et₂O (3 × 10 mL). The organic layer was dried over Na₂SO₄ and concentrated under reduced pressure. The crude material was then purified via a silica gel column using 80% Et₂O in hexanes as eluting solvent to give 43 mg (95% yield) of **3** as a yellowish liquid. For ¹H NMR data see reduction of **5** to **3** below.

Reduction of Nicotine *N,N*-Dioxide **5** to Nicotine *N*-Oxide **6**

In an oven-dried reaction vial equipped with a stirring bar was placed the nicotine *N,N*-dioxide **5** (50.0 mg, 0.257 mmol) in CH₃CN (0.25 mL). (PinB)₂ (65.0 mg, 0.257 mmol) was added, the vial was flushed with nitrogen gas, and the mixture was stirred at the room temperature. After 10 min, TLC indicated the reaction to be complete. The reaction mixture was directly loaded onto a silica gel column and purified using 10% MeOH in Et₂O as

eluting solvent to give 45 mg (96% yield) of **6** as a yellowish liquid. $[\alpha]_D^{25} = -146.7$ ($c = 3.5$ mg/mL, CHCl₃). R_f (SiO₂/40% MeOH in EtOAc) = 0.25. ¹H NMR (500 MHz, CDCl₃): δ 8.24 (br s, 1H, Ar-H), 8.10 (d, 1H, Ar-H, $J = 5.8$ Hz), 7.26 (m, 1H, Ar-H), 7.21 (app triplet, 1H, Ar-H, $J_{app} \sim 7.0$ Hz), 3.21 (t, 1H, $J = 7.8$ Hz), 3.08 (t, 1H, $J = 7.8$ Hz), 2.33 (app q, 1H, $J_{app} \sim 8.7$ Hz), 2.29–2.20 (s, 3H, CH₃ superimposed on m, 1H), 1.98–1.89 (m, 1H), 1.85–1.75 (m, 1H), 1.71–1.64 (m, 1H).

Reduction of Nicotine *N,N*-Dioxide **5** to (*S*)-(-)-Nicotine (**3**)

In an oven-dried reaction vial equipped with a stirring bar was placed the nicotine *N,N*-dioxide **5** (50.0 mg, 0.257 mmol) in CH₃CN (0.25 mL). (PinB)₂ (130.0 mg, 0.514 mmol) was added, the vial was flushed with nitrogen gas, and the mixture was stirred at the room temperature for 10 min, and then heated to 70 °C for 1 h. TLC indicated the reaction to be complete. The reaction mixture was directly loaded onto a silica gel column and purified using 80% Et₂O in hexanes as eluting solvent to yield 39 mg (93% yield) of **3** as a yellowish liquid. $[\alpha]_D^{25} = -165.5$ ($c = 1.5$ mg/mL, CHCl₃). For the commercial sample of nicotine used for this chemistry $[\alpha]_D^{25} = -167.2$ ($c = 1.5$ mg/mL, CHCl₃). ¹H NMR (500 MHz, CDCl₃): δ 8.55 (d, 1H, Ar-H, $J = 1.9$ Hz), 8.51 (dd, 1H, Ar-H, $J = 1.4, 4.8$ Hz), 7.71 (dt, 1H, Ar-H, $J = 1.9, 7.8$ Hz), 7.24 (dd, 1H, Ar-H, $J = 4.8, 7.8$ Hz), 3.24 (t, 1H, $J = 7.8$ Hz), 3.07 (t, 1H, $J = 8.3$ Hz), 2.30 (app q, 1H, $J_{app} \sim 8.7$ Hz), 2.23–2.16 (s, 3H, CH₃ superimposed on m, 1H), 1.98–1.91 (m, 1H), 1.85–1.78 (m, 1H), 1.74–1.70 (m, 1H).

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Infrastructural support at CCNY was provided by NIH/NCRR/RCMI Grant G12 RR03060. PFT was supported by Ruth L. Kirschstein Research Service Award 5F31 GM082025-02 from the NIH as well as the NIH RISE program. We are grateful to Allychem for a generous sample of bis(pinacolato)diboron and bis(catecholato)diboron. We thank Dr. P. Pradhan (CCNY) for assistance with the NMR experiments, Prof. B. Zajc (CCNY) for important discussions, as well as Ms. A. Joshi-Pangu and Prof. M. Biscoe (CCNY) for help with and use of their calorimeter.

REFERENCES

1. Albini A. *Synthesis*. 1993;263–277.

2. (a) Balicki R. *Synthesis*. 1989:645–646.(b) Berson JA, Cohen T. *J. Org. Chem.* 1955; 20:1461–1468.
3. Balicki R, Cybulski M, Maciejewski G. *Synth. Commun.* 2003; 33:4137–4141.
4. Jousseau B, Chanson F. *Synthesis*. 1987:55–57.
5. Zhang Y, Lin R. *Synth. Commun.* 1987; 17:329–332.
6. Ilanakumaran P, Chandrasekaran S. *Tetrahedron Lett.* 1995; 36:4881–4882.
7. Yoo BW, Choi JW, Yoon CM. *Tetrahedron Lett.* 2005; 47:125–126.
8. Yoo BW, Park MC. *Synth. Commun.* 2008; 38:1646–1650.
9. Yadav JS, Subba Reddy BV, Murlidhar Reddy M. *Tetrahedron Lett.* 2000; 41:2663–2665.
10. Ilias M, Barman DC, Prajapati D, Sandhu JS. *Tetrahedron Lett.* 2002; 43:1877–1879.
11. Barton DHR, Fekih A, Lusinch X. *Tetrahedron Lett.* 1985; 26:4603–4606.
12. Malinowski M. *Synthesis*. 1987; 8:732–734.
13. Chandrasekhar S, Reddy CR, Rao RJ, Rao JM. *Synlett.* 2002:349–351.
14. Han JH, Choi JW, Choi KI, Kim JH, Yoon CM, Yoo HW. *Bull. Korean Chem. Soc.* 2005; 26:1921–1922.
15. Alonso F, Radivoy G, Yus M. *Tetrahedron.* 2000; 56:8673–8678.
16. Han JH, Choi KI, Kim JH, Yoo BW. *Synth. Commun.* 2004; 34:3197–3201.
17. Saini A, Kumar S, Sandhu JS. *Synlett.* 2006:395–398.
18. Kano S, Tanaka Y, Sugino E, Hibino S. *Synthesis*. 1980:695–697.
19. Yoon NM. *Pure Appl. Chem.* 1996; 68:843–848.
20. Wang Y, Espenson JH. *Org. Lett.* 2000; 2:3525–3526. [PubMed: 11082025]
21. Kumar S, Saini A, Sandhu JS. *Tetrahedron Lett.* 2005; 46:8737–8739.
22. Hayashi E, Ijima Ch. *Yakugaku Zasshi.* 1962; 82:1093–1102. [PubMed: 13953261]
23. Daniher FA, Hackley BE Jr. *J. Org. Chem.* 1966; 31:4267–4268.
24. Bonini BF, Maccagnani G, Mazzanti G, Pedrini P. *Tetrahedron Lett.* 1979; 20:1799–1800.
25. Olah GA, Arvanaghi M, Vankar YD. *Synthesis*. 1980:660–661.
26. Lutz WB, Lazarus S, Klutchko S, Meltzer RI. *J. Org. Chem.* 1964; 29:1645–1647.
27. Howard E Jr, Olszewski WF. *J. Am. Chem. Soc.* 1959; 81:1483–1484.
28. Kagami H, Motoki S. *J. Org. Chem.* 1978; 43:1267–1268.
29. Yoshimura T, Asada K, Oae S. *Bull. Chem. Soc. Jpn.* 1982; 55:3000–3003.
30. Takeshita M, Yoshida S. *Heterocycles.* 1990; 30:871–874.
31. Bjørsvik H-R, Gambarotti C, Jensen VR, González RR. *J. Org. Chem.* 2005; 70:3218–3224. [PubMed: 15822984]
32. Bae S, Lakshman MK. *J. Org. Chem.* 2008; 73:1311–1319. [PubMed: 18205381]
33. Carter, CAG.; John, KD.; Mann, G.; Martin, RL.; Cameron, TM.; Baker, RT.; Bishop, KL.; Broene, RD.; Westcott, SA. Bifunctional Lewis Acid Reactivity of Diol-Derived Diboron Reagents. In: Shapiro, PJ.; Atwood, DA., editors. *Group 13 Chemistry/From Fundamentals to Applications*. Washington, DC: American Chemical Society; 2002. p. 70-87. ACS Symposium Series 822
34. Larkin JD, Bhat KL, Markham GD, James TD, Brooks BR, Block CW. *J. Phys. Chem. A.* 2008; 112:8446–8454. [PubMed: 18707068]
35. Sana M, Leroy G, Wilante C. *Organometallics.* 1991; 10:264–270.
36. Köster R, Morita Y. *Angew. Chem., Int. Ed.* 1966; 5:580.
37. Köster R, Morita Y. *Liebigs Ann. Chem.* 1967; 704:70–90.
38. Köster R, Arora S, Binger P. *Angew. Chem., Int. Ed.* 1969; 8:205.
39. Kabalka GW, Hedgecock HC Jr. *J. Org. Chem.* 1975; 40:1776–1779.
40. Tsunoda H, Ohkubo A, Taguchi H, Seio K, Sekine M. *J. Org. Chem.* 2008; 73:1217–1224. [PubMed: 18201098]
41. Uwai K, Sato H, Kazakami N, Matsuzaki H, Takeshita M. *Arkivoc.* 2003:viii, 211–219.
42. Phillipson JD, Handa SS. *Phytochemistry.* 1975; 14:2683–2690.

43. Johnson AW, King TJ, Turner JR. *J. Chem. Soc.* 1958:3230–3231.
44. Hawkeswood S, Stephan DW. *Dalton Trans.* 2005:2182–2187. [PubMed: 15957061]
45. Soderquist JA, Najafi MR. *J. Org. Chem.* 1986; 51:1330–1336.
46. Laitar DS, Müller P, Sadighi JP. *J. Am. Chem. Soc.* 2005; 127:17196–17197. [PubMed: 16332062]
47. Boron resonance of (pinB)₂ in the following solvents: benzene-*d*₆ δ = 34.2 ppm, CD₃CN δ = 30.6 ppm, CDCl₃ δ = 30.6 ppm, and THF-*d*₈ δ = 31.7 ppm. Boron resonance of (pinB)₂O in the following solvents: benzene-*d*₆ δ = 22.2 ppm (21.6 in ref 41), CD₃CN δ = 22.4 ppm, CDCl₃ δ = 21.7 ppm, and THF-*d*₈ δ = 22.5 ppm. Signals were referenced to BF₃•Et₂O.
48. (a) Bettinger HF, Filthaus M, Bornemann H, Oppel IM. *Angew. Chem. Int. Ed.* 2008; 47:4744–4747. (reported ¹¹B resonance of pinB–OH at δ = 22.5 ppm in benzene-*d*₆). (b) Salomon MA, Braun T, Penner A. *Angew. Chem., Int. Ed.* 2008; 47:8867–8871. (reported ¹¹B resonance of pinB–OH at δ = 21.6 ppm in PhMe-*d*₈).
49. Because of the similarities in the chemical shifts of (pinB)₂O and pinB–OH, we cannot definitively say that the former alone is formed in the previously described deoxygenation of nucleoside benzotriazol-1-yl ethers (reference 32). However, initial formation of (pinB)₂O is entirely plausible in that chemistry.
50. See for examples: Campeau L-C, Stuart DR, Leclerc J-P, Bertrand-Leperle M, Villemure E, Sun H-Y, Lasserre S, Guimond N, Lecavallier M, Fagnou K. *J. Am. Chem. Soc.* 2009; 131:3291–3306. [PubMed: 19215128] (b) Huestis MP, Fagnou K. *Org. Lett.* 2009; 11:1357–1360. [PubMed: 19226145] (c) Campeau L-C, Bertrand-Leperle M, Leclerc J-P, Villemure E, Gorelsky S, Fagnou K. *J. Am. Chem. Soc.* 2008; 130:3276–3277. [PubMed: 18302383] (d) Campeau L-C, Rousseaux S, Fagnou K. *J. Am. Chem. Soc.* 2005; 127:18020–18021. [PubMed: 16366550]
51. Gong X, Song G, Zhang H, Li X. *Org. Lett.* 2011; 13:1766–1769. [PubMed: 21388218]
52. (a) Londregan AT, Jennings S, Wei L. *Org. Lett.* 2011; 13:1840–1843. [PubMed: 21375291] (b) Londregan AT, Jennings S, Wei L. *Org. Lett.* 2010; 12:5254–5257. [PubMed: 20958085]
53. Ferrer M, Sánchez-Baeza F, Messeguer A. *Tetrahedron.* 1997; 53:15877–15888.
54. Rosenau T, Schmid P, Kosma P. *Tetrahedron.* 2005; 61:3483–3487.




Figure 1. Evaluating the reduction of 4-methylmorpholine *N*-oxide with (pinB)₂ by NMR (at 500 MHz in CDCl₃). **Panel A:** spectrum of 4-methylmorpholine *N*-oxide prior to addition of (pinB)₂. **Panel B:** immediately after addition of 1.0 molar equivalent (pinB)₂ (mixture was not shaken). **Panel C:** immediately after shaking the reaction mixture. **Panel D:** mixture of (pinB)₂ and 1.5 molar equivalents of 4-methylmorpholine *N*-oxide. **Panel E:** spectrum of pure 4-methylmorpholine.







Figure 2. Monitoring the reaction between 4-methoxypyridine *N*-oxide hydrate and (pinB)₂ in CD₃CN, by NMR.




Scheme 1.
Deoxygenation of *O*⁶-(benzotriazol-1-yl)inosine and 2'-deoxyinosine using (pinB)₂



Scheme 2.
Reduction of 2',3',5'-Tri-*O*-(*tert*-butyldimethylsilyl)adenosine N1-Oxide Using (pinB)₂



Scheme 3.
Reduction of Nicotine *N*-Oxides with (pinB)₂













Scheme 4.
Plausible Mechanism for the *N*-Oxide Reduction

Table 1





Generality of Amine *N*-Oxide Reduction with Diboron Reagents^a

entry	amine <i>N</i> -oxide	solvent	conditions	yield ^b
<i>Reductions using (pinB)₂:</i>				
1		CDCl ₃	rt, 5 min	Quant ^c
2		CDCl ₃	rt, 5 min	Quant ^c
3		CH ₃ CN	70 °C, 8 h	76%
4		CH ₃ CN Diglyme	70 °C, 21 h 120 °C, 12 h	64% ^d 72%
5		CH ₃ CN	70 °C, 5 h	91%

entry	amine <i>N</i> -oxide	solvent	conditions	yield ^b
6		CH ₃ CN	70 °C, 6 h	96%
7		CH ₃ CN	70 °C, 8 h	86%
8		Diglyme	120 °C, 10 h	67%
9		CH ₃ CN	70 °C, 8 h	82%
10		CH ₃ CN	70 °C, 8 h	84%

entry	amine <i>N</i> -oxide	solvent	conditions	yield ^b
11		1,4-Dioxane Diglyme	100 °C, 10 h 120 °C, 4 h	66% 72%
12		CH ₃ CN	70 °C, 8 h	80%
13		CH ₃ CN	rt, 10 min	95%
14		CH ₃ CN	rt, 5 min	97%
15		CH ₃ CN	rt, 10 min	92%

Reductions using (catB)₂:^e

entry	amine <i>N</i> -oxide	solvent	conditions	yield ^b
16		CH ₃ CN	120 °C, 12 h	78%
17		CH ₃ CN	120 °C, 12 h	79%
18		CH ₃ CN	120 °C, 12 h	88%
19		CH ₃ CN	120 °C, 24 h	65%

^aReactions were conducted at 1 M amine *N*-oxide concentration.





^bWhere reported, yield is of isolated and purified product.


^cNMR tube experiment showed complete disappearance of the *N*-oxide resonance(s) and appearance of product resonance(s). Some chemical shift differences and/or signal broadening was observed, likely due to coordination of the amine to (pinB)₂O in the reaction mixture (as reported in reference 33).

^dReaction was incomplete.

^eBis(catecholato)diboron: (catB)₂.

Table 2Use of Water as Reaction Medium for Reductions Using (pinB)₂^a

entry	amine <i>N</i> -oxide	conditions	yield ^b
1		70 °C, 6 h	78%
2	• H ₂ O 	110 °C, 10 h	72%
3		70 °C, 8 h	80%
4		70 °C, 7 h	82%

entry	amine <i>N</i> -oxide	conditions	yield ^b
5		rt, 15 min	94%

^aReactions were conducted at 1 M amine *N*-oxide concentration in water.

^bYields reported are of isolated and purified products.