

Regular Graphs with Maximum Forest Number

Avapa Chantasartrassmee*

Department of Mathematics, School of Science,
University of the Thai Chamber of Commerce,
Bangkok 10400, Thailand
avapa_cha@utcc.ac.th

Narong Punnim†

Department of Mathematics, Srinakharinwirot University,
Sukhumvit 23, Bangkok 10110, Thailand
narongp@swu.ac.th

Abstract

Let G be a graph and $F \subseteq V(G)$. Then F is called an induced forest of G if $\langle G \rangle$ contains no cycle. The *forest number*, $f(G)$, of G is defined by

$$f(G) := \max\{|F| : F \text{ is an induced forest of } G\}.$$

It was proved by the second author in [6] that if G is an r -regular graph of order n , then $f(G) \leq \lfloor \frac{nr-2}{2(r-1)} \rfloor$. It was also proved that the bound is sharp by constructing an r -regular graph H of order n with $f(H) = \lfloor \frac{nr-2}{2(r-1)} \rfloor$.

In this paper we consider the problem of determining which r -regular graphs G of order n have the forest number $\lfloor \frac{nr-2}{2(r-1)} \rfloor$. The problem was asked by Bau and Beineke [1] for $r = 3$ and, in this particular case, it was answered by the second author in [7]. We are able to answer the problem for all $r \geq 4$. More precisely, we are able to obtain an algorithm of finding all r -regular graphs G of order n with $f(G) = \lfloor \frac{nr-2}{2(r-1)} \rfloor$. Furthermore, we prove that if $\mathcal{R}(r^n; f = \lfloor \frac{nr-2}{2(r-1)} \rfloor)$ is the set of all r -regular graphs G of order n with $f(G) = \lfloor \frac{nr-2}{2(r-1)} \rfloor$ and $G_1, G_2 \in \mathcal{R}(r^n; f = \lfloor \frac{nr-2}{2(r-1)} \rfloor)$, then there exists a sequence of switchings $\sigma_1, \sigma_2, \dots, \sigma_t$ such that for each $i = 1, 2, \dots, t$, $G_1^{\sigma_1 \sigma_2 \dots \sigma_i} \in \mathcal{R}(r^n; f = \lfloor \frac{nr-2}{2(r-1)} \rfloor)$ and $G_1^{\sigma_1 \sigma_2 \dots \sigma_t} = G_2$.

Keywords: Degree sequence, Forest number, Regular graphs

AMS Mathematics Subject Classification (2000): 05C05, 05C07

*This work was carried out under financial support from the University of the Thai Chamber of Commerce.

†This work was carried out under financial support from the Thailand Research Fund.

References

- [1] S. Bau and L. W. Beineke, *The decycling number of graphs*, Austral. J. Combinatorics, **25**(2002), 285-298.
- [2] G. Chartrand and L. Lesniak, "GRAPHS & DIGRAPHS", 4th Edition, Chapman & Hall/CRC, A CRC Press Company, 2005.
- [3] R. B. Eggleton and D. A. Holton, *Graphic sequences, Combinatorial mathematics, VI (Proc. Sixth Austral. Conf., Univ. New England, Armidale, 1978)*, Lecture Notes in Math. **748**(1978),1–10.
- [4] S. Hakimi, *On the realizability of a set of integers as the degree of the vertices of a graph*, SIAM J. Appl. Math., **10**(1962), 496-506.
- [5] V. Havel, *A remark on the existence of finite graphs (Czech)*, Casopis Pest. Mat., **80** (1955), 477-480.
- [6] N. Punnim, *Decycling regular graphs*, Austral. J. Combinatorics, **32**(2005), 147-162.
- [7] N. Punnim, *The decycling number of cubic graphs*, Lecture Notes in Computer Science (LNCS), **3330**(2005), 141-145.
- [8] N. Punnim, *Decycling connected regular graphs*, Austral. J. Combinatorics, **35**(2006), 155-169.
- [9] N. Punnim, *The decycling number of cubic planar graphs*, Lecture Notes in Computer Science (LNCS), **4381**(2007), 149-161.