

Regularizations of general singular integral operators

Constanze Liaw

Texas A&M University

March 19th, 2011

This talk is based on joint work with Sergei Treil.

This work is accepted by 'Revista Matematica Iberoamericano' for publication and available on the arXiv:1010.6184. For further questions please do not hesitate to contact: conni 'at' math 'dot' tamu 'dot' edu

Outline

What is a singular integral operator (SIO)?

Smooth mollifying multipliers

- A trivial idea

- Families of smooth mollifying multipliers

- Uniform boundedness of regularized kernels

Uniform boundedness of truncations

The problem

- Consider kernel $K(s, t)$ which is singular near the diagonal, i.e. $K(s, \cdot)$ and $K(\cdot, t)$ are not in L^1_{loc} near $s = t$
- A SIO is an operator T on $L^2(\mu)$ is *formally* given by

$$Tf(s) = \int K(s, t)f(t) d\mu(t)$$

- This means that the above integral is not defined even for the simplest functions f
- What meaning do we give this formal expression?
- If T is the classical Hilbert Transform on the real line, then the integral exists in the sense of principal value

$$\lim_{\alpha \rightarrow 0^+} \int_{|s-t| > \alpha} \frac{f(t)}{s-t} dt \quad \text{for } f \in C_c^1(\mathbb{R})$$

- This approach is not possible, if $d\mu(t) \neq dt$

Some of literature's remedies

- In the general situation the boundedness in L^p is often defined as the uniform boundedness (independent of $\varepsilon \rightarrow 0$) of either

truncated operators:
$$T_\varepsilon f(s) = \int_{|s-t|>\varepsilon} K(s,t)f(t)d\mu(t),$$
 or

smooth regularizations, e.g.:
$$T_\varepsilon f(s) = \int_{\mathbb{R}} \frac{f(t)}{s-t+i\varepsilon} d\mu(t)$$

- The *corresponding SIO* is then the limit point (in WOT)
- T is unique up to '+multiplication by L^∞ -function'

Definitions

- Radon measures μ and ν in \mathbb{R}^N (no grow, doubling condition)
- A *singular kernel* in \mathbb{R}^N (wrt μ and ν) is locally $L^2(\mu \times \nu)$ off the diagonal $\{(s, t) \in \mathbb{R}^N \times \mathbb{R}^N : s = t\}$
- K is of order d , if the kernel \tilde{K} is locally $L^2(\mu \times \nu)$,

$$\tilde{K}(s, t) = \begin{cases} K(s, t)|s - t|^d, & s \neq t \\ 0 & s = t \end{cases}$$

- Formal singular integral operator with the kernel K is *restrictedly bounded* in L^p if

$$\left| \int K(s, t) f(t) g(s) d\mu(t) d\nu(s) \right| \leq C \|f\|_{L^p(\mu)} \|g\|_{L^{p'}(\nu)},$$

for all bounded f, g with separated compact supports

- The least C (for p, μ and ν fixed) is the *restricted norm* of K

A trivial idea

- If $K(s, t)$ has restricted L^p norm C , then so does the kernel

$$K(s, t)e^{-ia \cdot t}e^{ia \cdot s} \quad \text{for any } a \in \mathbb{R}^N$$

is also restrictedly bounded with the same constant

- Averaging over all a with weight $\rho \in L^1(dx)$, kernel

$$\int_{\mathbb{R}^N} \rho(a)K(s, t)e^{-ia \cdot t}e^{ia \cdot s} da = \widehat{\rho}(t - s)K(s, t)$$

is also restrictedly bounded with restricted norm $C\|\rho\|_1$

Lemma

Let K be a restrictedly L^p bounded kernel with a bound C .

Assume that $\rho \in L^1(dx)$ and let $M = 1 - \widehat{\rho}$, $M_\varepsilon(x) := M(x/\varepsilon)$.

Then the kernels $K_\varepsilon(s, t) := K(s, t)M_\varepsilon(t - s)$ are L^p restrictedly bounded with constant $(1 + \|\rho\|_1)C$.

Example: The Hilbert Transform

Lemma (repeated from previous slide)

Let K be a restrictedly L^p bounded kernel with a bound C . Assume that $\rho \in L^1(dx)$ and let $M = 1 - \widehat{\rho}$, $M_\varepsilon(x) := M(x/\varepsilon)$. Then the kernels $K_\varepsilon(s, t) := K(s, t)M_\varepsilon(t - s)$ are L^p restrictedly bounded with constant $(1 + \|\rho\|_1)C$.

- On \mathbb{R} consider Hilbert Transform $K(s, t) = \pi^{-1}(s - t)^{-1}$ and the weight $\rho(x) = e^{-x}\chi_{[0, \infty)}(x)$
- We get $\widehat{\rho}(s) = (1 + is)^{-1}$, so $M(s) = 1 - \widehat{\rho}(s) = \frac{s}{s - i}$ and

$$M_\varepsilon(s) := M(s/\varepsilon) = \frac{s}{s - i\varepsilon}$$

- Regularization with this mollifying factor gives the kernel

$$K_\varepsilon(s, t) = \frac{1}{\pi} \cdot \frac{1}{s - t} M_\varepsilon(t - s) = \frac{1}{\pi} \cdot \frac{1}{s - t + i\varepsilon}$$

Many smooth mollifying multipliers

Theorem

Let M be a function on \mathbb{R}^N such that $M \equiv 0$ near 0, and $1 - M \in H^k(\mathbb{R}^N) = W^{k,2}(\mathbb{R}^N)$, $k > N/2$.

Then the functions $M_\varepsilon(s, t) := M((t - s)/\varepsilon)$ is a family of smooth regularizing multipliers, meaning that:

- (i) $M_\varepsilon(s, t) \rightarrow 1$ as $\varepsilon \rightarrow 0$ uniformly on all sets $\{s, t \in \mathbb{R}^N : |s - t| > a\}$, $a > 0$.
- (ii) For any singular kernel K the regularized kernels $K_\varepsilon = KM_\varepsilon$ are in $L^2_{\text{loc}}(\mu \times \nu)$.
- (iii) If the kernel is restrictedly bounded in L^p , and the measures μ and ν do not have common atoms, then the regularized integral operators T_ε with kernels K_ε are uniformly (in ε) bounded as operators $L^p(\mu) \rightarrow L^p(\nu)$.

Main application: Uniform boundedness

- Let T be a limit point of T_ε , $\varepsilon \rightarrow 0$ in WOT

Theorem

Let μ and ν be Radon measures in \mathbb{R}^N without common atoms. Assume that a kernel $K \in L^2_{\text{loc}}(\mu \times \nu)$ is L^p restrictedly bounded, with the restricted norm C .

Then the integral operator with T kernel K is a bounded operator $L^p(\mu) \rightarrow L^p(\nu)$ with the norm at most $2C$.

- As in the classical case, T is unique up to ‘+multiplication by L^∞ -function’

Case of common atoms

When μ and ν do have common atoms, the boundedness of the singular integral operator can be defined as follows:

- Decompose

$$\mu = \tilde{\mu} + \mu_0, \quad \nu = \tilde{\nu} + \nu_0,$$

where μ_0 and ν_0 are the parts of μ and ν supported on their common atoms

- Use the Theorem to check the L^p boundedness as an operator $L^p(\mu) \rightarrow L^p(\tilde{\nu})$ or $L^p(\tilde{\mu}) \rightarrow L^p(\nu)$
- It remains to check the block acting $L^p(\mu_0) \rightarrow L^p(\nu_0)$. But the bilinear form of this block is well defined for functions supported at finitely many points, so there is no problem defining this block

Idea of Theorem

- Under some additional assumptions, the restricted L^p boundedness implies the uniform boundedness of the truncated operators T_ε ,

$$T_\varepsilon f(s) = \int_{|s-t|>\varepsilon} K(s,t)f(t)d\mu(t)$$

- These assumptions are satisfied for classical operators like generalized Riesz Transforms (treated as a vector-valued transformation), or Ahlfors–Beurling operator

Summary

What is a singular integral operator (SIO)?

Smooth mollifying multipliers

- A trivial idea

- Families of smooth mollifying multipliers

- Uniform boundedness of regularized kernels

Uniform boundedness of truncations

Thank you for your attention.