

REGULARLY VARYING SEQUENCES

J. GALAMBOS AND E. SENETA¹

ABSTRACT. A simple necessary and sufficient condition is developed for a sequence $\{\theta(n)\}$, $n=0, 1, 2, \dots$, of positive terms, to satisfy $\theta(n)=R(n)$, $n \geq 0$, where $R(\cdot)$ is a regularly varying function on $[0, \infty)$. The condition (2.1), below, leads to a Karamata-type exponential representation for $\theta(n)$. Various associated difficulties are also discussed. (The results are of relevance in connection with limit theorems in various branches of probability theory.)

1. Introduction. A function $R(\cdot)$, defined, finite, positive and measurable on $[A, \infty)$ for some $A \geq 0$, is said to be regularly varying if for each $\lambda > 0$

$$(1.1) \quad \lim_{x \rightarrow \infty} \frac{R(\lambda x)}{R(x)} = \phi(\lambda)$$

where $0 < \phi(\lambda) < \infty$. (In actual fact a weaker definition can be used, for the assumption that this positive finite limit property obtains for all λ in a subset of positive measure of $(0, \infty)$ implies that it obtains for all $\lambda \in (0, \infty)$.) Since $\phi(\lambda)$ is a positive measurable solution of the functional equation

$$(1.2) \quad \phi(uv) = \phi(u)\phi(v), \quad u, v > 0,$$

it is well known that $\phi(\lambda) = \lambda^\rho$ for some finite ρ , and so we can write $R(x) = x^\rho L(x)$ where

$$(1.3) \quad \lim_{x \rightarrow \infty} \frac{L(\lambda x)}{L(x)} = 1, \quad \text{for each } \lambda > 0;$$

such a regularly varying function, for which the index ρ of regular variation is zero, is called slowly varying.

Received by the editors January 30, 1973.

AMS (MOS) subject classifications (1970). Primary 26A12, 40A25; Secondary 10K30, 60F05.

Key words and phrases. Regularly varying functions, regularly varying sequences, imbedding sequences in functions, exponential representations of sequences and functions.

¹ Partial support by the Office of Naval Research on contract N00014-67-A-0151-0017 awarded to the Department of Statistics, Princeton University, is gratefully acknowledged.

© American Mathematical Society 1973

The two most important properties of regularly varying functions (from which others are easily deducible) are:

- (i) The convergence in (1.1) (or, equivalently, (1.3)) is uniform for λ in any fixed interval $[a, b]$, $0 < a < b < \infty$.
- (ii) For some $B \geq A$, a slowly varying function L has representation

$$(1.4) \quad L(x) = \exp\left\{\eta(x) + \int_B^x \frac{\varepsilon(t)}{t} dt\right\}, \quad x \geq B,$$

where $\eta(x) \rightarrow c$ ($|c| < \infty$) as $x \rightarrow \infty$ and $\varepsilon(t) \rightarrow 0$ as $t \rightarrow \infty$, both being measurable and bounded. Conversely, any function L having representation (1.4) is clearly slowly varying.

The systematic development of the notion of a regularly varying function, of great importance in probability theory, is due, for continuous functions, to Karamata ([1930], [1933]), and in the above setting to various later authors. A sketch of the basic history and theory is given in §§1 and 4 of the recent paper of Bojanić and Seneta [1971]. We pause to note only the result of de Bruijn [1959, §4] that in (1.4), $\varepsilon(t)$ may be taken as continuous (the less desirable properties of L being perpetuated by $\eta(x)$). This last remark enables us to deduce that as $x \rightarrow \infty$

$$R(x) = x^\rho L(x) \sim x^\rho L_1(x) = R_1(x)$$

where $(R_1 x) = x^\rho \exp\{c + \int_B^x \varepsilon(t)/t dt\}$, $x \geq B$, is a continuously differentiable regularly varying function such that

$$(1.5) \quad xR_1'(x)/R_1(x) \rightarrow \rho$$

as $x \rightarrow \infty$, since

$$(1.6) \quad xR_1'(x)/R_1(x) = \rho + \varepsilon(x), \quad x \geq B.$$

Conversely, any function R_1 satisfying (1.5) is regularly varying (with index ρ), as can be seen by defining $\varepsilon(x)$ from (1.6) and integrating for R_1 , to obtain the required representation.

More recently, a problem of the following *genre* has occurred in several probabilistic contexts. Given a sequence $\{\theta(n)\}$, $n=0, 1, 2, \dots$, of positive numbers, when is it possible to imbed it in a regularly varying function? In other words, when is it possible to find a regularly varying function $R(x)$ such that $R(n)=\theta(n)$? If it is possible, then it follows, for example, from either property (i) or (ii) of regularly varying functions, that

$$(1.7) \quad \theta(n+1)/\theta(n) \rightarrow 1$$

as $n \rightarrow \infty$. As examples of results obtained so far, we mention that of

de Haan [1970, pp. 6–8], who shows that the imbedding is possible if (a) $\{\theta(n)\}$ is monotone, and (b) $\theta(nm)/\theta(n) \rightarrow m^\rho$ for all positive m as $n \rightarrow \infty$, where ρ is finite, and that of R. S. Slack, which asserts that in (b), m^ρ may be replaced by $\phi(m)$, $0 < \phi(m) < \infty$, if (1.7) is imposed as an additional hypothesis, with the same conclusion.²

This type of problem, concerning regular behavior of sequences, was studied prior to the papers of Karamata mentioned above. The reader may want to consult the works of Schmidt [1925] and Schur [1930] in this regard.

There is some difficulty in attempting the obvious approach to the sequence problem along the lines of the elegant definition (1.1). Thus, it is possible to construct a sequence of positive numbers $\{\theta(n)\}$ satisfying simultaneously (for positive integer k), as $n \rightarrow \infty$,

$$(1.8) \quad \theta(nk)/\theta(n) \rightarrow 1, \quad \theta(n+1)/\theta(n) \nrightarrow 1,$$

so that the requirement (1.7) is broken.

To carry out such a construction, let $\omega(n)$ denote the number of prime divisors of n . Let $\theta(n) = \omega(n) + (\log \log n)^{1/2}$, $n \geq 2$. It is known (Kubilius [1964, p. 39]) that there exists a subsequence p_{i_1}, p_{i_2}, \dots of the primes such that $\omega(p_{i_n} - 1) \sim \log \log p_{i_n}$ as $n \rightarrow \infty$. If we consider the subsequence $\theta(p_{i_n})/\theta(p_{i_n} - 1)$ of the sequence $\theta(n+1)/\theta(n)$, we readily see that its limit is zero, since $\omega(p_{i_n}) = 1$, so that $\theta(n+1)/\theta(n) \nrightarrow 1$. If we consider $\theta^*(n) = \omega(n) + \log \log n$ instead, we obtain that $\theta^*(p_{i_n})/\theta^*(p_{i_n} - 1) \rightarrow \frac{1}{2}$ as $n \rightarrow \infty$, also a satisfactory result. On the other hand, we have for integer $k \geq 1$

$$\theta(nk) = \omega(nk) + (\log(\log n + \log k))^{1/2},$$

whence for large n ,

$$\begin{aligned} \omega(n) + (\log(\log n + \log k))^{1/2} \\ \leq \theta(nk) \leq \omega(n) + \omega(k) + (\log(\log n + \log k))^{1/2} \end{aligned}$$

from the definition ω ; and so $\theta(nk)/\theta(n) \rightarrow 1$, each integer $k \geq 1$. (It may be proved similarly that $\theta^*(nk)/\theta^*(n) \rightarrow 1$.)

To conclude this section, it is necessary to mention that Ibragimov and Linnik [1971, p. 397] seem to cite, as an example of a sequence of positive terms such that $\theta(nk)/\theta(n) \rightarrow 1$ as $n \rightarrow \infty$, but $\theta(n+1)/\theta(n) \nrightarrow 1$, the sequence given by $\theta(n) = \omega(n) + (\log n)^{1/2}$. Whereas it is easy to check that $\theta(nk)/\theta(n) \rightarrow 1$ for each positive integer k , the proposition regarding

² Mentioned in a letter to one of the authors from G. E. H. Reuter. Slack's result occurs in a branching process context, and his method is not known to us. Nevertheless, one of us has constructed a possibly different proof.

$\theta(n+1)/\theta(n)$ appears to be a deeper one whose validity or otherwise is not known; but note that $\theta(p_{j_n})/\theta(p_{j_n}-1) \rightarrow 1$ as $n \rightarrow \infty$.

2. Regularly varying sequences. We call a sequence $\{\theta(n)\}$ of positive terms *regularly varying* if there is a sequence of positive terms $\{\alpha(n)\}$ satisfying

$$(2.1a) \quad \theta(n) \sim K\alpha(n), \quad K \text{ a positive constant,}$$

$$(2.1b) \quad n(1 - \{\alpha(n-1)/\alpha(n)\}) \rightarrow \rho, \quad \rho \text{ finite.}$$

The number ρ will be called the index of regular variation. The case $\rho=0$ may be called slowly varying. It is first necessary to note that there exist regularly varying sequences $\{\theta(n)\}$ which themselves do not satisfy the condition (2.1b) with α replaced by θ (just as not all regularly varying functions can satisfy (1.5), although $R(x) \sim R_1(x)$ always). For example, if we take $\theta(n) = 1 + (-1)^n/n, n \geq 2$, then $\theta(n)$ is regularly varying with index 0, since appropriate $\alpha(n)$ is given by $\alpha(n) = 1$. However

$$n(1 - \{\theta(n-1)/\theta(n)\}) \rightarrow -2, \quad n \rightarrow \infty, n \text{ odd,} \\ \rightarrow 2, \quad n \rightarrow \infty, n \text{ even.}$$

We shall say that a sequence of positive terms $\{\theta(n)\}, n=0, 1, 2, \dots$, is *imbeddable* in a regularly varying function R on $[0, \infty)$ if $R(n) = \theta(n), n \geq 0$.

LEMMA. *If $\{\theta(n)\}, n=0, 1, 2, \dots$, is a regularly varying sequence of index ρ , then it has representation*

$$(2.2) \quad \theta(n) = n^\rho a(n) \exp\left\{ \sum_{j=1}^n \frac{\varepsilon(j)}{j} \right\}, \quad n \geq 1,$$

where, as $n \rightarrow \infty, a(n) \rightarrow$ positive limit, $\varepsilon(n) \rightarrow 0$.

PROOF. Since $\theta(n) \sim K\alpha(n)$, we may assume without loss of generality that $1 - \alpha(m-1)/\alpha(m) \equiv \rho/m + \varepsilon(m)/m, m \geq 1$, is less than unity in modulus for all $m \geq 1$, by changing the first few terms of $\{\alpha(n)\}$. Since for $|x| < 1, -\log(1-x) = \sum_{k=1}^\infty x^k/k$, we obtain

$$-\log\left\{ \frac{\alpha(m-1)}{\alpha(m)} \right\} - \sum_{k=2}^\infty \frac{1}{k} \left\{ 1 - \frac{\alpha(m-1)}{\alpha(m)} \right\}^k = \frac{\rho}{m} + \frac{\varepsilon(m)}{m}.$$

Summing over m from 1 to n ,

$$\log \alpha(n) - \log \alpha(0) - \sum_{m=1}^n \sum_{k=2}^\infty \frac{1}{k} \left\{ 1 - \frac{\alpha(m-1)}{\alpha(m)} \right\}^k = \rho \sum_{m=1}^n \frac{1}{m} + \sum_{m=1}^n \frac{\varepsilon(m)}{m}.$$

Now it is well known that $\sum_{m=1}^n m^{-1} - \log n = \gamma + o(1)$ as $n \rightarrow \infty$, where γ is a positive constant. Further since for each integer $k \geq 2$, from (2.1b),

$$|\{1 - \alpha(m - 1)/\alpha(m)\}^k| < (|\rho| + \delta_1)^k/m^k$$

for arbitrary fixed positive δ_1 , and positive integer m sufficiently large (independently of k), we have the upper bound

$$= ((|\rho| + \delta_1)/m^{1/4})^k/m^{3k/4} \leq \delta_2^k/m^{3/2}$$

for $k \geq 2$ and with $0 < \delta_2 < 1$, for m large (independent of k). Thus we obtain that the series $\sum_{m=1}^\infty \sum_{k=2}^\infty k^{-1} \{1 - \alpha(m - 1)/\alpha(m)\}^k$ is (absolutely) convergent. Hence it follows that

$$\alpha(n) = n^\rho a(n) \exp\left\{ \sum_{j=1}^n \frac{\varepsilon(j)}{j} \right\}$$

where, as $n \rightarrow \infty$, $a(n) \rightarrow$ positive limit, $\varepsilon(n) \rightarrow 0$. Since $K(n) \equiv \theta(n)/\alpha(n) \rightarrow$ pos. const. by (2.1a), it follows that $\theta(n)$ has the same kind of representation, as required.

THEOREM. *A sequence of positive terms $\{\theta(n)\}$, $n \geq 0$, is imbeddable in a regularly varying function R on $[0, \infty)$ if and only if the sequence is also regularly varying.*

PROOF. *Sufficiency.* If $\{\theta(n)\}$ is regularly varying with index ρ , we have representation (2.2) available for $\theta(n)$, $n \geq 1$. Put (where $[u]$ denotes the integer part of u)

$$R(0) = \theta(0),$$

$$R(x) = x^\rho a([x]) \exp\left\{ \int_0^x \frac{\varepsilon([t + 1])}{[t + 1]} dt \right\},$$

for $x > 0$, defining $a(0)$ by $a(0) = 1$, say. A glance at (1.4), or a direct verification using the definition of a regularly varying function, shows $R(x)$ is regularly varying with index ρ , and $R(n) = \theta(n)$, $n \geq 0$.

Necessity. Conversely, if $R(x)$ is a regularly varying function on $[0, \infty)$ of index ρ then we have, for $x \geq B \geq 0$,

$$R(x) = x^\rho \exp\left\{ \eta(x) + \int_B^x \frac{\varepsilon(t)}{t} dt \right\}$$

from (1.4), where $\varepsilon(t)$ may be taken as continuous for $x \geq B$, $\varepsilon(t) \rightarrow 0$ as $t \rightarrow \infty$, $\eta(x) \rightarrow c$ ($|c| < \infty$). For integer $n \geq 0$, we have $\theta(n) = R(n)$, so for

integer $n \geq B$

$$\theta(n) = n^\rho \exp\left\{\eta(n) + \int_B^n \frac{\varepsilon(t)}{t} dt\right\},$$

since we are assuming $\theta(n)$ is imbeddable in $R(x)$.

To verify (2.1a) and (2.1b), put $\alpha(n) = n^\rho \exp\left\{\int_B^n \varepsilon(t)/t dt\right\}$ for all sufficiently large n ; then $\theta(n) \sim K\alpha(n)$, K a positive constant; also

$$n\left(1 - \frac{\alpha(n-1)}{\alpha(n)}\right) = n\left(1 - (1-1/n)^\rho \exp\left\{-\int_{n-1}^n \frac{\varepsilon(t)}{t} dt\right\}\right) \rightarrow \rho$$

by using power series expansions, noting that $n \int_{n-1}^n \varepsilon(t)/t dt = n[\varepsilon(\xi_n)/\xi_n]$ where $n-1 < \xi_n < n$ by the mean value theorem ($\varepsilon(t)$ being continuous).

COROLLARY. *If a sequence of positive terms $\{\theta(n)\}$, $n \geq 0$, is regularly varying with index ρ , then so is the positive function $\theta(x)$, $x \in [0, \infty)$, defined in terms of the sequence by $\theta(x) \equiv \theta([x])$, $x \geq 0$.*

PROOF. Let $R(x)$, $x \in [0, \infty)$, be a regularly varying function of index ρ , in which $\{\theta(n)\}$ is imbeddable. Then, for $x > 0$,

$$\begin{aligned} \theta(x) &= \theta([x]) = R([x]) = R(x - \delta_x), \quad \text{where } 0 \leq \delta_x < 1; \\ &= R(x(1 - (\delta_x/x))), \quad \sim R(x), \quad \text{as } x \rightarrow \infty, \end{aligned}$$

by the uniform convergence property of regularly varying functions.

REFERENCES

- J. Aczél, *Vorlesungen über Funktionalgleichungen und ihre Anwendungen*, Lehrbücher und Monographien aus dem Gebiete der exakten Wissenschaften, Mathematische Reihe, Band 25, Birkhäuser Verlag, Basel, 1961; English transl., *Math. in Sci. and Engineering*, vol. 19, Academic Press, New York, 1966. MR 23 #A1959; 34 #8020.
- R. Bojanić and E. Seneta, *Slowly varying functions and asymptotic relations*, *J. Math. Anal. Appl.* 34 (1971), 302–315. MR 43 #438.
- N. G. de Bruijne, *Pairs of slowly oscillating functions occurring in asymptotic problems concerning the Laplace transform*, *Nieuw. Arch. Wisk.* (3) 7 (1959), 20–26. MR 21 #5847.
- L. de Haan, *On regular variation and its application to the weak convergence of sample extremes*, *Mathematical Centre Tracts*, 32, Mathematisch Centrum, Amsterdam, 1970. MR 44 #3370.
- I. A. Ibragimov and Ju. V. Linnik, *Independent and stationarily connected variables*, Nauka, Moscow, 1965; English transl., Noordhoff, Groningen, 1971. MR 34 #2049.
- J. Karamata, *Sur un mode de croissance régulière des fonctions*, *Mathematica (Cluj)* 4 (1930), 38–53.
- , *Sur un mode de croissance régulière. Théorèmes fondamentaux*, *Bull. Soc. Math. France* 61 (1933), 55–62.

J. P. Kubilius, *Probabilistic methods in the theory of numbers*, Gos. Izdat. Polit. Naučn. Lit. Litovsk. SSR, Vilna, 1962; English transl., Transl. Math. Monographs, vol. 11, Amer. Math. Soc., Providence, R.I., 1964. MR 26 #3691; 28 #3956.

R. Schmidt, *Über divergente Folgen und lineare Mittelbildungen*, Math. Z. 22 (1925), 89–152.

I. Schur, *Zur theorie der Césaroschen und Holderschen Mittelwerte*, Math. Z. 31 (1930), 391–407.

DEPARTMENT OF MATHEMATICS, TEMPLE UNIVERSITY, PHILADELPHIA, PENNSYLVANIA 19122 (Current address of J. Galambos)

DEPARTMENT OF STATISTICS, PRINCETON UNIVERSITY, PRINCETON, NEW JERSEY 08540

Current address (E. Seneta): Statistics Department, Australian National University, Canberra, Australia