

RESEARCH ARTICLE


Regulation of the JMJD3 (KDM6B) histone demethylase in glioblastoma stem cells by STAT3

Maureen M. Sherry-Lynes[☯], Sejuti Sengupta[☯], Shreya Kulkarni, Brent H. Cochran*

Graduate Program in Cell and Molecular Physiology, Sackler School of Graduate Biomedical Sciences and Dept. of Developmental, Molecular, and Chemical Biology Tufts University School of Medicine Boston, MA, United States of America

☯ These authors contributed equally to this work.

* Brent.cochran@tufts.edu



OPEN ACCESS

Citation: Sherry-Lynes MM, Sengupta S, Kulkarni S, Cochran BH (2017) Regulation of the JMJD3 (KDM6B) histone demethylase in glioblastoma stem cells by STAT3. PLoS ONE 12(4): e0174775. <https://doi.org/10.1371/journal.pone.0174775>

Editor: Anita B. Hjelmeland, University of Alabama at Birmingham, UNITED STATES

Received: February 16, 2017

Accepted: March 15, 2017

Published: April 6, 2017

Copyright: © 2017 Sherry-Lynes et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are presented in the paper or in the Supporting information files.

Funding: This work was supported in part by NIH R01NS072414 to BHC. Details can be found by searching <https://projectreporter.nih.gov/reporter.cfm>. Additional funding was provided by Tufts University. There was no additional external funding received for this study. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Abstract

The growth factor and cytokine regulated transcription factor STAT3 is required for the self-renewal of several stem cell types including tumor stem cells from glioblastoma. Here we show that STAT3 inhibition leads to the upregulation of the histone H3K27me2/3 demethylase Jmjd3 (KDM6B), which can reverse polycomb complex-mediated repression of tissue specific genes. STAT3 binds to the Jmjd3 promoter, suggesting that Jmjd3 is a direct target of STAT3. Overexpression of Jmjd3 slows glioblastoma stem cell growth and neurosphere formation, whereas knockdown of Jmjd3 rescues the STAT3 inhibitor-induced neurosphere formation defect. Consistent with this observation, STAT3 inhibition leads to histone H3K27 demethylation of neural differentiation genes, such as Myt1, FGF21, and GDF15. These results demonstrate that the regulation of Jmjd3 by STAT3 maintains repression of differentiation specific genes and is therefore important for the maintenance of self-renewal of normal neural and glioblastoma stem cells.

Introduction

Glioblastoma stem cells (GBM-SC) are highly tumorigenic, and share many properties with normal neural stem cells [1, 2]. GBM-SC have gene expression patterns that more closely resemble their tumor of origin than do matched serum-derived cell lines [1]. STAT3 is a transcription factor that is activated by many cytokines and growth factors, and has a demonstrated role in oncogenesis of many human tumors including glioblastoma [3, 4]. STAT3 is required for the maintenance of pluripotency of murine embryonic and neural stem cells and facilitates reprogramming of somatic cells to the pluripotent state [5–7].

We and others have previously shown that the transcription factor STAT3 is essential for glioblastoma stem cell proliferation and multipotency [8–11]. Inhibition or RNAi knockdown of STAT3 leads to a dramatic decrease in proliferation and neurosphere formation, as well as loss of stem cell markers [11]. Interestingly, this phenotype is irreversible. Transient treatment with STAT3 inhibitors for as little as four hours leads to a permanent loss of neurosphere

Competing interests: The authors have declared that no competing interests exist.

formation capacity, despite the fact that STAT3 signaling is restored upon drug removal [11]. This observation suggests that STAT3 regulates the epigenetic state of the cells, thereby causing a stable change in the ability of the cells to respond to stem cell growth factors.

In stem cells, including normal neural stem cells and GBM-SC, polycomb-mediated repression of differentiation specific genes is a major mechanism by which multipotency is maintained [12]. PRC2 adds methyl groups to histone H3K27, which leads to the recruitment of the PRC1 and the heritable inhibition of transcription [13, 14]. The H3K27me2/3 histone demethylase Jmjd3 (KDM6B) antagonizes the enzymatic activity of the polycomb repressive complex 2 (PRC2) [15–18]. Jmjd3 demethylates histone H3K27 at the promoters of neuronal-specific genes in mice [19], and is required for neural differentiation of murine embryonic stem cells and for proper neural differentiation in adult mice [20, 21]. Jmjd3 expression can also be rapidly induced in macrophages in response to LPS stimulation [16].

It is becoming increasingly apparent that H3K27 trimethylation is aberrantly regulated in several cancers. Inactivating mutations have been identified in the H3K27 demethylase *UTX* (*KDM6A*), suggesting that *UTX* acts as a tumor suppressor [22]. Overexpression of *EZH2* has been demonstrated in multiple cancer types as well, including medulloblastoma and glioblastoma brain tumors [23–25]. *EZH2* expression is necessary for the self-renewal of glioblastoma stem cells [24]. Mutations in histone H3 itself have been found in pediatric glioblastoma [26–28]. Recently, it has been shown that Jmjd3 is induced during glioblastoma stem cell differentiation and is mutated in some GBM-SC and that constitutive Jmjd3 expression in GBM-SC inhibits their tumorigenesis [29]. These findings suggest that the aberrant maintenance of H3K27 methylation contributes to oncogenesis. Based on these observations and the STAT3 inhibition phenotype, we investigated the possibility that the H3K27 demethylase Jmjd3 may mediate effects of STAT3 in glioblastoma stem cells.

Materials and methods

Culture of glioblastoma stem cells and normal neural stem cells

Glioblastoma stem cell lines and culture conditions were described previously [11]. Normal human neural stem cells derived from H9 embryonic stem cells were obtained from Gibco (Carlsbad, CA). Cells were grown according to instructions of the manufacturer. For neurosphere formation assays, cells were dissociated from the coated plates with accutase and plated in neural stem cell media in uncoated plates. STAT3 inhibitors S3I-201, STA-21 (NCI/DTP Open Chemical Repository) or BP1-102 (EMD Millipore, Billerica, MA) were used at the indicated concentrations. Recombinant BMP4 (Life technologies, Grand Island, NY) was added to the culture media when indicated.

Immunoblotting

Immunoblots were performed as previously described [11] with the exception that immunoblots to be probed with anti-Jmjd3 antibody were transferred in 10 mM CAPS (3-[Cyclohexylamino]-1-propanesulfonic acid, pH 10.5) buffer. Anti- β -Actin (1:2000) was obtained from Sigma (St. Louis, MO); anti-STAT3 (1:1000), anti pSer-STAT3 (1:200), anti-pTyr STAT3 (1:500) were obtained from Cell Signaling Technologies (Beverly, MA). Anti-Jmjd3 was obtained from Abgent (San Diego, CA) or generously provided by the lab of Yang Shi.

RT-qPCR

RT-qPCR was performed as previously described [11]. Fold changes were calculated relative to control by the $2^{-\Delta\Delta C_t}$ method [30]. Statistical analysis was performed according to the

comparative C_T method. All primer sets amplify a single product of the predicted size; primer sequences can be found in supplemental experimental procedures. β -actin was used as the constitutive control gene.

Lentiviral shRNA infection and overexpression retrovirus infection

Cells were infected with either PLKO.1 non-targeting control lentivirus (Sigma) or shSTAT3 PLKO.1 lentivirus (clones TRCN0000020843 and TRCN0000020842), shJmjd3 (V2LHS_139678, Open Biosystems), shJmjd3 2 (pSico-R-PGKPuro lentivirus from [20]), in the presence of 8 μ g/mL polybrene. 24 hours post-infection the virus containing media was removed and replaced with fresh stem cell media for 24 hours. Cells were then treated with 2.5 μ g/mL puromycin. After 2–3 days of puromycin selection, cells were plated for assays as described in the figure legends.

The MSCV-Jmjd3 (Addgene plasmid 21212, [31]) MSCV-Jmjd3 mutant (Addgene plasmid 21214, [31]) and MSCV-Tap Control (Addgene plasmid 12570, [32]) retroviral plasmids were obtained from Addgene (Cambridge, MA). The viruses were packaged in 293GPG cells as described by Ory et al [33]. Glioma stem cells were infected at an approximate MOI of 1 overnight, and then overnight again. 96 hours after the first infection cells were treated with 2.5 μ g/mL puromycin for 4 days. After 4 days of puromycin selection cells were plated for assays as described in the figure legends.

ChIP and ChIP-sequencing

Cells were lysed, fixed, and sonicated according to the methods of De Santa et. al [34]. Sonicated lysates were incubated overnight with either anti- STAT3 (Santa Cruz Biotechnology, Santa Cruz, CA), anti-H3K27me3 (Millipore, Billerica, MA), or IgG control. Immunoprecipitation was carried out using StaphA cells according to the protocol of Kirmizis, Bartley [13] or using Protein A-Sepharose beads according to the nano ChIP-seq protocol (Adli and Bernstein [35]). After washing and cross-link reversal, immunoprecipitated DNA was purified using the Qiagen PCR Purification kit (Qiagen, Valencia, CA) and subjected to either qPCR or PCR as indicated in figure legends. For nano ChIP sequencing and nano ChIP-qPCR, the immunoprecipitated DNA was subjected to two intermediate rounds of amplification (Adli and Bernstein [35]) with custom primers containing barcodes that allowed us to multiplex 3 replicates for each of the two treatment groups in a sequencing lane (supplemental procedure). It was then subjected to either library amplification using TruSeq ChIP Sample preparation kit (Illumina Inc, San Diego, CA) followed by next generation sequencing on HiSeq 2500 (Illumina Inc, San Diego, CA) or qPCR. Myt1 primers were first described in [36]. FGF21 and GDF15 primers were designed to span regions of H3K27me3 enrichment near the gene promoter from ChIP-Sequencing data analysis. Primers used appear in supplemental experimental procedures. ChIP qPCR data was analyzed according to the percent of input method described by Haring et. al [37].

ChIP-sequencing data analysis

ChIP Sequencing reads were aligned to the human genome (hg19) using Bowtie in GALAXY platform and visualized in the UCSC genome browser [38]. The average read count through the 5kb upstream promoter region of each gene was computed across the genome. The normalized read counts at the promoters of STAT3 inhibited and control treatment groups was compared and a list of genes that showed reduction in H3K27me3 mark upon STAT3 inhibition was obtained [39, 40].

Microarray

Microarray gene expression data was generated using GeneChip Human gene U133A 2.0 Arrays (Affymetrix, Santa Clara, CA). The CEL files were processed by Transcriptome Analysis Console (Affymetrix) software to analyze differentially expressed genes based on fold change.

Results

STAT3 represses Jmjd3 expression in glioblastoma stem cells

We first examined Jmjd3 expression in response to STAT3 inhibition in two glioblastoma stem cell lines, GS6-22 and GS7-2, which we have previously characterized [11]. Upon STAT3 inhibition with the SH2-domain-targeting small molecule inhibitors STA-21 and S3I-201 [41, 42], Jmjd3 expression is upregulated at both the protein and mRNA levels by 2–15 fold (Fig 1A, S1A and S1B Fig). Jmjd3 mRNA is also upregulated upon STAT3 knockdown using two different shRNAs (Fig 1B, S1D and S1E Fig). Additionally, Jmjd3 mRNA is upregulated by a highly potent next-generation STAT3 inhibitor BP-1-102 [43] and S3I-201 treatment in additional GBM-SC lines tested (S1B Fig). Thus, multiple STAT3 specific inhibitors lead to upregulation of Jmjd3 expression in different GBM-SC lines.

By chromatin immunoprecipitation assay, STAT3 was found to bind to the Jmjd3 promoter at a position containing a conserved STAT3 binding site (Fig 1C, S1C Fig), suggesting that Jmjd3 is a direct STAT3 target gene. Previously it has been shown that Jmjd3 is upregulated during murine neural stem cell differentiation, and that it targets several neural specific genes upon differentiation of neural stem cells [19, 20]. Consistent with this we have found that Jmjd3 is upregulated during GBM-SC differentiation as well (Fig 1D, S1F Fig). UTX, another member of the Jmjd3 family of H3K27me3 demethylases, is not consistently upregulated by the two STAT3 inhibitors, S3I-201 and STA-21, in both of our cell lines (S1G Fig).

Interestingly, STAT3 appears to repress Jmjd3 expression, rather than activate it as is typical for STAT3 target genes [44]. However, STAT3 has been shown to repress genes in other cell types, and is associated with both transcriptionally active and repressed genes in embryonic stem cells [45] and in neural stem cells [46]. In murine neural stem cells, Jmjd3 is repressed by the NCoR and SMRT nuclear co-repressors, which associate with the retinoic acid receptor to repress transcription [19]. STAT3 has been shown to associate with the retinoic acid receptor (RAR), so we considered the possibility that NCoR repression is involved in regulating Jmjd3 in glioblastoma stem cells [47, 48]. Retinoic acid treatment of GS6-22 cells leads to a rapid increase in Jmjd3 mRNA (Fig 1D). Additionally, the NCoR co-repressor can be found at the Jmjd3 promoter by chromatin immunoprecipitation (Fig 1E and 1F). NCoR can be immunoprecipitated using the same primers as STAT3, suggesting it binds a similar region of the Jmjd3 promoter. Furthermore, treatment with the STAT3 inhibitor S3I-201 decreases NCoR binding to the Jmjd3 promoter (Fig 1E and 1F). It is possible, then, that STAT3 represses Jmjd3 expression by interaction with the NCoR-RAR complex, and that treatment with S3I-201 abolishes this interaction, thereby relieving the repression of Jmjd3.

STAT3 regulates GBM-SC neurosphere formation and proliferation through repression of Jmjd3

Neurosphere formation is a hallmark of neural stem cells, and the ability to culture neurospheres from glioblastomas strongly correlates with poor patient prognosis [49]. Because STAT3 represses Jmjd3 expression in GBM-SC, we investigated whether knockdown of Jmjd3 expression could rescue the abrogation of neurosphere formation that accompanies STAT3




Fig 1. STAT3 represses Jmjd3 expression in glioblastoma stem cells. A. RT-qPCR of GS6-22 cells treated with S3I-201 or STA-21 demonstrates that Jmjd3 mRNA is upregulated at both 4 hours and 24 hours after inhibitor treatment. Values represent the fold change relative to DMSO treated cells for three experiments; bars SD (**p<0.01). B. Knockdown of STAT3 using an shRNA-containing lentivirus leads to the upregulation of Jmjd3 mRNA in GS6-22 cells, two days after selection. Values represent the fold change relative to DMSO treated cells for three experiments; bars SD (**p<0.01). C. Chromatin immunoprecipitation using anti-STAT3 antibody, followed by PCR using primers to the Jmjd3 promoter demonstrates STAT3 binding in GS7-2 and GS6-22 cells. Enrichment of DNA in STAT3 immunoprecipitation (relative to input) was quantified by ImageJ. D. Treatment of GS6-22 cells with retinoic acid increases Jmjd3 mRNA levels. GS6-22 cells were treated for 1 and 4 hours with 1 μM RA. Values represent the fold change relative to DMSO treated cells for three experiments; bars SD (*p<0.05, **p<0.01). E. S3I-201 treatment decreases NCOR binding at the Jmjd3 promoter. GS6-22 cells were treated with S3I-201 (50 μM) or DMSO for 4 hours, fixed, lysed, and sonicated. Sonicated lysates were incubated overnight with anti-NCOR or IgG control and subjected to chromatin immunoprecipitation. The resulting DNA was subjected to PCR using primers to the Jmjd3 promoter. F. Densitometry of E was performed using ImageJ.

<https://doi.org/10.1371/journal.pone.0174775.g001>

inhibition. GS6-22 and GS7-2 cells were infected with lentiviruses expressing shRNA to Jmjd3 or with control lentiviruses. Control infected cells exhibited decreased neurosphere formation in response to S3I-201 treatment as expected (Fig 2A, 2B and 2C). shJmjd3 infected cells, however, were able to form spheres in the presence of S3I-201 (Fig 2A, 2B and 2C). This was also true for cells infected with a second, distinct shRNA to Jmjd3, indicating that this is a specific effect of Jmjd3 knockdown (S2A and S2B Fig). This effect was more pronounced in the GS7-2 cells (Fig 2C) than the GS6-22 cells (Fig 2B) likely due to the enhanced degree of knockdown in the GS7-2 cells (78% knockdown in the GS7-2 cells versus 55% for GS6-22 cells) (S2C and S2D Fig). These data indicate that STAT3 repression of Jmjd3 is necessary for neurosphere formation.




Fig 2. STAT3 controls GBM-SC neurosphere formation and proliferation through repression of Jmjd3. A. Knockdown of Jmjd3 rescues neurosphere formation in the presence of S3I-201. Representative images of GS6-22 and GS7-2 cells infected with either control or shJmjd3 containing lentivirus treated with DMSO or S3I-201 (50 μ M). Images for GS6-22 cells were taken after 6 days and GS7-2 cells were taken after 4 days, based on differences in sphere formation rate in these lines. B. Quantification of neurosphere formation in GS6-22 cells. After 6 days of inhibitor treatment, the number of spheres per 100 cells was counted. Values represent the mean of triplicates within each treatment; bars SE (** p <0.01). C. After 4 days of inhibitor treatment, GS7-2 cell sphere formation was quantified as described for GS6-22 cells. Values represent the mean of triplicates within each treatment; bars SE (** p <0.01 relative to DMSO control, upon shJMJD3 infection). D. Representative images of GS6-22 and GS7-2 cells infected with overexpression retrovirus containing either a control (empty) plasmid, Jmjd3, or a catalytic domain mutant of Jmjd3. After three days of selection, cells were dissociated and replated at 100 cells per ml. Pictures were taken at 50X seven days after replating. E. Quantification of neurosphere formation capacity in GS6-22 and GS7-2 cells infected with the Jmjd3 overexpression retrovirus. Neurosphere formation assay was performed in triplicate (** p <0.01). F. GS6-22 and G. GS7-2 cells were infected with either control, Jmjd3, or Jmjd3 mutant retroviruses as previously described. Cells were pulsed with 30 μ M BrdU for 16 hours. After 24 hours, cells were fixed and stained with an anti-BrdU antibody. Cells were also stained with 7-AAD at this time. The percentage of BrdU positive cells was analyzed using flow cytometry. Values represent the mean of 3 experiments; bars SD of the mean (* p <0.05).

<https://doi.org/10.1371/journal.pone.0174775.g002>

Consistent with this, overexpression of *Jmjd3* recapitulates significant aspects of the STAT3 inhibition phenotype. Both GS6-22 and GS7-2 cells that had been engineered to constitutively express *Jmjd3* fail to form neurospheres in stem cell media (Fig 2D and 2E). Overexpression of a catalytic dead mutant of *Jmjd3*, however, does not inhibit sphere formation (Fig 2D and 2E). In addition, overexpression of *Jmjd3*, but not mutant *Jmjd3*, decreased proliferation in both GS6-22 and GS7-2 cells (Fig 2F and 2G). This effect was more pronounced in GS7-2 cells, but the degree of overexpression was also higher in these cells (S2E and S2F Fig). This indicates that the demethylase activity of *Jmjd3* is necessary for the inhibition of sphere formation and proliferation of GBM-SC. STAT3 repression of *Jmjd3*, then, is required for GBM-SC neurosphere formation and proliferation.

Interestingly, overexpression of both wildtype *Jmjd3* and the mutant increased apoptosis in GS6-22 cells (S2G Fig). It is possible the mutant *Jmjd3* may interfere with some cellular functions by substrate binding and sequestration. Consistent with this, *Jmjd3* has been found to have demethylase independent functions[50]. This effect was not observed in GS7-2 cells (S2H Fig). This may be because these cells lack expression of the *INK4A/ARF* locus (see below).

STAT3 regulates H3K27 trimethylation and neural gene induction in GBM-SC

To identify putative STAT3 regulated genes that show induction of gene expression concurrent with histone H3K27 demethylation in GS6-22, we performed microarrays and genome wide CHIP-sequencing analysis using an antibody specific to H3K27me3, following STAT3 inhibition. Differential analysis of STAT3 inhibited and control GS6-22 cells yielded a ranked list of genes that were likely to show reduction in H3K27me3 mark (S1 Table) or are induced upon STAT3 inhibition with a fold change >2 (S2 Table). The intersected list of genes showing both reduction in repressive H3K27me3 mark as well as 2 fold or higher gene expression upon S3i treatment is tabulated in S3 Table.

We subsequently confirmed that STAT3 inhibition by the next-generation STAT3 inhibitor BP-1-102 [43] upregulates mRNA expression of Fibroblast Growth Factor 21 (FGF21) and Growth Differentiation Factor 15 (GDF15) in GS6-22 and GS7-2 cells (Fig 3A, S3A Fig). These genes are known to play roles in neural development [51, 52]. GDF15 has also been shown to induce apoptosis and inhibit tumorigenesis in GBM [53, 54].

Consistent with the induction of gene expression, ChIP-sequencing data analysis suggested loss of H3K27me3 peaks at FGF21 and GDF15 promoters in STAT3 inhibited GS6-22 cells compared to control. This was confirmed using ChIP-qPCR analysis, which showed that STAT3 inhibition by BP-1-102 decreases H3K27 trimethylation at FGF21 and GDF15 promoters (Fig 3B), concurrent with JMJD3 upregulation in GS6-22 cells (Fig 3A). The amount of total histone H3 does not change upon BP-1-102 treatment indicating that there is a loss of H3K27 trimethylation at these promoters and not just a loss of histone H3 association (S3B and S3C Fig). These results indicate that inhibition of STAT3 induces H3K27 demethylation and subsequent expression of target genes involved in neural differentiation of GBM-SC.

Demethylation of H3K27 at the *Myt1* promoter confers BMP4 inducibility

Because *Jmjd3* inhibits sphere formation and proliferation of GBM-SC in a demethylase-dependent manner, we examined the H3K27me3 status at the promoter of a known *Jmjd3* and polycomb target gene, myelin transcription factor 1, *Myt1* [13], which is a gene that has been implicated in neuronal differentiation [55]. GS6-22 cells were treated with STAT3 inhibitor, BP-1-102, and subjected to chromatin immunoprecipitation (ChIP) using an antibody specific to H3K27me3. In GS6-22 cells, a decrease in H3K27 trimethylation was observed at the *Myt1*




Fig 3. STAT3 regulates H3K27 trimethylation and neural gene induction in GBM-SC. A. RT-qPCR analysis of FGF21, GDF15 and JMJD3 expression in GS6-22 cells treated with BP-1-102 (15uM) for 6 hours. Values represent the fold change relative to control cells for three experiments; bars SD (**p<0.01, ***p<0.005). B. GS6-22 cells were treated with BP-1-102 (15 μM) or DMSO for 6 hours, fixed, lysed, and subjected to chromatin immunoprecipitation with an antibody to H3K27 trimethylation or total histone H3. Quantitative PCR using primers to FGF21 or GDF15 or Myt1 genes was performed after immunoprecipitation, amplification and DNA purification. Data is displayed as percentage of input DNA of H3K27me3/H3 ChIP, normalized relative to control

cells for three experiments (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.005$). C. Demethylation of H3K27 at the Myt1 promoter confers BMP4 inducibility of the gene. RT-qPCR analysis of Myt1 and JMJD3 expression in GS6-22 cells treated with BP-1-102 (15 μ M), recombinant BMP4 (10 ng/ml) or a combination of both for 6 hours. Values represent the fold change relative to DMSO treated control cells for three experiments; bars SD (* $p < 0.05$, ** $p < 0.01$).

<https://doi.org/10.1371/journal.pone.0174775.g003>

promoter (Fig 3B, S3D Fig), concurrent with Jmjd3 upregulation. The degree of decrease in H3K27 trimethylation is consistent with that seen at other promoters demethylated by Jmjd3 [19, 56] and suggests that Myt1 is a Jmjd3 target gene in GBM-SC.

However, STAT3 inhibition or expression of Jmjd3 in stem cell growth conditions was not sufficient to induce Myt1 gene expression (Fig 3C, S4B Fig), suggesting that other factors or signals are required to activate expression of these neural specific genes. It has been previously shown that Bone Morphogenetic Protein 4 (BMP4) activates the Smad signaling cascade and induces differentiation of stem-like, tumor-initiating precursors of GBMs [57] and that SMADs and Jmjd3 co-regulate genes in NSCs [58]. Myt1 expression is strongly induced upon BMP4 treatment when STAT3 is inhibited in GS6-22 and GS7-2 cells (Fig 3C, S4A Fig). This suggests that STAT3 inhibition and subsequent demethylation of histone H3K27 at the promoter renders the chromatin accessible to BMP4 induced transcription factors at this locus.

Interestingly, we have not found a change in H3K27me3 at the *INK4A/Arf* locus (S4C Fig) in GS6-22 cells. This locus is a well-characterized Jmjd3 target [15, 56, 59], and its demethylation is necessary for the induction of growth arrest and senescence in several cell types. In GS7-2 cells, we failed to see PCR amplification of *INK4A/ARF* in genomic DNA, which suggests that this locus is deleted in GS7-2 cells (S4D Fig). This is not surprising given that over half of glioblastomas exhibit homozygous deletion of this locus [60]. Together, these observations suggest *INK4A/ARF* is not necessary for Jmjd3 regulation of GBM-SC proliferation and sphere formation. Ene et al (2012) came to a similar conclusion [29].

STAT3 regulates Jmjd3 expression in human neural stem cells

Finally, we examined whether STAT3 repression of Jmjd3 was specific to glioblastoma stem cells, or whether the STAT3 inhibition phenotype is recapitulated in normal human neural stem cells. In neural stem cells derived from H9 embryonic stem cells [61, 62], STAT3 is activated by phosphorylation on both pTyr705 and pSer727 (Fig 4A). S3I-201 treatment of these cells inhibited neurosphere formation (Fig 4B) and lead to upregulation of Jmjd3 mRNA (Fig 4C), as well as to dose-dependent inhibition of proliferation as judged by BrdU incorporation (Fig 4D). STAT3, then, regulates neurosphere formation and proliferation in normal embryonic neural stem cells as well as in glioblastoma stem cells. STAT3 repression of Jmjd3 is also maintained in normal neural stem cells. This indicates that STAT3 control of the epigenetic program through Jmjd3 repression is a key mechanism of stem cell maintenance in both normal and tumorigenic neural stem cells.

Discussion

We have shown that STAT3 maintains normal neural and glioblastoma stem cells in a proliferative, self-renewing state via the repression of the histone demethylase Jmjd3. STAT3 repression of Jmjd3 is consistent with our previous data that STAT3 inhibition is irreversible in GBM-SC. We have found that STAT3 inhibition or knockdown leads to rapid upregulation of Jmjd3. We have also found that STAT3 binds to the Jmjd3 promoter in human GBM-SC which is consistent with published genomic data from murine ES cells [45] and suggests that STAT3 is a direct regulator of JMJD3.




Fig 4. STAT3 inhibition in human neural stem cells leads to a decrease in neurosphere formation and proliferation, and an upregulation of Jmjd3 expression. A. STAT3 activation in human neural stem cells was assessed by immunoblotting with phospho-specific antibodies. B. Human neural stem cells were plated in serum free media supplemented with EGF and FGF2. 24 hours after plating, cells were treated with S3I-201 (50 μ M) or DMSO control. Pictures were taken 3 days after treatment. C. STAT3 regulation of Jmjd3 was assessed by qRT-PCR. Human neural stem cells were treated with S3I-201 (50 μ M) for 24 hours. Values represent the fold change relative to control cells for three experiments; bars SD (** p <0.01). D. Cells were treated with DMSO or S3I-201 for 24 hours, and then subjected to a 15 μ M BrdU pulse for 20 hours. Values represent the average of three experiments; bars SE (* p <0.05, ** p <0.01).

<https://doi.org/10.1371/journal.pone.0174775.g004>

While STAT3 is best known as a transcriptional activator, there is evidence that it can act as a repressor [63, 64]. ChIP-ChIP and microarray studies suggest that STAT3 represses genes in multiple cell types including in GBM-SC [45, 46, 65, 66]. Previous studies indicated that NCoR negatively regulates Jmjd3 [19]. We have found that following STAT3 inhibition NCoR binding is reduced at the same promoter fragment that binds STAT3, suggesting that the mechanism of repression is recruitment or stabilization of the NCoR repressor complex at the promoter. While Ene et al (2012)[29] have found evidence that DNA methylation can also contribute to Jmjd3 repression, the sites of that STAT3 binds to appear to be distinct from the regions of DNA methylation [29]. However, given recent evidence that STAT3 can control DNA methylation [67], this mechanism of regulation merits further investigation.

We have also found that Jmjd3 overexpression inhibits neurosphere formation and cell proliferation (Figs 2 and 3). Consistent with our data, Jmjd3 overexpression has been shown to decrease the tumorigenicity of GBM-SC *in vivo* [29]. By contrast, another study reported that GSK-J4 mediated inhibition of JMJD3 and UTX was shown to reduce the sphere forming capacity of GBM-SC [68]. However, GSK-J4 has nonspecific inhibitory effects on other Jumonji domain family members[69], which need to be assessed to resolve such context-dependent effects. Moreover, targeted knockout of JMJD3 in the mouse brain indicates that expression of JMJD3 and its demethylase activity is needed for adult brain neurogenesis consistent with our results here[21].

We have found that overexpression of Jmjd3, which opposes PRC2 activity, also induces apoptosis in GS6-22 cells. A similar effect on both proliferation and apoptosis of GBM-SC has been reported for knockdown of BMI-1, part of the PRC1 complex [70]. Similarly, inhibition of EZH2, the enzymatic part of the PRC2 complex, causes a decrease in sphere formation and proliferation of GBM-SC [24]. Interestingly, EZH2 also methylates and regulates STAT3 in

GBM-SC which further suggests a tight coordination between STAT3 and H3K27 trimethylation [71].

In this regard, it is also worth noting the role of histone H3K27 trimethylation in brain tumors is complex. While induction and maintenance of this mark is pro-oncogenic in medulloblastoma and adult glioblastomas [23–25, 71], global loss of H3K27 methylation is oncogenic in pediatric gliomas and in some adult GBMs [26, 27, 72]. Thus, the effects of H3K27 trimethylation on oncogenesis are context dependent. This is consistent with findings from other cancers as well [73, 74].

We have also found that a catalytic mutant of *Jmjd3* does not affect neurosphere formation or proliferation. Thus, *Jmjd3* upregulation and the subsequent demethylation of histone H3K27 on target genes is likely necessary for the decrease in GBM-SC growth. This induction of *Jmjd3* likely imparts widespread changes to the chromatin and sustained mRNA expression of target genes involved in neural differentiation of GBM-SC. This would explain why either the restoration of STAT3 signaling following removal of S3I-201 STAT3 inhibitor [11], or sustained STAT3 signaling when *Jmjd3* is constitutively expressed, does not rescue neurosphere formation. This difference in epigenetic state may also explain how STAT3 can induce astrocytic differentiation of neural stem cells under some conditions and preserve neural stem cell self-renewal in others [6, 75–77]. A similar persistence of a stable STAT3 phenotype has been demonstrated in breast epithelial cells, in which a transient expression of either of two STAT3-regulated microRNAs is sufficient to induce and maintain a stably transformed state that is also polycomb mediated [78, 79]. Thus, there is precedence for the transient upregulation of a single STAT3 target gene imparting permanent change to the cellular state.

Interestingly, although STAT3 inhibition induces neural genes, such as FGF21 and GDF15, neither *Jmjd3* overexpression or STAT3 inhibition are sufficient to drive differentiation of the glioblastoma stem cells (S4B Fig, [11]). While *Jmjd3* overexpression is sufficient to induce differentiation in some systems [19, 31], it is likely that additional positive differentiation signals are required both in these cases and in the case of the GBM-SC. Histone H3K27 demethylation may be sufficient to open the chromatin at specific genes, but positively acting transcription factors are likely needed in addition to bind to these promoters to activate transcription. Consistent with this, we have found that *Jmjd3* overexpression can demethylate histone H3K27 at certain differentiation-specific genes such as *Myt1*, but the gene is upregulated only in the presence of recombinant BMP4. This situation is similar to previous results for the GFAP gene where epigenetic changes must occur before factors like CNTF are able to induce the gene during astrocytic differentiation [80–82]. Interestingly, STAT3 regulates GFAP as well [80–82].

In addition, *Jmjd3* overexpression alone does not completely recapitulate the STAT3 inhibition phenotype, at least with respect to β III-tubulin expression (Fig 3D, S4B Fig, and [11]). Given that STAT3 signaling likely regulates many genes in addition to *Jmjd3* in GBM-SC, this is not surprising. However, the fact that constitutive *Jmjd3* expression substantially inhibits neurosphere formation as well as proliferation indicates that *Jmjd3* repression by STAT3 is required for glioblastoma stem cell maintenance in the cell lines tested. As STAT3 regulation of *Jmjd3* was found in GBM-SC derived from multiple patient tumors, it is likely that this finding indicates a broad role for STAT3/*Jmjd3* regulation of GBM-SC maintenance.

Strikingly, we have also found that STAT3 repression of *Jmjd3* is a mechanism shared by both glioma stem cells and normal neural stem cells (Fig 4). While this could complicate therapeutic targeting of STAT3 in glioblastoma, it does provide further evidence that similar signaling and epigenetic mechanisms that govern both cancer stem cells and tissue stem cells at least in this case. STAT3 is necessary for the self-renewal of a variety of stem cell types, including murine embryonic stem cells [5], and its regulation of the epigenetic program may prove to be a widespread and developmentally critical mechanism of maintaining stem cell self-renewal.

Supporting information

S1 Fig. STAT3 represses JMJD3 expression in glioblastoma stem cells.

(PDF)

S2 Fig. JMJD3 knockdown partially rescues neurosphere formation in S3I-201.

(PDF)

S3 Fig. Neural gene induction and H3K27 demethylation in BP-1-102-treated GBM-SC.

(PDF)

S4 Fig. MYT1 expression and INK4A/ARF H3K27 methylation in GBM-SC.

(PDF)

S1 Table. GS6-22 cells were treated with S3I-201 (50uM) or DMSO control for 3 days. ChIP Sequencing analysis using H3K27me3 antibody yielded a list of gene promoters that likely show a reduction in repressive H3K27me3 mark upon S3i treatment.

(XLSX)

S2 Table. GS6-22 cells were treated with S3I-201 (50uM) or DMSO control for 3 days. Microarray analysis yielded a list of genes that likely show 2 fold or higher gene expression upon S3i treatment.

(XLSX)

S3 Table. GS6-22 cells were treated with S3I-201 (50uM) or DMSO control for 3 days. Intersected list of genes showing both reduction in repressive H3K27me3 mark as well as 2 fold or higher gene expression upon S3i treatment.

(XLSX)

S1 File. Supplemental Experimental Procedures.

(PDF)

Acknowledgments

We thank G. Testa for shRNA to Jmjd3, Y. Shi for anti-Jmjd3 antibodies, P. Khavari for the Jmjd3 retroviral vectors obtained from Addgene, and H. Wakimoto for the MGG8 cell line. S3I-201 (NSC 74859) and STA-21 (NSC 628869) inhibitors were obtained from the NCI/DTP Open Chemical Repository (<http://dtp.cancer.gov>). The Tufts Neuroscience Core (P30 NS047243) provided the Stratagene qPCR machine. We thank J. Wu and A. Yee for critical reading of the manuscript. This work was supported in part by NIH R01NS072414 to BHC.

Author Contributions

Conceptualization: MS SS BHC.

Data curation: MS SS SK BHC.

Formal analysis: MS SS SK BHC.

Funding acquisition: BHC.

Investigation: MS SS SK.

Methodology: MS SS BHC.

Project administration: BHC.

Resources: MS SS SK.

Supervision: BHC.

Validation: MS SS SK BHC.

Visualization: MS SS BHC.

Writing – original draft: MS SS BHC.

Writing – review & editing: MS SS SK BHC.

References

1. Lee J, Kotliarova S, Kotliarov Y, Li A, Su Q, Donin NM, et al. Tumor stem cells derived from glioblastomas cultured in bFGF and EGF more closely mirror the phenotype and genotype of primary tumors than do serum-cultured cell lines. *Cancer Cell*. 2006; 9(5):391–403. <https://doi.org/10.1016/j.ccr.2006.03.030> PMID: 16697959
2. Vescovi AL, Galli R, Reynolds BA. Brain tumour stem cells. *Nature reviews*. 2006; 6(6):425–36. <https://doi.org/10.1038/nrc1889> PMID: 16723989
3. Yu H, Jove R. The STATs of cancer—new molecular targets come of age. *Nature reviews*. 2004; 4(2):97–105. <https://doi.org/10.1038/nrc1275> PMID: 14964307
4. Darnell JE. Validating Stat3 in cancer therapy. *Nat Med*. 2005; 11(6):595–6. <https://doi.org/10.1038/nm0605-595> PMID: 15937466
5. Niwa H, Burdon T, Chambers I, Smith A. Self-renewal of pluripotent embryonic stem cells is mediated via activation of STAT3. *Genes & development*. 1998; 12(13):2048–60.
6. Rajan P, McKay RD. Multiple routes to astrocytic differentiation in the CNS. *J Neurosci*. 1998; 18(10):3620–9. PMID: 9570793
7. Yang J, van Oosten AL, Theunissen TW, Guo G, Silva JC, Smith A. Stat3 activation is limiting for reprogramming to ground state pluripotency. *Cell stem cell*. 2010; 7(3):319–28. <https://doi.org/10.1016/j.stem.2010.06.022> PMID: 20804969
8. Li GH, Wei H, Lv SQ, Ji H, Wang DL. Knockdown of STAT3 expression by RNAi suppresses growth and induces apoptosis and differentiation in glioblastoma stem cells. *Int J Oncol*. 2010; 37(1):103–10. Epub 2010/06/02. PMID: 20514402
9. Carro MS, Lim WK, Alvarez MJ, Bollo RJ, Zhao X, Snyder EY, et al. The transcriptional network for mesenchymal transformation of brain tumours. *Nature*. 2010; 463(7279):318–25. Epub 2009/12/25. <https://doi.org/10.1038/nature08712> PMID: 20032975
10. Wang H, Lathia JD, Wu Q, Wang J, Li Z, Heddleston JM, et al. Targeting Interleukin 6 Signaling Suppresses Glioma Stem Cell Survival and Tumor Growth. *Stem Cells*. 2009; 27(10):2393–404. Epub 2009/08/07. <https://doi.org/10.1002/stem.188> PMID: 19658188
11. Sherry MM, Reeves A, Wu JK, Cochran BH. STAT3 is required for proliferation and maintenance of multipotency in glioblastoma stem cells. *Stem Cells*. 2009; 27(10):2383–92. Epub 2009/08/07. <https://doi.org/10.1002/stem.185> PMID: 19658181
12. Bracken AP, Helin K. Polycomb group proteins: navigators of lineage pathways led astray in cancer. *Nature reviews*. 2009; 9(11):773–84. <https://doi.org/10.1038/nrc2736> PMID: 19851313
13. Kirmizis A, Bartley SM, Kuzmichev A, Margueron R, Reinberg D, Green R, et al. Silencing of human polycomb target genes is associated with methylation of histone H3 Lys 27. *Genes & development*. 2004; 18(13):1592–605. Epub 2004/07/03. PubMed Central PMCID: PMC443521.
14. Kuzmichev A, Nishioka K, Erdjument-Bromage H, Tempst P, Reinberg D. Histone methyltransferase activity associated with a human multiprotein complex containing the Enhancer of Zeste protein. *Genes & development*. 2002; 16(22):2893–905.
15. Agger K, Cloos PA, Christensen J, Pasini D, Rose S, Rappsilber J, et al. UTX and JMJD3 are histone H3K27 demethylases involved in HOX gene regulation and development. *Nature*. 2007; 449(7163):731–4. Epub 2007/08/24. <https://doi.org/10.1038/nature06145> PMID: 17713478
16. De Santa F, Totaro MG, Prosperini E, Notarbartolo S, Testa G, Natoli G. The histone H3 lysine-27 demethylase Jmjd3 links inflammation to inhibition of polycomb-mediated gene silencing. *Cell*. 2007; 130(6):1083–94. <https://doi.org/10.1016/j.cell.2007.08.019> PMID: 17825402
17. Lan F, Bayliss PE, Rinn JL, Whetstone JR, Wang JK, Chen S, et al. A histone H3 lysine 27 demethylase regulates animal posterior development. *Nature*. 2007; 449(7163):689–94. Epub 2007/09/14. <https://doi.org/10.1038/nature06192> PMID: 17851529

18. Xiang Y, Zhu Z, Han G, Lin H, Xu L, Chen CD. JMJD3 is a histone H3K27 demethylase. *Cell research*. 2007; 17(10):850–7. <https://doi.org/10.1038/cr.2007.83> PMID: 17923864
19. Jepsen K, Solum D, Zhou T, McEvelly RJ, Kim HJ, Glass CK, et al. SMRT-mediated repression of an H3K27 demethylase in progression from neural stem cell to neuron. *Nature*. 2007; 450(7168):415–9. <https://doi.org/10.1038/nature06270> PMID: 17928865
20. Burgold T, Spreafico F, De Santa F, Totaro MG, Prosperini E, Natoli G, et al. The histone H3 lysine 27-specific demethylase Jmjd3 is required for neural commitment. *PLoS ONE*. 2008; 3(8):e3034. Epub 2008/08/22. PubMed Central PMCID: PMC2515638. <https://doi.org/10.1371/journal.pone.0003034> PMID: 18716661
21. Park DH, Hong SJ, Salinas RD, Liu SJ, Sun SW, Sgualdino J, et al. Activation of neuronal gene expression by the JMJD3 demethylase is required for postnatal and adult brain neurogenesis. *Cell reports*. 2014; 8(5):1290–9. PubMed Central PMCID: PMC4201382. <https://doi.org/10.1016/j.celrep.2014.07.060> PMID: 25176653
22. van Haafden G, Dalglish GL, Davies H, Chen L, Bignell G, Greenman C, et al. Somatic mutations of the histone H3K27 demethylase gene UTX in human cancer. *Nature genetics*. 2009; 41(5):521–3. <https://doi.org/10.1038/ng.349> PMID: 19330029
23. Robinson G, Parker M, Kranenburg TA, Lu C, Chen X, Ding L, et al. Novel mutations target distinct subgroups of medulloblastoma. *Nature*. 2012; 488(7409):43–8. <https://doi.org/10.1038/nature11213> PMID: 22722829
24. Suva ML, Riggi N, Janiszewska M, Radovanovic I, Provero P, Stehle JC, et al. EZH2 is essential for glioblastoma cancer stem cell maintenance. *Cancer Res*. 2009; 69(24):9211–8. Epub 2009/11/26. <https://doi.org/10.1158/0008-5472.CAN-09-1622> PMID: 19934320
25. Zhang J, Chen L, Han L, Shi Z, Zhang J, Pu P, et al. EZH2 is a negative prognostic factor and exhibits pro-oncogenic activity in glioblastoma. *Cancer Lett*. 2015; 356(2 Pt B):929–36. Epub 2014/12/03.
26. Venneti S, Garimella MT, Sullivan LM, Martinez D, Huse JT, Heguy A, et al. Evaluation of Histone 3 Lysine 27 Trimethylation (H3K27me3) and Enhancer of Zest 2 (EZH2) in Pediatric Glial and Glioneuronal Tumors Shows Decreased H3K27me3 in H3F3A K27M Mutant Glioblastomas. *Brain pathology (Zurich, Switzerland)*. 2013.
27. Bender S, Tang Y, Lindroth AM, Hovestadt V, Jones DT, Kool M, et al. Reduced H3K27me3 and DNA hypomethylation are major drivers of gene expression in K27M mutant pediatric high-grade gliomas. *Cancer Cell*. 2013; 24(5):660–72. <https://doi.org/10.1016/j.ccr.2013.10.006> PMID: 24183680
28. Wu G, Broniscer A, McEachron TA, Lu C, Paugh BS, Becksfors J, et al. Somatic histone H3 alterations in pediatric diffuse intrinsic pontine gliomas and non-brainstem glioblastomas. *Nature genetics*. 2012; 44(3):251–3. Epub 2012/01/31. PubMed Central PMCID: PMC3288377. <https://doi.org/10.1038/ng.1102> PMID: 22286216
29. Ene CI, Edwards L, Riddick G, Baysan M, Woolard K, Kotliarova S, et al. Histone demethylase Jumonji D3 (JMJD3) as a tumor suppressor by regulating p53 protein nuclear stabilization. *PLoS One*. 2012; 7(12):e51407. Epub 2012/12/14. PubMed Central PMCID: PMC3517524. <https://doi.org/10.1371/journal.pone.0051407> PMID: 23236496
30. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods (San Diego, Calif)*. 2001; 25(4):402–8.
31. Sen GL, Webster DE, Barragan DI, Chang HY, Khavari PA. Control of differentiation in a self-renewing mammalian tissue by the histone demethylase JMJD3. *Genes & development*. 2008; 22(14):1865–70.
32. Frias MA, Thoreen CC, Jaffe JD, Schroder W, Sculley T, Carr SA, et al. mSin1 is necessary for Akt/PKB phosphorylation, and its isoforms define three distinct mTORC2s. *Curr Biol*. 2006; 16(18):1865–70. <https://doi.org/10.1016/j.cub.2006.08.001> PMID: 16919458
33. Ory DS, Neugeboren BA, Mulligan RC. A stable human-derived packaging cell line for production of high titer retrovirus/vesicular stomatitis virus G pseudotypes. *Proc Natl Acad Sci U S A*. 1996; 93(21):11400–6. PMID: 8876147
34. De Santa F, Totaro M, Prosperini E, Notarbartolo S, Testa G, Natoli G. The histone H3 lysine-27 demethylase Jmjd3 links inflammation to inhibition of polycomb-mediated gene silencing. *Cell*. 2007; 130(6):1083–94. <https://doi.org/10.1016/j.cell.2007.08.019> PMID: 17825402
35. Adli M, Bernstein BE. Whole-genome chromatin profiling from limited numbers of cells using nano-ChIP-seq. *Nature protocols*. 2011; 6(10):1656–68. <https://doi.org/10.1038/nprot.2011.402> PMID: 21959244
36. Sarma K, Margueron R, Ivanov A, Pirrotta V, Reinberg D. Ezh2 requires PHF1 to efficiently catalyze H3 lysine 27 trimethylation in vivo. *Molecular and cellular biology*. 2008; 28(8):2718–31. <https://doi.org/10.1128/MCB.02017-07> PMID: 18285464

37. Haring M, Offermann S, Danker T, Horst I, Peterhansel C, Stam M. Chromatin immunoprecipitation: optimization, quantitative analysis and data normalization. *Plant methods*. 2007; 3:11. <https://doi.org/10.1186/1746-4811-3-11> PMID: 17892552
38. Karolchik D, Hinrichs AS, Kent WJ. The UCSC Genome Browser. *Current protocols in bioinformatics / editorial board, Andreas D Baxeavanis [et al]*. 2009;Chapter 1:Unit 1 4. PubMed Central PMCID: PMC2834533.
39. Hebenstreit D, Gu M, Haider S, Turner DJ, Lio P, Teichmann SA. EpiChIP: gene-by-gene quantification of epigenetic modification levels. *Nucleic Acids Res*. 2011; 39(5):e27. Epub 2010/12/07. PubMed Central PMCID: PMC3061070. <https://doi.org/10.1093/nar/gkq1226> PMID: 21131282
40. Maze I, Feng J, Wilkinson MB, Sun H, Shen L, Nestler EJ. Cocaine dynamically regulates heterochromatin and repetitive element unsilencing in nucleus accumbens. *Proc Natl Acad Sci U S A*. 2011; 108(7):3035–40. Epub 2011/02/09. PubMed Central PMCID: PMC3041122. <https://doi.org/10.1073/pnas.1015483108> PMID: 21300862
41. Siddiquee K, Zhang S, Guida WC, Blaskovich MA, Greedy B, Lawrence HR, et al. Selective chemical probe inhibitor of Stat3, identified through structure-based virtual screening, induces antitumor activity. *Proc Natl Acad Sci U S A*. 2007; 104(18):7391–6. <https://doi.org/10.1073/pnas.0609757104> PMID: 17463090
42. Song H, Wang R, Wang S, Lin J. A low-molecular-weight compound discovered through virtual database screening inhibits Stat3 function in breast cancer cells. *Proc Natl Acad Sci U S A*. 2005; 102(13):4700–5. <https://doi.org/10.1073/pnas.0409894102> PMID: 15781862
43. Zhang X, Yue P, Page BD, Li T, Zhao W, Namanja AT, et al. Orally bioavailable small-molecule inhibitor of transcription factor Stat3 regresses human breast and lung cancer xenografts. *Proc Natl Acad Sci U S A*. 2012; 109(24):9623–8. PubMed Central PMCID: PMC3386073. <https://doi.org/10.1073/pnas.1121606109> PMID: 22623533
44. Bromberg J, Darnell JE Jr. The role of STATs in transcriptional control and their impact on cellular function. *Oncogene*. 2000; 19(21):2468–73. <https://doi.org/10.1038/sj.onc.1203476> PMID: 10851045
45. Kidder BL, Yang J, Palmer S. Stat3 and c-Myc genome-wide promoter occupancy in embryonic stem cells. *PLoS One*. 2008; 3(12):e3932. <https://doi.org/10.1371/journal.pone.0003932> PMID: 19079543
46. Park DM, Jung J, Masjkur J, Makrogkikas S, Ebermann D, Saha S, et al. Hes3 regulates cell number in cultures from glioblastoma multiforme with stem cell characteristics. *Scientific reports*. 2013; 3:1095. PubMed Central PMCID: PMC3566603. <https://doi.org/10.1038/srep01095> PMID: 23393614
47. Yang L, Lian X, Cowen A, Xu H, Du H, Yan C. Synergy between signal transducer and activator of transcription 3 and retinoic acid receptor-alpha in regulation of the surfactant protein B gene in the lung. *Mol Endocrinol*. 2004; 18(6):1520–32. <https://doi.org/10.1210/me.2003-0458> PMID: 15044588
48. Asano H, Aonuma M, Sanosaka T, Kohyama J, Namihira M, Nakashima K. Astrocyte differentiation of neural precursor cells is enhanced by retinoic acid through a change in epigenetic modification. *Stem Cells*. 2009; 27(11):2744–52. Epub 2009/07/18. <https://doi.org/10.1002/stem.176> PMID: 19609931
49. Laks DR, Masterman-Smith M, Visnyei K, Angenieux B, Orozco NM, Foran I, et al. Neurosphere formation is an independent predictor of clinical outcome in malignant glioma. *Stem Cells*. 2009; 27(4):980–7. <https://doi.org/10.1002/stem.15> PMID: 19353526
50. Chen S, Ma J, Wu F, Xiong LJ, Ma H, Xu W, et al. The histone H3 Lys 27 demethylase JMJD3 regulates gene expression by impacting transcriptional elongation. *Genes & development*. 2012; 26(12):1364–75. PubMed Central PMCID: PMC3387663.
51. Huang X, Hu J, Li Y, Zhuyun Yang Z, Zhu H, Zhou L, et al. The cell adhesion molecule L1 regulates the expression of FGF21 and enhances neurite outgrowth. *Brain Res*. 2013; 1530:13–21. Epub 2013/08/07. <https://doi.org/10.1016/j.brainres.2013.07.043> PMID: 23916735
52. Carrillo-Garcia C, Prochnow S, Simeonova IK, Strelau J, Holzl-Wenig G, Mandl C, et al. Growth/differentiation factor 15 promotes EGFR signalling, and regulates proliferation and migration in the hippocampus of neonatal and young adult mice. *Development*. 2014; 141(4):773–83. Epub 2014/02/06. PubMed Central PMCID: PMC3930467. <https://doi.org/10.1242/dev.096131> PMID: 24496615
53. Lee SH, Krisanapun C, Baek SJ. NSAID-activated gene-1 as a molecular target for capsaicin-induced apoptosis through a novel molecular mechanism involving GSK3beta, C/EBPbeta and ATF3. *Carcinogenesis*. 2010; 31(4):719–28. Epub 2010/01/30. PubMed Central PMCID: PMC32847092. <https://doi.org/10.1093/carcin/bgg016> PMID: 20110283
54. Shimizu S, Kadowaki M, Yoshioka H, Kambe A, Watanabe T, Kinyamu HK, et al. Proteasome inhibitor MG132 induces NAG-1/GDF15 expression through the p38 MAPK pathway in glioblastoma cells. *Biochem Biophys Res Commun*. 2013; 430(4):1277–82. Epub 2012/12/25. PubMed Central PMCID: PMC3558552. <https://doi.org/10.1016/j.bbrc.2012.11.137> PMID: 23261467

55. Romm E, Nielsen JA, Kim JG, Hudson LD. Myt1 family recruits histone deacetylase to regulate neural transcription. *Journal of neurochemistry*. 2005; 93(6):1444–53. <https://doi.org/10.1111/j.1471-4159.2005.03131.x> PMID: [15935060](https://pubmed.ncbi.nlm.nih.gov/15935060/)
56. Barradas M, Anderton E, Acosta JC, Li S, Banito A, Rodriguez-Niedenfuhr M, et al. Histone demethylase JMJD3 contributes to epigenetic control of INK4a/ARF by oncogenic RAS. *Genes & development*. 2009; 23(10):1177–82. Epub 2009/05/20. PubMed Central PMCID: PMC2685533.
57. Piccirillo SG, Reynolds BA, Zanetti N, Lamorte G, Binda E, Broggi G, et al. Bone morphogenetic proteins inhibit the tumorigenic potential of human brain tumour-initiating cells. *Nature*. 2006; 444(7120):761–5. <https://doi.org/10.1038/nature05349> PMID: [17151667](https://pubmed.ncbi.nlm.nih.gov/17151667/)
58. Estaras C, Akizu N, Garcia A, Beltran S, de la Cruz X, Martinez-Balbas MA. Genome-wide analysis reveals that Smad3 and JMJD3 HDM co-activate the neural developmental program. *Development*. 2012; 139(15):2681–91. <https://doi.org/10.1242/dev.078345> PMID: [22782721](https://pubmed.ncbi.nlm.nih.gov/22782721/)
59. Agherbi H, Gaussmann-Wenger A, Verthuy C, Chasson L, Serrano M, Djabali M. Polycomb mediated epigenetic silencing and replication timing at the INK4a/ARF locus during senescence. *PLoS One*. 2009; 4(5):e5622. Epub 2009/05/23. PubMed Central PMCID: PMC2680618. <https://doi.org/10.1371/journal.pone.0005622> PMID: [19462008](https://pubmed.ncbi.nlm.nih.gov/19462008/)
60. Ivanchuk SM, Mondal S, Dirks PB, Rutka JT. The INK4A/ARF locus: role in cell cycle control and apoptosis and implications for glioma growth. *J Neurooncol*. 2001; 51(3):219–29. PMID: [11407594](https://pubmed.ncbi.nlm.nih.gov/11407594/)
61. Marei HE, Althani A, Afifi N, Michetti F, Pescatori M, Pallini R, et al. Gene expression profiling of embryonic human neural stem cells and dopaminergic neurons from adult human substantia nigra. *PLoS One*. 2011; 6(12):e28420. Epub 2011/12/14. PubMed Central PMCID: PMC3233561. <https://doi.org/10.1371/journal.pone.0028420> PMID: [22163301](https://pubmed.ncbi.nlm.nih.gov/22163301/)
62. Yan Y, Shin S, Jha BS, Liu Q, Sheng J, Li F, et al. Efficient and rapid derivation of primitive neural stem cells and generation of brain subtype neurons from human pluripotent stem cells. *Stem Cells Transl Med*. 2013; 2(11):862–70. Epub 2013/10/12. PubMed Central PMCID: PMC3808201. <https://doi.org/10.5966/sctm.2013-0080> PMID: [24113065](https://pubmed.ncbi.nlm.nih.gov/24113065/)
63. Fukada T, Ohtani T, Yoshida Y, Shirogane T, Nishida K, Nakajima K, et al. STAT3 orchestrates contradictory signals in cytokine-induced G1 to S cell-cycle transition. *Embo J*. 1998; 17(22):6670–7. <https://doi.org/10.1093/emboj/17.22.6670> PMID: [9822610](https://pubmed.ncbi.nlm.nih.gov/9822610/)
64. Rokavec M, Oner MG, Li H, Jackstadt R, Jiang L, Lodygin D, et al. IL-6R/STAT3/miR-34a feedback loop promotes EMT-mediated colorectal cancer invasion and metastasis. *J Clin Invest*. 2014; 124(4):1853–67. Epub 2014/03/20. PubMed Central PMCID: PMC3973098. <https://doi.org/10.1172/JCI73531> PMID: [24642471](https://pubmed.ncbi.nlm.nih.gov/24642471/)
65. Dauer DJ, Ferraro B, Song L, Yu B, Mora L, Buettner R, et al. Stat3 regulates genes common to both wound healing and cancer. *Oncogene*. 2005; 24(21):3397–408. Epub 2005/03/01. <https://doi.org/10.1038/sj.onc.1208469> PMID: [15735721](https://pubmed.ncbi.nlm.nih.gov/15735721/)
66. Androusellis-Theotokis A, Leker RR, Soldner F, Hoepfner DJ, Ravin R, Poser SW, et al. Notch signaling regulates stem cell numbers in vitro and in vivo. *Nature*. 2006; 442(7104):823–6. <https://doi.org/10.1038/nature04940> PMID: [16799564](https://pubmed.ncbi.nlm.nih.gov/16799564/)
67. Lee H, Zhang P, Herrmann A, Yang C, Xin H, Wang Z, et al. Acetylated STAT3 is crucial for methylation of tumor-suppressor gene promoters and inhibition by resveratrol results in demethylation. *Proc Natl Acad Sci U S A*. 2012; 109(20):7765–9. <https://doi.org/10.1073/pnas.1205132109> PMID: [22547799](https://pubmed.ncbi.nlm.nih.gov/22547799/)
68. Gallo M, Coutinho FJ, Vanner RJ, Gayden T, Mack SC, Murison A, et al. MLL5 Orchestrates a Cancer Self-Renewal State by Repressing the Histone Variant H3.3 and Globally Reorganizing Chromatin. *Cancer Cell*. 2015; 28(6):715–29. <https://doi.org/10.1016/j.ccell.2015.10.005> PMID: [26626085](https://pubmed.ncbi.nlm.nih.gov/26626085/)
69. Heinemann B, Nielsen JM, Hudlebusch HR, Lees MJ, Larsen DV, Boesen T, et al. Inhibition of demethylases by GSK-J1/J4. *Nature*. 2014; 514(7520):E1–2. <https://doi.org/10.1038/nature13688> PMID: [25279926](https://pubmed.ncbi.nlm.nih.gov/25279926/)
70. Abdouh M, Facchino S, Chato W, Balasingam V, Ferreira J, Bernier G. BMI1 sustains human glioblastoma multiforme stem cell renewal. *J Neurosci*. 2009; 29(28):8884–96. Epub 2009/07/17. <https://doi.org/10.1523/JNEUROSCI.0968-09.2009> PMID: [19605626](https://pubmed.ncbi.nlm.nih.gov/19605626/)
71. Kim E, Kim M, Woo DH, Shin Y, Shin J, Chang N, et al. Phosphorylation of EZH2 activates STAT3 signaling via STAT3 methylation and promotes tumorigenicity of glioblastoma stem-like cells. *Cancer Cell*. 2013; 23(6):839–52. Epub 2013/05/21. PubMed Central PMCID: PMC34109796. <https://doi.org/10.1016/j.ccr.2013.04.008> PMID: [23684459](https://pubmed.ncbi.nlm.nih.gov/23684459/)
72. de Vries NA, Hulsman D, Akhtar W, de Jong J, Miles DC, Blom M, et al. Prolonged Ezh2 Depletion in Glioblastoma Causes a Robust Switch in Cell Fate Resulting in Tumor Progression. *Cell reports*. 2015.
73. Ntziachristos P, Tsirigos A, Welstead GG, Trimarchi T, Bakogianni S, Xu L, et al. Contrasting roles of histone 3 lysine 27 demethylases in acute lymphoblastic leukaemia. *Nature*. 2014; 514:513–7. <https://doi.org/10.1038/nature13605> PMID: [25132549](https://pubmed.ncbi.nlm.nih.gov/25132549/)

74. Yamaguchi H, Hung M-C. Regulation and Role of EZH2 in Cancer. *Cancer research and treatment: official journal of Korean Cancer Association*. 2014; 46:209–22.
75. Lee J, Son MJ, Woolard K, Donin NM, Li A, Cheng CH, et al. Epigenetic-mediated dysfunction of the bone morphogenetic protein pathway inhibits differentiation of glioblastoma-initiating cells. *Cancer Cell*. 2008; 13(1):69–80. <https://doi.org/10.1016/j.ccr.2007.12.005> PMID: [18167341](https://pubmed.ncbi.nlm.nih.gov/18167341/)
76. Gu F, Hata R, Ma YJ, Tanaka J, Mitsuda N, Kumon Y, et al. Suppression of Stat3 promotes neurogenesis in cultured neural stem cells. *Journal of neuroscience research*. 2005; 81(2):163–71. <https://doi.org/10.1002/jnr.20561> PMID: [15948155](https://pubmed.ncbi.nlm.nih.gov/15948155/)
77. Bonni A, Sun Y, Nadal-Vicens M, Bhatt A, Frank DA, Rozovsky I, et al. Regulation of gliogenesis in the central nervous system by the JAK-STAT signaling pathway. *Science*. 1997; 278(5337):477–83. PMID: [9334309](https://pubmed.ncbi.nlm.nih.gov/9334309/)
78. Iliopoulos D, Hirsch HA, Struhl K. An epigenetic switch involving NF-kappaB, Lin28, Let-7 MicroRNA, and IL6 links inflammation to cell transformation. *Cell*. 2009; 139(4):693–706. Epub 2009/11/03. PubMed Central PMCID: PMC2783826. <https://doi.org/10.1016/j.cell.2009.10.014> PMID: [19878981](https://pubmed.ncbi.nlm.nih.gov/19878981/)
79. Iliopoulos D, Jaeger SA, Hirsch HA, Bulyk ML, Struhl K. STAT3 activation of miR-21 and miR-181b-1 via PTEN and CYLD are part of the epigenetic switch linking inflammation to cancer. *Mol Cell*. 2010; 39(4):493–506. Epub 2010/08/28. PubMed Central PMCID: PMC2929389. <https://doi.org/10.1016/j.molcel.2010.07.023> PMID: [20797623](https://pubmed.ncbi.nlm.nih.gov/20797623/)
80. Song MR, Ghosh A. FGF2-induced chromatin remodeling regulates CNTF-mediated gene expression and astrocyte differentiation. *Nat Neurosci*. 2004; 7(3):229–35. Epub 2004/02/11. <https://doi.org/10.1038/nn1192> PMID: [14770186](https://pubmed.ncbi.nlm.nih.gov/14770186/)
81. Urayama S, Semi K, Sanosaka T, Hori Y, Namihira M, Kohyama J, et al. Chromatin Accessibility at a STAT3 Target Site Is Altered Prior to Astrocyte Differentiation. *Cell Struct Funct*. 2013; 38(1):55–66. Epub 2013/02/27. PMID: [23439558](https://pubmed.ncbi.nlm.nih.gov/23439558/)
82. Takizawa T, Nakashima K, Namihira M, Ochiai W, Uemura A, Yanagisawa M, et al. DNA methylation is a critical cell-intrinsic determinant of astrocyte differentiation in the fetal brain. *Dev Cell*. 2001; 1(6):749–58. Epub 2001/12/13. PMID: [11740937](https://pubmed.ncbi.nlm.nih.gov/11740937/)