



Regulatory domain or CpG site variation in SLC12A5, encoding the chloride transporter KCC2, in human autism and schizophrenia

Citation

Merner, Nancy D., Madison R. Chandler, Cynthia Bourassa, Bo Liang, Arjun R. Khanna, Patrick Dion, Guy A. Rouleau, and Kristopher T. Kahle. 2015. "Regulatory domain or CpG site variation in SLC12A5, encoding the chloride transporter KCC2, in human autism and schizophrenia." *Frontiers in Cellular Neuroscience* 9 (1): 386. doi:10.3389/fncel.2015.00386. <http://dx.doi.org/10.3389/fncel.2015.00386>.

Published Version

doi:10.3389/fncel.2015.00386

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:23845213>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available. Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)



Regulatory domain or CpG site variation in *SLC12A5*, encoding the chloride transporter *KCC2*, in human autism and schizophrenia

Nancy D. Merner^{1,2}, Madison R. Chandler¹, Cynthia Bourassa², Bo Liang³, Arjun R. Khanna⁴, Patrick Dion², Guy A. Rouleau^{2*} and Kristopher T. Kahle^{4,5*†}

OPEN ACCESS

Edited by:

Ludovic MARTIN,
Université de Paris V, France

Reviewed by:

Melanie A Woodin,
University of Toronto, Canada
Wen-Jun Gao,
Drexel University College of Medicine,
USA

*Correspondence:

Guy A. Rouleau,
Department of Neurology and
Neurosurgery, Montreal Neurological
Hospital and Institute, Montréal,
QC, Canada
guy.rouleau@mcgill.ca;
Kristopher T. Kahle,
Department of Neurosurgery, Boston
Children's Hospital, Harvard Medical
School, 1309 Enders Research
Building, 320 Longwood Avenue,
Boston, MA 02115, USA
kkahle@enders.tch.harvard.edu

† Present address:

Kristopher T. Kahle,
Department of Neurosurgery and
Pediatrics, Yale School of Medicine,
Yale Program on Neurogenetics,
New Haven, CT, USA

Received: 12 July 2015

Accepted: 16 September 2015

Published: 12 October 2015

Citation:

Merner ND, Chandler MR,
Bourassa C, Liang B, Khanna AR,
Dion P, Rouleau GA and Kahle KT
(2015) Regulatory domain or CpG
site variation in *SLC12A5*, encoding
the chloride transporter *KCC2*, in
human autism and schizophrenia.
Front. Cell. Neurosci. 9:386.
doi: 10.3389/fncel.2015.00386

¹ Harrison School of Pharmacy, Department of Drug Discovery and Development, Auburn University, Auburn, AL, USA, ² Department of Neurology and Neurosurgery, Montreal Neurological Hospital and Institute, McGill University, Montréal, QC, Canada, ³ Department of Biological Chemistry and Molecular Pharmacology (BCMP), Harvard Medical School, Boston, MA, USA, ⁴ Department of Neurosurgery, Boston Children's Hospital and Harvard Medical School, Boston, MA, USA, ⁵ Manton Center for Orphan Disease Research, Boston Children's Hospital, Boston, MA, USA

Many encoded gene products responsible for neurodevelopmental disorders (NDs) like autism spectrum disorders (ASD), schizophrenia (SCZ), intellectual disability (ID), and idiopathic generalized epilepsy (IGE) converge on networks controlling synaptic function. An increase in *KCC2* (*SLC12A5*) Cl⁻ transporter activity drives the developmental GABA excitatory-inhibitory sequence, but the role of *KCC2* in human NDs is essentially unknown. Here, we report two rare, non-synonymous (NS), functionally-impairing variants in the *KCC2* C-terminal regulatory domain (CTRD) in human ASD (R952H and R1049C) and SCZ (R952H) previously linked with IGE and familial febrile seizures, and another novel NS *KCC2* variant in ASD (R1048W) with highly-predicted pathogenicity. Exome data from 2517 simplex families in the ASD Simon Simplex Collection (SSC) revealed significantly more *KCC2* CTRD variants in ASD cases than controls, and interestingly, these were more often synonymous and predicted to disrupt or introduce a CpG site. Furthermore, full gene analysis showed ASD cases are more likely to contain rare *KCC2* variants affecting CpG sites than controls. These data suggest genetically-encoded dysregulation of *KCC2*-dependent GABA signaling may contribute to multiple human NDs.

Keywords: *KCC2*, *NKCC1*, GABA, neurodevelopmental disorders, autism, schizophrenia

Introduction

Neurodevelopmental disorders (NDs) encompass a wide range of diseases, all of which feature some element of impaired brain development, and are associated with cognitive, neurological, and/or psychiatric dysfunction (Rubenstein, 2011). Common NDs include intellectual disability (ID), autism spectrum disorder (ASD), schizophrenia (SCZ), and epilepsy (Guilmatre et al., 2009; Bozzi et al., 2012), and although classified into distinct disease categories, these disorders show phenotypic overlap and shared genetic risk factors (Mitchell, 2011; Coe et al., 2012). The genetic architecture of NDs is complex, with oligogenic contributions converging on the disruption of the structure, function, and/or plasticity of neuronal networks (Pescosolido et al., 2012). Genomics has shed insight into mechanisms underlying the overlap among

the NDs, with copy number variant, exome sequencing, and genome-wide association study data suggesting a spectrum of neurodevelopmental pathology indexed by mutational load or severity (Krystal and State, 2014). Emerging evidence has shown many of the encoded gene products responsible for individual NDs converge on a relatively limited number of protein-protein interaction networks that operate in the same molecular processes, such as those controlling synaptic structure and function (Gilman et al., 2011; Voineagu et al., 2011; O'Roak et al., 2012; EPGP Collaborative et al., 2013; de Rubeis et al., 2014; Krumm et al., 2014).

Small, diverse populations of inhibitory GABAergic interneurons regulate the activity of excitatory neurons and their involved circuits. If inhibition is impaired, disturbance in excitatory/inhibitory balance can lead to the dysfunction of cognitive processes (Braat and Kooy, 2015). A common feature among different NDs is impaired GABAergic inhibition (Hashimoto et al., 2008; Kang and Macdonald, 2009; Schmidt and Mirnics, 2014), and neuronal hyperexcitability has been implicated in the pathogenesis of ASD (Coghlan et al., 2012), SCZ (Lewis et al., 2012), Rett syndrome (Medrihan et al., 2008; Chao et al., 2010), Tourette syndrome (Kalanithi et al., 2005), tuberous sclerosis (Talos et al., 2012), and neurofibromatosis type I (Cui et al., 2008). The strong bidirectional association between NDs and epilepsy suggests impaired GABAergic inhibition as a potential pathogenic mechanism of mutual susceptibility (Tuchman and Rapin, 2002; Brooks-Kayal, 2010; Chang et al., 2011; Deidda et al., 2014).

Ionotropic GABA_ARs are ligand-gated Cl⁻ channels, and the post-synaptic response to GABA_AR activation is significantly modulated by the intraneuronal concentration of Cl⁻ ([Cl⁻]_i). In immature neurons, [Cl⁻]_i is sufficiently high that GABA_AR activation triggers giant depolarizing potentials, which characterize early network activity and stimulate Ca²⁺-dependent synaptogenesis (Ben-Ari et al., 1989; Belhage et al., 1998). A post-natal increase in the functional expression of KCC2 (*SLC12A5*), a cation-Cl⁻ cotransporter (CCC) mediating Cl⁻ efflux, lowers [Cl⁻]_i in post-synaptic neurons such that GABA_AR activation elicits membrane hyperpolarization and fast synaptic inhibition (Kaila et al., 2014). KCC2 is required for the normal developmental GABA excitatory-inhibitory sequence, and KCC2 deficiency disrupts normal brain development and results in network hyperexcitability (Hubner et al., 2001; Hekmat-Safe et al., 2006, 2010; Tanis et al., 2009; Bellemer et al., 2011).

Abnormal functional expression of KCC2 or its Cl⁻ importing cousin NKCC1, and associated impairment of GABA inhibition, have been documented in rodent models of multiple different NDs [see Table 1 in Deidda et al. (2014)]. For example, in mouse models of Fragile X syndrome, the most common genetic cause of human autism, there is a significantly delayed developmental switch to GABAergic inhibition due to prolonged elevation in neuronal [Cl⁻]_i due in part to decreased KCC2 expression (Lemonnier et al., 2013), and blockade of up-regulated NKCC1-mediated Cl⁻ import in this

context normalizes [Cl⁻]_i, electrophysiological responses, and autistic-like behaviors in these mice (Tyzio et al., 2014), and improves autistic behaviors in humans (Lemonnier et al., 2012). However, the role of KCC2 in the human nervous system and in NDs is essentially unknown. Identification of such variants could elucidate the molecular pathophysiology of these diseases and identify therapeutic targets.

The KCC2 C-terminal regulatory domain (CTRD) is a critical region of transporter function, and contains multiple phosphorylated residues (such as Thr906/Thr1007 and Ser940 (reviewed in Blaesle et al., 2009; Chamma et al., 2012; Kahle et al., 2013; Medina et al., 2014); and functional domains (e.g., Acton et al., 2012) that establish context-appropriate transport activity (de Los Heros et al., 2014; see **Figure 1**). We recently identified an enrichment of KCC2 non-synonymous (NS) alleles in French Canadian (FC) idiopathic generalized epilepsy (IGE) cases compared to controls that included two rare and functional IGE risk alleles (KCC2 R952H and R1049C; Kahle et al., 2014). Both of these variants decrease KCC2-mediated Cl⁻ extrusion and render neuronal GABA activity more depolarized by decreasing the amount of Ser940 stimulatory phosphorylation (Kahle et al., 2014). KCC2 R952H was also identified as a cause of an inherited form of febrile seizures in an Australian family, and shown to impair dendritic spine formation (Puskarjov et al., 2014).

Here, we sought to determine if risk alleles in the KCC2 CTRD (amino acids 894–1086; NP_065759) were present in large FC cohorts of ASD, SCZ, or ID by utilizing a similar targeted genetic sequencing approach we successfully utilized for IGE (Kahle et al., 2014). We speculated that genetic mutations and/or functionally impairing variants in the KCC2 CTRD might contribute to NDs.

Materials and Methods

Clinical Sampling

The three disease cohorts of ASD, SCZ, and ID, were established as a part of the Synapse to Disease (S2D) project initiated by Dr. Guy Rouleau at Montreal Neurological Institute and McGill University in order to identify genes that cause or predispose to numerous disorders of brain development. The sample characteristics of these FC cohorts have previously been described (Gauthier et al., 2005; Hamdan et al., 2009; Awadalla et al., 2010).

KCC2 C-Terminus Targeted Screening

We implemented a targeted DNA Sanger sequencing approach to screen the 3' end of *SLC12A5* that encodes the CTRD of KCC2 (Kahle et al., 2014). Specifically, we targeted the coding nucleotides in exons 21–25 of *SLC12A5* [NM_020708.4 (NP_065759 amino acids 894–1086) or NM_001134771 (NP_001128243 amino acids 917–1110)] by following the same protocol outlined in Kahle et al. (2014); a total of 427 ASD, 143 SCZ and 190 ID cases were screened. A total of 1214 matched controls were previously screened for mutations in the targeted region of KCC2 (Kahle et al., 2014).

TABLE 1 | SLC12A5 variants detected in the FC ASD cohort through the targeted screening of the C-terminus.

Variants detected in the screened region			Detection of variants in Quebec ASD cohort			FC population controls (Total: 1214 controls)			
NM_020708.4 NP_065759	NM_001134771.1 NP_001128243	rs ID	Number of probands	Number of alleles	Allele frequency (%)	Number of alleles	Allele frequency (%)	p-value	Odds ratio
c.2855 G > A p.R952H	c.2924 G > A p.R975H	rs142740233	2/427	2/854	0.23	5/2428	0.21	1.00	1.14 Cl ₉₅ [0.1–7.0]
c.2961 G > A p.P987P	c.3030 G > A p.P1010P	rs550491448	1/427	1/854	0.12	0/2428	0.00	0.26	Inf Cl ₉₅ [0.1–Inf]
c.3142 C > T p.R1048W	c.3211C > T p.R1071W	rs369042030	1/427	1/854	0.12	0/2428	0.00	0.26	Inf Cl ₉₅ [0.1–Inf]
c.3145 C > T p.R1049C	c.3214 C > T p.R1072C	–	1/427	1/854	0.12	1/2428	4.12 × 10 ⁻⁴	0.45	2.84 Cl ₉₅ [0.0–223.0]
Total number of variants detected			5	5	–	6	–	0.17	2.37 Cl ₉₅ [0.6–9.4]

Protein Sequence Alignment and *In Silico* Prediction Programs

ClustalW (Larkin et al., 2007) and WebLogo (Crooks et al., 2004) were used to align different orthologues of the KCC2 to determine the evolutionary conservation of the novel KCC2 variant, R1048W (Figures 1B,C). Conservation of KCC2 R952H and R1049C were previously demonstrated (Kahle et al., 2014).

Conservation of the KCC2 protein was determined by aligning the following orthologues: *Homo sapiens* (NP_065759), *Macaca mulatta* (XM_001104494.2_prot), *Bos Taurus* (NP_001193309), *Rattus norvegicus* (NP_599190), *Mus musculus* (NP_065066), *Tetraodon nigroviridis* (ENSTNIT00000021299), and *Danio rerio* (ENSDART00000009569). Regarding the WebLogo output, the y-axis serves as a means of determining relative conservation




FIGURE 1 | KCC2 (SLC12A5) variants in human autism spectrum disorder (ASD) and schizophrenia (SCZ). (A) DNA chromatograms illustrating the detection of KCC2 variants in ASD (c.2855 G > A [p.R952H]; c.3145 C > T [p.R1049C]; and c.3142 C > T [p.R1048W]); and SCZ (p.R952H) via Sanger sequencing. (B,C) Evolutionary conservation of amino acid p.R1048; and conservation of amino acids p.R952 and p.R1049 shown in Kahle et al. (2014). (D) Schematic representation of KCC2 (human). Orange dots indicate the positions of the known critical phospho-regulatory residues p.T906, p.S940, p.T1007, and p.Y1087 (reviewed in Chamma et al., 2012; Kahle et al., 2013); Pink region denotes the KCC2 “ISO” domain, required for hyperpolarizing GABAergic transmission (Acton et al., 2012). Red dots depict the identified IGE mutations, p.R952 and p.R1049; green dots depict the identified ASD variants, p.R952H, p.R1049C, and p.R1048W; yellow dots indicate the identified SCZ variant p.R952H. (E) The modeled structure of the human KCC2 C-terminal domain (CTRD), based on homology modeling by I-TASSER (Roy et al., 2010) using a prokaryotic member of the CCC family (PDB code 3g40) (for details, see “Materials and Methods” Section). Color scheme same as in (D). Note the proximity of the novel p.R1048W ASD variant and the previously described KCC2 IGE variants, as well as their relation to important regulatory residues and domains. (F) Venn diagram showing overlap of KCC2 variants in multiple neurodevelopmental phenotypes (IGE, ASD, and SCZ) that exhibit dysfunctional GABA signaling (Deidda et al., 2014).

and is not an actual measurement; the height of each stack at each amino acid position is relative to the overall conservation at that position, the height of the letters within each stack indicate the relative frequencies for each amino acid possibility, and the width of each stack corresponds to the proportion of valid readings at that position (indicating if sequence gaps exist between the shown amino acids). The effects of amino acid substitutions on protein function were predicted using MutationTaster (Schwarz et al., 2010), Panther (Mi et al., 2005), and Polyphen-2 (Adzhubei et al., 2010).

Exome Sequencing Analysis

Iossifov et al. (2014) recently reported the whole exome sequencing of 2517 ASD simplex families from the Simon Simplex Collection (SSC; Fischbach and Lord, 2010). They carried out a *de novo* mutation analysis that generated an extensive list of *de novo* variants and recurrently hit genes that could be subdivided with different clinical phenotypes (Iossifov et al., 2014). These exome sequencing data are now available through the NDAR (National Database of Autism Research); we were granted access to variant calling files to further assess *SLC12A5*.

Our analysis involved filtering for sample type to include only probands (.p1), accession number (NM_001134771) to include only *SLC12A5*, and “_par-races_” (we either selected “white, white” or “african-amer, african-amer”) to include probands with either two Caucasian parents (EA ASD cases) or two African American parents (AA ASD cases), respectively. According to Supplementary Table 1 in Iossifov et al. (2014), there were 1892 EA ASD cases and 82 AA ASD cases. We carried out a rare variant (<1% MAF) analysis; therefore, we filtered out SNPs (single nucleotide polymorphisms) with a MAF >1% in each ethnic group. This resulted in a list of rare coding variants for the full gene and our targeted gene region. Furthermore, we used the EA and AA Exome Variant Server (EVS) data (NHLBI GO Exome Sequencing Project)¹ as our ethnic controls; there were a total of 4300 EA controls and 2203 AA controls exome sequenced. We generated lists of rare variants that were detected in each control cohort.

Statistical Analysis

All statistical genetic analysis was carried out using the program R (version 2.15.1). Fisher tests or Mantel-Haenzel Chi Squared

tests were carried out to generate the *p*-values and odds ratios where appropriate.

Results

Using Sanger sequencing, we examined the 3′ end of *SLC12A5* that encodes the KCC2 CTRD (amino acids 894–1086; NP_065759) in three large FC disease cohorts of ASD, SCZ, or ID that were collected as part of the S2D project (see “Materials and Methods” Section). In contrast to our previous analysis in IGE which identified an enrichment of KCC2 NS CTRD alleles in IGE cases compared to controls (*p*-value = 7.50×10^{-3} ; Kahle et al., 2014), analysis of our initial ASD, SCZ or ID sequencing results did not show an enrichment of NS KCC2 CTRD alleles in cases (Table 1). Interestingly, however, three different heterozygous and NS KCC2 variants were detected in the ASD cohort; these included the two previously-identified IGE risk variants, R952H and R1049C (Kahle et al., 2014), and R1048W (Figure 1 and Table 1).

R1048 in KCC2 is a highly conserved residue (Figure 1), and a substitution for a tryptophan at this position is predicted to be highly pathogenic using multiple *in silico* bioinformatics programs (Table 2). This variant is extremely rare; it was not detected in 2428 FC alleles, but this number of controls was too small to generate a significant *p*-value (Table 1). Indeed, power analysis determined that 50,000 control alleles would need to be genotyped, assuming the allele frequencies remain the same and a *p*-value of 0.0167 is significant (after a Bonferroni correction). This limitation is common when identifying rare genetic risk factors, and also relevant for the R1049C variant (Table 1), where 100,000 control alleles would need to be genotyped to reach significance. R952H was previously determined to have an allele frequency of 0.66% in the FC IGE cohort (Kahle et al., 2014) compared to 0.21% in FC controls and 0.23% in the FC ASD cohort (Table 1). One SCZ patient was also determined to carry R952H, corresponding to an allele frequency of 0.35% (Table 3). Two heterozygous synonymous variants were also detected in our screening, including P987P in an ASD patient (Table 1), and D935D in an SCZ patient (Table 3).

From a functional standpoint, both R952H and R1049C impair KCC2 transporter activity (Kahle et al., 2014); they significantly decrease KCC2-mediated Cl⁻ extrusion capacity in neurons, render EGly less hyperpolarized compared to WT KCC2, decrease the level of stimulatory phosphorylation of Ser940, act in a dominant-negative manner consistent with

¹<http://evs.gs.washington.edu/EVS/>

TABLE 2 | Predicted pathogenicity of the novel KCC2 (*SLC12A5*) variant.

Variant name		Prediction programs			
NM_020708.4 NP_065759	NM_001134771.1 NP_001128243	Mutation taster	Panther	Polyphen	
c.3142 C > T p.R1048W	c.3211C > T p.R1071W	Disease causing*	P (probability): 0.9999	P _{deleterious} = 0.80942	Possibly damaging (0.813)

*Predicted specifically to disrupt the function of the last cytoplasmic domain.

TABLE 3 | SLC12A5 variants detected in the SCZ cohort through the targeted screening of the C-terminus.

Variants detected in the screened region			Detection of variants in SCZ cohort			FC population controls (Total: 1214 controls)			
NM_020708.4 NP_065759	NM_001134771.1 NP_001128243	rs ID	Number of probands	Number of alleles	Allele frequency (%)	Number of alleles	Allele frequency (%)	p-value	Odds ratio
c.2805 T > C p.D935D	c.2874 T > C p.D958D	rs151293924	1/143	1/286	0.35	0/2428	0.00	0.11	Inf CI ₉₅ [0.2-Inf]
c.2855 G > A p.R952H	c.2924 G > A p.R975H	rs142740233	1/143	1/286	0.35	5/2428	0.21	0.50	1.70 CI ₉₅ [0.0-15.3]
Total number of variants detected			2	2	-	6*	-	0.20	2.85 CI ₉₅ [0.3-16.1]

*A total of 6 variants were detected in FC control cohort [5 R975H and 1 R1072C (which is not shown in this table)].

the known oligomerization of KCC2 molecules, and decrease transporter plasmalemmal expression (R952H) or lower the intrinsic activity of transporters at the cell surface (R1049C) (Kahle et al., 2014). We anticipate these variants function similarly in ASD; however, the phenotypic outcomes of any effect of these variants are likely dependent on the combination of other risk alleles within each individual patient. Considering the proximity of R1048 to R1049, and the fact that both variants substitute an arginine, we assume the functional effects of R1048W on KCC2 would be similar to that of R1049C (Kahle et al., 2014).

Overall, the data generated from the targeted screening of 3' end of *SLC12A5* in FC cases and controls did not reach statistical significance, possibly due to the rare nature of the variants and the size of the FC cohorts (Table 4). However, when the SSC and EVS exome sequencing data was considered, the combined analysis indicated there was an enrichment of all coding KCC2 CTRD variants in ASD cases compared to controls (*p-value* = 0.03; Table 4). In fact, when subdividing the variants into various groups, we determined that ASD cases actually had significantly more synonymous variants compared to controls (*p-value* = 0.02), as well as

TABLE 4 | SLC12A5 variants detected in the FC and SSC EA and AA cohorts in the targeted region of the C-terminus.

Variant type	Cohort			Number of rare variants transported	p value	Odds ratio	Combined	
	Ethnicity	Group	Size				p value	Odds ratio
All	FC	ASD Cases	427	5	0.17	2.37 CI ₉₅ [0.6-9.4]	0.03	2.00 CI ₉₅ [1.1-3.6]
		Controls	1214	6				
	EA	ASD Cases	1892	12	0.09	1.95 CI ₉₅ [0.8-4.5]		
		EVS Controls	4300	14				
	AA	ASD Cases	82	2	0.35	1.67 CI ₉₅ [0.2-6.8]		
EVS Controls		2203	32					
Non-synonymous	FC	ASD Cases	427	4	0.30	1.90 CI ₉₅ [0.4-8.0]	0.31	1.53 CI ₉₅ [0.8-3.1]
		Controls	1214	6				
	EA	ASD Cases	1892	9	0.37	1.46 CI ₉₅ [0.6-3.6]		
		EVS Controls	4300	14				
	AA	ASD Cases	82	0	1.00	0.00 CI ₉₅ [0.0-29.7]		
EVS Controls		2203	5					
Synonymous	FC	ASD Cases	427	1	0.26	Inf CI ₉₅ [0.1-Inf]	0.02	4.93 CI ₉₅ [1.7-14.8]
		Controls	1214	0				
	EA	ASD Cases	1892	3	0.02	Inf CI ₉₅ [0.9-Inf]		
		EVS Controls	4300	0				
	AA	ASD Cases	82	2	0.28	2.00 CI ₉₅ [0.2-8.1]		
EVS Controls		2203	27					
CpG site disrupted or gained	FC	ASD Cases	427	5	0.17	2.37 CI ₉₅ [0.6-9.4]	6.8 × 10 ⁻³	2.5 CI ₉₅ [1.3-4.7]
		Controls	1214	6				
	EA	ASD Cases	1892	11	0.02	2.78 CI ₉₅ [1.0-7.6]		
		EVS Controls	4300	9				
	AA	ASD Cases	82	2	0.31	1.85 CI ₉₅ [0.2-7.5]		
EVS Controls		2203	29					
CpG site not disrupted or gained	FC	ASD Cases	427	0	1.00	0.00 CI ₉₅ [0.0-Inf]	0.71	0.42 CI ₉₅ [0.0-3.6]
		Controls	1214	0				
	EA	ASD Cases	1892	1	0.67	0.45 CI ₉₅ [0.0-4.1]		
		EVS Controls	4300	5				
	AA	ASD Cases	82	0	1.00	0.00 CI ₉₅ [0.0-65.6]		
EVS Controls		2203	3					

TABLE 5 | Full gene rare variant (MAF < 1%) analysis of *SLC12A5* using the SSC exome sequencing data.

Variant type	Cohort			Number of rare variants transported	<i>p</i> value	Odds ratio	Combined	
	Ethnicity	Group	Size				<i>p</i> value	Odds ratio
All	EA	ASD Cases	1892	50	1.00	1.00 CI ₉₅ [0.7–1.4]	0.76	1.06 CI ₉₅ [0.8–1.4]
		EVS Controls	4300	114				
	AA	ASD Cases	82	9	0.30	1.44 CI ₉₅ [0.6–2.9]		
		EVS Controls	2203	168				
Non-synonymous	EA	ASD Cases	1892	21	0.68	1.11 CI ₉₅ [0.6–1.9]	0.98	1.04 CI ₉₅ [0.6–1.8]
		EVS Controls	4300	43				
	AA	ASD Cases	82	0	1.00	0.00 CI ₉₅ [0.0–4.8]		
		EVS Controls	2203	23				
Synonymous	EA	ASD Cases	1892	29	0.73	1.08 CI ₉₅ [0.7–1.7]	0.38	1.20 CI ₉₅ [0.8–1.8]
		EVS Controls	4300	61				
	AA	ASD Cases	82	9	0.18	1.67 CI ₉₅ [0.7–3.4]		
		EVS Controls	2203	145				
CpG site disrupted or gained	EA	ASD Cases	1892	41	0.22	1.29 CI ₉₅ [0.9–1.9]	0.09	1.37 CI ₉₅ [1.0–1.9]
		EVS Controls	4300	72				
	AA	ASD Cases	82	8	0.15	1.76 CI ₉₅ [0.7–3.7]		
		EVS Controls	2203	122				
CpG site not disrupted or gained	EA	ASD Cases	1892	9	0.05	0.5 CI ₉₅ [0.2–1.0]	0.06	0.50 CI ₉₅ [0.3–1.0]
		EVS Controls	4300	42				
	AA	ASD Cases	82	1	1.00	0.58 CI ₉₅ [0.0–3.5]		
		EVS Controls	2203	46				

variants that either disrupted or introduced a CpG site (*p*-value = 6.8×10^{-3} ; **Table 4**). Upon full gene analysis (using solely exome sequencing data), ASD cases were determined to have a higher percentage of rare *SLC12A5* variants that affect a CpG site compared to controls (**Tables 5, 6**), suggesting a possible epigenetic effect on gene expression through variation in methylation patterns.

Discussion

The identification of KCC2 NS genetic variants in ASD and SCZ cases that involve evolutionary conserved residues, are predicted to be pathogenic, and have either previously been shown [KCC2 R952H and R1049C (Kahle et al., 2014; Puskarjov et al., 2014)] or anticipated (KCC2 R1048W) to impact transporter function, trafficking, and/or regulatory phosphorylation, suggest that genetically-encoded impairment of KCC2 function may be risk factors for, or contribute to the pathogenesis of, human ASD and SCZ. Overall, these data are the first to describe functional KCC2 genetic variants in human psychiatric disease, and suggest a compelling genetic overlap among distinct NDs. Furthermore, we show that CpG sites in the targeted-screened 3' end of

SLC12A5, as well as in the entire gene, are more commonly affected (i.e., disrupted or gained) in ASD cases compared to controls.

Genetic links among the different NDs have been previously shown. For example, chromosome 1q21.1 microdeletions exist in both ID and SCZ, and, interestingly, are often inherited from an unaffected or a mildly affected parent (Christiansen et al., 2004; Brunetti-Pierri et al., 2008; International-Schizophrenia-Consortium, 2008; Mefford et al., 2008; Stefansson et al., 2008). Duplications of this region are also associated with mild to moderate ID and ASD (Brunetti-Pierri et al., 2008; Mefford et al., 2008). Deletions of chromosome 16p13.11 have been associated with ID and ASD (Ullmann et al., 2007), and epilepsy (Heinzen et al., 2010; de Kovel et al., 2010), and duplications of this region have been documented in ID (Ullmann et al., 2007; Mefford et al., 2009), ASD (Ullmann et al., 2007) and SCZ (Kirov et al., 2009). In addition, mutations in *MECP2* [OMIM 300005] (Lam et al., 2000) and *SLC6A8* [OMIM 300036] (Salomons et al., 2001) have been found in both ID and ASD; and, mutations in *SHANK3* [OMIM 606230] have been identified in ASD (Durand et al., 2007), SCZ (Gauthier et al., 2010) and nonsyndromic ID cohorts (Hamdan et al., 2011). The

TABLE 6 | ASD case/control comparisons of the total number of variants that affect a CpG site vs. do not affect a CpG site from the full gene analysis.

Cohort	Total number of rare variants in <i>SLC12A5</i>	Variants that disrupt or gain a CpG site		Variants that do not disrupt or gain a CpG site		<i>p</i> value	Odds ratio	Combined	
		Number	Percentage (%)	Number	Percentage (%)			<i>p</i> value	Odds ratio
EA	ASD Cases	50	41	82	9	0.02	2.64 CI ₉₅ [1.1–6.8]	0.01	2.71 CI ₉₅ [1.3–5.8]
	EVS Controls	114	72	63	42				
AA	ASD Cases	9	8	89	1	0.45	3.00 CI ₉₅ [0.4–136.6]		
	EVS Controls	168	122	73	46				

impact of *de novo* SNVs (single nucleotide variants) on sporadic forms of common NDs have also been increasingly appreciated (Vissers et al., 2010; Girard et al., 2011; O’Roak et al., 2011, 2012; Xu et al., 2011; Iossifov et al., 2012; Neale et al., 2012; Sanders et al., 2012), and *de novo* ASD mutations have been identified in genes previously associated with other NDs, such as *FOXP1* [OMIM 605515], *GRIN2B* [OMIM 138252], *SCN1A* [OMIM 182389], and *LAMC3* [OMIM 604349] (O’Roak et al., 2011). No *de novo* variants were detected in this study.

Identifying rare variants that are associated with a complex trait is most currently performed through whole-exome sequencing since there is a potential to identify many genes that underlie the trait at a substantially lower cost compared to whole-genome sequencing; plus, exome variants offer a clear-cut functional annotation that can be predicted through many *in silico* bioinformatics programs with an accuracy of ~80% (Kiezun et al., 2012). However, with exome sequencing, a *p*-value of less than 2.5×10^{-6} is required to reach genome-wide significance; this value accounts for multiple testing by making a Bonferroni correction for 20,000 independent tests (1 test for each gene in the genome). Such a threshold is very conservative and unless a study is extremely large, significant *p*-values will not be reached. In fact, current exome sequencing studies of complex traits are under-powered since, at minimum, 10,000 individuals with a distinct phenotype are needed in order to achieve the necessary power (Kryukov et al., 2009; Kiezun et al., 2012). Furthermore, rare variants should be combined in a gene (or pathway) during association tests in order to reach sufficient power (Purcell et al., 2003; Kiezun et al., 2012).

A true understanding of how risk alleles interact and contribute to disease is yet to come. In a polygenic or oligogenic disease model, causality cannot be assigned to any one variant, but rather results from an individual’s variant pattern (Klassen et al., 2011). A recent exome sequencing paper aimed to demonstrate the polygenic burden of rare disruptive mutations in SCZ by noting the disruptive mutations that were distributed across many genes as well as enriched gene sets (Purcell et al., 2014). The ultimate goal is to identify all risk variants and to establish a computational modeling of biological networks in order to improve risk predictions based on a combination of alleles. Interestingly, Kiezun et al. (2012) noted that no individual gene-based test achieved significance after correction for multiple testing, again reiterating the need to increase sample size and/or take more targeted screening approaches to minimize multiple testing.

Here, we utilized a hypothesis-driven, targeted sequencing approach to search for variation in the KCC2 CTRD in different NDs given the known neurophysiological importance of this gene (Gagnon and Delpire, 2013), and the demonstrated critical role of our targeted region for the functional regulation of synapses (Chamma et al., 2012; Kahle et al., 2013). This approach previously enabled us to simplify the statistical analysis to identify the enrichment of NS alleles in the KCC2 CTRD in IGE cases vs. controls (*p*-value = 7.50×10^{-3} ; Kahle et al., 2014). Rare variation is enriched for evolutionarily deleterious variants, and we demonstrated that the KCC2 IGE variants (R952H and R1049C) impaired KCC2 function (Kahle et al., 2014). The initial

targeted screening of the KCC2 CTRD in the FC cohort of ASD, SCZ, and ID did not generate any significant associations; however, it revealed that the functional variants overlap across different NDs, suggesting a role in disease pathogenesis, and the effects of these variants are likely dependent on the combination of other alleles within each individual. After combining our initial findings with the results of other exome sequencing projects (Iossifov et al., 2014), we were able to generate improved *p*-values due to the larger sample sets. It was after this analysis that a statistically significant excess of KCC2 CTRD variants in the targeted region was identified in ASD cases compared to controls, as well as an increase of variants that affected CpG sites. It should be noted that the authors recognize the disadvantages of comparing different exome sequencing data sets (SSC and EVS) that were analyzed using different pipelines; replication studies are warranted.

What is the functional impact of the discovered KCC2 variants? The KCC2 R952H and R1049C variants significantly decrease KCC2-mediated Cl⁻ extrusion capacity in neurons, render E_{Cl} less hyperpolarized compared to WT KCC2, decrease the level of KCC2 Ser940 phosphorylation, act in a dominant-negative manner, and decrease transporter plasmalemmal expression (R952H) or lower the intrinsic activity of transporters at the cell surface (R1049C). KCC2 R952H was also shown to substantially decrease dendritic spine density and alter spine morphology (Puskarjov et al., 2014). Given the evolutionary conservation of KCC2 R1048, the predicted pathogenicity of the R1048W substitution by multiple *in silico* algorithms, and the proximity of this variation to R1049, we anticipate the R1048W KCC2 variant detected in ASD, like R1049C, alters the intrinsic activity of KCC2 transporters at the cell surface. The clustering of the R1048 and R1049 variants suggests this region of KCC2 is particularly important for transporter regulation. Consistent with this is the proximity of the KCC2 “ISO” domain, encoded in amino acids 1022–1037, which is required for isotonic KCC2-mediated hyperpolarizing GABAergic transmission (Acton et al., 2012). These variants might change C-terminal protein structure and alter the function of the “ISO” domain, perhaps by disrupting the binding of key associated regulatory molecules.

From a pathophysiological standpoint, decreased KCC2-mediated Cl⁻ efflux in individuals carrying the KCC2 R952H, R1048W, and R1049C variants would be anticipated to increase intracellular [Cl⁻], raising the Cl⁻ reversal potential (E_{Cl}) to less hyperpolarized potentials, and compromising GABA_AR-mediated hyperpolarizing inhibition. In humans, KCC2 is developmentally upregulated, with low expression *in utero*, a rapid increase in expression around 40 postconceptional weeks, and progressively increasing levels of expression into adulthood (Dzhala et al., 2005). This pattern of expression drives the developmental switch of GABAergic signaling from depolarizing in early development to hyperpolarizing in adulthood, but the precise role of KCC2 or its disruption in neurodevelopment is unclear. The progressive developmental increase in KCC2 expression into adulthood might be expected to amplify the functional effects of KCC2 variants on CNS function over time. This may correlate with the age of onset of these neuropsychiatric disorders, as clinical symptomatology might manifest only after

years of neurodevelopment over which deficits in KCC2 activity relative to normal become more pronounced. Further studies are indicated to explore this hypothesis.

The effects of KCC2 variants identified in this study might be similar, though less potent in magnitude given their heterozygosity, to phenotypes observed in mice with complete knockout or mild dysfunction of KCC2. Mice with complete KCC2 knockout die at birth from profound motor deficits that abolish respiratory function (Hekmat-Scafe et al., 2006, 2010; Tanis et al., 2009; Bellemer et al., 2011), but, interestingly, mice expressing hypomorphic alleles that reduce but do not abolish KCC2 activity demonstrate anxiety-like behavior, impaired spatial learning, and decreased seizure threshold (Woo et al., 2002; Tornberg et al., 2005; Zhu et al., 2008). Together, these data indicate that regulation of Cl^- homeostasis and GABAergic signaling by KCC2 plays a role in multiple functional systems of the CNS. These studies have also demonstrated that genetic KCC2 knockdown is not accompanied by compensatory changes in the expression of NKCC1 or other KCC isoforms, although post-translational compensatory regulation (e.g., phosphorylation) cannot be ruled out (Hubner et al., 2001; Woo et al., 2002; Tornberg et al., 2005).

Compellingly, CpG sites in the screened 3' end of *SLC12A5*, as well as in the entire gene, are more commonly affected (i.e., disrupted or gained) in ASD cases compared to controls. Generally, DNA methylation occurs at CpG sites and plays a role in gene expression by suppressing gene transcription. Therefore, differences in methylation patterns between cases and controls could result in different gene expression patterns that increase risk to ASD. CpG islands, regions of CpG clusters that are associated with genes, are generally involved in transcription activation. Interestingly, there is a CpG island of ~2000bp

at the 3' end of *SLC12A5* that partially overlaps with our targeted-screened region. Furthermore, despite the fact that the variants detected in our targeted-screened region are outside that specific CpG island, most methylation differences between tissues and, perhaps even between patients and controls, actually occur at CpG sites at a short distance from the actual CpG island (Irizarry et al., 2009). It would be interesting to screen the entire *SLC12A5* locus (coding and non-coding regions) to determine if non-coding variants contribute towards a stronger association in ASD cases and controls. Additionally, carrying out the same analysis in an SCZ case/control cohort would be valuable.

Lastly, our results are interesting given the recent data from the Fragile X and valproate mouse models of autism, which demonstrate an abnormally prolonged elevation of $[\text{Cl}^-]_i$ in developing CNS neurons that delays the normal ontogenic switch to GABA inhibition (He et al., 2014). Maternal oxytocin during delivery normally mediates the hyperpolarization of E_{GABA} , but this effect is abolished in these models (Tyzio et al., 2014). Our data here suggest a compelling hypothesis that abnormally delayed KCC2-mediated Cl^- extrusion during the developmental GABA excitatory-inhibitory sequence might be a genetically programmed factor contributing to ASD. Further investigation into the roles of KCC2 in ASD and SCZ, and potentially other NDs, may offer new therapeutic strategies.

Acknowledgments

The authors thank JingJing Duan (Harvard Medical School, Boston, MA), Jin Zhang (Harvard Medical School, Boston, MA), and Jinwei Zhang (University of Dundee, Dundee, Scotland) for their technical support and assistance in conducting the experiments described in this study.

References

- Acton, B. A., Mahadevan, V., Mercado, A., Uvarov, P., Ding, Y., Pressey, J., et al. (2012). Hyperpolarizing GABAergic transmission requires the KCC2 C-terminal ISO domain. *J. Neurosci.* 32, 8746–8751. doi: 10.1523/jneurosci.6089-11.2012
- Adzhubei, I. A., Schmidt, S., Peshkin, L., Ramensky, V. E., Gerasimova, A., Bork, P., et al. (2010). A method and server for predicting damaging missense mutations. *Nat. Methods.* 7, 248–249. doi: 10.1038/nmeth0410-248
- Awadalla, P., Gauthier, J., Myers, R. A., Casals, F., Hamdan, F. F., Griffing, A. R., et al. (2010). Direct measure of the de novo mutation rate in autism and schizophrenia cohorts. *Am. J. Hum. Genet.* 87, 316–324. doi: 10.1016/j.ajhg.2010.07.019
- Belhage, B., Hansen, G. H., Elster, L., and Schousboe, A. (1998). Effects of gamma-aminobutyric acid (GABA) on synaptogenesis and synaptic function. *Perspect Dev. Neurobiol.* 5, 235–246.
- Bellemer, A., Hirata, T., Romero, M. F., and Koelle, M. R. (2011). Two types of chloride transporters are required for GABA(A) receptor-mediated inhibition in *C. elegans*. *EMBO. J.* 30, 1852–1863. doi: 10.1038/emboj.2011.83
- Ben-Ari, Y., Cherubini, E., Corradetti, R., and Gaiarsa, J. L. (1989). Giant synaptic potentials in immature rat CA3 hippocampal neurones. *J. Physiol.* 416, 303–325. doi: 10.1113/jphysiol.1989.sp017762
- Blaesse, P., Airaksinen, M. S., Rivera, C., and Kaila, K. (2009). Cation-chloride cotransporters and neuronal function. *Neuron.* 61, 820–838. doi: 10.1016/j.neuron.2009.03.003
- Bozzi, Y., Casarosa, S., and Caleo, M. (2012). Epilepsy as a neurodevelopmental disorder. *Frontiers in Psychiatry.* 3:19. doi: 10.3389/fpsy.2012.00019
- Braat, S., and Kooy, R. F. (2015). The GABA Receptor as a Therapeutic Target for Neurodevelopmental Disorders. *Neuron.* 86, 1119–1130. doi: 10.1016/j.neuron.2015.03.042
- Brooks-Kayal, A. (2010). Epilepsy and autism spectrum disorders: Are there common developmental mechanisms? *Brain and Development.* 32, 731–738. doi: 10.1016/j.braindev.2010.04.010
- Brunetti-Pierri, N., Berg, J. S., Scaglia, F., Belmont, J., Bacino, C. A., Sahoo, T., et al. (2008). Recurrent reciprocal 1q21.1 deletions and duplications associated with microcephaly or macrocephaly and developmental and behavioral abnormalities. *Nat. Genet.* 40, 1466–1471. doi: 10.1038/ng.279
- Chamma, I., Chevy, Q., Poncer, J. C., and Lévi, S. (2012). Role of the neuronal K-Cl co-transporter KCC2 in inhibitory and excitatory neurotransmission. *Front. Cell. Neurosci.* 6:5. doi: 10.3389/fncel.2012.00005
- Chang, Y. T., Chen, P. C., Tsai, I. J., Sung, F. C., Chin, Z. N., Kuo, H. T., et al. (2011). Bidirectional relation between schizophrenia and epilepsy: a population-based retrospective cohort study. *Epilepsia.* 52, 2036–2042. doi: 10.1111/j.1528-1167.2011.03268.x
- Chao, H. T., Chen, H., Samaco, R. C., Xue, M., Chahrour, M., Yoo, J., et al. (2010). Dysfunction in GABA signalling mediates autism-like stereotypies and Rett syndrome phenotypes. *Nature.* 468, 263–269. doi: 10.1038/nature09582
- Christiansen, J., Dyck, J. D., Elyas, B. G., Lilley, M., Bamforth, J. S., Hicks, M., et al. (2004). Chromosome 1q21.1 contiguous gene deletion is associated with congenital heart disease. *Circ. Res.* 94, 1429–1435. doi: 10.1161/01.res.0000130528.72330.5c
- Coe, B. P., Girirajan, S., and Eichler, E. E. (2012). The genetic variability and commonality of neurodevelopmental disease. *Am. J. Med. Genet. C. Semin. Med. Genet.* 160C, 118–129. doi: 10.1002/ajmg.c.31327

- Coghlan, S., Horder, J., Inkster, B., Mendez, M. A., Murphy, D. G., and Nutt, D. J. (2012). GABA system dysfunction in autism and related disorders: From synapse to symptoms. *Neurosci. Biobehav. Rev.* 36, 2044–2055. doi: 10.1016/j.neubiorev.2012.07.005
- Crooks, G. E., Hon, G., Chandonia, J. M., and Brenner, S. E. (2004). WebLogo: a sequence logo generator. *Genome Res.* 14, 1188–1190. doi: 10.1101/gr.849004
- Cui, Y., Costa, R. M., Murphy, G. G., Elgersma, Y., Zhu, Y., Gutmann, D. H., et al. (2008). Neurofibromin regulation of ERK signaling modulates GABA release and learning. *Cell.* 135, 549–560. doi: 10.1016/j.cell.2008.09.060
- de Kovel, C. G., Trucks, H., Helbig, I., Mefford, H. C., Baker, C., Leu, C., et al. (2010). Recurrent microdeletions at 15q11.2 and 16p13.11 predispose to idiopathic generalized epilepsies. *Brain.* 133, 23–32. doi: 10.1093/brain/awp262
- de Los Heros, P., Alessi, D. R., Gourlay, R., Campbell, D. G., Deak, M., Macartney, T. J., et al. (2014). The WNK-regulated SPAK/OSR1 kinases directly phosphorylate and inhibit the K⁺-Cl⁻ co-transporters. *Biochem. J.* 458, 559–573. doi: 10.1042/bj20131478
- de Rubeis, S., He, X., Goldberg, A. P., Poultney, C. S., Samocha, K., Cicek, A. E., et al. (2014). Synaptic, transcriptional and chromatin genes disrupted in autism. *Nature.* 515, 209–215. doi: 10.1038/nature13772
- Deidda, G., Bozarth, I. F., and Cancedda, L. (2014). Modulation of GABAergic transmission in development and neurodevelopmental disorders: investigating physiology and pathology to gain therapeutic perspectives. *Front. Cell. Neurosci.* 8:119. doi: 10.3389/fncel.2014.00119
- Durand, C. M., Betancur, C., Boeckers, T. M., Bockmann, J., Chaste, P., Fauchereau, F., et al. (2007). Mutations in the gene encoding the synaptic scaffolding protein SHANK3 are associated with autism spectrum disorders. *Nat. Genet.* 39, 25–27. doi: 10.1038/ng1933
- Dzhala, V. I., Talos, D. M., Sdrulla, D. A., Brumback, A. C., Mathews, G. C., Benke, T. A., et al. (2005). NKCC1 transporter facilitates seizures in the developing brain. *Nat. Med.* 11, 1205–1213. doi: 10.1038/nm1301
- EPGP Collaborative, Abou-Khalil, B., Alldredge, B., Bautista, J., Berkovic, S., Bluvstein, J., et al. (2013). The epilepsy phenome/genome project. *Clin. Trials.* 10, 568–586. doi: 10.1177/1740774513484392
- Fischbach, G. D., and Lord, C. (2010). The Simons Simplex Collection: a resource for identification of autism genetic risk factors. *Neuron.* 68, 192–195. doi: 10.1016/j.neuron.2010.10.006
- Gagnon, K. B., and Delpire, E. (2013). Physiology of SLC12 transporters: lessons from inherited human genetic mutations and genetically engineered mouse knockouts. *Am. J. Physiol. Cell. Physiol.* 304, C693–C714. doi: 10.1152/ajpcell.00350.2012
- Gauthier, J., Bonnel, A., St-Onge, J., Karemera, L., Laurent, S., Mottron, L., et al. (2005). NLGN3/NLGN4 gene mutations are not responsible for autism in the Quebec population. *Am. J. Med. Genet. B. Neuropsychiatr. Genet.* 132B, 74–75. doi: 10.1002/ajmg.b.30066
- Gauthier, J., Champagne, N., Lafrenière, R. G., Xiong, L., Spiegelman, D., Brustein, E., et al. (2010). De novo mutations in the gene encoding the synaptic scaffolding protein SHANK3 in patients ascertained for schizophrenia. *Proc. Natl. Acad. Sci. U. S. A.* 107, 7863–7868. doi: 10.1073/pnas.0906232107
- Gilman, S. R., Iossifov, I., Levy, D., Ronemus, M., Wigler, M., and Vitkup, D. (2011). Rare de novo variants associated with autism implicate a large functional network of genes involved in formation and function of synapses. *Neuron.* 70, 898–907. doi: 10.1016/j.neuron.2011.05.021
- Girard, S. L., Gauthier, J., Noreau, A., Xiong, L., Zhou, S., Jouan, L., et al. (2011). Increased exonic de novo mutation rate in individuals with schizophrenia. *Nat. Genet.* 43, 860–863. doi: 10.1038/ng.886
- Guilmare, A., Dubourg, C., Mosca, A. L., Legallie, S., Goldenberg, A., Drouin-Garraud, V., et al. (2009). Recurrent rearrangements in synaptic and neurodevelopmental genes and shared biologic pathways in schizophrenia, autism and mental retardation. *Arch. Gen. Psychiatry.* 66, 947–956. doi: 10.1001/archgenpsychiatry.2009.80
- Hamdan, F. F., Gauthier, J., Araki, Y., Lin, D. T., Yoshizawa, Y., Higashi, K., et al. (2011). Excess of de novo deleterious mutations in genes associated with glutamatergic systems in nonsyndromic intellectual disability. *Am. J. Hum. Genet.* 88, 306–316. doi: 10.1016/j.ajhg.2011.02.001
- Hamdan, F. F., Gauthier, J., Spiegelman, D., Noreau, A., Yang, Y., Pellerin, S., et al. (2009). Mutations in SYNGAP1 in autosomal nonsyndromic mental retardation. *N. Engl. J. Med.* 360, 599–605. doi: 10.1056/nejmoa0805392
- Hashimoto, T., Arion, D., Unger, T., Maldonado-Aviles, J. G., Morris, H. M., Volk, D. W., et al. (2008). Alterations in GABA-related transcriptome in the dorsolateral prefrontal cortex of subjects with schizophrenia. *Mol. Psychiatry.* 13, 147–161. doi: 10.1038/sj.mp.4002011
- He, Q., Nomura, T., Xu, J., and Contractor, A. (2014). The developmental switch in GABA polarity is delayed in fragile X mice. *J. Neurosci.* 34, 446–450. doi: 10.1523/jneurosci.4447-13.2014
- Heinzen, E. L., Radtke, R. A., Urban, T. J., Cavalleri, G. L., Depondt, C., Need, A. C., et al. (2010). Rare deletions at 16p13.11 predispose to a diverse spectrum of sporadic epilepsy syndromes. *Am. J. Hum. Genet.* 86, 707–718. doi: 10.1016/j.ajhg.2010.03.018
- Hekmat-Scafe, D. S., Lundy, M. Y., Ranga, R., and Tanouye, M. A. (2006). Mutations in the K⁺/Cl⁻ cotransporter gene *kazachoc (kcc)* increase seizure susceptibility in *Drosophila*. *J. Neurosci.* 26, 8943–8954. doi: 10.1523/jneurosci.4998-05.2006
- Hekmat-Scafe, D. S., Mercado, A., Fajilan, A. A., Lee, A. W., Hsu, R., Mount, D. B., et al. (2010). Seizure sensitivity is ameliorated by targeted expression of K⁺-Cl⁻ cotransporter function in the mushroom body of the *Drosophila* brain. *Genetics.* 184, 171–183. doi: 10.1534/genetics.109.109074
- Hubner, C. A., Stein, V., Hermans-Borgmeyer, I., Meyer, T., Ballanyi, K., and Jentsch, T. J. (2001). Disruption of KCC2 reveals an essential role of K-Cl cotransport already in early synaptic inhibition. *Neuron.* 30, 515–524. doi: 10.1016/s0896-6273(01)00297-5
- International-Schizophrenia-Consortium. (2008). Rare chromosomal deletions and duplications increase risk of schizophrenia. *Nature.* 455, 237–241. doi: 10.1038/nature07239
- Iossifov, I., O’roak, B. J., Sanders, S. J., Ronemus, M., Krumm, N., Levy, D., et al. (2014). The contribution of de novo coding mutations to autism spectrum disorder. *Nature.* 515, 216–221. doi: 10.1038/nature13908
- Iossifov, I., Ronemus, M., Levy, D., Wang, Z., Hakker, I., Rosenbaum, J., et al. (2012). De novo gene disruptions in children on the autistic spectrum. *Neuron.* 74, 285–299. doi: 10.1016/j.neuron.2012.04.009
- Irizarry, R. A., Ladd-Acosta, C., Wen, B., Wu, Z., Montano, C., Onyango, P., et al. (2009). The human colon cancer methylome shows similar hypo- and hypermethylation at conserved tissue-specific CpG island shores. *Nat. Genet.* 41, 178–186. doi: 10.1038/ng.298
- Kahle, K. T., Deeb, T. Z., Puskarjov, M., Silayeva, L., Liang, B., Kaila, K., et al. (2013). Modulation of neuronal activity by phosphorylation of the K-Cl cotransporter KCC2. *Trends Neurosci.* 36, 726–737. doi: 10.1016/j.tins.2013.08.006
- Kahle, K. T., Merner, N. D., Friedel, P., Silayeva, L., Liang, B., Khanna, A., et al. (2014). Genetically encoded impairment of neuronal KCC2 cotransporter function in human idiopathic generalized epilepsy. *EMBO Rep.* 15, 766–774. doi: 10.15252/embr.201438840
- Kaila, K., Price, T. J., Payne, J. A., Puskarjov, M., and Voipio, J. (2014). Cation-chloride cotransporters in neuronal development, plasticity and disease. *Nat. Rev. Neurosci.* 15, 637–654. doi: 10.1038/nrn3819
- Kalanithi, P. S. A., Zheng, W., Kataoka, Y., Difiglia, M., Grantz, H., Saper, C. B., et al. (2005). Altered parvalbumin-positive neuron distribution in basal ganglia of individuals with Tourette syndrome. *Proc. Natl. Acad. Sci. U S A* 102, 13307–13312. doi: 10.1073/pnas.0502624102
- Kang, J. Q., and Macdonald, R. L. (2009). Making sense of nonsense GABA(A) receptor mutations associated with genetic epilepsies. *Trends Mol. Med.* 15, 430–438. doi: 10.1016/j.molmed.2009.07.003
- Kiezun, A., Garimella, K., Do, R., Stitzel, N. O., Neale, B. M., McLaren, P. J., et al. (2012). Exome sequencing and the genetic basis of complex traits. *Nat. Genet.* 44, 623–630. doi: 10.1038/ng.2303
- Kirov, G., Grozeva, D., Norton, N., Ivanov, D., Mantripragada, K. K., Holmans, P., et al. (2009). Support for the involvement of large copy number variants in the pathogenesis of schizophrenia. *Hum. Mol. Genet.* 18, 1497–1503. doi: 10.1093/hmg/ddp043
- Klassen, T., Davis, C., Goldman, A., Burgess, D., Chen, T., Wheeler, D., et al. (2011). Exome sequencing of ion channel genes reveals complex profiles confounding personal risk assessment in epilepsy. *Cell.* 145, 1036–1048. doi: 10.3410/f.12207957.13373055
- Krumm, N., O’roak, B. J., Shendure, J., and Eichler, E. E. (2014). A de novo convergence of autism genetics and molecular neuroscience. *Trends Neurosci.* 37, 95–105. doi: 10.1016/j.tins.2013.11.005

- Krystal, J. H., and State, M. W. (2014). Psychiatric disorders: diagnosis to therapy. *Cell*. 157, 201–214. doi: 10.1016/j.cell.2014.02.042
- Kryukov, G. V., Shpunt, A., Stamatoyannopoulos, J. A., and Sunyaev, S. R. (2009). Power of deep, all-exon resequencing for discovery of human trait genes. *Proc. Natl. Acad. Sci. U. S. A.* 106, 3871–3876. doi: 10.1073/pnas.0812824106
- Lam, C. W., Yeung, W. L., Ko, C. H., Poon, P. M., Tong, S. F., Chan, K. Y., et al. (2000). Spectrum of mutations in the MECP2 gene in patients with infantile autism and Rett syndrome. *J. Med. Genet.* 37:E41. doi: 10.1136/jmg.37.12.e41
- Larkin, M. A., Blackshields, G., Brown, N. P., Chenna, R., Mcgettigan, P. A., McWilliam, H., et al. (2007). Clustal W and Clustal X version 2.0. *Bioinformatics*. 23, 2947–2948. doi: 10.1093/bioinformatics/btm404
- Lemonnier, E., Degrez, C., Phelep, M., Tyzio, R., Josse, F., Grandgeorge, M., et al. (2012). A randomised controlled trial of bumetanide in the treatment of autism in children. *Transl. Psychiatry*. 2:e197. doi: 10.1038/tp.2012.124
- Lemonnier, E., Robin, G., Degrez, C., Tyzio, R., Grandgeorge, M., and Ben-Ari, Y. (2013). Treating Fragile X syndrome with the diuretic bumetanide: a case report. *Acta Paediatrica*. 102, e288–e290. doi: 10.1111/apa.12235
- Lewis, D. A., Curley, A. A., Glausier, J. R., and Volk, D. W. (2012). Cortical parvalbumin interneurons and cognitive dysfunction in schizophrenia. *Trends Neurosci.* 35, 57–67. doi: 10.1016/j.tins.2011.10.004
- Medina, I., Friedel, P., Rivera, C., Kahle, K. T., Kourdougli, N., Uvarov, P., et al. (2014). Current view on the functional regulation of the neuronal K⁺-Cl⁻ cotransporter KCC2. *Front. Cell. Neurosci.* 8:27. doi: 10.3389/fncel.2014.00027
- Medrihan, L., Tantalaki, E., Aramuni, G., Sargsyan, V., Dudanova, I., Missler, M., et al. (2008). Early defects of GABAergic synapses in the brain stem of a MeCP2 mouse model of Rett syndrome. *J. Neurophysiol.* 99, 112–121. doi: 10.1152/jn.00826.2007
- Mefford, H. C., Cooper, G. M., Zerr, T., Smith, J. D., Baker, C., Shafer, N., et al. (2009). A method for rapid, targeted CNV genotyping identifies rare variants associated with neurocognitive disease. *Genome Res.* 19, 1579–1585. doi: 10.1101/gr.094987.109
- Mefford, H. C., Sharp, A. J., Baker, C., Itsara, A., Jiang, Z., Buysse, K., et al. (2008). Recurrent rearrangements of chromosome 1q21.1 and variable pediatric phenotypes. *N. Engl. J. Med.* 359, 1685–1699. doi: 10.1056/nejmoa0805384
- Mi, H., Lazareva-Ulitsky, B., Loo, R., Kejarawal, A., Vandergriff, J., Rabkin, S., et al. (2005). The PANTHER database of protein families, subfamilies, functions and pathways. *Nucleic. Acids. Res.* 33, D284–288. doi: 10.1093/nar/gki078
- Mitchell, K. J. (2011). The genetics of neurodevelopmental disease. *Curr Opin Neurobiol.* 21, 197–203. doi: 10.1002/9781118524947
- Neale, B. M., Kou, Y., Liu, L., Ma'ayan, A., Samocha, K. E., Sabo, A., et al. (2012). Patterns and rates of exonic de novo mutations in autism spectrum disorders. *Nature*. 485, 242–245. doi: 10.1038/nature11011
- O'Roak, B. J., Deriziotis, P., Lee, C., Vives, L., Schwartz, J. J., Girirajan, S., et al. (2011). Exome sequencing in sporadic autism spectrum disorders identifies severe de novo mutations. *Nat. Genet.* 43, 585–589. doi: 10.3410/f.10819956.11732055
- O'Roak, B. J., Vives, L., Girirajan, S., Karakoc, E., Krumm, N., Coe, B. P., et al. (2012). Sporadic autism exomes reveal a highly interconnected protein network of de novo mutations. *Nature*. 485, 246–250. doi: 10.3410/f.716397823.791302874
- Pescosolido, M. F., Yang, U., Sabbagh, M., and Morrow, E. M. (2012). Lighting a path: genetic studies pinpoint neurodevelopmental mechanisms in autism and related disorders. *Dialogues Clin. Neurosci.* 14, 239–252.
- Purcell, S., Cherny, S. S., and Sham, P. C. (2003). Genetic Power Calculator: design of linkage and association genetic mapping studies of complex traits. *Bioinformatics*. 19, 149–150. doi: 10.1093/bioinformatics/19.1.149
- Purcell, S. M., Moran, J. L., Fromer, M., Ruderfer, D., Solovieff, N., Roussos, P., et al. (2014). A polygenic burden of rare disruptive mutations in schizophrenia. *Nature*. 506, 185–190. doi: 10.1038/nature12975
- Puskarjov, M., Seja, P., Heron, S. E., Williams, T. C., Ahmad, F., Iona, X., et al. (2014). A variant of KCC2 from patients with febrile seizures impairs neuronal Cl⁻ extrusion and dendritic spine formation. *EMBO Rep.* 15, 723–729. doi: 10.1002/embr.201438749
- Roy, A., Kucukural, A., and Zhang, Y. (2010). I-TASSER: a unified platform for automated protein structure and function prediction. *Nat Protoc.* 5, 725–738. doi: 10.1038/nprot.2010.5
- Rubenstein, J. L. R. (2011). Annual Research Review: Development of the cerebral cortex: implications for neurodevelopmental disorders. *J. Child. Psychol. Psychiatry*. 52, 339–355. doi: 10.1111/j.1469-7610.2010.02307
- Salomons, G. S., Van Dooren, S. J., Verhoeven, N. M., Cecil, K. M., Ball, W. S., Degrauw, T. J., et al. (2001). X-linked creatine-transporter gene (SLC6A8) defect: a new creatine-deficiency syndrome. *Am. J. Hum. Genet.* 68, 1497–1500. doi: 10.1086/320595
- Sanders, S. J., Murtha, M. T., Gupta, A. R., Murdoch, J. D., Raubeson, M. J., Willsey, A. J., et al. (2012). De novo mutations revealed by whole-exome sequencing are strongly associated with autism. *Nature*. 485, 237–241. doi: 10.1038/nature10945
- Schmidt, M. J., and Mirnics, K. (2014). Neurodevelopment, GABA System Dysfunction and Schizophrenia. *Neuropsychopharmacology*. 40, 190–206. doi: 10.1038/npp.2014.95
- Schwarz, J. M., Rodelsperger, C., Schuelke, M., and Seelow, D. (2010). MutationTaster evaluates disease-causing potential of sequence alterations. *Nat. Methods*. 7, 575–576. doi: 10.1038/nmeth0810-575
- Stefansson, H., Rujescu, D., Cichon, S., Pietilainen, O. P., Ingason, A., Steinberg, S., et al. (2008). Large recurrent microdeletions associated with schizophrenia. *Nature*. 455, 232–236. doi: 10.1038/nature07229
- Talos, D. M., Sun, H., Kosaras, B., Joseph, A., Folkerth, R. D., Poduri, A., et al. (2012). Altered inhibition in tuberous sclerosis and type IIb cortical dysplasia. *Ann. Neurol.* 71, 539–551. doi: 10.1002/ana.22696
- Tanis, J. E., Bellemer, A., Moresco, J. J., Forbush, B., and Koelle, M. R. (2009). The potassium chloride cotransporter KCC-2 coordinates development of inhibitory neurotransmission and synapse structure in *Caenorhabditis elegans*. *J. Neurosci.* 29, 9943–9954. doi: 10.1523/jneurosci.1989-09.2009
- Tornberg, J., Voikar, V., Savilahti, H., Rauvala, H., and Airaksinen, M. S. (2005). Behavioural phenotypes of hypomorphic KCC2-deficient mice. *Eur. J. Neurosci.* 21, 1327–1337. doi: 10.1111/j.1460-9568.2005.03959.x
- Tuchman, R., and Rapin, I. (2002). Epilepsy in autism. *Lancet Neurol.* 1, 352–358. doi: 10.1007/978-1-4614-0361-6_24
- Tyzio, R., Nardou, R., Ferrari, D. C., Tsintsadze, T., Shahrokhi, A., Eftekhari, S., et al. (2014). Oxytocin-Mediated GABA Inhibition During Delivery Attenuates Autism Pathogenesis in Rodent Offspring. *Science*. 343, 675–679. doi: 10.1126/science.1247190
- Ullmann, R., Turner, G., Kirchoff, M., Chen, W., Tonge, B., Rosenberg, C., et al. (2007). Array CGH identifies reciprocal 16p13.1 duplications and deletions that predispose to autism and/or mental retardation. *Hum. Mutat.* 28, 674–682. doi: 10.1002/humu.20546
- Visser, L. E., De Ligt, J., Gilissen, C., Janssen, I., Stehouwer, M., De Vries, P., et al. (2010). A de novo paradigm for mental retardation. *Nat. Genet.* 42, 1109–1112. doi: 10.1038/ng.712
- Voineagu, I., Wang, X., Johnston, P., Lowe, J. K., Tian, Y., Horvath, S., et al. (2011). Transcriptomic analysis of autistic brain reveals convergent molecular pathology. *Nature*. 474, 380–384. doi: 10.1038/nature10110
- Woo, N. S., Lu, J., England, R., McClellan, R., Dufour, S., Mount, D. B., et al. (2002). Hyperexcitability and epilepsy associated with disruption of the mouse neuronal-specific K-Cl cotransporter gene. *Hippocampus*. 12, 258–268. doi: 10.1002/hipo.10014
- Xu, B., Roos, J. L., Dexheimer, P., Boone, B., Plummer, B., Levy, S., et al. (2011). Exome sequencing supports a de novo mutational paradigm for schizophrenia. *Nat. Genet.* 43, 864–868. doi: 10.1038/ng.902
- Zhu, L., Polley, N., Mathews, G. C., and Delpire, E. (2008). NKCC1 and KCC2 prevent hyperexcitability in the mouse hippocampus. *Epilepsy Res.* 79, 201–212. doi: 10.1016/j.eplepsyres.2008.02.005

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2015 Merner, Chandler, Bourassa, Liang, Khanna, Dion, Rouleau and Kahle. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution and reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.