

Research

Open Access

## Regulatory role of E-NTPase/E-NTPDase in $\text{Ca}^{2+}/\text{Mg}^{2+}$ transport via gated channel

Hans M Schreiber<sup>1</sup> and Subburaj Kannan\*<sup>1,2</sup>

Address: <sup>1</sup>Division of Gastroenterology, School of Medicine, University of Pennsylvania, Philadelphia, PA 19104, USA and <sup>2</sup>Departments of Microbiology and Immunology, School of Medicine, PO Box 25056, University of Texas Medical Branch, 300 University Boulevard, Galveston, Texas, 77550 USA

Email: Hans M Schreiber - skannan22@hotmail.com; Subburaj Kannan\* - skannan22@hotmail.com

\* Corresponding author

Published: 12 August 2004

Received: 31 May 2004

*Theoretical Biology and Medical Modelling* 2004, 1:3 doi:10.1186/1742-4682-1-3

Accepted: 12 August 2004

This article is available from: <http://www.tbiomed.com/content/1/1/3>

© 2004 Schreiber and Kannan; licensee BioMed Central Ltd.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Abstract

**Background:** E-NTPase/E-NTPDase is activated by millimolar concentrations of  $\text{Ca}^{2+}$  or  $\text{Mg}^{2+}$  with a pH optimum of 7.5 for the hydrolysis of extracellular NTP and NDP. It has been generally accepted that E-NTPase/E-NTPDase plays regulatory role in purinergic signalling, but other functions may yet be discovered.

**Results:** In this article it is proposed on the basis of published data that E-NTPase/E-NTPDase could play a role in the influx and efflux of  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  in vivo.

**Conclusions:** Attenuation of extracellular  $\text{Ca}^{2+}$  influx by rat cardiac sarcoplasmic anti-E-NTPase antibodies and oligomerization studies on mammalian CD39 conclusively point towards the existence of a new channel in the membrane. Further studies on these properties of the E-NTPase/E-NTPDase may provide detailed mechanisms and identify the potential patho-physiological significance.

### Background

The mechanism by which  $[\text{Ca}^{2+}]_i$  is increased in excitable cells differs from that obtaining in non-excitable cells. Excitable cells exhibit an action potential, a substantial general depolarization of the plasma membrane, in response to depolarizing stimuli; influx of  $\text{Ca}^{2+}$  occurs via plasma membrane  $\text{Ca}^{2+}$  channels and/or release from sarco (endo) plasmic reticulum via ryanodine-receptor  $\text{Ca}^{2+}$  channels which regulate the excitation – contraction coupling [1,2]. The factors that determine the extent of  $\text{Ca}^{2+}$  entry are (i) magnitude of the membrane potential and (ii) magnitude of the transmembrane  $\text{Ca}^{2+}$  gradient. These two factors also determine whether  $\text{Ca}^{2+}$  or  $\text{Mg}^{2+}$  enters and the time (probably milliseconds) that elapses

between channel opening and termination of  $\text{Ca}^{2+}$  or  $\text{Mg}^{2+}$  transport [3].

In non-excitable cells, the increase in  $[\text{Ca}^{2+}]_i$  results from influx of  $\text{Ca}^{2+}$  across the plasma membrane and  $\text{Ca}^{2+}$  release from the endoplasmic reticulum.  $\text{Ca}^{2+}$  release from the SER depends on the binding of inositol 1,4,5-triphosphate ( $\text{InsP}_3$ ) to its receptor  $\text{Ca}^{2+}$  channels, and also on  $\text{Ca}^{2+}$  binding to ryanodine receptor –  $\text{Ca}^{2+}$  channels.

$\text{Ca}^{2+}$  is removed from the cell by the following means. i: the sarco (endo) plasmic reticular  $\text{Ca}^{2+}$  pump ATPase (SERCA), which transports  $\text{Ca}^{2+}$  from the cytoplasm into the SER lumen (~70% of the activator  $\text{Ca}^{2+}$ ); ii: The plasma membrane  $\text{Ca}^{2+}$  pump ATPase (PMCA), which

exports  $\text{Ca}^{2+}$  across the plasma membrane ( $\sim 1\%$  of the activator  $\text{Ca}^{2+}$ ); **iii**: Mitochondrial  $\text{Ca}^{2+}$ -Uniporters (m $\text{Ca}^{2+}$  uniporters), which transport  $\text{Ca}^{2+}$  into mitochondria ( $\sim 1\%$  of the activator  $\text{Ca}^{2+}$ ); **iv**: the  $\text{Na}^+/\text{Ca}^{2+}$  exchanger (28% of the activator  $\text{Ca}^{2+}$ ). This last transport system is reversible but under normal physiological conditions, in the  $\text{Ca}^{2+}$  extrusion mode, it exhibits a stoichiometry of 3  $\text{Na}^+$  influx/1  $\text{Ca}^{2+}$  efflux [4].

$\text{Ca}^{2+}$  enters animal cells via **(i)** voltage-operated  $\text{Ca}^{2+}$  channels (VOCC), **(ii)** ligand gated non-specific cation channels (LGCCS), and **(iii)** stretch/receptor activated non-specific  $\text{Ca}^{2+}$  channels (RACC) [4,5]. A "receptor operated  $\text{Ca}^{2+}$  channel" (ROCC) is defined as a plasma membrane  $\text{Ca}^{2+}$  channel other than VOCC or RACC. VOCC opening depends on membrane depolarization, whereas RACC opening depends on both direct and indirect activation of membrane bound receptors. In contrast, ROCC opening depends solely on agonist-receptor interaction. It has also been suggested that mobile intracellular messengers such as elevated  $[\text{Ca}^{2+}]_i$  play a role in ROCC opening [5,6]. Different types of ROCC are activated (opened) by diverse cell signaling mechanisms such as ligand specificity, increase in  $[\text{Ca}^{2+}]_i$ , increase in  $[\text{cAMP}]_i$  [7] and activation/inactivation of specific trimeric G proteins [8].

Opening of  $\text{Ca}^{2+}$  channels must be a highly regulated event involving physical movement of channel components inclusive of the alteration in channel protein conformation; Also, an extracellular source of free energy ( $\Delta G$ ) could be of critical importance. This might be supplied by E-NTPase/E-NTPDase mediated hydrolysis of NTP/NDP. Co-ordination of this process might play a role in the opening of  $\text{Ca}^{2+}$  channels, independently of membrane depolarization or other factors.

The biochemical, structural, and functional properties of E-type nucleotidases have been covered in several excellent reviews: **i**. Extracellular metabolism [9]; **ii**. purine signalling [10,11]; **iii**. adhesion [12]; **iv**. transporter functions [13]; **v**. pathophysiology [14,15].

#### **Rationale for the proposed hypothesis: E-NTPase/E-NTPDase mediated $\text{Ca}^{2+}/\text{Mg}^{2+}$ transport**

It has been suggested that  $\text{Ca}^{2+}$  entry during the slow inward current in normal myocardium involves membrane-bound channels potentially controlled and/or regulated by metabolic energy transfer from unknown sources, though  $\text{Ca}^{2+}$  enters the cell down its concentration gradient [16]. Electrical stimulation and membrane phosphorylation by cAMP-dependent protein kinase have been shown to increase E-NTPase/E-NTPDase activity. Metal ions such as  $\text{Mn}^{2+}$ ,  $\text{Co}^{2+}$ ,  $\text{Ni}^{2+}$  and  $\text{La}^{2+}$  that attenuate  $\text{Ca}^{2+}$  influx also inhibit the E-NTPase. In the late stages of heart failure the E-NTPase is down regulated. Activation

of E-NTPase by various concentrations of  $\text{Ca}^{2+}$  has been shown to correlate linearly with cardiac contractile force development [17].

"Calcium paradox" is defined as irreversible functional and structural protein loss in the isolated heart that is first perfused with  $\text{Ca}^{2+}$ -free buffer and then reperfused with  $\text{Ca}^{2+}$ -containing buffer [18]. E-NTPase activity is highest during the initial phases of reperfusion, which might favour the initial  $\text{Ca}^{2+}$  influx that causes  $\text{Ca}^{2+}$  overload. During the later stages of reperfusion with  $\text{Ca}^{2+}$ -containing buffer there is a loss of E-NTPase activity. During mild stages of  $\text{Ca}^{2+}$  paradox, E-NTPase retains its function and continues to favour  $\text{Ca}^{2+}$  influx, resulting in the development of intracellular  $\text{Ca}^{2+}$  overloads. However, during severe stages of calcium paradox, impaired E-NTPase activity may contribute to irreversible failure of contractile force recovery [19].

To date there is no report describing the detailed mechanism of E-NTPase/E-NTPDase-mediated channel gating and its role in  $\text{Ca}^{2+}/\text{Mg}^{2+}$  transport. In this article an attempt is made to delineate the molecular mechanism of  $\text{Ca}^{2+}/\text{Mg}^{2+}$  transport, identifying the source of energy and the activation and termination of the process. The central issues are:

- How the metabolic energy from nucleotide hydrolysis is effectively utilized in channel opening;
- What stage of the opening/closing cycle requires energy;
- By what (probable) mechanism the proposed scheme is completed;
- How, if at all, homeostasis is affected

The current hypothetical proposal is set out in three sections with appropriate illustrations.

#### **Phase I: Activation**

identifies the evidence that leads to the current proposal and describes how the metabolic energy from nucleotide triphosphate hydrolysis is utilised to assemble a functional homo-oligomer of the E-NTPase/E-NTPDase, forming a channel that is subsequently opened.


#### **Phase II: Suggested: $\text{Ca}^{2+}/\text{Mg}^{2+}$ Transport**

Describes, with supporting evidence, how the energy released from  $[\text{NTP}]_o$  or  $[\text{NDP}]_o$  hydrolysis might be utilized for opening the channel formed by the homo-oligomeric E-NTPase/E-NTPDase.

### PHASE I: ACTIVATION

#### E-NTPase: ROLE AS A Ca<sup>2+</sup>/Mg<sup>2+</sup> TRANSPORTER VIA CHANNEL GATING:

Electrical Stimulation (see Fig 1;2;3-Mol.Cell.Biochem, 77;135-141(1987))



**Figure 1**

**Phase I: Activation.** Based on direct experimental evidence, suppose that in response to electrical stimuli, an increased phosphatidylinositol turnover leads to elevated intracellular phospholipid. This in turn could induce post-translational modification of the monomer subunits of E-NTPase/E-NTPDase in the intracellular milieu. Subsequently, the monomers are translocated to the membrane, depending on the tissue type(s) and functional requirement(s).

#### Phase III: Termination of the transport processes


outlines the intracellular and extracellular factors that would influence the termination of the Ca<sup>2+</sup>/Mg<sup>2+</sup> transport processes, and the experimental evidence obtained in favor of the whole proposal.

#### Phase I: Activation of E-NTPase/E-NTPDase and channel formation

Membrane depolarization could locally alter protein conformation. This in turn could potentially induce post-translational modification in the (intracellular) monomer subunits of the E-NTPase/E-NTPDase, followed by translocation to the membrane (depending on the tissue type(s) and functional requirement(s)) (Fig. 1). Fig. 2

shows the proposed functional state of the E-NTPase/E-NTPDase after oligomerization and assembly in the membrane to form a gated Ca<sup>2+</sup>/Mg<sup>2+</sup> channel. Fig. 3, indicates that the oligomerized E-NTPase/E-NTPDase is likely to possess sensors to control the opening and closing of the Ca<sup>2+</sup>/Mg<sup>2+</sup> channel gate. Fig. 4, represents an interior view of the E-NTPase/E-NTPDase in the functional state after oligomerization and assembly in the membrane.

Probable energy sources and other significant factors are as follows. The source of extracellular nucleotides could be spontaneous release from dead cells or exocytosis from live/damaged cells [20]. In ocular ciliary epithelial cells, ATP is released in hypotonic conditions, and this release


**Figure 2**

**Phase I: Activation.** Proposed model for E-NTPase/E-NTPDase in a functional state after oligomerization and assembly in the membrane, functioning as a gated channel.

is inhibited by NPPB (5-nitro-2-(3-phenyl propylamine benzoic acid), a potent inhibitor of CFTR (cystic fibrosis transmembrane receptor) and p-glycoprotein mediated ATP release [21]. On the other hand, the endogenous CD39 of oocytes transforms under hypertonic conditions to a conformation mediating ATP transport to the extracellular environment, either by exocytosis or by acting as an ion channel [22,23]. However, under what conditions (hyper- or hypotonic) might CD39 assume an extracellular nucleotide hydrolyzing activity; and under those conditions, can this property be coupled to ion influx? This question remains unanswered.

At normal physiological temperature in presence of divalent succinyl CoA, Con A mediates the oligomerization of E-NTPase monomers/dimers to form a holoenzyme with enhanced activity. Eosin iodoacetamide (EIAA), a fluorescein iodoacetamide that forms thioester bonds with cysteine at neutral pH, enhances chicken gizzard ecto-ATPase activity [24].

There are ten conserved cysteine residues in E-NTPase (with additional cysteine residues in the N-terminal region that are known to mediate disulfide bond formation, essential in oligomerization). CD39, an ecto- $\text{Ca}^{2+}/\text{Mg}^{2+}$  apyrase that hydrolyses ATP and ADP [25], forms tetramers and might act as a bivalent cation channel.



**Figure 3**

**Phase I: Activation.** The oligomerized E-NTPase/E-NTPDase would probably possess hypothetical sensors acting to open/close the gates.


However, the precise mechanism and functional properties are not known at present. CD39 expression is associated with ATP release; it was speculated that ATP release (along with drugs) into the extracellular milieu is followed by the hydrolysis of the extracellular nucleotides by CD39 [26].

Furthermore, native CD39 (ecto-ATP/Dase/apyrase) forms tetramers upon oligomerization. Loss of either of the two transmembrane domains of rat CD39 ecto-ATP/Dase impairs enzyme activity. It has been suggested that the functional (holoenzyme) E-NTPase/E-NTPDase is a homotrimer in mammals.

Differences in enzyme activity among different species have been attributed to variations in the interaction among the monomers resulting in homotrimeric holoenzyme formation (66 kDa-ATPase) [27]. It seems clear that changes in the conformation of the E-NTPase/E-NTPDase could mediate changes in the channel transport function.

**Phase II:  $Ca^{2+}/Mg^{2+}$  Transport**

Fig. 5a, illustrates the possible utilization of the energy released from  $[NTP]_o/[NDP]_o$  hydrolysis ( $-7.3 \text{ kcal mol}^{-1}$  or by formation of AMP,  $-10.9 \text{ kcal/mol}^{-1}$ ) for opening the channel formed by the homo-oligomeric E-NTPase/E-NTPDase. This channel is postulated to open and close in




**Figure 4**  
**Phase I: Activation.** Interior view of E-NTPase/E-NTPDase in a functional state in the membrane.

response to energy availability (Fig. 5b). Fig. 6A, is an artist's impression of the three-dimensional configuration of the E-NTPase/E-NTPDase in vivo.  $\text{Ca}^{2+}$  might enter the cell and excess  $\text{Mg}^{2+}$  might leave by the influx and efflux mechanisms depicted in Fig 6b.

The opening of the slow inward  $\text{Ca}^{2+}$  current channel in cardiac sarcolemma during the plateau phase of the action potential requires ATP [28]. Furthermore, protein kinase-A (PKA) dependent phosphorylation appears to mediate the increase in  $\text{Ca}^{2+}$  influx in hormonal modulation of that process [29]. A similar model has been proposed for sodium channels in nerve membranes, in which a cycle of phosphorylation and dephosphorylation is proposed for opening and closing [30].

Other corroborating evidence implicating E-NTPase in  $\text{Ca}^{2+}/\text{Mg}^{2+}$  transport via the gated channel is briefly summarised. Rat cardiac sarcolemmal E-NTPase has considerable sequence homology with the human platelet thrombospondin receptor CD36 [31]. An antibody directed against the purified E-NTPase blocked the increase in intracellular calcium concentration, implying that the E-NTPase plays an unknown but significant role in the delayed  $\text{Ca}^{2+}$  influx or  $\text{Mg}^{2+}$  efflux during the plateau phase of the action potential (Unpublished observation). Activation of E-NTPase by millimolar concentrations of  $\text{Ca}^{2+}$  and electrical stimulation is linearly related to the contractile force developed in the myocardium [32]. Gramicidin S inhibits the E-NTPase activity and it attenuates the slow channel efflux in perfused frog left ventricles.



**Figure 5**


**Phase II:  $\text{Ca}^{2+}/\text{Mg}^{2+}$  Transport.** (A) Free energy released from ATP hydrolysis by E-NTPase on the outer membrane surface would yield  $-7.3 \text{ kcal mol}^{-1}$  or by formation of AMP by E-NTPDase would yield  $-10.9 \text{ kcal mol}^{-1}$ . (B) The energy is utilized for opening the channel formed by the E-NTPase/E-NTPDase, by altering the conformation of the sensors. This altered conformation has an inherent channel-opening effect; loss of the energy source causes the sensors to revert to the resting state, which corresponds to channel closing.

Based on these observations, we propose that E-NTPase might be involved in providing energy for  $\text{Ca}^{2+}/\text{Mg}^{2+}$  influx-efflux in the cardiac sarcolemma, opening the channel formed by the E-NTPase/E-NTPDase protein by altering the conformation of the sensors. The altered channel sensor conformation opens the channel; loss of the energy source allows the sensors to revert to the resting state, which corresponds to channel closing.

There are at least two  $\text{Mg}^{2+}$  transport systems: (a) rapid transport down the concentration gradient and (b) efflux in low  $\text{Ca}^{2+}$  Ringer during ventricular perfusion in vitro. In rat liver mitochondria, 50 nM cAMP or 250  $\mu\text{M}$  ADP

induced rapid loss of 6 mmol of  $\text{Mg}^{2+}/\text{mg}$  protein coupled with the stimulation of ATP efflux. This effect was specific and was blocked by adenosine nucleotide translocase inhibitors. Evidently cAMP acts as a mobilizer of  $\text{Mg}^{2+}$  in isolated rat liver mitochondria. Adenine nucleotide translocase is the cAMP target [33].

Myocardial  $\text{Mg}^{2+}$  content is maintained at physiological level by the sarcolemmal transport system, which pumps  $\text{Mg}^{2+}$  across the plasma membrane when the extracellular  $[\text{Mg}^{2+}]_o$  concentration is  $<1 \text{ mM}$  and restores  $[\text{Mg}^{2+}]_i$  when the heart is perfused with Ringer buffer containing  $5 \times 10^{-7} \text{ M}$   $\text{Mg}^{2+}$ . Failure of either of these two transport



**Figure 6**  
**Phase II: Ca<sup>2+</sup>/Mg<sup>2+</sup> Transport.** (A) Three-dimensional impression of the E-NTPase/E-NTPDase in vivo. (B) It is possible that Ca<sup>2+</sup> can enter the cell and excess Mg<sup>2+</sup> can leave via the influx/efflux mechanisms depicted in the figure.

mechanisms may result in a rise in  $[Mg^{2+}]_i$ , impairing the contractile machinery of the myocardium [34].

Gramicidin S inhibits total  $Mg^{2+}$  efflux in the myocardium, while epinephrine restores  $Mg^{2+}$  efflux and contractile force development in the frog ventricle perfused with 10 mM  $Mg^{2+}$ . It should be pointed out that both E-NTPase activity and myocardial contraction and relaxation are inhibited by gramicidin S [35].

In the light of the evidence surveyed here, there would appear to be a significant functional role for activated E-NTPase in  $Ca^{2+}$  influx and  $Mg^{2+}$  efflux (or vice versa) in the myocardium.

### **Phase III: Termination of the transport process**

Fig. 7 summarizes the possible means by which the transport process is terminated. There are several potential contributing factors that can be grouped into two categories, extracellular and intracellular. Additional experimental evidence is indicated. Based on the heterologous expression of ecto-apyrase in COS cells in the presence of tunicamycin, glycosylation might be required for homooligomerization and nucleotidase activity. Conversely, deglycosylation might impair the E-type nucleotidase activity by weakening the monomer-monomer interaction and altering the tertiary and quaternary structures, result in the loss of holoenzyme. Essentially, glycosylation and deglycosylation of the ecto apyrase (HB6) monomer



## Potential contributing factors for the termination of $\text{Ca}^{2+}/\text{Mg}^{2+}$ transporter function via channel gating :

### Extracellular...

- i. Decreased extracellular nucleotide(s) concentration
- ii. Loss of free energy availability on the extracellular surface
- iii. Catalytically active ecto-kinase maintain the E-NTP level

### Intracellular...

- i. Increased intracellular  $\text{Ca}^{2+}/\text{Mg}^{2+}$  concentration
- ii. Alteration in intracellular pH
- iii. Deglycosylation of the E-NTPase holoenzyme
- iv. Increased activation of intracellular cholesterol oxidase

### Experimental Evidences...

- i. Verapamil mediated inhibition of E-NTPase activity and  $\text{Ca}^{2+}/\text{Mg}^{2+}$  influx
- ii. Attenuation of  $\text{Ca}^{2+}$  influx by anti-rat cardiac sarcolemmal  $\text{Ca}^{2+}/\text{Mg}^{2+}$  ectoATPase (IgG Fraction)

#### Figure 7


**Phase III: Termination of the transport processes.** (A) Several factors might contribute to the termination of  $\text{Ca}^{2+}/\text{Mg}^{2+}$  transport via channel gating by E-NTPase/E-NTPDase: extracellular and Intracellular. Additional experimental evidence is mentioned. Decreased flow of  $\text{Ca}^{2+}/\text{Mg}^{2+}$  due to closing of the channel gate.

and the consequences for homodimer formation have been regarded as an on-off switch for ecto nucleotidase activity [36].

Fig. 8a is a three-dimensional impression of the ecto-ATPase in vivo at the termination of ion transport. Fig. 8b illustrates how biochemical modifications such as deglycosylation of the E-NTPase/E-NTPDase oligomers might cause dissociation of the homo-oligomers to individual monomers. This is a potential mechanism for the disassembly of the functional channel and closure of  $\text{Ca}^{2+}$  influx and  $\text{Mg}^{2+}$  efflux. Also, an increase in membrane fluidity induced by cholesterol oxidation might cause defec-

tive association or disassociation due to weak interaction among the E-NTPase monomers, whereas increased membrane cholesterol might sustain higher E-NTPase activity. Oligomerization of E-NTPase and associated increase of activity could also be responsible for the rapid termination of the purinergic response mediated by extracellular ATP [37].

The extracellular nucleotide mediated activation of channel gating could be terminated by ecto (extracellular)-adenylate kinase, which catalyzes trans-phosphorylase activity ( $\text{ADP}+\text{ADP}\rightarrow\text{ATP}+\text{AMP}$ ). This enzyme has a higher affinity for extracellular nucleotides than the



**Figure 8**

**Phase III: Termination of the transport processes.** (A) Three-dimensional impression of the E-NTPase/E-NTPDase *in vivo* when termination of the ion transport function commences. (B) Biochemical modifications of the E-NTPase/E-NTPDase oligomers such as deglycosylation would probably cause instability, leading to dissociation of the homo-oligomers. Disassembly of the functional molecule would ensue, closing the  $\text{Ca}^{2+}$  influx and  $\text{Mg}^{2+}$  efflux processes, as portrayed in the figure.

dephosphorylating enzyme (E-NTPase/E-NTPDase) or ecto-nucleotide pyrophosphatase/phospho-diesterase ( $\text{ATP} \rightarrow \text{AMP} + \text{ppi}$ ) [38].

As the transport process winds down, ecto-adenylate kinase mediated ATP generation might maintain the extracellular nucleotide level. However, the precise biochemical kinetic process by which this process is completed remains to be elucidated [39].

#### **Pathophysiological Significance of E-type nucleotidase mediated $\text{Ca}^{2+}/\text{Mg}^{2+}$ transport**

Impairment of E-Type nucleotidases during  $\text{Ca}^{2+}$  paradox in isolated rat heart model warrants investigation of the

molecular mechanism(s) involved. Knowledge obtained from these studies will elucidate the observed protective effects of anti-rat cardiac  $\text{Ca}^{2+}/\text{Mg}^{2+}$ -ecto-ATPase antibodies in ischemia reperfusion induced damage, which is a corollary of organ transplantation. Furthermore, the anti-proliferative effect(s) of these antibodies in left anterior descending coronary artery smooth muscle cell(s) emphasize the need to explore more fully the hypothesis proposed in this article.

#### **Authors' contributions**

HMS participated and provided the hypothetical scheme of the gating mechanism with appropriate literature. SK

conceived and carried out experimental part of the investigation and formulating the hypotheses.

### Abbreviations

E-NTPase = Ecto or Extracellular Nucleotide triphosphatase; E-NTPDase = Ecto or Extracellular Nucleotide triphosphate diphosphohydrolase;  $[Ca^{2+}]_i$  = Intracellular  $Ca^{2+}$ ;  $[NTP]_0$  = Extracellular Nucleotide triphosphate;  $[NDP]_0$  = Extracellular Nucleotide diphosphate; ROCC = Receptor Operated  $Ca^{2+}$  channel; SER = Sarco (Endo) plasmic reticulum;  $[cAMP]_i$  = Cytoplasmic or intracellular cAMP; PMCA = Plasma membrane  $Ca^{2+}$ pump ATPase. VOCC = voltage-operated  $Ca^{2+}$  channel; LGCCS = Ligand gated non-specific cation channels. RACC = Stretch/Receptor activated non-specific  $Ca^{2+}$ channels. SUR = Sulfonylurea Receptor Proteins. CD36 = Thrombospondin receptor on platelets. CD39 = Ecto  $Ca^{2+}/Mg^{2+}$ apyrase.

### Acknowledgements

Data were obtained during graduate (Ph.D.) work (1993–1998) supported by a graduate fellowship (to S.K) from St. Boniface General Hospital Research Foundation, Winnipeg, Canada. T.R. Smith, Medicinal Chemistry, University of Warwick, Leamington Spa, and Warwickshire, United Kingdom prepared most of the models. This manuscript was prepared during the tenure of a post-doctoral fellowship (September 2000 – January 2001) supported by grants DK-52216 and DK-44237 from the National Institutes of Health, Bethesda, MD, and USA. T.L. Kirley, A.F. Knowles, L. Plesner, A.R. Beaudoin, A.Z. Herzberg, M. Handa, K.A. Jacobson, A. Froese, P. Zahradka, L.J. Murphy, N.N. Tandon, N. Abumrad, A. Ibrahim, R. Lipsky, D. Perlmutter, S.H. Lin, H. Zimmermann and N.N. Tandon are acknowledged for donating reagents, advises, and expert opinion on investigation during the period 1991–1998. Dr. Hans M. Schreiber has passed away during the preparation of this manuscript.

The authors do not have any competing financial or intellectual properties interests with CIBA-GEIGY Canada or Novartis Pharmaceuticals Inc, Switzerland. Modified reagents are part of the impending U.S. or International patent application(s).

### References

- Catterall WA: **Structure and function of voltage-gated ion channels.** *Annu Rev Biochem* 1995, **64**:493-531.
- Dunlap K, Luebke JI, Turner TJ: **Exocytotic  $Ca^{2+}$  channels in mammalian central neurons.** *Trends Neurosci* 1995, **18**:89-98.
- Rutecki PA: **Neuronal excitability: voltage-dependent currents and synaptic transmission.** *J Clin Neurophysiol* 1992, **9**:195-211.
- Clapham DE: **Calcium signaling.** *Cell* 1995, **80**:259-268.
- Berridge MJ: **Elementary and global aspects of calcium signalling.** *J Exp Biol* 1997, **200**:315-319.
- Putney JW Jr, Bird GS: **The inositol phosphate-calcium signaling system in nonexcitable cells.** *Endocr Rev* 1993, **14**:610-631.
- Applegate TL, Karjalainen A, Bygrave FL: **Rapid  $Ca^{2+}$  influx induced by the action of dibutylhydroquinone and glucagon in the perfused rat liver.** *Biochem J* 1997, **323**:463-467.
- Macrez-Lepretre N, Kalkbrenner F, Schultz G, Mironneau J: **Distinct functions of Gq and G11 proteins in coupling alpha1-adrenoceptors to  $Ca^{2+}$  release and  $Ca^{2+}$  entry in rat portal vein myocytes.** *J Biol Chem* 1997, **272**:5261-5268.
- Zimmermann H: **Nucleotides and cd39: principal modulatory players in hemostasis and thrombosis.** *Nat Med* 1999, **5**:987-988.

- Gendron FP, Benrezzak O, Krugh BW, Kong Q, Weisman GA, Beaudoin AR: **Purine signaling and potential new therapeutic approach: possible outcomes of NTPDase inhibition.** *Curr Drug Targets* 2003, **3**:229-245.
- Westfall DP, Todorov LD, Mihaylova-Todorova ST: **ATP as a cotransmitter in sympathetic nerves and its inactivation by releasable enzymes.** *J Pharmacol Exp Ther* 2002, **303**:439-444.
- Roberto Meyer-Fernandes J: **Ecto-ATPases in protozoa parasites: looking for a function.** *Parasitol Int* 2002, **51**:299-303.
- Kannan S: **E-NTPase /E-NTPDase: a potential regulatory role in E-kinase/PKA-mediated CD36 activation.** *Cell Biol Int* 2003, **27**:153-163.
- Marcus AJ, Broekman MJ, Drosopoulos JH, Islam N, Pinsky DJ, Sesti C, Levi R: **Metabolic control of excessive extracellular nucleotide accumulation by CD39/ecto-nucleotidase-1: implications for ischemic vascular diseases.** *J Pharmacol Exp Ther* 2003, **305**:9-16.
- Linden J: **Molecular approach to adenosine receptors: receptor-mediated mechanisms of tissue protection.** *Annu Rev Pharmacol Toxicol* 2001, **41**:775-787.
- Dhalla NS, Yates JC, Proveda V: **Calcium-linked changes in myocardial metabolism in the isolated perfused rat heart.** *Can J Physiol Pharmacol* 1977, **55**:925-933.
- Dhalla NS, Pierce GN, Panagia V, Singal PK, Beamish RE: **Calcium movements in relation to heart function.** *Basic Res Cardiol* 1982, **77**:117-139.
- Zimmerman AN, Hulsmann WC: **Paradoxical influence of calcium ions on the permeability of the cell membranes of the isolated rat heart.** *Nature* 1966, **211**:646-647.
- Alto L, Elimban VE, Lukas A, Dhalla NS: **Modification of heart sarcolemmal Na<sup>+</sup>/K<sup>+</sup>-ATPase activity during development of the calcium paradox.** *Mol Cell Biochem* 2000, **207**:87-94.
- Gordon JL: **Extracellular ATP: effects, sources and fate.** *Biochem J* 1986, **233**:309-319.
- Mitchell CH, Carre DA, McGlenn AM, Stone RA, Civan MM: **A release mechanism for stored ATP in ocular ciliary epithelial cells.** *Proc Natl Acad Sci U S A* 1998, **95**:7174-7178.
- Aleu J, Martin-Satue M, Navarro P, Lara IP, Bahima L, Marsal J, Solsona C: **Release of ATP induced by hypertonic solutions in Xenopus oocytes.** *J Physiol* 2003, **547**:209-219.
- Bodas E, Aleu J, Pujol G, Martin-Satue M, Marsal J, Solsona C: **ATP crossing the cell plasma membrane generates an ionic current in xenopus oocytes.** *J Biol Chem* 2000, **275**:20268-20273.
- Caldwell CC, Hornyak SC, Pendleton E, Campbell D, Knowles AF: **Regulation of chicken gizzard ecto-ATPase activity by modulators that affect its oligomerization status.** *Arch Biochem Biophys* 2001, **387**:107-116.
- Wang TF, Ou Y, Guidotti G: **The transmembrane domains of ectoapyrase (CD39) affect its enzymatic activity and quaternary structure.** *J Biol Chem* 1998, **273**:24814-24821.
- Abraham EH, Sterling KM, Kim RJ, Salikhova AY, Huffman HB, Crockett MA, Johnston N, Parker HW, Boyle WE Jr, Hartov A, Demidenko E, Efrid J, Kahn SA, Grubman DM, Jefferson J, Robson SC, Thakar JH, Lorico A, Rappa G, Sartorelli AC, Okunieff P: **Erythrocyte membrane ATP binding cassette (ABC) proteins: MRP1 and CFTR as well as CD39 (ecto-apyrase) involved in RBC ATP transport and elevated blood plasma ATP of cystic fibrosis.** *Blood Cells Mol Dis* 2001, **27**:165-180.
- Stout JG, Kirley TL: **Control of cell membrane ecto-ATPase by oligomerization state: intermolecular cross-linking modulates ATPase activity.** *Biochemistry* 1996, **35**:8289-8298.
- Sperelakis N, Schneider JA: **A metabolic control mechanism for calcium ion influx that may protect the ventricular myocardial cell.** *Am J Cardiol* 1976, **37**:1079-1085.
- Ziegelhoffer A, Anand-Srivastava MB, Khandelwal RL, Dhalla NS: **Activation of heart sarcolemmal  $Ca^{2+}/Mg^{2+}$  ATPase by cyclic AMP-dependent protein kinase.** *Biochem Biophys Res Commun* 1979, **89**:1073-1081.
- Schoffeniels E, Dandriofosse G: **Protein phosphorylation and sodium conductance in nerve membrane.** *Proc Natl Acad Sci U S A* 1980, **77**:812-816.
- Kannan S, Lalonde C, Zahradka P, Dhalla NS: **Molecular cloning of rat cardiac sarcolemmal  $Ca^{2+}/Mg^{2+}$  ectoATPase (Myoglein).** *J Mol Cell Cardiol* 1998, **30**:2261-2268.

32. Ziegelhoffer A, Dhalla NS: **Activation of Ca<sup>2+</sup>/Mg<sup>2+</sup> ATPase in heart sarcolemma upon electrical stimulation.** *Mol Cell Biochem* 1987, **77**:135-141.
33. Buchweitz O, Bianchi CP: **Myocardial magnesium transport: effect of gramicidin S and epinephrine.** *Life Sci* 1994, **55**:1853-1861.
34. Romani A, Dowell E, Scarpa A: **Cyclic AMP-induced Mg<sup>2+</sup> release from rat liver hepatocytes, permeabilized hepatocytes, and isolated mitochondria.** *J Biol Chem* 1991, **266**:24376-24384.
35. Bianchi CP, Liu D: **Calcium dependent magnesium uptake in myocardium.** *Life Sci* 1993, **52**:1225-1229.
36. Zhao D, Dhalla NS: **Characterization of rat heart plasma membrane Ca<sup>2+</sup>/Mg<sup>2+</sup> ATPase.** *Arch Biochem Biophys* 1988, **263**:281-292.
37. Smith TM, Kirley TL: **Glycosylation is essential for functional expression of a human brain ecto-apyrase.** *Biochemistry* 1999, **38**:1509-1516.
38. Stout JG, Kirley TL: **Control of cell membrane ecto-ATPase by oligomerization state: intermolecular cross-linking modulates ATPase activity.** *Biochemistry* 1996, **35**:8289-8298.
39. Picher M, Boucher RC: **Human airway ecto-adenylate kinase. A mechanism to propagate ATP signaling on airway surfaces.** *J Biol Chem* 2003, **278**:11256-12264.

Publish with **BioMed Central** and every scientist can read your work free of charge

*"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."*

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

