

# Relative contribution of monsoon precipitation and pumping to changes in groundwater storage in India

Akarsh Asoka<sup>1</sup>, Tom Gleeson<sup>2</sup>, Yoshihide Wada<sup>3,4,5,6</sup> and Vimal Mishra<sup>1\*</sup>

**The depletion of groundwater resources threatens food and water security in India. However, the relative influence of groundwater pumping and climate variability on groundwater availability and storage remains unclear. Here we show from analyses of satellite and local well data spanning the past decade that long-term changes in monsoon precipitation are driving groundwater storage variability in most parts of India either directly by changing recharge or indirectly by changing abstraction. We find that groundwater storage has declined in northern India at the rate of  $2\text{ cm yr}^{-1}$  and increased by  $1$  to  $2\text{ cm yr}^{-1}$  in southern India between 2002 and 2013. We find that a large fraction of the total variability in groundwater storage in north-central and southern India can be explained by changes in precipitation. Groundwater storage variability in northwestern India can be explained predominantly by variability in abstraction for irrigation, which is in turn influenced by changes in precipitation. Declining precipitation in northern India is linked to Indian Ocean warming, suggesting a previously unrecognized teleconnection between ocean temperatures and groundwater storage.**


Significant depletion of groundwater storage in a number of regions around the world, including northwest India<sup>1,2</sup>, has been shown with Gravity Recovery Climate Experiment (GRACE) observational data as well as global hydrologic and water use models<sup>3,4</sup>, and attributed to groundwater pumping (abstraction) for irrigation<sup>1,2,5,6</sup>. In India, irrigated agriculture produces over 70% of food grain, and groundwater plays a major role<sup>7</sup>, with annual groundwater abstraction increasing from  $10\text{--}20\text{ km}^3\text{ yr}^{-1}$  to  $240\text{--}260\text{ km}^3\text{ yr}^{-1}$  between 1950 and 2009<sup>8</sup>. India is a global leader in groundwater-fed irrigation due to intensive agriculture driven by multiple crops in a year<sup>9</sup>, especially after the green revolution<sup>1,2</sup>, with the largest non-renewable groundwater abstraction ( $68\text{ km}^3\text{ yr}^{-1}$ ) in the world<sup>7</sup>. Persistent droughts can reduce groundwater recharge and enhance groundwater pumping for irrigation, leading to lowered groundwater levels. For instance, due to a continuous deficit in precipitation,  $80\text{ km}^3$  of groundwater has been depleted in southern California since 1960<sup>5</sup>. Over the Gangetic Plain and other parts of north India, the monsoon season (June to September) precipitation has declined since 1950<sup>10–12</sup>, which has led to increased frequency and intensity of droughts<sup>13</sup>, possibly contributing to enhanced abstraction and/or reduced recharge of groundwater. Using multiple data sources (GRACE, well observations, model (PCR-GLOBWB<sup>14</sup>), precipitation, and sea surface temperature (SST)) and methods (regression and dominance analysis), we explore two related hypotheses: that precipitation deficit may have an impact on declining groundwater levels in northwestern India, which have previously been largely attributed to abstraction for irrigation<sup>2</sup>, and that groundwater storage variability may be partially associated with large-scale climate effects<sup>15</sup>, since weakening of the monsoon season precipitation is linked to large-scale climate variability<sup>10,12</sup>.

## Changes in groundwater storage

We estimated groundwater storage anomalies from GRACE for 2002–2013 to evaluate the spatial patterns of changes in groundwater in north and south India (Fig. 1). Consistent with previous analysis, and further supported for the first time by comparison to a large data set of water-level observations, GRACE groundwater anomalies show significant declines ( $2\text{ cm yr}^{-1}$ ,  $p\text{-value} < 0.05$ ) in the majority of north India in January, May, August, and November for which observations from Central Groundwater Board (CGWB) are available (Fig. 1a–d and Supplementary Fig. 3). Moreover, changes in groundwater anomalies from GRACE show increases ( $\sim 1\text{--}2\text{ cm yr}^{-1}$ , change in linear units) in south India (Fig. 1a–d and Supplementary Fig. 3). We find that changes in groundwater level from the observation wells and GRACE are consistent for 2002–2013 (Fig. 2e–h). However, GRACE-based estimates of trends are lower than those of observation wells, as GRACE examines larger spatial domains ( $\sim 100\text{ km}$  grid), whereas well observations are for point scale and represent very local depletion, which is not visible at GRACE resolution. However, standardized anomalies of groundwater level and GRACE-based groundwater storage change showed a close correspondence for north and south India, with correlation coefficients of **0.46** and **0.77**, respectively (Fig. 1i,j). GRACE groundwater anomalies show a large pattern of declining groundwater in north India, but increasing groundwater level in south India. However, it is unclear if these patterns of changes in groundwater anomalies in north and south India are driven by groundwater abstraction for irrigation or long-term changes in precipitation.

Previous studies<sup>1,2,11</sup> reported declines in groundwater storage in north India based on GRACE data, which are available for 2002 onwards; however, quantification of groundwater storage


<sup>1</sup>Civil Engineering and Earth Sciences, Indian Institute of Technology (IIT), Gandhinagar, India. <sup>2</sup>Department of Civil Engineering and School of Earth and Ocean Sciences, University of Victoria, Canada. <sup>3</sup>NASA Goddard Institute for Space Studies, New York, USA. <sup>4</sup>Center for Climate Systems Research, Columbia University, New York, USA. <sup>5</sup>Department of Physical Geography, Utrecht University, Utrecht, The Netherlands. <sup>6</sup>International Institute for Applied Systems Analysis, Laxenburg, Austria. \*e-mail: vmishra@iitgn.ac.in



**Figure 1 | Changes in groundwater storage from observation well and GRACE data during 2002–2013.** **a–h**, Monthly trends in groundwater anomaly are from GRACE (in  $\text{cm yr}^{-1}$ ) (**a–d**) and *in situ* well observations from the CGWB (**e–h**) for 2002–2013. Stippling in **a–d** indicates statistically significant changes at the 5% level. **e–h**, Wells that experienced significant declines and increases in groundwater levels ( $\text{cm yr}^{-1}$ ) during 2002–2013. Trends were estimated using the non-parametric Mann–Kendall test and Sen’s slope method. Monthly anomalies for January, May, August, and November were estimated from GRACE and *in situ* observations after removing the monthly mean. *In situ* groundwater well observations from the CGWB are available only for four months (January, May, August, and November). **i, j**, Area-averaged standardized departure (after removing mean and dividing by the standard deviation) from GRACE and *in situ* well observations for north (above  $23^\circ \text{N}$ ) and south (below  $23^\circ \text{N}$ ) India, respectively. Correlation coefficients between standardized anomalies of GRACE and groundwater wells for north and south India are 0.46 and 0.77, respectively.

1 variability in India beyond the GRACE period is limited. We  
 2 estimated changes (using linear trend) in the groundwater table  
 3 depth (m) using well observations from the CGWB for 1996–2013  
 4 and applied the non-parametric Mann–Kendall trend test and Sen’s

slope method. Moreover, we used the field significance test<sup>16</sup> to  
 5 evaluate trends at a regional scale considering the influence of spatial  
 6 and temporal correlations. Results show a significant decline ( $\sim 15$ –  
 7  $25 \text{ cm yr}^{-1}$ ,  $p$ -value  $< 0.05$ ) in groundwater table depth during  
 8




**Figure 2 | Changes in groundwater level in observation wells during 1996–2013 and their linkage with precipitation. a–d,** Observed trend in groundwater table for the months of January, May, August, and November for 1996–2013. Trends were estimated using the non-parametric Mann-Kendall trend test and Sen’s slope (wells that show statistical significant changes at the 5% level are shown). **e,f,** Relationship between standardized groundwater table anomaly and 12-month standardized precipitation index (SPI) for January, May, August, and November for northern India (above 23° N) and for southern India (below 23° N), respectively.

1 1996–2013 in a majority of observation wells located in north  
 2 India (23° north, Fig. 2a–d). Moreover, we find that the number  
 3 of wells with significant ( $p$ -value < 0.05) declines is higher for the  
 4 non-monsoon season than for the monsoon season, which may be  
 5 due to increased pumping during the non-monsoon season as it  
 6 is a major crop-growing period (Supplementary Fig. 2). In India,  
 7 the monsoon season overlaps with a major crop-growing season  
 8 (Kharif, June to September), in which groundwater pumping may  
 9 be high during monsoon deficit years. In the Rabi (October to  
 10 April) season, however, a majority of crops (for example, wheat)  
 11 mostly rely on groundwater-based irrigation. Observation wells


with significant water-level increases ( $\sim 5$ – $20$  cm yr<sup>-1</sup>) are mainly  
 located in south India, which is consistent with GRACE data (Fig. 1).  
 However, a minority of wells in each region show opposite trends  
 of decreasing groundwater levels in southern India and increasing  
 groundwater levels in northern India, highlighting the complexity  
 and heterogeneity of the data and localized influence of groundwater  
 pumping and recharge (Fig. 2).

Standardized groundwater level anomalies averaged over  
 northwest, north-central, and south India for all four months  
 (January, May, August, and November) represent annual variability  
 and show a close relationship (correlation coefficients 0.55, 0.54,



**Figure 3 | Changes in precipitation in irrigated and non-irrigated areas.** **a**, Changes in the monsoon season precipitation (mm) during 1980–2013. Changes were estimated using the Mann-Kendall trend test and Sen's slope method. **b**, Cumulative departure of precipitation from long-term mean (1980–2013) for 2002–2013. **c**, Area (%) irrigated with groundwater in India according to data obtained from the Food and Agricultural Organization (FAO). **d**, Areas irrigated with more than 40% contribution from groundwater (from **c**) and significantly increasing (blue) and decreasing (pink) precipitation during 1980–2013; red and blue dots represent locations of observation wells with significant trends in groundwater levels. **e–h**, Median trend in water-level change (m) in groundwater wells that are located in the region that experienced significant positive (blue bars, 63 wells) or negative changes (red bars, 170 wells) in precipitation and more than 40% area irrigated (as shown in **d**).


1 and 0.80, respectively) with the 12-month (Supplementary Table 1) 4  
 2 standardized precipitation index (SPI) for 1996–2013. Precipitation 5  
 3 deficit in north India influences soil moisture, groundwater 6  
 abstraction, and evaporative demands, as shown for the drought  
 year of 2009 (Supplementary Section 1 and Supplementary  
 Fig. 3). Evaporative stress index (ESI, ratio of evapotranspiration



**Figure 4 | Groundwater recharge from water-level observations and the PCR-GLOBWB model for 1996–2010. a,** Mean annual (climatology) groundwater recharge (cm) estimated using the water-table fluctuation method (see Methods for details) for 1996–2010. **b,** Same as **a**, but using recharge data from the PCR-GLOBWB model. **c,** Change (trend/year multiplied by the total duration (1996–2010)) in groundwater recharge for observation wells estimated using the non-parametric Mann–Kendall test and Sen’s slope method for 1996–2010. **d,** Same as **c**, but for the recharge estimates from the PCR-GLOBWB model.

1 (ET)/potential evapotranspiration (PET)) estimated using  
 2 Moderate Resolution Imaging Spectroradiometer (MODIS) satellite  
 3 data for 2002–2013 (Supplementary Fig. 2) shows a significant

increase during the post-monsoon season in the majority of  
 4 northern India, which may be due to increased groundwater  
 5 abstraction for irrigation as a precipitation contribution to  
 6



**Figure 5 | Linkage between groundwater storage variability and Indian Ocean SST. a,** Trend ( $\text{cm yr}^{-1}$ ) in annual groundwater anomaly from GRACE data for 2002–2013. The trend was estimated using the non-parametric Mann-Kendall test and Sen’s slope method. Stippling shows areas that experienced statistically significant increases/declines in annual groundwater anomaly. **b,** Leading mode (EOF-1) of variability obtained using the Empirical Orthogonal Function (EOF) analysis of the annual groundwater anomaly data from GRACE. **c,** Principal component (PC, PC-1) corresponding to the EOF-1. **d,** Correlation between the Indian Ocean SST and PC-1 for 2002–2013.

1 increased ET is less in the dry season (Supplementary Fig. 1i).  
 2 Moreover, positive SST anomalies (El-Niño) in the central Pacific  
 3 Ocean result in precipitation deficit in the monsoon season in  
 4 north and south India (Supplementary Table 6) and precipitation  
 5 deficit in 2002 and 2009 can be partially attributed to El-Niño.

6 **Precipitation and groundwater storage variability**

7 Groundwater storage could be affected by significant declines in the  
 8 monsoon season precipitation in India after 1950<sup>11–13</sup> if changes in  
 9 precipitation lead to changes in recharge or groundwater pumping.  
 10 Declines in the monsoon season precipitation have been observed  
 11 since 1950, and have continued during 1980–2013 (Fig. 3a,b).

Moreover, cumulative deficit in the monsoon season precipitation  
 showed substantial reductions in precipitation during 2002–2013  
 in north India (Fig. 3b). Long-term changes in precipitation  
 may affect groundwater storage in north India due to high  
 groundwater persistence, as groundwater levels respond slowly  
 to recharge (Supplementary Fig. 4). We notice that parts of the  
 Gangetic Plain, semi-arid western India (including Gujarat in  
 west-central India), and peninsular India are heavily irrigated  
 with groundwater (Fig. 3b). To evaluate the role of long-term  
 changes in precipitation on groundwater storage, we separated  
 the wells located in the regions with significant increases/declines  
 in precipitation (1980–2013) and heavily irrigated (more than

12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23

40% irrigation from groundwater) with groundwater (Fig. 3c,d). We find that wells that are located in the areas that witnessed significant increases in precipitation showed positive median trends in groundwater level (1996–2013) despite these wells being located in the area that is heavily irrigated with groundwater (Fig. 3d–h and Supplementary Fig. 5). On the other hand, wells that are located in the areas with significant declines in precipitation showed declines in groundwater tables, although there is a large variability in trends in both cases (Fig. 3e–h). The analysis was repeated for 2002–2013 with consistent results, suggesting that changes in precipitation substantially influence groundwater storage in India. Positive trends in groundwater storage change in south India are consistent with the long-term increase in precipitation<sup>10,12</sup>.

### Changes in groundwater recharge

We estimated annual groundwater recharge from well data using the water-table fluctuation method<sup>17</sup> and from the PCR-GLOBWB model. We found a substantial fluctuation in water-table depth in the observation wells during 1996–2010, which may be associated with the seasonal variability in precipitation and abstraction (Supplementary Figs 6 and 7). Consistent with model results, mean annual groundwater recharge estimated using the water-table fluctuation method for 1996–2010 showed high recharge in north-central India and Gujarat (Fig. 4a,b), primarily due to higher specific yields (Supplementary Fig. 8). Groundwater wells in north India are located in alluvial (unconsolidated sediment) plains, whereas wells in south India are primarily in bedrock (primarily consolidated sediment or igneous rock), which can affect the time for groundwater recharge in response to precipitation. Moreover, groundwater pumping can substantially reduce the well levels in the low-recharge areas, while in high-recharge areas, stream-aquifer interaction can also raise water levels<sup>15,16</sup>. A significant decline in precipitation in the north-central region (Supplementary Fig. 9) resulted in reduced groundwater recharge, as shown by both observation wells and model data (Fig. 4c,d). However, recharge in north and south India may be variable and not always directly related to precipitation. There might be other factors affecting groundwater recharge in India that are not considered in our analysis. For instance, groundwater systems have been modified by the large-scale canal network<sup>18</sup> for water diversions; however, the influence of canals and other surface water storage structures are not considered in our groundwater recharge estimates, which can be substantial in the drier parts of aquifers<sup>18</sup>. Water losses from unlined and lined canals can be substantial<sup>19</sup> in the areas where an extensive canal network is present (for example, the Gangetic Plain) contributing to groundwater recharge and water logging<sup>19</sup>. Since the area irrigated by groundwater wells in north and south India is far larger than that irrigated by canals (Supplementary Fig. 10), recharge from canals may not be sufficient to compensate groundwater declines due to abstraction<sup>20</sup>. Moreover, in north India (especially in the Indo-Gangetic Plain), the contribution of glacier melt to streamflow is within only 5–10% (refs 21,22); therefore, groundwater recharge due to stream-aquifer interactions may not be sufficient to balance the losses due to groundwater abstraction for irrigation in downstream regions.

### Relative importance of precipitation and abstraction

We analysed 12-month SPI and standardized abstraction index (SAI, estimated using recharge from the PCR-GLOBWB model) to investigate the relative contributions of precipitation and groundwater abstraction on changes in groundwater storage (Supplementary Fig. 11B). We separated observation wells located in northwest, north-central, and south India, which showed field significant declines (northwest and north-central) and increases (south) in groundwater level during 1996–2013 (Supplementary Fig. 11A). Long-term mean groundwater abstraction for 1996–2010

was substantially high ( $\sim 50 \text{ cm yr}^{-1}$ ) in northwestern India, which is consistent with the findings of Rodell and colleagues<sup>2</sup>. We found significant increases (5–10 cm) in groundwater abstraction in northwest India for 1996–2010, whereas significant declines in the monsoon season precipitation (Supplementary Fig. 9) and groundwater recharge (Fig. 4) can be noticed in the north-central India (Supplementary Fig. 9), indicating different driving factors such as the monsoon season precipitation, recharge, and groundwater abstraction in the northwest, north-central, and south India (Supplementary Fig. 9 and Fig. 4). Annual SPI and SAI are strongly related in northwest and south India, with correlation coefficients of  $-0.80$  ( $p$ -value  $< 0.05$ ) and  $-0.72$  ( $p$ -value  $< 0.05$ ), respectively. However, a relatively weaker (correlation =  $-0.46$ ,  $p$ -value  $< 0.05$ ) relationship between annual SPI and SAI was found in the north-central region (Supplementary Fig. 12 and Supplementary Table 1). Our results show that a precipitation deficit can lead to higher groundwater abstraction in India, as modelled abstraction is strongly related to precipitation (Supplementary Fig. 11B). Correlation between annual SAI and groundwater level anomalies from observation wells is strong in northwest and south India, with correlation coefficients of  $-0.62$  ( $p$ -value  $< 0.05$ ) and  $-0.55$  ( $p$ -value  $< 0.05$ ), respectively (Supplementary Fig. 12D,F). However, we did not find a strong relationship (correlation =  $0.31$ ) between groundwater abstraction and groundwater levels in north-central India.

Linear regression was performed for 1996–2010 using groundwater levels from observation wells, SPI and SAI to evaluate the relative importance (contribution) of precipitation and abstraction on groundwater variability. We find that SPI (12-month) explains 29, 30, and 64% of total groundwater storage variability in northwest, north-central, and south India, respectively (Supplementary Table 2). Annual groundwater abstraction (12 month-SAI) explains 38, 10, and 30% of total groundwater storage variability in northwest, north-central, and south India (Supplementary Table 2). However, looking at individual contributions (in total variability of groundwater storage) of annual (12-month) precipitation and abstraction, we find groundwater abstraction (SAI) explains more variability (38%) in northwest India, whereas SPI explains more variability in the north-central (30%) and south India (64%) (Supplementary Table 2). To understand if the groundwater abstraction is driven by precipitation in India, we estimated the fraction of total variability in annual abstraction (SAI) explained by annual precipitation (SPI). Our results showed that 65% variability of groundwater abstraction (SAI) in northwest India is explained by the annual precipitation (SPI), indicating that groundwater abstraction for irrigation is higher during precipitation deficit. It is important to note that about 35% of the variability of groundwater abstraction in northwestern India is contributed by other factors (such as choice of crops, intensive agriculture, subsidized electricity, and market driven prices). Moreover, the model results for abstraction may have a relatively higher uncertainty in the north-central region than in southern India<sup>33</sup>. We evaluated the relative importance of SPI and SAI using dominance analysis<sup>24</sup> to predict groundwater level anomalies because SPI and SAI are correlated. Results from linear regression and dominance analysis were consistent, indicating a larger role of SPI in groundwater storage variability in south and north-central India (Supplementary Tables 2 and 3). Similar to groundwater storage; we estimated the relative importance of SPI and SAI in groundwater recharge for all three regions. We found that annual precipitation (12-month SPI) explains 50, 91, and 83% of the total variability of annual groundwater recharge in northwestern, north-central, and south India (Supplementary Table 4). Our results from the regression and dominance analysis showed that the relative contribution from SPI in annual groundwater recharge is higher than SAI in all three regions (northwest, north-central,

and south) (Supplementary Table 4 and Supplementary Table 5), which further highlights the importance of the role of precipitation on groundwater recharge and abstraction in India.

Year-to-year variability in the monsoon season precipitation is linked to the large-scale climate<sup>10,12</sup>, suggesting large-scale climate may also influence groundwater variability in India. Annual changes in the groundwater anomaly from GRACE showed significant ( $\sim 2 \text{ cm yr}^{-1}$ ,  $p$ -value  $< 0.05$ ) declines in north India and increases in south India (Fig. 5). The leading mode obtained from the empirical orthogonal function (EOF-1), which explained about 46% of total squared covariance, exhibited a similar spatial structure to that obtained from trend analysis (Fig. 5a,b). The principal component (PC-1) of the leading mode obtained from the EOF analysis showed consistent declines during 2002–2013, indicating the leading mode represents the trend in groundwater anomaly. Negative correlation between PC-1 and SST anomalies indicates that warmer SST anomalies in the Indian Ocean result in declines in groundwater levels in northern India (Fig. 4d), which can be explained on the basis of the relationship between rainfall and SST<sup>12,13</sup>, and rainfall and groundwater levels, as shown above. Moreover, the ENSO affects the Indian monsoon rainfall in India<sup>10</sup>, which can also indirectly lead to enhanced warming over the Indian Ocean<sup>10,25</sup>. Consistent with previous studies<sup>10,26,27</sup>, we found that a positive SST anomaly over the central Pacific Ocean results in a similar impact (decline in precipitation) in north and south India (Supplementary Table 6), indicating that contrasting trends in groundwater storage in north and south India are more strongly linked to the SST variability in the Indian Ocean. The role of ENSO on groundwater storage variability, which affects SST over the Indian Ocean<sup>10,25,28</sup>, can be separated<sup>29</sup> if long-term GRACE data are available.

### Significance of untangling impacts on groundwater storage

Groundwater storage plays a key role in Indian agriculture, on which a large population rely directly or indirectly<sup>8</sup>. Although groundwater-based irrigated area has increased in northwest, north-central, and south India during 2002–2013 (Supplementary Table 7), contrasting trends in groundwater storage in north and south India highlight the importance of precipitation variability. Our results show that the contributions of anthropogenic pumping and precipitation to groundwater variability vary regionally in India—in north-central and south India precipitation is the major contributing factor, whereas in northwest India groundwater pumping is more important. We show that precipitation variability controls groundwater storage and recharge directly or indirectly in the majority of India, which has implications for water management in current and projected climate conditions<sup>30–32</sup>. Although groundwater-based irrigated area has increased in northwest, north-central, and south India (Supplementary Table 7), contrasting trends in groundwater storage in north and south India highlight the importance of precipitation variability. Importantly, other factors impacting groundwater storage (choice of crops, type of irrigation methods, intensive agriculture, subsidized electricity, and increasing trend in irrigated area) and groundwater recharge (aquifer characteristics<sup>16</sup>, depth of water table, presence of canals and surface storage structures<sup>23,33</sup>, pumping-induced recharge<sup>34</sup> and abstraction<sup>18</sup>, and stream–aquifer interaction<sup>16</sup> of glacier-fed rivers<sup>22</sup>) may affect the linkage between groundwater storage and precipitation in India. Moreover, several other factors related to irrigation practices and methods, uncertainties in recharge<sup>23,33–35</sup>, and management practices related to agriculture can influence variability of groundwater storage in the current and future climate<sup>28</sup>. For instance, improving irrigation methods (for example, sprinkler, drip) possibly reduces the return flow from irrigation to groundwater and baseflow, which may be another important factor for irrigation development and groundwater storage change

in India. Understanding the relative contribution from precipitation and anthropogenic pumping provides insight into better water management approaches for food and water security in India.

### Methods

Methods, including statements of data availability and any associated accession codes and references, are available in the online version of this paper.

Received 13 June 2016; accepted 2 December 2016; published online XX Month XXXX

### References

1. Tiwari, V. M., Wahr, J. & Swenson, S. Dwindling groundwater resources in northern India, from satellite gravity observations. *Geophys. Res. Lett.* **36**, L18401 (2009).
2. Rodell, M., Velicogna, I. & Famiglietti, J. S. Satellite-based estimates of groundwater depletion in India. *Nature* **460**, 999–1002 (2009).
3. Wada, Y., van Beek, L. P. H. & Bierkens, M. F. P. Nonsustainable groundwater sustaining irrigation: a global assessment. *Wat. Resour. Res.* **48**, W00L06 (2012).
4. Döll, P. & Siebert, S. Global modeling of irrigation water requirements. *Wat. Resour. Res.* **38**, 8–1–8–10 (2002).
5. Scanlon, B. R. *et al.* Groundwater depletion and sustainability of irrigation in the US high plains and central valley. *Proc. Natl Acad. Sci. USA* **109**, 9320–9325 (2012).
6. Taylor, R. G. *et al.* Evidence of the dependence of groundwater resources on extreme rainfall in East Africa. *Nat. Clim. Change* **3**, 374–378 (2013).
7. Gandhi, V. P. *et al.* *Groundwater Irrigation in India: Gains, Costs, and Risks* (Indian Institute of Management Ahmedabad, 2009).
8. Shah, T. Climate change and groundwater: India's opportunities for mitigation and adaptation. *Environ. Res. Lett.* **4**, 035005 (2009).
9. Siebert, S. *et al.* Groundwater use for irrigation—a global inventory. *Hydrol. Earth Syst. Sci.* **14**, 1863–1880 (2010).
10. Mishra, V., Smoliak, B. V., Lettenmaier, D. P. & Wallace, J. M. A prominent pattern of year-to-year variability in Indian summer monsoon rainfall. *Proc. Natl Acad. Sci. USA* **109**, 7213–7217 (2012).
11. Bollasina, M. A., Ming, Y. & Ramaswamy, V. Anthropogenic aerosols and the weakening of the South Asian summer monsoon. *Science* **334**, 502–505 (2011).
12. Roxy, M. K. *et al.* Drying of Indian subcontinent by rapid Indian Ocean warming and a weakening land–sea thermal gradient. *Nat. Commun.* **6**, 7423 (2015).
13. Vimal Mishra, R. S. Soil moisture droughts under the retrospective and projected climate in India. *J. Hydrometeorol.* **2267–2292** (2014).
14. Wada, Y., Wisser, D. & Bierkens, M. F. P. Global modeling of withdrawal, allocation and consumptive use of surface water and groundwater resources. *Earth Syst. Dyn.* **5**, 15–40 (2014).
15. Whittemore, D. O., Butler, J. J. Jr & Wilson, B. B. Assessing the major drivers of water-level declines: new insights into the future of heavily stressed aquifers. *Hydrol. Sci. J.* **61**, 134–145 (2016).
16. Yue, S. & Wang, C. Y. Regional streamflow trend detection with consideration of both temporal and spatial correlation. *Int. J. Climatol.* **22**, 933–946 (2002).
17. Government of India (Ministry of Water Resources): *Report of the Groundwater Resource Estimation Committee, Groundwater Resource Estimation Methodology* (2009).
18. MacDonald, A. M. *et al.* *Groundwater Resources in the Indo-Gangetic Basin: Resilience to Climate Change and Abstraction* OR/15/047 (British Geological Survey, 2015).
19. Basharat, M., Hassan, D., Bajkani, A. & Sultan, S. J. Surface water and groundwater Nexus: groundwater management options for Indus basin irrigation system. *Int. Waterlogging Salin. Res. Inst. IWASRI Lahore Pak. Water Power Dev. Auth. Publ. Vol. 155* (2014).
20. MacDonald, A. M. *et al.* Groundwater quality and depletion in the Indo-Gangetic Basin mapped from *in situ* observations. *Nat. Geosci.* **9**, 762–766 (2016).
21. Schaner, N., Voisin, N., Nijssen, B. & Lettenmaier, D. P. The contribution of glacier melt to streamflow. *Environ. Res. Lett.* **7**, 034029 (2012).
22. Immerzeel, W. W., Van Beek, L. P. & Bierkens, M. F. Climate change will affect the Asian water towers. *Science* **328**, 1382–1385 (2010).
23. Harvey, F. E. & Sibray, S. S. Delineating ground water recharge from leaking irrigation canals using water chemistry and isotopes. *Ground Water* **39**, 408–421 (2001).
24. Azen, R. & Budescu, D. V. The dominance analysis approach for comparing predictors in multiple regression. *Psychol. Methods* **8**, 129–148 (2003).
25. Roxy, M. K., Ritika, K., Terray, P. & Masson, S. The curious case of Indian Ocean warming. *J. Clim.* **27**, 8501–8509 (2014).



1	26. Kumar, K. K., Rajagopalan, B., Hoerling, M., Bates, G. & Cane, M. Unraveling the mystery of Indian monsoon failure during El Niño. <i>Science</i> <b>314</b> , 115–119 (2006).	25
2		26
3		
4	27. <del>Shah, H. L. &amp; Mishra, V. Hydrologic changes in Indian sub-continental river basins (1901–2012). <i>J. Hydrometeorol.</i> <b>2667–2687</b> (2016).</del>	
5		
6	28. Ashok, K., Guan, Z., Saji, N. H. & Yamagata, T. Individual and combined influences of ENSO and the Indian Ocean dipole on the Indian summer monsoon. <i>J. Clim.</i> <b>17</b> , 3141–3155 (2004).	
7		
8		
9	29. Compo, G. P. & Sardeshmukh, P. D. Removing ENSO-related variations from the climate record. <i>J. Clim.</i> <b>23</b> , 1957–1978 (2010).	
10		
11	30. Taylor, R. G. <i>et al.</i> Ground water and climate change. <i>Nat. Clim. Change</i> <b>3</b> , 322–329 (2013).	
12		
13	31. Fishman, R. M., Siegfried, T., Raj, P., Modi, V. & Lall, U. Over-extraction from shallow bedrock versus deep alluvial aquifers: reliability versus sustainability considerations for India's groundwater irrigation. <i>Wat. Resour. Res.</i> <b>47</b> , W00L05 (2011).	
14		
15		
16		
17	32. Döll, P. Vulnerability to the impact of climate change on renewable groundwater resources: a global-scale assessment. <i>Environ. Res. Lett.</i> <b>4</b> , 035006 (2009).	
18		
19		
20	33. Raza, A., Latif, M. & Shakir, A. S. Long-term effectiveness of lining tertiary canals in the Indus basin of Pakistan. <i>Irrig. Drain.</i> <b>62</b> , 16–24 (2013).	
21		
22	34. Shamsudduha, M., Taylor, R. G., Ahmed, K. M. & Zahid, A. The impact of intensive groundwater abstraction on recharge to a shallow regional aquifer system: evidence from Bangladesh. <i>Hydrogeol. J.</i> <b>19</b> , 901–916 (2011).	
23		
24		
	35. Wada, Y. <i>et al.</i> Global depletion of groundwater resources. <i>Geophys. Res. Lett.</i> <b>37</b> , L20402 (2010).	25
		26
	<b>Acknowledgements</b>	27
	The authors acknowledge funding from the ITRA-Water project. Data availability from the Central Groundwater Board (CGWB), Gravity Recovery and Climate Experiment (GRACE), and India Meteorological Department (IMD) is greatly appreciated.	28
		29
		30
	<b>Author contributions</b>	31
	V.M. conceived the idea. A.A. collected, analysed the data and developed the methodology. T.G. and Y.W. contributed to discussions of the findings. Y.W. provided groundwater recharge and abstraction data from the PCR-GLOBWB model. V.M. and A.A. wrote the manuscript with contributions from T.G. and Y.W.	32
		33
		34
		35
	<b>Additional information</b>	36
	Supplementary information is available in the <a href="#">online version of the paper</a> . Reprints and permissions information is available online at <a href="http://www.nature.com/reprints">www.nature.com/reprints</a> .	37
	Correspondence and requests for materials should be addressed to V.M.	38
		39
	<b>Competing financial interests</b>	40
	The authors declare no competing financial interests.	41

## 1 Methods

2 We used Gravity Recovery Climate Experiment (GRACE) groundwater anomaly,  
3 groundwater well observations from the Central Ground Water Board (CGWB),  
4 daily precipitation<sup>36</sup> from India Meteorological Department (IMD), and irrigated  
5 area map from the Food and Agriculture Organization (FAO) to understand the  
6 driving factors of groundwater variability in India. We derived the groundwater  
7 anomaly (GWA) at 1° spatial resolution after subtracting surface water storage  
8 (sum of soil moisture, canopy storage, and surface water) from GRACE terrestrial  
9 water storage anomaly (TWSA) for 2002–2013. Monthly TWSA version 05<sup>37,38</sup>  
10 was obtained from the Centre for Space Research (CSR) at the University of Texas,  
11 Austin. A 300 km Gaussian filter was applied to reduce the random errors in the  
12 data<sup>37</sup>. We applied scaling factors to minimize the attenuation caused due to  
13 sampling and post processing. We used monthly surface water storage from the  
14 Noah, CLM, VIC, and MOSAIC land surface models, which are available from the  
15 Global Land Data Assimilation System (GLDAS)<sup>39</sup>. The ensemble of groundwater  
16 anomalies based on GLDAS models (Noah, CLM, VIC, and MOSAIC) was used for  
17 the analysis. We used observations from more than 19,000 groundwater wells from  
18 CGWB, which are available for the months of January, May, August, and November  
19 (frequency of measurements is four times a year) for 1996–2013. However, there  
20 are significant data gaps and inconsistencies in the observed records. We selected  
21 groundwater observation wells for the analysis that have long-term data and are  
22 free from substantial missing data and inconsistencies. We finally selected 2,458  
23 wells with a minimum 15 yr (out of the entire record of 17 yr) of observations in  
24 each month (January, May, August, and September). Gridded daily precipitation<sup>36</sup>  
25 at 0.25° was obtained from IMD for 1980–2014. For areas outside India, monthly  
26 precipitation data were obtained from the Tropical Rainfall Measurement Mission  
27 (TRMM 3B43 V7)<sup>40</sup>. To understand groundwater variability in irrigated regions,  
28 the fraction of total area irrigated with groundwater was obtained from the 'Global  
29 Map of Irrigation Areas' (GMIA) version 5<sup>41</sup>. The water-table fluctuation method  
30 as suggested by the CGWB was used for recharge estimation using the difference  
31 between maximum and minimum depths (or fluctuation) of the water table at the  
32 observation wells and specific yield (recharge = fluctuation in water table ×  
33 specific yield). Specific yield for aquifers in India was obtained by digitizing a map  
34 provided by the CGWB<sup>42</sup>.

35 We used the satellite-derived volumetric soil moisture product from the  
36 European Space Agency Climate Change Initiative (ESACCI SMv02.1)<sup>43</sup> for  
37 1980–2013. Soil moisture trends using the ESACCI data were evaluated with the  
38 other products, and consistent results were found<sup>45</sup>. We estimated the evaporative  
39 stress index (ESI), which is a ratio of evapotranspiration (ET) and potential  
40 evapotranspiration (PET), using data from the Moderate Resolution Imaging  
41 Spectroradiometer (MODIS) monthly ET and PET (MOD16) products<sup>44</sup> at 5 km  
42 spatial resolution for 2002–2013. Higher ESI indicates water-stressed conditions as  
43 ET approaches PET.

44 We performed trend analysis based on the non-parametric Mann-Kendall  
45 trend test<sup>45</sup> with Sen's slope estimator<sup>46</sup>. For groundwater level anomalies, trends  
46 were estimated for each month for which observations were available to avoid the  
47 influence of seasonality. Spatial and temporal correlations in the data set in trend  
48 analysis were considered, and a field significance test was performed to evaluate  
49 changes at regional scales using the methodology described in Yue and Wang<sup>16</sup>.  
50 Please note that trend analysis can be influenced by the start and end year as well as  
51 length of the record. Changes for 2002–2013 in the selected variables were  
52 estimated by multiplying the trend slope by the number of years. Changes in  
53 groundwater anomaly from GRACE, groundwater table from the CGWB,  
54 precipitation, soil moisture, maximum temperature, and ESI were estimated for  
55 2002–2013 as well as for the other periods based on the data availability. To  
56 represent meteorological drought, the standardized precipitation index (SPI)<sup>47</sup> was  
57 used after fitting the Gamma distribution to monthly precipitation data. Similarly,  
58 we estimated the standardized abstraction index (SAI) using the model  
59 (PCR-GLOBWB) simulated abstraction data (in linear units) considering  
60 cumulative abstraction for a given period. For instance, an  $n$ -month SPI or SAI  
61 represents a standardized anomaly for cumulative precipitation or abstraction for  
62 the same period (that is,  $n$  months). Daily abstraction data were simulated from the  
63 PCR-GLOBWB<sup>44</sup> model for 1950–2013. The PCR-GLOBWB model simulates  
64 water storage for each grid cell at 0.5° spatial and daily temporal resolutions using  
65 two soil layers and an underlying groundwater layer. The model considers  
66 groundwater recharge from precipitation and irrigation water, while abstraction is  
67 estimated using requirements for irrigation and other sectors.

68 To evaluate the influence of climate variability and human intervention on  
69 groundwater, we selected the regions with increasing and declining trends in  
70 precipitation and significant groundwater-based irrigation. Areas were selected  
71 that are irrigated more than 40% with groundwater and have significant increasing  
72 or declining trends in the monsoon season rainfall. Groundwater wells falling in  
73 these areas were selected and their median trends were evaluated to understand if  
74 the monsoon season precipitation is a major driver of groundwater variability in  
75 India. To check consistency between groundwater anomalies from GRACE and  
76 observation wells, we used aggregated standardized departure fields (standardized  
77 anomaly) for wells located in northern (above 23°) and southern India (below 23°).

We used GRACE groundwater anomalies to estimate persistence (autocorrelation)  
for northern and southern India, which may influence estimated recharge rates  
during precipitation deficit years.

To evaluate the relative contribution of SPI and SAI on groundwater storage  
variability, linear regression was used. The relative contribution was estimated on  
the basis of the fraction of total variability ( $R^2$ ) in groundwater storage (represented  
by well level anomalies) explained by SPI or SAI. The relative contribution of SPI or  
SAI on groundwater storage variability was estimated by using just one of these  
(SPI or SAI) as a predictor of groundwater storage anomaly. The relative  
contribution of SPI and SAI was estimated for 3–24-month accumulation periods  
for precipitation and abstraction (using 3–24-month SPI for the same month for  
which groundwater anomaly was used) on groundwater storage anomaly. Similarly,  
we estimated the relative contribution of SPI and SAI on model-simulated  
groundwater recharge for north-central, northwestern, and south India. Since SPI  
and SAI may be correlated, we used dominance analysis<sup>24,48,49</sup> to estimate the  
relative importance of SPI and SAI on groundwater storage or recharge (where  
both are estimated using linear units rather than volumes). In dominance analysis  
the overall coefficient of determination ( $R^2$ ) of a predictor variable is computed  
after evaluating all the possible ( $p-1$ ) sub-models. The conditional dominance of a  
variable for each sub-model (0 to  $p-1$ ) is evaluated and the predictor with highest  
average conditional dominance is identified as the largest contributor<sup>24,48,49</sup>.

To evaluate the role of large-scale climate variability on groundwater, we used  
empirical orthogonal function (EOF) analysis<sup>10</sup> using GRACE groundwater  
anomaly for 2002–2013. The leading mode obtained from the EOF analysis  
(EOF-1) and the corresponding principal component (PC-1) were obtained.  
Correlation between the detrended PC-1 of annual groundwater anomalies from  
GRACE and SST anomalies over the Indian Ocean was estimated. SST data were  
obtained from the National Climatic Data Center's Extended Reconstructed SST  
(ERSSTv3b)<sup>50</sup>. We also estimated the correlation between precipitation in north  
and south India and the Nino 3.4 ENSO index.

**Data availability** The data used in the study are publicly available and can be  
directly obtained from the source websites. For instance, GRACE TWS data were  
obtained from JPL NASA ([ftp://podaac-ftp.jpl.nasa.gov/allData/tellus/L3/land\\_mass/RL05](ftp://podaac-ftp.jpl.nasa.gov/allData/tellus/L3/land_mass/RL05)). GLDAS surface water storage data are available from GSFC  
NASA (<http://disc.sci.gsfc.nasa.gov/services/grads-gds/gldas>). Data of satellite  
(TRMM) and gridded precipitation were obtained from GSFC NASA ([http://disc.gsfc.nasa.gov/ui/datasets/TRMM\\_3B43\\_7/summary](http://disc.gsfc.nasa.gov/ui/datasets/TRMM_3B43_7/summary)) and India Meteorological  
Department (IMD, <http://www.imd.gov.in/Welcome%20To%20IMD/Welcome.php>), respectively. Soil moisture data used in this study can be obtained from  
European Space Agency's Climate Change Initiative (ESACCI, <http://www.esa-soilmoisture-cci.org>). Satellite-based evapotranspiration from MODIS (MOD 16)  
can be obtained from the University of Montana (<http://www.natsg.umt.edu/project/mod16>). Groundwater well data from CGWB are available through Water  
Resources Information System of India (India-WARIS, <http://www.india-wris.nrsc.gov.in/wris.html>). Global Map of Irrigation Area (GMIA v5) can be obtained from  
Food and Agricultural Organization (FAO, <http://www.fao.org/nr/water/aquastat/irrigationmap>), while state-level irrigated area information can be obtained from  
India Stat (<http://www.indiastat.com/default.aspx>).

## References

36. Pai, D. S. *et al.* Development of a new high spatial resolution (0.25° × 0.25°)  
long period (1901–2010) daily gridded rainfall data set over India and its  
comparison with existing data sets over the region. *MAUSAM* **65**, 1–18 (2014).
37. Landerer, F. W. & Swenson, S. C. Accuracy of scaled GRACE terrestrial water  
storage estimates: ACCURACY OF GRACE-TWS. *Wat. Resour. Res.* **48**,  
W04531 (2012).
38. Swenson, S. & Wahr, J. Post-processing removal of correlated errors in GRACE  
data. *Geophys. Res. Lett.* **33**, L08402 (2006).
39. Rodell, M. *et al.* The global land data assimilation system. *Bull. Am. Meteorol.*  
*Soc.* **85**, 381–394 (2004).
40. Huffman, G. J. *et al.* The TRMM multisatellite precipitation analysis (TMPA):  
quasi-global, multyear, combined-sensor precipitation estimates at fine scales.  
*J. Hydrometeorol.* **8**, 38–55 (2007).
41. Siebert, S., Döll, P., Feick, S., Hoogeveen, J. & Frenken, K. *Global Map of  
Irrigation Areas* (Version, 2007).
42. *Central Groundwater Board, Ministry of Water Resources, Government of India:  
Aquifer Systems of India* (CGWB, 2012).
43. Dorigo, W. *et al.* Evaluating global trends (1988–2010) in harmonized  
multi-satellite surface soil moisture. *Geophys. Res. Lett.* **39**, L18405 (2012).
44. Mu, Q., Heinsch, F. A., Zhao, M. & Running, S. W. Development of a global  
evapotranspiration algorithm based on MODIS and global meteorology data.  
*Remote Sens. Environ.* **111**, 519–536 (2007).
45. Mann, H. B. Nonparametric tests against trend. *Econ. J. Econ. Soc.*  
**245**–259 (1945).
46. Sen, P. K. Estimates of the regression coefficient based on Kendall's tau. *J. Am.*  
*Stat. Assoc.* **63**, 1379–1389 (1968).

Q.14

Q.15

- 1 47. McKee, T. B. *et al.* The relationship of drought frequency and duration to time  
2 scales. *Proc. 8th Conf. Appl. Clim.* Vol. 17, 179–183 (American Meteorological  
3 Society, 1993).
- 4 48. Budescu, D. V. Dominance analysis: a new approach to the problem of relative  
5 importance of predictors in multiple regression. *Psychol. Bull.* **114**,  
6 542–551 (1993).
49. Nimon, K. F. & Oswald, F. L. Understanding the results of multiple linear  
7 regression beyond standardized regression coefficients. *Organ. Res. Methods*  
8 **16**, 650–674 (2013).  
9
50. Smith, T. M., Reynolds, R. W., Peterson, T. C. & Lawrimore, J. Improvements to  
10 NOAA's historical merged land-ocean surface temperature analysis  
11 (1880–2006). *J. Clim.* **21**, 2283–2296 (2008).  
12