

## RELATIVE NULLITY FOLIATIONS AND INDEFINITE ISOMETRIC IMMERSIONS

KINETSU ABE AND MARTIN MAGID

**This paper investigates the relative nullity distribution of an indefinite Riemannian manifold isometrically immersed into an indefinite space form.**

**Introduction.** In this paper we investigate the relative nullity distribution of an indefinite Riemannian manifold isometrically immersed into an indefinite space form. This distribution is totally geodesic and gives rise to a Riccati-type differential equation along a geodesic in a leaf of the distribution.

This differential equation is applied in several ways to estimate the index of relative nullity  $\nu$  for geodesically complete, connected, Lorentzian submanifolds  $M_1^n$  of  $\tilde{M}_1^{n+1}(c)$ , the Lorentzian sphere. These applications extend the work of Abe [1], [2], [3], Ferus [7], [8], and others to the setting of indefinite manifolds. Some of the work in §2 was obtained previously by Graves [10] in the codimension one case and by M. Dajczer. In particular Theorem 2 was conjectured by Dajczer [5].

Sections 1 and 2 lay the groundwork and derive the Riccati-type differential equation. In §3 an integer  $\nu_n$  is defined and it is shown that if  $M_1^n$  is as above and if  $\nu > \nu_n$  then  $M_1^n$  is totally geodesic. This integer is used to formulate a geometric condition which guarantees that a complete connected hypersurface of  $S_1^n(c)$  is totally geodesic. We also estimate  $\nu$  given a natural condition on the space-like Ricci curvature of the submanifold. In [6] other conditions on Ricci curvature are given.

The general scheme of our investigation is very similar to that of the Riemannian case as formulated in the papers mentioned above. However, there are a few basic and non-trivial differences from the Riemannian case. These differences are due to the indefinite metric and are to be overcome. Therefore, we think it worthwhile to include the details of the proofs for most of our results.

**1. Preliminaries.** An indefinite Riemannian manifold  $M_s^n$  of dimension  $n$  is a connected manifold with a non-degenerate metric in each

tangent space. The metric can be written

$$ds^2 = -(dx^1)^2 - \dots - (dx^s)^2 + (dx^{s+1})^2 + \dots + (dx^n)^2$$

at every point of  $M^n$ . In this case, we say the signature of  $M^n$  is  $(s, n - s)$  and write  $M_s^n$ .

If  $f: M_s^n \rightarrow \tilde{M}_t^{n+k}$  is an immersion and the metric induced on  $M^n$  is non-degenerate,  $f$  is an isometric immersion of  $M_s^n$  with this metric into  $\tilde{M}_t^{n+k}$ . Denote by  $D$  the torsion-free metric connection on  $\tilde{M}_t^{n+k}$ .  $D$  induces a torsion-free, metric connection  $\nabla$  on  $M$  as follows:

$$(1.1) \quad D_X Y = f_*(\nabla_X Y) + \alpha(X, Y),$$

where  $X$  and  $Y$  are tangent vectors on  $M$ ,  $f_*(\nabla_X Y)$  is the tangential component and  $\alpha(X, Y)$  is the normal component.  $\alpha$  is called the second fundamental form of  $f$ .

Given a field of unit normal vectors  $\xi$  on  $f(M_s^n)$  we can define a field of endomorphisms  $A_\xi$  on  $M$  by

$$(1.2) \quad D_X \xi = -f_*(A_\xi X) + \nabla_X^\perp \xi,$$

where  $-f_*(A_\xi X)$  is the tangential component.  $A_\xi$  is called the shape operator associated to  $\xi$ .

We denote by  $N(x)$  the set of all normal vectors to  $f(M_s^n)$  at  $f(x)$ . The metric on  $\tilde{M}_t^{n+k}$  and  $M_s^n$  is denoted by  $\langle \cdot, \cdot \rangle$ . As usual,  $R(\cdot, \cdot)$  denotes the curvature tensor of  $M_s^n$ . For the sake of future use, we list the following:

$$(1.3) \quad R(X, Y)Z = \tilde{R}(X, Y)Z + \sum_{p=1}^k \langle \xi_p, \xi_p \rangle \left[ (A_{\xi_p} X) \wedge (A_{\xi_p} Y) \right] Z$$

$$= \tilde{R}(X, Y)Z + A_{\alpha(Y, Z)}X - A_{\alpha(X, Z)}Y$$

(Gauss equation);

$$(1.4) \quad S(X, Y) = \sum_{i=1}^n \langle X_i, X_i \rangle \langle R(X_i, X)Y, X_i \rangle \quad (\text{Ricci tensor});$$

$\tilde{R}(\cdot, \cdot)$  is the curvature tensor of  $\tilde{M}_t^{n+k}$ ;  $\xi_p$ 's form an orthonormal base for  $N(x)$ ; and  $X_i$ 's form an orthonormal base for  $T_x M$ , the tangent space of  $M_s^n$  at  $x$ .

**2. Relative nullity.** If  $f: M_s^n \rightarrow \tilde{M}_t^{n+k}$  is an isometric immersion between indefinite Riemannian manifolds we define the relative nullity space at  $x$ ,  $T^0(x)$ , to be

$$(2.1) \quad T^0(x) = \{ X \in T_x(M) : A_\xi X = 0 \quad \forall \xi \in N(x) \}.$$

The orthogonal complement  $[T^0(x)]^\perp$  of  $T^0(x)$  in  $T_x M$  is denoted by  $T^1(x)$ .

**PROPOSITION 1.**  $T^0(x) = \{X \in T_x(M) : \alpha(X, Y) = 0 \forall Y \in T_x(M)\}$ .

*Proof.* It is obvious.

**PROPOSITION 2.**  $T^1(x) = \text{span}\{A_\xi Y\}$  for  $\xi \in N(x)$ ,  $Y \in T_x(M)$ .

*Proof.* Given any  $\xi \in N(x)$ ,  $Y \in T_x(M)$  and  $X \in T^0(x)$ ,  $\langle X, A_\xi Y \rangle = 0$ , so  $A_\xi Y \in T^1(x)$ .

On the other hand, suppose  $Z \in T_x(M)$  satisfies  $\langle Z, A_\xi Y \rangle = 0$  for all  $\xi, Y$  as above. Then  $\alpha(Y, Z) = 0$  for all  $Y$  and  $Z \in T^0(x)$ . This means that  $[\text{span}\{A_\xi Y\}]^\perp \subset T^0(x)$  so that  $[\text{span}\{A_\xi Y\}] \supset T^1(x)$ .

The dimension  $\nu(x)$  of  $T^0(x)$  is called the relative nullity of the immersion at  $x$ . The minimum value of  $\nu(x)$  on  $M$  is called the index of relative nullity and is denoted by  $\nu_0$ .

**THEOREM 1.** *Assume that  $\tilde{M}$  is a space form and let  $G$  denote the set of points in  $M$  where  $\nu(x) = \nu_0$ . Then*

- (1)  $G$  is an open subset of  $M$ ;
- (2)  $x \rightarrow T^0(x)$ ,  $x \in G$  is a differentiable and involutive distribution in  $G$ ;
- (3) the foliation  $T^0$  is totally geodesic in  $M$ ; and
- (4) each leaf of  $T^0$  is immersed as a totally geodesic submanifold of  $\tilde{M}$ .

*Proof.* (1) Pick a point  $x_0 \in G$  and a basis  $T_{x_0}(M)$ ,  $\{Y_1, \dots, Y_{\nu_0}, Y_{\nu_0+1}, \dots, Y_n\}$ , such that, for some  $\xi_j(x_0)$ ,  $\{A_{\xi_j} Y_{\nu_0+j}\}$  forms a basis of  $T^1(x_0)$ . Extend  $Y_1, \dots, Y_n$  and  $\xi_1, \dots, \xi_{n-\nu_0}$  smoothly in a neighborhood of  $x_0$ . The set  $\{A_{\xi_j} Y_{\nu_0+j}\}$  remains linearly independent in a neighborhood of  $x_0$ . Therefore in a neighborhood of  $x_0$  the dimension of  $T^0(x_0)$  must be less than or equal to  $\nu_0$ , and so equals  $\nu_0$ .

(2) It can be shown that  $T^0$  is a smooth distribution on  $G$  by noting that  $T^0 = \bigcap_{j=1}^k \ker A_{\xi_j}$  and that this intersection has constant rank on  $G$ .

(3) We use Codazzi's equation to see that  $T^0$  is totally geodesic. Let  $Y, Z$  be vector fields in  $T^0$ . For all  $X$  in  $TM$  and normal vectors  $\xi$  we have

$$\nabla_X(A_\xi Y) - A_\xi(\nabla_X Y) - A_{\nabla_X \xi} Y = \nabla_Y(A_\xi X) - A_\xi(\nabla_Y X) - A_{\nabla_Y \xi} X.$$

This reduces to

$$-A_\xi(\nabla_X Y) = \nabla_Y(A_\xi X) - A_\xi(\nabla_Y X) - A_{\nabla_Y^\perp \xi} X.$$

Taking the inner product of both sides with  $Z$  yields  $0 = \langle \nabla_Y(A_\xi X), Z \rangle$ .  $\langle A_\xi X, Z \rangle = 0$  can be differentiated in the direction of  $Y$ .

$$0 = Y\langle A_\xi X, Z \rangle = \langle \nabla_Y(A_\xi X), Z \rangle + \langle A_\xi X, \nabla_Y Z \rangle.$$

This gives  $\langle A_\xi X, \nabla_Y Z \rangle = 0$ , i.e.,  $\nabla_Y Z \in T^0$ . This shows  $T^0$  is totally geodesic. Examining Codazzi's equation with  $Y$  in  $T^0$  and  $\xi, X$  as above gives

$$-A_\xi(\nabla_X Y) = \nabla_Y(A_\xi X) - A_\xi(\nabla_Y X) - A_{\nabla_Y^\perp \xi} X$$

so that

$$\nabla_Y(A_\xi X) = A_\xi(\nabla_Y X) + A_{\nabla_Y^\perp \xi} X - A_\xi(\nabla_X Y).$$

This shows that  $T^1$  is also parallel along  $T^0$ .

For (4), notice that

$$D_Y f_*(Z) = f_*(\nabla_Y Z) + \alpha(Y, Z) = f_*(\nabla_Y Z). \quad \square$$

Next we define a complementary distribution  $T^c(x)$  in a neighborhood of  $y(t)$ ,  $t \in [a, b)$  where  $y(t)$  is a geodesic in a leaf of  $T^0$ . The distribution is complementary in the sense that  $T^c(x) \oplus T^0(x) = T_x(M)$ .

If, at a fixed point  $y(0)$ ,  $T^0(y(0))$  is non-degenerate then  $T^0(x)$  is non-degenerate for all points  $x$  near  $y(0)$ . Along a geodesic  $y(t)$  in a leaf of  $T^0$ ,  $T^0(y(t))$  remains non-degenerate since  $T^0(y(t))$  is parallel along the geodesic. Therefore,  $T^0(x)$  is non-degenerate in a neighborhood of the geodesic. In this case set  $T^c(x) = T^1(x)$ .

If  $T^0(y(0))$  is degenerate we use the following procedure. At  $y(0)$  choose a pseudo-orthonormal basis

$$\{L_1(0), \dots, L_r(0), E_1(0), \dots, E_{\nu_0-r}(0)\} \text{ of } T^0(y(0))$$

and

$$\{L_1(0), \dots, L_r(0), F_1(0), \dots, F_{n-\nu_0-r}(0)\} \text{ of } T^1(y(0)),$$

so that

$$\langle L_i(0), L_j(0) \rangle = 0 = \langle L_i(0), E_k(0) \rangle = \langle L_i(0), F_l(0) \rangle$$

and the  $E_k(0)$  and  $F_l(0)$  form an orthonormal set. Add  $\{\hat{L}_1(0), \dots, \hat{L}_r(0)\}$  so that each  $\hat{L}_i(0)$  is perpendicular to  $E_k(0)$  and  $F_l(0)$ ,  $\langle \hat{L}_i(0), \hat{L}_j(0) \rangle = 0$  and  $\langle L_i(0), \hat{L}_j(0) \rangle = -\delta_{ij}$ .

Denote the parallel extension of this basis

$$\{L_1(0), \dots, L_r(0), E_1(0), \dots, E_{v_0-r}(0), \\ F_1(0), \dots, F_{n-v_0-r}(0), \hat{L}_1(0), \dots, \hat{L}_r(0)\}$$

along  $y(t)$  by  $\{L_1(t), \dots, \hat{L}_r(t)\}$ .

Assume, without loss of generality, that  $\bar{y}(0)$  is one of

$$\{L_1(0), \dots, L_r(0), E_1(0), \dots, E_{v_0-r}(0)\},$$

say  $E_{v_0-r}(0)$ . Generalizing an argument in [10], define  $h: \mathbf{R}^n \rightarrow M^n$  by

$$h(t, x_1, \dots, x_r, y_1, \dots, y_{v_0-r-1}, u_1, \dots, u_{n-v_0-r}, v_1, \dots, v_r) \\ = \exp_{y(t)}(\sum x_j L_j(t) + \sum y_k E_k(t) + \sum u_l F_l(t) + \sum v_j \hat{L}_j(t)).$$

Since

$$\begin{aligned} (h_*)_{(t,0)}(\partial/\partial t) &= \bar{y}(t) \\ (h_*)_{(t,0)}(\partial/\partial x_j) &= L_j(t) \\ (h_*)_{(t,0)}(\partial/\partial y_k) &= E_k(t) \\ (h_*)_{(t,0)}(\partial/\partial u_l) &= F_l(t) \\ (h_*)_{(t,0)}(\partial/\partial v_j) &= \hat{L}_j(t) \end{aligned}$$

for each  $t$ , there is a neighborhood  $U$  of  $(t, \bar{0})$  such that  $h$  is an imbedding on  $U$ . By shrinking, if necessary, we can find a neighborhood  $V$  of  $\{(t, \bar{0}) \mid t \in \mathbf{R}\}$  such that  $h_*(\partial/\partial u_l)$  and  $h_*(\partial/\partial v_j)$  are extensions of  $F_l(t)$  and  $\hat{L}_j(t)$  respectively to  $h(V)$ . By making  $V$  smaller and restricting  $t$  to  $[a, b]$  we can assume  $h_*(\partial/\partial u_l)$  and  $h_*(\partial/\partial v_j)$  span a complement in a neighborhood of  $y(t)$ ,  $t \in [a, b]$ .

In this neighborhood we let  $Q$  be the projection defined by the decomposition  $T_x(M) = T^0(x) \oplus T^c(x)$

$$(2.2) \quad Q: T_x(M) \rightarrow T^c(x).$$

For any  $Y \in T^0$  and  $X \in TM$  we can define

$$(2.3) \quad C_Y X = -Q(\nabla_X Y).$$

$C$  is called the conullity operator.

We need the following simple, technical lemma.

**LEMMA 1.** *Let  $C$  and  $Q$  be defined by (2.2) and (2.3). If  $Y$  is in  $T^0$  and  $U$  and  $V$  are in  $TM$  then*

- (1)  $Q(\nabla_Y U) = Q(\nabla_Y (QU))$
- (2)  $Q(\nabla_{U-QU} Y) = 0$
- (3)  $\alpha(U, V) = \alpha(QU, V)$
- (4)  $C$  is a tensor.

*Proof* (1) Suppose  $U = U_0 + U_c$ , where  $U_0 \in T^0$  and  $U_c \in T^c$ . Then  $Q(\nabla_Y U) = Q(\nabla_Y(U_0 + U_c)) = Q(\nabla_Y U_c)$ , since  $\nabla_Y U_0$  is in  $T^0$ . This means that  $Q(\nabla_Y U) = Q(\nabla_Y(QU))$ .

(2) If  $U = U_0 + U_c$  then  $QU = U_c$ . This says that  $U - QU \in T^0$ , so that  $Q(\nabla_{U-QU} Y) = 0$ .

(3) Again  $\alpha(U, V) = \alpha(U_0 + QU, V) = \alpha(QU, V)$ .

(4) It is sufficient to show that  $C_{\varphi Y} = \varphi C_Y U$  for  $\varphi: M \rightarrow \mathbf{R}$ .

$$\begin{aligned} C_{\varphi Y} U &= -Q(\nabla_U \varphi Y) = -Q[(U\varphi)Y + \varphi(\nabla_U Y)] \\ &= \varphi[-Q(\nabla_U Y)] = \varphi C_Y U. \end{aligned} \quad \square$$

We now define a connection  $\nabla'$  in the complementary local bundle  $T^c$ . If  $U \in TM$ ,  $V \in T^c$ , then

$$(2.4) \quad \nabla'_U V = Q(\nabla_U V).$$

Using this connection we can differentiate  $C$ .

$$(2.5) \quad (\nabla'_Y C_Y) X = \nabla'_Y (C_Y X) - C_Y (\nabla'_Y X).$$

Another expression can be found for  $(\nabla'_Y C_Y) X$ . The first term is

$$\nabla'_Y (C_Y X) = Q(\nabla_Y (C_Y X)) = -Q(\nabla_Y (Q(\nabla_X Y))) = -Q(\nabla_Y \nabla_X Y).$$

The second term is, by Lemma (1.2),

$$-C_Y (\nabla'_Y X) = Q(\nabla_{\nabla'_Y X} Y) = Q(\nabla_{Q(\nabla_Y X)} Y) = Q(\nabla_{\nabla_Y X} Y).$$

Combining both terms gives

$$\nabla'_Y (C_Y X) - C_Y (\nabla'_Y X) = -Q(R(Y, X)Y + \nabla_X \nabla_Y Y - \nabla_{\nabla_X Y} Y).$$

If  $W \in T^1$  then  $\langle \nabla_Y Y, W \rangle = 0$  and so  $0 = X \langle \nabla_Y Y, W \rangle = \langle \nabla_X \nabla_Y Y, W \rangle + \langle \nabla_Y Y, \nabla_X W \rangle$ . Along a geodesic  $y_i$  in  $T^0$  let  $Y = \bar{y}_i$ . Then  $\nabla_Y Y = 0$ , and along  $y_i$  we have  $\langle \nabla_X \nabla_Y Y, W \rangle = 0$  and  $Q(\nabla_X \nabla_Y Y) = 0$ .

Next we claim that  $Q(\nabla_{\nabla_X Y} Y) = C_Y (C_Y X)$ . In fact  $C_Y (C_Y X) = -Q(\nabla_{C_Y X} Y) = Q(\nabla_{Q(\nabla_X Y)} Y) = Q(\nabla_{\nabla_X Y} Y)$  by Lemma 1.

Finally then, if  $Y$  is an extension of the tangent vectors  $\bar{y}_i$  along a geodesic in  $T^0$  then

$$(2.6) \quad (\nabla'_Y C_Y) X = Q(R(X, Y)Y) + C_Y^2 X.$$

**THEOREM 2.** *If  $f: M_s^n \rightarrow M_t^{n+k}(c)$  is an isometric immersion and  $M^n$  is complete, then the relative nullity foliation is geodesically complete.*

We first sketch a proof of Theorem 2.

Let  $y_t$  be a geodesic in a leaf  $L$  of  $T^0$ . It must be shown that  $y_t$  can be extended indefinitely in  $L$ . Since  $L$  is totally geodesic in  $M$  we know that  $y_t$  is a geodesic in  $M$  which can be extended indefinitely in  $M$  because  $M$  is geodesically complete. It must be proven to lie entirely in  $L$ . Assume that  $y_t$  is in  $L$  for  $t$  in  $[a, b)$ . If we can show that  $y_b$  is in  $G$ , then we can take a coordinate system  $\{y^1, \dots, y^n\}$  adapted to the foliation with origin  $y_b$ , that is, with the property that the integral manifolds of  $T^0$  are given by  $y^{\nu_0+j} = c_j$ . Now all points  $y_t$ , for  $t$  less than and close to  $b$ , belong to one slice. As  $t$  approaches  $b$ ,  $y_t$  approaches  $y_b$  with coordinates  $(0, \dots, 0)$ , so that  $c_1, \dots, c_{n-\nu_0}$  are all zero. Thus,  $y_b \in L$  and we are done.

To show that  $y_b \in G$  we need the following lemma, which will be proved after the proof of the theorem.

**LEMMA 2.** *For any  $Z$  in  $T_{y_a}(M)$  there exist  $Z_t \in T_{y_t}(M)$ ,  $a \leq t < b$ , such that  $Z_a = Z$  and*

$$(2.7) \quad \nabla'_t(QZ_t) + C_{\vec{y}_t}(QZ_t) = 0 \quad \text{for } a \leq t < b.$$

Moreover,  $Z_t$  can be extended differentiably to  $t = b$ . Here  $\nabla'_t$  stands for  $\nabla'_{\vec{y}_t}$ .

The extension part of Lemma 2 will be proved using 2.6. Let  $X_t$  be a parallel vector field along  $y_t$ ,  $a \leq t < b$ , such that  $X_b \in T^0(y_b)$ . We will prove that  $X_a \in T^0(y_a)$  so  $\nu(y_a) \geq \nu(y_b) \geq \nu_0$  and  $\nu(y_b) = \nu_0$ . Take  $Z_t$  as in Lemma 2. For each point  $y_t$ ,  $t < b$ , extend  $Y_t = \vec{y}_t$ ,  $X_t$  and  $Z_t$  to vector fields  $Y$ ,  $X$  and  $Z$  with  $Y$  in  $T^0$ .

Examine Codazzi's equation with  $Y$ ,  $X$  and  $Z$ .

$$\begin{aligned} & \nabla_Y^\perp \alpha(Z, X) - \alpha(\nabla_Y Z, X) - \alpha(Z, \nabla_Y X) \\ & = \nabla_Z^\perp \alpha(Y, X) - \alpha(\nabla_Z Y, X) - \alpha(Y, \nabla_Z X) \end{aligned}$$

Along  $y_t$   $\nabla_Y X = 0$  and  $\alpha(Y, X) = 0$  and  $\alpha(Y, \nabla_Z X) = 0$ . The equation reduces to

$$\nabla_Y^\perp \alpha(Z, X) - \alpha(\nabla_Y Z, X) = -\alpha(\nabla_Z Y, X) = \alpha(C_Y Z, X)$$

by Lemma (1.3). This gives

$$\begin{aligned} \nabla_Y^\perp \alpha(Z, X) & = \alpha(\nabla_Y Z, X) + \alpha(C_Y Z, X) \\ & = \alpha(Q(\nabla_Y(QZ)), X) + \alpha(C_Y Z, X) \\ & = \alpha(\nabla'_t QZ, X) + \alpha(C_{\vec{y}_t} Z, X) \end{aligned}$$

along  $y_t$ .

This is true for  $t < b$ , so by continuity it holds for  $t = b$ . That is to say that  $\alpha(Z, X)$  is parallel along  $y_t$ ,  $a \leq t \leq b$ .  $\alpha(Z, X) = 0$  at  $t = b$

which means  $\alpha(Z, X) = 0$  at  $y_a$ . Since  $Z_a$  is arbitrary,  $X_a \in T^0(y_a)$ , as desired.

*Proof of Lemma 2.* In what follows we use the same notation that was introduced in the definition of  $T^c(x)$ . Along  $y_t$  write

$$QZ_t = \Sigma\varphi_j(t)\hat{L}_j(t) + \Sigma\psi_p(t)F_p(t).$$

We want to solve  $Q(\nabla_t QZ_t) + C_{\bar{y}_t}(QZ_t) = 0$ . Rewriting, this is

$$Q\left(\nabla_t(\Sigma\varphi_j(t)\hat{L}_j(t) + \Sigma\psi_p(t)F_p(t))\right) + C_{\bar{y}_t}(QZ_t) = 0;$$

or

$$Q\left(\Sigma\varphi_j'(t)\hat{L}_j(t) + \Sigma\psi_p'(t)F_p(t)\right) + C_{\bar{y}_t}(QZ_t) = 0;$$

or

$$\Sigma\varphi_j'(t)\hat{L}_j(t) + \Sigma\psi_p'(t)F_p(t) + C_{\bar{y}_t}(QZ_t) = 0.$$

Since  $C_{\bar{y}_t}(QZ_t)$  can be written in terms of  $\hat{L}_j(t)$  and  $F_p(t)$  this is a system of ordinary differential equations which can be solved for  $a \leq t < b$ . To see that the solution can be extended: By (2.6)

$$(\nabla_Y' C_Y)X = C_Y^2 X + Q(R(X, Y)Y) \quad \text{for all } X \text{ in } TM.$$

$Q(R(X, Y)Y) = c\langle Y, Y\rangle QX$ , so that along  $y_t$ ,  $a \leq t < b$ ,

$$(2.8) \quad (\nabla_Y' C_Y) = C_Y^2 + c\langle Y, Y\rangle Q.$$

If  $Z_t$  satisfies (2.7), then by differentiating once more we have:

$$\nabla_t'^2(QZ_t) + \nabla_t' C_{\bar{y}_t}(QZ_t) = 0$$

or

$$\nabla_t'^2(QZ_t) + (\nabla_t' C_{\bar{y}_t})QZ_t + C_Y(\nabla_t'(QZ_t)) = 0.$$

This yields, using (2.8),

$$\nabla_t'^2(QZ_t) + C_Y^2(QZ_t) + c\langle Y, Y\rangle QZ_t + C_Y(\nabla_t'(QZ_t)) = 0.$$

Plugging in (2.7) gives

$$\nabla_t'^2(QZ_t) + C_Y^2(QZ_t) + c\langle Y, Y\rangle QZ_t - C_Y^2(QZ_t) = 0,$$

i.e.,

$$(2.9) \quad \nabla_t'(QZ_t) - c\langle Y, Y\rangle QZ_t = 0 \quad \text{for } a \leq t < b.$$

In terms of the parallel basis for  $T^c$  along  $y_t$  (2.9) can be written

$$\frac{d^2\varphi_j(t)}{dt^2} + c\langle Y, Y\rangle\varphi_j = 0, \quad \frac{d^2\psi_p(t)}{dt^2} + c\langle Y, Y\rangle\psi_p = 0.$$

Solutions to these equations can be extended differentiably beyond  $b$ .  $\square$



To see that (2.6) is an equation of Ricatti-type write

$$C_Y \hat{L}_j(t) = \sum_{i=1}^r C_{ij}^L(t) \hat{L}_i(t) + \sum_{p=1}^{n-\nu_0-r} C_{pj}^F(t) F_p(t)$$

$$C_Y F_q(t) = \sum_{i=1}^r D_{iq}^L(t) \hat{L}_i(t) + \sum_{p=1}^{n-\nu_0-r} D_{pq}^F(t) F_p(t).$$

Then

$$\begin{aligned} (\nabla'_Y C_Y) \hat{L}_j(t) &= \nabla'_Y (C_Y \hat{L}_j(t)) - C_Y (\nabla'_Y \hat{L}_j(t)) \\ &= \nabla'_Y \left[ \sum_i C_{ij}^L(t) \hat{L}_i(t) + \sum_p C_{pj}^F(t) F_p(t) \right] \\ &= \sum_i \frac{dC_{ij}^L(t)}{dt} \hat{L}_i(t) + \sum_p \frac{dC_{pj}^F(t)}{dt} F_p(t). \end{aligned}$$

Similarly,

$$\begin{aligned} (\nabla'_Y C_Y) F_q(t) &= \sum_i \frac{dD_{iq}^L(t)}{dt} \hat{L}_i(t) + \sum_p \frac{dD_{pq}^F(t)}{dt} F_p(t). \\ Q(R(\hat{L}_j(t), Y)Y) &= c \langle Y, Y \rangle \hat{L}_j(t), \\ Q(R(F_q(t), Y)Y) &= c \langle Y, Y \rangle F_q(t). \end{aligned}$$

Let

$$C(t) = \begin{bmatrix} C_{ij}^L(t) & D_{iq}^L(t) \\ C_{pj}^F(t) & D_{pq}^F(t) \end{bmatrix} \quad \text{and} \quad K(t) = c \langle Y, Y \rangle I_n.$$

Then, (2.6) can be written as

$$(2.10) \quad \frac{dC(t)}{dt} = C^2(t) + K(t).$$

**3. Applications of the Ricatti-type differential equation.** Let  $M_1^n$  be a geodesically complete Lorentzian submanifold of  $\tilde{M}_1^{n+k}(c)$  ( $c > 0$ ), where  $\tilde{M}_1^{n+k}(c)$  is the Lorentzian space form of constant curvature  $c$ . Let  $\nu_0$  be the index of relative nullity of  $M_1^n$  in  $\tilde{M}_1^{n+k}(c)$ .

It is well-known that for  $K(t) > 0$ , the equation (2.10) has no global solution with an initial condition  $C(0)$  which has a real eigenvalue. This implies, in our case, that there is no global solution of (2.10) if  $\langle Y, Y \rangle > 0$ . If  $\langle Y, Y \rangle = 0$ , 0 is the only global solution under an initial condition  $C(0)$

with a real eigenvalue. In particular, if  $C(t)$  has a real eigenvalue for some  $t$ ,  $Y$  must be 0, provided that  $Y$  is space-like.

Here let us state an elementary fact on a Lorentzian vector space.

**PROPOSITION 3.** *Let  $L^n$  be a Lorentzian inner product space with the inner product  $\langle \cdot, \cdot \rangle$ . Let  $V^\nu$  be a linear subspace of dimension  $\nu$ . Then,*

- (i)  $V^\nu = L^\nu$  and  $\langle \cdot, \cdot \rangle|_{V^\nu}$  is non-degenerate, or
- (ii)  $V^\nu = E^\nu$  and  $\langle \cdot, \cdot \rangle|_{V^\nu}$  is positive definite, or
- (iii)  $\langle \cdot, \cdot \rangle|_{V^\nu}$  is degenerate and  $V^\nu = E^{\nu-1} + \text{span}\{\xi\}$ , where  $\langle \xi, \xi \rangle = 0$  and  $\xi \perp E^{\nu-1}$ .

(Proof). See Graves [9].

Now let  $Y_1, \dots, Y_{\nu_0-1}$  be a set of  $(\nu_0 - 1)$ , linearly independent space-like vectors in  $T^0(x)$  such that  $\text{span}\{Y_1, \dots, Y_{\nu_0-1}\}$  is a positive definite subspace of  $T^0(x)$ . Proposition 1 tells us the choice of  $Y_1, \dots, Y_{\nu_0-1}$  is possible. Denote  $C_i = C_{Y_i}$  ( $i = 1, \dots, \nu_0 - 1$ ) for simplicity.

**LEMMA 3.** *The set of vectors  $\{X, C_1(X), \dots, C_{\nu_0-1}(X)\}$  forms a  $\nu_0$ -frame in  $T^c(x)$  for  $X \neq 0 \in T^c(x)$ .*

*Proof.* Let  $\alpha X + \alpha_1 C_1(X) + \dots + \alpha_{\nu_0-1} C_{\nu_0-1}(X) = 0$ . Then

$$\alpha_1 C_1(X) + \dots + \alpha_{\nu_0-1} C_{\nu_0-1}(X) = C_{\alpha_1 Y_1 + \dots + \alpha_{\nu_0-1} Y_{\nu_0-1}}(X) = (-\alpha)X.$$

Hence,  $-\alpha$  is a real eigenvalue of  $C_{\alpha_1 Y_1 + \dots + \alpha_{\nu_0-1} Y_{\nu_0-1}}$ . Since  $\alpha_1 Y_1 + \dots + \alpha_{\nu_0-1} Y_{\nu_0-1}$  is a space-like vector,  $\alpha_1 Y_1 + \dots + \alpha_{\nu_0-1} Y_{\nu_0-1} = 0$  from the above remark. This implies  $\alpha_1 = \dots = \alpha_{\nu_0-1} = 0$  and  $\alpha = 0$ .

As was done in [7], denote by  $V_{n,r}$  the Stiefel manifold of ordered  $r$ -frames in  $E^n$ . It is well known that  $V_{n,r} \rightarrow V_{n,1}$  is a principal fiber bundle in a natural way. Denote by  $\rho(n)$  the largest integer such that the fibration  $V_{n,\rho(n)} \rightarrow V_{n,1}$  has a global cross-section. Define by  $\nu_n$  the largest integer such that  $\rho(n - \nu_n) \geq \nu_n$ .

**THEOREM 3.** *Let  $M_1^n$  be a geodesically complete, connected submanifold of  $\tilde{M}_1^{n+k}(c)$ ,  $c > 0$ . If the index of relative nullity  $\nu_0 > \nu_n$ , then  $M_1^n$  is totally geodesic in  $\tilde{M}_1^{n+k}(c)$  and  $\nu_0 = n$ .*

*Proof.* For any  $x \in G$ ,  $T^0(x)$  always contains a copy of  $E^{\nu_0-1}$ . By Lemma 3,  $V_{n-\nu_0,\nu_0} \rightarrow V_{n-\nu_0,1}$  has a global cross-section  $\varphi$  defined by  $\varphi(X) = (X, C_1 X, \dots, C_{\nu_0-1} X)$  for  $\forall X \neq 0 \in T^c(x)$ . Hence,  $\rho(n - \nu_0) \geq \nu_0$ ; therefore,  $\nu_0 \leq \nu_n$ .

REMARK. If one of the relative nullity leaves is a Riemannian manifold relative to the induced metric, the above  $\nu_n$  can be improved by defining  $\bar{\nu}_n$  to be the largest integer such that  $\rho(n - \bar{\nu}_n) \geq \bar{\nu}_n + 1$ . Clearly  $\bar{\nu}_n \leq \nu_n$ .

Some of the numerical values for  $\nu_n$  are as follows:  $\nu_1 = 0$ ,  $\nu_2 = 1$ ,  $\nu_3 = 1$ ,  $\nu_4 = 2$ ,  $\nu_5 = 1$ ,  $\nu_6 = 2$ ,  $\nu_7 = 3$ ,  $\nu_8 = 4$ ,  $\nu_9 = 1$ .

The argument used here can be applied to obtain a similar result for more general indefinite metrics. Unlike the Riemannian case, it is known that the above  $\nu_n$ 's are often the best possible value. For example, Graves and Nomizu [11] constructed an isometric immersion of  $S_1^2$  into  $S_1^3$  with the index of relative nullity 1.

Our next result states:

THEOREM 4. *Let  $f: M_1^n \rightarrow \tilde{M}_1^{n+p}(c)$  be an isometric immersion between two Lorentzian manifolds, where  $\tilde{M}_1^{n+p}(c)$  is the Lorentzian space form of positive curvature  $c$ . Suppose that the Ricci curvature  $S$  of  $M_1^n$  satisfies  $S(X, X) \geq (n - 1)c\langle X, X \rangle$  for all space-like vectors  $X$ .*

(1) *If  $T^0(x)$  is Lorentzian for some  $x \in G$ , then the index of relative nullity is either 0 or  $n$ .*

(2) *If  $T^0(x)$  is degenerate for some  $x \in G$ , then the index of relative nullity is 0, 1,  $n - 1$  or  $n$ .*

(3) *If  $T^0(x)$  is Riemannian for all  $x \in G$  and if  $p = 1$ , then the index of relative nullity is 0,  $n - 2$ ,  $n - 1$ , or  $n$ .*

We will prove Theorem 4 after a sequence of propositions and lemmas.

PROPOSITION 4. *Let  $(\ , \ )$  be a symmetric bilinear form on an  $n - \nu_0$  dimensional vector space  $V$  over  $\mathbf{R}$  with signature  $(m_1, m_2, m_3)$ . If  $m_1 \neq m_2$  and  $T: V \rightarrow V$  is a symmetric linear operator with respect to  $(\ , \ )$ , then  $T$  has a real eigenvalue. Note a symmetric bilinear form of signature  $(m_1, m_2, m_3)$  has  $m_1$   $(-1)$ 's,  $m_2$   $(+1)$ 's and  $m_3$  0's in the canonical form.*

*Proof.* Choose a canonical basis  $\{e_1, \dots, e_{m_1}, f_1, \dots, f_{m_2}, g_1, \dots, g_{m_3}\}$  of  $V$  for the symmetric bilinear form, so that  $(e_\alpha, e_{\alpha'}) = -\delta_{\alpha\alpha'}$ ,  $(f_\beta, f_{\beta'}) = \delta_{\beta\beta'}$ ,  $(g_\gamma, g_{\gamma'}) = 0$  and all other products are zero. The matrix of the

symmetric operator  $T$  with respect to this basis has the form:

$$\begin{array}{c} m_1 \\ m_2 \\ m_3 \end{array} \left[ \begin{array}{c|c|c} m_1 & m_2 & m_3 \\ \hline A & -B' & 0 \\ \hline B & C & 0 \\ \hline D & E & F \end{array} \right].$$

Here  $A' = A$  and  $C' = C$ . Denote this matrix by  $M_T$ . The matrix  $M_s = \begin{bmatrix} A & -B' \\ B & C \end{bmatrix}$  represents a symmetric linear transformation on the span of  $\{e_1, \dots, e_{m_1}, f_1, \dots, f_{m_2}\}$ . By a change of basis,  $M_s$  can be put into a standard form [15]. If  $m_1 \neq m_2$ , there is one block of the form:

$$\begin{bmatrix} \lambda & 1 & \cdots & 0 \\ \vdots & & \ddots & \vdots \\ \vdots & & & 1 \\ 0 & \cdots & & \lambda \end{bmatrix}.$$

Thus,  $M_s$  has an eigenvalue  $\lambda$ . Noting that

$$\det[\lambda I - M_T] = \det[\lambda I - M_s] \det[\lambda I - F] = 0,$$

we see that  $T$  has a real eigenvalue  $\lambda$ . □

Now let  $\eta = \sum_{i=1}^n \langle X_i, X_i \rangle \alpha(X_i, X_i)$ , where  $\{X_1, \dots, X_n\}$  is an orthonormal basis of  $T_x(M)$ . Define a bilinear form  $(, ) : T^c(x) \times T^c(x) \rightarrow \mathbf{R}$  by, for any  $X$  and  $Y \in T^c(x)$ ,

$$(X, Y) = \langle \alpha(X, Y), \eta \rangle.$$

It is clear that  $(, )$  is symmetric, since  $\alpha$  is symmetric. We may find an alternative expression for  $(, )$  using the Gauss equation and the Ricci curvature of  $M_1^n$ .

$$\begin{aligned} (X, Y) &= \langle \alpha(X, Y), \eta \rangle \\ &= \sum_{i=1}^n \langle X_i, X_i \rangle \langle \alpha(X, Y), \alpha(X_i, X_i) \rangle \\ &= S(X, Y) - nc \langle X, Y \rangle + c \langle X, Y \rangle \\ &\quad + \sum_{i=1}^n \langle X_i, X_i \rangle \langle \alpha(X_i, X), \alpha(X_i, Y) \rangle. \end{aligned}$$

Clearly, the definition of  $(, )$  and the above expression for  $(X, Y)$  do not depend on the choice of the orthonormal basis  $\{X_1, \dots, X_n\}$ .

If  $\{\xi_1, \dots, \xi_p\}$  is an orthonormal basis for  $N(x)$ , then the last term of the above expression can be rewritten

$$\begin{aligned} & \sum_{i,j} \langle X_i, X_i \rangle \langle \xi_j, \xi_j \rangle \langle \alpha(X_i, X), \xi_j \rangle \langle \alpha(X_i, Y), \xi_j \rangle \\ &= \sum_{i,j} \langle X_i, X_i \rangle \langle \xi_j, \xi_j \rangle \langle A_{\xi_j} X, X_i \rangle \langle A_{\xi_j} Y, X_i \rangle \\ &= \sum_j \langle \xi_j, \xi_j \rangle \langle A_{\xi_j} X, A_{\xi_j} Y \rangle. \end{aligned}$$

Since the normal space is positive definite,

$$(X, Y) = S(X, Y) - c(n-1)\langle X, Y \rangle + \sum_{j=1}^p \langle A_{\xi_j}^2 X, Y \rangle.$$

For convenience, let  $\tau = c(n-1)$ , and  $A_{\xi_j}$  by  $A_j$ . We examine  $(\ , \ )$  by looking at

$$h(X, Y) := S(X, Y) - \tau \langle X, Y \rangle \quad \text{and} \quad k(X, Y) := \sum_{j=1}^p \langle A_j^2 X, Y \rangle.$$

**LEMMA 4.** *If the Ricci curvature satisfies the hypothesis in the theorem, then  $h(X, Y)$  is positive semi-definite or Lorentzian.*

*Proof.* We show that  $h$  cannot have two  $(-1)$ 's in its signature. If it did, we could find a pair of linearly independent vectors  $e$  and  $f$  in  $T_x M$  such that  $h(e, e) = -1$ ,  $h(f, f) = -1$  and  $h(e, f) = 0$  i.e.,  $S(e, e) = -1 + \tau \langle e, e \rangle$ ,  $S(f, f) = -1 + \tau \langle f, f \rangle$  and  $S(e, f) = \tau \langle e, f \rangle$ . By the hypothesis, it would then be the case that  $\langle e, e \rangle \leq 0$  and  $\langle f, f \rangle \leq 0$ .

We now examine the various possibilities for the lengths of  $e$  and  $f$ . In each case, we will find a space-like vector which violates the condition on the Ricci curvature. We will use the reverse Cauchy-Schwarz inequality, i.e.  $0 \leq \langle e, e \rangle \langle f, f \rangle \leq \langle e, f \rangle^2$  in this case. In the following argument, denote  $\langle e, f \rangle$  by  $a$  for brevity.  $\langle e, f \rangle = a$  can be positive.

If both  $e$  and  $f$  were light-like,  $e + f$  would be space-like and  $\langle e + f, e + f \rangle = 2a$ .

$$\begin{aligned} S(e + f, e + f) &= S(e, e) + 2S(e, f) + S(f, f) \\ &= (-1) + 2\tau \langle e, f \rangle + (-1) < 2\tau \langle e, f \rangle = 2\tau a \\ &= \tau \langle e + f, e + f \rangle. \end{aligned}$$

This contradicts the hypothesis.

If  $e$  were light-like and  $f$  were time-like, we could also assume that  $\langle f, f \rangle = -1$ . Then  $e + af$  would be space-like, for  $\langle e + af, e + af \rangle = a^2$ . But

$$\begin{aligned} S(e + af, e + af) &= S(e, e) + 2aS(e, f) + a^2S(f, f) \\ &= -1 + 2a\tau a + a^2(-1 - \tau) < \tau a^2, \end{aligned}$$

a contradiction.

If  $e$  and  $f$  are both time-like, we could assume that  $\langle e, e \rangle = -1 = \langle f, f \rangle$ . Then,  $e + af$  is space-like with  $\langle e + af, e + af \rangle = -1 + a^2 > 0$  by the reverse Cauchy-Schwarz inequality.

$$S(e + af, e + af) = -1 - \tau + 2a^2\tau - a^2 - a^2\tau < (-1 + a^2)\tau,$$

a contradiction.

Next we show that if there are any vectors which are time-like with respect to  $h$ , then  $h$  is non-degenerate and therefore Lorentzian. If this is not the case, there are linearly independent  $e$  and  $g$  such that  $h(e, e) = -1$ ,  $h(g, g) = 0$  and  $h(e, g) = 0$ , i.e.,  $S(e, e) = -1 + \tau\langle e, e \rangle$ ,  $S(g, g) = \tau\langle g, g \rangle$  and  $S(e, g) = \tau\langle e, g \rangle$ . We know that  $\langle e, e \rangle \leq 0$  and can assume that  $\langle e, g \rangle \geq 0$ . If  $\langle g, g \rangle > 0$ , then  $\langle g + e, g + e \rangle = \langle g, g \rangle + 2\langle e, g \rangle > 0$ , so that  $g + e$  is space-like. However,

$$\begin{aligned} S(g + e, g + e) &= \tau\langle g, g \rangle + 2\tau\langle e, g \rangle - 1 \\ &\quad + \tau\langle e, e \rangle < \tau[\langle g, g \rangle + 2\langle e, g \rangle], \end{aligned}$$

a contradiction. The only remaining possibility is that  $\langle e, e \rangle \leq 0$  and  $\langle g, g \rangle \leq 0$ . The span of  $e$  and  $g$  is non-degenerate, so for some  $k \in \mathbf{R}$ ,  $e + kg$  is space-like.

$$\begin{aligned} S(e + kg, e + kg) &= S(e, e) + 2kS(e, g) + k^2S(g, g) \\ &= -1 + \tau\langle e, e \rangle + 2k\tau\langle e, g \rangle + k^2\tau\langle g, g \rangle \\ &< \tau[\langle e, e \rangle - 2k\langle e, g \rangle + k^2\langle g, g \rangle], \end{aligned}$$

a contradiction.

This completes the proof of Lemma 4.

We now turn to  $k(X, Y) = \sum_{j=1}^p \langle A_j^2 X, Y \rangle$  on  $T^c(x)$ .

**LEMMA 5.** *If  $T^c(x)$  is positive definite, then  $k$  restricted to  $T^c(x)$  is positive definite.*

*Proof.*

$$k(X, X) = \sum_{j=1}^p \langle A_j^2 X, X \rangle = \sum_{j=1}^p \langle A_j X, A_j X \rangle.$$

If  $k(X, X) = 0$ , then  $A_j X = 0$  for  $j = 1, \dots, p$ , since  $T^c(x) = T^1(x) = \text{span}\{A_j Y\}$  is positive definite. This means that  $X \in T^0(x) \cap T^1(x)$ ; therefore,  $X = 0$ . Hence,  $k$  is positive definite.

**PROPOSITION 5.** *If  $T^c(x)$  is positive definite, then  $(\cdot, \cdot)$  is positive definite on  $T^c(x)$ .*

*Proof.* For no non-zero  $e \in T^c(x)$  is  $h(e, e) = -1$ , since this implies  $\langle e, e \rangle \leq 0$ . Thus, the form  $(\cdot, \cdot)$  is the sum of a positive semi-definite form and a positive definite form.

**LEMMA 6.** *If  $T^c(x)$  is a degenerate subspace with respect to  $\langle \cdot, \cdot \rangle$ , then  $h$  is positive semi-definite on  $T^c(x)$  for  $\dim T^c(x) > 1$ .*

*Proof.*  $T^c(x)$  is a positive semi-definite subspace with respect to  $\langle \cdot, \cdot \rangle$  by hypothesis. If there were an  $e \in T^c(x)$  with  $h(e, e) = -1$ , then  $\langle e, e \rangle = 0$ . By Lemma 4, there would be  $g \in T^c(x)$  such that  $h(g, g) = 1$  and  $h(e, g) = 0$ . By hypothesis we would have  $\langle g, g \rangle > 0$  and  $\langle e, g \rangle = 0$ . For all  $t \in \mathbf{R}$   $\langle g + te, g + te \rangle = \langle g, g \rangle > 0$ , but

$$S(g + te, g + te) = 1 + \tau \langle g, g \rangle - t^2 < \tau \langle g, g \rangle.$$

This is a contradiction.

**LEMMA 7.** *If  $T^c(x)$  is a degenerate subspace with respect to  $\langle \cdot, \cdot \rangle$  and if  $\dim T^c(x) > 1$ , then  $k$  is positive semi-definite on  $T^c(x)$  and for some  $Y \in T^c(x)$   $k(Y, Y) > 0$ .*

*Proof.* If  $T^0(x)$  is degenerate, so is  $T^1(x) = \text{span}[A_\xi Y]$ . Thus, there is a light-like vector  $L$  such that, for any normal vector  $\xi$ ,  $\text{Im } A_\xi \subseteq [L]^\perp$ . The metric on  $[L]^\perp$  is positive semi-definite, so  $\langle A_\xi Y, A_\xi Y \rangle \geq 0$  for all  $\xi, Y$ . This implies that  $k$  is positive semi-definite. If  $k(Y, Y) = 0$  for all  $Y$ , then each  $A_\xi Y$  is light-like. We also have  $\langle A_\xi Y, L \rangle = 0$  for  $\xi, Y$ . Recalling that perpendicular light-like vectors are linearly dependent, we see that  $T^1(x)$  would be one-dimensional, which is not the case.

**PROPOSITION 6.** *If  $T^c(x)$  is a degenerate subspace, then  $(\cdot, \cdot)$  is positive semi-definite on  $T^c(x)$  and for some  $Y \in T^c(x)$ ,  $(Y, Y) > 0$ .*

Finally, we assume  $M_1^n$  is a hypersurface in  $\tilde{M}_1^{n+1}$  and that  $T^c = T^1$  is Lorentzian. Here  $k(X, Y) = \langle AX, AY \rangle$  where  $A$  is a transformation of  $T^1$  which is one-to-one.

**PROPOSITION 7.** *If  $M_1^n$  is a hypersurface in  $\tilde{M}_1^{n+1}(c)$ ,  $T^1$  is Lorentzian and  $n - \nu_0 > 2$ , then the signature  $(m_1, m_2, m_3)$  of  $(, )$  has  $m_2 > m_1$ .*

*Proof.* We know that  $h$  is positive semi-definite or Lorentzian. The shape operator  $A$  can be put into one of four canonical forms [15] and  $k$  can be explicitly calculated using these forms.

$$(i) \quad A = \begin{bmatrix} \lambda & 0 & & & & \\ 1 & \lambda & & & & \\ & & \lambda_1 & & & \\ & & & \ddots & & \\ & & & & \lambda_{n-\nu_0-2} & \end{bmatrix},$$

where  $A$  is given with respect to a pseudo-orthonormal basis  $\{\hat{L}, L, f_1, \dots, f_{n-\nu_0-2}\}$ .  $k$  then has the following matrix with respect to this basis:

$$\begin{bmatrix} -2\lambda & -\lambda^2 & & & & \\ -\lambda^2 & 0 & & & & \\ & & \lambda_1^2 & & & \\ & & & \ddots & & \\ & & & & \lambda_{n-\nu_0-2}^2 & \end{bmatrix}.$$

Note that  $\lambda\lambda_1 \cdots \lambda_{n-\nu_0-2} \neq 0$  and the signature of  $k$  is  $(1, n - \nu_0 - 1)$ . We can find a space-like vector  $U$  in  $\text{span}\{\hat{L}, L\}$  such that  $k(U, U) > 0$ . In fact, there are choices for  $b \in \mathbf{R}$  such that  $\langle \hat{L} + bL, \hat{L} + bL \rangle = -2b$  and  $k(\hat{L} + bL, \hat{L} + bL) = -2\lambda(1 + b\lambda)$  are both positive. Set  $U = \hat{L} + bL$ . We claim that  $(, )$  is positive definite on  $\text{span}\{U, f_1, \dots, f_{n-\nu_0-2}\}$ .

$$h(cU + \sum c_j f_j, cU + \sum c_j f_j) \geq 0$$

since

$$\langle cU + \sum c_j f_j, cU + \sum c_j f_j \rangle = c^2 \langle U, U \rangle + \sum c_j^2 \geq 0.$$

It is also clear that  $k(cU + \sum c_j f_j, cU + c_j f_j) \geq 0$  and is equal to 0 if and only if  $c = 0 = c_j$  for  $j = 1, \dots, n - \nu_0 - 2$ . Hence, if  $n - \nu_0 > 2$ , the signature of  $(, )$  will have more plus signs than minus signs.

$$(ii) \quad A = \begin{bmatrix} \lambda & 0 & 0 & & & \\ 0 & \lambda & 1 & & & \\ -1 & 0 & \lambda & & & \\ & & & \lambda_1 & & \\ & & & & \ddots & \\ & & & & & \lambda_{n-\nu_0-3} \end{bmatrix}$$





The following result concerns the over simplified ‘‘The axiom of sphere’’ for the hypersurfaces in the Lorentzian space form of positive constant curvature  $c$ .

Let  $f: M_1^n \rightarrow \tilde{M}_1^{n+1}(c)$  be a geodesically complete Lorentzian hypersurface, where  $\tilde{M}_1^{n+1}(c)$  is the Lorentzian space form of positive curvature  $c$ . Let us assume:

(\*) Through each point  $x$  of  $M$  exists a  $k(x)$ -dimensional local submanifold  $S_x$  of  $M$  which is mapped under  $f$  isometrically into a  $k(x)$ -dimensional totally geodesic submanifold of  $\tilde{M}_1^{n+1}(c)$ .

**THEOREM 5.** *If  $2k(x) - n \geq v_n$ ,  $M_1^n$  is totally geodesic in  $\tilde{M}_1^{n+1}(c)$ . Here  $v_n$  is the numerical value determined in Theorem 3.*

**LEMMA 8.** *If  $2k(x) - n > 0$ , the relative nullity  $\nu(x)$  at  $x \geq 2k(x) - n$ . In particular, the index of relative nullity  $\nu_0 \geq 2k - n$ , where  $k = \min_{x \in M_1^n} k(x)$ .*

*Proof.* If  $S_x$  is non-degenerate, the result of Lemma 8 will be obtained in the same manner as in [3]. Now let us assume that  $S_x$  is degenerate. Let  $\hat{e}_1(x), \dots, \hat{e}_{n-k-1}(x)$ ,  $L$  form a pseudo-orthonormal basis for  $(TS_x)^\perp$ , which is the orthogonal complement of  $TS_x$  in  $T_x M_1^n$ . Extend it to a basis of  $T_x M$  by adding  $\{e_1(x), \dots, e_{k-1}(x), \hat{L}(x)\}$ , which is also pseudo-orthonormal. Then,  $e_1(x), \dots, e_{k-1}(x), L(x)$  form a basis for  $TS_x$ . Since  $S_x$  is totally geodesic,  $\alpha(e_i(x), e_j(x)) = \alpha(e_i(x), L) = 0$ , where  $\alpha$  is the second fundamental form. Set

$$A_\xi(e_i) = \sum_j a_{ji} e_j + b_i L + \text{a linear combination of } \hat{e}_k \text{ and } \hat{L}$$

and

$$A_\xi(L) = \sum_j c_j e_j + bL + \text{a linear combination of } \hat{e}_k \text{ and } \hat{L}.$$

Similarly,  $A_\xi(\hat{e}_k)$  and  $A_\xi(L)$  can be given as a linear combination of the basis elements  $e_1, \dots, e_{k-1}, L, \hat{L}, \hat{e}_1, \dots, \hat{e}_{n-k-1}$ . With respect to this basis,  $A_\xi$  is represented by an  $n \times n$ -matrix, which we also denote by the same symbol. In fact, we have

$$A_\xi(e_i) = -\langle A_\xi(e_i), \hat{L} \rangle L + \text{a linear combination of } \hat{e}_k \text{'s}$$

and

$$A_\xi(L) = -\langle A_\xi(L), \hat{L} \rangle L + \text{a linear combination of } \hat{e}_k \text{'s}.$$



## REFERENCES

- [1] K. Abe, *Application of a Riccati type differential equation to Riemannian manifolds with totally geodesic distributions*, Tohoku Math. J., **25** (1973), 425–444.
- [2] ———, *Some remarks on a class of submanifolds in space forms of non-negative curvature*, Math. Ann., **247** (1980), 275–280.
- [3] ———, *A characterization of totally geodesic hypersurfaces of  $S^{n+1}$  and  $CP^{n+1}$* , Proc. Amer. Math. Soc., **81** No. 4 (1981), 603–606.
- [4] S. S. Chern and N. H. Kuiper, *Some theorems on the isometric imbedding of compact Riemannian manifolds in Euclidean space*, Ann. of Math., **56** (1952), 422–430.
- [5] M. Dajczer and P. Dombrowski, *Examples of 1-codimensional non-totally geodesic isometric immersions of pseudo-Riemannian space forms with the same positive constant curvature and the same spacelike rank*, (to appear).
- [6] M. Dajczer and K. Nomizu, *On the boundedness of Ricci curvature of an indefinite metric*, Bol. Soc. Bras. Mat., **11**, #1 (1980), 25–30.
- [7] D. Ferus, *Totally geodesic foliations*, Math. Ann., **188** (1970), 313–316.
- [8] ———, *Isometric immersions of constant curvature manifolds*, Math. Ann., **217** (1975), 155–156.
- [9] L. Graves, *Codimension one isometric immersions between Lorentzian spaces*, Trans. Amer. Math. Soc., **252** (1979), 367–392.
- [10] ———, *On codimension one isometric immersions between indefinite space forms*, Tsukuba J. Math., **3**, No. 2 (1979), 17–29.
- [11] L. Graves and K. Nomizu, *Isometric immersions of Lorentzian space forms*, Math. Ann., **233** (1978), 123–136.
- [12] M. Magid, *Isometric immersions of Lorentz space with parallel second fundamental forms*, Tsukuba J. Math., **8**, No. 1 (1984), 31–54.
- [13] ———, *Shape operators of Einstein hypersurfaces in indefinite space forms*, Proc. Amer. Math. Soc., **84** No. 2 (1982), 237–242.
- [14] ———, *Lorentzian isoparametric hypersurfaces*, Pacific J. Math., **118** (1985), 165–197.
- [15] A. Z. Petrov, *Einstein space*, Pergamon Press, Hungary, Oxford, and New York (1969).

Received April 2, 1985.

UNIVERSITY OF CONNECTICUT  
STORRS, CT 06268

AND

WELLESLEY COLLEGE  
WELLESLEY, MA 02181