

Relativistic Closed-Form Hamiltonian for Many-Body Gravitating Systems in the Post-Minkowskian Approximation

Tomáš Ledvinka,^{1,2} Gerhard Schäfer,^{2,1} and Jiří Bičák^{1,2}

¹*Institute of Theoretical Physics, Faculty of Mathematics and Physics, Charles University, Prague, Czech Republic*

²*Theoretisch-Physikalisches Institut, Friedrich-Schiller-Universität, Max-Wien-Platz 1, 07743 Jena, Germany*

(Received 27 January 2008; published 23 June 2008)

The Hamiltonian for a system of relativistic bodies interacting by their gravitational field is found in the post-Minkowskian approximation, including all terms linear in the gravitational constant. It is given in a surprisingly simple closed form as a function of canonical variables describing the bodies only. The field is eliminated by solving inhomogeneous wave equations, applying transverse-traceless projections, and using the Routh functional. By including all special relativistic effects our Hamiltonian extends the results described in classical textbooks of theoretical physics. As an application, the scattering of relativistic objects is considered.

DOI: [10.1103/PhysRevLett.100.251101](https://doi.org/10.1103/PhysRevLett.100.251101)

PACS numbers: 04.25.Nx

Introduction.—The problem of motion of bodies under gravitational interaction has been the central issue in general relativity from its birth. As early as 1916, Einstein, Droste, Lorentz, and de Sitter started to develop post-Newtonian (PN) approximation methods. These are based on the weak-field limit in which the metric is close to the Minkowski metric and the assumption that the typical velocity v in a system divided by the speed of light c is very small, $v/c \sim \epsilon$. The deviation from the flat metric can be characterized by the Newtonian potential Φ ; so for a binary system, for example, $\Phi/c^2 \sim v^2/c^2 \sim \epsilon^2$. In an appropriate limit (as $\epsilon \rightarrow 0$), the PN approximation yields Newton's equations. In 1916 Einstein also worked out “the linearized approximation” to general relativity in which the flat-space wave equations for the deviations $h_{\mu\nu}$ from Minkowski metric with an energy-momentum tensor $T_{\mu\nu}$ of matter as the source are considered. This work marks the beginning of the post-Minkowskian (PM) approximation methods: the weakness of the gravitational field is assumed—the deviations of the metric from flat are small, but no assumption about slowness of motion is made. In an appropriate limit, when a suitable parameter, usually identified with Newton's gravitational constant G , is approaching zero, the PM approximation yields equations of special relativity. The “historical” period culminated in 1938 when Einstein, Infeld, and Hoffmann, by formal expansions in c^{-1} , investigated the n th iterated field equations of the PN scheme.

Since the 1950s numerous investigations of both PN and PM approximations have appeared; see, for example, refined comprehensive reviews [1,2]. Because of the evidence that the orbit of the binary pulsar PSR 1913 + 16 decays as a consequence of the emission of gravitational waves, the most promising candidates for the detectors such as LIGO, VIRGO, and GEO600 became binary neutron stars or black holes. This led to new studies of higher-order equations of motion. Extensive reviews summarizing these developments were prepared recently [3,4].

The Hamiltonian methods have been widely used in general relativity in such problems as perturbation of black holes, dynamics of anisotropic cosmological models, etc. In the problem of motion of gravitating systems the Hamiltonian yields equations of motion, expressions for the mechanical energy and angular momentum; in the case of binaries, the result for the last stable circular orbit follows from $\partial^2 H / \partial r^2 = 0$. The Hamiltonian approach to the PN approximation was initiated by Kimura and Toiya [5] and developed recently in the work of Schäfer, Jaranowski, and Damour (see [4] for references). The canonical formalism of Arnowitt, Deser, and Misner (ADM) [6] is commonly used.

Before we enter into details we wish to indicate why our final result—the closed form Hamiltonian (11) including all terms linear in G —is of importance in the theory of gravity and can be relevant in other branches of theoretical physics and astrophysics: (i) Since PM approximation can be restricted to slow motions, the Hamiltonian describes PN approximations to *any* order in $1/c$ when terms linear in G are considered. When considering only two particles and terms up to the second order ($\leq v^2/c^2$) in center-of-mass system we precisely recover the Hamiltonian given by Landau and Lifshitz [7], except for the last term which is $\sim G^2$. Moreover, we checked that, after a suitable canonical transformation, it yields precisely all the terms linear in G in the 3PN approximation of [8], and we calculated corresponding terms in 4PN order which have not been given thus far. These will be published elsewhere. (ii) An electromagnetic counterpart of our Hamiltonian was derived by completely different methods by Kennedy [9]. The first terms in the expansion in c^{-2} of the Kennedy Hamiltonian correspond to the well-known Darwin Lagrangian (see, e.g., [7,10]), but Kennedy's treatment goes beyond Darwin's approximation. Our procedure shows how the Darwin Hamiltonian or Lagrangian can be generalized in both gravitational and electromagnetic cases. The Darwin Lagrangian or Hamiltonian has many

applications in both classical and quantum domains (see, e.g., [10]). The extension for two point charges considered up to order c^{-4} (included) has been given by the Golubenkov-Smorodinsky Lagrangian, which is described in [7] and discussed in [11]. (iii) As our Hamiltonian can describe particles with ultrarelativistic velocities or with zero rest mass, it is well suited for treating gravitational scattering of such objects. These are closely related to gravitational scattering of shock waves. At high energies in quantum scattering processes, the interaction is well approximated by the classical collision at the speed of light, as first shown by the influential work by 't Hooft (see [12] for a detailed description and methods). (iv) Because of its simplicity, our Hamiltonian yields a convenient starting point in investigations of gravitational lensing by extended gravitationally interacting objects since it provides a unified description of both gravitating bodies and deflected photons or neutrinos. (v) Our method of deriving the Hamiltonian by using the Routhian and the induced canonical transformation, etc. should be of interest in other branches of theoretical physics.

The Hamiltonian approach to the PM approximation was first undertaken in 1986 [13]. Following the ADM canonical formalism, the gravitational field is described by h_{ij}^{TT} , the transverse-traceless part of $h_{ij} = g_{ij} - \delta_{ij}$ ($h_{ii}^{\text{TT}} = 0$, $h_{ij,j}^{\text{TT}} = 0$, $i, j = 1, 2, 3$), and by conjugate momenta $\bar{\pi}^{ij\text{TT}}$. The system of bodies located at \mathbf{x}_a , $a = 1, \dots, N$, with rest masses m_a and momenta \mathbf{p}_a , has the energy and linear momentum densities $\gamma^{1/2} T^{\mu\nu} n_\mu n_\nu = \sum_a (g^{ij} p_{ai} p_{aj} + m_a^2)^{1/2} \delta(\mathbf{x} - \mathbf{x}_a)$, $-\gamma^{1/2} T_i^\mu n_\mu = \sum_a p_{ai} \delta(\mathbf{x} - \mathbf{x}_a)$, where $\gamma = \det(g_{ij})$, g^{ij} is inverse to g_{ij} , n^ν is a unit timelike normal to hypersurface $x^0 = \text{const}$, and $T^{\mu\nu}$ is the energy-momentum tensor of the matter system. Hereafter, we call its constituents the ‘‘particles,’’ but they may well represent neutron stars or black holes. This is substantiated by ‘‘general relativity’s adherence to the strong equivalence principle’’: black holes and other bodies obey the same laws of motion as test bodies; see, e.g., [12].

It is convenient to choose four coordinate conditions $\bar{\pi}^{ii} = 0$ and $g_{ij} = (1 + \frac{1}{8}\phi)^4 \delta_{ij} + h_{ij}^{\text{TT}}$. The standard ADM Hamiltonian (cf. [6]), $H = (1/16\pi G) \oint dS_i (g_{ij,i} - g_{jj,i})$, then becomes, using the Gauss theorem, $H = -(1/16\pi G) \int d^3x \Delta\phi$. The integrand $\Delta\phi = \partial_i \partial_i \phi$ can be expressed in terms of \mathbf{x}_a , \mathbf{p}_a , h_{ij}^{TT} , and $\bar{\pi}^{ij\text{TT}}$ from the constraint equations. By expansions in powers of G and after adopting suitable regularization procedures of integrals involved (see the appendix in [13]), one can determine the Hamiltonian. The Hamiltonian worked out in [13] includes terms $\sim G^2$.

Here we start from the same Hamiltonian neglecting, however, terms $\sim G^2$. Still, the form of the Hamiltonian is quite complicated [see Eq. (1)]: rather than just quantities associated with particles it involves field variables, non-local TT projections, and integrals. The main purpose of

this Letter is to show that due to somewhat magical simplifications the fields h_{ij}^{TT} , assuming they are generated just by particles, can be expressed entirely in terms of the particles’ variables and, after two canonical transformations, the Hamiltonian can be cast into the closed form (11). It is local in the particles’ canonical variables, involving no field variables. It is exact up to terms linear in G , and within the same accuracy it yields equations of motion for particles moving with possibly ultrarelativistic speeds, including those with zero rest mass.

1PM Hamiltonian.—The Hamiltonian describing N particles (\mathbf{x}_a , \mathbf{p}_a) and their gravitation field (h_{ij}^{TT} , $\bar{\pi}^{ij\text{TT}}$) accurate up to the terms linear in G reads

$$H_{\text{lin}} = \sum_a \bar{m}_a - \frac{1}{2} G \sum_{a,b \neq a} \frac{\bar{m}_a \bar{m}_b}{r_{ab}} \left(1 + \frac{p_a^2}{\bar{m}_a^2} + \frac{p_b^2}{\bar{m}_b^2} \right) + \frac{1}{4} G \sum_{a,b \neq a} \frac{1}{r_{ab}} (7\mathbf{p}_a \cdot \mathbf{p}_b + (\mathbf{p}_a \cdot \mathbf{n}_{ab})(\mathbf{p}_b \cdot \mathbf{n}_{ab})) - \frac{1}{2} \sum_a \frac{p_{ai} p_{aj}}{\bar{m}_a} h_{ij}^{\text{TT}}(\mathbf{x} = \mathbf{x}_a) + \frac{1}{16\pi G} \int d^3x \left(\frac{1}{4} h_{ij,k}^{\text{TT}} h_{ij,k}^{\text{TT}} + \bar{\pi}^{ij\text{TT}} \bar{\pi}^{ij\text{TT}} \right), \quad (1)$$

where $\bar{m}_a = (m_a^2 + \mathbf{p}_a^2)^{1/2}$, $\mathbf{n}_{ab} r_{ab} = \mathbf{x}_a - \mathbf{x}_b$, $|\mathbf{n}_{ab}| = 1$. [This is Eq. 18 in [13]. However, although we also put $c = 1$, we keep explicitly G and do not put $16\pi G = 1$; the ‘‘geometric’’ momenta $\bar{\pi}^{ij\text{TT}}$ are thus different from the true $\bar{\pi}^{ij\text{TT}} = (16\pi G)^{-1} \pi^{ij\text{TT}}$. Notice that the regularization is also needed in the terms containing h_{ij}^{TT} , $\bar{\pi}^{ij\text{TT}}$, and $h_{ij}^{\text{TT}}(\mathbf{x} = \mathbf{x}_a)$.] The equations of motion for particles are standard Hamilton equations. The Hamilton equations for the field are

$$\dot{\bar{\pi}}^{ij\text{TT}} = -\delta_{kl}^{\text{TT}ij} \frac{\delta H}{\delta h_{kl}^{\text{TT}}}, \quad \dot{h}_{ij}^{\text{TT}} = \delta_{ij}^{\text{TT}kl} \frac{\delta H}{\delta \bar{\pi}^{kl\text{TT}}}; \quad (2)$$

here the variational derivatives and the TT-projection operator $\delta_{kl}^{\text{TT}ij} = \frac{1}{2} (\Delta_{ik} \Delta_{jl} + \Delta_{il} \Delta_{jk} - \Delta_{ij} \Delta_{kl}) \Delta^{-2}$, $\Delta_{ij} = \delta_{ij} \Delta - \partial_i \partial_j$, appear. These equations imply the equations for gravitational field in the first PM approximation to be the following wave equations:

$$\square h_{ij}^{\text{TT}} = -16\pi G \delta_{ij}^{\text{TT}kl} \sum_a \frac{p_{ak} p_{al}}{\bar{m}_a} \delta^{(3)}(\mathbf{x} - \mathbf{x}_a). \quad (3)$$

As is usual in the system of particles and field, the field is the source of the particles’ accelerations and particles are sources of the field. Once $O(G^2)$ terms can be neglected, however, this coupling simplifies. Since both the field and the accelerations $\dot{\mathbf{p}}_a$ are proportional to G , the changes of the field due to the accelerations of particles are of the order $O(G^2)$. Therefore, we can assume the field to be generated by unaccelerated particles; it is thus given only by their instantaneous positions and velocities. If such a field is used in equations of motion of particles, the gravitational interaction becomes an ‘‘action at a distance.’’

Given the linearity of the field equations (3), we can write $h_{ij}^{\text{TT}}(\mathbf{x}) = \sum_b h_{ij}^{\text{TT}}(\mathbf{x}; \mathbf{x}_b, \mathbf{p}_b, \dot{\mathbf{x}}_b)$, where $h_{ij}^{\text{TT}}(\mathbf{x}; \mathbf{x}_b, \mathbf{p}_b, \dot{\mathbf{x}}_b)$ is the contribution to the field at point \mathbf{x} generated by the particle moving at \mathbf{x}_b with velocity $\dot{\mathbf{x}}_b$. Because the d'Alembertian operator commutes with the TT-projection operator, the field from each particle involves the solution of the scalar wave equation with point source—the boosted static spherical field $\sim 1/r$. Denoting $\mathbf{x} - \mathbf{x}_a = \mathbf{n}_a |\mathbf{x} - \mathbf{x}_a|$ and $\cos\theta_a = \mathbf{n}_a \cdot \dot{\mathbf{x}}_a / |\dot{\mathbf{x}}_a|$, the solution of (3) can thus be written as

$$\Delta^2(|\mathbf{x} - \mathbf{x}_a| \sqrt{1 - v^2 \sin^2 \theta_a})^3 = 3(1 - v^2)^2 \left[8 + 7v \frac{d}{dv} + v^2 \frac{d^2}{dv^2} \right] \left(\frac{1}{|\mathbf{x} - \mathbf{x}_a|} \frac{1}{\sqrt{1 - v^2 \sin^2 \theta_a}} \right). \quad (5)$$

Hence, instead of working out Δ^{-2} , we can evaluate $[8 + 7vd/dv + v^2 d^2/dv^2]^{-1}$; i.e., the elliptic partial differential equation of the fourth order can be simplified into an inhomogeneous linear second order ordinary differential equation. It has a unique solution which is regular at $v = 0$. Even though this solution is quite complicated and contains logarithmic terms such as $\ln \cos\theta_a$, they cancel out when partial derivatives are combined. After somewhat lengthy calculations (the details of which will be given elsewhere), we find the field of a moving source

$$h_{ij}^{\text{TT}}(\mathbf{x}; \mathbf{x}_b, \mathbf{p}_b, \dot{\mathbf{x}}_b) = \frac{G}{|\mathbf{x} - \mathbf{x}_b|} \frac{1}{\bar{m}_b} \frac{1}{y(1+y)^2} \{ [\mathbf{y} \mathbf{p}_b^2 - (\mathbf{n}_b \cdot \mathbf{p}_b)^2 (3y+2)] \delta_{ij} + 2[1 - \dot{\mathbf{x}}_b^2 (1 - 2\cos^2 \theta_b)] p_{bi} p_{bj} \\ + [(2+y)(\mathbf{n}_b \cdot \mathbf{p}_b)^2 - (2+3y - 2\dot{\mathbf{x}}_b^2) \mathbf{p}_b^2] n_{bi} n_{bj} + 2(\mathbf{n}_b \cdot \mathbf{p}_b)(1 - \dot{\mathbf{x}}_b^2 + 2y)(n_{bi} p_{bj} + p_{bi} n_{bj}) \} \\ + O(\bar{m}_b \dot{\mathbf{x}}_b - \mathbf{p}_b)G + O(G^2); \quad (6)$$

here $y = y_b \equiv \sqrt{1 - \dot{\mathbf{x}}_b^2 \sin^2 \theta_b}$. As indicated by the symbol $O(\bar{m}_b \dot{\mathbf{x}}_b - \mathbf{p}_b)$, the last expression gets simplified by using $\dot{\mathbf{x}}_b = \mathbf{p}_b / \bar{m}_b$ and we anticipate that later $O(\bar{m}_b \dot{\mathbf{x}}_b - \mathbf{p}_b)G$ will turn into terms $\sim O(G^2)$.

In order to later suppress field degrees of freedom, we shall turn to the Routh functional (see, e.g., [8])

$$R(\mathbf{x}_a, \mathbf{p}_a, h_{ij}^{\text{TT}}, \dot{h}_{ij}^{\text{TT}}) = H - \frac{1}{16\pi G} \int d^3x \pi^{\text{TT}ij} \dot{h}_{ij}^{\text{TT}}, \quad (7)$$

which is “the Hamiltonian for the particles but the Lagrangian for the field.” While the functional derivatives of the Hamiltonian yield the time derivatives of the canonically conjugated field, the functional derivatives of Routhian vanish if the field equations (3) hold. Their solution (4) is nonradiative and can thus be substituted into the Routh functional without changing the Hamilton equations for the particles. Hereafter the symbol h_{ij}^{TT} is a shortcut for the solution of (3) depending on coordinates, momenta, and velocities of the particles. So the reduced Routhian (7) which becomes Hamiltonian $H(\mathbf{x}_a, \mathbf{p}_a, \dot{\mathbf{x}}_a)$ is

$$H_{\text{lin}}(\mathbf{x}_c, \mathbf{p}_c, \dot{\mathbf{x}}_c) = \sum_a \bar{m}_a - \frac{1}{2} G \sum_{a,b \neq a} \frac{\bar{m}_a \bar{m}_b}{r_{ab}} \left(1 + \frac{p_a^2}{\bar{m}_a^2} + \frac{p_b^2}{\bar{m}_b^2} \right) + \frac{1}{4} G \sum_{a,b \neq a} \frac{1}{r_{ab}} (7\mathbf{p}_a \cdot \mathbf{p}_b + (\mathbf{p}_a \cdot \mathbf{n}_{ab})(\mathbf{p}_b \cdot \mathbf{n}_{ab})) \\ - \frac{1}{4} \sum_a \frac{p_{ai} p_{aj}}{\bar{m}_a} h_{ij}^{\text{TT}}(\mathbf{x} = \mathbf{x}_a; \mathbf{x}_b, \mathbf{p}_b, \dot{\mathbf{x}}_b). \quad (10)$$

Dropping out the total time derivatives in (9) means a canonical transformation, but the new canonical coordinates will not be denoted by primes. In fact, another change of coordinates has to follow since the Hamiltonian (10) is a function of $\dot{\mathbf{x}}_a$. We define new momenta by putting $p'_{ai} = p_{ai} - \partial H / \partial \dot{x}_{ai}$, and then eliminate $\dot{\mathbf{x}}_a$ by introducing the new Hamiltonian $H'(\mathbf{x}_a, \mathbf{p}'_a) = H(\mathbf{x}_a, \mathbf{p}_a(\mathbf{p}'_a), \dot{\mathbf{x}}_a(\mathbf{p}'_a)) - \sum_b \dot{x}_{bi}(\mathbf{p}'_a) \partial H / \partial \dot{x}_{bi}$. Since $\partial H / \partial \dot{x}_a \sim G$, the only change in the Hamiltonian which

$$h_{ij}^{\text{TT}}(\mathbf{x}) = \delta_{ij}^{\text{TT}kl} \sum_b \frac{4G}{\bar{m}_b} \frac{1}{|\mathbf{x} - \mathbf{x}_b|} \frac{p_{bk} p_{bl}}{\sqrt{1 - \dot{\mathbf{x}}_b^2 \sin^2 \theta_b}}. \quad (4)$$

The action of $\delta_{ijkl}^{\text{TT}}$ in Eq. (4) consists of two steps: first, one has to solve the Poisson equation twice and then evaluate a number of partial derivatives. The first step is feasible due to the form of the boosted spherical potential. It can be shown to satisfy the relation (here $v = |\dot{\mathbf{x}}_a|$)

obtained. The field part of the Routhian ,

$$R_f = \frac{1}{16\pi G} \int d^3x \frac{1}{4} (h_{ij,k}^{\text{TT}} h_{ij,k}^{\text{TT}} - \dot{h}_{ij}^{\text{TT}} \dot{h}_{ij}^{\text{TT}}), \quad (8)$$

still needs to be transformed into an explicit function of the particles' variables. Using Gauss's law in the first term and integrating by parts the second term, we arrive at

$$R_f = -\frac{1}{16\pi G} \int d^3x \frac{1}{4} h_{ij}^{\text{TT}} \left(\Delta h_{ij}^{\text{TT}} - \frac{\partial^2}{\partial t^2} h_{ij}^{\text{TT}} \right) + \frac{1}{64\pi G} \\ \times \oint dS_k (h_{ij}^{\text{TT}} h_{ij,k}^{\text{TT}}) - \frac{1}{64\pi G} \frac{d}{dt} \int d^3x (h_{ij}^{\text{TT}} \dot{h}_{ij}^{\text{TT}}). \quad (9)$$

The field equations (3) imply that the first integral (in which the self-interaction term is discarded) directly combines with the “interaction” term containing $\sum \bar{m}_a^{-1} p_{ai} p_{aj} h_{ij}^{\text{TT}}(\mathbf{x}_a)$, so only its coefficient is changed. Remaining terms do not modify the dynamics of the system in our approximation. The Hamiltonian thus takes the form

is linear in G comes into the kinetic term $\sum \bar{m}_a$ from the last change of momenta. This change is exactly cancelled by the sum $\sum_b \dot{x}_{bi}(\mathbf{p}'_a) \partial H / \partial \dot{x}_{bi}$ in $H(\mathbf{x}_a, \mathbf{p}'_a)$. We now make simple substitutions $\dot{x}_{ai} = \frac{\partial \bar{m}_a}{\partial p_{ai}} = \frac{p_{ai}}{\bar{m}_a}$, $\cos \theta_a = \frac{\mathbf{n}_a \cdot \mathbf{p}_a}{|\mathbf{p}_a|}$, $y_a^2 \equiv 1 - \dot{x}_a^2 \sin^2 \theta_a = 1 + \bar{m}_a^{-2} [(\mathbf{n}_a \cdot \mathbf{p}_a)^2 - \mathbf{p}_a^2]$ and, again, omit primes. At this moment we can substitute for h_{ij}^{TT} in (10) the solution (6), in which the above substitutions turn the term $O(\bar{m}_b \dot{x}_b - \mathbf{p}_b)G$ into term $\sim O(G^2)$. In this way, using the shortcut $y_{ba} = \bar{m}_b^{-1} [m_b^2 + (\mathbf{n}_{ba} \cdot \mathbf{p}_b)^2]^{1/2}$, we finally arrive at the Hamiltonian in the first post-Minkowskian approximation:

$$\begin{aligned}
H_{\text{lin}} = & \sum_a \bar{m}_a - \frac{1}{2} G \sum_{a,b \neq a} \frac{\bar{m}_a \bar{m}_b}{r_{ab}} \left(1 + \frac{p_a^2}{\bar{m}_a^2} + \frac{p_b^2}{\bar{m}_b^2} \right) + \frac{1}{4} G \sum_{a,b \neq a} \frac{1}{r_{ab}} (7 \mathbf{p}_a \cdot \mathbf{p}_b + (\mathbf{p}_a \cdot \mathbf{n}_{ab})(\mathbf{p}_b \cdot \mathbf{n}_{ab})) \\
& - \frac{1}{4} G \sum_{a,b \neq a} \frac{1}{r_{ab}} \frac{(\bar{m}_a \bar{m}_b)^{-1}}{(y_{ba} + 1)^2 y_{ba}} [2(2(\mathbf{p}_a \cdot \mathbf{p}_b)^2 (\mathbf{p}_b \cdot \mathbf{n}_{ba})^2 - 2(\mathbf{p}_a \cdot \mathbf{n}_{ba})(\mathbf{p}_b \cdot \mathbf{n}_{ba})(\mathbf{p}_a \cdot \mathbf{p}_b) \mathbf{p}_b^2 + (\mathbf{p}_a \cdot \mathbf{n}_{ba})^2 \mathbf{p}_b^4 \\
& - (\mathbf{p}_a \cdot \mathbf{p}_b)^2 \mathbf{p}_b^2) \frac{1}{\bar{m}_b^2} + 2[-\mathbf{p}_a^2 (\mathbf{p}_b \cdot \mathbf{n}_{ba})^2 + (\mathbf{p}_a \cdot \mathbf{n}_{ba})^2 (\mathbf{p}_b \cdot \mathbf{n}_{ba})^2 + 2(\mathbf{p}_a \cdot \mathbf{n}_{ba})(\mathbf{p}_b \cdot \mathbf{n}_{ba})(\mathbf{p}_a \cdot \mathbf{p}_b) \\
& + (\mathbf{p}_a \cdot \mathbf{p}_b)^2 - (\mathbf{p}_a \cdot \mathbf{n}_{ba})^2 \mathbf{p}_b^2] + [-3\mathbf{p}_a^2 (\mathbf{p}_b \cdot \mathbf{n}_{ba})^2 + (\mathbf{p}_a \cdot \mathbf{n}_{ba})^2 (\mathbf{p}_b \cdot \mathbf{n}_{ba})^2 + 8(\mathbf{p}_a \cdot \mathbf{n}_{ba})(\mathbf{p}_b \cdot \mathbf{n}_{ba})(\mathbf{p}_a \cdot \mathbf{p}_b) \\
& + \mathbf{p}_a^2 \mathbf{p}_b^2 - 3(\mathbf{p}_a \cdot \mathbf{n}_{ba})^2 \mathbf{p}_b^2] y_{ba}]. \tag{11}
\end{aligned}$$

This is our main result. The Hamiltonian for a many-particle gravitating system in post-Minkowskian approximation, i.e., including all terms linear in G , was derived in the closed form entirely in terms of the variables of the particles. Putting $G = 0$ it becomes the standard Hamiltonian of N noninteracting particles in special relativity.

Let us yet note that in [14] the post-Minkowskian action for a helically symmetric binary solution was constructed; however, it turns out that due to the restriction to helical worldlines ambiguities may arise. We will return to this issue elsewhere. It should also be useful to derive the Hamiltonian (11) within the effective field theory approach to gravity [15].

Scattering.—As an application of the Hamiltonian obtained we calculate gravitational scattering of two possibly ultrarelativistic or zero-rest-mass particles. When only terms linear in G are considered, the transferred momentum can be computed as $\Delta \mathbf{p}_1 = \int_{-\infty}^{+\infty} \dot{\mathbf{p}}_1 dt$, integrating along the straight line trajectories of noninteracting particles [$\dot{\mathbf{p}}_1$ is determined from the Hamilton equations using (11)]. If perpendicular separation \mathbf{b} of trajectories ($|\mathbf{b}|$ is the impact factor) in center-of-mass system ($\mathbf{p}_1 = -\mathbf{p}_2 \equiv \mathbf{p}$) is used, $\mathbf{p} \cdot \mathbf{b} = 0$, we find, after evaluating a few simple integrals, that the exchanged momentum in the system is given by

$$\begin{aligned}
\Delta \mathbf{p} = & -2 \frac{\mathbf{b}}{\mathbf{b}^2} \frac{G}{|\mathbf{p}|} \frac{\bar{m}_1^2 \bar{m}_2^2}{\bar{m}_1 + \bar{m}_2} \\
& \times \left[1 + \left(\frac{1}{\bar{m}_1^2} + \frac{1}{\bar{m}_2^2} + \frac{4}{\bar{m}_1 \bar{m}_2} \right) \mathbf{p}^2 + \frac{\mathbf{p}^4}{\bar{m}_1^2 \bar{m}_2^2} \right]. \tag{12}
\end{aligned}$$

The quartic term is all that remains from the field part h_{ij}^{TT} . It is not difficult to show that (12) agrees with the result [16] obtained by a very different method.

The authors benefitted from the exchange program between Charles University, Prague, and Friedrich Schiller University, Jena. T.L. and J.B. acknowledge the partial

support from SFB/TR7 in Jena, from the Grant GAČR 202/06/0041 of the Czech Republic, and of Grants No. LC 06014 and No. MSM 0021620860 of the Ministry of Education. J.B. is also grateful for the support of the Alexander von Humboldt Foundation.

- [1] T. Damour, in *Gravitational Radiation*, edited by N. Deruelle and T. Piran (North-Holland, Amsterdam, 1983), pp. 59–144; in *Three Hundred Years of Gravitation*, edited by S. Hawking and W. Israel (Cambridge University Press, Cambridge, England, 1987), pp. 128–198.
- [2] K. Thorne, in *Three Hundred Years of Gravitation* (Ref. [1]), pp. 330–458.
- [3] L. Blanchet, *Living Rev. Relativity* **9**, 4 (2006) <http://www.livingreviews.org/lrr-2006-4>.
- [4] T. Futamase and Y. Itoh, *Living Rev. Relativity* **10**, 2 (2007) <http://www.livingreviews.org/lrr-2007-2>.
- [5] T. Kimura and T. Toiya, *Prog. Theor. Phys.* **48**, 316 (1972).
- [6] R. Arnowitt, S. Deser, and C. W. Misner, in *Gravitation: An Introduction to Current Research*, edited by L. Witten (Wiley, New York, 1962), pp. 227–265.
- [7] L. D. Landau and E. M. Lifshitz, *The Classical Theory of Fields* (Pergamon, New York, 1975), 4th ed.
- [8] T. Damour, P. Jaranowski, and G. Schäfer, *Phys. Rev. D* **62**, 021501 (2000); *Phys. Rev. D* **63**, 029903(E) (2000); *Phys. Lett. B* **513**, 147 (2001).
- [9] F. J. Kennedy, *J. Math. Phys. (N.Y.)* **16**, 1844 (1975).
- [10] J. D. Jackson, *Classical Electrodynamics* (John Wiley, New York, 1975), 2nd ed.
- [11] T. Damour and G. Schäfer, *J. Math. Phys. (N.Y.)* **32**, 127 (1991).
- [12] P. D. D’Eath, *Black Holes: Gravitational Interaction* (Clarendon Press, Oxford, England, 1996).
- [13] G. Schäfer, *Gen. Relativ. Gravit.* **18**, 255 (1986).
- [14] J. L. Friedman and K. Uryu, *Phys. Rev. D* **73**, 104039 (2006).
- [15] W. D. Goldberger and I. Z. Rothstein, *Phys. Rev. D* **73**, 104029 (2006); *Gen. Relativ. Gravit.* **38**, 1537 (2006).
- [16] K. Westpfahl, *Fortschr. Phys.* **33**, 417 (1985).