

## 

**Citation:** Grund B, Baker JV, Deeks SG, Wolfson J, Wentworth D, Cozzi-Lepri A, et al. (2016) Relevance of Interleukin-6 and D-Dimer for Serious Non-AIDS Morbidity and Death among HIV-Positive Adults on Suppressive Antiretroviral Therapy. PLoS ONE 11(5): e0155100. doi:10.1371/journal.pone.0155100

Editor: Esaki M. Shankar, University of Malaya, MALAYSIA

Received: December 21, 2015

Accepted: April 25, 2016

Published: May 12, 2016

**Copyright:** © 2016 Grund et al. This is an open access article distributed under the terms of the <u>Creative Commons Attribution License</u>, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: Participant level data are not provided as open access due to ethical and legal obligations to the participants in the clinical trials. Data will be available to all qualified researchers upon request to the INSIGHT Executive Steering Committee; contact James D. Neaton at jim@ccbr.umn.edu.

**Funding:** The trials analyzed in this manuscript were supported by grants from the National Institutes of Health (NIH) and Chiron-Novartis. The National Institute of Allergy and Infectious Diseases (NIAID) funded the INSIGHT SMART [U01-Al068641, U01**RESEARCH ARTICLE** 

# Relevance of Interleukin-6 and D-Dimer for Serious Non-AIDS Morbidity and Death among HIV-Positive Adults on Suppressive Antiretroviral Therapy

Birgit Grund<sup>1</sup>\*, Jason V. Baker<sup>2,3</sup>, Steven G. Deeks<sup>4,5</sup>, Julian Wolfson<sup>6</sup>, Deborah Wentworth<sup>6</sup>, Alessandro Cozzi-Lepri<sup>7</sup>, Calvin J. Cohen<sup>8</sup>, Andrew Phillips<sup>7</sup>, Jens D. Lundgren<sup>9</sup>, James D. Neaton<sup>6</sup>, INSIGHT SMART/ESPRIT/SILCAAT Study Group<sup>1</sup>

1 School of Statistics, University of Minnesota, Minneapolis, MN, United States of America, 2 Hennepin County Medical Center, Minneapolis, MN, United States of America, 3 Department of Medicine, University of Minnesota, Minneapolis, MN, United States of America, 4 University of California San Francisco, San Francisco, CA, United States of America, 5 San Francisco General Hospital, San Francisco, CA, United States of America, 6 Division of Biostatistics, School of Public Health, University of Minnesota, Minneapolis, MN, United States of America, 7 University College London, London, United Kingdom, 8 Medical Affairs Department, Gilead Sciences, Foster City, CA, United States of America, 9 Department of Infectious Diseases, Rigshospitalet, University of Copenhagen, Copenhagen, Denmark

 $\$  Complete membership of the author group can be found in the Acknowledgments. \* <u>birgit@ccbr.umn.edu</u>

## Abstract

## Background

Despite effective antiretroviral treatment (ART), HIV-positive individuals are at increased risk of serious non-AIDS conditions (cardiovascular, liver and renal disease, and cancers), perhaps due in part to ongoing inflammation and/or coagulation. To estimate the potential risk reduction in serious non-AIDS conditions or death from any cause that might be achieved with treatments that reduce inflammation and/or coagulation, we examined associations of interleukin-6 (IL-6), D-dimer, and high-sensitivity C-reactive protein (hsCRP) levels with serious non-AIDS conditions or death in 3 large cohorts.

## Methods

In HIV-positive adults on suppressive ART, associations of IL-6, D-dimer, and hsCRP levels at study entry with serious non-AIDS conditions or death were studied using Cox regression. Hazard ratios (HR) adjusted for age, gender, study, and regression dilution bias (due to within-person biomarker variability) were used to predict risk reductions in serious non-AIDS conditions or death associated with lower "usual" levels of IL-6 and D-dimer.

## Results

Over 4.9 years of mean follow-up, 260 of the 3766 participants experienced serious non-AIDS conditions or death. IL-6, D-dimer and hsCRP were each individually associated with



AI046362, U01-AI042170] and ESPRIT [U01-AI068641, U01AI46957] trials and the biomarker analyses [3U 01-Al068641-0451, 3U01-Al06841-0551]. The SILCAAT and ESPRIT trials were supported by grants from Chiron-Novartis. These funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. Individual authors: Gilead Sciences currently provides salary support for CJC, but did not have any additional role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript. CJC was previously employed by the Community Research Initiative New England, Boston, MA, and joined Gilead Sciences only after the data analysis and the first draft of the manuscript were completed. Otherwise, none of the authors received any third-party funding for the study design, data analysis or writing of this manuscript. The specific roles of the authors are articulated in the 'author contributions' section.

**Competing Interests:** The authors have the following potentially competing interests: AP received fees for speaking from Gilead Sciences, consulting from GSK Biologicals, and advisory board membership from AbbVie. CJC is currently receiving salary support from Gilead Sciences. The other authors reported no competing interests. While the SILCAAT and ESPRIT trials were supported by Chiron-Novartis, none of the authors received salary support from Chiron-Novartis, or have any competing interests with respect to ChironNovartis. The potentially competing interests do not alter the authors' adherence to PLOS ONE policies on sharing data and materials. risk of serious non-AIDS conditions or death, HR = 1.45 (95% CI: 1.30 to 1.63), 1.28 (95% CI: 1.14 to 1.44), and 1.17 (95% CI: 1.09 to 1.26) per 2x higher biomarker levels, respectively. In joint models, IL-6 and D-dimer were independently associated with serious non-AIDS conditions or death, with consistent results across the 3 cohorts and across serious non-AIDS event types. The association of IL-6 and D-dimer with serious non-AIDS conditions or death was graded and persisted throughout follow-up. For 25% lower "usual" IL-6 and D-dimer levels, the joint biomarker model estimates a 37% reduction (95% CI: 28 to 46%) in the risk of serious non-AIDS conditions or death if the relationship is causal.

## Conclusions

Both IL-6 and D-dimer are independently associated with serious non-AIDS conditions or death among HIV-positive adults with suppressed virus. This suggests that treatments that reduce IL-6 and D-dimer levels might substantially decrease morbidity and mortality in patients on suppressive ART. Clinical trials are needed to test this hypothesis.

## Introduction

Among HIV-positive adults with high CD4 cell counts on effective antiretroviral therapy (ART), serious non-AIDS conditions (SNA) are the primary cause of severe morbidity and mortality [1–4]. Among HIV-positive adults, higher levels of interleukin-6 (IL-6), D-dimer, and high-sensitivity C-reactive protein (hsCRP) are associated with an increased risk of cardio-vascular disease (CVD) [5–7], cancer [8], and all-cause mortality [9–11]. While ART decreases IL-6, D-dimer and hsCRP levels [10, 12], these biomarkers remain elevated relative to the general population even when the plasma HIV RNA is suppressed [13]. Markers of inflammation and coagulation have been widely studied in the general population, and related to higher risk of CVD [14–22], cancer [23–25], kidney function decline [26] and all-cause mortality [27–29]. These data collectively suggest that chronic inflammation and/or hyper-coagulation contribute to the pathogenesis of these serious non-AIDS events during otherwise effective ART.

Given the assumed, albeit unproven, role of these pathways in causing disease, both vascular and non-vascular, there is intense interest in studying interventions that reduce inflammation and/or coagulation [30, 31]. To date, no single biomarker has been formally validated as a surrogate marker for non-AIDS morbidity. We developed a biomarker score (the "IL-6 & D-dimer score"), which optimally combines markers of inflammation and coagulation to predict the risk of serious non-AIDS conditions or death from any cause (SNA/death) among HIV-infected adults on suppressive ART, and a biomarker model to estimate the effect of sustained long-term reductions in IL-6 and D-dimer levels on the risk of SNA/death. The IL-6 & D-dimer score could be a suitable biomarker endpoint for phase II trials aiming to compare treatments side-by-side for their potential to reduce serious non-AIDS conditions, even if the treatments have different mechanisms of action, or target inflammation and/or coagulation to varying degrees.

## Methods

## Population

Our analysis includes 3766 HIV-positive participants on suppressive ART from the control arms of three studies performed by the INSIGHT trials network: the Strategies for

Management of Antiretroviral Therapy (SMART; 1748 participants) [32, 33], Evaluation of Subcutaneous Proleukin® in a Randomized International Trial (ESPRIT; 1446 participants) and Subcutaneous Recombinant, Human Interleukin-2 in HIV-Infected Patients with Low CD4+ Counts under Active Antiretroviral Therapy (SILCAAT; 572 participants) [34] trials. All participants were using ART at study entry, had HIV RNA levels  $\leq$  500 copies/mL, had been randomized to continuing suppressive ART, and provided consent to store blood for future research. We restricted our analysis to participants who were treated and virally suppressed, because early treatment and viral suppression is increasingly the norm, and because effective ART also decreases levels of inflammation and coagulation [10, 12]. These three studies comprised all large HIV trials conducted by the INSIGHT network in the past 15 years that enrolled ART-experienced participants.

The study was conducted in accordance with the declaration of Helsinki. The SMART, ESPRIT and SILCAAT trials were approved by the Institutional Review Board at the University of Minnesota and the institutional review boards or institutional ethics committees at each of the participating clinical sites worldwide. The Institutional Review Board at the University of Minnesota also approved plans for the analysis of stored specimens for consenting participants. All participants provided written informed consent. Clinicaltrials.gov identifiers: NCT00027352, NCT00004978 and NCT00013611.

#### Outcome measures

The primary outcome was SNA/death, a composite endpoint comprising cardiovascular disease (myocardial infarction, stroke, or death due to cardiovascular disease including sudden death not attributed to other causes), end-stage renal disease, decompensated liver cirrhosis, non-AIDS cancer (excluding non-fatal non-melanoma skin cancer), and death due to any cause. In the SMART and ESPRIT trials, clinical events were assessed by an independent Endpoint Review Committee (ERC), blinded to randomization assignment. Only outcomes assessed as confirmed or probable were included. In the SILCAAT study, SNA events were identified from serious adverse event reports. Non-fatal events were coded according to the Medical Dictionary for Regulatory Activities (MedDRA®) version 12.0 [35], causes of death according to the CoDe system [36].

### Biomarkers

We focused on IL-6, hsCRP, and D-dimer for several reasons. First, in a panel of 6 plasma biomarkers measured in the INSIGHT SMART study, these 3 biomarkers were strongly associated with all-cause mortality, which motivated us to determine these markers also for the ESPRIT and SILCAAT trials, using stored baseline plasma specimens.[6] Second, higher levels of IL-6, hsCRP and D-dimer have been associated with increased risk of CVD, cancer, and all-cause mortality among people living with HIV [5–11], as well as in the general population [14–25, 27–29]. Third, the biomarkers have high laboratory and biological reproducibility. [37]

Levels of IL-6, D-dimer and hsCRP at study entry were measured on plasma collected at baseline for all patients who provided written consent, and stored at -70 degrees Celsius. For the SMART study, the biomarkers were determined centrally at the Laboratory for Clinical Biochemistry Research at the University of Vermont (Burlington), for the ESPRIT and SIL-CAAT trials, by SAIC-Frederick (Frederick, MD). IL-6 was measured by ELISA assays by R&D Systems, Minneapolis, MN, using the QuantiGlo® Chemiluminescent ELISA (range 0.48– 1500 pg/mL) for SMART and the high-sensitivity Quantikine® HS ELISA (range 0.16–10 pg/ mL) for ESPRIT and SILCAAT samples. The QuantiGlo® Chemiluminescent ELISA was chosen for SMART because HIV-positive populations have higher levels of IL-6 than the general population, and this assay's range is wider while still giving sufficient precision at the lower limit. Our laboratories validated that the two R&D ELISA assays provided similar measurements, using split samples. D-dimer was measured by ELISA on the Sta-R analyzer, Liatest D-DI (Diagnostic Stago, Parsippany, NJ, USA) for the SMART study, and on a VIDAS instrument (BioMerieux Inc., Durham, NC, USA) for ESPRIT and SILCAAT. HsCRP was measured by ELISA by both laboratories, using a NBTMII nephelometer, N Antiserum to Human CRP (Siemens Diagnostics) for the SMART study, and an R&D Systems ELISA assay for ESPRIT and SILCAAT. The D-dimer and hsCRP assays, while different, compared very well on duplicate samples.[6] Lower limits of detection for D-dimer and hsCRP were 0.01 and 0.16 µg/mL, respectively, for SMART, and 0.045 and 0.078 µg/mL for ESPRIT and SILCAAT. All samples were analyzed blinded to clinical information about the participants.

### **Statistical Methods**

Associations of baseline biomarker levels with SNA/death. In order to describe associations of individual biomarkers with SNA/death, we computed Kaplan-Meier estimates of the cumulative risk of SNA/death by quartiles of baseline biomarker levels, and estimated hazard ratios (HRs) per 2x higher baseline biomarker levels in separate Cox proportional hazards models. All biomarkers were analyzed on the log<sub>2</sub> scale, because their distributions were skewed on the original scale. All proportional hazards models were adjusted for age (categories 18–40, 41–50 and 51+ years), sex, and study (ESPRIT, SILCAAT, SMART), because these covariates were independently associated with SNA/death in a joint biomarker model.

In order to investigate whether IL-6, D-dimer and hsCRP are independently associated with the risk of SNA/death, we started with a model that contained all three biomarkers, as well as age, sex, race, study indicator, baseline CD4 cell counts (in categories  $\leq$ 500, 500–800, 801 + cells/µL), nadir CD4, and indicator variables for the use of protease inhibitors (PIs) and non-nucleoside reverse transcriptase inhibitors (NNRTIs) at baseline, and stepwise eliminated covariates that did not improve the AIC model fit [38]. The final model contained log<sub>2</sub> IL-6, log<sub>2</sub> D-dimer, age, sex, and the study indicator, each with p-values  $\leq$ 0.05. Because in the general population BMI is associated with the risk of CVD and death, we also tested for an association between BMI and the risk of SNA/death by adding BMI to the final biomarker model. We did not include BMI in the formal backwards variable selection procedure, because we were missing BMI values for 52 participants.

The IL-6 & D-dimer score was determined as the log HR estimate for given IL-6 and Ddimer levels using the final proportional hazards model, which included the two log<sub>2</sub>-transformed biomarkers, age, sex and the study indicator. Therefore, the score is a linear combination of the log<sub>2</sub> biomarkers, weighted by the estimated regression coefficients. We tested the proportional hazards assumption using the product of the score with log-transformed time.

As sensitivity analyses, we estimated associations of IL-6, D-dimer, and the IL-6 & D-dimer score with each of the components of SNA/death in proportional hazards models, again adjusted for age, sex and study indicator. We compared the associations of the IL-6 & D-dimer score with the risk of SNA/death across subgroups by age, sex, and study cohort by testing for the interaction effects between score and subgroup indicator. In order to assess whether the association between the IL-6 & D-dimer score and the risk of SNA/death might have been driven by the presence of subclinical disease at baseline, we performed two analyses. First, we estimated the associations with events during the first 3 years (censoring later follow-up) and with later events (by excluding participants with events prior to year 3), and compared the associations during the earlier versus later time periods by adding the time-updated indicator function of years 4–10 to the proportional hazards model. Second, we compared the

associations of the IL-6 & D-dimer score with the risk of SNA/death in participants who had experienced non-fatal SNA events prior to enrollment to the associations between score and risk in participants without prior SNA events, by testing for an interaction effect between the prior event status and the score. The latter analysis was restricted to the SMART and ESPRIT participants, because the history of prior non-AIDS events was not collected in the SILCAAT study.

**Correction for regression dilution to estimate the effect of reducing the "usual" biomarker levels.** Associations between a person's "usual" biomarker levels (within-person longterm average levels) with the risk of SNA/death are stronger than those of single, baseline biomarker measurements, due to measurement error and random within-person fluctuations in the biomarkers over time. To correct for this regression dilution, the reliability coefficients for IL-6, D-dimer, and the IL-6 & D-dimer score were estimated using biomarker data collected at baseline, 1 and 3 years for 235 participants on stable ART in the ESPRIT study [39, 40]. To estimate the reliability coefficients, variance components were estimated using longitudinal mixed models for the biomarkers and the score, adjusted for visit by treatment group, age and sex. Similar adjustments for regression dilution bias have been carried out in previous studies of IL-6 and D-dimer [14–16]. After adjusting for regression dilution bias, we estimated the potential effect of a therapy-induced, sustained reduction of 0%-40% in IL-6 and/or D-dimer levels on the risk of SNA/death using proportional hazards models, and summarized the estimates and pointwise 95% confidence limits in contour plots.

We estimated the mean change in CD4 cell counts through follow-up using a longitudinal mixed model with random intercept and fixed slope, adjusted for study indicator.

Analyses used SAS version 9.3 (SAS Institute, Cary, NC, USA) and R version 3.1 [41]. All p-values are two-sided.

## Results

#### **Baseline characteristics**

Baseline characteristics of the 3766 participants are summarized in Table 1. The median age was 42 years and 21% were women. At study entry, median time since HIV diagnosis was 7.3 years and participants had used ART for a median of 4.9 years. All participants had plasma HIV RNA levels  $\leq$  500 copies/mL. The median CD4 cell count was 500 cells/µL, median nadir 181 cells/µL. Twenty-seven percent had a prior AIDS diagnosis. Median (IQR) for IL-6, D-dimer and hsCRP were 1.7 (1.1–2.7) pg/mL, 0.22 (0.15–0.35) µg/mL and 1.54 (0.67–3.60) µg/mL, respectively (Table 1). The three biomarkers were pairwise correlated; Spearman's correlations between IL-6 and D-dimer, IL-6 and hsCRP, and D-dimer and hsCRP were 0.31, 0.46, and 0.24, respectively (p<0.001 each).

#### Associations of baseline biomarker levels with SNA/death

Over a mean follow-up of 4.9 years, 260 participants experienced SNA/death (1.44 per 100 person-years, 95% CI: 1.27 to 1.63). The first event was non-AIDS cancer for 97 participants (37%), CVD for 82 (32%; 45 myocardial infarction, 19 stroke, 18 deaths due to CVD), decompensated liver cirrhosis for 19 (7%), end-stage renal disease for 8 (3%), and death due to other causes for 54 (21%) participants, including 14 deaths due to AIDS, and 8 due to violence, accident or suicide (Fig 1).

Fig 2, panels A-C, show Kaplan-Meier curves for time to SNA/death by quartiles of IL-6, Ddimer and hsCRP at study entry. High baseline levels of IL-6 and D-dimer predict an elevated risk of SNA/death over 5 years and longer; by year 5, the cumulative rate of SNA/death was 12.0% among participants in the highest quartile of IL-6, compared with 3.5% for participants

#### Table 1. Baseline characteristics <sup>a</sup>.

	Median (IQR) or %							
Characteristic	Overall		SMART	ESPRIT (n = 1446)	SILCAAT (n = 572)			
	(n = 3766)		(n = 1748)					
Age (years)	42	(37, 49)	45	40	41			
Female (%)	21.2		25.8	17.4	16.6			
Race/Ethnicity (%)								
Black	16.4		24.9	8.6	10.1			
White/Latino/Other	83.6		75.1	91.4	89.9			
Location of Enrollment (%)								
Asia	3.4		0.7	8.0	-			
Australia, Europe, Israel	46.9		37.0	52.3	63.1			
North America	37.4		49.0	23.5	36.9			
South America	11.8		12.8	15.3	-			
Africa	0.6		0.5	0.9	-			
HIV-related factors								
CD4 cell count (cells/mm <sup>3</sup> )	500	(365, 690)	654	452	205			
Nadir CD4 (cells/mm <sup>3</sup> )	181	(70, 290)	230	176	57			
Prior AIDS (%)	27.1		24.9	28.1	31.3			
Years since HIV diagnosis	7.3	(4.0, 11.4)	8.4	6.0	6.3			
Duration of ART use (years)	4.9	(2.5, 8,0)	6.0	4.2	3.9			
Current use of a PI (%)	47.8		42.1	47.9	64.9			
Current use of a NNRTI (%)	50.8		52.0	51.7	44.8			
BMI (kg/m²)	24.1	(22.0, 26.6)	24.7	23.7	23.7			
Biomarkers								
IL-6 (pg/mL)	1.7	(1.1, 2.7)	1.6	1.9	1.7			
D-dimer (µg/mL)	0.22	(0.15, 0.35)	0.18	0.26	0.25			
hsCRP (µg/mL)	1.54	(0.67, 3.60)	1.72	1.42	1.34			

Abbreviations: ART = antiretroviral therapy, hsCRP = high-sensitivity C-reactive protein, IL-6 = Interleukin-6, IQR = interquartile range, NNRTI = nonnucleoside reverse transcriptase inhibitor, PI = protease inhibitor.

<sup>a</sup> All 3766 participants were using suppressive ART at baseline (HIV RNA  $\leq$  500 cp/mL), and were randomized to continue their ART regimen.

doi:10.1371/journal.pone.0155100.t001

in the lowest two quartiles (Fig 2A). The spread of the curves for D-dimer and hsCRP quartiles was less dramatic but still significant (Fig 2B and 2C).

Estimated hazard ratios of SNA/death per 2x higher baseline levels of IL-6, D-dimer and hsCRP are summarized in Fig.3A. In separate models, IL-6, D-dimer and hsCRP were each individually associated with risk of SNA/death; the estimated hazard ratios per 2x higher biomarker levels were 1.45 (95% CI: 1.30 to 1.63) for IL-6, 1.28 (95% CI: 1.14 to 1.44) for D-dimer, and 1.17 (95% CI: 1.09 to 1.26) for hsCRP; p<0.001 for each. All models were adjusted for age, sex and study.

Mean CD4 cell count levels increased slightly through follow-up, by 14 cells/mm<sup>3</sup> per year.

## Development of the IL-6 & D-dimer biomarker score

We obtained a parsimonious joint biomarker model by applying backwards variable selection to a proportional hazards regression model containing all three biomarkers along with several other baseline factors. In the resulting *joint biomarker model*, both IL-6 and D-dimer were independently associated with the risk of SNA/death (Fig 3A, joint model), and the log hazard



## Type of the first SNA/death event




Fig 1. Composition of the serious non-AIDS disease or death (SNA/death) endpoint. Only the first event per participant is counted.

doi:10.1371/journal.pone.0155100.g001


ratio for SNA/death per log<sub>2</sub> IL-6 and log<sub>2</sub> D-dimer units higher is estimated by the

 $IL-6\&D - dimer \text{ score}: 0.33 \times \log_2 IL-6 + 0.16 \times \log_2 D - dimer.$ 


Higher levels of the IL-6 & D-dimer score were associated with increased risk of SNA/death (Figs 2D and 4) and each of its components (Fig 4). In particular, 2x higher levels of both IL-6 and D-dimer correspond to 1 unit higher for each of the two log<sub>2</sub> biomarker components of the score, an IL-6 & D-dimer score of 0.49, and an estimated hazard ratio of 1.64 (95% CI: 1.43 to 1.89; p<0.001) for SNA/death.

In the joint biomarker model, the estimated risk of SNA/death was higher for older participants (HR = 2.0 for ages 41–50 versus 18–40 years, 3.3 for older than 50 years versus 18–40 years, both p<0.001), men compared with women (HR = 1.62, p = 0.01), and for participants in the SILCAAT study, which enrolled subjects with lower CD4 T cell counts (Table 1). The associations of the IL-6 & D-dimer score with the risk of SNA/death, however, were homogeneous across age, sex, or study (p >0.99, 0.94, and 0.70, respectively, for the interactions between the score and the factors). Baseline hsCRP, CD4, CD4 nadir, and use of PIs or NNRTIs at baseline did not contribute independently to the risk and were excluded by the variable selection procedure. Similarly, BMI was not independently associated with the risk of SNA/death in the joint biomarker model (p = 0.08).

Associations of the IL-6 & D-dimer score with the risk of SNA/death remained stable over time (p = 0.45 for a test of proportional hazards using log time). During the first 3 years, 2x higher baseline levels of both IL-6 and D-dimer were associated with a hazard ratio of 1.65 (95% CI: 1.37 to 2.00; p < 0.001), while the hazard ratio was 1.61 (95% CI: 1.31 to 1.98; p < 0.001) after the first 3 years (p = 0.69 for a difference in hazard ratios by time period). Associations were also consistent across the SMART, ESPRIT and SILCAAT studies (p = 0.86 for interaction between the IL-6 & D-dimer score and study indicator), across age groups (p = 0.95), and for men compared with women (p = 0.89).



q1:944



q1:1038

C. hsCRP Quartiles Quartiles: Quartiles: Cum. % of pts with event Cum. % of pts with event q4 (highest) q4 (highest) q3 q3 q2 q2 q1 (lowest) q1 (lowest) Years Years No. at risk: No. at risk: q4:941 q4:942 q3:940 q3:941 q2:938 q2:941 q1:947 q1:942 

Fig 2. Kaplan-Meier estimates of the cumulative proportion of participants who experienced serious non-AIDS disease or death (SNA/death), by quartiles of IL-6 (A), D-dimer (B), hsCRP (C), and the IL-6 & D-dimer score (D). Biomarkers were measured

PLOS ONE | DOI:10.1371/journal.pone.0155100 May 12, 2016

## D. IL-6 & D-Dimer Score Quartiles

at study entry. At 5 years, the risk of SNA/death was 12.0% for participants in the highest quartile of IL-6, compared with 3.5% for participants in the two lowest quartiles. Abbreviations: IL-6 = Interleukin-6; hsCRP = high-sensitivity C-reactive protein.

doi:10.1371/journal.pone.0155100.g002

Non-fatal non-AIDS events prior to study entry were reported in the SMART and ESPRIT studies. When excluding the 146 (4.6%) participants who had a prior myocardial infarction (35), stroke (28), non-AIDS cancer (42), liver cirrhosis or hepatic steatosis (42), or end-stage kidney disease (5), the hazard ratio for SNA/death among SMART and ESPRIT participants changed from 1.62 to 1.57 (95% CI: 1.31 to 1.87, p<0.001) per 2x higher baseline levels of IL-6 and D-dimer; associations between the IL-6 and D-dimer score and the risk of SNA events were similar for those with and without a history of prior SNA (p = 0.48 for the interaction).

**Predicting the effect of reducing "usual" levels of IL-6 and/or D-dimer through therapy.** Estimated reliability coefficients were 0.39 (95% CI: 0.28 to 0.50) for IL-6, 0.53 (95% CI: 0.44 to 0.62) for D-dimer, and 0.43 (95% CI: 0.33 to 0.54) for the IL-6 & D-dimer score. Associations of "usual" (within-person long-term average) biomarker levels with the risk of SNA/ death are summarized in Fig 3B; in the joint biomarker model, the estimated hazard ratio per 2 times higher "usual" biomarker levels was 2.16 (95% CI: 1.64 to 2.84) for IL-6, 1.45 (95% CI: 1.09 to 1.93) for D-dimer, and 3.16 (95% CI: 2.28 to 4.37) for both. These hazard ratios were estimated in proportional hazards models, adjusted for regression dilution, and for age, sex and study.

Based on the joint biomarker model, after adjustment for regression dilution, we estimate that a sustained, long-term reduction in levels of IL-6 and D-dimer by 25% would result in a 37% reduction in the risk of SNA/death (95% CI: 28% to 46%) (Fig\_5). The contour plot in Fig 5A shows the estimated reduction in the risk of SNA/death for combinations of different amounts (0%-40%) of sustained reductions in the "usual" IL-6 and/or D-dimer levels. A similar risk reduction by 37% would also be predicted for a sustained reduction in IL-6 by 30% and D-dimer by 12%, for example, or for a reduction in IL-6 by 20% and D-dimer by 33%. Limits of 95% confidence intervals are provided in Fig 5B and 5C.

#### Discussion

Clinical endpoint trials are needed to establish whether lowering levels of inflammation and/or coagulation among HIV-positive individuals will decrease the risk of serious non-AIDS conditions, the leading cause of severe morbidity and mortality in virally well-controlled HIV infection. [1-4] We have developed a novel IL-6 & D-dimer biomarker score, which could be used as a primary biomarker outcome in phase II trials aiming to compare side-by-side drugs with different mechanisms of action (decrease of inflammation and/or coagulation), by evaluating treatments with respect to their model-predicted potential to reduce SNA/death.

We have shown that a single measurement of IL-6, D-dimer or hsCRP at baseline predicted the risk of SNA/death over 5 years and longer in a large cohort of HIV-positive individuals on suppressive ART regimens with moderate or high CD4 cell counts. Higher baseline levels of IL-6, D-dimer and hsCRP were associated with increased risk of SNA/death when considered separately, and IL-6 and D-dimer were independently associated with the risk in a joint biomarker model.

We combined CVD, cancer, serious renal and hepatic disease, and all-cause mortality into the composite outcome of SNA/death, because each of these conditions had been linked individually to high levels of inflammation and/or coagulation in the general population as well as among HIV-positive individuals. Markers of inflammation and coagulation have been most widely studied for their association with CVD [6, 7, 14–22] and all-cause mortality [10, 11, 27–29]. Currently, a clinical trial is ongoing to test whether reduction in inflammation through

<b>PLOS</b>	ONE
-------------	-----

			Biomarker R	β egr. Coef.	HR per 2–fold higher biomarker (95% Cl)	P−val.
Α.	ine at) <sup>a</sup>	0	log2 <b>IL-6</b>	0.37	1.45	<0.001
	t baseli uremen	Separate models	log <sub>2</sub> D-dimer	0.25	1.28	<0.001
	r level a e meas	Ø	log <sub>2</sub> hsCRP	0.16	1.17	<0.001
	Biomarker level at baseline (one-time measurement) <sup>³</sup>	Joint <sub>b</sub> model	log <sub>2</sub> IL-6	0.33	1.40	<0.001
	Bi	or	log <sub>2</sub> D-dimer	0.16	1.17	0.01
В.	'Usual' biomarker levels (within–person average) <sup>c</sup>	Separate models	log₂ <b>IL−6</b>	0.96	2.60	<0.001
	omarke erson á	omarke erson a Se n	log <sub>2</sub> D-dimer	0.46	1.59	<0.001
	sual' bio ithin-p	Joint model	log2 <b>IL-6</b>	0.77	2.16	<0.001
	ň گ	oL	log <sub>2</sub> D-dimer	0.37	1.45	0.01
					1 1.5 2 2.5 3	

<sup>a</sup> HRs were estimated in Cox proportional hazards models, adjusted for age, sex and study.

- <sup>b</sup> The joint model was obtained through backwards variable selection with AIC, starting with a proportional hazards model for SNA/death that contained log2-transformend IL-6, D-dimer, and hsCRP, age, sex, study indicator, and the following baseline variables: CD4 and nadir CD4 cell counts, use of protease inhibitors (PI), use of non-nucleoside reverse transcriptase inhibitors (NNRTI). HsCRP, CD4, nadir CD4, PI and NNRTI use were not independently associated with SNA/death and were eliminated by the automatic variable selection procedure from the joint model.
- <sup>c</sup> HR estimates were adjusted for regression dilution, age, sex and study. Reliability coefficients for IL–6, D–dimer, and the IL–6 & D–dimer score (joint model) were 0.39, 0.53, and 0.43, respectively, estimated from 235 ESPRIT participants with IL–6 and D–dimer data at baseline, year 1 and year 3.

Fig 3. Hazard ratios (HRs) for serious non-AIDS conditions or death (SNA/death) per 2x higher levels of IL-6, D-dimer, and hsCRP. Panel A shows HRs for differences in baseline biomarker levels (a one-time measurement), estimated using proportional hazards models for each biomarker separately, and in a joint biomarker model containing both IL-6 and D-dimer. Panel B shows HRs for differences in "usual" (the within-subject long-term average) biomarker levels, estimated in separate and joint biomarker models after adjustment for regression dilution bias. All models were adjusted for age, sex, and study indicator (SMART, ESPRIT, SILCAAT). Abbreviations: IL-6 = Interleukin-6; hsCRP = high-sensitivity C-reactive protein.

doi:10.1371/journal.pone.0155100.g003

low-dose methotrexate reduces CVD or death in the general population [42], as well as a trial to test whether statin therapy prevents CVD among HIV-infected individuals using ART [43]. However, systemic inflammation and activation of coagulation appear to have broader effects beyond CVD. Elevated levels of IL-6, hsCRP and/or D-dimer were predictive of cancer, in particular death due to cancer [16, 23, 24, 44]; high D-dimer levels were linked to cancer mortality more strongly than to CVD [28]. Higher levels of inflammation and coagulation are also predictive of kidney function decline [26], and liver disease has been linked to increased coagulation [45]. In our cohort, associations of elevated IL-6 & D-dimer scores with the risk of cancer, kidney and liver disease, and all-cause mortality individually were at least as strong as associations with CVD. Associations of the IL-6 & D-dimer score with the risk of SNA/death were also consistent across studies (SMART, ESPRIT, and SILCAAT), across age groups, similar for men and women, and across baseline ART regimens.

Our biomarker models suggest that lowering levels of inflammation and/or coagulation might substantially decrease non-AIDS morbidity and mortality among HIV-positive patients on suppressive ART regimens, if inflammation and coagulation are on the causal pathway. Suitable agents to test this hypothesis in a clinical trial could target inflammation or coagulation to varying degrees, but would have to be safe for long-term use in combination with ART. Phase II trials would be needed to evaluate safety along with the potential to decrease inflammation or coagulation. The IL-6 & D-dimer score combines markers of inflammation and coagulation optimally to estimate the risk of SNA/death, and such could be used as biomarker outcome in pilot trials to compare several candidate agents.

Within the limitations of a prospective cohort study, we performed analyses to address the possibility that the associations of elevated baseline levels of IL-6 and D-dimer with the risk of SNA/death might have been due to subclinical disease which increased the biomarker levels as well as the subsequent risk of clinical disease. In our cohort, the strength of the associations between baseline biomarkers and risk of SNA/death remained stable through follow-up in, and associations of the IL-6 & D-dimer score with SNA/death were similar for participants with and without a prior history of non-fatal SNA events. These findings suggest that the associations are not explained by subclinical disease.

The main strength of our study was the large cohort of well-characterized participants, centrally determined biomarkers, the standardized definition of SNA events across the cohort, and the rigorous adjudication of the events in the SMART and ESPRIT cohorts by an independent ERC. The study has several limitations. First, we considered only three biomarkers; other markers might provide even stronger risk predictions. Second, the biomarkers were determined in two different central laboratories. Analyses of 20 repeated samples across the laboratories showed similar values, however [6]; also, all models included the study indicator, and thereby adjusted for potential differences between the two laboratories. Third, biomarker levels are highly variable within subjects, both due to assay variability as well as due to natural fluctuations. While we adjusted our estimates for the resulting regression dilution bias, the statistical uncertainty inherent in the biomarker measures resulted in large confidence intervals for the hazard ratios. Lastly, the SMART, ESPRIT and SILCAAT trials were conducted in 1997–2008, and it is possible that results would be slightly different in a population receiving more current ART regimens.

P-value

< 0.001

0.009

< 0.001

< 0.001

< 0.001

< 0.001

0.03

0.06

<0.001

< 0.001

< 0.001

0.003

< 0.001

< 0.001

< 0.001

Biomarker	Event type	No. of pts with event	HR per 2x higher biomarker( with 95% Cl
IL-6	SNA or death	260	1.45
	CVD	84	
	Cancer	99	1.39
	Renal or hepatic	31	2.35
	All-cause death	144	<u>1.7</u> 0
D-dimer	SNA or death	260	1.28
	CVD	84	1.26
	Cancer	99	
	Renal or hepatic	31	<b>1.71</b>
	All-cause death	144	1.43
IL−6 & D−dimer	SNA or death	260	1.64
	CVD	84	↓ <u>1.48</u>
	Cancer	99	1152
	Renal or hepatic	31	, <mark>-3</mark> .1
	All-cause death	144	<u>2.03</u>

Fig 4. Estimated hazard ratios (HR) for the composite endpoint of serious non-AIDS or death (SNA/death) and its components. Hazard ratios are estimated per 2x higher level of IL-6 and D-dimer, with 95% confidence intervals. Two times higher biomarker levels corresponds to 1 unit higher on the log<sub>2</sub> biomarker scale, and 0.49 units higher for the IL-6 & D-dimer score. Hazard ratios were estimated in proportional hazards models adjusted for age, sex and study. *Footnotes*: Of the 144 deaths, 37 were due to non-AIDS cancer, 33 CVD, 11 non-cancer hepatic disease, 1 renal failure, 15 AIDS-related, 9 due to violence, accident or suicide, 38 due to other or unknown causes. Abbreviations: CI = confidence interval; CVD = cardiovascular disease (myocardial infarction, stroke, and death due to CVD); HR = hazard ratio.

0.8

1

1.5

2


3

4

doi:10.1371/journal.pone.0155100.g004

PLOS ONE

In summary, both IL-6 and D-dimer independently predict the risk of serious non-AIDS conditions or death among HIV-positive patients with suppressed virus and moderate to high




A. Estimated Reduction in the

Reduction in IL-6 as % of baseline

## B. Lower Limit of the 95% CI

## C. Upper Limit of the 95% CI



**Fig 5. Estimated risk reduction.** Panel A: Contour plot of the estimated reduction in the risk of serious non-AIDS conditions or death (SNA/death) when "usual" (within-person long-term average) levels of IL-6 and/or D-dimer are decreased by 0–40%, possibly through an intervention. Solid lines show predicted risk reductions of 5%, 10%, etc. Panels B and C show the lower and upper 95% confidence limits, respectively. For a 25% reduction in IL-6 and D-dimer, the risk of SNA/death is predicted to decline by 37% (95% CI: 28 to 46%). *Footnotes:* The reduction in the risk was estimated using Cox proportional hazards models, adjusted for regression dilution bias, and for age, sex, and study. Reliability coefficients for the regression dilution adjustment were estimated using longitudinal mixed models for biomarker data collected at baseline, years 1 and 3 for 235 participants on stable ART in the ESPRIT study. Abbreviations: CI = confidence interval; IL-6 = Interleukin-6.

doi:10.1371/journal.pone.0155100.g005

CD4 cell counts. Treatments to decrease IL-6 and D-dimer might substantially reduce morbidity and mortality in HIV-positive patients on suppressive ART, since non-AIDS conditions dominate morbidity in this population. Clinical trials are needed to test this hypothesis. The IL-6 & D-dimer score could be a suitable biomarker outcome in phase II trials to compare drugs with different mechanisms of action (targeting inflammation, coagulation, or both) for their estimated, model-predicted potential to reduce non-AIDS morbidity and mortality.

## **Supporting Information**

**S1 File. Abbreviations.** Abbreviations in alphabetical order. (DOCX)

## Acknowledgments

We gratefully acknowledge the participants, staff and clinical investigators of the SMART, ESPRIT and SILCAAT trials. The INSIGHT SMART/ESPRIT/SILCAAT Study Group represents the investigators of the three trials. The investigators of the SMART study are listed in [32] and [33], ESPRIT and SILCAAT investigators in [34]. We thank Gary Collins and Jacqueline Neuhaus, University of Minnesota, Minneapolis, MN, USA, for statistical support, and Lewis Kuller, University of Pittsburg, Pittsburgh, PA, USA, and Russell Tracy, Univ. of Vermont, Burlington, VT, USA, for helpful discussions.

## **Author Contributions**

Conceived and designed the experiments: BG JVB SGD JW CJC AP JDL JDN. Analyzed the data: BG DW. Wrote the paper: BG JVB SGD JW CJC AC-L AP JDL JDN.

## References

- Achhra AC, Amin J, Law MG, Emery S, Gerstoft J, Gordin FM, et al. Immunodeficiency and the risk of serious clinical endpoints in a well studied cohort of treated HIV-infected patients. AIDS. 2010; 24 (12):1877–86. doi: 10.1097/QAD.0b013e32833b1b26 PMID: 20588170; PubMed Central PMCID: PMCPMC2902669.
- Antiretroviral Therapy Cohort Collaboration. Causes of death in HIV-1-infected patients treated with antiretroviral therapy, 1996–2006: collaborative analysis of 13 HIV cohort studies. Clinical Infectious Diseases. 2010; 50(10):1387–96. doi: <u>http://dx.doi.org/10.1086/652283</u> PMID: <u>20380565</u>; PubMed Central PMCID: PMCNIHMS308097 PMC3157754.
- INSIGHT START Study Group, Lundgren JD, Babiker AG, Gordin F, Emery S, Grund B, et al. Initiation of Antiretroviral Therapy in Early Asymptomatic HIV Infection. N Engl J Med. 2015; 373(9):795–807. doi: <u>10.1056/NEJMoa1506816</u> PMID: <u>Medline:26192873</u>.
- Mocroft A, Phillips AN, Gatell J, Horban A, Ledergerber B, Zilmer K, et al. CD4 cell count and viral loadspecific rates of AIDS, non-AIDS and deaths according to current antiretroviral use. AIDS. 2013; 27 (6):907–18. PMID: Medline:23698060. doi: 10.1097/QAD.0b013e32835cb766
- Duprez DA, Neuhaus J, Kuller LH, Tracy R, Belloso W, De Wit S, et al. Inflammation, coagulation and cardiovascular disease in HIV-infected individuals. PLoS One. 2012; 7(9):e44454. Epub 2012/09/13. doi: <u>10.1371/journal.pone.0044454</u> PMID: <u>22970224</u>; PubMed Central PMCID: PMCPmc3438173.
- Nordell AD, McKenna M, Borges AH, Duprez D, Neuhaus J, Neaton JD, et al. Severity of cardiovascular disease outcomes among patients with HIV is related to markers of inflammation and coagulation. J Am Heart Assoc. 2014; 3(3). PMID: Medline:24870935.
- Triant VA, Meigs JB, Grinspoon SK. Association of C-reactive protein and HIV infection with acute myocardial infarction. J Acquir Immune Defic Syndr. 2009; 51(3):268–73. PMID: <u>Medline:19387353</u>. doi: <u>10.1097/QAI.0b013e3181a9992c</u>
- Borges AH, Silverberg MJ, Wentworth D, Grulich AE, Fatkenheuer G, Mitsuyasu R, et al. Predicting risk of cancer during HIV infection: the role of inflammatory and coagulation biomarkers. AIDS. 2013; 27 (9):1433–41. Epub 2013/08/16. doi: <u>10.1097/QAD.0b013e32835f6b0c</u> PMID: <u>23945504</u>; PubMed Central PMCID: PMCPmc4046103.
- Achhra AC, Amin J, Sabin C, Chu H, Dunn D, Kuller LH, et al. Reclassification of risk of death with the knowledge of D-dimer in a cohort of treated HIV-infected individuals. AIDS. 2012; 26(13):1707–17. Epub 2012/05/23. doi: <u>10.1097/QAD.0b013e328355d659</u> PMID: <u>22614887</u>.

- Kuller LH, Tracy R, Belloso W, De Wit S, Drummond F, Lane HC, et al. Inflammatory and coagulation biomarkers and mortality in patients with HIV infection. PLoS Med. 2008; 5(10):e203. doi: <u>http://dx.doi.org/10.1371/journal.pmed.0050203</u> PMID: <u>18942885</u>.
- Tien PC, Choi AI, Zolopa AR, Benson C, Tracy R, Scherzer R, et al. Inflammation and mortality in HIVinfected adults: analysis of the FRAM study cohort. J Acquir Immune Defic Syndr. 2010; 55(3):316–22. PMID: Medline:20581689. doi: 10.1097/QAI.0b013e3181e66216
- Baker JV, Neuhaus J, Duprez D, Kuller LH, Tracy R, Belloso WH, et al. Changes in inflammatory and coagulation biomarkers: a randomized comparison of immediate versus deferred antiretroviral therapy in patients with HIV infection. J Acquir Immune Defic Syndr. 2011; 56(1):36–43. PMID: <u>Medline:20930640</u>. doi: <u>10.1097/QAI.0b013e3181f7f61a</u>
- Neuhaus J, Jacobs DR, Baker JV, Calmy A, Duprez D, La Rosa A, et al. Markers of inflammation, coagulation, and renal function are elevated in adults with HIV infection. The Journal of Infectious Diseases. 2010; 201(12):1788–95. doi: <u>10.1086/652749</u> PMID: <u>20446848</u>; PubMed Central PMCID: PMCPMC2872049.
- Danesh J, Kaptoge S, Mann AG, Sarwar N, Wood A, Angleman SB, et al. Long-term interleukin-6 levels and subsequent risk of coronary heart disease: two new prospective studies and a systematic review. PLoS Med. 2008; 5(4):e78. PMID: <u>18399716</u>; PubMed Central PMCID: PMCPMC2288623. doi: <u>10.</u> <u>1371/journal.pmed.0050078</u>
- Danesh J, Whincup P, Walker M, Lennon L, Thomson A, Appleby P, et al. Fibrin D-dimer and coronary heart disease: prospective study and meta-analysis. Circulation. 2001; 103(19):2323–7. PMID: <u>11352877</u>.
- 16. Emerging Risk Factors Collaboration, Kaptoge S, Di Angelantonio E, Lowe G, Pepys MB, Thompson SG, et al. C-reactive protein concentration and risk of coronary heart disease, stroke, and mortality: an individual participant meta-analysis. Lancet. 2010; 375(9709):132–40. PMID: 20031199; PubMed Central PMCID: PMCPMC3162187. doi: 10.1016/S0140-6736(09)61717-7
- Pai JK, Pischon T, Ma J, Manson JE, Hankinson SE, Joshipura K, et al. Inflammatory markers and the risk of coronary heart disease in men and women. N Engl J Med. 2004; 351(25):2599–610. PMID: Medline:15602020.
- Di Castelnuovo A, Agnoli C, de Curtis A, Giurdanella MC, Sieri S, Mattiello A, et al. Elevated levels of Ddimers increase the risk of ischaemic and haemorrhagic stroke. Findings from the EPICOR Study. Thrombosis and Haemostasis. 2014; 112(2014-07-17). doi: 10.1160/TH14-04-0297
- Luc G, Bard J-M, Juhan-Vague I, Ferrieres J, Evans A, Amouyel P, et al. C-reactive protein, interleukin-6, and fibrinogen as predictors of coronary heart disease: the PRIME Study. Arterioscler Thromb Vasc Biol. 2003; 23(7):1255–61. PMID: Medline:12775578.
- Ridker PM, Rifai N, Stampfer MJ, Hennekens CH. Plasma concentration of interleukin-6 and the risk of future myocardial infarction among apparently healthy men. Circulation. 2000; 101(15):1767–72. PMID: Medline:10769275.
- Smith A, Patterson C, Yarnell J, Rumley A, Ben-Shlomo Y, Lowe G. Which hemostatic markers add to the predictive value of conventional risk factors for coronary heart disease and ischemic stroke? The Caerphilly Study. Circulation. 2005; 112(20):3080–7. PMID: <u>Medline:16286603</u>.
- Kuller LH, Tracy RP, Shaten J, Meilahn EN. Relation of C-reactive protein and coronary heart disease in the MRFIT nested case-control study. Multiple Risk Factor Intervention Trial. American Journal of Epidemiology. 1996; 144(6):537–47. PMID: <u>Medline:8797513</u>.
- Allin KH, Bojesen SE, Nordestgaard BG. Baseline C-reactive protein is associated with incident cancer and survival in patients with cancer. Journal of Clinical Oncology: official journal of the American Society of Clinical Oncology. 2009; 27(13):2217–24. doi: <u>10.1200/JCO.2008.19.8440</u> PMID: <u>Medline:19289618</u>.
- Il'yasova D, Colbert LH, Harris TB, Newman AB, Bauer DC, Satterfield S, et al. Circulating levels of inflammatory markers and cancer risk in the health aging and body composition cohort. Cancer Epidemiol Biomarkers Prev. 2005; 14(10):2413–8. PMID: Medline:16214925.
- Tian G, Mi J, Wei X, Zhao D, Qiao L, Yang C, et al. Circulating interleukin-6 and cancer: A meta-analysis using Mendelian randomization. Sci Rep. 2015; 5:11394. PMID: <u>Medline:26096712</u>. doi: <u>10.1038/</u> <u>srep11394</u>
- Bash LD, Erlinger TP, Coresh J, Marsh-Manzi J, Folsom AR, Astor BC. Inflammation, hemostasis, and the risk of kidney function decline in the Atherosclerosis Risk in Communities (ARIC) Study. Am J Kidney Dis. 2009; 53(4):596–605. PMID: <u>Medline:19110358</u>. doi: <u>10.1053/j.ajkd.2008.10.044</u>
- Di Castelnuovo A, de Curtis A, Costanzo S, Persichillo M, Olivieri M, Zito F, et al. Association of Ddimer levels with all-cause mortality in a healthy adult population: findings from the MOLI-SANI study. Haematologica. 2013; 98(9):1476–80. doi: <u>10.3324/haematol.2012.083410</u> PMID: <u>23645692</u>

- Folsom AR, Delaney JAC, Lutsey PL, Zakai NA, Jenny NS, Polak JF, et al. Associations of factor VIII (c), D-dimer, and plasmin-antiplasmin with incident cardiovascular disease and all-cause mortality. Am J Hematol. 2009; 84(6):349–53. doi: <u>10.1002/Ajh.21429</u> PMID: <u>Medline:19472201</u>; PubMed Central PMCID: PMC2950108.
- Harris TB, Ferrucci L, Tracy RP, Corti MC, Wacholder S, Ettinger WH Jr., et al. Associations of elevated interleukin-6 and C-reactive protein levels with mortality in the elderly. Am J Med. 1999; 106(5):506–12. PMID: Medline:10335721.
- Rajasuriar R, Khoury G, Kamarulzaman A, French MA, Cameron PU, Lewin SR. Persistent immune activation in chronic HIV infection: do any interventions work? AIDS. 2013; 27(8):1199–208. doi: <u>10.</u> <u>1097/QAD.0b013e32835ecb8b</u> PMID: <u>WOS:000318156400002</u>.
- Butler SL, Valdez H, Westby M, Perros M, June CH, Jacobson JM, et al. Disease-modifying therapeutic concepts for HIV in the era of highly active antiretroviral therapy. J Acquir Immune Defic Syndr. 2011; 58(3):297–303. doi: 10.1097/QAI.0b013e31822ccfcc PMID: 21792065.
- El-Sadr WM, Lundgren JD, Neaton JD, Gordin F, Abrams D, Arduino RC, et al. CD4+ count-guided interruption of antiretroviral treatment. N Engl J Med. 2006; 355(22):2283–96. PMID: <u>17135583</u>.
- 33. EI-Sadr W, Grund B, Neuhaus J, Babiker A, Cohen C, Darbyshire J, et al. Risk for opportunistic disease and death after reinitiating continuous antiretroviral therapy in patients with HIV previously receiving episodic therapy—a randomized trial. Annals of Internal Medicine. 2008; 149(5):289–W62. PMID: ISI:000259229300001.
- Abrams D, Levy Y, Losso MH, Babilker A, Collins G, Cooper DA, et al. Interleukin-2 Therapy in Patients with HIV Infection. New England Journal of Medicine. 2009; 361(16):1548–59. PMID: <u>ISI:000270777000008</u>. doi: <u>10.1056/NEJMoa0903175</u>
- CIOMS Working Group. Introductiory Guide for Standardised MedDRA Queries (SMQs) Version 12.0. In: International Federation of Pharmaceutical Manufacturers and Associations (IFPMA), editor. 2009.
- Kowalska JD, Friis-Moller N, Kirk O, Bannister W, Mocroft A, Sabin C, et al. The Coding Causes of Death in HIV (CoDe) Project: initial results and evaluation of methodology. Epidemiology. 2011; 22 (4):516–23. PMID: Medline:21522013. doi: 10.1097/EDE.0b013e31821b5332
- Sakkinen PA, Macy EM, Callas PW, Cornell ES, Hayes TE, Kuller LH, et al. Analytical and biologic variability in measures of hemostasis, fibrinolysis, and inflammation: assessment and implications for epidemiology. Am J Epidemiol. 1999; 149(3):261–7. PMID: Medline:9927222.
- Burnham K, Anderson D. Multimodel inference—understanding AIC and BIC in model selection. Sociological Methods & Research. 2004; 33(2):261–304. doi: <u>10.1177/0049124104268644</u> PMID: WOS:000224706300004.
- Hughes MD. Regression dilution in the proportional hazards model. Biometrics. 1993; 49(4):1056–66. PMID: <u>8117900</u>.
- Knuiman MW, Divitini ML, Buzas JS, Fitzgerald PE. Adjustment for regression dilution in epidemiological regression analyses. Ann Epidemiol. 1998; 8(1):56–63. S1047279797001075 [pii]. PMID: 9465995.
- **41.** R Development Core Team. R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing; 2014. Available: <u>http://www.r-project.org</u>.
- Ridker PM. Testing the inflammatory hypothesis of atherothrombosis: scientific rationale for the cardiovascular inflammation reduction trial (CIRT). J Thromb Haemost. 2009; 7 Suppl 1:332–9. PMID: <u>Medline:19630828</u>. doi: <u>10.1111/j.1538-7836.2009.03404.x</u>
- 43. National Institute of Allergy and Infectious Diseases (NIAID). NIH Launches Largest Clinical Trial Focused on HIV-Related Cardiovascular Disease. NIH News 2015 [cited 2015 26 September]. Available: <u>http://www.niaid.nih.gov/news/newsreleases/2015/Pages/REPRIEVE.aspx</u>.
- Borges AH, Silverberg MJ, Wentworth D, Grulich AE, Fatkenheuer G, Mitsuyasu R, et al. Predicting risk of cancer during HIV infection: the role of inflammatory and coagulation biomarkers. AIDS. 2013; 27 (9):1433–41. PMID: Medline:23945504. doi: 10.1097/QAD.0b013e32835f6b0c
- Tripodi A, Mannucci PM. The coagulopathy of chronic liver disease. N Engl J Med. 2011; 365(2):147– 56. PMID: <u>Medline:21751907</u>. doi: <u>10.1056/NEJMra1011170</u>