

Remarks about the inviscid limit of the Navier-Stokes system.

Nader MASMOUDI*

Courant Institute of Mathematical Sciences

251 Mercer Street

New York, NY 10012-1185, USA

masmoudi@cims.nyu.edu

September 19, 2006

Abstract

In this paper we prove two results about the inviscid limit of the Navier-Stokes system. The first one concerns the convergence in H^s of a sequence of solutions to the Navier-Stokes system when the viscosity goes to zero and the initial data is in H^s . The second result deals with the best rate of convergence for vortex patch initial data in 2 and 3 dimensions. We present here a simple proof which also works in the 3D case. The 3D case is new.

1 The inviscid limit

The Navier-Stokes system is the basic mathematical model for viscous incompressible flows. In a bounded domain, it reads

$$\begin{cases} \partial_t u + u \cdot \nabla u - \nu \Delta u + \nabla p = 0, \\ \operatorname{div}(u) = 0, \\ u = 0 \quad \text{on} \quad \partial\Omega, \end{cases} \quad (1)$$

where u is the velocity, p is the pressure and ν is the kinematic viscosity. We can define a typical length scale L and a typical velocity U . The dimensionless

*Courant Institute, New York University 251 Mercer St, New York, NY 10012

parameter $Re = \frac{UL}{\nu}$ is very important to compare the properties of different flows. When Re is very large (ν very small), we expect that the Navier-Stokes system (NS_ν) behaves like the Euler system

$$\begin{cases} \partial_t u + u \cdot \nabla u + \nabla p = 0, \\ \operatorname{div}(u) = 0, \\ u \cdot n = 0 \quad \text{on} \quad \partial\Omega. \end{cases} \quad (2)$$

The zero-viscosity limit of the incompressible Navier-Stokes equation in a bounded domain, with Dirichlet boundary conditions, is one of the most challenging open problems in Fluid Mechanics (see [19] and the references therein). This is due to the formation of a boundary layer which appears because, we can not impose a Dirichlet boundary condition for the Euler equation. This boundary layer satisfies formally the Prandtl equations, which seem to be ill-posed in general.

In this paper we only deal with the inviscid limit in the whole space. All the results presented here can easily be extended to the periodic case.

2 Convergence in H^s .

The inviscid limit in the whole space case was performed by several authors, we can refer for instance to Swann [20] and Kato [16, 17] (see also Constantin [10]). Here, we would like to improve slightly the convergence stated in the previous works by proving the convergence in the H^s space as long as the solution of the Euler system exists. Indeed, in most of the previous results only convergence in $H^{s'}$ for $s' < s$ was proved. We also point out that in [17], Kato proved the convergence in H^s for a short time, by using a general theory about quasi-linear equations. So, we do not claim that theorem 2.1 is really new.

Take the Navier-Stokes system in the whole space \mathbb{R}^d

$$\partial_t u^n + \operatorname{div}(u^n \otimes u^n) - \nu_n \Delta u^n = -\nabla p^n \quad \text{in} \quad \mathbb{R}^d \quad (3)$$

$$\operatorname{div}(u^n) = 0 \quad \text{in} \quad \mathbb{R}^d \quad (4)$$

$$u^n(t=0) = u_0^n \quad \text{with} \quad \operatorname{div}(u_0^n) = 0 \quad (5)$$

where ν_n goes to 0 when n goes to infinity.

Theorem 2.1 *Let $s > d/2 + 1$, and $u_0^n \in H^s(\mathbb{R}^d)$ such that u_0^n goes to u_0 in $H^s(\mathbb{R}^d)$ when n goes to infinity. Let T^* be the time of existence and $u \in C_{loc}([0, T^*]; H^s)$ be the solution of the Euler system*

$$\partial_t u + \operatorname{div}(u \otimes u) = -\nabla p \quad \text{in } \mathbb{R}^d \quad (6)$$

$$\operatorname{div}(u) = 0 \quad \text{in } \mathbb{R}^d \quad (7)$$

$$u(t=0) = u_0 \quad \text{with } \operatorname{div}(u_0) = 0. \quad (8)$$

Then, for all $0 < T_0 < T^$, there exists $\nu_0 > 0$ such that for all $\nu_n \leq \nu_0$, the Navier-Stokes system (3 - 5) has a unique solution $u^n \in C([0, T_0]; H^s(\mathbb{R}^d))$. Moreover,*

$$\|u^n - u\|_{L^\infty(0, T_0; H^s)} \rightarrow 0, \quad n \rightarrow \infty \quad (9)$$

$$\|(u^n - u)(t)\|_{H^{s-2}} \leq C(\nu_n t + \|u_0^n - u_0\|_{H^{s-2}}) \quad (10)$$

$$\|(u^n - u)(t)\|_{H^{s'}} \leq C((\nu_n t)^{(s-s')/2} + \|u_0^n - u_0\|_{H^{s'}}) \quad (11)$$

for all $0 \leq t \leq T_0$, $s - 2 \leq s' \leq s - 1$ and C depends only on u and T_0 .

Remark 2.2 *1) The only relatively new part in theorem 2.1 is the convergence in H^s stated in (9) which holds for all $T_0 < T^*$.*

2) Interpolating between (10) and the uniform bound for w^n in $C([0, T]; H^s(\mathbb{R}^d))$, we deduce that u^n converges to u in $H^{s'}$ for any $s' < s$ and for $s - 2 < s' < s$, we have

$$\|(u^n - u)(t)\|_{H^{s'}} \leq C(\nu_n t + \|u_0^n - u_0\|_{H^{s-2}})^{\frac{s-s'}{2}}. \quad (12)$$

for all $0 \leq t \leq T_0$.

Proof:

The proof of this theorem is based on a standard Grönwall inequality (see also [20, 16, 10]). Let us start by proving (10). First, we see that we can solve the Navier-Stokes system and Euler system in $C([0, T]; H^s(\mathbb{R}^d))$ on some time interval independent of $\nu_n \leq \nu_0$ with bounds which are independent of n . This is because there is no boundary. Moreover, $w^n = u^n - u$ satisfies

$$\partial_t w^n + u^n \nabla w^n + w^n \cdot \nabla u - \nu_n \Delta w^n + \nu_n \Delta u + \nabla(p^n - p) = 0 \quad (13)$$

Then, we can write an energy estimate in H^{s-2} for $w^n = u^n - u$, namely

$$\begin{aligned} & \partial_t \|w^n\|_{H^{s-2}}^2 + \nu_n \|\nabla w^n\|_{H^{s-2}}^2 \\ & \leq \left(C(\|u\|_{H^s} + \|w^n\|_{H^s}) \|w^n\|_{H^{s-2}} + \nu_n \|\Delta u\|_{H^{s-2}} \right) \|w^n\|_{H^{s-2}} \end{aligned} \quad (14)$$

and by Grönwall lemma, we can deduce that (10) holds for $T_0 = T$. It was proved in [10] that the convergence holds as long as we can solve the Euler system and hence we can take any T_0 such that $T_0 < T^*$ (see [10]). Notice that in [10], the regularity required is $s - 2 > d/2 + 1$. However, this is not necessary modulo the regularization argument which is used to prove the convergence in H^s .

To prove (11), we write an energy estimate in $H^{s'}$, $s - 2 \leq s' \leq s - 1$

$$\begin{aligned} & \partial_t \|w^n\|_{H^{s'}}^2 + \nu_n \|\nabla w^n\|_{H^{s'}}^2 \\ & \leq \left(C(\|u\|_{H^s} + \|w^n\|_{H^s}) \|w^n\|_{H^{s'}}^2 + \nu_n \|\nabla u\|_{H^{s-1}} \|\nabla w^n\|_{H^{2s'-(s-1)}} \right). \end{aligned} \quad (15)$$

Then using an interpolation inequality and Holder inequality, we deduce that

$$\|\nabla w^n\|_{H^{2s'-s+1}} \leq C \|w^n\|_{H^{s'}}^{s-s'-1} \|\nabla w^n\|_{H^{s'}}^{2-(s-s')} \leq \frac{1}{C} \|\nabla w^n\|_{H^{s'}}^2 + C^2 \|w^n\|_{H^{s'}}^{2-\frac{2}{s-s'}}.$$

Hence,

$$\partial_t \|w^n\|_{H^{s'}}^2 \leq C \|w^n\|_{H^{s'}}^2 + C \nu_n \|w^n\|_{H^{s'}}^{2-\frac{2}{s-s'}} \quad (16)$$

and (11) follows.

Getting the convergence in H^s requires a regularization of the initial data. For all $\delta > 0$, we take u_0^δ such that $\|u_0^\delta\|_{H^s} \leq C \|u_0\|_{H^s}$, $\|u_0^\delta\|_{H^{s+1}} \leq \frac{C}{\delta}$, $\|u_0^\delta\|_{H^{s+2}} \leq \frac{C}{\delta^2}$ and for some s' such that $d/2 < s' < s - 1$, we have $\|u_0^\delta - u_0\|_{H^{s'}} \leq C \delta^{s-s'}$. Such a u_0^δ can be easily constructed by taking $u_0^\delta = \mathcal{F}^{-1}(1_{\{|\xi| \leq 1/\delta\}} \mathcal{F}u_0)$.

Let v^δ be the solution of the Euler system (6,7,8) with the initial data $v^\delta(t=0) = u_0^\delta$. It is easy to see that v^δ exists on some time interval $[0, T]$, $T < T^*$ which only depends on $\|u_0\|_{H^s}$ and such that for $0 \leq t \leq T$, we have $\|v^\delta(t)\|_{H^s} \leq C$ and $\|v^\delta(t)\|_{H^{s+2}} \leq \frac{C}{\delta^2}$ uniformly in δ . Indeed, the energy estimates at the level H^s and H^{s+2} read

$$\partial_t \|v^\delta\|_{H^s}^2 \leq C \|v^\delta\|_{H^s}^3 \quad (17)$$

$$\partial_t \|v^\delta\|_{H^{s+2}}^2 \leq C \|v^\delta\|_{H^s} \|v^\delta\|_{H^{s+2}}^2 \quad (18)$$

from which the uniform estimates follow.

Then, setting $w^\delta = v^\delta - u$, we have

$$\partial_t w^\delta + w^\delta \cdot \nabla v^\delta + u \cdot \nabla w^\delta = -\nabla(p^\delta - p). \quad (19)$$

Taking the energy estimate in H^s yields

$$\partial_t \|w^\delta\|_{H^s}^2 \leq C(\|u\|_{H^s} + \|v^\delta\|_{H^s}) \|w^\delta\|_{H^s}^2 + C\|v^\delta\|_{H^{s+1}} \|w^\delta\|_{H^s} \|w^\delta\|_{L^\infty}. \quad (20)$$

Then, we notice that on the time interval $[0, T]$, we have $\|v^\delta\|_{H^{s+1}} \leq \frac{C}{\delta}$. Moreover, taking the energy estimate at the regularity s' , we get

$$\partial_t \|w^\delta\|_{H^{s'}}^2 \leq C(\|u\|_{H^{s'}} + \|v^\delta\|_{H^{s'+1}}) \|w^\delta\|_{H^{s'}}^2 \quad (21)$$

and since $s' + 1 < s$, we get easily that $\|w^\delta\|_{L^\infty(0, T; H^{s'})} \leq C\delta^{s-s'}$ and by Sobolev embedding, we have $\|w^\delta\|_{L^\infty(0, T; L^\infty)} \leq C\|w^\delta\|_{L^\infty(0, T; H^{s'})} \leq C\delta^{s-s'}$. Hence, (20) gives

$$\partial_t \|w^\delta\|_{H^s} \leq C(\|u\|_{H^s} + \|v^\delta\|_{H^s}) \|w^\delta\|_{H^s} + C\delta^{s-s'-1}. \quad (22)$$

Hence w^δ goes to zero in $L^\infty(0, T; H^s)$, namely v^δ goes to v in $L^\infty(0, T; H^s)$ when δ goes to zero and we have

$$\|v^\delta - u\|_{L^\infty(0, T; H^s)} \leq C(\|u_0^\delta - u_0\|_{H^s} + \delta^{s-s'-1}T) \quad (23)$$

Writing an energy estimate for $w^{n, \delta} = u^n - v^\delta$, we get (here we drop the n and δ)

$$\begin{aligned} & \partial_t \|w\|_{H^s}^2 + \nu_n \|\nabla w\|_{H^s}^2 \\ & \leq C(\|w\|_{L^\infty} \|v^\delta\|_{H^{s+1}} \|w\|_{H^s} + (\|v^\delta\|_{H^s} + \|u^n\|_{H^s}) \|w\|_{H^s}^2) + \\ & \quad \nu_n \|v^\delta\|_{H^{s+2}} \|w\|_{H^s}. \end{aligned} \quad (24)$$

Hence, we get

$$\begin{aligned} \partial_t \|w\|_{H^s} & \leq C\|u^n - u\|_{L^\infty} \|v^\delta\|_{H^{s+1}} + C\|v^\delta - u\|_{L^\infty} \|v^\delta\|_{H^{s+1}} + \\ & \quad \nu_n \|v^\delta\|_{H^{s+2}} + C(\|v^\delta\|_{H^s} + \|u^n\|_{H^s}) \|w\|_{H^s}. \end{aligned} \quad (25)$$

Since u^n converges to u in H^{s-1} , we deduce that

$$\|u^n - u\|_{L^\infty(0, T; L^\infty)} \leq C\|u^n - u\|_{L^\infty(H^{s-1})} \leq C((\nu_n T)^{1/2} + \|u_0^n - u_0\|_{H^{s-1}}). \quad (26)$$

Taking $\delta = \delta_n$ such that δ_n , $\frac{\|u_0^n - u_0\|_{H^{s-1}}}{\delta_n}$ and $\frac{\nu_n}{\delta_n^2}$ go to zero when n goes to infinity, we deduce that

$$\partial_t \|w^{n, \delta}\|_{H^s} \leq C\left(\frac{(\nu T)^{1/2} + \|u_0^n - u_0\|_{H^{s-1}}}{\delta} + \delta^{s-s'-1} + \frac{\nu}{\delta^2} + \|w^{n, \delta}\|_{H^s}\right) \quad (27)$$

Hence, by Grönwall lemma, we deduce that $w^{n,\delta}$ goes to zero in $L^\infty(0, T; H^s)$ and that u^n goes to u in $L^\infty(0, T; H^s)$. Moreover,

$$\|u^n - u\|_{L^\infty(0, T; H^s)} \leq CT \left(\frac{(\nu T)^{1/2} + \|u_0^n - u_0\|_{H^{s-1}}}{\delta} + \delta^{s-s'-1} + \frac{\nu}{\delta^2} \right) + C(\|u_0^n - u_0\|_{H^s} + \|u_0^\delta - u_0\|_{H^s} + \delta^{s-s'-1}T). \quad (28)$$

We notice here that the rate of convergence gets better if we have a better approximation of u_0 by u_0^δ . This will be studied in the next subsection.

Since, we have proved the convergence in H^s till the time T , we can iterate the previous argument. Indeed, taking T as a new initial time and noticing that $u^n(T)$ goes to $u(T)$ in H^s , we see that we can iterate the previous argument on some time interval $[T, T+T_1]$ where $T_1 = T_1(\|u(T)\|_{H^s})$ only depends on $\|u(T)\|_{H^s}$ and $T_1 \geq C/\|u(T)\|_{H^s}$. Then, we can construct a sequence of times T_k , $k \geq 1$ by this procedure. Now, it is clear that $T + T_1 + \dots + T_k$ goes to T^* when k goes to infinity. Indeed, the time T_{k+1} goes to zero only if $\|u(T + T_1 + \dots + T_k)\|_{H^s}$ goes to infinity, which means that $T + T_1 + \dots + T_k$ goes to T^* . This iteration argument allows us to get the convergence on any time interval $[0, T_0]$, $T_0 < T^*$. \square

Remark 2.3 1) We notice that the time T^* is related to the existence time for the Euler system (6). If $d = 2$ it is known [22, 21] that the Euler system (6) has a global solution and hence one can take any time $T_0 < \infty$ in the above theorem.

2) The idea of using a regularization of the initial data was also used by Beirão da Veiga [2, 3] to prove a similar result in the compressible-incompressible limit. It is also used to prove the continuity of the solution with respect to the initial data in hyperbolic equations (see for instance Bona and Smith [5]). In the inviscid limit, this idea was used by Constantin and Wu [12] to prove some estimates on the rate of convergence of the vorticity.

2.1 Rate of convergence in H^s

Take β such that $1 < \beta \leq 2$ and $d/2 < s - \beta$ and for $0 \leq \delta < \infty$, $T > 0$ we define $u_0^\delta = \mathcal{F}^{-1}(1_{\{|\xi| \leq 1/\delta\}} \mathcal{F}u_0)$, $\varepsilon_T(\delta) = \|u_0^\delta - u_0\|_{H^s} + T\delta^{\beta-1}$, $f_T(\delta) = \delta\varepsilon_T(\delta)$ and $g_T(\delta) = \delta^2\varepsilon_T(\delta)$. We can see easily that for $T > 0$, f_T and g_T are increasing on $[0, \infty)$. We denote by f_T^{-1} and g_T^{-1} their inverse. From the proof of theorem 2.1, we can deduce the following corollary

Corollary 2.4 *Under the same hypotheses of theorem 2.1, we have the following rate of convergence*

$$\|(u^n - u)(t)\|_{H^s} \leq C \frac{\nu t}{(g_t^{-1}(\nu t))^2} + C \frac{t((\nu t)^{\beta/2} + \|u_0^n - u_0\|_{H^{s-\beta}})}{f_t^{-1}(t((\nu t)^{\beta/2} + \|u_0^n - u_0\|_{H^{s-\beta}}))} + C \|u_0^n - u_0\|_{H^s} \quad (29)$$

for all $0 \leq t \leq T_0$ and C depends only on u and T_0 .

Proof:

Going back to the proof of theorem 2.1, we see that (26) can be replaced by

$$\|(u^n - u)(t)\|_{L^\infty} \leq C \|u^n - u(t)\|_{H^{s-\beta}} \leq C((\nu_n t)^{\beta/2} + \|u_0^n - u_0\|_{H^{s-\beta}}). \quad (30)$$

Hence (28) can be replaced by

$$\|(u^n - u)(t)\|_{H^s} \leq Ct \left(\frac{(\nu t)^{\beta/2} + \|u_0^n - u_0\|_{H^{s-\beta}}}{\delta} + \frac{\nu}{\delta^2} \right) + C(\varepsilon_t(\delta) + \|u_0^n - u_0\|_{H^s}). \quad (31)$$

Taking the optimum in δ and applying lemma 2.5, we deduce easily that (29) holds. \square

Lemma 2.5 *For $a, b, t > 0$, we have*

$$\inf_{\delta > 0} \frac{a}{\delta} + \frac{b}{\delta^2} + \varepsilon_t(\delta) \leq 2 \frac{a}{f_t^{-1}(a)} + \frac{b}{(g_t^{-1}(b))^2} \quad (32)$$

The proof of this lemma is simple and is left for the reader

If we assume that u_0 is more regular, we can give a more precise rate.

Corollary 2.6 *We take the same hypotheses as in theorem 2.1 and assume in addition that $u_0 \in H^{s+\alpha}$ for some $0 < \alpha \leq 2$.*

If $1 \leq \alpha \leq 2$, we have

$$\|(u^n - u)(t)\|_{H^s} \leq C((\nu t)^{\alpha/2} + \|u_0^n - u_0\|_{H^s}) \quad (33)$$

for all $0 \leq t \leq T_0$ and C depends only on u and T_0 .

If $0 < \alpha < 1$, then for all β such that $1 \leq \alpha + \beta \leq 2$ and $s - \beta > d/2$, we have

$$\|(u^n - u)(t)\|_{H^s} \leq C(t \|u_0^n - u_0\|_{H^{s-\beta}})^\alpha + C((\nu t)^{\alpha/2} + \|u_0^n - u_0\|_{H^s}) \quad (34)$$

for all $0 \leq t \leq T_0$ and C depends only on u , β and T_0 .

Proof:

First, notice that from the extra regularity of u_0 , we deduce that $\|v^\delta\|_{H^{s+1}} \leq C(1 + \delta^{\alpha-1})$, $\|v^\delta\|_{H^{s+2}} \leq C\delta^{\alpha-2}$ and $\|v^\delta - u\|_{H^s} \leq C\delta^\alpha$.

If $1 \leq \alpha \leq 2$, then (24) yields

$$\partial_t \|w\|_{H^s} \leq C\|w\|_{H^s} + C\nu_n \delta^{\alpha-2}. \quad (35)$$

Hence

$$\|(u^n - u)(t)\|_{H^s} \leq C\left(\nu t \delta^{\alpha-2} + \delta^\alpha + \|u_0^n - u_0\|_{H^s}\right) \quad (36)$$

Taking the optimum in δ , namely $\delta = \sqrt{\nu t}$, we deduce that (33) holds.

If $0 < \alpha < 1$, then arguing as in (15), we have

$$\begin{aligned} & \partial_t \|w^n\|_{H^{s-\beta}}^2 + \nu_n \|\nabla w^n\|_{H^{s-\beta}}^2 \\ & \leq \left(C(\|u\|_{H^{s+\alpha}} + \|w^n\|_{H^s}) \|w^n\|_{H^{s-\beta}}^2 + \nu_n \|\nabla u\|_{H^{s+\alpha-1}} \|\nabla w^n\|_{H^{s+1-2\beta-\alpha}} \right). \end{aligned} \quad (37)$$

Then using an interpolation inequality and Holder inequality, we deduce that

$$\|\nabla w^n\|_{H^{s+1-2\beta-\alpha}} \leq C \|w^n\|_{H^{s-\beta}}^{\beta+\alpha-1} \|\nabla w^n\|_{H^{s-\beta}}^{2-\beta-\alpha} \leq \frac{1}{C} \|\nabla w^n\|_{H^{s-\beta}}^2 + C^2 \|w^n\|_{H^{s-\beta}}^{2-\frac{2}{\beta+\alpha}}.$$

Hence, we deduce that

$$\|(u^n - u)(t)\|_{H^{s-\beta}} \leq C((\nu_n t)^{(\beta+\alpha)/2} + \|u_0^n - u_0\|_{H^{s-\beta}}). \quad (38)$$

In the proof of theorem 2.1, we see that (26) can be replaced by

$$\|(u^n - u)(t)\|_{L^\infty} \leq C\|u^n - u(t)\|_{H^{s-\beta}} \leq C((\nu_n t)^{(\beta+\alpha)/2} + \|u_0^n - u_0\|_{H^{s-\beta}}). \quad (39)$$

Moreover, $\|v^\alpha - u\|_{L^\infty} \leq C\delta^{\beta+\alpha}$. Hence (28) can be replaced by

$$\begin{aligned} \|(u^n - u)(t)\|_{H^s} & \leq Ct \left(((\nu t)^{(\beta+\alpha)/2} + \|u_0^n - u_0\|_{H^{s-\beta}}) \delta^{\alpha-1} + \nu \delta^{\alpha-2} \right) + \\ & C(\delta^\alpha + t\delta^{\beta+2\alpha-1} + \|u_0^n - u_0\|_{H^s}). \end{aligned} \quad (40)$$

Taking the optimum in δ , we deduce that

$$\|(u^n - u)(t)\|_{H^s} \leq C \left(t((\nu t)^{(\beta+\alpha)/2} + \|u_0^n - u_0\|_{H^{s-\beta}}) \right)^\alpha + C(\nu t)^{\alpha/2} + C\|u_0^n - u_0\|_{H^s}. \quad (41)$$

Hence (34) holds. \square

3 Vortex patches

In this section $d = 2$ or 3 . For $2D$ vortex patches, namely the case where $\text{curl}(u_0)$ is the characteristic function of a $C^{1+\alpha}$ domain $\alpha > 0$, it was proved in [7] (see also [4]) that the Euler system (6,7,8) has a unique solution u such that the characteristic function of $\text{curl}(u)$ remains a $C^{1+\alpha}$ domain and that the velocity u is in $L_{loc}^\infty(\mathbb{R}; Lip)$. A similar result holds for $3D$ vortex patches, but only on a bounded interval, namely $u \in L_{loc}^\infty(0, T^*; Lip)$ (see [14]).

For vortex patches, theorem 2.1 does not apply. Indeed, the velocity is not in H^s for any $s > d/2 + 1$. For $2D$ vortex patches, it was proved in [11, 12] that the convergence to the Euler system still holds and that

$$\|u^n - u\|_{L^\infty(0,T;L^2)} \leq C(\nu_n T)^{\frac{1}{2}}. \quad (42)$$

In [12], the authors also prove some estimate in L^p spaces for the difference between the vorticities, in particular they prove for $p \geq 2$ that $\|\text{curl}(u^n - u)\|_{L^\infty(0,T;L^p)} \leq C\nu_n^{\frac{1}{4p}-\varepsilon}$ for some short time T and $\varepsilon > 0$.

Also, in [1], a better rate of convergence is given for $2D$ vortex patches, namely

$$\|u^n - u\|_{L^\infty(0,T;L^2)} \leq C(\nu_n T)^{\frac{3}{4}} \quad (43)$$

which is optimal.

Here, we would like to extend the result of Abid and Danchin [1] to the $3D$ case and also give a slight improvement of their $2D$ result by allowing $u_0^n - u_0$ to be just in L^2 . Moreover, the proof we present is much simpler.

Let us recall the definition of a vortex patch

Definition 3.1 *Take $0 < r < 1$. A vector field u is called a C^r vortex patch if the following decomposition holds*

$$\text{curl}(u) = \chi_P \omega_i + \chi_{P^c} \omega_e \quad (44)$$

where $P \subset \mathbb{R}^d$ is an open set of class C^{1+r} and $\omega_i, \omega_e \in C^r(\mathbb{R}^d) \cap L^1(\mathbb{R}^d)$ are compactly supported.

Here χ_P denotes the characteristic function of P . Notice that since $\text{curl}(u)$ is divergence-free, we have $\omega_i \cdot n = \omega_e \cdot n$ on ∂P . This condition is always satisfied if $d = 2$.

First, we recall that Gamblin and Saint-Raymond [14] proved the existence of a local solution $u \in L_{loc}^\infty(0, T^*; Lip)$ to the vortex patch problem in $3D$ (see also [13] and [15]). Moreover, u remains a C^r vortex patch.

Hence, $\text{curl}(u) \in L_{loc}^\infty(0, T^*; \dot{B}_{2,\infty}^\alpha)$ where $\alpha = \min(r, 1/2)$ (see the appendix).

Theorem 3.2 *Here $d = 2$ or 3 . We assume that $u_0^n - u_0$ goes to 0 in $L^2(\mathbb{R}^d)$ when n goes to infinity. We also assume that u_0 is a C^r vortex patch.*

Then, if T^ is the time of existence and $u \in C_{loc}([0, T^*]; Lip)$ is the solution of the Euler system with initial data u_0 , then for all $0 < T < T^*$, there exists ν_0 such that for all $\nu_n \leq \nu_0$ and for all sequence of weak (Leray) solutions to the Navier-Stokes system (3 - 5), we have for $0 < t < T$,*

$$\|(u^n - u)(t)\|_{L^2} \leq C((t\nu_n)^{\frac{1+\alpha}{2}} + \|u_0^n - u_0\|_{L^2}) \quad (45)$$

where $\alpha = \min(1/2, r)$ and where C depends only on u and T .

Remark 3.3 *In the 2D case, knowing that $\text{curl}(u_0) \in L^1 \cap L^\infty$ does not imply that $u \in L^2$ and in general u_0 is not in L^2 unless $\int \text{curl} u = 0$. In particular in the classical 2D vortex patch problem [18], namely the case $\text{curl}(u)$ is the characteristic function of a C^{r+1} domain, u_0 is not in L^2 . However, in the 3D case, the fact that $\text{curl} u_0 \in L^1 \cap L^\infty$ implies that $u_0 \in L^2$ from Biot-Savart formula.*

Proof: Let us denote $w^n = u^n - u$, hence

$$\partial_t w^n + w^n \cdot \nabla u + u \cdot \nabla w^n - \nu_n \Delta w^n - \nu_n \Delta u = -\nabla(p^n - p). \quad (46)$$

Taking the L^2 product with w^n , we get (at least formally) for $0 < t < T$,

$$\begin{aligned} \frac{1}{2} \|w^n(t)\|_{L^2}^2 + \nu_n \int_0^t \|\nabla w^n\|_{L^2}^2 &\leq \frac{1}{2} \|w^n(0)\|_{L^2}^2 + \\ &\int_0^t C \|\nabla u\|_{L^\infty} \|w^n\|_{L^2}^2 - \int_0^t \int \nu_n \nabla u \cdot \nabla w^n. \end{aligned} \quad (47)$$

In the 2D case, this computation is fully justified. We only point that in the 2D case, u and u^n are not in general in L^2 but their difference is in L^2 .

To prove (47) rigorously in the 3D case, we just add the energy inequality (48) and energy equality (49)

$$\frac{1}{2} \|u^n(t)\|_{L^2}^2 + \nu_n \int_0^t \|\nabla u^n\|_{L^2}^2 \leq \frac{1}{2} \|u^n(0)\|_{L^2}^2 \quad (48)$$

$$\frac{1}{2}\|u(t)\|_{L^2}^2 = \frac{1}{2}\|u(0)\|_{L^2}^2 \quad (49)$$

and subtract the weak formulation

$$\int u^n u(t) - \int u^n u(0) + \int_0^t \int u^n u \cdot \nabla u + u u^n \cdot \nabla u^n + \nu_n \nabla u^n \cdot \nabla u = 0. \quad (50)$$

Besides, using the duality between $\dot{B}_{2,1}^{-\alpha}$ and $\dot{B}_{2,\infty}^\alpha$, the divergence-free property of u and lemma 5.1 (see the appendix), we have

$$|\int \nabla u \cdot \nabla w^n| \leq C \|\nabla u\|_{\dot{B}_{2,\infty}^\alpha} \|\nabla w^n\|_{\dot{B}_{2,1}^{-\alpha}} \quad (51)$$

$$\leq C \|\operatorname{curl} u\|_{\dot{B}_{2,\infty}^\alpha} \|w^n\|_{\dot{B}_{2,1}^{1-\alpha}} \quad (52)$$

$$\leq C \|\operatorname{curl} u\|_{\dot{B}_{2,\infty}^\alpha} \|w^n\|_{L^2}^\alpha \|\nabla w^n\|_{L^2}^{1-\alpha}. \quad (53)$$

By Holder inequality, we have

$$\nu_n \|\operatorname{curl} u\|_{\dot{B}_{2,\infty}^\alpha} \|w^n\|_{L^2}^\alpha \|\nabla w^n\|_{L^2}^{1-\alpha} \leq C \nu_n \|\operatorname{curl} u\|_{\dot{B}_{2,\infty}^\alpha}^{\frac{2}{1+\alpha}} \|w^n\|_{L^2}^{\frac{2\alpha}{1+\alpha}} + \frac{\nu_n}{2} \|\nabla w^n\|_{L^2}^2. \quad (54)$$

Hence, we get from (47)

$$\|w^n(t)\|_{L^2}^2 \leq \|w^n(0)\|_{L^2}^2 + C \int_0^t \|\nabla u\|_{L^\infty} \|w^n\|_{L^2}^2 + C \nu_n \|\operatorname{curl} u\|_{\dot{B}_{2,\infty}^\alpha}^{\frac{2}{1+\alpha}} \|w^n\|_{L^2}^{\frac{2\alpha}{1+\alpha}} ds. \quad (55)$$

And by Grönwall lemma, we deduce that

$$\|w^n(t)\|_{L^2}^{\frac{2}{1+\alpha}} \leq C \|w^n(0)\|_{L^2}^{\frac{2}{1+\alpha}} + C \nu_n t \quad (56)$$

and (45) follows. \square

From the proof, we can see that the only information we used about u is that $u \in L_{loc}^\infty(0, T^*; Lip)$ and $\operatorname{curl} u \in L_{loc}^\infty(0, T^*; \dot{B}_{2,\infty}^\alpha)$. Moreover, it is easy to see that if $u \in L_{loc}^\infty([0, T^*]; Lip)$ then the $\dot{B}_{2,\infty}^\alpha$, $0 < \alpha < 1$ regularity of $\operatorname{curl} u$ is propagated by the flow, namely if $\operatorname{curl} u^0 \in \dot{B}_{2,\infty}^\alpha$, then $\operatorname{curl} u \in L_{loc}^\infty([0, T^*]; \dot{B}_{2,\infty}^\alpha)$ (see [8]). Hence, we have the following theorem

Theorem 3.4 *We assume that $u_0^n - u_0$ goes to 0 in $L^2(\mathbb{R}^d)$ when n goes to infinity and that $\operatorname{curl}u_0 \in \dot{B}_{2,\infty}^\alpha$, $0 < \alpha < 1$. We also assume that the Euler system with the initial data u_0 has a unique solution $u \in L_{loc}^\infty([0, T^*]; Lip)$. Then for all $0 < T < T^*$, there exists ν_0 such that for all $\nu_n \leq \nu_0$ and for all sequence of weak (Leray) solutions to the Navier-Stokes system (3 - 5), we have for $0 < t < T$,*

$$\|(u^n - u)(t)\|_{L^2} \leq C((t\nu_n)^{\frac{1+\alpha}{2}} + \|u_0^n - u_0\|_{L^2}) \quad (57)$$

where C depends only on u and T .

This theorem is an improvement of theorem 1.1 of [1] since we only assume that the solution of the Euler system is Lipschitz. We would like to give two applications of this theorem which yield a better convergence rate than the simple application of theorem 3.2.

Consider a vector field u_0 which is a C^r vortex patch with $0 < r < 1/2$ and assume in addition that $\operatorname{curl}u_0 \in \dot{B}_{2,\infty}^{1/2}$, then theorem 3.4 allows us to prove that

$$\|(u^n - u)(t)\|_{L^2} \leq C((t\nu_n)^{\frac{3}{4}} + \|u_0^n - u_0\|_{L^2}) \quad (58)$$

which is better than the rate we get from (45).

There are several situations where u_0 is a C^r vortex patch with $0 < r < 1/2$ and $\operatorname{curl}u_0 \in \dot{B}_{2,\infty}^{1/2}$. For instance this is the case if $\operatorname{curl}(u_0) = \chi_P \omega_{i0} + \chi_{P^c} \omega_{e0}$ is such that $P \subset \mathbb{R}^d$ is an open set of class C^{1+r} and $\omega_{i0}, \omega_{e0} \in C^{1/2}(\mathbb{R}^d) \cap L^1(\mathbb{R}^d)$. We notice here that for $t > 0$ we only know that u is a C^r vortex patch, namely $\operatorname{curl}(u) = \chi_{P(t)} \omega_i(t) + \chi_{P(t)^c} \omega_e(t)$ with $P(t)$ of class C^{1+r} and $\omega_i(t), \omega_e(t) \in L_{loc}^\infty(0, T^*; C^r(\mathbb{R}^d) \cap L^1(\mathbb{R}^d))$. Hence, $\operatorname{curl}u \in L_{loc}^\infty([0, T^*]; \dot{B}_{2,\infty}^r)$. However, propagating the initial $\dot{B}_{2,\infty}^{1/2}$ yields that $\operatorname{curl}u \in L_{loc}^\infty([0, T^*]; \dot{B}_{2,\infty}^{1/2})$ and gives the better rate (58).

In particular theorem 3.4 applies to the classical 2D vortex patch, namely the case $\operatorname{curl}u_0 = \chi_P$ and P is of class C^{1+r} , $r > 0$ in which case (58) holds even if $r < 1/2$.

Remark 3.5 *In the 2D case, one can lower the regularity of the initial data. Indeed Yudovich [22] proved that if $\omega_0 = \operatorname{curl}(u_0) \in L^\infty \cap L^p$ for some $1 < p < \infty$ then the Euler system (6) has a unique global solution (see also [8]). It was proved in [9] that the solution to the Navier-Stokes system converges*

in $L^\infty((0, T); L^2)$ to the solution of the Euler system if we only assume that $\omega_0 = \text{curl}(u_0) \in L^\infty \cap L^p$. More precisely, Chemin [9] proves that

$$\|u^n - u\|_{L^\infty(0, T; L^2)} \leq C \|\text{curl}(u_0)\|_{L^\infty \cap L^2} (\nu_n T)^{\frac{1}{2} \exp(-C \|\text{curl}(u_0)\|_{L^\infty \cap L^2} T)}. \quad (59)$$

Notice that here, the rate of convergence deteriorates with time. This does not happen if we also know that u is in $L^\infty(0, T; Lip)$ as was proved by Constantin and Wu [11].

4 Acknowledgments

The author would like to thank Raphael Danchin for many remarks about an earlier version of the paper. He also would like to thank the referee for suggesting the study of the rate of convergence in subsection 2.1.

5 Appendix

We define \mathcal{C} to be the ring of center 0, of small radius 1/2 and great radius 2. There exist two nonnegative radial functions χ and φ belonging respectively to $\mathcal{D}(B(0, 1))$ and to $\mathcal{D}(\mathcal{C})$ so that

$$\chi(\xi) + \sum_{q \geq 0} \varphi(2^{-q}\xi) = 1, \quad (60)$$

$$|p - q| \geq 2 \Rightarrow \text{Supp } \varphi(2^{-q}\cdot) \cap \text{Supp } \varphi(2^{-p}\cdot) = \emptyset. \quad (61)$$

For instance, one can take $\chi \in \mathcal{D}(B(0, 1))$ such that $\chi \equiv 1$ on $B(0, 1/2)$ and take

$$\varphi(\xi) = \chi(2\xi) - \chi(\xi).$$

Then, we are able to define the Littlewood-Paley decomposition. Let us denote by \mathcal{F} the Fourier transform on \mathbb{R}^d . Let $h, \tilde{h}, \Delta_q, S_q$ ($q \in \mathbb{Z}$) be defined as follows:

$$\begin{aligned} h &= \mathcal{F}^{-1}\varphi \quad \text{and} \quad \tilde{h} = \mathcal{F}^{-1}\chi, \\ \Delta_q u &= \mathcal{F}^{-1}(\varphi(2^{-q}\xi)\mathcal{F}u) = 2^{qd} \int h(2^q y) u(x - y) dy, \\ S_q u &= \mathcal{F}^{-1}(\chi(2^{-q}\xi)\mathcal{F}u) = 2^{qd} \int \tilde{h}(2^q y) u(x - y) dy. \end{aligned}$$

Then, we define the non-homogeneous and homogeneous Besov norms

$$\|u\|_{B_{2,r}^s} = \left(\|S_0 u\|_{L^2}^r + \sum_{q \geq 0} 2^{rsq} \|\Delta_q u\|_{L^2}^r \right)^{1/r}$$

$$\|u\|_{\dot{B}_{2,r}^s} = \left(\sum_{q \in \mathbb{Z}} 2^{rsq} \|\Delta_q u\|_{L^2}^r \right)^{1/r}$$

for $s \in \mathbb{R}$ and $1 \leq r \leq \infty$. If $r = \infty$, then the summation over q is replaced by the L^∞ norm.

Lemma 5.1 *For $0 < \alpha < 1$, we have*

$$\|w\|_{\dot{B}_{2,1}^{1-\alpha}} \leq C \|w\|_{L^2}^\alpha \|\nabla w\|_{L^2}^{1-\alpha}. \quad (62)$$

Proof:

This inequality can be easily deduced from real interpolation. We give here a direct proof. Actually, we will prove a stronger estimate, namely

$$\|w\|_{\dot{B}_{2,1}^{1-\alpha}} \leq C \|w\|_{\dot{B}_{2,\infty}^0}^\alpha \|\nabla w\|_{\dot{B}_{2,\infty}^0}^{1-\alpha}. \quad (63)$$

Indeed, we have

$$\|\Delta_q w\|_{L^2} \leq C \|w\|_{\dot{B}_{2,\infty}^0} \quad (64)$$

$$\|\Delta_q w\|_{L^2} \leq C 2^{-q} \|w\|_{\dot{B}_{2,\infty}^1}. \quad (65)$$

We take N such that

$$2^N \|w\|_{\dot{B}_{2,\infty}^0} \leq \|w\|_{\dot{B}_{2,\infty}^1} \leq 2^{N+1} \|w\|_{\dot{B}_{2,\infty}^0}.$$

Hence

$$\begin{aligned} \sum_{q=-\infty}^{\infty} 2^{(1-\alpha)q} \|\Delta_q w\|_{L^2} &\leq C \sum_{q \leq N} 2^{(1-\alpha)q} \|w\|_{\dot{B}_{2,\infty}^0} + C \sum_{q \geq N} 2^{-\alpha q} \|w\|_{\dot{B}_{2,\infty}^1} \\ &\leq C 2^{(1-\alpha)N} \|w\|_{\dot{B}_{2,\infty}^0} + C 2^{-\alpha N} \|w\|_{\dot{B}_{2,\infty}^1}. \end{aligned} \quad (66)$$

From which (63) follows. \square

In the next two lemmas, we prove that if u_0 is a C^r vortex patch then $\text{curl}(u) \in L_{loc}^\infty(0, T^*; B_{2,\infty}^\alpha)$ where $\alpha = \min(r, 1/2)$.

Lemma 5.2 *If P is bounded open set of \mathbb{R}^d of class C^{1+r} then $\chi_P \in \dot{B}_{2,\infty}^{1/2}$*

The proof is based on interpolation. Indeed, since P is C^{1+r} , it is Lipschitz and hence $\chi_P \in L^\infty \cap BV$. Then

$$\|\Delta_q \chi_P\|_{L^\infty} \leq C \|\chi_P\|_{L^\infty} \quad (67)$$

$$\|\Delta_q \chi_P\|_{L^1} \leq C 2^{-q} \|\chi_P\|_{BV}. \quad (68)$$

interpolating between L^1 and L^∞ , we deduce that

$$\|\Delta_q \chi_P\|_{L^2} \leq C 2^{-q/2} \|\chi_P\|_{BV} \|\chi_P\|_{L^\infty}. \quad (69)$$

and hence, $\chi_P \in \dot{B}_{2,\infty}^{1/2}$. \square

Lemma 5.3 *If u is a C^r vortex patch then $\text{curl}(u) \in \dot{B}_{2,\infty}^\alpha$ where $\alpha = \min(r, 1/2)$.*

The proof uses the para-product decomposition of Bony ([6])

$$uv = T_u v + T_v u + R(u, v)$$

where

$$T_u v = \sum_{q \in \mathbb{Z}} S_{q-1} u \Delta_q v \quad \text{and} \quad R(u, v) = \sum_{|q-q'| \leq 1} \Delta_{q'} u \Delta_q v.$$

We decompose

$$\chi_P \omega_i = T_{\omega_i} \chi_P + R(\chi_P, \omega_i) + T_{\chi_P} \omega_i \quad (70)$$

and notice that since χ_P and ω_i are both in L^∞ , we get

$$\|T_{\omega_i} \chi_P + R(\chi_P, \omega_i)\|_{\dot{B}_{2,\infty}^\alpha} \leq C \|\omega_i\|_{L^\infty} \|\chi_P\|_{\dot{B}_{2,\infty}^\alpha} \quad (71)$$

$$\|T_{\chi_P} \omega_i\|_{\dot{B}_{2,\infty}^\alpha} \leq C \|\chi_P\|_{L^\infty} \|\omega_i\|_{\dot{B}_{2,\infty}^\alpha} \quad (72)$$

Since, ω_i is in C^r and is compactly supported, we deduce that $\omega_i \in \dot{B}_{2,\infty}^\alpha$. Hence $\chi_P \omega_i \in \dot{B}_{2,\infty}^\alpha$

The same proof holds for $\chi_{P^c} \omega_e$ and hence, $\text{curl}(u) \in \dot{B}_{2,\infty}^\alpha$. \square

References

- [1] H. Abidi and R. Danchin. Optimal bounds for the inviscid limit of Navier-Stokes equations. *Asymptot. Anal.*, 38(1):35–46, 2004.
- [2] H. Beirão da Veiga. On the singular limit for slightly compressible fluids. *Calc. Var. Partial Differential Equations*, 2(2):205–218, 1994.
- [3] H. Beirão da Veiga. Singular limits in compressible fluid dynamics. *Arch. Rational Mech. Anal.*, 128(4):313–327, 1994.
- [4] A. L. Bertozzi and P. Constantin. Global regularity for vortex patches. *Comm. Math. Phys.*, 152(1):19–28, 1993.
- [5] J. L. Bona and R. Smith. The initial-value problem for the Korteweg-de Vries equation. *Philos. Trans. Roy. Soc. London Ser. A*, 278(1287):555–601, 1975.
- [6] J.-M. Bony. Calcul symbolique et propagation des singularités pour les équations aux dérivées partielles non linéaires. *Ann. Sci. École Norm. Sup. (4)*, 14(2):209–246, 1981.
- [7] J.-Y. Chemin. Persistence de structures géométriques dans les fluides incompressibles bidimensionnels. *Ann. Sci. École Norm. Sup. (4)*, 26(4):517–542, 1993.
- [8] J.-Y. Chemin. Fluides parfaits incompressibles. *Astérisque*, (230):177, 1995.
- [9] J.-Y. Chemin. A remark on the inviscid limit for two-dimensional incompressible fluids. *Comm. Partial Differential Equations*, 21(11-12):1771–1779, 1996.
- [10] P. Constantin. Note on loss of regularity for solutions of the 3-D incompressible Euler and related equations. *Comm. Math. Phys.*, 104(2):311–326, 1986.
- [11] P. Constantin and J. Wu. Inviscid limit for vortex patches. *Nonlinearity*, 8(5):735–742, 1995.
- [12] P. Constantin and J. Wu. The inviscid limit for non-smooth vorticity. *Indiana Univ. Math. J.*, 45(1):67–81, 1996.

- [13] A. Dutrifoy. On 3-D vortex patches in bounded domains. *Comm. Partial Differential Equations*, 28(7-8):1237–1263, 2003.
- [14] P. Gamblin and X. Saint Raymond. On three-dimensional vortex patches. *Bull. Soc. Math. France*, 123(3):375–424, 1995.
- [15] C. Huang. Singular integral system approach to regularity of 3D vortex patches. *Indiana Univ. Math. J.*, 50(1):509–552, 2001.
- [16] T. Kato. Nonstationary flows of viscous and ideal fluids in R^3 . *J. Functional Analysis*, 9:296–305, 1972.
- [17] T. Kato. Quasi-linear equations of evolution, with applications to partial differential equations. In *Spectral theory and differential equations (Proc. Sympos., Dundee, 1974; dedicated to Konrad Jörgens)*, pages 25–70. Lecture Notes in Math., Vol. 448. Springer, Berlin, 1975.
- [18] A. J. Majda and A. L. Bertozzi. *Vorticity and incompressible flow*, volume 27 of *Cambridge Texts in Applied Mathematics*. Cambridge University Press, Cambridge, 2002.
- [19] N. Masmoudi. The Euler limit of the Navier-Stokes equations, and rotating fluids with boundary. *Arch. Rational Mech. Anal.*, 142(4):375–394, 1998.
- [20] H. S. G. Swann. The convergence with vanishing viscosity of nonstationary Navier-Stokes flow to ideal flow in R^3 . *Trans. Amer. Math. Soc.*, 157:373–397, 1971.
- [21] W. Wolibner. Un théorème sur l’existence du mouvement plan d’un fluide parfait, homogène, incompressible, pendant un temps infiniment long. *Math. Z.*, 37(1):698–726, 1933.
- [22] V. I. Yudovich. Non-stationary flows of an ideal incompressible fluid. *u Z. Vy cisl. Mat. i Mat. Fiz.*, 3:1032–1066, 1963.