

Remarks on the Development and Recent Results in the Theory of Gibbs Derivatives

Radomir S. Stanković and Jaakko Astola

Abstract: This tutorial paper discusses the development of the theory of Gibbs differential operators, and highlights some important older and more and recent results in this area.

Keywords: Gibbs derivatives, Walsh functions, finite dyadic group.

1 Introduction

FOURIER ANALYSIS and differential calculus are two very important tools in signal processing and related areas. From a group theoretic point of view, these two essential concepts can be related by the observation that classical Newton-Leibniz derivative can be viewed as a closed linear operator on the real line R that maps each group character of R $e^{2\pi ikx}$ into the scalar multiple $2\pi ike^{2\pi ikx}$ of that character, and by recalling that exponential functions, eigenfunctions of differential operators, are at the same time kernels in the Fourier representations of signals.

Attempts towards extending classical Fourier analysis to function systems different than group characters of R emerged in part also due to convergence problems in approximation of real valued functions. In this context, different function systems, such as the Walsh functions [77] and the Haar functions [35], have been intentionally invented to resolve related problems.

Fine [15] and Vilenkin [74] observed independently that Walsh functions can be identified with group characters of the dyadic or Cantor group. In this way, the Walsh functions have been fully incorporated in the Walsh-Fourier analysis as a particular example of harmonic analysis on groups.

Manuscript received on October 15, 2008.

J. Astola is with Dept. of Signal Processing, Tampere University of Technology, FI-33101 Tampere, Finland (e-mail Jaakko.Astola@tut.fi). R. Stanković is with Dept. of Computer Science, Faculty of Electronics, 18 000 Niš, Serbia (e-mail rstankovic@bankerinter.net)

In late 60's and early 70' there emerged a high interest in the discrete counterpart of Walsh functions due to their compatibility with binary signals and bistable circuits, and for the simplicity of the calculation of the spectral Walsh coefficients without multiplication of complex numbers that is necessary in the discrete Fourier analysis.

Ideas to extend the theory to the discrete Walsh functions by linking differential operators and group characters naturally emerged in the research community. In particular, such an initiative was suggested to J. Edmund Gibbs by Dr Alastair Gebbie, at that time the Head of the Basic Physic Division of National Physical Laboratory in Teddington, Middlessex, England. The first complete formulation of the definition of the Gibbs derivatives has been made by James Edmund Gibbs on January 13, 1967. It has been completely presented in [19].

The especially simple case of the Gibbs derivative on a finite dyadic group, called initially logic derivative or finite dyadic derivative, is of importance not only for its own sake but also how this conceptually new approach to differentiation first emerged [25–28]. The subsequent development of this theory and highlighting some important results are the subject of this paper. For the limited space, we have to restrict the selection of references to the publications where certain results were reported for the first time and in exceptional cases to some other relevant publications. An extensive bibliography of Gibbs derivatives can be found in [69]. In [69], we provide some insight into the bibliography on Gibbs derivatives.

2 Notation and Definitions

For each positive integer n , let G_n denote the finite dyadic group whose elements are n -tuples $x = (x_0, x_1, \dots, x_{n-1})$ with $x \in \{0, 1\}$. The group operation \oplus is coordinatewise addition modulo 2, that is, for each $x, \xi \in G_n$, $x \oplus \xi = (x_0 \oplus \xi_0, \dots, x_{n-1} \oplus \xi_{n-1})$. Denote by B_n the set of non-negative integers less than 2^n : $B_n = \{0, 1, \dots, 2^n - 1\}$.

With each element x of G_n we associate a unique element of B_n by means of a function $V_n : G_n \rightarrow B_n$ defined by

$$V_n(x) = \sum_{i=0}^{n-1} 2^{n-i-1} x_i. \quad (1)$$

The element of B_n thus defined is usually called, in switching theory, the "decimal index" of x . The function V_n is bijective, so to each integer of B_n corresponds under V_n^{-1} a unique element of G_n whose coordinates are the coefficients in the dyadic expansion of that integer. The existence of the bijection V_n enables us safely to ignore the distinction between G_n and B_n . Thus we shall denote by x indifferently the n -tuple (x_0, \dots, x_{n-1}) or its decimal index $V_n((x_0, \dots, x_{n-1}))$. We denote by

\oplus indifferently the operation already defined in G_n or the operation in B_n defined by

$$x \oplus \xi = V_n(V_n^{-1}(x) \oplus V_n^{-1}(\xi)), \quad (x, \xi \in B_n).$$

In the right member of this equality, \oplus is the operation in G_n ; in the left member, \oplus is the operation in B_n . Thus $V_n : G_n \rightarrow B_n$ is a group isomorphism.

The space of all complex-valued functions f on G_n (or on B_n) will be denoted by L_n , or, since it is not usually necessary to express the dependence on n , simply L .

Notice that the finite dyadic group of order 2^n is equal to the direct n -th power of the cyclic group $G_1 = (\{0, 1\}, \oplus)$. This enables us to write (the value-vectors of) the discrete Walsh functions $wal(w, \cdot) : B_n \rightarrow \{1, -1\}$, ($w \in B_n$) (the characters of the group G_n) as the rows (or columns) of a matrix \mathbf{W}_n defined by

$$\mathbf{W}_n = \left[\begin{array}{cc} 1 & 1 \\ 1 & -1 \end{array} \right]^n, \tag{2}$$

where $[\cdot]^n$ denotes the continued Kronecker product of n copies of the matrix $[\cdot]$.

Using the discrete Walsh functions as an orthogonal basis in the space L_n , we may define the discrete Walsh transform as a particular case of the Fourier transform on groups. Thus, the Walsh transform coefficients $S_f(w)$ ($w \in B_n$) of $f \in L$ are defined by

$$S_f(w) = \sum_{x=0}^{2^n-1} f(x)wal(w,x). \tag{3}$$

The inverse Walsh transform is given by

$$f(x) = 2^{-n} \sum_{w=0}^{2^n-1} S_f(w)wal(w,x). \tag{4}$$

If we express the Walsh spectral coefficients $S_f(w)$ and the original function f as vectors of order 2^n , $\mathbf{S}_f = [S_f(0), \dots, S_f(2^n - 1)]^T$ and $\mathbf{F} = [f(0), \dots, f(2^n - 1)]^T$, then the equalities (3) and (4) may be written

$$\mathbf{S}_f = \mathbf{W}_n \mathbf{F}, \tag{5}$$

$$\mathbf{F} = 2^{-n} \mathbf{W}_n \mathbf{S}_f. \tag{6}$$

The discrete dyadic convolution product $f * g$ of two functions $f, g \in L$ is defined by

$$(f * g) = \sum_{u=0}^{2^n-1} f(x \oplus u)g(u), \quad (x \in B_n).$$

The dyadic convolution theorem states that, if the Walsh transforms of f and g are S_f and S_g , respectively, then the transform of $f \cdot g$ is $S_f * S_g$ and the transform of $f * g$ is $S_f \cdot S_g$.

3 The Gibbs Derivative on Finite Dyadic Groups

The initial definition of the Gibbs derivative has been formulated for functions on finite dyadic group G_n as follows [19].

Definition 1. To each function $f \in L$ we assign a function $f^{[1]} \in L$ defined by

$$f^{[1]}(x) = -\frac{1}{2} \sum_{r=0}^{n-1} (f(x \oplus 2^r) - f(x))2^r, \quad (x \in B_n). \quad (7)$$

We call $f^{[1]}$ the (first-order) Gibbs derivative of f . The corresponding operator $D_n : L \rightarrow L$, defined by $D_n f = f^{[1]}$ ($f \in L$), will be called the Gibbs differentiator $B_n \rightarrow C$, where C is the field of complex numbers.

Nomenclature of this type enables us to distinguish each of the various species of Gibbs differentiator, by indicating the domain and range of the functions on which it operates. To simplify the notation we shall generally abbreviate D_n as D .

If $f : B_n \rightarrow C$ is identified with the corresponding function ¹ defined on G_n , and if we use the same symbol f for this function $f : G_n \rightarrow \mathbf{K}$, then Definition 1 may be written

$$f^{[1]}(x) = -\frac{1}{2} \sum_{i=0}^{n-1} 2^{n-i-1} (D_i f)(x), \quad (8)$$

where $x = (x_0, \dots, x_{n-1})$, and D_i ($i = 0, \dots, n-1$) denotes the (i -th coordinate) partial Gibbs differentiator, defined ² by

$$(D_i f)((x_0, \dots, x_{n-1})) = f((x_0, \dots, x_i \oplus 1, \dots, x_{n-1})) - f((x_0, \dots, x_{n-1})).$$

The following theorem summarises some of the chief properties of the Gibbs differentiator $B_n \rightarrow \mathbf{K}$ (or $G_n \rightarrow C$).

Theorem 1. The operator $D : L \rightarrow L$ has the following properties:

¹This function should strictly be denoted by the symbol fV_n , where juxtaposition denotes composition.

²Let the complement $x \oplus 1$ of $x \in \{0, 1\}$ be denoted by \bar{x} . Then the i -th coordinate partial Gibbs derivative may be written

$$(D_i f)((x_0, \dots, x_{n-1})) = f((x_0, \dots, \bar{x}_i, \dots, x_{n-1})) - f((x_0, \dots, x_i, \dots, x_{n-1})).$$

1. *Linearity:* $D(a_1f_1 + a_2f_2) = a_1f_1 + a_2f_2$, $(a_1, a_2 \in C, f_1, f_2 \in L)$.
2. $Df = O \in L$ iff $f \in L$ is a constant function.
3. If the Walsh transform of $f \in L$ is S_f , then the Walsh transform of Df is given by

$$S_{Df} = wS_f. \quad (9)$$

4. $D(f_1 * f_2) = (Df_1) * f_2 = f_1 * (Df_2)$, $(f_1, f_2 \in L)$.
5. For each $\alpha \in B_n$, the operator D commutes with the translation operator T_α defined by $(T_\alpha f)(x) = f(x \oplus \alpha)$, $(f \in L, x \in B_n)$:

$$DT_\alpha = T_\alpha D.$$

6. The operator D does not obey the product rule satisfied by the Newton- Leibniz differentiator D_{NL} , namely,

$$\begin{aligned} D_{NL}(f_1f_2) &= f_1(D_{NL}f_2) + (D_{NL}f_1)f_2, \\ &(f_1, f_2 : R \rightarrow R, \text{ say}). \end{aligned}$$

In other words, for some $f_1, f_2 \in L$,

$$D(f_1f_2) \neq f_1(Df_2) + (Df_1)f_2.$$

Proof. The properties 1 to 6 are easily derived from Definition 1 and properties of the Walsh transform.

By using the property 3 of Theorem 1 it is easy to prove that the set of 2^n discrete Walsh functions defined on B_n is the set of eigenfunctions of the Gibbs differentiator $B_n \rightarrow \mathbf{K}$; that is, if we write f_w for $wal(w, \cdot)$ ($w \in B_n$), then

$$f^{[1]} = wf_w. \quad (10)$$

In other words, the discrete Walsh functions are the 2^n solutions of the eigenvalue problem

$$D_n f = wf, \quad (11)$$

of which the eigenvalues are $w = 0, 1, \dots, 2^n - 1$, and the corresponding eigenfunctions are $wal(0, \cdot), wal(1, \cdot) \dots wal(2^n - 1, \cdot)$, respectively [21].

In this context we distinguish Gibbs derivatives of the first and second kinds, corresponding respectively to the Walsh-Paley [59] and Walsh-Kaczmarz [41] orderings of the Walsh functions [20].

The Gibbs derivative can be extended to arbitrary complex order p by way of the definition of the discrete δ -function

$$\delta(x) = 2^{-n} \sum_{w=0}^{2^n-1} wal(w,x) \quad (x \in B_n).$$

The function thus defined has the properties

$$\delta(x) = \begin{cases} 1, & (x = 0), \\ 0, & (x \neq 0), \end{cases}$$

and

$$f = \delta * f, \quad (f \in L).$$

The derivative of order $p \in \mathbf{K}$ of the δ -function is defined by an extension of the equality (10):

$$\delta^{[p]}(x) = 2^{-n} \sum_{w=0}^{2^n-1} w^p wal(w,x). \quad (12)$$

The derivative of order p of an arbitrary function $f \in L$ may be conveniently obtained as the dyadic convolution product of $\delta^{[p]}$ and f . For, by an easily proved extension of the property 4,

$$D^p f = f^{[p]} = \delta * f^{[p]} = \delta^{[p]} * f. \quad (13)$$

The *Gibbs derivative of fractional order* is obtained in the case that p is real.

4 Generalizations of the Gibbs Derivative Through Spectral Interpretation

We note first that the property 3 of Theorem 1 serves as an alternative definition of the Gibbs derivative. More precisely, the Gibbs derivative of a function $f \in L$ is the function $f^{[1]} \in L$ such that

$$S_{f^{[1]}}(w) = wS_f(w), \quad (w \in B_n).$$

The differentiator operating on the space of complex-valued functions on each locally compact Abelian group has the characters of that group as eigenfunctions.

The first attempt at such a definition, restricted to finite Abelian groups, was made by Gibbs and Ireland [29]. These authors regarded such a group as the direct product of indecomposable cyclic groups, and tried to define the differentiator

on such a cyclic group consistently with the recapture, as a limiting case, of the classical derivative (see [27]). The differentiator on a finite direct product of indecomposable cyclic groups is defined as a linear combination of the partial differentiators with respect to the direct factors, the numerical coefficients being chosen to yield a convenient set of eigenvalues (essentially, for a group of order g , the least g non-negative integers).

Onneweer [46] independently gave the analogous definition on a countable direct product of groups of prime order, effectively amending and extending the definition of the earlier authors.

Ren Fu-xian, Su Weiyi, and Zheng Wei-xing [16] in 1978 and Zheng Wei-jing and Su Weiyi in 1981 [79] gave a definition (reproduced as **Definition B** by Zelin He in 1983 [36]) on the p -adic groups. This result by Zelin He, has been referred in the further work by the same author [37].

Moraga [44], has discussed the particular case of Gibbs differentiation on finite direct powers of a group of prime order: in this case the eigenfunctions are the discrete Levy-Vilenkin-Chrestenson functions [10, 42, 74].

The extension of Gibbs differentiation to non-Abelian groups. Such a definition is obtained [65] where the dyadic derivative of a function is expressed in terms of its Walsh-Fourier coefficients. The role of the group characters in the definition of the Gibbs derivatives on Abelian groups is taken over, in the case of the derivatives on non-Abelian groups, by the unitary irreducible representations of these groups [65]. The definition applies also to functions with codomain a finite field, provided that this field admits a Fourier transform [67]. The Gibbs derivatives on finite non-Abelian groups may be regarded as linear harmonic translation-invariant systems [64]. An extension of the notion of Gibbs differentiation to matrix-valued functions on finite non-Abelian groups has been done in [72].

4.1 Butzer-Wagner Dyadic Derivative

A major contribution to the theory of dyadic differentiation has been made by Butzer and Wagner in a sequence of publications starting with [6] and [9]. These authors were foremost in recognising the value of the Gibbs derivative in analysis, just as Pichler, from 1970 onwards, led the way in applications to system theory [61].

The dyadic group G is (isomorphic to) the set of all sequences $x = (x_1, x_2, \dots)$ with the group operation \oplus being termwise addition modulo 2. Let $X(G)$ denote one of the spaces $C(G)$ (the space of dyadic continuous functions on G) and $L^p(G)$ ($1 \leq p < \infty$) (that of p -th power Lebesgue-integrable functions on G). The series

used to define the derivative in either the pointwise or the norm sense [9], [75] is

$$(Df)(x) = \frac{1}{2} \sum_{j=0}^{\infty} 2^j (f(x) - f(x \oplus e_{j+1})),$$

where $f \in X(G)$, $x \in G$, $e_j = (\delta_{j,1}, \delta_{j,2}, \dots, \delta_{j,k}, \dots)$. For a function $f \in X(G)$, if there exist $g \in X(G)$, such that

$$\lim_{m \rightarrow \infty} \left\| \frac{1}{2} \sum_{j=0}^m (f(x) - f(x \oplus e_{j+1})) - g(x) \right\|_{X(G)} = 0,$$

then $g(x)$ is the first strong dyadic derivative of f .

For a function f defined on $[0, 1)$, if

$$\frac{1}{2} \sum_{j=0}^{\infty} 2^j (f(x) - f(x \oplus e_{j+1})) = c < \infty,$$

for $x \in [0, 1)$, then c is called the first pointwise dyadic derivative of f at x .

Butzer and Wagner [6] use the Fine map to transfer the definition of the strong derivative to periodic functions on the real group, of period 1.

In a further paper, the same authors [7] define Gibbs derivatives of the second kind (corresponding to the Walsh-Kaczmarz ordering of the Walsh functions) for functions defined on G , in both the norm sense and the pointwise sense. In the following year [8], they extend to functions on the non-negative real line the definitions of both strong and pointwise derivatives (of the first kind, corresponding to the Walsh-Paley ordering). A further paper [60] provides a readable summary of the results up to that time. The booklet of Wagner [75] gives a helpful introduction to many analytical aspects of dyadic differentiation, treating in parallel the differentiators of the first and second kinds. A generalization of Butzer-Wagner differentiation was considered in [57, 58] and few other publications [69].

The class of dyadically differentiable functions was greatly extended by Butzer, Engels, and Wipperfurth [4], by their definition of an *extended dyadic (ED) derivative*. They defined the ED-derivative as the result of applying the Euler summation process to the series obtained by applying a certain sequence of multipliers to the coefficients of the Walsh-Fourier series of the conventional dyadic derivative. This remarkable definition is justified by its success in extending the class of differentiable functions, for example, to piecewise polynomial functions and to the products of such functions with Dirichlet's function. Butzer and Engels [3] subsequently defined the ED-derivative directly in the original function space, not in the transform space.

4.2 Dyadic derivative on R_+

Differentiation of functions on R_+ has been initiated in [53], and [56, 76].

The dyadic differentiators on $[0, 1)$ and on R_+ may be defined by using the Walsh-Fourier coefficients in a manner that is an extension of the equality (9) above. Zelin He in 1983 [36] has generalised this approach to define the dyadic derivatives of fractional orders $\alpha \in R$. This definition, in case $\alpha = r \in N$, reduces to that of Butzer and Wagner for the dyadic derivative of integer order; and, in case $\alpha = -r$, ($r \in N$), to that of Butzer and Wagner for the dyadic antiderivative of integer order. The extended dyadic derivative of Butzer, Engels, and Wipperfurth [4] is defined at the outset for fractional orders $\alpha \in R$. The definition therefore covers, by the case $\alpha < 0$, antidifferentiation.

Zelin He [38] further related the Gibbs derivative and integral to Walsh-Fourier integral operators, and defined the concept of the p -adic differential-integral-type operator with order-type (λ_1, λ_2) . This is a far-reaching generalisation, in a certain direction, of the concepts of Gibbs derivative and integral, and leads to further applications of the Gibbs derivative to p -adic approximation theory. See, for instance, a related discussion in [69].

The definition of dyadic differentiation on the real half-line [8] has been extended by Engels and Splettstösser [14] to square-integrable stochastic processes, in particular, dyadic-stationary processes. Iff such a derivative exists for a process X , X is said to be dyadically differentiable i.m. (in the mean). A necessary and sufficient condition for a Walsh-harmonisable³ dyadic-stationary process to be dyadically differentiable has been given by Endow in 1987 [13].

F. Weisz studied in [82, 83] a general summability method of different orthogonal series by using an integral function θ . The results derived have been verified also for multidimensional dyadic derivatives [84].

4.3 Modified dyadic derivatives and dyadic distributions

A modified definition of the strong dyadic derivative has been discussed in [30]. This research has been continued in [32], where also the fractional derivative of order $\alpha > 0$ have been considered. There have been shown criteria for the existence of these differential operators and their inverse integral operators, and determined a countable set of eigenfunctions of these operators. Some related applications of these results were reported in [31].

The paper [33], reviews results related to the pointwise and strong dyadic

³A dyadic-stationary process is said to be Walsh harmonisable iff it assumes its spectral representation in terms of the generalised Walsh functions. A characterization of such processes has been given by Endow [12] in 1984.

derivative and integral for functions on R_+ . Based on these results introduced are the modified dyadic strong and pointwise derivatives and integrals of fractional orders on R_+ and some of their properties studied. Study of properties of modified fractional dyadic derivative and integral has been continued in [34].

4.4 Differentiation on p -adic groups and generalizations

The definition of dyadic differentiation has been further discussed by Onneweer in 1977, 1978, and 1979 [46, 47, 49], [52] and [53] and [54]. Every acceptable definition has the property that the eigenfunctions of the dyadic differentiator are the Walsh functions, but the corresponding eigenvalues are, and therefore the definition is, to some extent arbitrary, depending on, among other things, the chosen enumeration (Paley or Kaczmarz) of the Walsh functions. Onneweer gives a definition such that the set of Walsh functions is partitioned into classes each having a common eigenvalue: this partition is independent of the choice of enumeration, and consequently so is the definition. The eigenvalues of the respective classes of Walsh functions are analogues of frequency [22–24]. This suggests that this type of definition may be better adapted to applications in the exact sciences. This definition has subsequently been extended to functions defined on p -adic and p -series fields [48, 50], p -adic groups [49], and a local field by Onneweer [51], [52].

In [80] and [81], presented is the definition of a derivative in terms of pseudo-differential operators.

There have been defined also L' -weak p -adic derivative, the adjacent p -adic derivative, the partial p -adic derivative [16, 39, 78]. Extensions of Gibbs differentiation to a -adic groups has been provided in [40], and [43].

Further contributions to the theory of Gibbs differential operators on locally compact Vilenkin groups are done in [85, 86].

4.5 Gibbs derivatives with respect to Haar functions and generalizations

By analogy with the conventional dyadic calculus (related to Walsh functions), derivatives and antiderivatives have been introduced related to the Haar system of orthogonal functions. These definitions enable a calculus to be built up that plays the same role in Haar-Fourier analysis as the conventional dyadic calculus plays in Walsh-Fourier analysis [5, 55].

A Haar derivative for complex-valued functions on a finite dyadic group has been defined by Stanković and Stojić [73], using an analogue of the matrix representation of the discrete dyadic derivative, see, for instance, [71]. These authors (1987) use the same approach to define a family of differentiators having the dis-

crete generalised Haar functions [1] as eigenfunctions. A generalisation of the Gibbs derivative to an arbitrary orthogonal basis in the space of functions from an arbitrary finite Abelian group to the complex, or a finite (Galois), field is given in [66, 68]. Extension of Gibbs differentiation to functions with finite fields as codomains had been discussed earlier by Cohn-Sfetcu and Gibbs [11]. The Gibbs derivatives for multiple-valued logic functions have been discussed in [70].

5 Recent Results

Recent development of Gibbs differentiation have been done in two-parameter differentiation, as for instance, the fundamental theorem of two-parameter pointwise derivative on Vilenkin groups [17] has been presented in [18, 45]. The monograph [82] is the first monograph which considers the theory of more-parameter dyadic and classical Hardy spaces, including also d -dimensional dyadic derivatives. See, also [83, 84]. Different generalizations of the Gibbs dyadic derivative have been used in study of local fields [87–89].

From the point of view of practical applications, research have been concentrated in applications of various generalizations of Gibbs differential operators in theory of fractals and fractal functions. Notice, for instance, that in [62], it is introduced a kind of Weierstrass-like function in p -series local field and discussed their its p -adic derivative with targeted applications in determination of the "rate of change" of fractal functions in local fields. In [63], the 3-adic Cantor functions on 3-series field is constructed and its 3-adic derivative evaluated, and shown that it has at most $\ln 2/\ln 3$ order. A recently compiled bibliography on Gibbs derivatives comprises 313 bibliographic items [69].

Acknowledgments

The authors are grateful to Prof. Claudio Moraga for comments and suggestions that improved the presentation in the paper.

This work was supported by the Academy of Finland, Finnish Center of Excellence Programme, Grant No. 213462.

References

- [1] N.N. Aizenberg, V.P. Rudko, and E.V. Sisuev. Haar functions and discrete mappings. *Tekhn. Kibernetika*, 6:86–94, 1975. (in Russian).
- [2] P.L. Butzer. *Theory and Applications of Gibbs Derivatives*. Matematički Institut, Beograd, Serbia, 1990.

- [3] P.L. Butzer and W. Engels. Background to an extension of Gibbs differentiation in Walsh analysis. In P.L. Butzer and R. S. Stanković, editors, *Theory and Applications of Gibbs Derivatives*, pages 19–57. Matematički Institut, Belgrade, Serbia, 1990.
- [4] P.L. Butzer, W. Engels, and U. Wipperfurth. An extension of the dyadic calculus with fractional order derivatives: general theory. *Comp. & Math. with Appls.*, 12B(5/6):1073–1090, 1986.
- [5] P.L. Butzer, W. Splettstösser, and H.J. Wagner. On the role of Walsh and Haar functions in dyadic analysis. In H. Hubner, editor, *Proc. Spec. Sess. Sequency Techniques*, pages 1–6, Montreux, Darmstadt, May 20–22, 1975. (in co-operation with the 1st Sympos. & Tech. Exhib. EMC).
- [6] P.L. Butzer and H.J. Wagner. Approximation by Walsh polynomials and the concept of a derivative. In *Proc. Sympos. Applic. Walsh Functions*, pages 388–392, Washington, D.C., 1972.
- [7] P.L. Butzer and H.J. Wagner. On a Gibbs-type derivative in Walsh-Fourier analysis with applications. In *Proc. Nat. Electron. Conf.*, volume 27, pages 393–398, 1972.
- [8] P.L. Butzer and H.J. Wagner. A calculus for Walsh functions defined on R_+ . In *Proc. Sympos. Applic. Walsh Functions*, pages 75–81, Washington, D.C., 1973.
- [9] P.L. Butzer and H.J. Wagner. Walsh-Fourier series and the concept of a derivative. *Applicable Anal.*, 3:29–46, 1973.
- [10] H.E. Chrestenson. A class of generalized Walsh functions. *Pacific J. Math.*, 5:17–31, 1955.
- [11] S. Cohn-Sfetcu and J.E. Gibbs. Harmonic differential calculus and filtering in Galois fields. In *Proc. IEEE Int. Conf. Acoust. Speech & Signal Processing*, pages 148–153, Philadelphia, PA, April 12–14, 1976.
- [12] Y. Endow. Analysis of dyadic stationary processes using the generalized Walsh functions. *Tohoku Math. J., Ser. 2*, 36(4):485–503, 1984.
- [13] Y. Endow. On a condition for the dyadic differentiability of dyadic stationary processes. *Repts Statist. Appl. Res., JUSE*, 34:1–5, 1987.
- [14] W. Engels and W. Splettstösser. On Walsh differentiate dyadically stationary random processes. *IEEE Trans.*, IT-28(4):612–619, 1982.
- [15] N.J. Fine. On the Walsh functions. *Trans. Amer. Math. Soc.*, 65:372–414, 1949.
- [16] Ren Fu-xian, Su Wei-yi, and Zheng Wei-xing. The generalized logical derivative and its applications. *J. Nanjing Univ.*, 3:1–8, 1978. (in Chinese, English Abstract).
- [17] Gy. Gát. On the two-dimensional pointwise dyadic calculus. *J. Approx. Theory*, 92:191–215, 1998.
- [18] Gy. Gát and J. Nagy. The fundamental theorem of two-parameter pointwise derivative on Vilenkin groups. *Analysis Math.*, 25:33–55, 1999.
- [19] J.E. Gibbs. Walsh spectrometry, a form of spectral analysis well suited to binary digital computation. Technical Report 24, Nat. Phys. Lab., Teddington, Middx, UK, 1967.
- [20] J.E. Gibbs. Sine waves and Walsh waves in physics. In *Proc. Sympos. Applic. Walsh Functions*, pages 260–274, Washington, D.C., March 31/April 3, 1970.
- [21] J.E. Gibbs. Walsh functions and the Gibbs derivative. Technical Report 10, NPL DES Memo., 1973.
- [22] J.E. Gibbs. The concept of frequency in dyadic analysis. In *Spec. Sess. Sequency Techniques, EMC Sympos. Exhib.*, page 5, Montreux, 1975.
- [23] J.E. Gibbs. Differentiation and frequency on the dyadic group. Technical Report 18, NPL DES Rept., 1975.

- [24] J.E. Gibbs. Eine Erweiterung der harmonischen Analyse auf den dyadischen Körper als Funktionenraum. In *Kurzfass. Kolloq. Theor. Anwend. diskret. Signale*, pages 135–138, Aachen, October 7–9, 1976.
- [25] J.E. Gibbs. Harmonic analysis in the dyadic field regarded as a function space. In *Seminar, Royal Signals & Radar Estab.*, pages ii+24, October 28, 1976.
- [26] J.E. Gibbs. Towards a pure dyadic differentiator. Technical report, NPL rept of discussion held at Lehrstuhl A für Math., RWTH Aachen, October 4, 1979.
- [27] J.E. Gibbs. Local and global views of differentiation. In [2], pages 1–18. 1990.
- [28] J.E. Gibbs and B. Ireland. Significance of the logical differential calculus. In *Proc. Sympos. Digital Filtering*, London, U.K., 1971. Imperial Coll.
- [29] J.E. Gibbs and B. Ireland. Walsh functions and differentiation. In H. Schreiber and G.F. Sandy, editors, *Applications of Walsh Functions and Sequency Theory*, pages 147–176, New York, 1974. IEEE.
- [30] B.I. Golubov. On modified strong dyadic integral and derivative. *Ukrainian Mathematical Journal*, 54(5):770–784, 2002.
- [31] B.I. Golubov. A dyadic analogue of Wiener’s Tauberian theorem and related questions. *Izv. Math.*, 67(1):29–53, 2003. translation from *Izv. Ross. Akad. Nauk, Ser. Mat.*, 67, 1, 33–58, 2003.
- [32] B.I. Golubov. Dyadic fractional differentiation and integration of Walsh transform. In Shyam L. Kalla et al., editors, *Proceedings of the International Conference on Mathematics and Its Applications, ICMA*, pages 274–284, Kuwait, April 5–7, 2004.
- [33] B.I. Golubov. Modified dyadic derivative and integrals of fractional order on R_+ . *Acta Mathematica Academiae Paedagogicae Nyiregyháziensis*, 20:153–163, 2004.
- [34] B.I. Golubov. On some properties of fractional dyadic derivative and integral. *Analysis Mathematica*, 32(3):173–205, September 2006.
- [35] A. Haar. Zur Theorie der Orthogonalen Funktionensysteme. *Math. Ann.*, 69:331–371, 1910.
- [36] Zelin He. The derivatives and integrals of fractional order in Walsh-Fourier analysis, with applications to approximation theory. *J. Approx. Theory*, 39:361–373, 1983.
- [37] Zelin He. General linear summation of the Vilenkin-Fourier series. *Acta Math. Hungar.*, 46(3-4):285–295, 1985.
- [38] Zelin He. The p -adic differential-integral type operator. *Anal. Math.*, 19:65–84, 1993. Fasc. 1.
- [39] H.K. Jiang. The kernels of de la Vallee Poussin type on p -adic fields. *Approx. Theory & Appl.*, 6(1):65–79, 1990.
- [40] H.K. Jiang. The derivatives and integrals of fractional order on a -adic groups. *Chin. Ann. of Math., Ser. B*, 14(4):515–526, 1993.
- [41] S. Kaczmarz. Über ein Orthogonal Systeme. In *Comptes rendus du premier congrès des mathématiciens des pays Slaves*, pages 189–192, Varsovie, 1929, 1930.
- [42] P. Levy. Sur une généralisation des fonctions orthogonales de M. Rademacher. *Comment. Math. Helv.*, 16:146–152, 1944. Translation by Bramhall, Janet N., On a generalization of the Rademacher orthogonal functions, Translation T-2659, The R.E. Gibson Library Bulletin, The Johns Hopkins University Applied Physics Laboratory, 8621 Georgia Avenue, Silver Spring, MD 20910, i+8, 1972.
- [43] Y. Li and Weiyi Su. Random a -adic groups and random net fractals. *Chaos, Soliton & Fractals*, to appear.

- [44] C. Moraga. Introduction to linear p -adic systems. In R. Trappl, editor, *Cybernetics and Systems Research 2*. North-Holland, 1984.
- [45] K. Nagy and Gy. Gát. The fundamental theorem of two-parameter pointwise derivative on Vilenkin groups. *Analysis Math.*, 25:33–55, 1999.
- [46] C.W. Onneweer. Differentiability for Rademacher series on groups. *Acta Sci. Math.*, 39:121–128, 1977.
- [47] C.W. Onneweer. Fractional differentiation on the group of integers of a p -adic or p -series field. *Anal. Math.*, 3:119–130, 1977.
- [48] C.W. Onneweer. On the pointwise differentiability of functions on a p -adic or p -series field. Technical Report 342, Univ. New Mexico, 1977.
- [49] C.W. Onneweer. Differentiation of fractional order on p -groups, approximation properties. *Anal. Math.*, 4:297–302, 1978.
- [50] C.W. Onneweer. Differentiation on a p -adic or p -series field. In P. L., editor, *Linear Spaces and Approximation*, pages 187–198. Birkhauser, Basel, 1978.
- [51] C.W. Onneweer. On the definition of dyadic differentiation. *Applicable Anal.*, 9:267–278, 1979.
- [52] C.W. Onneweer. On pointwise differentiability of functions on a p -adic or p -series field. *Ann. Univ. Sci. Budapest. R. Eotvos Nomin., Sect. Math.*, (49-57):22–23, 1979–1980.
- [53] J. Pál. On the dyadic calculus for functions defined on R_+ . In *Kurzfass. Kolloq. Theor. Anwend. diskret. Signale*, pages 250–253, Aachen, October 7–9, 1976.
- [54] J. Pál. On almost everywhere differentiability of dyadic integral functions on R_+ . In *Colloq. Math. Soc. J. Bolyai, Budapest 1976*, volume II of *Fourier Analysis and Approximation Theory*, pages 591–601. 1978.
- [55] J. Pál. On the dyadic Haar derivative. In *Constructive Theory of Functions '84, Sofia*, pages 658–663, 1984.
- [56] J. Pál and F. Schipp. On the a.e. dyadic differentiability of dyadic integral on R_+ . In [2], pages 103–113. Matematički institut, Belgrade, Serbia, 1990.
- [57] J. Pál and P. Simon. On a generalization of the concept of derivative. *Acta Math. Acad. Scient. Hungar.*, 29(1-2):155–164, 1977.
- [58] J. Pál and P. Simon. On the generalized Butzer-Wagner type a.e. differentiability of integral function. *Ann. Univ. Scient. Budapest. R. Eötvös Nomin., Sect. Math.*, 20:157–165, 1977.
- [59] R.E.A.C. Paley. A remarkable series of orthogonal functions. *Proc. Lond. Math. Soc.*, 34:241–279, 1932.
- [60] F.R. Pichler. Some aspects of a theory of correlation with respect to Walsh harmonic analysis. Technical Report Rept R-70-11, Univ. Maryland Tech. Res., 1970.
- [61] F.R. Pichler. Walsh functions and linear system theory. In *Proc. Sympos. Applic. Walsh Functions*, pages 175–182, Washington, D.C., 1970.
- [62] H. Qiu and Su Weiyi. Weierstrass-like functions on local fields and their p -adic derivatives. *Chaos, Solitons & Fractals*, 28(4):958–965, 2006.
- [63] H. Qiu and Su Weiyi. 3-adic Cantor function on local fields and its p -adic derivative. *Chaos, Solitons and Fractals*, 33:1625–1634, 2007.
- [64] R.S. Stanković. Linear harmonic translation invariant systems on finite non-Abelian groups. In R. Trappl, editor, *Cybernetics and Systems '86*, pages 103–110. Reidel, Dordrecht, 1986.

- [65] R.S. Stanković. A note on differential operators on finite non-Abelian groups. *Applicable Anal.*, 21(1-2):31–41, 1986.
- [66] R.S. Stanković. A note on differential operators on finite Abelian groups. *Cybernetics & Systems*, 18:221–231, 1987.
- [67] R.S. Stanković. Gibbs derivatives on finite non-Abelian groups. In [2], pages 269–297. 1990.
- [68] R.S. Stanković. Some remarks on fourier transform and differential operators for digital functions. In *Proc. 22nd Int. Symp. on Multiple-Valued Logic*, pages 365–370, Sendai, Japan, May 27–29, 1992.
- [69] R.S. Stanković and J. Astola. *Gibbs Derivatives - the First Forty Years*. TICSP Series #39. Tampere International Center for Signal Processing, Tampere, Finland, 2008.
- [70] R.S. Stanković, C. Moraga, and J.T. Astola. Derivatives for multiple-valued functions induced by Galois field and Reed-Muller-Fourier expressions. In *Proc. 34th Int. Symp. on Multiple-Valued Logic*, pages 184–189, Toronto, Canada, May 19–22, 2004.
- [71] R.S. Stanković, C. Moraga, and J.T. Astola. *Fourier Analysis on Finite Groups with Applications in Signal Processing and System Design*. IEEE Press, Wiley & Sons, Inc., 2005. Electronics book edition 1, ISBN=0471745421.
- [72] R.S. Stanković, M.S. Stanković, and C. Moraga. Remarks on systems and differential operators on groups. *Facta. Univ., Ser.: Elect. Energ.*, 18(3):531–545, 2005.
- [73] R.S. Stanković and M.R. Stojić. A note on the discrete Haar derivative. In *Colloq. Math. Soc. J.*, pages 897–907, Bolyai Budapest, 1985.
- [74] N.Ya. Vilenkin. On a class of complete orthonormal systems. *Izv. Akad. Nauk, Ser. Mat.*, 11:363–400, 1947. Translation by Boas, R.P., *Amer. Math. Soc. Transl.*, 28, Ser. 2, 1-35.
- [75] H.J. Wagner. *Ein Differential- und Integralkalkül in der Walsh-Fourier-Analyse mit Anwendungen*. PhD thesis, Math.-Naturwiss. Fakultät, RWTH, 1972. publ. as Forschungsber. Landes Nordrhein-Westfalen, Nr 2334, Westdeutscher Verlag, Köln-Opladen, 71, 1973.
- [76] H.J. Wagner. On dyadic calculus for functions defined on R_+ . In *Proc. Sympos. Theory & Applic. Walsh & Other Non-sinus. Functions*, pages ii+27, RWTH Aachen, 1974. Arbeitsber., Lehrstuhl A für Math. also in Hatfield, UK, 1975 July 1-3, 1975.
- [77] J.L. Walsh. A closed set of orthogonal functions. *Amer. J. Math.*, 55:5–24, 1923.
- [78] Zheng Wei-xing. The generalized Walsh transform and an extremum problem. *Acta Math. Sinica*, 22(3):362–374, 1979. (in Chinese, English summary).
- [79] Zheng Wei-xing and Su Wei-yi. The logical derivatives and integrals. *J. Math. Res. Exposition*, 1:79–90, 1981. (in Chinese, English summary).
- [80] Su Wei-yi. Pseudo-differential operators and derivatives on locally compact vilenkin groups. *Science in China, Ser. A*, 35(7):826–836, 1992.
- [81] Su Wei-yi. Operators-derivatives-spaces-differential equations on locally compact Vilenkin groups. In M. Cheng, D.G. Deng, S. Gong, and C.C. Yang, editors, *Harmonic Analysis in China*, pages 240–255. Kluwer Academic Publisher, Hong Kong, 1995.
- [82] F. Weisz. *Symmability of Multi-Dimensional Fourier Series and Hardy Spaces*, volume 541 of *Mathematic and Its Applications*. Kluwer Academic Publishers, 2002.
- [83] F. Weisz. Summation of Fourier series. *Acta Mathematica Academiae Paedagogicae Nyiregyháziensis*, 20:239–266, 2004.

- [84] F. Weisz. Summability of Walsh-Fourier series and the dyadic derivative. In R.S. Stanković, editor, *Walsh and Dyadic Analysis*, pages 109–135. Elektronski fakultet, Niš, Serbia, 2008.
- [85] Su Weiyi. Gibbs-Butzer differential operators on locally compact Vilenkin groups. *Science in China, Ser. A*, 39(7):718–727, 1996.
- [86] Su Weiyi. Gibbs derivatives and differential equations on Vilenkin groups. In R.S. Stanković, M. Stojić, and M.S. Stanković, editors, *Recent Developments in Abstract Harmonic Analysis with Applications in Signal Processing*, pages 79–94. Nauka and Elektronski fakultet, Niš, Serbia, 1996.
- [87] Su Weiyi. Calculus on fractals based upon local fields. *Approx. Theory and Its Applications*, 16(1):92–100, 2000.
- [88] Su Weiyi and G.X. Chen. Lipschitz classes on local fields. *Science in China, Ser. A, Math.*, 50(7):1005–1014, 2007.
- [89] Su Weiyi and Xu Qiang. Function spaces on local fields. *Science in China*, 49(1):66–74, 2006.