

Editorial

'Aqualisation' of neuraxis: Wondrous neuraqua CSF 1

Manu Kothari, Atul Goel

1

View and Review

Organization of neurology services in India: Unmet needs and the way forward

Mandaville Gourie-Devi

4

Original Articles

Endoscopic management of brain abscesses

Yad Ram Yadav, Mallika Sinha, Neha, Vijay Parihar

13

Pattern of cerebellar perfusion on single photon emission computed tomography in subcortical hematoma: A clinical and computed tomography correlation

Jayantee Kalita, Usha K. Misra, Prasen Ranjan, P. K. Pradhan

17

Imaging features in Hirayama disease

Hemant A. Sonwalkar, Rakesh S. Shah, Firosh K. Khan, Arun K. Gupta, Narendra K. Bodhey, Surjith Vottath, Sukalyan Purkayastha

22

Delayed habituation in Behcet's disease

Sefa Gulturk, Melih Akyol, Hulusi Kececi, Sedat Ozelik, Ziyet Cinar, Ayse Demirkazik

27

Erythrocyte indicators of oxidative changes in patients with graded traumatic head injury

Chandrika D. Nayak, Dinesh M. Nayak, Annaswamy Raja, Anjali Rao

31

Repeat gamma knife radiosurgery for recurrent or refractory trigeminal neuralgia

Liang Wang, Zhen-wei Zhao, Huai-zhou Qin, Wen-tao Li, Hua Zhang, Jian-hai Zong, Jian-Ping Deng, Guo-dong Gao

36

Taste dysfunction in vestibular schwannomas

Rabi Narayan Sahu, Sanjay Behari, Vimal K. Agarwal, Pramod J. Giri, Vijendra K. Jain

42

Surgical management of traumatic intracranial pseudoaneurysms: A report of 12 cases

Xiang Wang, Jin-Xiu Chen, Chao You, Min He

47

Expression of truncated dystrophin cDNAs mediated by a lentiviral vector

Sun Shunchang, Chen Haitao, Chen Weidong, He Jingbo, Peng Yunsheng

52

Gamma knife radiosurgery for glomus jugulare tumors: Therapeutic advantages of minimalism in the skull base

Manish S. Sharma, A. Gupta, S. S. Kale, D. Agrawal, A. K. Mahapatra and B. S. Sharma

57

Case Reports

Subarachnoid hemosiderin deposition after subarachnoid hemorrhage on T2*-weighted MRI correlates with the location of disturbed cerebrospinal fluid flow on computed tomography cisternography <i>Yoshifumi Horita, Toshio Imaizumi, Yuji Hashimoto, Jun Niwa</i>	62
Anesthesia management of awake craniotomy performed under asleep-awake-asleep technique using laryngeal mask airway: Report of two cases <i>Gadhinglajkar Shrinivas Vitthal, Rupa Sreedhar, Mathew Abraham</i>	65
High cervical C3-4 'disc' compression associated with basilar invagination <i>Atul Goel</i>	68
Short-lasting unilateral neuralgiform headache with conjunctival injection and tearing: Response to antiepileptic dual therapy <i>Ravi Gupta, Manjeet S. Bhatia</i>	71
Correlation of autism with temporal tubers in tuberous sclerosis complex <i>Kavitha Kothur, Munni Ray, Prahbjot Malhi</i>	74
Non-traumatic carotid dissection and stroke associated with anti-phospholipid antibody syndrome: Report of a case and review of the literature <i>Benzi M. Kluger, Richard L. Hughes, C. Alan Anderson, Kathryn L. Hassell</i>	77
Osteoma of anterior cranial fossa complicated by intracranial mucocele with emphasis on its radiological diagnosis <i>Jinhu Ye, Hui Sun, Xin Li, Jianping Dai</i>	79
Vasospasm after transsphenoidal pituitary surgery: A case report and review of the literature <i>Manish Kumar Kasliwal, Ravinder Srivastava, Sumit Sinha, Shashank S. Kale, Bhawani S. Sharma</i>	81
Chondromyxoid fibroma of the seventh cervical vertebra <i>Ashish Jonathan, Vedantam Rajshekhar, Geeta Chacko</i>	84
Acute progressive midbrain hemorrhage after topical ocular cyclopentolate administration <i>Tarkan Calisaneller, Ozgur Ozdemir, Erkin Sonmez, Nur Altinars</i>	88

Letters to Editor

Digital subtraction angiography laboratory with inbuilt CT (DynaCT): Application during intracranial aneurysm embolization	90
Concomitant tuberculous and pyogenic cerebellar abscess in a patient with pulmonary tuberculosis	91
Drug compliance after stroke and myocardial infarction: Is complementary medicine an issue?	93

Multiple intracranial developmental venous anomalies associated with complex orbitofacial vascular malformation	93
Nitrofurantoin-induced peripheral neuropathy:A lesson to be re-learnt	94
Posterior longitudinal ligament cyst as a rare cause of lumbosacral radiculopathy with positive straight leg raising test	96
Aqueductal stenosis caused by an atypical course of a deep collector vein draining bilateral cerebellar developmental venous anomalies	97
Recovery of increased signal intensity of the cervical cord on magnetic resonance imaging after surgery for spontaneous spinal epidural hematoma causing hemiparesis	98
Simultaneous thalamic and cerebellar hypertensive hemorrhages	100

Neuroimages

MRI and MRA in spontaneous intracranial arterial dissection <i>S. Raghavendra, Sanjeev V. Thomas, Krishnamoorthy Thamburaj, Bejoy Thomas</i>	102
Shunt catheter migration into pulmonary arteries <i>Miikka Korja, Matti K. Karvonen, Arto Haapanen, Reijo J. Marttila</i>	103
Susceptibility weighted imaging in holohemispheric venous angioma with cerebral hemiatrophy <i>Sivaraman Somasundaram, Chandrasekharan Kesavadas, Bejoy Thomas</i>	104

Forthcoming Events	105
--------------------------	-----

Instructions to Authors	106
-------------------------------	-----

Referees List - 2007	000???
----------------------------	--------

The copies of the journal to members of the association are sent by ordinary post. The editorial board, association or publisher will not be responsible for non-receipt of copies. If any of the members wish to receive the copies by registered post or courier, kindly contact the journal's / publisher's office. If a copy returns due to incomplete, incorrect or changed address of a member on two consecutive occasions, the names of such members will be deleted from the mailing list of the journal. Providing complete, correct and up-to-date address is the responsibility of the members. Copies are sent to subscribers and members directly from the publisher's address; it is illegal to acquire copies from any other source. If a copy is received for personal use as a member of the association/society, one cannot resale or give-away the copy for commercial or library use.

Repeat gamma knife radiosurgery for recurrent or refractory trigeminal neuralgia

Liang Wang, Zhen-wei Zhao, Huai-zhou Qin, Wen-tao Li¹, Hua Zhang, Jian-hai Zong, Jian-Ping Deng, Guo-dong Gao

Department of Neurosurgery, Tangdu Hospital, Institute for Functional Brain Disorders, Fourth Military Medical University, ¹Neurosurgery, First Hospital of Xi'an Jiaotong University, Xi'an China

Background: Repeat gamma knife radiosurgery (GKRS) is considered to be an effective treatment for refractory or recurrent trigeminal neuralgia (TN). **Aims:** The purpose of this report was to demonstrate the relationship between the outcome of repeat GKRS and prior operative procedures on patients with recurrent or refractory TN. **Materials and Methods:** A retrospective analysis was performed on 34 patients with refractory or recurrent TN who had undergone repeat GKRS; 21 patients had undergone other types of procedures, 11 of whom had undergone more than three such procedures prior to radiosurgery. The maximum dose of the repeat procedure was between 60 and 75 Gy. The mean follow-up time was 21.6 months. **Statistical Analysis Used:** The log-rank test and Fisher's exact test were used to analyze the data. **Results:** Excellent pain relief was achieved in 14 patients (41.2%) after repeat GKRS, while a successful outcome occurred in 29 of 34 patients (85.3%). Better pain relief occurred in the patients who did not have a prior procedure or who had undergone fewer than three prior procedures ($P = 0.042$). Twenty-four of 25 patients (96.0%) who had recurrent pain had a successful operation and five of nine patients (55.6%) who did not have significant relief of pain after the first procedure had a successful operation. The difference was statistically significant ($P < 0.01$). Only four patients had mild complications. **Conclusion:** It is more likely to relieve pain in patients with recurrent or refractory TN who did not have a prior procedure or who had fewer than three procedures before undergoing their first GKRS. Moreover, it seems that patients who had a good response following the initial GKRS had better results after a repeat procedure.

Key words: Gamma knife surgery, pain, radiosurgery, trigeminal neuralgia

Since Leksell^[1] performed gamma knife radiosurgery (GKRS) on 63 patients with trigeminal neuralgia (TN), GKRS is considered to be an effective therapy and a minimally invasive surgical approach for medically

or surgically intractable TN, especially for the patient who has failed other surgical procedures.^[2-8] Although the reports^[3,9-15] show that there is a high rate of pain relief (77-95%) after the first GKRS, there is a definite percentage of patients who require repeat GKRS and who experience persistent or recurrent pain after the first procedure. In this paper, we retrospectively analyzed the clinical data of 34 patients who underwent repeat GKRS with persistent, particularly severe, medically or surgically refractory or recurrent TN. We divided the patient cohort according to whether they had previously undergone other operative procedures before radiosurgery to ascertain the treatment effect.

Materials and Methods

Patient population

Between June 1999 and December 2005, a total of 322 patients with medically unresponsive TN underwent GKRS at the Tangdu Hospital of the Fourth Military Medical University and the First Hospital of Xi'an Jiaotong University. Thirty-eight patients required further intervention for delayed, recurrent symptoms or inadequate pain relief and 34 patients accepted repeat GKRS, all of whom had failed medical management before GKRS. Of these patients, 21 (61.8%) had undergone previous surgical interventions and 11 patients (32.3%) had undergone more than three procedures. The remaining 13 patients selected GKRS as their first and second surgical procedures. Two patients (5.9%) had previously been diagnosed with multiple sclerosis (MS). We excluded patients who achieved good pain relief by microvascular decompression (MVD) after initial GKRS. Tumor-related TN was excluded from this analysis.

The patient characteristics are summarized in Table 1. The male to female ratio was 14:20. The median age at the time of the first GKRS of the retreated patients was

Guo-dong Gao

Department of Neurosurgery, Tangdu Hospital, Institute for Functional Brain Disorders, Fourth Military Medical University, 1 Xinsi Road, Xi'an - 710038 China.
E-mail: gguodong@fmmu.edu.cn

Table 1: Patient characteristics

Characteristics	Number
Age at first GKRS (median range years)	68 (42-87)
First onset age (median range years)	55 (36-71)
Male/female	14/20
Right/left	22/12
Pain duration prior to initial GKRS (median range months)	63 (5-169)
Interval between initial and repeat GKRS (mean range months)	17.4 (8-34)
Follow-up time after repeat GKRS (mean range months)	21.6 (7-41)
Number of procedures prior to GKRS	
0	13
1-2	10
>3	11
Multiple sclerosis	2
Recurrent/refractory	25/9
Maximum dose of repeat GKRS	
60-70 Gy	13
70-75 Gy	21

GKRS - Gamma knife radiosurgery

68 years (range, 42-87 years). The median age of the initial symptom onset was 55 years (range, 36-71 years). The median pain duration prior to initial GKRS was 63 months (range, 5-169 months).

Radiosurgical techniques

All patients were treated with a ^{60}Co gamma unit by a team consisting of a radiation neurosurgeon and a medical engineer. Under local anesthesia, a stereotactic coordinate frame was attached to each patient's head. Then, MRI (Philip 1.5-tesla magnetic resonance scanner) scans were acquired by use of axial T1-weighted sequence and 0.8-mm slice thickness within the frame to identify the TN root entry zone. The MRI data were entered into the treatment planning computer, through which the conformal radiation plan was produced.

Using a 4-mm collimator, a single isocenter or two isocenters were positioned 3-4 mm anterior to the region where the sensory root of the TN exits the pons. The number of isocenters selected was related to the length of the TN root entry zone for reducing the influencing region. Five patients were treated with two isocenters. The isocenter was placed so that the 60% isodose line covered the entire target segment and there was no greater than 20 Gy radiations at the brainstem. At the first radiosurgical procedure, the mean dose was 73.4 Gy (range, 60-80 Gy) to the cisternal TN. At the second radiosurgical procedure, the patients were divided into two groups. In the first group of 13 patients, we selected a mean dose of 66.2 Gy (range, 60-70 Gy) to decrease the complications of the operation and the isocenter was the same as in the first procedure [Figure 1-left]. Later, we placed the isocenter 1-2 mm anterior to the first isocenter as Hasegawa *et al.*^[12] reported [Figure 1-right]. In this way, the first target and the additional length of the nerve received adequate additional irradiation, while




Figure 1: Axial T1-weighted MRI scan demonstrated the designated target of the second group (right, two isocenters, dose 70-75 Gy) was anterior to the first group (left, one isocenter, dose 60-70 Gy). The red ring represents the designing target. The yellow cross represents the target isocenter. The margin dose on the surface of the pons was less than 20 Gy

the brainstem received less irradiation. In the second group, the mean dose was 73.1 Gy (range, 70-75 Gy).

Follow-up and statistical analysis

Patients were discharged within 24 h after radiosurgery. All patients were subsequently evaluated by physicians who did not participate in the treatment via telephone or questionnaires six months after GKRS and then yearly. The mean interval of time between the initial and repeat GKRS was 17.4 months (range, 8-34 months) and the follow-up time was 21.6 months (range, 7-41 months) after the second radiosurgical procedure.

Patients were studied according to their degree of pain relief, use of medications and sensory disturbances, including corneal anesthesia and ocular complications. The Barrow Neurological Institute (BNI) scoring system was used to label the degree of pain relief as follows: Grade I, no pain and no medication required; Grade II, occasional pain and no medication required; Grade IIIa, no pain and continued use of medications required; Grade IIIb, some pain which was adequately controlled on medication; Grade IV, pain improved but not adequately controlled on medication; and Grade V, no pain relief whatsoever. If pain returned to the level of the poor category (i.e., BNI IV or V) after an initial significant improvement, we considered it to be recurrent pain. And if pain keeps at the level of the poor category (BNI IV or V) after surgical improvement, we considered it as refractory TN. Patients who achieved complete or partial pain relief, but continued to take some medications for fear of recurrent pain or achieved mild pain control (i.e., better than BNI IIIb) after radiosurgery were categorized as having a successful operation and the patients who obtained the best pain relief (i.e., BNI I) were categorized as having an excellent outcome.

The toxicity grade was rated on a four-point scale: Grade I, no facial numbness and no related neurological symptoms (including headache, vertigo, vomiting and other cranial nerve deficits) and symptomatic radiological changes; Grade II, mild facial numbness

but not bothersome and/or mild neurological symptoms (including headache, vertigo, vomiting and other cranial nerve deficits) and symptomatic radiological changes which could be controlled by some outpatient medications; Grade III, facial numbness that was somewhat bothersome and/or moderate neurological symptoms (including headache, vertigo, vomiting and other cranial nerve deficits) and symptomatic radiological changes which could be alleviated by some outpatient medications; and Grade IV, facial numbness that was very bothersome and/or severe neurological symptoms and symptomatic radiological changes which required surgical intervention. The medications used or the surgical procedures after repeat GKRS for each patient were recorded in the computer for analysis.

Statistical analysis was performed using the Statistical Package for Social Sciences (SPSS), Version 11.5. Fisher's exact test, the log-rank test and the χ^2 -test were used to analyze the data. All tests were two-tailed and a probability value <0.05 was considered significant.

Results

The outcome of patients with TN undergoing repeat GKRS is summarized in Table 2. Pain relief occurred in nearly all 34 patients after repeat GKRS. The median time of pain relief was eight weeks (range, 1-17 weeks). Fourteen patients (41.2%) were completely pain-free (BNI Grade I), four patients (11.8%) stopped all medications, but occasionally experienced slight pain (BNI Grade II), nine patients (26.5%) were pain-free, but still used medication for fear of pain recurrence (BNI Grade IIIa) and two patients (5.9%) continued to rely on medications to control the syndrome adequately (BNI Grade IIIb), yet the dose of medicine was less than before. A successful operation occurred in 29 of the 34 patients (85.3%) after repeat GKRS; the remaining four patients (11.8%), including two patients with

recurrent symptoms three and seven months after the repeat procedure, experienced some symptomatic improvement, but still needed additional treatment (BNI Grade IV). One patient (2.9%) had no pain relief and had the same symptoms as before radiosurgery (BNI Grade V). Three of those patients who achieved poor results underwent MVD and partial rhizotomy after the repeat procedure. One patient had good pain relief (BNI Grade II) and the other two had a fair response, but needed some medications (BNI Grade IIIb). One patient who underwent repeat MVD, two separate glycerol rhizotomies and one radiofrequency rhizotomy after the repeat GKRS, achieved a little pain relief, but not completely after these procedures and endured permanent facial numbness (Grade III). One patient, who had no pain relief after the repeat procedure, refused any additional treatments and was lost to follow-up after 10 months. Two MS-TN patients had pain relief after repeat GKRS (BNI Grade IIIb) and required decreased doses of medication for pain control.

Among the 13 patients with no previous procedures before the initial GKRS, 12 patients (92.3%) achieved successful outcomes, including seven patients (53.9%) with excellent pain relief after repeat radiosurgery. By way of comparison, of the 21 patients with one or more prior surgical procedures before the repeat GKRS, 17 patients (81.0%) achieved successful outcomes, including seven patients (33.3%) with excellent pain relief. However, the difference between the two groups was not statistically significant ($P = 0.377$, log-rank test, $\nu = 1$). Compared with the 11 patients with more than three prior surgical procedures before the repeat GKRS, of which seven patients achieved successful outcomes, the difference was statistically significant ($P = 0.042$, log-rank test, $\nu = 2$).

Twenty-four of 25 patients (96.0%) who had recurrent pain after the first procedure achieved a successful operation and five of the nine patients (55.6%) who had no significant relief after the first procedure achieved a successful operation. The difference was statistically significant ($P = 0.006$, Fisher's exact test; $P = 0.003$, log-rank test, $\nu = 1$). Two patients with no pain relief after the initial GKRS reported unsuccessful outcomes after repeat GKRS as well. The outcome of patients who had 60-70 Gy radiation in the second treatment had no difference in outcome from those who had 70-75 Gy radiation ($P = 0.17$, $\nu = 1$).

Only four patients (11.8%) developed facial numbness after a repeat procedure. No other neurological symptoms and symptomatic radiological changes appeared. One patient reported continued trigeminal numbness after the initial GKRS, which did not worsen after the repeat procedure and was described as somewhat bothersome (Grade III). The remaining three patients with new facial numbness were described as mild, but not bothersome (Grade II) and the symptoms gradually resolved

Table 2: The outcome of patients with trigeminal neuralgia who underwent repeat gamma knife radiosurgery

Outcome after repeat GKRS	Recurrent/ refractory		No. of procedure		
	Recurrent	Refractory	0	1-2	>3
Grade I	13	1	7	4	3
Grade II	4	0	1	2	1
Grade IIIa	4	3	4	1	2
Grade IIIb	3	1	0	3	1
Grade IV	1	3	1	0	3
Grade V	0	1	0	0	1
Excellent	13	1	7	4	3
Success	24	5	12	10	7
Poor	1	4	1	0	4
Total	25	9	13	10	11

The outcome of the operation was evaluated by BNI scoring system. Pain returning to the poor level (BNI IV or V) was considered as recurrent pain. And if pain keeps at the level of the poor category (BNI IV or V) after surgical improvement, we considered it as refractory TN. Patients who achieved outcome better than BNI IIIb were categorized as success

spontaneously in one to six months. Moreover, one of the patients with mild facial numbness had experienced similar symptoms after the first GKRS, but it was not bothersome and gradually resolved spontaneously in three months. No severe complications (i.e. more severe than Grade III) were reported.

Discussion

In this paper, we describe a retrospective analysis of clinical data from 34 patients who had refractory or recurrent TN and underwent repeat GKRS. The results indicated that the more surgical procedures the patients underwent prior to repeat GKRS, the worse the clinical outcome. The patients who had a good response following initial GKRS may therefore achieve a better result after a repeat procedure.

Being a newer treatment alternative since Leksell's^[1] report, GKRS has more than an 80% probability of significant pain relief with a 10% or less risk of subsequent mild facial numbness or dysesthesia.^[2,5,10,14-25] Thus, radiosurgery has gradually been regarded as a more efficacious procedure than MVD, radiofrequency rhizotomy, glycerol rhizotomy, balloon compression or radiofrequency ablation as an initial treatment for idiopathic trigeminal neuralgia.^[14,26-34] Nevertheless, recurrent or refractory pain inevitably occurs. Thus, choosing the optimal modality after treatment failure continues to pose a dilemma for patients and physicians.

Hasegawa *et al.*^[12] described 23 of 27 patients who achieved a successful outcome after repeat GKRS. Two of four patients who had recurrent pain obtained relief following additional procedures. The results were similar to the results of Pollock *et al.*^[35-36] in which 14 of 19 patients with persistent TN continued to have an excellent outcome after repeat GKRS. Two patients had recurrent pain seven and 22 months after a repeat procedure. Both the patients had undergone more than five procedures, including GKRS, before or after repeat GKRS. Moreover, the results reported by the other authors are remarkably similar, as summarized in Table 3. These results demonstrated that repeat GKRS may be useful for some patients who have recurrent or refractory pain.^[12,35-39]

According to the previous study, there may be some relationship between the outcome of repeat radiosurgery

and the number of times the procedure was performed. In this study, we used 60-75 Gy for repeat GKRS. The difference between the successful patients (92.3%) who had no prior procedures before the initial GKRS and the successful patients (80.1%) who underwent more than one procedure before the initial GKRS was not significant. However, when compared with the successful patients (63.6%) who underwent more than three procedures before repeat GKRS, the difference was statistically significant ($P = 0.042$). Because the initial GKRS could be considered as a prior procedure to the repeat procedure, the latter result carries more weight. These patients were similar to Pollock *et al.*'s cohort;^[40] excellent outcomes were achieved after radiosurgery in 67% of the patients who did not undergo previous surgery, compared with 51% of patients who underwent prior surgery. It was demonstrated that patients who had not undergone surgery previously had a significantly better outcome than patients who had undergone prior operations before the repeat GKRS for idiopathic trigeminal neuralgia.

In the present study, success occurred in 24 of 25 patients (96%) who had recurrence after the first GKRS, compared with five of nine patients (55.6%) who were refractory after the initial treatment. Both the Fisher exact and log-rank tests were statistically significant ($P < 0.01$), suggesting that the patients who achieved good responses following the first procedure received a better result than those who were refractory for the first procedure after repeat treatment, but some controversy persists. Herman *et al.*^[38] reported that among the 15 patients (83%) with recurrent pain after initially successful GKRS, 14 patients (93%) achieved excellent or good outcomes after repeat radiosurgery; all three patients with poor responses after initial GKRS reported poor outcomes after repeat GKRS. However, Brisman^[37] reported that the difference was not statistically significant. In his cohort, a successful second GKRS occurred in 21 of 29 patients who had excellent or fair pain relief following the first GKRS and in seven of 15 patients who had a poor result following the first GKRS. He suggested the result may indicate that patients who have better pain relief initially after the first procedure have no or fewer prior procedures before radiosurgery. In addition, Pollock *et al.*^[36] found that there is no relationship between excellent outcomes

Table 3: Series reports of repeat gamma knife radiosurgery for trigeminal neuralgia

Series	No.	Repeat dose (Gy)	Follow-up (mon)	Excellent (%)	Success (%)	Toxicity (%)	Prior surgery (%)
Hasegawa ^[12]	27	64.4	20.4	19	85	11	82
Pollock ^[36]	19	76	24	74	95	58	/
Brisman ^[37]	44	40	15	22	62	13	52
Herman ^[38]	18	70	24.5	45	78	11	28
Shetter ^[39]	19	46.6	13.5	53	85	42	/
Present	34	70	21.6	41	85	15	62

No. - Number. Patients who achieved outcome better than BNI IIIb after radiosurgery were categorized as having a successful operation and the patients who obtained the best pain relief (BNI I) were categorized as having excellent outcome

after repeat radiosurgery and excellent outcomes after the initial procedure; however, in their data, there were only two patients who had persistent facial pain after the first procedure and the different conclusion from Brisman's report^[37] may result from the different radiation dose (40 Gy vs. 60-75 Gy in the current report) delivered to the patients for repeat GKRS.

The radiation dose of the repeat GKRS treatment in the available clinic reports varies from 35 to 90 Gy.^[1,5,12,13,35-39] According to the previous studies, pain relief and the incidence of facial numbness after GKRS correlate with the dose of treatment. The higher the treatment dose delivered, the better the pain relief achieved and unfortunately the higher the associated toxicity. Pollock *et al.*^[35] indicated that patients treated with 70 Gy required additional surgery (37%), which is more often than patients treated with >70 Gy (13%); however, the incidence of new facial numbness or paresthesias was 8% compared with 32%. The result was similar to the other reports.^[23,36,37] On the other hand, the patients who underwent multiple surgical prior procedures would have obtained a worse pain outcome and enhanced the difficulty of treatment. These might not be all related to the dose of the radiation of the first and repeat GKRS procedures.

Finally, it is worth pointing out that although similar reliable statistical conclusions have been achieved in the present study and in other previous reported results, the investigation on the GKRS treatment for refractory and recurrent TN is not over, especially considering that the numbers of the cohorts reported so far are not big enough. Extended research with large sample size is necessary to further clarify the key factors affecting the outcomes of repeated GKRS and optimize the treatment procedure.

Conclusion

Repeat GKRS for TN has been shown to be an efficient noninvasive surgical procedure with a low complication rate and is more likely to relieve pain in those who did not have a prior procedure or who had fewer than three procedures prior to their first GKRS. The dose selection is important such that the higher the dose used in treatment, the better the result can be achieved and the higher the incidence of facial numbness one may endure. Moreover, it seems that patients who had a good response following initial GKRS would achieve better results after a repeat procedure.

References

- Leksell L. Stereotactic radiosurgery. *J Neurol Neurosurg Psychiatry* 1983;46:797-803.
- Brisman R. Microvascular decompression vs. gamma knife radiosurgery for typical trigeminal neuralgia: Preliminary findings. *Stereotact Funct Neurosurg* 2007;85:94-8.
- Kondziolka D, Lunsford LD, Flickinger JC, Young RF, Vermeulen S, Duma CM, *et al.* Stereotactic radiosurgery for trigeminal neuralgia: A multiinstitutional study using the gamma unit. *J Neurosurg* 1996;84:940-5.
- Maesawa S, Salame C, Flickinger JC, Pirris S, Kondziolka D, Lunsford LD. Clinical outcome after stereotactic radiosurgery for idiopathic trigeminal neuralgia. *J Neurosurg* 2001;94:14-20.
- Massager N, Lorenzoni J, Devriendt D, Levivier M. Radiosurgery for trigeminal neuralgia. *Prog Neurol Surg* 2007;20:235-43.
- Regis J, Bartolomei F, Metellus P, Rey M, Genton P, Dravet C, *et al.* Radiosurgery for trigeminal neuralgia and epilepsy. *Neurosurg Clin North Am* 1999;10:359-77.
- Rogers CL, Shetter AG, Fiedler JA, Smith KA, Han PP, Speiser BL. Gamma knife radiosurgery for trigeminal neuralgia: The initial experience of the Barrow Neurological Institute. *Int J Radiat Oncol Biol Phys* 2000;47:1013-9.
- Young RF, Vermeulen S, Posewitz A. Gamma knife radiosurgery for the treatment of trigeminal neuralgia. *Stereotact Funct Neurosurg* 1998;70:192-9.
- Huang E, Teh BS, Zeck O, Woo SY, Lu HH, Chiu JK, *et al.* Gamma knife radiosurgery for treatment of trigeminal neuralgia in multiple sclerosispatients. *Stereotact Funct Neurosurg* 2002;79:44-50.
- Petit J, Herman J, Nagda S, DiBiase SJ, Chin LS. Radiosurgical treatment of trigeminal neuralgia: Evaluating quality of life and treatment-outcomes. *Int J Radiat Oncol Biol Phys* 2003;56:1147-53.
- Regis J, Metellus P, Dufour H, Roche PH, Muracciole X, Pellet W, *et al.* Long-term outcome after gamma knife surgery for secondary trigeminal neuralgia. *J Neurosurg* 2001;95:199-205.
- Hasegawa T, Kondziolka D, Spiro R, Flickinger JC, Lunsford LD. Repeat radiosurgery for refractory trigeminal neuralgia. *Neurosurgery* 2002;50:494-502.
- Massager N, Murata N, Tamura M, Devriendt D, Levivier M, Régis J. Influence of nerve radiation dose in the incidence of trigeminal dysfunction after trigeminal neuralgia radiosurgery. *Neurosurgery* 2007;60:681-8.
- Brisman R. Microvascular decompression vs. gamma knife radiosurgery for typical trigeminal neuralgia: Preliminary findings. *Stereotact Funct Neurosurg* 2007;85:94-8.
- Fountas KN, Lee GP, Smith JR. Outcome of patients undergoing gamma knife stereotactic radiosurgery for medically refractory idiopathic trigeminal neuralgia: Medical College of Georgia's experience. *Stereotact Funct Neurosurg* 2006;84:88-96.
- Brisman R. Gamma knife radiosurgery for primary management for trigeminal neuralgia. *J Neurosurg* 2000;93:159-61.
- Brisman R. Gamma knife surgery with a dose of 75 to 76.8 Gray for trigeminal neuralgia. *J Neurosurg* 2004;100:848-54.
- Chang JW, Chang JH, Park YG, Chung SS. Gamma knife radiosurgery for idiopathic and secondary trigeminal neuralgia. *J Neurosurg* 2000;93:147-51.
- Kondziolka D, Lunsford LD, Flickinger JC. Stereotactic radiosurgery for the treatment of trigeminal neuralgia. *Clin J Pain* 2002;18:42-7.
- McNatt SA, Yu C, Giannotta SL, Zee CS, Apuzzo ML, Petrovich Z. Gamma knife radiosurgery for trigeminal neuralgia. *Neurosurgery* 2005;56:1295-303.
- Nicol B, Regine WF, Courtney C, Meigooni A, Sanders M, Young B. Gamma knife radiosurgery using 90 Gy for trigeminal neuralgia. *J Neurosurg* 2000;93:152-4.
- Pollock BE, Phuon LK, Foote RL, Stafford SL, Gorman DA. High-dose trigeminal neuralgia radiosurgery associated with increased risk of trigeminal nerve dysfunction. *Neurosurgery* 2001;49:58-64.
- Tawk RG, Duffy-Fronckowiak M, Scott BE, Alberico RA, Diaz AZ, Podgorsak MB, *et al.* Stereotactic gamma knife surgery for trigeminal neuralgia: detailed analysis of treatment response. *J Neurosurg* 2005;102:442-9.
- Lopez BC, Hamlyn PJ, Zakrzewska JM. Systematic review of ablative neurosurgical techniques for the treatment of trigeminal neuralgia. *Neurosurgery* 2004;54:973-83.
- Massager N, Lorenzoni J, Devriendt D, Levivier M. Radiosurgery for trigeminal neuralgia. *Prog Neurol Surg* 2007;20:235-43.
- Barker FG 2nd, Jannetta PJ, Bissonette DJ, Larkins MV, Jho HD. The long-term outcome of microvascular decompression for trigeminal neuralgia. *N Engl J Med* 1996;334:1077-83.

27. Brown JA, McDaniel MD, Weaver MT. Percutaneous trigeminal nerve compression for treatment of trigeminal neuralgia: Results in 50 patients. *Neurosurgery* 1993;32:570-3.
28. Henson CF, Goldman HW, Rosenwasser RH, Downes MB, Bednarz G, Pequinot EC, *et al.* Glycerol rhizotomy versus gamma knife radiosurgery for the treatment of trigeminal neuralgia: An analysis of patients treated at one institution. *Int J Radiat Oncol Biol Phys* 2005;63:82-90.
29. Kondziolka D, Lunsford LD. Percutaneous retrogasserian glycerol rhizotomy for trigeminal neuralgia: Technique and expectations. *Neurosurg Focus* 2005;18:E7.
30. Taha JM, Tew JM Jr, Buncher CR. A prospective 15-year followup of 154 consecutive patients with trigeminal neuralgia treated by percutaneous stereotactic radiofrequency rhizotomy. *J Neurosurg* 1995;83:989-93.
31. Young RF. Glycerol rhizolysis for treatment of trigeminal neuralgia. *J Neurosurg* 1988;69:39-45.
32. Sanchez-Mejia RO, Limbo M, Cheng JS, Camara J, Ward MM, Barbaro NM. Recurrent or refractory trigeminal neuralgia after microvascular decompression, radiofrequency ablation or radiosurgery. *Neurosurg Focus* 2005;18:e12.
33. Ramnarayan R, Mackenzie I. Brain-stem auditory evoked responses during microvascular decompression for trigeminal neuralgia: Predicting post-operative hearing loss. *Neurol India* 2006;54:250-4.
34. Natarajan M. Percutaneous trigeminal ganglion balloon compression: Experience in 40 patients. *Neurol India* 2000;48:330-2.
35. Pollock BE, Foote RL, Stafford SL, Link MJ, Gorman DA, Schomberg PJ. Results of repeated gamma knife radiosurgery for medically unresponsive trigeminal neuralgia. *J Neurosurg* 2000;93:162-4.
36. Pollock BE, Foote RL, Link MJ, Link MJ, Gorman DA, Schomberg PJ. Repeat radiosurgery for idiopathic trigeminal neuralgia. *Int J Radiat Oncol Biol Phys* 2005;61:192-5.
37. Brisman R. Repeat gamma knife radiosurgery for trigeminal neuralgia. *Stereotact Funct Neurosurg* 2003;81:43-9.
38. Herman JM, Petit JH, Amin P, Kwok Y, Dutta PR, Chin LS. Repeat gamma knife radiosurgery for refractory or recurrent trigeminal neuralgia: treatment outcomes and quality-of-life assessment. *Int J Radiat Oncol Biol Phys* 2004;59:112-6.
39. Shetter AG, Rogers CL, Ponce F, Fiedler JA, Smith K, Speiser BL. Gamma knife radiosurgery for recurrent trigeminal neuralgia. *J Neurosurg* 2002;97:536-8.
40. Pollock BE, Phuong LK, Gorman DA, Foote RL, Stafford SL. Stereotactic radiosurgery for idiopathic trigeminal neuralgia. *J Neurosurg* 2002;97:347-53.

Accepted on 16-08-2007

Source of Support: Nil, **Conflict of Interest:** None declared.