

Title	Representing elements of stable homotopy groups by symmetric maps
Author(s)	Mimura, Mamoru; Mukai, Juno; Nishida, Goro
Citation	Osaka Journal of Mathematics. 11(1) P.105-P.111
Issue Date	1974
Text Version	publisher
URL	<a href="https://doi.org/10.18910/11878">https://doi.org/10.18910/11878</a>
DOI	10.18910/11878
rights	
Note	

*Osaka University Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

Osaka University

## REPRESENTING ELEMENTS OF STABLE HOMOTOPY GROUPS BY SYMMETRIC MAPS

MAMORU MIMURA, JUNO MUKAI AND GORO NISHIDA

(Received June 14, 1973)

### 0. Introduction

Let  $S^m$  be the unit  $m$ -sphere. Let  $p$  be a prime and  $\pi$  the cyclic group of order  $p$ . Denote by  $B\pi^{(r)}$  the  $r$ -skeleton of the classifying space  $B\pi$ . Recall that  $B\pi$  is the infinite real projective space for  $p=2$  and the infinite lens space for  $p>2$ . Let  $X$  be a space. Let  $m$  be a positive integer for the case  $p=2$  and  $m$  an odd integer for the case  $p>2$ . Then a map  $f: S^m \rightarrow X$  is called *symmetric* if there exists a map  $\tilde{f}: B\pi^{(m)} \rightarrow X$  such that the following diagram is commutative:

$$(1) \quad \begin{array}{ccc} S^m & \xrightarrow{f} & X \\ \omega \searrow & & \nearrow \tilde{f} \\ & B\pi^{(m)} & \end{array}$$

, where  $\omega: S^m \rightarrow B\pi^{(m)}$  is the canonical projection.

An element of the homotopy group  $\pi_m(X)$  is called *symmetric* if it is represented by a symmetric map. For  $p=2$ , the definition of a symmetric map is due to J. H. C. Whitehead [14], in which he showed that if an essential element of  $\pi_m(S^{m-1})$  is symmetric, then  $m \equiv 3 \pmod{4}$ . Some results about the symmetry of the elements of  $\pi_m(X)$  are found in [4], [8], [10], [21] and [13].

Let  $X$  be an  $(l-1)$ -connected, finite CW-complex. Then our purpose is to show the following

**Theorem 1.** *Every element of  $\pi_m(X)$  is symmetric for any  $m$  satisfying  $2 \dim X - l < m < 2l - 2$  and*

- i)  $m \equiv -1 \pmod{2^{\phi(k+1)}} \quad \text{for } p=2,$
- ii)  $m \equiv -1 \pmod{2p^{\lfloor (k+1)/2 \rfloor}} \quad \text{for } p>2,$

where  $k=m-l$ ,  $\phi(s)$  is the number of integers  $i$  such that  $0 < i \leq s$  and  $i \equiv 0, 1, 2$  or  $4 \pmod{8}$  and  $\lfloor s \rfloor$  indicates the integer part of a rational  $s$ .

**Corollary 2.** *For an arbitrary  $k > 0$ , every element of the  $k$ -stem of the stable*

homotopy groups of spheres is symmetric.

To prove the above theorem we use the  $S$ -duality [11] and the Kahn-Priddy theorem [6] which is stated as follows for our use. Denote by  ${}^p\{X, Y\}$  the  $p$ -primary component of  $\{X, Y\} = \lim_{n \rightarrow \infty} [S^n X, S^n Y]$ .

**Theorem 3.** [Kahn-Priddy]. *Let  $N$  be a sufficiently large integer and  $h: S^N B\pi^{(s)} \rightarrow S^N$  a map such that the functional  $\mathfrak{P}^1(Sq^2)$ -operation is non-trivial (respectively). Then for a connected, finite CW-complex  $X$  of dimension  $< s$ ,  $h_*: \{X, B\pi^{(s)}\} \rightarrow {}^p\{X, S^0\}$  is an epimorphism. Furthermore, assume that the functional  $\mathfrak{P}^{[(s+1)/2(p-1)]}(Sq^{s+1})$ -operation of  $h$  is non-trivial for odd  $s$  (respectively), then  $h_*$  is an epimorphism for  $X$  of dimension  $\leq s$ .*

We express our thanks to H. Toda who suggested us to use the  $S$ -duality.

### 1. A proof of the Kahn-Priddy theorem

First we shall prove Theorem 3 for  $p=2$ . The notations of [6] are carried over to the present section unless otherwise stated.

Roughly speaking, the proof of Theorem 3 is to replace the infinite dimensional real projective space  $P^\infty$  with the  $s$ -dimensional one  $P^s$  and the map  $\phi: P^\infty \rightarrow (QS^0)_0$  with a map  $\text{adj}(h): P^s \rightarrow (QS^0)_0$  (cf. p. 985 of [6] and Theorem 7.3 of [9]) in the proof of Theorem 3.1 of [6].

$$\begin{aligned} \text{Let } t: B\mathcal{E}_{2^k}^{(s)} &\rightarrow \hat{Q}_m(B\mathcal{E}_{2^k}(2))^{(s)} \\ &= \hat{Q}_m(\underbrace{\hat{Q}_2 \cdots \hat{Q}_2}_{k-1} B\mathcal{E}_2)^{(s)} \subset \hat{Q}_m(\underbrace{\hat{Q}_2 \cdots \hat{Q}_2}_{k-1} P^s) \end{aligned}$$

be a restriction of the pretransfer  $T: B\mathcal{E}_{2^k} \rightarrow \hat{Q}_m(B\mathcal{E}_{2^k}(2))$  (Definition 3.1 of [6]) on the  $s$ -skeleton  $B\mathcal{E}_{2^k}^{(s)}$ . Let  $g_2': \hat{Q}_m(\underbrace{\hat{Q}_2 \cdots \hat{Q}_2}_{k-1} P^s) \rightarrow \hat{Q}_{m2^{k-1}}(P^s)$  be induced by the

wreath product and  $g_3': \hat{Q}_{m2^{k-1}}(P^s) \rightarrow Q(P^s)$  a Dyer-Lashof map. Then we obtain a commutative diagram

$$\begin{array}{ccc} \sum^\infty B\mathcal{E}_{2^k}^{(s)} & \xrightarrow{b} & \sum^\infty (QS^0)_0 \\ \downarrow a & & \downarrow r' \\ \sum^\infty P^s & \xrightarrow{h} & \sum^\infty S^0 \end{array}$$

, where  $a = \text{adj}(g_2' g_3' t)$ ,  $b$  is a restriction of  $G_\phi$  (p. 985 of [6]) on  $\sum^\infty \mathcal{E}_{2^k}^{(s)}$  and  $r'$  is defined by  $r'(x \wedge f) = f(x)$  for  $x \in \sum^\infty S^0$  and  $f \in (QS^0)_0$ . Remark that  $b$  is a restriction of  $\sum^\infty \bar{\phi} \circ g_3 g_2 f_1$  on  $\sum^\infty B\mathcal{E}_{2^k}^{(s)}$ .

For large  $k$ ,  $b_*: H_i(B\mathcal{E}_{2^k}^{(s)}; Z_2) \rightarrow H_i(Q(S^0)_0; Z_2)$  is an isomorphism if  $i < s$  (p. 985 of [6]). So, by the Whitehead-Serre theorem,  $b_*: {}^2\{X, B\mathcal{E}_{2^k}^{(s)}\} \rightarrow$

$\{X, (QS^0)_0\}$  is an isomorphism for a finite CW-complex  $X$  of dimension  $< s - 1$  and an epimorphism for  $X$  of dimension  $< s$ . It is clear that  $r_*': \{X, (QS^0)_0\} \rightarrow \{X, S^0\}$  is an epimorphism if  $X$  is connected. Thus  $(r'b)_*$  is an epimorphism on the 2-component and hence so is  $h_*$ . This proves the first part of Theorem 3 for  $p=2$ .

Under the first assumption of Theorem 3, the functional  $\mathfrak{B}^i(Sq^{2^i})$ - and  $\beta\mathfrak{B}^i(Sq^{2^i+1})$ - operations are non-trivial for  $2i(p-1) \leq s$  ( $2i \leq s$ , respectively). This is easily seen by use of the cohomology structure of  $B\pi^{(s)}$  and the Adem relation. So, by adding the second assumption,  $b_*: H_i(B\mathcal{C}_2^{(2)}; Z_2) \rightarrow H_i(Q(S^0)_0; Z_2)$  is an isomorphism for  $i < s$  and an epimorphism for  $i \leq s$ . This completes the proof of Theorem 3 for  $p=2$ .

For  $p > 2$ , the argument is quite parallel (cf. Remark 3.5 of [6] and Theorem 7.5 of [9]) and we omit it.

### 2. The S-duality

From now on we shall devote ourselves to the proof of Theorem 1. Denote by  $B\pi_s^r = B\pi^{(r)}/B\pi^{(s-1)}$ , where  $B\pi_0^r$  means  $B\pi^{(r)} \cup$  (one point). Let  $X$  be an  $(l-1)$ -connected, finite CW-complex of dimension  $j$ . Then  $f: S^m \rightarrow X$  is symmetric if and only if there is a map  $\tilde{f}: B\pi_n^m \rightarrow X$  for  $1 \leq n \leq l$  such that the following diagram is commutative:

$$(2) \quad \begin{array}{ccc} S^m & \xrightarrow{f} & X \\ \omega' \searrow & & \nearrow \tilde{f}' \\ & B\pi_n^m & \end{array}$$

, where  $\omega'$  is the map  $\omega$  of (1) followed by the collapsing map from  $B\pi^{(m)}$  to  $B\pi_n^m$ .

Let  $N$  be so large that  $N \geq \max(2j+1, 2m+1)$  and take  $N$ -duals of everything in (2):

$$(2') \quad \begin{array}{ccc} D_N S^m & \xleftarrow{\Delta_N f} & D_N X \\ \Delta_N \omega' \swarrow & & \nwarrow \Delta_N(\tilde{f}') \\ & D_N(B\pi_n^m) & \end{array}$$

, where  $D_N Y$  and  $\Delta_N g$  are  $N$ -duals of a finite CW-complex  $Y$  and a map  $g$  [11].

If  $m \leq 2n-2$ , then we work in the stable range. So, we obtain the following

**Proposition 4.** *Let  $X$  be an  $(l-1)$ -connected, finite CW-complex,  $N \geq \max(2j+1, 2m+1)$  and  $m \leq 2n-2$ . Then a map  $f: S^m \rightarrow X$  represents a symmetric element if and only if there is a map  $\tilde{f}: D_N X \rightarrow D_N(B\pi_n^m)$  for  $1 \leq n \leq l$  such that the following diagram is homotopy commutative:*

$$(3) \quad \begin{array}{ccc} & D_N f & \\ S^{N-m-1} \longleftarrow & & D_N X \\ D_N \omega' \swarrow & & \searrow \tilde{f} \\ & D_N(B\pi_n^m) & \end{array}$$

### 3. The S-dual of $B\pi_n^m$

Take  $N=N(a,s)=a2^{\phi(s)}$  for  $p=2$  and  $2ap^{\lfloor s/2(p-1) \rfloor}$  for  $p>2$ , where  $a$  is a sufficiently large integer.

Put  $s=m-n$ . Let  $\varepsilon=\varepsilon(s)=0$  if  $s\equiv -1 \pmod{2(p-1)}$  and  $\varepsilon=1$  if  $s\not\equiv -1 \pmod{2(p-1)}$  for  $p>2$  and  $\varepsilon=0$  for  $p=2$ . Then we have the following

**Proposition 5.**  $D_N(B\pi_n^m)$  has the same homotopy type as  $B\pi_{N-m-1}^{N-n-1}$  for  $N=N(a, s+\varepsilon)$  with  $s=m-n$ .

*Proof.* For  $p>2$ , recall from Theorem 1 of [7] that the stunted lens space  $B\pi_{2n}^{2m+1}=L^m(p)/L^{n-1}(p)$  is the Thom complex  $(L^s(p))^{\pi_1^*r(\xi)}$ , where  $L^r(p)=B\pi^{(2r+1)}$  is the  $(2r+1)$ -dimensional lens space,  $\xi$  is the canonical line bundle over the complex projective  $s$ -space  $CP^s$ ,  $r(\xi)$  is the real restriction of  $\xi$  and  $\pi_1^*r(\xi)=r\pi_1^*(\xi)$  is the bundle induced by the natural projection  $\pi_1:L^s(p)\rightarrow CP^s$ .

First we shall show

$$(4) \quad \begin{aligned} D_N(B\pi_{2n}^{2m+1}) &\simeq (L^s(p))^{(N/2-(m+1))\pi_1^*r(\xi)} \\ &\simeq B\pi_{N-2m-2}^{N-2n-1} \quad \text{for } N=N(a, 2s+1). \end{aligned}$$

According to Theorem 3.3 of [3], the S-dual of  $B\pi_{2n}^{2m+1}\simeq(L^s(p))^{\pi_1^*r(\xi)}$  is  $(L^s(p))^{-n\pi_1^*r(\xi)-\tau}$ , where  $\tau$  is the tangent bundle over  $L^s(p)$ . As is well known,  $\tau+1=(s+1)\pi_1^*r(\xi)$ . So  $(L^s(p))^{-n\pi_1^*r(\xi)-\tau}\simeq(L^s(p))^{1-(m+1)\pi_1^*r(\xi)}$ . By Theorem 2 of [7] the  $J$ -order of  $\pi_1^*r(\xi)-2$  is  $p^{\lfloor s/(p-1) \rfloor}$ . Obviously  $\lfloor s/p-1 \rfloor=\lfloor 2s+1/2(p-1) \rfloor$  holds. So by Theorem 3 of [7],  $(L^s(p))^{-(m+1)\pi_1^*r(\xi)}$  and  $(L^s(p))^{(N/2-(m+1))\pi_1^*r(\xi)}$  have the same stable homotopy type. Therefore we have obtained (4).

Observe that  $D_N(B\pi_{2n}^{2m})=D_N(B\pi_{2n}^{2m+1})/S^{N-2m-2}$  for  $N=N(a, 2s)$  and also that  $D_N(B\pi_{2n+1}^{2m+1})$  is obtained from  $D_N(B\pi_{2n}^{2m+1})$  by deleting the top dimensional cell for  $N=N(a, 2s)$ . By the same way as above we obtain  $D_N(B\pi_{2n+1}^{2m})$  from  $D_N(B\pi_{2n}^{2m})$  for  $N=N(a, 2s)$ .

Similarly and more simply we have the assertion for  $p=2$  (Theorem 6.1 of [3]). We note that the  $J$ -order of  $\xi-1$  is  $2^{\phi(s)}$ , where  $\xi$  is the canonical line bundle over the  $s$ -dimensional real projective space  $P^s$  ([1] and [2]).

### 4. On the Kahn-Priddy map

Consider the cofibring sequence

$$(5) \quad \dots \rightarrow S^m \xrightarrow{\omega'} B\pi_n^m \xrightarrow{i'} B\pi_n^{m+1} \xrightarrow{q'} S^{m+1} \rightarrow \dots,$$

where  $i'$  and  $q'$  are the canonical inclusion and projection respectively.

Put  $s=m-n$  and take  $N(a, s+2)$ -duals of everything in (5) and use Theorem 6.2 of [11] and Proposition 5, then we have the following

**Proposition 6.** *There is a cofibring sequence*

$$\dots \leftarrow S^{N-m-1} \xleftarrow{h} B\pi_{N-m-1}^{N-n-1} \xleftarrow{q} B\pi_{N-m-2}^{N-n-1} \xleftarrow{i} S^{N-m-2} \leftarrow \dots$$

for  $N=N(a, s+2)$ , where  $h: B\pi_{N-m-1}^{N-n-1} \simeq D_N(B\pi_n^m) \xrightarrow{D_N\omega'} S^{N-m-1}$ ,  $q=D_Ni'$  and  $i=D_Nq'$ .

We note that the cofibre of  $h$  is  $SB\pi_{N-m-2}^{N-n-1}$ .

**Proposition 7.** *Let  $m+1 \equiv 0 \pmod{2^{\phi(s+1)}}$  for  $p=2$  and  $m+1 \equiv 0 \pmod{2^{p^{\lceil (s+1)/2 \rceil}}}$  for  $p>2$ . Then*

i)  $B\pi_{N-m-1}^{N-n-1} \simeq S^{N-m-1}B\pi_0^s \simeq S^{N-m-1}B\pi^{(s)} \vee S^{N-m-1}$ .

ii)  $h|S^{N-m-1}$  is of degree  $p$  and  $h|S^{N-m-1}B\pi^{(s)}$  has non-trivial functional  $\mathfrak{B}^i(Sq^i)$ -operations for  $2i(p-1) \leq s+1$  ( $2 \leq i \leq s+1$ , respectively).

*Proof.* Recall that  $N=N(a, s+2)$  with sufficiently large  $a$ . Put  $m+1=N(b, s+1)$  for any  $b$  with  $0 < b < a-1$ . Then  $N-m-1=N(c, s+1)$  and  $N-n-1=N(c, s+1)+s$  for some integer  $c$ . So by the James periodicity for  $p=2$  ([5]) and by Theorem 4 of [7] for  $p>2$ , we have  $B\pi_{N-m-1}^{N-n-1} \simeq S^{N-m-1}B\pi_0^s = S^{N-m-1}B\pi^{(s)} \vee S^{N-m-1}$ . This leads us to i).

Since  $N-m-1$  is even,  $h|S^{N-m-1}$  is of degree  $p$ . For  $i > N-m-2$ , there is the natural isomorphism  $H^i(SB\pi_{N-m-2}^{N-n-1}; Z_p) \cong H^{i-1}(B\pi^{(N-n-1)}; Z_p)$ . Let  $u$  and  $v$  be generators of  $H^{i-1}(B\pi^{(N-n-1)}; Z_p)$  for  $i=2$  and 3 respectively. Then the non-triviality of the functional  $\mathfrak{B}^i$ -operation follows directly from the following relation:

$$\mathfrak{B}^i(uv^{(N-m-3)/2}) = u\mathfrak{B}^i(v^{(N-m-3)/2}) = \binom{cp^{\lceil (s+1)/2 \rceil} - 1}{i} uv^{(N-m-3)/2+i(p-1)} \neq 0$$

for  $2i(p-1) \leq s+1$ .

Similarly the functional  $Sq^i$ -operation is non-trivial for  $2 \leq i \leq s+1$ .

This completes the proof.

## 5. A proof of the main theorem

Obviously we have the following

**Lemma 8.** *If  $Y$  is an  $(r-1)$ -connected CW-complex of dimension  $r+s$  with  $r>s$ , then there exists an  $(s-1)$ -connected CW-complex  $W$  of dimension  $2s$  such that  $Y \simeq S^{r-s}W$ .*

Now we are ready to prove Theorem 1 supposing Theorem 3. If  $X$  is

$(l-1)$ -connected and  $\dim X=j$ , then  $D_N X$  is  $(N-j-2)$ -connected and  $\dim D_N X=N-l-1$ . Therefore, by the above lemma, there exists a  $(j-l-1)$ -connected and  $2(j-l)$ -dimensional  $CW$ -complex  $W$  such that  $D_N X \simeq S^{N+l-2j-1}W$ . If  $m+l>2j$ , then  $S^{N+l-2j-1}W=S^{N-m-1}(S^{m+l-2j}W)$  and  $\dim S^{m+l-2j}W=m-l=k$ . Hence Propositions 4 and 7 for  $n=l$  and Theorem 3 complete the proof of Theorem 1.

**6. An example**

Theorem 1 does not hold without the assumption  $2 \dim X < m+l$ . This is shown as follows.

Let  $\iota \in \{S^0, S^0\}$ ,  $\eta \in \{S^1, S^0\}$  and  $\nu \in \{S^3, S^0\}$  be generators. Put  $\alpha = \nu \vee 2\iota$  and  $X = (S^{m-5} \vee S^{m-2}) \cup_{\omega} e^{m-1}$ . Then it is clear that  $\pi_m(X) = \{\tilde{\eta}\} \cong Z_4$  for  $m > 11$ , where  $\tilde{\eta}$  is a co-extension of  $\eta$ . It is shown as follows that  $\tilde{\eta}$  is not symmetric for any  $m > 10$ .

If  $\tilde{\eta}$  is represented by a symmetric map  $f: S^m \rightarrow X$ , then  $f$  is decomposed as (2) for  $n=m-5$ . It is easily seen that  $m$  is odd and  $(f')^*: H^{m-1}(X; Z_2) \rightarrow H^{m-1}(P_{m-5}^m; Z_2)$  is an isomorphism. Put  $m \equiv k \pmod 8$ , where  $k=1, 3, 5$  or  $7$ . Since  $Sq^4$  is non-trivial in  $H^*(X; Z_2)$ , we have  $k=1$  or  $3$  and  $(f')^*: H^{m-5}(X; Z_2) \rightarrow H^{m-5}(P_{m-5}^m; Z_2)$  is an isomorphism. The operation  $Sq^2: H^{m-1}(P_{m-5}^{m+1}; Z_2) \rightarrow H^{m+1}(P_{m-5}^{m+1}; Z_2)$  is non-trivial and so we have  $k=3$ .

Consider the diagram (2)' for  $n=m-5$ . Then we have

$$\begin{array}{ccc}
 S^{N-m-1} & \xleftarrow{\Delta_N f} & D_N X \\
 \Delta_N \omega' \swarrow & & \searrow \Delta_N(f') \\
 & P_{N-m-1}^{N-m+4} &
 \end{array}$$

, where  $N = a2^{\phi(5)} = 8a$  for sufficiently large  $a$  and  $D_N X = S^{N-m} \cup e^{N-m+1} \cup e^{N-m+4}$ . Put  $N-m = 8t+5$  and let  $q: P_{8t+4}^{8t+9} \rightarrow S^{8t+9}$  be the collapsing map. Then it is clear that  $q\Delta_N(f'): D_N X \rightarrow S^{8t+9}$  is also the collapsing map and  $(q\Delta_N(f'))^*: \widetilde{KO}(S^{8t+9}) \rightarrow \widetilde{KO}(D_N X) \cong Z_2$  is an isomorphism. On the other hand, we have  $\widetilde{KO}(P_{8t+4}^{8t+9}) \cong Z + Z_4$  by Theorem 7.4 of [1]. This is a contradiction. Hence  $\tilde{\eta}$  is not symmetric for  $m > 10$ .

KYOTO UNIVERSITY  
 OSAKA UNIVERSITY  
 KYOTO UNIVERSITY

## References

- [1] J. F. Adams: *Vector fields on spheres*, Ann. of Math. **75** (1962), 603–632.
- [2] ———: *On the groups  $J(X)$ -II*, Topology. **3** (1965), 137–172.
- [3] M. F. Atiyah: *Thom complexes*, Proc. London Math Soc. **11** (1961), 291–310.
- [4] G. E. Bredon: *Equivariant homotopy*, Proc. Conference on Transformation Groups, New Orleans, Springer Verlag (1968), 281–292.
- [5] I. M. James: *Cross-sections of Stiefel manifolds*, Proc. London Math. Soc. **8** (1958), 536–547.
- [6] D. S. Kahn and S. Priddy: *Applications of the transfer to stable homotopy theory*, Bull. Amer. Math. Soc. **78** (1972), 981–987.
- [7] T. Kambe, H. Matsunaga and H. Toda: *A note on stunted lens space*, J. Math. Kyoto Univ. **5** (1966), 143–149.
- [8] J. Mukai: *Even maps from spheres to spheres*, Proc. Japan Acad. **47** (1971), 1–5.
- [9] G. Nishida: *The nilpotency of elements of the stable homotopy groups of spheres*, J. Math. Soc. Japan **25** (1973), 703–732.
- [10] E. Rees: *Symmetric maps*, J. London Math. Soc. **3** (1971), 267–272.
- [11] E. H. Spanier and J. H. C. Whitehead: *Duality in homotopy theory*, Mathematika. **2** (1955), 56–80.
- [12] J. Strutt: *Projective homotopy classes of spheres in the stable range*, Bol. Soc. Mat. Mexicana **16** (1971), 15–25.
- [13] ———: *Projective homotopy classes of Stiefel manifolds*, Canad. J. Math. **14** (1972), 465–476.
- [14] J. H. C. Whitehead: *On the groups  $\pi_r(V_{n,m})$  and sphere bundles*, Proc. London Math. Soc. **48** (1944), 243–291.



