

Rescue of Amyloid-Beta-Induced Inhibition of Nicotinic Acetylcholine Receptors by a Peptide Homologous to the Nicotine Binding Domain of the Alpha 7 Subtype

Arthur A. Nery¹*, Margaret H. Magdesian²*, Cleber A. Trujillo¹, Luciana B. Sathler², Maria A. Juliano³, Luiz Juliano³, Henning Ulrich^{1*}, Sergio T. Ferreira^{2*}

1 Department of Biochemistry, Chemistry Institute, São Paulo University, São Paulo, SP, Brazil, **2** Institute of Medical Biochemistry, Federal University of Rio de Janeiro, Rio de Janeiro, RJ, Brazil, **3** Department of Biophysics, Federal University of São Paulo, São Paulo, SP, Brazil

Abstract

Alzheimer's disease (AD) is characterized by brain accumulation of the neurotoxic amyloid- β peptide (A β) and by loss of cholinergic neurons and nicotinic acetylcholine receptors (nAChRs). Recent evidence indicates that memory loss and cognitive decline in AD correlate better with the amount of soluble A β than with the extent of amyloid plaque deposits in affected brains. Inhibition of nAChRs by soluble A β 40 is suggested to contribute to early cholinergic dysfunction in AD. Using phage display screening, we have previously identified a heptapeptide, termed IQ, homologous to most nAChR subtypes, binding with nanomolar affinity to soluble A β 40 and blocking A β -induced inhibition of carbamylcholine-induced currents in PC12 cells expressing α 7 nAChRs. Using alanine scanning mutagenesis and whole-cell current recording, we have now defined the amino acids in IQ essential for reversal of A β 40 inhibition of carbamylcholine-induced responses in PC12 cells, mediated by α 7 subtypes and other endogenously expressed nAChRs. We further investigated the effects of soluble A β , IQ and analogues of IQ on α 3 β 4 nAChRs recombinantly expressed in HEK293 cells. Results show that nanomolar concentrations of soluble A β 40 potently inhibit the function of α 3 β 4 nAChRs, and that subsequent addition of IQ or its analogues does not reverse this effect. However, co-application of IQ makes the inhibition of α 3 β 4 nAChRs by A β 40 reversible. These findings indicate that A β 40 inhibits different subtypes of nAChRs by interacting with specific receptor domains homologous to the IQ peptide, suggesting that IQ may be a lead for novel drugs to block the inhibition of cholinergic function in AD.

Citation: Nery AA, Magdesian MH, Trujillo CA, Sathler LB, Juliano MA, et al. (2013) Rescue of Amyloid-Beta-Induced Inhibition of Nicotinic Acetylcholine Receptors by a Peptide Homologous to the Nicotine Binding Domain of the Alpha 7 Subtype. PLoS ONE 8(7): e67194. doi:10.1371/journal.pone.0067194

Editor: Stefano L. Sensi, University G. D'Annunzio, Italy

Received: February 1, 2013; **Accepted:** May 15, 2013; **Published:** July 22, 2013

Copyright: © 2013 Nery et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by grants from National Institute for Translational Neuroscience/Brazil (STF), Conselho Nacional de Desenvolvimento Científico e Tecnológico (STF and HU), Fundação de Amparo à Pesquisa do Estado do Rio de Janeiro (STF and MHM), Fundação Universitária José Bonifácio (MHM), and Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) (HU) (project No. 2006/61285-9 and 2012/50880-4). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have read the journal's policy and have the following conflict: corresponding authors Henning Ulrich and Sergio T. Ferreira are PLOS ONE Editorial Board members. This does not alter the authors' adherence to all the PLOS ONE policies on sharing data and materials.

* E-mail: henning@iq.usp.br (HU); ferreira@bioqmed.ufrj.br (STF)

† These authors contributed equally to this work.

Introduction

Alzheimer's disease (AD) is the most common age-related neurodegenerative disorder and the seventh leading cause of death in the United States [1]. Currently, no effective treatment is available to slow down or stop deterioration of nerve cells in AD. This irreversible disease appears to be initiated by synapse failure, resulting in impairment of cognitive and other cerebral functions [2]. A large body of evidence indicates that the primary agent of neurodegeneration in AD is a 39–43 amino acid residues long peptide known as amyloid- β (A β). The majority of secreted A β is 40 amino acids long (A β 40), while the longer, 42-amino acid species (A β 42) has a high propensity to nucleate and drive the formation of soluble aggregates (e.g., oligomers, protofibrils) and insoluble amyloid fibrils [3,4,5]. Substantial evidence indicates that soluble A β oligomers are the proximal neurotoxins responsible for synapse dysfunction in AD (for reviews, see [2,6,7]. However, the

mechanisms linking A β 40 to synapse dysfunction and neuronal loss remain to be fully elucidated.

A prominent feature of AD pathology is the loss of cholinergic neurons and nicotinic acetylcholine receptors (nAChRs) throughout the brain [8,9]. With nearly 30 subtypes of brain nAChRs having been described, the three most abundant nAChR subtypes in the mammalian brain are composed of α 7, α 4 β 2, and α 3 β 4 subunits [10], expressed in major brain areas including cortex and hippocampus [11]. Although the direct binding of A β to α 7 receptors has been questioned [12], high-affinity association of A β 42 with α 7 and α 4 β 2 nAChRs has been observed in amyloid plaques and in neurons of AD patients [13,14,15,16]. There is also considerable evidence that A β affects the function of nAChRs (for reviews, see [17,18]). Nanomolar concentrations of A β 42 or A β 40 have been reported to inhibit both human and rat homomeric α 7 receptors [19,20,21,22,23,24]. Moreover, A β has been shown to exert subtype-specific actions, activating non- α 7 nAChRs in rat basal forebrain neurons [25] and inhibiting non- α 7 nAChR

subtypes ($\alpha 4\beta 2$, $\alpha 2\beta 2$, $\alpha 4\alpha 5\beta 2$) in rodent hippocampal slices [26]. In studies employing heterologously expressed human nAChRs, A β has been shown to inhibit $\alpha 7$ and $\alpha 4\beta 2$ subtype function without affecting $\alpha 3\beta 4$ nAChRs [23]. Those effects, however, are still somewhat controversial, as other reports show that picomolar concentrations of A β have no effect [23] or even activate whole-cell current responses of $\alpha 7$ nAChRs ([27,28,29,30]; for a review, see [31]).

Using phage-display screening of a peptide library, we previously reported that soluble A β binds with nanomolar affinity to a heptapeptide with amino acid sequence IQTTWSR, henceforth denoted IQ, which is homologous to an amino acid sequence located at the nicotine and acetylcholine (ACh) binding pocket in most subtypes of human nAChRs [24]. Nanomolar concentrations of IQ block A β -induced inhibition of carbamylcholine-induced currents in neuronal-differentiated PC12 cells expressing $\alpha 7$ nAChRs, suggesting that inhibition of nAChRs by A β results from its binding to the nicotine/ACh binding domain in the receptor. Our previous results further indicated that A β interacts with several nAChR subunits homologous to IQ, such as α subunits [24]. Crystallographic studies and alignment of nAChR sequences reveal that the location of the ligand binding site is highly conserved in nAChRs, but the actual ligand binding residues may vary, creating specificities for different ligands [32].

Here, we have used a combination of alanine scanning mutagenesis and rapid kinetic whole-cell current recording [33,34,35] to define the amino acid residues in IQ that are essential for alleviating blockade of the inhibition of $\alpha 7$ nAChRs by A $\beta 40$. In addition, we examined the effects of soluble A $\beta 40$, IQ and IQ peptide analogues on $\alpha 3\beta 4$ nAChRs, which are present in human brain but exhibit low homology to $\alpha 7$ in terms of amino acid sequences at the nicotine/ACh binding site. Results show that nanomolar concentrations of soluble A $\beta 40$ inhibit $\alpha 3\beta 4$ nAChRs. In contrast with our previous observations on $\alpha 7$ nAChRs [24], IQ and its analogues do not block A $\beta 40$ inhibition of $\alpha 3\beta 4$ nAChRs. However, simultaneous exposure to IQ and A $\beta 40$ makes the inhibition of $\alpha 3\beta 4$ nAChRs by A $\beta 40$ reversible. These results suggest that A β binds to distinct binding sites on different nAChR subtypes and point to the region homologous to IQ in nAChRs as a relevant target for A $\beta 40$ neurotoxicity in AD.

Results

Amino acid residues of IQ involved in blocking inhibition of $\alpha 7$ and other endogenously expressed nAChRs by A $\beta 40$

In order to identify key amino acid residues of the IQ peptide involved in blockade of A β -induced inhibition of nAChRs, whole-cell recordings of nAChR currents were carried out in neuronal-differentiated PC12 cells. RT-PCR analysis revealed that such cells express $\alpha 3$, $\alpha 5$, $\alpha 7$, $\beta 2$ and $\beta 4$ nAChR subunits, and measurements in the presence of methyllycaconitine (MLA) indicated that, on day 2 following induction to neuronal differentiation, $\alpha 7$ receptors contributed about 50% of cholinergic receptor-mediated whole cell currents [36].

We have previously shown that soluble A $\beta 40$ (200 nM) caused a marked (~60%) inhibition of nAChR currents and that addition of 500 nM IQ completely blocked this effect. Control measurements showed that addition of IQ alone (up to 500 nM) did not elicit any current in differentiated PC12 cells and (up to 750 nM) did not interfere with currents evoked by carbamylcholine (CCh) [24]. However, at higher concentrations (>1 μ M) IQ inhibited nAChR-mediated whole cell currents (I_{CCh}) and induced cell death (data not shown), indicating a relatively narrow concentra-

tion range in which IQ could be safely used to prevent nAChR inhibition by A $\beta 40$ in the absence of cell toxicity.

We have now investigated a series of IQ analogues for their abilities to block A β -induced inhibition of nAChRs in the absence of cell toxicity. A number of peptides were synthesized corresponding to a full alanine scan of the IQ heptapeptide or to truncated tetrapeptides. The impact of those peptides on cell viability was initially tested by the MTT assay, and none of them exhibited cytotoxicity at concentrations of 1 or 100 μ M in PC12 cells (Fig. S1). Moreover, no toxic effects exerted by these peptides were observed in HEK cells transfected for $\alpha 3$, $\beta 4$ receptor expression (data not shown).

Each of the IQ analogue peptides (at a fixed concentration of 500 nM, based on our previous results with IQ; ref 24) was then tested for its capacity to alleviate A β -induced inhibition of nAChRs in PC12 cells (measured in the presence of 0.2 mM CCh and 200 nM A $\beta 40$ in order to assess maximum inhibition rates; ref 24). Among the tetrapeptides tested, TTWS best mimicked the effect of full-length IQ (Fig. 1), completely reversing A $\beta 40$ inhibition of nAChR-mediated whole cell currents (I_{CCh} 95 \pm 2%), followed by TWSR (I_{CCh} 84 \pm 4%), IQTT (I_{CCh} 80 \pm 5%) and QTTW (I_{CCh} 72 \pm 6%). Representative current traces are shown in Fig. S2. Alanine scanning of the IQ sequence showed that replacement of Ile eliminated the capacity to block A β -induced inhibition of nAChRs (Ile \rightarrow Ala, I_{CCh} 57 \pm 4%). Moreover, replacement of Trp or Ser residues by Ala resulted in significantly reduced abilities to block A $\beta 40$ inhibition (Trp \rightarrow Ala, I_{CCh} 72 \pm 3%; Ser \rightarrow Ala, I_{CCh} 79 \pm 3%).

Effects of A β , IQ and IQ analogues on $\alpha 3\beta 4$ nAChRs

Given the abundance of $\alpha 3\beta 4$ receptors in the human brain, we next investigated the inhibition of $\alpha 3\beta 4$ nAChRs by A $\beta 40$ in the presence of 200 nM A $\beta 40$ at effective 0.2 mM CCh concentration. Co-application of 200 nM A $\beta 40$ and 0.2 mM CCh resulted in approximately 35% inhibition of $\alpha 3\beta 4$ nAChR currents in transformed HEK cells (Fig. 2). Successive shots of 0.2 mM CCh on the same cell at 5 min intervals (Fig. 2, white bars, shots 1–6) had no significant effect in the response to CCh, indicating lack of receptor desensitization under these conditions. However, application of three successive shots of 0.2 mM CCh plus 200 nM A $\beta 40$ (Fig. 2, light grey bars, applications 1–3) reduced the cellular response to CCh to approximately 60% of the control level. Subsequent application of three additional shots of 0.2 mM CCh alone to the same cell did not recover the original response to CCh (Fig. 2, light grey bars, shots 4–6), indicating that $\alpha 3\beta 4$ nAChRs remained inhibited even after washout of A β . In fact, A β -induced inhibition of $\alpha 3\beta 4$ nAChRs persisted even after 6 successive shots of CCh (4 minutes between each shot, comprising approximately 30 minutes for each experiment) following a single initial exposure to 0.2 mM CCh plus 200 nM A $\beta 40$ (Fig. S3).

Irreversible inhibition of $\alpha 3\beta 4$ receptors by A β was also observed following three shots of 0.2 mM CCh plus 200 nM A $\beta 40$ and 0.5 μ M SQI (a control scrambled peptide that has the same amino acid composition as IQ but does not bind to A β), followed by three shots of 0.2 mM CCh alone (Fig. 2, black bars). Cells that had been exposed to three shots of 0.2 mM CCh plus 200 nM A $\beta 40$ in the presence of 0.5 or 2 μ M IQ presented reduced response to CCh stimulation (~60% and 70% of control currents, respectively; Fig. 2, grey bars, shots 1–3). Thus, in contrast with its ability to block inhibition of $\alpha 7$ nAChRs [24], IQ was not capable of preventing the inhibition of $\alpha 3\beta 4$ nAChRs by A β . Interestingly, however, the response of $\alpha 3\beta 4$ receptors to CCh (Fig. 2, grey bars, shots 4–6) returned to control levels during the washout period after the co-application of CCh, A $\beta 40$ and IQ.




Figure 1. IQ and selected analogues reverse $A\beta$ 40 inhibition of nAChRs in PC12 cells. (A) Current responses (normalized by the maximal current evoked by 0.2 mM CCh) of neuronal-differentiated PC12 cells exposed for 2 s to 0.2 mM CCh plus 200 nM $A\beta$ 40 in all experimental conditions, except for the control measurement with CCh alone, and, as indicated, 500 nM of different IQ analogues. Bars represent means \pm S.D. of at least 3 replicate measurements performed in 4–6 different cells (**, $p < 0.01$; ***, $p < 0.001$ in the comparison with the control current evoked by CCh alone).
doi:10.1371/journal.pone.0067194.g001




Figure 2. IQ makes $A\beta$ 40 inhibition of $\alpha 3\beta 4$ nAChR currents in transformed HEK cells reversible. HEK cells expressing $\alpha 3\beta 4$ nAChRs received consecutive shots (at 5 min intervals) of 0.2 mM CCh plus 200 nM $A\beta$, in the absence or presence of IQ (0.5 μ M) as indicated. Shots 1–3 contained 0.2 mM CCh alone (white bars), 0.2 mM CCh plus 200 nM $A\beta$ (light grey bars), 0.2 mM CCh plus 200 nM $A\beta$ and 0.5/2 μ M IQ (grey bars) or 0.5 μ M SQI (black bars), used as an inactive control. Shots 4–6 contained 0.2 mM CCh alone for evaluation of reversibility of receptor inhibition. Bars represent mean values \pm S.D. of at least 3 replicate measurements (normalized by the maximal current evoked by 0.2 mM CCh) obtained from 4–6 different cells. (***, $p < 0.001$, in comparison with 0.2 mM CCh plus 200 nM $A\beta$).
doi:10.1371/journal.pone.0067194.g002

This indicates that, in the presence of IQ, the inhibition of $\alpha 3\beta 4$ receptors by $A\beta$ 40 becomes reversible following $A\beta$ 40 washout.

As a control, we tested whether IQ, QI (a peptide with a reverse sequence compared to IQ) or SQI induced activation of $\alpha 3\beta 4$ nAChR currents or had any impact on cellular response to CCh. Results showed that none of the three peptides by themselves elicited currents or had any detectable effect on whole-cell current responses of PC12 cells (Fig. S4), supporting the notion that rescue of cellular $\alpha 3\beta 4$ nAChR response to CCh by IQ is due to its interaction with $A\beta$.

Finally, we evaluated the effects of IQ and selected peptide analogues on the inhibition of $\alpha 3\beta 4$ nAChRs by $A\beta$. To this end, cells received three shots of each peptide as shown in Fig. 3. For all cells analyzed (at least 3 cells per experimental condition), currents measured in the presence of the peptides were compared to those measured in the presence of CCh alone or CCh+ $A\beta$. For each cell, 3 shots (with a 4 minute interval between them) of CCh were applied to elicit maximum responses, then 3 shots of CCh+ $A\beta$ to induce inhibition, followed by 3 shots of CCh+ $A\beta$ +peptide, and finally 3 more shots of CCh alone in order to verify the persistence of inhibition.

In the absence of peptides, inhibition by $A\beta$ 40 was found to be persistent when CCh alone was applied after the shots of CCh+ $A\beta$. Interestingly, when shots included CCh+ $A\beta$ +IQTTWSR (0.5 or 2 μ M), $\alpha 3\beta 4$ nAChR currents were rescued from inhibition when measured in the presence of CCh alone (after washout of $A\beta$). We next tested the effects of the TTWS and TWSR tetrapeptides, which had shown the best protective actions against $A\beta$ -induced inhibition of nAChRs, and IQTTASR, which lacks the highly conserved Trp residue in the agonist-binding domain of nAChRs and presented the lowest capacity to alleviate




Figure 3. Effects of IQ and analogues on A β 40 inhibition of α 3 β 4 nAChRs in transformed HEK cells. (A) HEK cells expressing recombinant α 3 β 4 nAChRs received 3 consecutive shots (at 4 min intervals) of 0.2 mM CCh plus 200 nM A β 40 in the absence or presence of 0.5 and 2 μ M IQTTWSR. QI and SQI (500 nM) were used as ineffective control peptides. Recovery of current response was evaluated after washout and 3 shots of CCh alone. (B) HEK cells expressing recombinant α 3 β 4 nAChRs received 3 consecutive shots of 0.2 mM CCh plus 200 nM A β 40 in the absence or presence of 500 nM TTWS, TWSR or IQTTASR. QI and SQI (500 nM) were used as ineffective control peptides. Recovery of current response was evaluated after washout and 3 shots of CCh alone. Bars represent mean values \pm S.D. of current responses (normalized by the maximal current evoked by 0.2 mM CCh) of at least 3 measurements performed in at least 3 different cells. (***, $p < 0.001$). doi:10.1371/journal.pone.0067194.g003

A β 40 inhibition of α 7 currents (Fig. 1). A slight increase in A β -induced inhibition was observed in the presence of SQI, which, however, was not statistically significant.

When tested on α 3 β 4 nAChR-expressing cells, all peptides tested failed in preventing the inhibition of α 3 β 4 nAChR-

mediated currents by A β . However, when A β 40 was added to cells in conjunction with IQ, TTWS or IQTTASR (0.5 μ M each), α 3 β 4 nAChR-mediated currents in response to CCh alone returned to approximately 100%, 88% and 94%, respectively, of control levels. On the other hand, TWSR, QI and SQI peptides

were unable to rescue the inhibition of CCh-induced $\alpha 3\beta 4$ nAChR-mediated responses by A $\beta 40$ (Fig. 3).

Discussion

We have previously identified an A $\beta 40$ ligand, a peptide termed IQ, that blocks A β -induced inhibition of nAChRs at nanomolar concentrations [24]. IQ is homologous to the ligand-binding domain of nAChRs. The location of the ligand-binding site is conserved among different pentameric ligand gated ion channel receptors, but the actual ligand binding residues may vary, creating specificities for different ligands [32]. Therefore, we proposed that A $\beta 40$ might interact with the ligand-binding domain of distinct nAChRs subunits, preferentially binding to those with higher homology to IQ, such as $\alpha 7$. This is consistent with previous reports of higher affinity interactions between A $\beta 40$ and $\alpha 7$ than with $\alpha 4\beta 2$ nAChRs from rat and guinea pig cerebral cortex and from hippocampal synaptic membranes [13,14]. Neuronal nAChRs are assembled as homomeric or heteromeric combinations of α ($\alpha 2$ –10) and β ($\beta 2$ –4) subunits. The majority of human CNS nAChRs is of the $\alpha 4\beta 2$ subtype and the remainder is largely made up of $\alpha 7$ subunit homopentamers and $\alpha 3\beta 4$ heteromers, although several other combinations are also known [10,23].

Here, we have asked which amino acid residues of IQ are essential for blockade of A $\beta 40$ inhibition of receptor currents in cells containing $\alpha 7$ or heteromeric nicotinic receptors, and specifically tested the effects of soluble A β , IQ and IQ analogues in cells expressing only the $\alpha 3\beta 4$ nAChR subtype. We used a whole-cell current-recording approach in combination with the cell-flow technique [33] to briefly expose differentiated PC12 cells or HEK cells expressing $\alpha 3\beta 4$ nAChRs to A $\beta 40$ and other ligands. This procedure minimizes receptor desensitization and avoids long periods of incubation with A β , assuring preservation of A $\beta 40$ in soluble state during the experiments, as previously described [24].

Co-application of 0.2 mM CCh, 200 nM A $\beta 40$ and 500 nM of different IQ analogues to differentiated PC12 cells showed that, among the tetrapeptides tested, TTWS was the analogue that best emulated the protective effect of full-length IQ, completely preventing A β -induced inhibition of nAChRs (I_{CCh} 95 \pm 2%). Next in terms of effectiveness were TWSR, IQTT and QTTW. Both TTWS and TWSR contain Trp57, a highly conserved residue present in the sequences of all nAChRs described so far [37]. Trp57 has been shown to be important for binding of d-tubocurarine (a competitive antagonist of nAChRs) to *Torpedo* nAChR [38]. Both peptides also contain a Ser residue (Ser58) present in 1 of the 12 human nAChR subunit sequences and conservatively replaced by Thr in 5 of the remainder 11 sequences. Ala-scanning of the IQ sequence indicated that the Trp and Ser residues of IQ are essential for efficacy in preventing A $\beta 40$ inhibition of nicotinic receptors. Ala substitutions also pointed to the importance of Ile in the IQ sequence. Significantly, Ile53 (or its highly conserved substitution Leu) is present in 11 of the 12 human nAChR subunits known to date.

On the other hand, replacement of Gln, Thr or Arg residues by Ala did not significantly affect the efficacy of IQ analogues (Fig. 1), despite the fact that mutations in Gln56 (numbering according to the $\alpha 7$ nAChR sequence) affect the affinities for ACh and nicotine [39]. Collectively, these results show that Ile, Trp and Ser residues in the amino acid sequence of IQ (IQTTWSR) are essential to block A $\beta 40$ inhibition of nAChRs. Based on these findings, we propose that protection by longer peptides (containing 6 amino acid residues or more) can be explained on the basis of a sequence motif in which Ile, Trp and Ser residues at positions 1, 5 and 6,


respectively, are conserved (i.e., IxxxWS). A similar model can be developed for shorter peptides (of 4 amino acid residues or less) and also for protection against inhibition of $\alpha 3\beta 4$ receptors and possibly other nicotinic subtypes by A $\beta 40$ (Fig. 4). For $\alpha 7$ nicotinic receptors and other subtypes expressed by PC12 cells, carboxy-terminal Trp and Ser residues must be conserved to preserve efficacy of tetrapeptides in blocking A $\beta 40$ inhibition. On the other hand, our results show that for $\alpha 3\beta 4$ receptors the Trp residue can be replaced by a nonpolar (aliphatic or aromatic) amino acid residue without loss in activity (Fig. 3B). Defining these structural motifs may prove useful for development of novel IQ analogues with improved efficacy in protection against A $\beta 40$ inhibition of nAChRs and/or recovery from such inhibition, and as a molecular backbone for development of non-peptide drugs.

The fact that IQ is homologous to several nAChR subunits [24] suggests that A $\beta 40$ binds to this highly conserved domain in different nAChR subtypes. Although direct binding was not tested in the present study, we assume a similar mechanism of action for the other tested peptides. In order to test this hypothesis, we tested the effects of A β , IQ and IQ analogues in a cell line expressing a single subtype of nAChR, the $\alpha 3\beta 4$ subtype, characterized by large whole-cell current responses and widely used as model for binding and activity screening on nicotinic receptors [40,41,42,43,44]. Moreover, this receptor subtype was recently employed to characterize the mechanism of action of the Alzheimer drug tacrine [42] and has also been shown to be involved in disease states such as nicotine-induced seizures and hypolocomotion in mice [45]. Results showed that 200 nM soluble A $\beta 40$ persistently blocked the response of $\alpha 3\beta 4$ nAChRs to CCh remaining 60 \pm 14% of currents induced by CCh alone. To our knowledge, there is only one other study testing the effects of A $\beta 40$ on $\alpha 3\beta 4$ nAChRs [23]. That study showed that A β failed to elicit changes in amplitude of ACh-evoked currents mediated by human $\alpha 3\beta 4$ nAChRs expressed in *Xenopus laevis* oocytes. It should be noted, however, that A $\beta 40$ was bath applied at a significantly lower concentration (10 nM) than used in the present study (200 nM). Moreover, Pym et al. [23] pre-incubated A $\beta 40$ with cells for 3 min, which might lead to aggregation and, consequently, to a decrease in the concentration of soluble A $\beta 40$ species that directly interact with nAChRs. Although the concentration of A $\beta 40$ in the cerebrospinal fluid of AD patients has been reported to be between 1 and 10 nM [46], the concentration of A $\beta 40$ at cholinergic synapses is unknown.

In contrast to their effects in cells expressing $\alpha 7$ and heteromeric nAChRs, IQ and analogues did not block A $\beta 40$ inhibition of $\alpha 3\beta 4$ nAChRs expressed in HEK cells, suggesting that IQ binding to A $\beta 40$ is not sufficient to prevent A β from interacting with and inhibiting $\alpha 3\beta 4$ receptors. However, addition of IQ made the inhibition of $\alpha 3\beta 4$ nAChRs by A $\beta 40$ reversible, suggesting that IQ binding to A $\beta 40$ modifies its interaction with $\alpha 3\beta 4$ receptors, likely facilitating A $\beta 40$ dissociation and receptor re-activation by the agonist. Activity screening of IQ analogues indicated that only TTWS and IQTTASR were able to mimic IQ and make A $\beta 40$ inhibition of $\alpha 3\beta 4$ nAChRs reversible, emphasizing the importance of the TTWS tetrapeptide in A $\beta 40$ interaction with nAChRs. Unexpectedly, the Trp residue (present in all nAChRs) that is essential in IQ to block A $\beta 40$ inhibition of $\alpha 7$ nAChRs was not necessary to alleviate the effects of A $\beta 40$ on $\alpha 3\beta 4$ nAChRs, suggesting that different amino acid residues or different protein domains are involved in A $\beta 40$ interaction with distinct subtypes of nAChRs.

Current results support the notion that A $\beta 40$ binds with distinct affinities to and has different effects on various subtypes of nAChRs [6,27]. Indeed, it has been reported that A β binds with

Large Peptides



Small Peptides

nAChRs



$\alpha 3\beta 4$



Figure 4. Suggested conserved amino acid sequence for reversal of $\alpha 3\beta 4$ nAChR inhibition by A β .
doi:10.1371/journal.pone.0067194.g004

high affinity (in the picomolar range) to $\alpha 7$ nAChRs in cortical regions and in the hippocampus in AD, and with about 5,000 times lower affinity to $\alpha 4\beta 2$ nAChRs [13,14]. However, as a general mechanism, blockade of nAChRs by A β may also affect, at concentrations similar to those used in the present study, the cholinergic control of neurotransmitter release, including glycine, glutamate, aspartate and GABA [47,48].

Distinct effects of A β on different subtypes of nAChRs reinforce the idea that A β binding to different receptor subtypes may involve different binding sites, occasionally increasing but more often blocking agonist response. Moreover, the difference in effects of IQ and analogues in cells expressing $\alpha 7$ and other heteromeric receptors versus in cells solely expressing $\alpha 3\beta 4$ nAChRs may be due to the fact that IQ presents higher homology to the ligand binding pocket of $\alpha 7$ than of $\alpha 3\beta 4$ nAChRs. In nAChRs, the ligand-binding site is located at the interface between two subunits [46,49]. Numerous biochemical studies have shown that the principal part of the binding site is formed by α -subunit residues [50,51,52,53], whereas neighboring subunit residues contribute to form the complementary part of the binding pocket. Thus, heteropentamers such as $\alpha 3\beta 4$ subtype contain two different ligand-binding sites with distinct affinities, whereas the homopentameric $\alpha 7$ receptor contains five identical ligand-binding sites [32]. The most vulnerable neurons in AD seem to be those expressing high levels of nAChRs, particularly those containing the $\alpha 7$ subunit [54], and levels of nAChRs as well as some of their associated proteins decrease in AD [55,56]. An interesting recent study reported that deletion of the $\alpha 7$ nAChR gene prevents cognitive deficits and synaptic pathology in a mouse model of

Alzheimer's disease [57]. Our current results provide novel information to drive further progress in AD drug design. Drugs like IQ, capable of disrupting A β - $\alpha 7$ nAChR interactions, might alleviate A β -mediated toxicity and block AD development.

In conclusion, our finding that A β exerts subtype-specific inhibitory effects on $\alpha 7$ and $\alpha 3\beta 4$ nAChRs suggests that receptor subunit composition might account for some of the different actions reported for A $\beta 40$ on neurons *in vivo*. Furthermore, we show that the region homologous to IQ in nAChRs is a relevant target to alleviate blockade of $\alpha 7$ and $\alpha 3\beta 4$ nAChRs by A β . Our results identify, for the first time, the amino acid residues probably involved in binding and inhibition of nAChRs by A β and may provide a valuable platform for drug design of novel AD therapeutics. The potential relevance of our findings to drug design and development of novel AD treatments is further underscored by a recent string of disappointing clinical trials on A β antibodies (Bapineuzumab and Solanezumab), which have cast a shadow over anti-A β immunotherapy strategies [58].

Materials and Methods

Peptide synthesis

Peptides IQ (IQTTWSR), AQTTSWR, IATTWSR, IQATWSR, IQTAWSR, IQTTASR, IQTTWAR, IQTTWSA, IQTT, QTTW, TTWS, TWSR and scrambled IQ (SQI; TIWQSTR) were synthesized as detailed elsewhere [24].

Cell culture

PC12 cells (ATCC, catalogue # CRL-1721) were cultured and induced to neuronal differentiation as described [24,36]. Briefly, PC12 cells were cultured in DMEM (Invitrogen, Life Technologies, Carlsbad, CA, USA) in the presence of 10% FBS (Cultilab, Campinas, São Paulo, Brazil), 5% horse serum (Invitrogen, Life Technologies, Carlsbad, CA, USA), streptomycin (100 μ g/ml), penicillin (100 U/ml – Sigma-Aldrich, St. Louis, MO, USA) and 1 mM sodium pyruvate (Invitrogen, Life Technologies, Carlsbad, CA, USA). $N^5,2'$ -O-dibutyl cAMP (dibutyl cAMP) and FGF-2 (45 ng/ml) were added to cultures to induce differentiation into mature sympathetic neurons expressing increased numbers of neuronal nAChRs [59]. For differentiation, 2.5×10^5 cells/ml, as determined by Neubauer chamber counting, were induced to neuronal differentiation for up to 6 days in DMEM containing 30 ng/ml FGF-2 (Sigma-Aldrich, St. Louis, MO, USA) and 250 μ M dibutyl cAMP (Sigma-Aldrich, St. Louis, MO, USA). Under these conditions, differentiated PC12 cells express $\alpha 3$, $\alpha 5$, $\alpha 7$, $\beta 2$ and $\beta 4$ nicotinic receptor subunits [36]. For evaluation of cell viability, PC12 cells on day 3 of neuronal differentiation were exposed for 48 h to different peptides at 1 and 100 μ M concentrations, then washed with PBS and stained with trypan blue. Five fields were photographed per well and live and dead cells were counted. Statistical analysis was based on the Student's t-test.

Human embryonic kidney cells (HEK293 cells) stably expressing rat $\alpha 3$ and $\beta 4$ nAChR subunits [60] were obtained from Dr. Yingxian Xiao, Georgetown University. Transfected cells were cultured in DMEM (Invitrogen, Life Technologies, Carlsbad, CA, USA) supplemented with 10% FBS, 100 U/ml penicillin G, 100 μ g/ml streptomycin and 0.3 mg/ml geneticin (Sigma-Aldrich, St. Louis, MO, USA) at 37°C and 5% CO₂. Cells were allowed to attach to 35 mm cell culture dishes for 48 h prior to being used in whole-cell recording experiments.

Whole-cell current recording and rapid application of ligand solutions (cell flow technique)

PC12 cells following 3–6 days of neuronal differentiation were cultured at a density of 20–100 cells/mm² on 35 mm cell culture dishes. Whole-cell recordings were performed at room temperature at a transmembrane voltage of –70 mV. The solution in the recording pipette contained 145 mM KCl, 10 mM NaCl, 2 mM MgCl₂, 1 mM EGTA, 25 mM HEPES, pH 7.4. The bath solution was composed of 145 mM NaCl, 5.3 mM KCl, 1.8 mM CaCl₂, 1.2 mM MgCl₂, 10 mM glucose, 25 mM HEPES, pH 7.4. Further details were previously reported [24].

Using carbamylcholine (CCh), a stable analog of ACh, we have previously shown that whole-cell current (I_{CCh}) data in neuronal-differentiated PC12 cells could be well described by a single binding site model, yielding a K_d of 259 ± 58 μ M for CCh. A K_d value of 2 mM has already been determined for $\alpha 3\beta 4$ nAChRs expressed by HEK cells [61]. According to previous work published by Niu et al., 1995 and Hess et al. 2000 [62,63], the equilibrium between open and closed channel forms is defined by the concentration of the agonist, thereby the closed channel form reveals higher affinity for the inhibitor as the open channel form does. Therefore, higher percentages of inhibition by A β 40 are expected at low CCh (0.2 mM) concentration. Because the density of receptors in the plasmamembrane (i.e., the total number of binding sites) differs somewhat from cell to cell, all I_{CCh} values were normalized to the currents measured in the presence of 0.2 mM CCh [24]. CCh-induced currents were recorded by whole-cell recording in combination with a rapid kinetic ligand delivery system, denominated the cell-flow technique, which

provides a time resolution of 10 ms [24,33,34,35]. Briefly, a U-shaped stainless steel capillary tube (250 μ m i.d.) with a circular porthole of 150 μ m in diameter at the base of the U was connected to pumps on both ends so the solution containing ligand could be driven into the tube at one end and removed through the other end at twice the entry flow rate [33]. The porthole was placed about 100 μ m away from each cell clamped by the recording pipette. Upon closing a solenoid valve between the U-tube and the suction pump by an electric trigger, CCh, A β 40 and/or different peptides were applied to the cell in a laminar flow. A β 40 and peptide solutions were mixed prior to co-application with CCh. Recorded signals were amplified using an Axopatch 200B amplifier (Molecular Devices, LLC, Sunnyvale, CA, USA) and filtered at 2 KHz using a 40-pole low-pass Bessel filter. The filtered signals were digitized using a Digidata 1322A interface, recorded using the pCLAMP software package (Molecular Devices) and analyzed using Microcal Origin software (Microcal Software, Inc., North Hampton, MA, USA). Statistical analysis was performed by comparing mean values using one-way analysis of variance (ANOVA) with Bonferroni's correction.

Correction for receptor desensitization in cell-flow measurements

The maximum current amplitude is proportional to the density of open channels. As receptor desensitization may occur while the ligand solution is equilibrating with the cell surface, observed current amplitudes are corrected for desensitization using the equation [33,35]:

$$I(t) = I_1 e^{-t/\tau_1} + I_2 e^{-t/\tau_2} + I_e, \quad (1)$$

where $I(t)$ is the maximum current amplitude at time t ; I_1 , I_2 , I_e are the maximum current amplitudes for the first, second, and equilibrium current decay components, respectively; and τ_1 and τ_2 are the time constants for the first and second components (fast and slow receptor desensitization, respectively). Origin software (Microcal Software, Inc.) was used to estimate the rate of current decay in the presence of agonist. Equation 1 was fitted to the decreasing part of the recording and the observed maximum current amplitude was corrected for receptor desensitization accordingly [33].

Supporting Information

Figure S1 Cytotoxicity assay selected IQ analogues. PC12 cells induced to neuronal differentiation were incubated in the presence of different peptides for 48 hours, washed with PBS and stained with trypan blue. The percentages of live and dead cells of five fields per well were counted and compared to those of control cells incubated in the absence of peptides. (TIFF)

Figure S2 Current traces of different peptides tested for reversion of $\alpha 3\beta 4$ nAChR inhibition by A β 40. Current responses (normalized by the maximal current evoked by 0.2 mM CCh) of neuronal differentiated PC12 cells exposed for 2 s to 0.2 mM CCh plus 200 nM A β 40 in all experimental conditions, except for the control measurement with CCh alone, and, as indicated, 500 nM of different IQ analogues. The here shown original data are illustrative for mean values \pm S.D. reported in Fig. 1. (TIF)

Figure S3 A β 40-induced inhibition of $\alpha 3\beta 4$ nAChR currents in transformed HEK cells persists after

washout. Following six consecutive applications of 0.2 mM CCh, 0.2 mM CCh was co-applied once in the presence of 200 nM A β 40. Following washout of A β , inhibition persisted in six consecutive applications of CCh ($p < 0.005$, when compared to control currents measured prior to A β administration). (TIF)

Figure S4 IQ, QI or SQI alone do not instigate nAChR currents in PC12 cells. The initial whole-cell response induced by 0.2 mM CCh was normalized to 100% of activity. None of the peptides (IQ, QI, SQI, tested at 2 μ M) induced changes in CCh-evoked currents nor activated receptor responses in the absence of agonist. Arrows indicate time points of ligand application. (TIF)

References

- Alzheimer's Association (2010) Alzheimer's disease facts and figures. *Alzheimers Dement* 6: 158–194.
- Ferreira ST, Klein WL (2011) The A β oligomer hypothesis for synapse failure and memory loss in Alzheimer's disease. *Neurobiol Learn Mem* 96: 529–543.
- Lambert MP, Barlow AK, Chromy BA, Edwards C, Freed R, et al. (1998) Diffusible, nonfibrillar ligands derived from Abeta1-42 are potent central nervous system neurotoxins. *Proc Natl Acad Sci USA* 95: 6448–6453.
- Riek R (2006) Cell biology: infectious Alzheimer's disease? *Nature* 444: 429–431.
- Greenwald J, Riek R (2010) Biology of amyloid: structure, function, and regulation. *Structure* 18: 1244–1260.
- Ferreira ST, Vieira MN, De Felice FG (2007) Soluble protein oligomers as emerging toxins in Alzheimer's and other amyloid diseases. *IUBMB Life* 59: 332–345.
- Selkoe DJ (2008) Soluble oligomers of the amyloid beta-protein impair synaptic plasticity and behavior. *Behav Brain Res* 192: 106–113.
- Nordberg A (2001) Nicotinic receptor abnormalities of Alzheimer's disease: therapeutic implications. *Biol Psychiatry* 49: 200–210.
- Kadir A, Almkvist O, Wall A, Långström B, Nordberg A (2006) PET imaging of cortical 11C-nicotine binding correlates with the cognitive function of attention in Alzheimer's disease. *Psychopharmacology (Berl)* 188: 509–520.
- Lindstrom JM (2003) Nicotinic acetylcholine receptors of muscles and nerves: comparison of their structures, functional roles, and vulnerability to pathology. *Ann N Y Acad Sci* 998: 41–52.
- Clementi F, Fornasari D, Gotti C (2000) Neuronal nicotinic receptors, important new players in brain function. *Eur J Pharmacol* 393: 3–10.
- Small DH, Maksel D, Kerr ML, Ng J, Hou X, et al. (2007) The beta-amyloid protein of Alzheimer's disease binds to membrane lipids but does not bind to the $\alpha 7$ nicotinic acetylcholine receptor. *J Neurochem* 101: 1527–1538.
- Wang HY, Lee DH, D'Andrea MR, Peterson PA, Shank RP, et al. (2000) Beta-Amyloid (1-42) binds to alpha7 nicotinic acetylcholine receptor with high affinity. Implications for Alzheimer's disease pathology. *J Biol Chem* 275: 5626–5632.
- Wang HY, Lee DH, Davis CB, Shank RP (2000). Amyloid peptide A β 1–42 binds selectively and with picomolar affinity to $\alpha 7$ nicotinic acetylcholine receptors. *J Neurochem* 75: 1155–1161.
- Nagele RG, D'Andrea MR, Anderson WJ, Wang HY (2002) Intracellular accumulation of beta-amyloid (1-42) in neurons is facilitated by the alpha 7 nicotinic acetylcholine receptor in Alzheimer's disease. *Neurosci* 110: 199–211.
- Wang HY, Bakshi K, Shen C, Frankfurt M, Trocmé-Thibierge C, et al. (2010) S 24795 limits beta-amyloid-alpha7 nicotinic receptor interaction and reduces Alzheimer's disease-like pathologies. *Biol. Psychiatry* 67: 522–530.
- Dineley KT (2007) Beta-amyloid peptide-nicotinic acetylcholine receptor interaction: the two faces of health and disease. *Front Biosci* 12: 5030–5038.
- Jürgensen S, Ferreira ST (2010) Nicotinic receptors, amyloid-beta, and synaptic failure in Alzheimer's disease. *J Mol Neurosci* 40: 221–229.
- Petit DL, Shao Z, Yakel JL (2001) Beta-Amyloid1–42 peptide directly modulates nicotinic receptors in the rat hippocampal slice. *J Neurosci* 21, RC120.
- Liu QS, Kawai H, Berg DK (2001) Beta-amyloid peptide blocks the response of alpha 7-containing nicotinic receptors on hippocampal neurons. *Proc Natl Acad Sci USA* 98: 4734–9.
- Grassi F, Palma E, Tonini R, Amici M, Ballivet V, et al. (2003) Amyloid beta (1-42) peptide alters the gating of human and mouse alpha-bungarotoxin-sensitive nicotinic receptors. *J Physiol* 547: 147–157.
- Wu J, Kuo YP, George AA, Xu L, Hu J, et al. (2004) beta-Amyloid directly inhibits human alpha4beta2-nicotinic acetylcholine receptors heterologously expressed in human SH-EP1 cells. *J Biol Chem* 279: 37842–37851.
- Pym L, Kemp M, Raymond-Delpech V, Buckingham S, Boyd CA, et al. (2005) Subtype-specific actions of beta-amyloid peptides on recombinant human neuronal nicotinic acetylcholine receptors ($\alpha 7$, $\alpha 4\beta 2$, $\beta 3\beta 4$) expressed in *Xenopus laevis* oocytes. *Br J Pharmacol* 146: 964–971.
- Magdesian MH, Nery AA, Martins AH, Juliano MA, Juliano L, et al. (2005) Peptide blockers of the inhibition of neuronal nicotinic acetylcholine receptors by amyloid beta. *J Biol Chem* 280: 31085–31090.
- Fu W, Jhamandas JH (2003) Beta-amyloid peptide activates non-alpha7 nicotinic acetylcholine receptors in rat basal forebrain neurons. *J Neurophysiol* 90: 3130–3136.
- Lamb PW, Melton MA, Yakel JL (2005) Inhibition of neuronal nicotinic acetylcholine receptor channels expressed in *Xenopus* oocytes by betaamyloid1–42 peptide. *J Mol Neurosci* 27: 13–21.
- Dineley KT, Bell KA, Bui D, Sweatt JD (2002) Beta-Amyloid peptide activates alpha 7 nicotinic acetylcholine receptors expressed in *Xenopus* oocytes. *J Biol Chem* 277: 25056–25061.
- Dougherty JJ, Wu J, Nichols RA (2003) Beta-amyloid regulation of presynaptic nicotinic receptors in rat hippocampus and neocortex. *J Neurosci* 23: 6740–6747.
- Wu J, Khan GM, Nichols RA (2007) Dopamine release in prefrontal cortex in response to beta-amyloid activation of alpha7 nicotinic receptors. *Brain Res* 1182: 82–89.
- Puzzo D, Privitera L, Leznik E, Fà M, Staniszewski A, et al. (2008) Picomolar amyloid-beta positively modulates synaptic plasticity and memory in hippocampus. *J Neurosci* 28: 14537–14545.
- Buckingham SD, Jones AK, Brown LA, Sattelle DB (2009) Nicotinic acetylcholine receptor signalling: roles in Alzheimer's disease and amyloid neuroprotection. *Pharmacol Rev* 61: 39–61.
- Brejčič K, van Dijk WJ, Klaassen RV, Schuurmans M, van Der Oost J et al. (2001) Crystal structure of an ACh-binding protein reveals the ligand-binding domain of nicotinic receptors. *Nature* 411: 269–276.
- Udgaonkar JB, Hess GP (1987) Chemical kinetic measurements of a mammalian acetylcholine receptor by a fast-reaction technique. *Proc Natl Acad Sci U S A* 84: 8758–8762.
- Ulrich H, Ippolito JE, Pagan OR, Eterović VA, Hann RM, et al. (1998) In vitro selection of RNA molecules that displace cocaine from the membrane-bound nicotinic acetylcholine receptor. *Proc Natl Acad Sci U S A* 95: 14051–14056.
- Ulrich H, Akk G, Nery AA, Trujillo CA, Rodriguez AD, et al. (2008) Mode of cembranoid action on embryonic muscle acetylcholine receptor. *J Neurosci Res* 86: 93–107.
- Nery AA, Resende RR, Martins AH, Trujillo CA, Eterović VA, et al. (2010) Alpha 7 nicotinic acetylcholine receptor expression and activity during neuronal differentiation of PC12 pheochromocytoma cells. *J Mol Neurosci* 4: 329–339.
- Williams DK, Stokes C, Horenstein NA, Papke RL (2009) Differential regulation of receptor activation and agonist selectivity by highly conserved tryptophans in the nicotinic acetylcholine receptor binding site. *J Pharmacol Exp Ther* 330: 40–53.
- Chiara DC, Cohen JB (1997) Identification of amino acids contributing to high and low affinity d-tubocurarine sites in the Torpedo nicotinic acetylcholine receptor. *J Biol Chem* 272: 32940–32950.
- Corringier PJ, Galzi JL, Eisele JL, Bertrand S, Changeux JP, et al. (1995) Identification of a new component of the agonist binding site of the nicotinic alpha 7 homooligomeric receptor. *J Biol Chem* 270: 11749–11752.
- Cheffer A, Mustafá EV, T-do Amaral A, Ulrich H (2012) Lipophilicity as a determinant of binding of procaine analogs to rat $\alpha 3\beta 4$ nicotinic acetylcholine receptor. *J Neurosci Res* 90: 1607–1614.
- Moaddel R, Jozwiak K, Yamaguchi R, Cobello C, Whittington K, et al. (2004) On-line screening of conformationally constrained nicotines and anabasines for agonist activity at the alpha3beta4- and alpha4beta2-nicotinic acetylcholine receptors using immobilized receptor-based liquid chromatographic stationary phases. *J Chromatogr B Analyt Technol Biomed Life Sci* 813: 235–240.
- Cheffer A, Ulrich H (2011). Inhibition mechanism of rat $\alpha 3\beta 4$ nicotinic acetylcholine receptor by the Alzheimer therapeutic tacrine. *Biochemistry* 50: 763–770.
- Nunes-Alves A, Nery AA, Ulrich H (2013) Tobacco nitrosamine N-nitrosornicotine as inhibitor of neuronal nicotinic acetylcholine receptors. *J Mol Neurosci* 49: 52–61.

Acknowledgments

We are grateful to Prof. Yingxian Xiao (Department of Pharmacology, Georgetown University, Washington, DC) for giving us the KX $\alpha 3\beta 4$ R2 cell line expressing the rat neuronal $\alpha 3\beta 4$ nAChR.

Author Contributions

Conceived and designed the experiments: AAN MHM MAJ LJ HU STF. Performed the experiments: AAN MHM CAT LBS. Analyzed the data: AAN MHM HU STF. Contributed reagents/materials/analysis tools: MHM MAJ LJ HU STF. Wrote the paper: AAN MHM HU STF.

44. Azam L, McIntosh JM (2009) Alpha-conotoxins as pharmacological probes of nicotinic acetylcholine receptors. *Acta Pharmacol Sin* 30: 771–783.
45. Salas R, Cook KD, Bassetto L, De Biasi M (2004) The alpha3 and beta4 nicotinic acetylcholine receptor subunits are necessary for nicotine-induced seizures and hypolocomotion in mice. *Neuropharmacology* 47: 401–417.
46. Mehta PD, Pirttila T, Patrick BA, Barshatzky M, Mehta SP (2001) Amyloid beta protein 1-40 and 1-42 levels in matched cerebrospinal fluid and plasma from patients with Alzheimer disease. *Neurosci Lett* 304: 102–106.
47. Zappettini S, Grilli M, Olivero G, Mura E, Preda S, et al. (2012) Beta amyloid differently modulate nicotinic and muscarinic receptor subtypes which stimulate in vitro and in vivo the release of Glycine in the rat hippocampus. *Front Pharmacol* 3:146.
48. Mura E, Zappettini S, Preda S, Biundo F, Lanni C, et al. (2012) Dual effect of beta-amyloid on $\alpha 7$ and $\alpha 4\beta 2$ nicotinic receptors controlling the release of glutamate, aspartate and GABA in rat hippocampus. *PLoS One* 2012;7(1): e29661.
49. Corringer PJ, LeNov e N, Changeux JP (2000). Nicotinic receptors at the amino-acid level. *Annu Rev Pharmacol Toxicol* 40: 431–458.
50. Arias HR (2000) Localization of agonist and competitive antagonist binding sites on nicotinic acetylcholine receptors. *Neurochem Int* 36: 595–645.
51. Dennis M, Giraudat J, Kotzby-Hibert F, Goeldner M, Hirth C, et al. (1988) Amino acids of the Torpedo marmorata acetylcholine receptor alpha subunit labeled by a photoaffinity ligand for the acetylcholine binding site. *Biochemistry* 27: 2346–2357.
52. Galzi JL, Revah F, Black D, Goeldner M, Hirth C, et al. (1990) Identification of a novel amino acid alpha-tyrosine 93 within the cholinergic ligand binding sites of the acetylcholine receptor by photoaffinity labeling. Additional evidence for a three loop model of the cholinergic ligands-binding sites. *J Biol Chem* 265: 10430–10437.
53. Fu DX, Sine SM (1994) Competitive antagonists bridge the alpha-gamma subunit interface of the acetylcholine receptor through quaternary ammonium-aromatic interactions. *J Biol Chem* 269: 26152–26157.
54. D'Andrea MR, Nagele RG (2006) Targeting the alpha7 nicotinic acetylcholine receptor to reduce amyloid accumulation in Alzheimer's disease pyramidal neurons. *Curr Pharm Des* 12: 677–684.
55. Martin-Ruiz CM, Court JA, Molnar E, Lee M, Gotti C, et al. (1999) Alpha4 but not alpha3 and alpha7 nicotinic acetylcholine receptor subunits are lost from the temporal cortex in Alzheimer's disease. *J Neurochem* 73: 1635–1640.
56. Gotti C, Moretti M, Bohr I, Ziabreva I, Vailati S, et al. (2006) Selective nicotinic acetylcholine receptor subunit deficits identified in Alzheimer's disease, Parkinson's disease and dementia with Lewy bodies by immunoprecipitation. *Neurobiol Dis* 23: 481–489.
57. Dziewczapolski G, Glogowski CM, Masliah E, Heinemann SF (2009) Deletion of the alpha 7 nicotinic acetylcholine receptor gene improves cognitive deficits and synaptic pathology in a mouse model of Alzheimer's disease. *J Neurosci* 29: 8805–8815.
58. Tayeb HO, Murray ED, Price BH, Tarazi FI (2013) Bapineuzumab and solanezumab for Alzheimer's disease: is the 'amyloid cascade hypothesis' still alive? *Expert Opin Biol Ther*. 2013 Apr 10. [Epub ahead of print]
59. Ho PL, Raw I (1992) Cyclic AMP potentiates bFGF-induced neurite outgrowth in PC12 cells. *J Cell Physiol* 150: 647–656
60. Xiao Y, Meyer EL, Thompson JM, Surin A, Wroblewski J, et al. (1998) Rat alpha3/beta4 subtype of neuronal nicotinic acetylcholine receptor stably expressed in a transfected cell line: pharmacology of ligand binding and function. *Mol Pharmacol* 54: 322–333.
61. Krivoshein AV, Hess GP (2004) Mechanism-based approach to the successful prevention of cocaine inhibition of the neuronal (alpha 3 beta 4) nicotinic acetylcholine receptor. *Biochemistry* 43: 481–489.
62. Niu L, Abood LG, Hess GP (1995) Cocaine: mechanism of inhibition of a muscle acetylcholine receptor studied by a laser-pulse photolysis technique. *Proc Natl Acad Sci U S A* 92: 12008–12012.
63. Hess GP, Ulrich H, Breitinger HG, Niu L, Gameiro AM, et al. (2000) Mechanism-based discovery of ligands that counteract inhibition of the nicotinic acetylcholine receptor by cocaine and MK-801. *Proc Natl Acad Sci U S A* 97: 13895–13900.