

Research Article

Global Asymptotic Stability in a Class of Difference Equations

Xiaofan Yang, Limin Cui, Yuan Yan Tang, and Jianqiu Cao

Received 29 April 2007; Accepted 5 November 2007

Recommended by John R. Graef

We study the difference equation $x_n = [(f \times g_1 + g_2 + h)/(g_1 + f \times g_2 + h)](x_{n-1}, \dots, x_{n-r})$, $n = 1, 2, \dots$, $x_{1-r}, \dots, x_0 > 0$, where $f, g_1, g_2 : (R_+)^r \rightarrow R_+$ and $h : (R_+)^r \rightarrow [0, +\infty)$ are all continuous functions, and $\min_{1 \leq i \leq r} \{u_i, 1/u_i\} \leq f(u_1, \dots, u_r) \leq \max_{1 \leq i \leq r} \{u_i, 1/u_i\}$, $(u_1, \dots, u_r)^T \in (R_+)^r$. We prove that this difference equation admits $c = 1$ as the globally asymptotically stable equilibrium. This result extends and generalizes some previously known results.

Copyright © 2007 Xiaofan Yang et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Ladas [1] suggested investigating the nonlinear difference equation

$$x_n = \frac{x_{n-1} + x_{n-2}x_{n-3}}{x_{n-1}x_{n-2} + x_{n-3}}, \quad n = 1, 2, \dots, \quad x_{-2}, x_{-1}, x_0 > 0. \quad (1.1)$$

Since then, it has been proved that $c = 1$ is the common globally asymptotically stable equilibrium of this difference equation and all of the following difference equations (where a and b are nonnegative constants):

$$x_n = \frac{x_{n-2} + x_{n-1}x_{n-3}}{x_{n-1}x_{n-2} + x_{n-3}}, \quad n = 1, 2, \dots, \quad x_{-2}, x_{-1}, x_0 > 0 \quad (\text{see [1]}), \quad (1.2)$$

$$x_n = \frac{x_{n-1}x_{n-2} + x_{n-3} + a}{x_{n-1} + x_{n-2}x_{n-3} + a}, \quad n = 1, 2, \dots, \quad x_{-2}, x_{-1}, x_0 > 0 \quad (\text{see [6, 12]}), \quad (1.3)$$

$$x_n = \frac{x_{n-2} + x_{n-1}x_{n-3} + a}{x_{n-1}x_{n-2} + x_{n-3} + a}, \quad n = 1, 2, \dots, \quad x_{-2}, x_{-1}, x_0 > 0 \quad (\text{see [6]}), \quad (1.4)$$

2 Advances in Difference Equations

$$x_n = \frac{x_{n-1} + x_{n-2}x_{n-3} + a}{x_{n-1}x_{n-2} + x_{n-3} + a}, \quad n = 1, 2, \dots, \quad x_{-2}, x_{-1}, x_0 > 0 \quad (\text{see [14]}), \quad (1.5)$$

$$x_n = \frac{x_{n-1}x_{n-2} + x_{n-3} + a}{x_{n-2} + x_{n-1}x_{n-3} + a}, \quad n = 1, 2, \dots, \quad x_{-2}, x_{-1}, x_0 > 0 \quad (\text{see [14]}), \quad (1.6)$$

$$x_n = \frac{x_{n-1}^b x_{n-3} + x_{n-4}^b + a}{x_{n-1}^b + x_{n-3}x_{n-4}^b + a}, \quad n = 1, 2, \dots, \quad x_{-3}, x_{-2}, x_{-1}, x_0 > 0 \quad (\text{see [7]}), \quad (1.7)$$

$$x_n = \frac{x_{n-k}^b x_{n-m} + x_{n-l}^b + a}{x_{n-k}^b + x_{n-m}x_{n-l}^b + a}, \quad n = 1, 2, \dots, \quad x_{1-\max\{k,m,l\}}, \dots, x_0 > 0 \quad (\text{see [8]}). \quad (1.8)$$

Motivated by the above work and the work by Sun and Xi [2], this article addresses the difference equation

$$x_n = \left[\frac{f \times g_1 + g_2 + h}{g_1 + f \times g_2 + h} \right] (x_{n-1}, \dots, x_{n-r}), \quad n = 1, 2, \dots, \quad x_{1-r}, \dots, x_0 > 0, \quad (1.9)$$

where $f, g_1, g_2 : (R_+)^r \rightarrow R_+$ and $h : (R_+)^r \rightarrow [0, +\infty)$ are all continuous functions, and

$$\min_{1 \leq i \leq r} \{u_i, 1/u_i\} \leq f(u_1, \dots, u_r) \leq \max_{1 \leq i \leq r} \{u_i, 1/u_i\}, \quad (u_1, \dots, u_r)^T \in (R_+)^r. \quad (1.10)$$

It can be seen that (1.9) subsumes (1.1) and (1.8). For example, if we let $r = \max\{k, l, m\}$, $f(x_{n-1}, \dots, x_{n-r}) = x_{n-m}$, $g_1(x_{n-1}, \dots, x_{n-r}) = x_{n-k}^b$, $g_2(x_{n-1}, \dots, x_{n-r}) = x_{n-l}^b$, and $h(x_1, \dots, x_r) \equiv a$, then (1.9) reduces to (1.8).

We prove that (1.9) admits $c = 1$ as the globally asymptotically stable equilibrium. As a consequence, our result includes all of the above-mentioned results.

2. Preliminary knowledge

For two functions, $f(x_1, \dots, x_n)$ and $g(x_1, \dots, x_n)$, we adopt the following notations:

$$\begin{aligned} [f + g](x_1, \dots, x_n) &:= f(x_1, \dots, x_n) + g(x_1, \dots, x_n), \\ [f \times g](x_1, \dots, x_n) &:= f(x_1, \dots, x_n) \times g(x_1, \dots, x_n), \\ \left[\frac{f}{g} \right](x_1, \dots, x_n) &:= \frac{f(x_1, \dots, x_n)}{g(x_1, \dots, x_n)} \quad \text{if } g(x_1, \dots, x_n) \neq 0. \end{aligned} \quad (2.1)$$

Let R_+ denote the whole set of positive real numbers. The part metric (or Thompson's metric) [3, 4] is a metric defined on $(R_+)^r$ in the following way: for any $X = (x_1, \dots, x_r)^T \in (R_+)^r$ and

$$Y = (y_1, \dots, y_r)^T \in (R_+)^r, \quad p(X, Y) := -\log_2 \min_{1 \leq i \leq r} \{x_i/y_i, y_i/x_i\}. \quad (2.2)$$

THEOREM 2.1 (see [5, Theorem 2.2], see also [3]). Let $T : (R_+)^r \rightarrow (R_+)^r$ be a continuous mapping with an equilibrium $C \in (R_+)^r$. Consider the following difference equation:

$$X_n = T(X_{n-1}), \quad n = 1, 2, \dots, \quad X_0 \in (R_+)^r. \quad (2.3)$$

Suppose there is a positive integer k such that $p(T^k(X), C) < p(X, C)$ holds for all $X \neq C$. Then C is globally asymptotically stable.

THEOREM 2.2 (see [6, page 1]). Let $a_1, \dots, a_n, b_1, \dots, b_n, c_1, \dots, c_n$ be positive numbers. Then

$$\min \left\{ \frac{a_i}{b_i} : 1 \leq i \leq n \right\} \leq \frac{\sum_{i=1}^n c_i a_i}{\sum_{i=1}^n c_i b_i} \leq \max \left\{ \frac{a_i}{b_i} : 1 \leq i \leq n \right\}. \quad (2.4)$$

Moreover, one of the two equalities holds if and only if $a_1/b_1 = a_2/b_2 = \dots = a_n/b_n$.

3. Main result

The main result of this article is the following.

THEOREM 3.1. Consider the difference equation

$$x_n = \left[\frac{f \times g_1 + g_2 + h}{g_1 + f \times g_2 + h} \right] (x_{n-1}, \dots, x_{n-r}), \quad n = 1, 2, \dots, \quad x_{1-r}, \dots, x_0 > 0, \quad (3.1)$$

where $f, g_1, g_2 : (R_+)^r \rightarrow R_+$ and $h : (R_+)^r \rightarrow [0, +\infty)$ are all continuous functions, and

$$\min_{1 \leq i \leq r} \{u_i, 1/u_i\} \leq f(u_1, \dots, u_r) \leq \max_{1 \leq i \leq r} \{u_i, 1/u_i\}, \quad (u_1, \dots, u_r)^T \in (R_+)^r. \quad (3.2)$$

Let $\{x_n\}$ be a solution of (3.1). Then the following assertions hold:

(i) for all $n \geq 1$ and $j \geq 0$, one has

$$\min_{1 \leq i \leq r} \{x_{n-i}, 1/x_{n-i}\} \leq x_{n+j} \leq \max_{1 \leq i \leq r} \{x_{n-i}, 1/x_{n-i}\}; \quad (3.3)$$

- (ii) there exist $n \geq 1$ and $j \geq 0$ such that one of the two equalities in chain (3.3) holds if and only if $(x_{n-1}, \dots, x_{n-r}) = (1, \dots, 1)$;
- (iii) $c = 1$ is the globally asymptotically stable equilibrium of (3.1).

Proof. (i) For any given $n \geq 1$, we prove the assertion by induction on j . By Theorem 2.2 and chain (3.3), we have

$$\begin{aligned}
 x_n &= \left[\frac{f \times g_1 + g_2 + h}{g_1 + f \times g_2 + h} \right] (x_{n-1}, \dots, x_{n-r}) \geq \min \left\{ f(x_{n-1}, \dots, x_{n-r}), \frac{1}{f(x_{n-1}, \dots, x_{n-r})} \right\} \\
 &\geq \min \left\{ \min_{1 \leq i \leq r} \{x_{n-i}, 1/x_{n-i}\}, \frac{1}{\max_{1 \leq i \leq r} \{x_{n-i}, 1/x_{n-i}\}} \right\} = \min_{1 \leq i \leq r} \{x_{n-i}, 1/x_{n-i}\}, \\
 x_n &= \left[\frac{f \times g_1 + g_2 + h}{g_1 + f \times g_2 + h} \right] (x_{n-1}, \dots, x_{n-r}) \leq \max \left\{ f(x_{n-1}, \dots, x_{n-r}), \frac{1}{f(x_{n-1}, \dots, x_{n-r})} \right\} \\
 &\leq \max \left\{ \max_{1 \leq i \leq r} \{x_{n-i}, 1/x_{n-i}\}, \frac{1}{\max_{1 \leq i \leq r} \{x_{n-i}, 1/x_{n-i}\}} \right\} = \max_{1 \leq i \leq r} \{x_{n-i}, 1/x_{n-i}\}.
 \end{aligned} \tag{3.4}$$

So the assertion is true for $j = 0$.

Suppose the assertion is true for all integer k ($0 \leq k \leq j-1$), that is,

$$\min_{1 \leq i \leq r} \{x_{n-i}, 1/x_{n-i}\} \leq x_{n+k} \leq \max_{1 \leq i \leq r} \{x_{n-i}, 1/x_{n-i}\}, \quad 0 \leq k \leq j-1. \tag{3.5}$$

By Theorem 2.2, chain (3.2), and the inductive hypothesis, we get

$$\begin{aligned}
 x_{n+j} &= \left[\frac{f \times g_1 + g_2 + h}{g_1 + f \times g_2 + h} \right] (x_{n+j-1}, \dots, x_{n+j-r}) \\
 &\geq \min \left\{ f(x_{n+j-1}, \dots, x_{n+j-r}), \frac{1}{f(x_{n+j-1}, \dots, x_{n+j-r})} \right\} \\
 &\geq \min \left\{ \min_{1 \leq i \leq r} \{x_{n+j-i}, 1/x_{n+j-i}\}, \frac{1}{\max_{1 \leq i \leq r} \{x_{n+j-i}, 1/x_{n+j-i}\}} \right\} = \min_{1 \leq i \leq r} \{x_{n+j-i}, 1/x_{n+j-i}\} \\
 &\geq \min \left\{ \min_{1 \leq i \leq r} \{x_{n-i}, 1/x_{n-i}\}, \frac{1}{\max_{1 \leq i \leq r} \{x_{n-i}, 1/x_{n-i}\}} \right\} = \min_{1 \leq i \leq r} \{x_{n-i}, 1/x_{n-i}\};
 \end{aligned} \tag{3.6}$$

$$\begin{aligned}
 x_{n+j} &= \left[\frac{f \times g_1 + g_2 + h}{g_1 + f \times g_2 + h} \right] (x_{n+j-1}, \dots, x_{n+j-r}) \\
 &\leq \max \left\{ f(x_{n+j-1}, \dots, x_{n+j-r}), \frac{1}{f(x_{n+j-1}, \dots, x_{n+j-r})} \right\} \\
 &\leq \max \left\{ \max_{1 \leq i \leq r} \{x_{n+j-i}, 1/x_{n+j-i}\}, \frac{1}{\min_{1 \leq i \leq r} \{x_{n+j-i}, 1/x_{n+j-i}\}} \right\} = \max_{1 \leq i \leq r} \{x_{n+j-i}, 1/x_{n+j-i}\} \\
 &\leq \max \left\{ \max_{1 \leq i \leq r} \{x_{n-i}, 1/x_{n-i}\}, \frac{1}{\min_{1 \leq i \leq r} \{x_{n-i}, 1/x_{n-i}\}} \right\} = \max_{1 \leq i \leq r} \{x_{n-i}, 1/x_{n-i}\}.
 \end{aligned} \tag{3.7}$$

Thus the assertion is true for j . The inductive proof of this assertion is complete.

(ii) The sufficiency follows immediately from the first assertion of this theorem. *Necessity.* Suppose there exist $n \geq 1$ and $j \geq 0$ such that $x_{n+j} = \min_{1 \leq i \leq r} \{x_{n-i}, 1/x_{n-i}\}$. Then all of the equalities in chain (3.6) hold. This chain of equalities plus Theorem 2.2 yield $f(x_{n+j-1}, \dots, x_{n+j-r}) = 1$ and, hence, $(x_{n-1}, \dots, x_{n-r}) = (1, \dots, 1)$. Likewise, one can show that $(x_{n-1}, \dots, x_{n-r}) = (1, \dots, 1)$ if $x_{n+j} = \max_{1 \leq i \leq r} \{x_{n-i}, 1/x_{n-i}\}$.

(iii) The system of first-order difference equations associated with (3.1) is

$$Y_n = T(Y_{n-1}), \quad n = 1, 2, \dots, \quad (3.8)$$

where $T : (R_+)^r \rightarrow (R_+)^r$ is a mapping defined by

$$T((y_1, \dots, y_r)^T) = \left(y_2, \dots, y_r, \left[\frac{f \times g_1 + g_2 + h}{g_1 + f \times g_2 + h} \right] (y_r, \dots, y_1) \right)^T. \quad (3.9)$$

By chain (3.2), we have $f(1, \dots, 1) = 1$. Hence, $C = (1, \dots, 1)^T$ is an equilibrium of system (3.8). Consider an arbitrary $X = (x_1, \dots, x_r)^T \in (R_+)^r$, $X \neq C$. Then

$$T^r((x_1, \dots, x_r)^T) = (x_{r+1}, \dots, x_{2r})^T, \quad (3.10)$$

where $x_j = [(f \times g_1 + g_2 + h)/(g_1 + f \times g_2 + h)](x_{j-1}, \dots, x_{j-r})$, $r+1 \leq j \leq 2r$. By the first two assertions of this theorem, we induce

$$\min_{r+1 \leq i \leq 2r} \{x_i, 1/x_i\} > \min \left\{ \min_{1 \leq i \leq r} \{x_i, 1/x_i\}, \frac{1}{\max_{1 \leq i \leq r} \{x_i, 1/x_i\}} \right\} = \min_{1 \leq i \leq r} \{x_i, 1/x_i\}. \quad (3.11)$$

Hence,

$$p(T^r(X), C) = -\log_2 \min_{r+1 \leq i \leq 2r} \{x_i, 1/x_i\} < -\log_2 \min_{1 \leq i \leq r} \{x_i, 1/x_i\} = p(X, C). \quad (3.12)$$

By Theorem 2.1, we conclude that C is the globally asymptotically stable equilibrium of system (3.8). This implies that $c = 1$ is the globally asymptotically stable equilibrium of (3.1). \square

4. Applications

Example 4.1. Consider the difference equation

$$x_n = \left[\frac{f \times g_1 + g_2 + h}{g_1 + f \times g_2 + h} \right] (x_{n-1}, \dots, x_{n-r}), \quad n = 1, 2, \dots, \quad x_{1-r}, \dots, x_0 > 0, \quad (4.1)$$

where $g_1, g_2 : (R_+)^r \rightarrow R_+$ and $h : (R_+)^r \rightarrow [0, +\infty)$ are all continuous functions, $1 \leq p \leq r$, $1 \leq q \leq r$, $1 \leq s \leq r$, $f(u_1, \dots, u_r) = (u_p + u_q + u_s)/3$, and $(u_1, \dots, u_r)^T \in (R_+)^r$.

As $f(u_1, \dots, u_r)$ is the arithmetic mean of u_p, u_q , and u_s , we get

$$\begin{aligned} f(u_1, \dots, u_r) &\leq \max \{u_p, u_q, u_s\} \leq \max_{1 \leq i \leq r} \{u_i, 1/u_i\}, \\ f(u_1, \dots, u_r) &\geq \min \{u_p, u_q, u_s\} \geq \min_{1 \leq i \leq r} \{u_i, 1/u_i\}. \end{aligned} \quad (4.2)$$

By Theorem 3.1, $c = 1$ is the globally asymptotically stable equilibrium of (4.1).

Example 4.2. Consider the difference equation

$$x_n = \left[\frac{f \times g_1 + g_2 + h}{g_1 + f \times g_2 + h} \right] (x_{n-1}, \dots, x_{n-r}), \quad n = 1, 2, \dots, \quad x_{1-r}, \dots, x_0 > 0, \quad (4.3)$$

where $g_1, g_2 : (R_+)^r \rightarrow R_+$ and $h : (R_+)^r \rightarrow [0, +\infty)$ are all continuous functions, $1 \leq p \leq r$, $1 \leq q \leq r$, $1 \leq s \leq r$, $f(u_1, \dots, u_r) = (u_p + u_q + 1/u_s)/3$, and $(u_1, \dots, u_r)^T \in (R_+)^r$.

As $f(u_1, \dots, u_r)$ is the arithmetic mean of u_p , u_q , and $1/u_s$, we get

$$\begin{aligned} f(u_1, \dots, u_r) &\leq \max \{u_p, u_q, 1/u_s\} \leq \max_{1 \leq i \leq r} \{u_i, 1/u_i\}, \\ f(u_1, \dots, u_r) &\geq \min \{u_p, u_q, 1/u_s\} \geq \min_{1 \leq i \leq r} \{u_i, 1/u_i\}, \end{aligned} \quad (4.4)$$

By Theorem 3.1, $c = 1$ is the globally asymptotically stable equilibrium of (4.3).

Example 4.3. Consider the difference equation

$$x_n = \left[\frac{f \times g_1 + g_2 + h}{g_1 + f \times g_2 + h} \right] (x_{n-1}, \dots, x_{n-r}), \quad n = 1, 2, \dots, \quad x_{1-r}, \dots, x_0 > 0, \quad (4.5)$$

where $g_1, g_2 : (R_+)^r \rightarrow R_+$ and $h : (R_+)^r \rightarrow [0, +\infty)$ are all continuous functions, $1 \leq p \leq r$, $1 \leq q \leq r$, $1 \leq s \leq r$, $f(u_1, \dots, u_r) = \sqrt[3]{u_p u_q / u_s}$, and $(u_1, \dots, u_r)^T \in (R_+)^r$.

As $f(u_1, \dots, u_r)$ is the geometric mean of u_p , u_q , and $1/u_s$, we get

$$\begin{aligned} f(u_1, \dots, u_r) &\leq \max \{u_p, u_q, 1/u_s\} \leq \max_{1 \leq i \leq r} \{u_i, 1/u_i\}, \\ f(u_1, \dots, u_r) &\geq \min \{u_p, u_q, 1/u_s\} \geq \min_{1 \leq i \leq r} \{u_i, 1/u_i\}. \end{aligned} \quad (4.6)$$

By Theorem 3.1, $c = 1$ is the globally asymptotically stable equilibrium of (4.5).

5. Conclusions

This article has studied the global asymptotic stability of a class of difference equations. The result obtained extends and generalizes some previous results. We are attempting to apply the technique used in this article to deal with other generic difference equations which include some well-studied difference equations such as those in [7, 8].

Acknowledgments

The authors are grateful to the anonymous reviewers for their valuable comments and suggestions. This work is supported by Natural Science Foundation of China (Grant no. 10771227), Program for New Century Excellent Talent of Educational Ministry of China (Grant no. NCET-05-0759), Doctorate Foundation of Educational Ministry of China (Grant no. 20050611001), and Natural Science Foundation of Chongqing (Grants no. CSTC 2006BB2231, 2005BB2191).

References

- [1] G. Ladas, "Open problems and conjectures," *Journal of Difference Equations and Applications*, vol. 2, no. 4, pp. 449–452, 1996.
- [2] T. Sun and H. Xi, "Global attractivity for a family of nonlinear difference equations," *Applied Mathematics Letters*, vol. 20, no. 7, pp. 741–745, 2007.
- [3] N. Kruse and T. Nesemann, "Global asymptotic stability in some discrete dynamical systems," *Journal of Mathematical Analysis and Applications*, vol. 235, no. 1, pp. 151–158, 1999.
- [4] T. Nesemann, "Positive nonlinear difference equations: some results and applications," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 47, no. 7, pp. 4707–4717, 2001.
- [5] X. Yang, D. J. Evans, and G. M. Megson, "Global asymptotic stability in a class of Putnam-type equations," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 64, no. 1, pp. 42–50, 2006.
- [6] J. Kuang, *Applied Inequalities*, Shandong Science and Technology Press, Jinan, China, 2004.
- [7] K. S. Berenhaut and S. Stević, "The global attractivity of a higher order rational difference equation," *Journal of Mathematical Analysis and Applications*, vol. 326, no. 2, pp. 940–944, 2007.
- [8] H. Xi and T. Sun, "Global behavior of a higher-order rational difference equation," *Advances in Difference Equations*, vol. 2006, Article ID 27637, 7 pages, 2006.

Xiaofan Yang: College of Computer Science, Chongqing University, Chongqing 400044, China;
 School of Computer and Information, Chongqing Jiaotong University, Chongqing 400074, China
Email address: xf_yang1964@yahoo.com

Limin Cui: Department of Computer Science, Hong Kong Baptist University, Kowloon, Hong Kong
Email address: clm628@hotmail.com

Yuan Yan Tang: College of Computer Science, Chongqing University, Chongqing 400044, China
Current address: Department of Computer Science, Hong Kong Baptist University,
 Kowloon, Hong Kong
Email address: yytang@cqu.edu.cn

Jianqiu Cao: School of Computer and Information, Chongqing Jiaotong University,
 Chongqing 400074, China
Email address: caojq86@cquc.edu.cn

Special Issue on Modeling Experimental Nonlinear Dynamics and Chaotic Scenarios

Call for Papers

Thinking about nonlinearity in engineering areas, up to the 70s, was focused on intentionally built nonlinear parts in order to improve the operational characteristics of a device or system. Keying, saturation, hysteretic phenomena, and dead zones were added to existing devices increasing their behavior diversity and precision. In this context, an intrinsic nonlinearity was treated just as a linear approximation, around equilibrium points.

Inspired on the rediscovering of the richness of nonlinear and chaotic phenomena, engineers started using analytical tools from “Qualitative Theory of Differential Equations,” allowing more precise analysis and synthesis, in order to produce new vital products and services. Bifurcation theory, dynamical systems and chaos started to be part of the mandatory set of tools for design engineers.

This proposed special edition of the *Mathematical Problems in Engineering* aims to provide a picture of the importance of the bifurcation theory, relating it with nonlinear and chaotic dynamics for natural and engineered systems. Ideas of how this dynamics can be captured through precisely tailored real and numerical experiments and understanding by the combination of specific tools that associate dynamical system theory and geometric tools in a very clever, sophisticated, and at the same time simple and unique analytical environment are the subject of this issue, allowing new methods to design high-precision devices and equipment.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

José Roberto Castilho Piqueira, Telecommunication and Control Engineering Department, Polytechnic School, The University of São Paulo, 05508-970 São Paulo, Brazil; piqueira@lac.usp.br

Elbert E. Neher Macau, Laboratório Associado de Matemática Aplicada e Computação (LAC), Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil ; elbert@lac.inpe.br

Celso Grebogi, Center for Applied Dynamics Research, King's College, University of Aberdeen, Aberdeen AB24 3UE, UK; grebogi@abdn.ac.uk