

 Open access • Journal Article • DOI:10.1038/NG.733

Resequencing of positional candidates identifies low frequency IL23R coding variants protecting against inflammatory bowel disease — [Source link](#)

Yukihide Momozawa, Myriam Mni, Kayo Nakamura, Wouter Coppieters ...+25 more authors

Institutions: University of Liège, Linköping University, Université libre de Bruxelles, Catholic University of Leuven ...+7 more institutions

Published on: 01 Jan 2011 - Nature Genetics (Nature Publishing Group)

Topics: Common disease-common variant, Genome-wide association study, Inflammatory bowel disease, NOD2 and Medical genetics

Related papers:

- [Deep resequencing of GWAS loci identifies independent rare variants associated with inflammatory bowel disease.](#)
- [Genome-wide meta-analysis increases to 71 the number of confirmed Crohn's disease susceptibility loci](#)
- [A Genome-Wide Association Study Identifies IL23R as an Inflammatory Bowel Disease Gene](#)
- [Rare variants of IFIH1, a gene implicated in antiviral responses, protect against type 1 diabetes.](#)
- [Host-microbe interactions have shaped the genetic architecture of inflammatory bowel disease](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/resequencing-of-positional-candidates-identifies-low-87t40kslzo>

1 **Resequencing of positional candidates identifies low frequency *IL23R* coding variants**
2 **protecting against inflammatory bowel disease.**

3 *Yukihide Momozawa*¹, *Myriam Mni*¹, *Kayo Nakamura*¹, *Wouter Coppieters*¹, *Sven Almer*², *Leila*
4 *Amininejad*³, *Isabelle Cleyne*⁴, *Jean-Frédéric Colombel*⁵, *Peter de Rijk*⁶, *Olivier Dewit*⁷, *Yigael Finkel*⁸,
5 *Miquel A. Gassull*⁹, *Dirk Goossens*⁶, *Debby Laukens*¹⁰, *Marc Lémann*¹¹, *Cécile Libiouille*¹, *Colm*
6 *O'Morain*¹², *Catherine Reenaers*¹³, *Paul Rutgeerts*⁴, *Curt Tysk*¹⁴, *Diana Zelenika*¹⁵, *Mark Lathrop*¹⁵,
7 *Jurgen Del-Favero*⁶, *Jean-Pierre Hugot*¹⁶, *Martine de Vos*¹⁰, *Denis Franchimont*³, *Severine Vermeire*⁴,
8 *Edouard Louis*¹³ & *Michel Georges*¹.

9 ¹Unit of Animal Genomics, GIGA-R & Faculty of Veterinary Medicine, University of Liège (B34), 1
10 Avenue de l'Hôpital, 4000-Liège, Belgium. ²Division of Gastroenterology and Hepatology, IMK
11 Linköpings Universitet, Linköping, Sweden. ³Department of Gastroenterology, Erasme Hospital, ULB,
12 1070 Brussels, Belgium. ⁴Department of Pathophysiology, Gastroenterology Section, Catholic
13 University of Leuven, Leuven, Belgium. ⁵Registre EPIMAD, Hôpital Calmette, Lille, France. ⁶Applied
14 Molecular Genomics, Department of Molecular Genetics, VIB, University of Antwerp,
15 Universiteitsplein 1, 2610 Antwerp, Belgium. ⁷Department of Gastroenterology, Clinique
16 Universitaire St Luc, UCL, Brussels, Belgium. ⁸Department of Gastroenterology, Karolinska Children's
17 Hospital, Stockholm, Sweden. ⁹Gastroenterology Department, Hospital Universitari Germans Trias i
18 Pujol, 5^a planta, edifici general, Ctra del Canyet, s/n, 08916 Badalona, Spain. ¹⁰Department
19 Gastroenterology, University Hospital, Ghent University, 9000 Gent, Belgium. ¹¹Department of
20 Gastroenterology, AP-HP, Hôpital Saint-Louis, Université Paris Diderot Paris-VII, Paris, France.
21 ¹²Adelaide and Meath Hospital, Dublin, Ireland. ¹³Unit of Hepato-gastroenterology, GIGA-R & Faculty
22 of Medicine, University of Liège (B34), 1 Avenue de l'Hôpital, 4000-Liège, Belgium. ¹⁴Department of
23 Gastroenterology, Örebro Medical Center Hospital, Örebro, Sweden. ¹⁵Centre National de
24 Génotypage, Evry, France. ¹⁶INSERM U843, Hopital Robert Debré, 75019 Paris, France.

25

26 Correspondence: michel.georges@ulg.ac.be

1 **Genome-wide association studies (GWAS) have identified tens of risk loci for many complex**
2 **disorders including Crohn's disease (CD)^{1,2}. However, common disease-associated SNPs explain at**
3 **most ~20% of genetic variance. Several factors may account for the missing heritability³⁻⁵,**
4 **including rare risk variants not adequately tagged in GWAS⁶⁻⁸. That rare susceptibility variants**
5 **indeed contribute to the variation of multifactorial phenotypes has been demonstrated for**
6 **colorectal cancer⁹, plasma levels of HDL cholesterol¹⁰, blood pressure¹¹, type I diabetes¹²,**
7 **hypertriglyceridemia¹³ and - in the case of CD – for the *NOD2* gene^{14,15}. We herein describe the use**
8 **of high-throughput resequencing of DNA pools to search for rare coding variants influencing**
9 **susceptibility to CD in 63 GWAS-identified positional candidate genes. We report low frequency**
10 **coding variants conferring protection against inflammatory bowel disease (IBD) in the *IL23R* gene,**
11 **yet conclude that rare coding variants in positional candidates don't make a large contribution to**
12 **inherited predisposition to CD.**

13 A meta-analysis of three GWAS resulted in the identification of 30 significant and 10 suggestive
14 susceptibility loci for CD². The average confidence interval was 233 Kb (range: 20 to 1,140)
15 encompassing 4.1 genes (range: 0 to 37) for a total of 153 positional candidates (Supplemental Table
16 1). We decided to sequence the open reading frame (ORF) and intron-exon boundaries of the 51
17 genes mapping to loci containing between one and five genes. For loci with more than six
18 candidates, we retained 15 genes that mapped to significant networks identified when analyzing all
19 candidates with Ingenuity Pathways (v8.5) (Supplemental Table 2). To these 66 genes, we added the
20 *SLC22A4* candidate¹⁶, as well as *PTGER4*, *ORMDL3* and *GSDMB* on the basis of reported cis-eQTL
21 effects^{2,17}. The list of 70 selected genes is provided in Supplemental Table 3.

22 After extensive optimization (cfr. Supplemental note 1), we selected a protocol involving (i)
23 constitution of equimolar pools of genomic DNA from sets of 32 cases or controls, (ii) amplification,
24 using Phusion Hot Start High Fidelity DNA Polymerase (Finnzymes Oy), of the 70 targeted ORFs and
25 intron-exon boundaries as a series of 1,045 amplicons averaging 222 bp (range: 136-337 bp) (iii)
26 equimolar pooling of ~300 amplicons, (iv) massive parallel pyrosequencing using the Roche FLX

1 system¹⁸ targeting an average sequence depth of 500 for both the Watson and Crick (W&C) strands,
2 (v) detection of DNA sequence variants (DSV) using the Amplicon Variant Analyzer (AVA) software
3 (Roche) augmented with custom-made scripts (Methods).

4 We opted for a staged design in which all 70 candidate genes would first (stage I) be sequenced on
5 112 cases and 112 controls. This provides 98.5% and 73.3% nominal power ($p \leq 0.05$) to detect the
6 12% and 7% excess of rare *NOD2* variants reported by Hugot *et al.*¹⁴ and Lesage *et al.*¹⁵ respectively.
7 The most promising genes would be further evaluated on additional pools of cases and controls
8 (stage II). To increase the impact of genetic effects other than *NOD2*, the 112 stage I cases did not
9 carry either of three known *NOD2* susceptibility variants (*p.R702W*, *p.G908R* and *p.A1007fs*). To
10 avoid subtle stratification, the corresponding 112 controls underwent the same selection. All
11 analyzed cases and controls were of European descent.

12 92.9% of amplicons, corresponding to 63/70 genes (Supplemental Table 3) and 108.3 kb, could be
13 sequenced with coverage ≥ 200 for both W&C strand in at least one case and one control pool.
14 Simulations indicate that this coverage provides $\geq 83.4\%$ power to detect singletons (i.e. one variant
15 chromosome in DNA pool of 32 individuals) given the settings of the AVA software and self-imposed
16 curation filters (Supplemental Fig.1). Average sequence depth (\pm SD) of retained amplicons was
17 $1,471 \pm 849$ in cases and $1,420 \pm 822$ in controls (Supplemental Table 3). Analysis of the flowgrams
18 yielded 372 DSV (Table 1, Supplemental Table 4). Transitions accounted for 82.5%, transversions for
19 16.1%, dinucleotide substitutions for 0.3% and indels for 1.1% of the variants. Synonymous (S)
20 variants accounted for 41.7%, missense variants for 55.9%, nonsense variants for 0.8%, in-frame
21 indels for 1.1%, and “boundary” (intronic within 2 bp from exon) (β) variants for 0.5%. DSV with
22 estimated MAF < 0.05 amounted to 78.5%, while singletons represented 50.3% of the total (Fig. 1).

23 As expected and reflecting purifying selection on (mildly) deleterious variants, the frequency
24 spectrum of non-synonymous (NS) variants was shifted towards lower frequencies when compared
25 to S variants. NS variants represented 60% of the total for MAF < 0.05 versus 40% for MAF ≥ 0.05
26 (Table 1 & Fig. 1). The high transition/transversion ratio (5.1) is thought to be due to (i) the analysis

1 of ORF, as transversions are more likely to be NS and selected against, (ii) idiosyncrasies of the
2 analyzed set of genes as their transition/transversion ratio tended to be higher than that of other
3 ORF in HapMap data, and (iii) the elimination of low frequency ($< 2.5\%$) C-A=G-T variants
4 (Supplemental Table 5 and Supplemental note 1).

5 We evaluated our protocol in terms of sensitivity (fraction of true variants called), positive predictive
6 value (PPV; fraction of true variants amongst called variants), and accuracy in estimating allelic
7 frequency, focusing first on common variants ($MAF \geq 0.05$). Analysis of the HapMap data revealed 62
8 bona fide SNPs with $MAF \geq 0.05$ that were covered by the 879 retained amplicons. Five of these (=
9 8.1%) would lie within 6 bp from a homopolymer track and ignored (Methods). The remaining 57
10 were all detected, pointing towards excellent sensitivity. The 24 called SNPs with $MAF \geq 0.05$ that
11 were not genotyped in HapMap were inventoried in dbSNP (22/24 SNPs) or confirmed by the 1,000
12 Genomes Project (www.1000genomes.org/) (2 remaining), indicating excellent PPV. To evaluate the
13 accuracy in estimating allelic frequencies we took advantage of 31 common SNPs that had been
14 genotyped on the same individuals as part of other projects. Supplemental Fig. 2 shows the
15 correlation between allelic frequency estimated from the genotyping data and read counts. The
16 regression coefficient was 0.975 and the correlation 0.993.

17 To obtain similar estimates for rare variants, we manually (Sanger sequencing on ABI3730)
18 sequenced 2,283bp of the *NOD2* ORF on the same 112 cases and 112 controls. Sanger sequencing
19 revealed 38 variants with $MAF < 0.05$. Assuming faultless Sanger sequencing, sensitivity and PPV of
20 the massive parallel resequencing protocol were 82.4% and 97.9%. Frequency estimates from read
21 counts tended to underestimate actual frequencies of rare variants (regression coefficient: 0.822;
22 correlation: 0.578) (Supplemental Fig. 2). We observed no difference in sequence depth between
23 amplicon x DNA pool combinations in which rare DSV were detected and those in which no such DSV
24 were found (Supplemental Fig. 3).

25 Having evaluated the performances of our protocol, we initiated the search for differences in
26 cumulative frequencies of rare variants ($MAF < 0.05$) between cases and controls. Statistical

1 significance of the observed differences was estimated on a gene-by-gene basis using a permutation
2 test (Methods). P-values were computed for (i) S variants, (ii) all NS+ β variants, and (iii) NS variants
3 predicted by SIFT¹⁹ to be damaging. For each gene x DSV-type combination we computed two p-
4 values corresponding, respectively, to an enrichment of rare variants in cases (i.e. risk variants), or an
5 enrichment of rare variants in controls (i.e. protective variants). Thus, we made the hypothesis that
6 disruptive variants would increase disease-risk in some genes, while decreasing disease-risk in others.
7 When applying a Bonferroni correction, none of the 63 sequenced genes showed a significant ($p <$
8 7.94×10^{-4}) enrichment of rare variants neither in cases nor in controls, whether S, NS+ β or damaging
9 (Supplemental Table 6). There was no evidence for a difference in the distribution of p-values
10 between S and NS+ β variants, whether considering variants independently or on a gene-by-gene
11 basis (Supplemental note 2). However, *NOD2* was showing the expected enrichment of rare NS
12 variants (excluding the well known *p.R702W*, *p.G908R*, and *p.A1007fs* DSV) in cases (nominal $p = 5.94$
13 $\times 10^{-3}$; rank 3).

14 We therefore decided to pursue the sequencing (stage II) of the top 20% (i.e. 12) genes on 288 to 928
15 additional cases and 288 to 1,216 additional controls, depending on intermediate results. The
16 procedure was identical to stage I: high-throughput resequencing of pooled amplicons obtained from
17 DNA pools of cases or controls (32 individuals/pool; up to 29 case and 38 control pools/gene).
18 Amplicons were appended with DNA pool-specific tags allowing simultaneous sequencing of multiple
19 DNA pools. Average sequence depth in stage II was 988 ± 512 (range: 411 – 13,506) in cases, and
20 $1,019 \pm 415$ (range: 405 – 10,414) in controls (Supplemental Fig. 1 & Supplemental Table 3). We
21 detected 2 new common and 112 new rare variants (Supplemental Table 7). We observed no
22 difference in sequence depth between amplicon x DNA pool combinations in which rare DSV were
23 detected and those in which no such DSV were found (Supplemental Fig. 3).

24 We tested for differences in cumulative frequencies of rare variants in cases and controls using the
25 same permutation test as above except that we only tested the significance of the enrichment with
26 same polarity as in stage I, i.e. enrichment either for rare risk variants in cases (*FGFR1OP*, *GSDMB*,

1 *IKZF3*, *IL1RL1*, *NOD2*, *SLC9A4*, *TNFSF8*) or rare protective variants in controls (*CCL8*, *CDKAL1*, *ENOX1*,
2 *IL23R*, *SLC22A5*). At the outcome of stage II (Table 2), one gene yielded a suggestive association
3 (*FGFR10P*; nominal $p = 0.040$; Bonferonni-corrected $p = 0.386$), and one gene yielded a significant
4 association (*IL23R*; nominal $p = 2.67 \times 10^{-3}$; Bonferonni-corrected $p = 0.0314$).
5 Closer examination of *FGFR10P* revealed that three NS variants (*p.T184I*, *p.K251N*, *p.S281P*) located
6 within 2,436 bp from each other were segregating identically across DNA pools. This strongly
7 suggested that they were in complete LD. When considering them as a single event, nominal p -
8 values dropped to 0.124 in stage I and 0.081 in stage II. Hence, *FGFR10P* was not considered for
9 further analysis.

10 The *IL23R* signal was entirely due to three variants (*p.R86Q*, *p.G149R* and *p.V362I*) with cumulative
11 frequency of 0.0052 in cases versus 0.0370 in controls in stage I and 0.0088 in cases versus 0.0193 in
12 controls in stage II (Table 3 and Supplemental Table 8). The observation of an enrichment in controls
13 of these presumably protective *IL23R* variants was consistent with the protective effect of *p.R381Q*
14 that lead to the discovery of *IL23R* by GWAS²⁰. *p.R381Q* was enriched in our controls as expected (p
15 = 8.49×10^{-9}).

16 Being low frequency rather than very rare DSV⁴ allowed targeted genotyping in independent
17 samples. We developed TaqMan assays for *p.R86Q*, *p.G149R* and *p.V362I* in addition to *p.R381Q*.
18 We first genotyped the sequenced individuals, which confirmed the enrichment of the *p.R86Q*,
19 *p.G149R* and *p.V362I* variants in controls (Table 3 and Supplemental Fig. 2). We then genotyped an
20 additional 1,565 CD patients, 2,000 controls and 3,101 familial samples (740 affected, 2,361 non-
21 affected) (stage III). All analyzed individuals were of European decent and most of them previously
22 used in GWAS replications. *p.G149R* ($p = 0.022$) and *p.V362I* ($p = 1.51 \times 10^{-3}$) were significantly
23 enriched in controls in the replication cohort while a similar trend, albeit not strictly significant, was
24 observed for the rarer *p.R86Q* variant ($p = 0.057$) (Table 4).

25 *p.R381Q* confers protection against ulcerative colitis (UC)²⁰ (as well as ankylosing spondylitis²¹). We
26 thus genotyped a cohort of 1,251 European decent UC patients for the same four *IL23R* DSV. Both

1 *p.R381Q* ($p = 9.03 \times 10^{-9}$) and *p.V362I* ($p = 8.31 \times 10^{-3}$) were significantly depleted in UC patients,
2 while the expected trend was observed for *p.G149R* ($p = 0.087$), but not for *p.R86Q* ($p = 0.613$) (Table
3 5).

4 We herein describe the systematic search for rare coding variants influencing inherited
5 predisposition to CD in 63 positional candidates identified by GWAS. We report three novel low
6 frequency *IL23R* variants protecting against CD: *p.R86Q*, *p.G149R* and *p.V362I*. The three same
7 variants were found to be protective in an independent study, hence strengthening our claims (M.
8 Rivas & M. Daly, personal communication). We present preliminary evidence that *p.G149R* and
9 *p.V362I* act protectively against UC as well, as would be expected from the equivalent effect of
10 *p.R381Q*.

11 As for the previously described *p.R381Q*, the novel *p.R86Q*, *p.G149R* and *p.V362I* variants are
12 assumed to be hypomorphs dampening *IL23R* signaling. *p.G149R* and *p.R381Q* affect extremely
13 conserved residues in the extracellular and intracellular domain of the receptor, respectively, and are
14 predicted by SIFT¹⁹ to be damaging. *p.R86Q* and *p.V362I*, on the contrary, affect poorly conserved
15 residues and are predicted to be “tolerated” by SIFT¹⁹ (using sequence information only) and
16 “benign” by PolyPhen²² (using sequence and structural information). Moreover, the reference *IL23R*
17 sequences of some mammals carry the *Q* and *I* residues associated with IBD in human. While we
18 cannot exclude that *p.R86Q* and *p.V362I* are enriched in cases because of their association with
19 causative variants lying outside the ORF, we consider it more parsimonious that they affect *IL23R*
20 signaling directly. Of note, relative protection conferred by the “damaging” *p.G149R* and *p.R381Q*
21 (2.98 and 2.75) tended to be higher than that conferred by the “tolerated” *p.R86Q* and *p.V362I* (2.50
22 and 1.76), and the same tendency was observed for UC.

23 Relative protection against CD conferred by the newly detected low frequency variants was ~2.4 on
24 average. Although possible overestimated (winner’s curse), this value appears considerably larger
25 than the ~1.2 relative risk conferred by the bulk of common effects detected in GWAS, and supports
26 an increase in effect size with decreasing frequency⁶. However, the newly detected variants jointly

1 explain only ~0.18% of the variance of the underlying liability, to be compared with ~0.85% for the
2 more common *p.R381Q* and *rs7517847* variants¹⁹ (Supplemental note 3). Haplotype analysis
3 indicates that protection conferred by *p.R86Q*, *p.G149R* and *p.V362I* is largely independent of the
4 more common *p.R381Q* and *rs7517847*, i.e. we provide no evidence for “synthetic association”²³ at
5 the *IL23R* locus (Supplemental note 3).

6 Although not significant when accounting for multiple testing, we obtained evidence suggesting an
7 enrichment of rare NS *NOD2* risk variants in cases in stage I, supporting previous reports^{14,15}. This
8 enrichment was not confirmed in stage II despite the sequencing of 928 cases and 992 controls. This
9 discrepancy may be related to the selection of stage I samples carrying neither of the previously
10 described *p.R702W*, *p.G908R* or *p.A1007fs* *NOD2* susceptibility variants, which were consequently
11 enriched in stage II samples. Considering stage I and II samples jointly, however, indicates that the
12 excess *NOD2* mutation load in CD cases is likely to be lower than previously assumed^{14,15}, more in line
13 with recent estimates from a similarly conducted North-American scan for rare variants (M. Rivas &
14 M. Daly, personal communication).

15 Our findings are highly reminiscent of those of Nejentsev et al.¹², who resequenced the ORF and
16 regulatory regions of ten candidates for type I diabetes in 480 cases and 480 controls and reported
17 four low frequency protective variants in *IFIH1*. These modest success rates contrast with Johansen
18 et al.¹³ who reported an enrichment of rare variants associated with hypertriglyceridemia (HTG;
19 defined as fasting plasma triglyceride concentrations above the 95th percentile) in all four
20 resequenced (438 cases and 327 controls) candidate genes from GWAS (*APOA5*, *GCKR*, *LPL*, *APOB*).
21 We performed simulations indicating that this discrepancy is more likely to result from a difference in
22 genomic architecture of the studied traits rather than from methodological ideosyncrasies (high-
23 throughput sequencing of DNA pools in two stages versus Sanger sequencing of individual samples in
24 one stage) (Supplemental note 4).

25 This study confirms the enrichment of low frequency variants (either in cases or controls) in at least
26 some genes underlying inherited predisposition to complex diseases. Our results support an increase

1 in effect size with decreasing variant frequency. However, because of their frequency rare variants
2 explain less of the heritability than their common counterparts. Achieving adequate power to reliably
3 detect low frequency variants will require resequencing of cohorts larger than in this study. This will
4 become increasingly feasible as sequencing technology continues to improve. The demonstration of
5 an enrichment of rare or low frequency variants in candidate genes could then become an effective
6 way to demonstrate the causality of candidate genes from GWAS.

7

8 **Acknowledgments**

9 This work was funded by the Walloon Region (IPSEQ, Crohn & CIBLES projects), by the Politique
10 Scientifique Fédérale (IAP GENFUNC), by the Fonds National de la Recherche Scientifique (FNRS), by
11 the Communauté Française de Belgique (ARC BIOMOD). EL is a senior research associate from the
12 FNRS. KN benefitted from a post-doctoral fellowship from the University of Liège. We are grateful to
13 Alex Kvasz, Frédéric Farnir and Denis Baurain for their help with bioinformatic analyses. We thank
14 the personnel of the Flanders Institute for Biotechnology (VIB) Genetic Service
15 Facility (www.vibgeneticservicefacility.be) and the GIGA-R genotranscriptomic platform for their
16 contribution to sequencing.

17

18 **AUTHOR CONTRIBUTIONS**

19 YM, MM, KN, LA, DG, DZ performed experiments; YM, WC, PdR, MG analyzed data; MLa, JDF
20 supervised experiments; SA, LA, JFC, OD, YF, MAG, MLé, CO, CR, PR, CT, JPH, MdV, DF, SV, EL
21 examined patients and collected samples; IC, DL, CL prepared and organized samples; YM and MG
22 wrote the manuscript.

23

24 **Figure 1:** Frequency distribution of MAF for (i) synonymous (blue), (ii) all non-synonymous (red), (iii)
25 damaging (SIFT¹⁹) non-synonymous DSV (orange).

26

1 **Methods**

2 **High-throughput pyrosequencing on Roche FLX.** Genomic DNA concentrations were determined by
3 Quant-iT PicoGreen dsDNA Reagent and Kits (Invitrogen) for the constitution of equimolar pools of
4 32 cases or controls (except one pool of 48 in stage I). Primer pairs for PCR were selected using
5 Primer 3²⁴, avoiding known SNP positions. Amplicon-specific PCR reactions were set up in 30 μ l
6 volumes containing 6 μ l of 5x Phusion HF buffer, 200 μ M of each dNTP, 0.5 μ M of each primer, and 0.6
7 U of Phusion High-Fidelity DNA Polymerase (Finnzymes Oy). Cycling conditions were 98 °C for 2 min,
8 32 cycles at 98 °C for 10 sec, 60 °C for 30 sec and 72 °C for 15 sec, followed by 72 °C for 10 min on a
9 GeneAmp PCR System 2700 thermal cycler (Applied Biosystems). PCR products were purified using
10 MultiScreen PCR _{μ 96} Filter Plates (Millipore), and quantified with the Quant-iT PicoGreen dsDNA
11 reagent and kit. Up to 300 amplicons were combined in equimolar ratios. Pooled amplicons were
12 concentrated using the Montage PCR Filter Units (Millipore) and purified using the AMPure kit
13 (Agencourt Biosciences). Final concentration and length distribution were measured using the
14 Experion DNA 1K Analysis kit (Bio-Rad) on Experion Automated Electrophoresis Station (Bio-Rad).
15 High-throughput pyrosequencing was carried out using both primer A and B on a Roche 454 Genome
16 Sequencer FLX instrument following the recommendations of the manufacturer¹⁸.

17 **DSV detection.** Image and data were processed with the Genome Sequencer FLX System Software
18 Package (Roche). DSV were extracted from sff files using the AVA software. AVA reports DSV if
19 observed at least four times and representing $\geq 0.5\%$ of the reads. From the AVA-generated list, we
20 eliminated DSV (i) unless observed on both W&C strand, (ii) with flanking DSV within 2-bp on both
21 sides, (iii) in or within 6-bp from a homopolymer ($\geq 5x$) track, (iv) corresponding to C/A or G/T
22 substitutions with frequency < 0.025 (cfr. Supplemental note 1).

23 **Testing for a differential load of rare variants in cases and controls from resequencing data.** Excess
24 load of rare variants in cases (risk variants) or controls (protective variants) was tested on a gene-by-
25 gene basis, and - within gene - by DSV variant type (S, NS+ β , damaging). Rare variants were defined
26 as DSV with MAF < 0.05 . The results were essentially unaffected by the threshold frequency used to

1 define rare variants (0.02 - 0.05) (data not shown). DSV read counts (number of reads with the
2 DSV/total number of reads for that amplicon) were converted to the closest chromosome counts (>0)
3 (number of chromosomes with the DSV/total number of chromosomes in the pool = 64) and these
4 were summed over DNA pools, separately for cases and controls. The p-value of the difference in
5 DSV chromosome counts between cases and controls was then computed using two one-tailed
6 Fisher's exact tests, one testing an excess in cases (risk), the other in controls (protective). For a
7 given gene, we then multiplied hypothesis-specific (risk and protective) p-values across rare variants,
8 to generate two gene-specific summary p-values. The statistical significance of these summary
9 statistics was estimated by permutation testing. For each rare DSV, case vs control status of mutant
10 chromosomes were assigned randomly yet accounting for the possibility that the number of
11 successfully sequenced chromosomes (i.e. DNA pools) might differ between cases and controls. The
12 same two gene-specific summary p-values were generated for 1,000,000 permutations, and the
13 significance of the p-values obtained with the real data estimated as the proportion of permutations
14 with lower, hypothesis-specific, summary p-value.

15 ***Case-control and familial association test based on individual genotypes.*** SNPs were tested on
16 individual DNA using custom TaqMan assays (Applied Biosystems). The statistical significance of the
17 difference in DSV frequency between cases and controls was estimated using one-side Fisher's exact
18 test. The familial cohort was used to evaluate the significance of the distorted segregation of DSV
19 *p.R381Q* and *p.V362I* (TDT) from heterozygous parents to affected offspring using a custom-made
20 script. As no heterozygous parents were available in the familial cohort for DSV *p.R86Q* and *p.G149R*,
21 one affected individual per family was added to the case cohort in the case-control analysis for the
22 analysis of these variants. Combining test statistics were done across resequencing, case-control and
23 TDT experiments using a permutation test akin to the one described above.

24

25 **References**

- 1 1. Altshuler, D., Daly, M.J. & Lander, E.S. Genetic mapping in human disease. *Science* **322**, 881-8
2 (2008).
- 3 2. Barrett, J.C. et al. Genome-wide association defines more than 30 distinct susceptibility loci
4 for Crohn's disease. *Nat Genet* **40**, 955-62 (2008).
- 5 3. Yang, J. et al. Common SNPs explain a large proportion of the heritability for human height.
6 *Nat Genet* **42**, 565-9 (2010).
- 7 4. Manolio, T.A. et al. Finding the missing heritability of complex diseases. *Nature* **461**, 747-53
8 (2009).
- 9 5. Kong, A. et al. Parental origin of sequence variants associated with complex diseases. *Nature*
10 **462**, 868-74 (2009).
- 11 6. Pritchard, J.K. Are rare variants responsible for susceptibility to complex diseases? *Am J Hum*
12 *Genet* **69**, 124-37 (2001).
- 13 7. Bodmer, W. & Bonilla, C. Common and rare variants in multifactorial susceptibility to
14 common diseases. *Nat Genet* **40**, 695-701 (2008).
- 15 8. Kruglyak, L. The road to genome-wide association studies. *Nat Rev Genet* **9**, 314-8 (2008).
- 16 9. Fearnhead, N.S. et al. Multiple rare variants in different genes account for multifactorial
17 inherited susceptibility to colorectal adenomas. *Proc Natl Acad Sci U S A* **101**, 15992-7 (2004).
- 18 10. Cohen, J.C. et al. Multiple rare alleles contribute to low plasma levels of HDL cholesterol.
19 *Science* **305**, 869-72 (2004).
- 20 11. Ji, W. et al. Rare independent mutations in renal salt handling genes contribute to blood
21 pressure variation. *Nat Genet* **40**, 592-9 (2008).
- 22 12. Nejentsev, S., Walker, N., Riches, D., Egholm, M. & Todd, J.A. Rare variants of IFIH1, a gene
23 implicated in antiviral responses, protect against type 1 diabetes. *Science* **324**, 387-9 (2009).
- 24 13. Johansen, C.T. et al. Excess of rare variants in genes identified by genome-wide association
25 study of hypertriglyceridemia. *Nat Genet* **42**, 684-7 (2010).

- 1 14. Hugot, J.P. et al. Association of NOD2 leucine-rich repeat variants with susceptibility to
2 Crohn's disease. *Nature* **411**, 599-603 (2001).
- 3 15. Lesage, S. et al. CARD15/NOD2 mutational analysis and genotype-phenotype correlation in
4 612 patients with inflammatory bowel disease. *Am J Hum Genet* **70**, 845-57 (2002).
- 5 16. Peltekova, V.D. et al. Functional variants of OCTN cation transporter genes are associated
6 with Crohn disease. *Nat Genet* **36**, 471-5 (2004).
- 7 17. Libioulle, C. et al. Novel Crohn disease locus identified by genome-wide association maps to a
8 gene desert on 5p13.1 and modulates expression of PTGER4. *PLoS Genet* **3**, e58 (2007).
- 9 18. Margulies, M. et al. Genome sequencing in microfabricated high-density picolitre reactors.
10 *Nature* **437**, 376-80 (2005).
- 11 19. Kumar, P., Henikoff, S. & Ng, P.C. Predicting the effects of coding non-synonymous variants
12 on protein function using the SIFT algorithm. *Nat Protoc* **4**, 1073-81 (2009).
- 13 20. Duerr, R.H. et al. A genome-wide association study identifies IL23R as an inflammatory bowel
14 disease gene. *Science* **314**, 1461-3 (2006).
- 15 21. Burton, P.R. et al. Association scan of 14,500 nonsynonymous SNPs in four diseases identifies
16 autoimmunity variants. *Nat Genet* **39**, 1329-37 (2007).
- 17 22. Ramensky, V., Bork, P. & Sunyaev, S. Human non-synonymous SNPs: server and survey.
18 *Nucleic Acids Res* **30**, 3894-900 (2002).
- 19 23. Dickson, S.P., Wang, K., Krantz, I., Hakonarson, H. & Goldstein, D.B. Rare variants create
20 synthetic genome-wide associations. *PLoS Biol* **8**, e1000294 (2010).
- 21 24. Rozen, S. & Skaletsky, H. Primer3 on the WWW for general users and for biologist
22 programmers. *Methods Mol Biol* **132**, 365-86 (2000).

23

24