



Review

Resolving the Ionotropic P2X4 Receptor Mystery Points towards a New Therapeutic Target for Cardiovascular Diseases

Bruno Bragança^{1,2} and Paulo Correia-de-Sá^{1,*} 

¹ Laboratório de Farmacologia e Neurobiologia, Center for Drug Discovery and Innovative Medicines (MedInUP), Instituto de Ciências Biomédicas de Abel Salazar, Universidade do Porto, R. Jorge Viterbo Ferreira, 228, 4050-313 Porto, Portugal; bbraganca14@hotmail.fr

² Department of Cardiology, Centro Hospitalar Tâmega e Sousa, 4564-007 Penafiel, Portugal

* Correspondence: farmacol@icbas.up.pt; Tel.: +351-220-428-212; Fax: +351-220-428-090

Received: 30 June 2020; Accepted: 13 July 2020; Published: 15 July 2020



Abstract: Adenosine triphosphate (ATP) is a primordial versatile autacoid that changes its role from an intracellular energy saver to a signaling molecule once released to the extracellular milieu. Extracellular ATP and its adenosine metabolite are the main activators of the P2 and P1 purinoceptor families, respectively. Mounting evidence suggests that the ionotropic P2X4 receptor (P2X4R) plays pivotal roles in the regulation of the cardiovascular system, yet further therapeutic advances have been hampered by the lack of selective P2X4R agonists. In this review, we provide the state of the art of the P2X4R activity in the cardiovascular system. We also discuss the role of P2X4R activation in kidney and lungs vis a vis their interplay to control cardiovascular functions and dysfunctions, including putative adverse effects emerging from P2X4R activation. Gathering this information may prompt further development of selective P2X4R agonists and its translation to the clinical practice.

Keywords: ATP; cardiovascular system; P2X4 receptor

1. The “Purinome”

The concept of “purinome” encompasses the molecular machinery necessary for purinergic signaling. The core of this system is adenine nucleotides and nucleosides and their activity over a wide range of transmembrane purinoceptors. Terms like “purinergic nerves” and “purinoceptors” were coined in the early 1970s after Geoffrey Burnstock, one of the most widely cited scientists who recently (2 June 2020) passed away at the age of 91. The purinoceptor family is further subdivided into P1 and P2 receptor subtypes according to their sensitivity to adenosine and adenine and uracil nucleotides (e.g., ATP, ADP, UTP, UDP, UDP-glucose), respectively. Four subtypes of adenosine-sensitive G protein-coupled P1 receptors have been cloned, A₁, A_{2A}, A_{2B}, and A₃. Regarding the P2 receptor family, it is sub-classified into seven ionotropic P2X receptors (P2X1-7) and eight G-protein-coupled metabotropic P2Y receptors (P2Y_{1,2,4,6,11,12,13,14}). Each purinoceptor has a specific pharmacological profile and biological fingerprint, which allows its identification overtaking the scarcity of selective pharmacological tools [1]. Besides lytic mechanisms that occur after cellular damage, the release of ATP and other purines/pyrimidines may be elicited by multiple stimuli (e.g., hypoxia, inflammation, mechanical stimuli) and occur via highly regulated processes, such as vesicle exocytosis and diffusion through plasma membrane channels [2,3]. Once in the extracellular milieu, ATP is sequentially hydrolyzed to adenosine by a cascade of plasma membrane-bound nucleotidases (e.g., NTPDase1/CD39, ecto-5'-nucleotidase/CD73), shifting cellular signaling from P2- to P1-mediated responses. Then, adenosine follows one of two removal pathways: (1) cellular reuptake by nucleoside transporters or (2) enzymatic deamination to inosine by adenosine deaminase [4].

The relevance of purinergic signaling to control cardiovascular functions is indisputable, spanning from the pioneering work of Drury and Szent-Györgyi in 1929 on the role of adenosine in the mammalian heart [5] to more recent reports leading to the use of P2Y₁₂ receptor antagonists to reduce platelet aggregation in thrombotic diseases (e.g., myocardial infarction, stroke) [6,7]. Notwithstanding this, drugs other than adenosine (and its derivatives) and P2Y₁₂ receptor antagonists have already entered clinical trials for cardiovascular diseases, with some other drugs still under development and awaiting clearance to enter the clinical trial pipeline [8].

Nowadays, there is an increasing enthusiasm for the modulation of P2X purinoceptors functions in health and disease, with the P2X7R as the most promising candidate confirmed by successfully completed clinical trials. More recently, the P2X4R has emerged as another promising drug target. Despite the lack of selective P2X4R agonists available so far, modulation of the P2X4R has been demonstrated to be beneficial in cardiac diseases, pain sensation, cancer, and inflammatory diseases [8,9]. Due to these putative therapeutic hints, we expect a boost in the development of selective P2X4R agonists in the forthcoming years. This prompted us to review the recent advances in deciphering the pathophysiological relevance of the P2X4R in the cardiovascular system. Considering that impairment of other body systems (e.g., renal and respiratory) might have a huge impact in the management and prognosis of cardiac patients, we thought it was useful to discuss the (beneficial and/or detrimental) repercussions of the P2X4R activation in these territories.

2. The P2X4 Receptor: From Molecular Characteristics to Its Modes of Action

The P2X4R was the first crystal structure of a P2X receptor to be resolved [10]. Sequence homology of P2X4R in mammals exceeds 85%, with some genetic variants been described for this receptor [11,12]. The P2X4R results from the assembly of three subunits organized in a dolphin-like structure. Each subunit has two membrane-spanning domains, with C- and N-terminals located intracellularly, and a large ectodomain for ATP binding [10]. Structural homology among P2X receptors allows different assemblies of P2X subunits. Heterotrimeric P2X receptors have different pharmacological and functional properties from homotrimeric versions [13]. Heterotrimers have been described for P2X4R, particularly in combination with P2X6R subunits; however, it remains to be clarified whether they exist in native systems [13,14].

P2X4R orthosteric binding sites can be found between two adjacent subunits, which means that each P2X4R can harbor up to three ATP molecules [9,15]; the pEC₅₀ for ATP of the Human P2X4R is in the low micromolar range (Table 1) [16]. The P2X4R ATP-binding site lacks some specificity for the purine base allowing binding of other nucleotides including non-adenine compounds, such as cytidine triphosphate (CTP); the P2X4R exhibits low affinity (millimolar range) for this pyrimidine [17]. Similarly to other P2X receptors, the P2X4R has two main conformational states: closed and open. Occupancy of the orthosteric binding pocket by ATP and other agonists induces structural changes that open a permeation pore for Na⁺, K⁺, and Ca²⁺ ions [15,18]. Opening of P2X4R generates a depolarizing inward current, with a fourfold permeability for Ca²⁺ in relation to Na⁺ ($P_{Ca}/P_{Na} \approx 4$), contributing to the activation of several Ca²⁺-dependent intracellular signaling processes [18,19]. The P2X4R reversion potential is around 0 millivolts, which means that P2X4R-mediated currents easily become outward at positive potentials [20,21]. Interestingly, two P2X4R permeant conformations seem to exist regarding the dimensions of the channel pore (small vs. large) [19,20,22]. Indeed, the P2X4R may share some permeant characteristics with the P2X7R, including the formation of a high-permeability pore that allows passage of large molecules, including large fluorescent dyes. This high-permeant state of the P2X4R exists on its own, without needing the accessory proteins (e.g., pannexin 1) required to form the high-permeability pore together with the P2X7R [22,23]. Like its P2X7R sibling, the P2X4R large pore formation is favored by low extracellular Ca²⁺ and is regulated by phosphatidylinositol 4,5-bisphosphate (PIP2) content in the plasma membrane [23,24].

Table 1. Pharmacology of the P2X4 receptor.

	Compound	EC ₅₀ or IC ₅₀ (μM) *			P2X Cross-Reactivity/Fold Selectivity	References
		Human	Rat	Mouse		
Agonist	ATP	0.7–5.7	1.7–16.2	0.3–2.3		[12,17,21,22, 25–27]
	ATPγS	10.9				[28]
	2-meSATP	0.3–2.2	~ 1–10	1.4		[21,27–29]
	CTP		~0.1–1 mM		P2X1,2,3,7	[21,30–32]
	BzATP	0.5–9.4	>100	2.9	P2X7 (h)	[12,27,28]
	α,β-meATP	0.8–19	≥100	7–>100	P2X1,3	[26–28]
	β,γ-meATP	>100	>100	>100		[27]
	AP4A [‡]	0.1–3	20–>100	2.6–>100		[26–28]
	AR-C67085MX	2.5	>100	>100		[27]
	8-Azido-ATP	>100	>100	>100		[27]
	β,γ-Imido-ATP	6.5	>100	>100		[27]
	MRS-2339					[33]
	MRS-2978					[34]
PSB-0412	2.1				[27]	
Antagonist	PPADS	9.6–>100	>300	>100		[26,28]
	Suramin	>100	>100	>100		[26,28]
	KN-62	>10	>10	>10		[26]
	TNP-ATP	1.5	1.3–4.71	1.3–4.2	P2X1,2,3,7	[27,35–38]
	Brilliant Blue G	3–100			1000-fold more potent at P2X7 (r)	[39]
	5-BDBD	0.3–1.2	0.75–3.5	2.04	10-fold P2X1 (r), 3-fold P2X3 (r)	[27,36,40]
	BX-430	0.54–1			10 to 100-fold P2X1,3,5,7	[41,42]
	Carbamazepine der. [†]	3.44	54.6	14.9	2 to 30-fold P2X1,2,3,7 (h)	[43]
	PSB-12054	0.19	2.10	1.8	≥50-fold P2X1,2,3,7 (h)	[35]
	PSB-12062	1.4	0.9	1.8	≥35-fold P2X1,2,3,7 (h)	[35]
	PSB-15417	10 [‡]			>5-fold P2X7 (h)	[44]
	NP-1815-PX	0.26			P2X1,2,3,7	[45]
	NC-2600 [‡]					[46]
	UoS14919	61 nM				[47]
	Paroxetine	1.87–4.8	1.64–2.45	0.7		[27,48]
Duloxetine	1.59–17				[49,50]	
BAY-1797	0.2				[51]	
IgG#151-LO	0.7			10,000-fold more potent over other P2X	[52]	
PAM	Ivermectin	0.1			P2X7 (h)	[53,54]
	Cibacron blue	>300				[55]
	Ginsenosides	7.5–10.5			P2X7	[56]
	Testosterone butyrate		30 [‡]		P2X2	[57]
	Zn ²⁺		1.8			[58]
	Cd ²⁺		7.3			[58]
	Alfaxalone		0.4–1.6			[59]
	Allopregnanolone		0.4			[59]
	Propofol	56				[39]
NM	Ethanol		5–200 mM [‡]		PAM at rat P2X3 inhibits P2X1,3,4; stimulates P2X2	[60,61]
	H ⁺	pKa ~6.8				[62]
	Hg ²⁺		9			[58]
	Cu ²⁺		8.6			[58]
	Fluvastatin [‡]	10 for 1 h				[63]
	Filipin III [‡]	10 for 30 min				[63]
	Methyl-β-cyclodextrin [‡]	10 mM, 1 h				[63]
t-DCA [§]	160				[64]	

Data are mostly from functional studies (e.g., electrophysiological currents, intracellular calcium oscillations) in heterologous systems expressing human (h), rat (r) or mouse (m) P2X4 receptors. * Potency of agonists and positive allosteric modulators (PAM) is represented as half maximal effective concentration (EC₅₀), whereas for antagonists and negative modulators (NM) indicated is the half maximal inhibitory concentration (IC₅₀). EC₅₀ and IC₅₀ values are in μM, unless stated otherwise. [‡] AP4A, adenosine tetraphosphate. [†] N,N-diisopropyl-5H-dibenzo[b,f]azepine-5-carboxamide. [‡] NC-2600 is a specific P2X4 antagonist from Chemifar[®], with unknown structure and selectivity, currently in phase I clinical trial for neuropathic pain. [‡] Potency not calculated, it indicates the concentration of the compound used to produce the functional effect. [‡] Activity suppressed by cholesterol depletion. [§] Tauro-deoxycholic acid (t-DCA). ^{||} Mouse antibody anti-P2X4 IgG#151-LO.

The role of the P2X receptor in several physiological processes has been hindered by the current lack of selective agonists and antagonists. This limitation has been partially overridden by the pharmacological characterization of agonist profiles, gating kinetics, and function in comparison with other P2X receptors [20]. The P2X4R is a fast purinergic excitatory receptor, and in comparison with other P2X receptors, it exhibits intermediate sensitivity for ATP ($\approx 10 \mu\text{M}$) and desensitization time course in rodents [20,22,65]. Synthetic ATP derivatives (e.g., $\alpha\beta$ -methylene ATP ($\alpha\beta$ -meATP); $\beta\gamma$ -methyleneATP ($\beta\gamma$ -meATP); adenosine 5'-O-(3-thio)triphosphate (ATP γ S), 2-MethylthioATP (2-meSATP), 2,3-O-(4-benzoyl-benzoyl)ATP (BzATP)) activate all P2X receptors, yet, contrary to other P2X receptor subtypes, the P2X4R is relatively insensitive to $\alpha\beta$ -meATP in rodents (Table 1). The agonist rank potency order for the P2X4R is as follows: ATP > ATP γ S > 2-meSATP \geq CTP > $\alpha\beta$ -meATP [20,66]. The list of available P2X4R antagonists is gradually expanding with some compounds presenting well-acceptable selectivity over other P2X receptor subtypes, like 5-BDBD (5-(3-bromophenyl)-1,3-dihydro-(1)benzofuro(3,2-e)(1,4)diazepin-2-one; pIC₅₀~5-6) [11,20,64]. Relative insensitivity to pyridoxalphosphate-6-azophenyl-2',4'-disulfonic acid (PPADS), and suramin, two broad-spectrum P2 antagonists, is another pharmacological characteristic of the P2X4R (Table 1) [20].

Regulation of the P2X4R activity is complex and not limited to binding and unbinding of ATP [9,20,67]. Trace metals are important modulators of the P2X4R activity, and their role may be used as a strategy for this receptor characterization (Table 1). For instance, zinc ions (Zn^{2+}) bind to the "M1 site" of the P2X4R and increase its affinity for ATP [66,68]. Conversely, copper ions (Cu^{2+}) inhibit the P2X4R maximal response without affecting its affinity for ATP [20]. This receptor is also sensitive to pH, a property derived from the surface lining of protonable histidines. In acidic conditions, like those occurring during ischemia or inflammation, protons (H^+) inhibit the P2X4R activity [67,69]. Allosteric modulators are other important pharmacological tools. This is headed by the prototypical positive allosteric modulator of the P2X4R, ivermectin, which highly increases the ATP agonist potency [67,70]. Other allosteric modulators have been described, which include several commonly used drugs like paroxetine, alcohol, and propofol [20,41,57,64]. The C-terminal of the P2X4R is a target for kinase regulation, namely by protein kinase A (PKA), which increases the receptor activity. Rather than a direct regulation by phosphorylation, PKA stimulation recruits accessory proteins to bind to the C-terminal of the P2X4R [71]. The P2X4R is also sensitive to the composition of plasma membrane; for instance, depletion of membrane PIP2 by stimulation of phospholipase C (PLC) decreases the current induced by the receptor and slows its recovery from desensitization [72].

Gene expression and cellular localization represent another level of regulation of the P2X4R. This receptor is present in multiple cells throughout the body, but its relative expression depends on cell types, development status, and health/disease conditions [20,21,66]. Expression of the P2X4R is highly sensitive to certain stimuli, including hypoxia [73,74] and inflammation [75], and is controlled by specific transcription factors. STAT1, GATA-2, and IRF-5 transcription factors are known activators of the P2X4R gene promoter [49,75,76]. Trafficking of the P2X4R to the plasma membrane is paramount to sense extracellular ATP, while internalization represents an important negative feedback mechanism to avoid overactivation. The C-terminal tail of the P2X4R is a key element in desensitization and internalization processes; it functions as an endocytic-sorting signal to lysosomes [77]. Intracellular localization of the P2X4R has increasingly been demonstrated in certain cell types; localization of this receptor in vesicles and lysosomes supports its role in vesicle turnover, as demonstrated concerning exocytosis [11,78,79].

3. Benefits from the P2X4 Receptor Activation in the Heart

Cardiac output is a function of heart rate and stroke volume, with the latter being influenced by pre-load, after-load, and inotropy. ATP and other purines regulate all these features. For decades it was generally accepted that intravenous infusion of ATP decreased cardiac works by means of its breakdown to adenosine via activation of the inhibitory A₁ receptor. With experimental advances, the "true"

ATP effects soon became resolved, and now there is plenty of evidence to support involvement of P2 receptors in acute and chronic physiological processes in the heart. P2 receptors are expressed not only in the working myocardium, but also in other cardiac cells [80–82]. Among P2X receptors, the P2X4R is the most expressed receptor subtype in the heart, particularly in supraventricular tissues [83,84]. The P2X4R is distributed in various cardiac cells, including cardiomyocytes, endothelial and smooth muscle cells of cardiac blood vessels, cells of the cardiac conduction system [83,85], intracardiac neurons [86], and cardiac fibroblasts [87].

ATP, through activation of P2 purinoceptors, increases cardiac inotropy. Although some conflicting results exist regarding the receptor involved in this effect, accumulating evidence suggests an involvement of P2X receptors, particularly the P2X4R subtype [81]. In 2001, a seminal study from B.T. Liang's group demonstrated that cardiac preparations overexpressing the human P2X4R exhibited increased contractile performance when challenged with the P2X receptor agonist, 2-meSATP. Remarkably, baseline cardiac output was also incremented, but no effects were observed regarding chronotropy and/or ventricular remodeling [88]. Later on, the same group showed that P2X4R overexpression was indeed protective by increasing survival both in ischemic and in non-ischemic models of heart failure. This theory was strengthened by finding that overexpression of the P2X4R favors rescuing of several hallmarks of heart failure (HF), namely ventricular systolic dysfunction, β -adrenergic desensitization, hypertrophy, and maladaptive remodeling [89–91]. Interestingly, HF and coronary artery disease are on their own associated with upregulation of the P2X4R expression in the heart [33,83]. In a rodent model of hypoxia-induced pulmonary hypertension, the P2X4R was found to be upregulated in the right ventricle [73]. Likewise, a compensatory upregulation of P2X currents was observed in the heart of animals exhibiting cardiomyopathy due to calsequestrin overexpression; in this animal model stimulation of P2X receptors with MRS-2339, a charged methanocarpa derivative of 2-Cl-AMP resulted in reversion of maladaptive cardiac remodeling and prolonged survival [33]. The recently developed diester-masked uncharged phosphonate, MRS-2978, a compound structurally related with MRS-2339 but chemically modified to be orally bioavailable and more resistant to enzymatic hydrolysis, presents benefits in improving cardiac dysfunction in both ischemic and pressure-overload HF animal models [34,53]. Conversely, knocking down the P2X4R is normally associated with deterioration of cardiac ventricular function and progression to HF [92]. Data, thus, indicate that stimulation of the P2X4R ameliorates cardiac contractile performance by increasing inotropy and by favoring the recruitment of protective signaling mechanisms.

As aforementioned, the P2X4R activation increases Ca^{2+} influx into the cells [18,19,66]. Ca^{2+} transients induced by P2X4R activation further stimulate massive Ca^{2+} release from sarcoplasmic reticulum through a mechanism of calcium-induced calcium release, which might explain the positive inotropic effect of this receptor [93]. It has also been demonstrated that the P2X4R modulates the activity of the Na^+ - Ca^{2+} exchanger (NCX), which contributes to prolonging intracellular Ca^{2+} accumulation and signaling. In ventricular myocytes from both wild-type and P2X4R-overexpressing mice, 2-meSATP caused an inward Na^+ current which was high enough (about 1 mM) to potentiate Ca^{2+} -entry via the NCX functioning in the reverse mode [94]. These findings led to the conclusion that Na^+ influx via the P2X4R may putatively interfere with normal function of NCX, thus favoring intracellular Ca^{2+} accumulation (Figure 1).

Recently, our group demonstrated that in the sinoatrial node (SAN), the first element of the cardiac conduction system, the P2X4R interferes with the Ca^{2+} -extrusion (forward) mode of the NCX [85], which constitutes a fundamental pacemaking mechanism to drive spontaneous SAN depolarization and to set sinus rhythm [95]. Our study demonstrated for the first time that rather than requiring its breakdown to adenosine, ATP on its own is able to decrease chronotropy via P2X4R activation. The dual role of the P2X4R in decreasing chronotropy while increasing inotropy strikingly resembles the effect of Na^+/K^+ -ATPase inhibition by cardiac glycosides (e.g., digoxin) [85,96], expanding the clinical usefulness of P2X4R agonists to the treatment of tachyarrhythmias and congestive HF [97–99]. In support of this mechanism, istaroxime, a newer intracellular Na^+ enhancer, has been demonstrated

to improve both diastolic and systolic function in patients with decompensated HF with reduced ejection fraction (HFrEF), along with increasing systolic blood pressure while reducing heart rate, with no evidence of increases in the probability of cardiac arrhythmias [100–103]. Notwithstanding this, a note of caution must be taken concerning intracellular Na^+ overload as a consequence of chronic P2X4R activation, since increases in diastolic Ca^{2+} concentration may worsen the diastolic dysfunction in HF conditions [104]. Another caveat is related to rhythm disturbances that may arise from perturbations of Na^+ and Ca^{2+} -handling resulting from prolonged activation of the P2X4R. As a matter of fact, some reports ascribed a pro-arrhythmic potential to ATP [105,106], which deserves to be further explored in future studies.




Figure 1. Schematic overview of systemic P2X4R stimulation. An overall cardioprotective effect is observed with P2X4R stimulation. Permeation of Na^+ and Ca^{2+} ions through P2X4R interferes with the forward mode of the Na^+ - Ca^{2+} exchanger (NCX) and Ca^{2+} -dependent activation of endothelial nitric oxide synthase (eNOS) (right panels). In the heart, stimulation of P2X4R improves cardiac muscle contraction and optimizes myocardial energy supply vs. demand by reducing chronotropy. The diuretic effect of P2X4R activation together with its vasodilation properties in both systemic and pulmonary arteries optimizes preload and afterload and, thereby, cardiac output in HF. Chronic activation of the P2X4R reverses maladaptive cardiac remodeling and improves the outcome of HF. Future P2X4R agonist therapies must deal with the wide P2X4R distribution throughout the human body resulting in putative unwanted side-effects (marked in orange). Figure composition used elements from Servier Medical Art (<https://smart.servier.com>).

Apart from its key role in cardiac contraction and heartbeat initiation, Ca^{2+} is an important intracellular second messenger [107]. Yang and coworkers demonstrated that cardiac protection triggered by the P2X4R involves stimulation of endothelial synthesis of nitric oxide (NO) [92]. Synthesis of NO is a downstream target of Ca^{2+} signaling. The beneficial effects of NO in the cardiovascular system are well-known, which include vasodilation, inhibition of platelet aggregation, and regulation Ca^{2+} homeostasis [108]. Putative reversion of pathological cardiac remodeling may also be owed to NO production (Figure 1), though the P2X4R activation has been linked to proliferation and migration of cultured human cardiac fibroblasts, but this was only reported once several years ago [87]. Clearly, further studies are also needed to clarify the role of P2X4R in cardiac remodeling.

4. Cardiovascular Risk Factors: Is the P2X4 Receptor Friend or Foe?

Hypertension, diabetes, obesity, and dyslipidemia have a close relationship with heart diseases, major cardiovascular events, and death [109]. Strict control of these modifiable cardiovascular risk factors has a major impact on patients' prognosis [30,110,111]. Not controlling these risk factors engage the cardiovascular disease continuum driven by atherosclerosis and inflammation [112].

Purinergic receptors have been targeted for the treatment of diabetes, hypertension, and other conditions associated with the metabolic syndrome [113–115]. Purines regulate multiple functions in blood vessels, ranging from short-lived mechanisms to more sustained processes, including those involved in maladaptive remodeling of hypertension and atherosclerosis [81]. The net vascular response depends on where and when the purinergic transmission occurs. Like acetylcholine (ACh), ATP has a dual response in controlling blood vessel tone, with both vasodilatory or vasoconstrictor responses [81,116]. One important source of ATP is provided from sympathetic nerves surrounding blood vessels. Stimulation of sympathetic nerves releases norepinephrine and ATP in the smooth muscle layer of blood vessels; these two potent vasoconstrictors are upregulated in hypertension [117]. ATP exerts transient vasoconstriction mostly through activation of the P2X1R on vascular smooth muscle cells (VSMCs) [118,119], followed by a smaller yet sustained contraction via P2X4R activation [120]. Both P2X1R and P2X4R localized on VSMCs contribute to vasospasm occurring as a consequence of hemorrhagic stroke [120].

When acting at the luminal side of intact blood vessels, circulating ATP causes vasodilation through activation of endothelial metabotropic P2Y₁, P2Y₂, P2Y₄, and P2Y₆ receptor subtypes; while the P2Y₁ receptor is more sensitive to the ATP metabolite, ADP, the P2Y₂, P2Y₄, and P2Y₆ receptor subtypes are mainly activated by uracil nucleotides, namely UTP and UDP, respectively. Increasing evidence shows that endothelial ATP-sensitive ionotropic P2X4R also participates in vasodilation. These receptors are particularly abundant in coronary and cerebral arteries [119]. Noteworthy is that endothelial cells release huge amounts of ATP in response to increased blood pressure/flow [121], thus creating a "self-regenerative ATP activation wave" spreading along endothelial cells in the vicinity [81]. In this sense, endothelial P2X4R may act as immediate mechano-transducers to re-establish local blood pressure/flow by promoting the endothelial release of vasodilation signaling mediators like NO [122]. This hypothesis is strengthened by findings linking hypertensive and vascular remodeling phenotypes to knockdown or loss-of-function mutations of the P2X4R [12,122]. For instance, the loss-of-function tyrosine-to-cysteine polymorphism in the 315 position of the human P2X4R ectodomain (Tyr 315 > Cys) was associated with higher blood pressure and reduced arterial compliance as a result of impaired endothelium-dependent vasodilation in large arteries [12]. Pathological alterations of renal P2X4R may also contribute to arterial hypertension (see below).

Apart from the local control, vascular tone can be also regulated centrally through cardiovascular reflexes involving carotid bodies (CBs) and aortic bodies (ABs). In CB, a cluster of chemoreceptor cells releases excitatory ATP in response to several chemostimulants such as hypoxia and hypercapnia, which then stimulates P2X_{2/3R} located in sensory afferents of the carotid sinus nerve (CSN), a branch of the petrosal nerve. Firing of the CSN will then be integrated centrally at the brainstem to increase ventilation and to cause bradycardia and peripheral vasoconstriction [123]. Feedback inhibition of CB-mediated responses involves nitrenergic nerves from the glossopharyngeal nerve (GPN). Interestingly, these inhibitory nitrenergic nerves can be stimulated by several ATP-sensitive P2X receptors, namely P2X_{2/3R}, P2X_{4R}, and P2X_{7R}, but contrary to the P2X_{2/3R} that is present both in afferent and in efferent nerves, the P2X_{4R} exclusively mediates the inhibitory pathway [123,124]. Hence, selective stimulation of the CB inhibitory drive with P2X_{4R} agonists or blockage of P2X_{2/3R}-mediated CB excitation may represent putative targets to control cardiorespiratory reflexes, particularly in patients who are refractory to common anti-hypertensive drugs [125,126]. Noteworthy, overactivation of CB is also observed in obstructive sleep apnea and type 2 diabetes and may represent a pivotal element in the vicious cycle causing drug-refractory hypertension and type 2 diabetes [126].

Atherosclerosis is a low-grade chronic inflammatory process that develops upon persistence of one or more of the aforementioned vascular risk factors [127]. ATP, acting mainly via metabotropic P2Y receptors, has potent mitogenic and pro-inflammatory effects involved in stepwise formation of atheroma plaques [81]. Evidence also exists on direct involvement of P2X receptors in atherosclerosis [81,128,129]. For instance, upregulation of endothelial P2X4R and/or P2X7R has been observed in response to cellular damage inflicted by dyslipidemia and hyperglycemia; overactivation of these receptors has been linked to the release of pro-inflammatory cytokines and increases in endothelial permeability [130]. Moreover, overexpression of the P2X4R in medial and neointima of rabbit aorta may ascribe the pro-atherosclerotic role of ATP in this vessel [131].

In primary monocyte-derived human macrophages, the P2X4R stimulates the release of CXCL5, a pro-inflammatory chemokine involved in neutrophil recruitment [132]. Nevertheless, caution must be taken in attributing a deleterious role to the P2X4R in atherosclerosis development, because increased production of endothelial CXCL5 was protective against macrophage foam cell accumulation in atherosclerotic plaques by a mechanism that involves cholesterol efflux from macrophages in a rodent model of atherosclerosis [133]. Moreover, the P2X4R was responsible for increases in Krüppel-like factor 2 (KLF2), an atheroprotective transcription factor [134].

As atherosclerotic plaques grow, the lumen of blood vessels becomes narrow, thus limiting blood flow and distal delivery of oxygen and nutrients. At this stage of atherosclerosis development, blood-flow-mediated shear stress dramatically increases along with overactivation of endothelial P2X4R (see above). In this context, activation of endothelial P2X4R engages protective mechanisms against ischemia in blood vessels where its density is higher, like coronary and cerebral arteries; this results in the increased expression of osteopontin, a well-known neuroprotective factor [135]. In the late stage of atherosclerosis, erosion and plaque rupture occurs leading to endothelial cells damage and inability to prevent thrombotic events, thus prompting acute coronary and stroke events. Endothelial damage exposes not only the procoagulant and platelet-aggregating features of subendothelial matrix proteins [6] but also the vasoconstrictive P2X1R and P2X4R on VSMCs to surplus amounts of ATP derived from platelet aggregation and damaged cells [119,120]. Thus, one may hypothesize that the P2X4R exerts a protective role in blood vessels when the endothelium is functional, but its action is detrimental once blood vessels are severely damaged, such as in acute coronary and aortic syndromes, as well as in stroke.

5. Role Played by the P2X4 Receptor in Renal Function and Dysfunction

Renal integrity and function are of utmost importance in patients with cardiovascular diseases. Acute and chronic kidney diseases have a negative impact on the prognosis of patients with HF and coronary artery diseases [136–138]. Multiple factors are involved in the cardiorenal axis regulation. These include neurohormonal mediators and the autonomic nervous system [139]. Renal purinoceptors control blood flow and glomerular filtration, which are main determinants of the water-electrolyte balance [140].

In healthy conditions, myogenic and tubule–glomerular feedback (TFG) mechanisms ensure that renal blood flow matches glomerular filtration rate (GFR) for proper water and solute handling. TFG is orchestrated by macula densa cells in distal tubules. These cells react to low NaCl levels in the lumen of distal tubules by favoring dilation of afferent arterioles and renin–angiotensin-dependent constriction of efferent arterioles (reviewed in [141]). Infusion of ATP into the renal artery exerts a dual effect in afferent arterioles that is dependent on the vascular tone. Upon increasing renal vascular resistance, ATP effects shift from P2X1R-mediated vasoconstriction to P2Y/P2X4R-induced NO-mediated vasodilation [140,142,143]. Blood filtration yields the urinary ultrafiltrate, and its composition changes as it moves throughout renal tubules. Multiple P2 receptors are expressed in renal tubules, and their activation promotes natriuresis through inhibition of several sodium transporters [144]. Interestingly, after surgical renal denervation for treatment of drug-refractory hypertension, the P2-mediated natriuresis is significantly favored [145]. Although the P2X4R is

abundantly expressed in all renal segments, its functions are better characterized in distal segments of the nephron [140]. Activation of the P2X4R in this region contributes to modulate the activity of epithelial sodium channels (ENaC) [146]. However, regulation of ENaC activity by the P2X4R is not linear, and ATP may exert a dual role depending on sodium and body fluid composition. In low-salt conditions, stimulation of the P2X4R increases ENaC activity, while in salt-overload conditions the P2X4R exerts an opposite effect and promotes natriuresis [147,148]. As mentioned before, P2X4R⁻null mice exhibit a hypertensive phenotype [148]. Therefore, it is tempting to speculate that in patients with essential hypertension, P2X4R agonists may contribute to promote natriuresis and to lower blood pressure.

Acute kidney injury (AKI) inflicts cardiovascular damage by means of local inflammation, altered mitochondrial function, and the development of local fibrosis [138]. Inflammation-induced P2X7R receptor activation display an important role in renal disease progression, with some properties being shared by its counterpart P2X4R [149]. In a rodent model of ischemic AKI, activation of the P2X4R exacerbates tubular necrosis and NOD-like receptor 3 (NLRP3)-dependent inflammation [150]. In contrast to the cardiac P2X4R, the activation of which putatively favors fibrosis [87], the renal P2X4R seems to have an antifibrotic role [151].

Chronic elevation of intraglomerular pressure caused by arterial hypertension is accompanied by proteinuria, as a consequence of podocyte damage. Proteinuria is a surrogate marker of cardiovascular diseases [152]. Podocytes are sensitive to intraglomerular pressure, and they are important elements of the glomerular filtration barrier. Interestingly, podocytes are endowed with P2X4Rs operating mechanotransduction by interacting with podocin and highly enriched cholesterol domains [153]. Microalbuminuria is a hallmark of diabetic nephropathy, the progression of which has been associated with P2X4R activation and downstream stimulation of NLRP3 inflammasome in podocytes [154]. However, it still remains to be clarified whether the mechanosensitive role of the P2X4R in podocytes has a protective or deleterious effect in the progression to glomerulosclerosis secondary to hypertension and diabetes.

6. Is the P2X4 Receptor Activation Beneficial or Detrimental in the Lung?

Heart and lungs operate as a coupled unit, with numerous hemodynamic and neurohormonal interactions between them. Pulmonary diseases increase morbidity and worsen survival in patients with cardiovascular diseases. Pulmonary arterial hypertension (PAH) is a prototypic disease model of pulmonary and cardiac functions' interdependence. In PAH, there is a disproportionate increase in pulmonary vascular resistance (PVR), resulting in maladaptive remodeling of the right ventricle (RV) and HF [155]. Pulmonary vasodilators decrease PVR and improve PAH, particularly in group 1 PAH patients [156]. ATP and its metabolite, adenosine, have important roles in controlling pulmonary vascular tone. Similarly to that occurring in other vascular beds, ATP has both vasodilatory and vasoconstrictive effects in pulmonary arteries. Pulmonary vasoconstriction caused by ATP depends on activation of P2X₁, P2Y₂, and P2Y₄ receptors on VSMCs [157–160], while vasodilation prevails when it activates endothelial-located metabotropic P2Y₁, P2Y₂, P2Y₄, and/or ionotropic P2X4R [157–160]. As aforementioned, the pulmonary arteries express high levels of the P2X4R [161,162], where it seems to mediate shear-stress-induced ATP vasodilation [119,160]. Despite the fact that it has been demonstrated that intravenous delivery of P2X4R agonists has the potential to cause pulmonary vasodilation and to decrease RV overload, this warrants further investigations in disease models of PAH.

Respiratory infections and exacerbation of chronic lung diseases are common causes of acute decompensated heart failure (ADHF) [162,163]. Moreover, pulmonary congestion in ADHF often accompanies with ventilation/perfusion mismatch, which intensifies respiratory insufficiency with hypoxemia and hypercapnia, a situation that is already compromised in patients with underlying chronic pulmonary disease. As mentioned above, chronic hypoxemia, as well as respiratory acidosis, stimulate peripheral chemoreceptors and trigger several cardiovascular reflexes, accompanied by deleterious sympathetic overactivation and abnormal handling of water and natriuresis [123,155].

Most patients presenting with ADHF usually require diuretic, vasodilatory, and inotropic therapies to reduce congestion and optimize respiratory function and cardiac output [99]. Considering the proposed framework for the P2X4R in cardiovascular diseases, development of new and better-tolerated agonists for this receptor may represent an opportunity to improve ADHF patients' symptomatology, as they would reduce pulmonary congestion subsequently to increases in cardiac inotropy, diuresis, and vasodilation.

Apart from endothelial-induced vasodilation of pulmonary vessels and decrease of right ventricle afterload, prolonged P2X4R activation has been associated with inflammation, hyperreactivity, and mucus production in the airways of susceptible individuals [164–166]. Chronic lung diseases usually have in their background an inflammatory response that precipitates deterioration of both cardiac and respiratory functions [167,168]. Extracellular ATP raises dramatically during inflammatory conditions to levels high enough to activate P2X7, P2Y₂, and P2Y₆ receptors, which normally exhibit low affinity for the nucleotide (see Section 2). Activation of these receptors is often associated with tissue fibrosis, polymorphonuclear leukocytes infiltration, and other immune cell activation, which are hallmarks of maladaptive pulmonary remodeling [167]. Indeed, termination of P2 purinoceptors-mediated signaling by enzymatic conversion of ATP to adenosine revealed to be protective in lung injury [169].

Although the P2X4R is present in immune cells from both myeloid and lymphoid lineages [166], and its expression is upregulated in inflammatory conditions [170], there is still no consensus about its role in inflammation given that both pro- and anti-inflammatory effects have been described in the literature [166,171–174]. In a murine experimental model of asthma, a positive correlation between P2X4R levels and inflammation severity has been found; blockage of P2X4R activation reversed some asthma remodeling hallmarks, including bronchoalveolar lavage fluid eosinophilia, peri-bronchial inflammation, Th2 cytokine production, and bronchial hyper-responsiveness [173,175]. Recently, it was demonstrated that stimulation of the P2X4R favors T cell migration to the lungs, whereas blockage of this receptor is protective against acute rejection of lung allotransplants [176]. The pro-inflammatory roles of the P2X4R are, in part, due to its expression in leucocytes, namely lymphocytes and eosinophils [176,177]. Moreover, it seems that ATP-induced mast cell degranulation involves the P2X4R activation [42], and this receptor is also implicated in the contraction of airway smooth muscle cells [178]. Overall, these features may contribute to favor bronchoconstriction and allergic responses in inflammatory lung diseases.

ATP-induced airway secretion has gained special attention in the last decade. The P2X4R is involved in mucous secretion and in the control of lung surfactant production [165,170]. Mechanical stretch of alveoli increases the release of ATP to the extracellular milieu along with increases in lung surfactant production [179,180]. Alveolar type II (AT II) cells are specialized in the secretion of lung surfactant [181]; these cells harbor the P2X4R as the most abundant P2X receptor subtype [182]. The P2X4R has a unique subcellular location in AT II cells, being found in lamellar bodies that contain lung surfactant. The release of lung surfactant is an exocytotic process designated by fusion-activated Ca²⁺ entry (FACE), which is operated by P2X4 (and possibly P2Y₂) receptors activation [182]. The P2X4R inward current also drives water reabsorption across the alveolar epithelium, which is crucial to maintain alveoli relatively dry to increase gas exchange efficiency [183]. Although not completely resolved, it is believed that the main extracellular ATP source for P2X4R activation is alveolar type I (AT I) cells [179,180], yet extracellular release of ATP may be further amplified by P2X4R-induced lysosome fusion, resulting in the release of ATP stored inside lamellar bodies [184].

Acute lung inflammation causes a disproportionate increase in extracellular ATP, which contributes to dysregulation of surfactant production and initiation of the inflammatory response [185]. Although speculative, from a mechanistic point of view, alveolar P2X4Rs may have some role in acute respiratory distress syndrome (ARDS), a life-threatening condition frequently requiring emergent positive-pressure ventilation support, so it is pursued nowadays in the SARS-CoV-2/COVID-19 pandemic [186]. In this context, selective modulation of P2X4Rs activation may control lung surfactant production and protect patients from barotrauma and alveolar collapse, which are common features in ARDS [187,188].

Likewise, patients presenting with ADHF usually require non-invasive positive-pressure ventilation to resolve signs and symptoms of congestion. It is, thus, tempting to hypothesize that activation of alveolar P2X4Rs may also be beneficial to these patients' outcomes.

7. Is there a Dark Side of P2X4 Receptor Activation Outside the Cardiovascular System?

In this section, we will briefly address putative side-effects of P2X4R activation outside the cardiovascular system, which might be relevant when translating the use of P2X4R agonists to the clinical practice.

The P2X4R was first identified in the central nervous system [21], with the following reports showing its widespread expression both in central and in peripheral neurons (reviewed in [9,189,190]). Great attention has been given to the modulation of P2X4R activity to control chronic pain. The P2X4R is upregulated in spinal microglia cells following peripheral nerve injury; genetic ablation or pharmacological blockage of overexpressed P2X4Rs can reverse allodynia occurring in this condition [191]. Increasing evidence demonstrates that blockade of the P2X4R alleviates neuropathic and inflammatory pain [45,192,193], as well as other causes of chronic pain, including migraine [194]. Hyperalgesia caused by P2X4R activation seems to depend on brain-derived neurotrophic factor (BDNF) release [192,195]. In this context, experimental data indicate that the P2X4R may play a role in long-term potentiation and synaptic plasticity [196,197], which may underlie its effect on chronic pain. Notwithstanding the involvement of the P2X4R in neuropathic and inflammatory pain conditions, its activation did not affect acute pain sensation in animals not prone to pain hypersensitivity [191]. Although changes in nociception may be a problem for the clinical use of P2X4R agonists in cardiovascular diseases, this warrants confirmation in clinical trials due to contradictory results using animal models. The same holds true regarding involvement of the P2X4R in neuroinflammatory conditions, as conflicting results, either protective [135,172] or deleterious [198,199], have emerged. With this in mind, forthcoming P2X4R agonists targeting the cardiovascular system should not cross the blood–brain barrier; otherwise, prolonged stimulation may have negative impacts in the course of some neurodegenerative diseases, chronic pain conditions, and epilepsy in susceptible individuals [8,9,199].

Rather than deleterious, the P2X4R expression and function in the brain reward system have been associated with salutary effects in alcohol abuse. Ethanol is a preferential inhibitor of the P2X4R, a situation that can be overcome by positive allosteric modulators, like ivermectin or derivatives, which have been shown to reduce ethanol drinking behavior (reviewed in [200,201]).

Inflammatory flairs may also occur with P2X4Rs activation, particularly in patients with proinflammatory backgrounds, such as rheumatoid arthritis and asthma [8,9,166,173,202]. Pruritus and hypersensitivity reactions can also be a consequence of stimulation of P2X4R in mast cells and other immunocytes [42,166]. Cancer is another type of inflammatory condition where the P2X4R might be involved, although its role remains uncertain [8].

In the opposite direction of the inflammatory spectrum, activation of P2X4R in macrophages exhibited both anti-inflammatory properties and bactericidal activity against Gram-negative bacteria in the cecal ligation and puncture-induced sepsis rodent model. This P2X4R-mediated protection improved multi-organ failure and sepsis-related death [174], which supports future therapeutics with P2X4R agonists to treat patients with ADHF secondary to sepsis.

8. Conclusions

The development of novel purinergic-based medications has been struggling with the ubiquitous nature of P1 and P2 purinoceptors and lack of drug selectivity. Pushed by the success of P2Y₁₂ receptor antagonists as anti-thrombotic drugs, last decades have provided new experimental and pharmacological tools to dissect the purinergic signaling cascade, including the purinoceptors function, which have granted a series of promising druggable targets, and some of them were already admitted to clinical trials. Selective drugs targeting the P2X4R are limited to few antagonists and positive allosteric

modulators, without any selective agonist being available so far. Despite this, researchers worldwide have been unlocking the potential of the ionotropic P2X4R to treat several diseases. By reviewing recent data in the literature, we have highlighted here the benefits putatively arising from activation of the P2X4R, either by directly ameliorating the cardiac function or by indirectly improving the function of other systems impacting cardiovascular disease conditions. We also provide some hints about problems to be solved regarding unwanted side effects that may emerge upon activation of P2X4R to stimulate the cardiovascular system, such as arrhythmias and worsening of diastolic dysfunction in the heart; water–electrolyte imbalance and remodeling in renal injury; inflammation, mucous secretion, and hyperreactivity in airway diseases; and nociceptive-related changes. Taken together, we provide input for forthcoming development of medications based on selective activation of the P2X4R to reduce the burden of cardiovascular diseases.

Author Contributions: Conceptualization, B.B. and P.C.-d.-S.; writing—original draft preparation, B.B. and P.C.-d.-S.; writing—review and editing, B.B. and P.C.-d.-S.; supervision, P.C.-d.-S.; funding acquisition, P.C.-d.-S. All authors have read and agreed to the published version of the manuscript.

Funding: This work was supported by Foundation for Science and Technology (FCT) (UID/BIM/4308/2019 and UIDB/04308/2020). BB is in receipt of a PhD studentship from FCT (FEDER funding SFRH/BD/104114/2014).

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

References

1. Burnstock, G. Purine and purinergic receptors. *Brain Neurosci. Adv.* **2018**, *2*, 2398212818817494. [[CrossRef](#)]
2. Lazarowski, E.R. Vesicular and conductive mechanisms of nucleotide release. *Purinergic Signal.* **2012**, *8*, 359–373. [[CrossRef](#)] [[PubMed](#)]
3. Taruno, A. ATP Release Channels. *Int. J. Mol. Sci.* **2018**, *19*, 808. [[CrossRef](#)] [[PubMed](#)]
4. Yegutkin, G.G. Enzymes involved in metabolism of extracellular nucleotides and nucleosides: Functional implications and measurement of activities. *Crit. Rev. Biochem. Mol. Biol.* **2014**, *49*, 473–497. [[CrossRef](#)] [[PubMed](#)]
5. Drury, A.N.; Szent-Gyorgyi, A. The physiological activity of adenine compounds with especial reference to their action upon the mammalian heart. *J. Physiol.* **1929**, *68*, 213–237. [[CrossRef](#)] [[PubMed](#)]
6. Van der Meijden, P.E.J.; Heemskerk, J.W.M. Platelet biology and functions: New concepts and clinical perspectives. *Nat. Rev. Cardiol.* **2019**, *16*, 166–179. [[CrossRef](#)] [[PubMed](#)]
7. Roffi, M.; Patrono, C.; Collet, J.-P.; Mueller, C.; Valgimigli, M.; Andreotti, F.; Bax, J.J.; Borger, M.A.; Brotons, C.; Chew, D.P.; et al. 2015 ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation: Task Force for the Management of Acute Coronary Syndromes in Patients Presenting without Persistent ST-Segment Elevation of the European Society of Cardiology (ESC). *Eur. Heart J.* **2016**, *37*, 267–315. [[CrossRef](#)]
8. Burnstock, G. Purinergic Signalling: Therapeutic Developments. *Front. Pharmacol.* **2017**, *8*, 661. [[CrossRef](#)] [[PubMed](#)]
9. Suurvali, J.; Boudinot, P.; Kanellopoulos, J.; Ruutel Boudinot, S. P2X4: A fast and sensitive purinergic receptor. *Biomed. J.* **2017**, *40*, 245–256. [[CrossRef](#)]
10. Kawate, T.; Michel, J.C.; Birdsong, W.T.; Gouaux, E. Crystal structure of the ATP-gated P2X(4) ion channel in the closed state. *Nature* **2009**, *460*, 592–598. [[CrossRef](#)]
11. Sophocleous, R.A.; Berg, T.; Finol-Urdaneta, R.K.; Sluyter, V.; Keshiya, S.; Bell, L.; Curtis, S.J.; Curtis, B.L.; Seavers, A.; Bartlett, R.; et al. Pharmacological and genetic characterisation of the canine P2X4 receptor. *Br. J. Pharmacol.* **2020**, *177*, 2812–2829. [[CrossRef](#)] [[PubMed](#)]
12. Stokes, L.; Scurrah, K.; Ellis, J.A.; Cromer, B.A.; Skarratt, K.K.; Gu, B.J.; Harrap, S.B.; Wiley, J.S. A loss-of-function polymorphism in the human P2X4 receptor is associated with increased pulse pressure. *Hypertension* **2011**, *58*, 1086–1092. [[CrossRef](#)]
13. Saul, A.; Hausmann, R.; Kless, A.; Nicke, A. Heteromeric assembly of P2X subunits. *Front. Cell Neurosci* **2013**, *7*, 250. [[CrossRef](#)]

14. Antonio, L.S.; Stewart, A.P.; Varanda, W.A.; Edwardson, J.M. Identification of P2X2/P2X4/P2X6 heterotrimeric receptors using atomic force microscopy (AFM) imaging. *FEBS Lett.* **2014**, *588*, 2125–2128. [[CrossRef](#)] [[PubMed](#)]
15. Hattori, M.; Gouaux, E. Molecular mechanism of ATP binding and ion channel activation in P2X receptors. *Nature* **2012**, *485*, 207–212. [[CrossRef](#)] [[PubMed](#)]
16. Jacobson, K.A.; Jarvis, M.F.; Williams, M. Purine and pyrimidine (P2) receptors as drug targets. *J. Med. Chem.* **2002**, *45*, 4057–4093. [[CrossRef](#)] [[PubMed](#)]
17. Kasuya, G.; Fujiwara, Y.; Tsukamoto, H.; Morinaga, S.; Ryu, S.; Touhara, K.; Ishitani, R.; Furutani, Y.; Hattori, M.; Nureki, O. Structural insights into the nucleotide base specificity of P2X receptors. *Sci. Rep.* **2017**, *7*, 45208. [[CrossRef](#)]
18. Egan, T.M.; Samways, D.S.; Li, Z. Biophysics of P2X receptors. *Pflugers Arch.* **2006**, *452*, 501–512. [[CrossRef](#)]
19. Samways, D.S.; Li, Z.; Egan, T.M. Principles and properties of ion flow in P2X receptors. *Front. Cell Neurosci.* **2014**, *8*, 6. [[CrossRef](#)]
20. Coddou, C.; Yan, Z.; Obsil, T.; Huidobro-Toro, J.P.; Stojilkovic, S.S. Activation and regulation of purinergic P2X receptor channels. *Pharmacol. Rev.* **2011**, *63*, 641–683. [[CrossRef](#)]
21. Soto, F.; Garcia-Guzman, M.; Gomez-Hernandez, J.M.; Hollmann, M.; Karschin, C.; Stuhmer, W. P2X4: An ATP-activated ionotropic receptor cloned from rat brain. *Proc. Natl. Acad. Sci. USA* **1996**, *93*, 3684–3688. [[CrossRef](#)] [[PubMed](#)]
22. Khakh, B.S.; Bao, X.R.; Labarca, C.; Lester, H.A. Neuronal P2X transmitter-gated cation channels change their ion selectivity in seconds. *Nat. Neurosci.* **1999**, *2*, 322–330. [[CrossRef](#)]
23. Bernier, L.P.; Ase, A.R.; Boue-Grabot, E.; Seguela, P. P2X4 receptor channels form large noncytolytic pores in resting and activated microglia. *Glia* **2012**, *60*, 728–737. [[CrossRef](#)]
24. Shinozaki, Y.; Sumitomo, K.; Tsuda, M.; Koizumi, S.; Inoue, K.; Torimitsu, K. Direct observation of ATP-induced conformational changes in single P2X(4) receptors. *PLoS Biol.* **2009**, *7*, e1000103. [[CrossRef](#)] [[PubMed](#)]
25. Bo, X.; Zhang, Y.; Nassar, M.; Burnstock, G.; Schoepfer, R. A P2X purinoceptor cDNA conferring a novel pharmacological profile. *FEBS Lett.* **1995**, *375*, 129–133. [[CrossRef](#)]
26. Jones, C.A.; Chessell, I.P.; Simon, J.; Barnard, E.A.; Miller, K.J.; Michel, A.D.; Humphrey, P.P. Functional characterization of the P2X4 receptor orthologues. *Br. J. Pharmacol.* **2000**, *129*, 388–394. [[CrossRef](#)] [[PubMed](#)]
27. Abdelrahman, A.; Namasivayam, V.; Hinz, S.; Schiedel, A.C.; Köse, M.; Burton, M.; El-Tayeb, A.; Gillard, M.; Bajorath, J.; De Ryck, M.; et al. Characterization of P2X4 receptor agonists and antagonists by calcium influx and radioligand binding studies. *Biochem. Pharmacol.* **2017**, *125*, 41–54. [[CrossRef](#)] [[PubMed](#)]
28. Bianchi, B.R.; Lynch, K.J.; Touma, E.; Niforatos, W.; Burgard, E.C.; Alexander, K.M.; Park, H.S.; Yu, H.; Metzger, R.; Kowaluk, E.; et al. Pharmacological characterization of recombinant human and rat P2X receptor subtypes. *Eur. J. Pharmacol.* **1999**, *376*, 127–138. [[CrossRef](#)]
29. He, M.-L.; Gonzalez-Iglesias, A.E.; Stojilkovic, S.S. Role of Nucleotide P2 Receptors in Calcium Signaling and Prolactin Release in Pituitary Lactotrophs. *J. Biol. Chem.* **2003**, *278*, 46270–46277. [[CrossRef](#)] [[PubMed](#)]
30. Yusuf, S.; Bosch, J.; Dagenais, G.; Zhu, J.; Xavier, D.; Liu, L.; Pais, P.; Lopez-Jaramillo, P.; Leiter, L.A.; Dans, A.; et al. Cholesterol Lowering in Intermediate-Risk Persons without Cardiovascular Disease. *N. Engl. J. Med.* **2016**, *374*, 2021–2031. [[CrossRef](#)] [[PubMed](#)]
31. Browne, L.E.; North, R.A. P2X receptor intermediate activation states have altered nucleotide selectivity. *J. Neurosci.* **2013**, *33*, 14801–14808. [[CrossRef](#)] [[PubMed](#)]
32. Roberts, J.A.; Evans, R.J. Atp Binding at Human P2x1 Receptors: Contribution of Aromatic and Basic Amino Acids Revealed Using Mutagenesis and Partial Agonists. *J. Biol. Chem.* **2004**, *279*, 9043–9055. [[CrossRef](#)] [[PubMed](#)]
33. Shen, J.B.; Cronin, C.; Sonin, D.; Joshi, B.V.; Gongora Nieto, M.; Harrison, D.; Jacobson, K.A.; Liang, B.T. P2X purinergic receptor-mediated ionic current in cardiac myocytes of calsequestrin model of cardiomyopathy: Implications for the treatment of heart failure. *Am. J. Physiol. Heart Circ. Physiol.* **2007**, *292*, H1077–H1084. [[CrossRef](#)]
34. Kumar, T.S.; Yang, T.; Mishra, S.; Cronin, C.; Chakraborty, S.; Shen, J.B.; Liang, B.T.; Jacobson, K.A. 5'-Phosphate and 5'-phosphonate ester derivatives of (N)-methanocarba adenosine with in vivo cardioprotective activity. *J. Med. Chem.* **2013**, *56*, 902–914. [[CrossRef](#)]

35. Hernandez-Olmos, V.; Abdelrahman, A.; El-Tayeb, A.; Freudendahl, D.; Weinhausen, S.; Müller, C.E. N-Substituted Phenoxazine and Acridone Derivatives: Structure–Activity Relationships of Potent P2X4 Receptor Antagonists. *J. Med. Chem.* **2012**, *55*, 9576–9588. [[CrossRef](#)]
36. Balázs, B.; Danko, T.; Kovacs, G.; Köles, L.; Hediger, M.A. Zsembery, Ákos Investigation of the Inhibitory Effects of the Benzodiazepine Derivative, 5-BDBD on P2X4Purinergerg Receptors by two Complementary Methods. *Cell. Physiol. Biochem.* **2013**, *32*, 11–24. [[CrossRef](#)]
37. Bo, X.; Liu, M.; Schoepfer, R.; Burnstock, G. Characterization and expression of ATP P2X4 receptor from embryonic chick skeletal muscle. *Drug Dev. Res.* **2001**, *53*, 22–28. [[CrossRef](#)]
38. Virginio, C.; Robertson, G.; Surprenant, A.; North, R.A. Trinitrophenyl-Substituted Nucleotides Are Potent Antagonists Selective for P2x₁, P2x₃, and Heteromeric P2x_{2/3} Receptors. *Mol. Pharmacol.* **1998**, *53*, 969–973. [[PubMed](#)]
39. Tomioka, A.; Ueno, S.; Kohama, K.; Goto, F.; Inoue, K. Propofol potentiates ATP-activated currents of recombinant P2X4 receptor channels expressed in human embryonic kidney 293 cells. *Neurosci. Lett.* **2000**, *284*, 167–170. [[CrossRef](#)]
40. Coddou, C.; Sandoval, R.; Hevia, M.J.; Stojilkovic, S.S. Characterization of the antagonist actions of 5-BDBD at the rat P2X4 receptor. *Neurosci. Lett.* **2019**, *690*, 219–224. [[CrossRef](#)] [[PubMed](#)]
41. Ase, A.R.; Honson, N.S.; Zaghdane, H.; Pfeifer, T.A.; Seguela, P. Identification and characterization of a selective allosteric antagonist of human P2X4 receptor channels. *Mol. Pharmacol.* **2015**, *87*, 606–616. [[CrossRef](#)] [[PubMed](#)]
42. Yoshida, K.; Tajima, M.; Nagano, T.; Obayashi, K.; Ito, M.; Yamamoto, K.; Matsuoka, I. Co-Stimulation of Purinergerg P2X4 and Prostanoid EP3 Receptors Triggers Synergistic Degranulation in Murine Mast Cells. *Int. J. Mol. Sci.* **2019**, *20*, 5157. [[CrossRef](#)] [[PubMed](#)]
43. Tian, M.; Abdelrahman, A.; Weinhausen, S.; Hinz, S.; Weyer, S.; Dosa, S.; El-Tayeb, A.; Müller, C.E. Carbamazepine derivatives with P2X4 receptor-blocking activity. *Bioorganic Med. Chem.* **2014**, *22*, 1077–1088. [[CrossRef](#)] [[PubMed](#)]
44. Schneider, M.; Prudic, K.; Pippel, A.; Klapperstück, M.; Braam, U.; Müller, C.E.; Schmalzing, G.; Markwardt, F. Interaction of Purinergerg P2x4 and P2x7 Receptor Subunits. *Front. Pharmacol.* **2017**, *8*, 860. [[CrossRef](#)] [[PubMed](#)]
45. Matsumura, Y.; Yamashita, T.; Sasaki, A.; Nakata, E.; Kohno, K.; Masuda, T.; Tozaki-Saitoh, H.; Imai, T.; Kuraishi, Y.; Tsuda, M.; et al. A novel P2X4 receptor-selective antagonist produces anti-allodynic effect in a mouse model of herpetic pain. *Sci. Rep.* **2016**, *6*, 32461. [[CrossRef](#)] [[PubMed](#)]
46. Inoue, K.; Tsuda, M. Microglia in neuropathic pain: cellular and molecular mechanisms and therapeutic potential. *Nat. Rev. Neurosci.* **2018**, *19*, 138–152. [[CrossRef](#)] [[PubMed](#)]
47. Beswick, P.; Wahab, B.; Honey, M.A.; Paradowski, M.; Jiang, K.; Lochner, M.; Murrell-Lagnado, R.D.; Thompson, A.J. A challenge finding P2X1 and P2X4 ligands. *Neuropharmacology* **2019**, *157*, 107674. [[CrossRef](#)] [[PubMed](#)]
48. Nagata, K.; Imai, T.; Yamashita, T.; Tsuda, M.; Tozaki-Saitoh, H.; Inoue, K. Antidepressants inhibit P2X4 receptor function: a possible involvement in neuropathic pain relief. *Mol. Pain* **2009**, *5*, 20. [[CrossRef](#)]
49. Masuda, T.; Iwamoto, S.; Yoshinaga, R.; Tozaki-Saitoh, H.; Nishiyama, A.; Mak, T.W.; Tamura, T.; Tsuda, M.; Inoue, K. Transcription factor IRF5 drives P2X4R⁺-reactive microglia gating neuropathic pain. *Nat. Commun.* **2014**, *5*, 3771. [[CrossRef](#)]
50. Yamashita, T.; Yamamoto, S.; Zhang, J.; Kometani, M.; Tomiyama, D.; Kohno, K.; Tozaki-Saitoh, H.; Inoue, K.; Tsuda, M. Duloxetine Inhibits Microglial P2X4 Receptor Function and Alleviates Neuropathic Pain after Peripheral Nerve Injury. *PLoS ONE* **2016**, *11*, e0165189. [[CrossRef](#)]
51. Werner, S.; Mesch, S.; Hillig, R.C.; Laak, A.T.; Klint, J.; Neagoe, I.; Laux-Biehlmann, A.; Dahllhof, H.; Brauer, N.; Puetter, V.; et al. Faculty Opinions recommendation of Discovery and Characterization of the Potent and Selective P2X4 Inhibitor N-[4-(3-Chlorophenoxy)-3-sulfamoylphenyl]-2-phenylacetamide (BAY-1797) and Structure-Guided Amelioration of Its CYP3A4 Induction Profile. *Fac. Opin. Post Publ. Peer Rev. Biomed. Lit.* **2019**, *62*, 11194–11217. [[CrossRef](#)]
52. Williams, W.A.; Linley, J.E.; Jones, C.A.; Shibata, Y.; Snijder, A.; Button, J.; Hatcher, J.P.; Huang, L.; Taddese, B.; Thornton, P.; et al. Antibodies binding the head domain of P2X4 inhibit channel function and reverse neuropathic pain. *Pain* **2019**, *160*, 1989–2003. [[CrossRef](#)] [[PubMed](#)]

53. Shen, J.B.; Toti, K.S.; Chakraborty, S.; Kumar, T.S.; Cronin, C.; Liang, B.T.; Jacobson, K.A. Prevention and rescue of cardiac dysfunction by methanocarba adenosine monophosphonate derivatives. *Purinergic Signal*. **2020**, *16*, 61–72. [[CrossRef](#)] [[PubMed](#)]
54. Norenberg, W.; Sobottka, H.; Hempel, C.; Plötz, T.; Fischer, W.; Schmalzing, G.; Schaefer, M. Positive allosteric modulation by ivermectin of human but not murine P2X7 receptors. *Br. J. Pharmacol.* **2012**, *167*, 48–66. [[CrossRef](#)]
55. Miller, K.J.; Michel, A.D.; Chessell, I.; Humphrey, P. Cibacron blue allosterically modulates the rat P2X4 receptor. *Neuropharmacol.* **1998**, *37*, 1579–1586. [[CrossRef](#)]
56. Dhuna, K.; Felgate, M.; Bidula, S.; Walpole, S.; Bibic, L.; Cromer, B.A.; Angulo, J.; Sanderson, J.; Stebbing, M.J.; Stokes, L. Ginsenosides Act As Positive Modulators of P2X4 Receptors. *Mol. Pharmacol.* **2018**, *95*, 210–221. [[CrossRef](#)] [[PubMed](#)]
57. Sivcev, S.; Slavikova, B.; Rupert, M.; Ivetic, M.; Nekardova, M.; Kudova, E.; Zemkova, H. Synthetic testosterone derivatives modulate rat P2X2 and P2X4 receptor channel gating. *J. Neurochem.* **2019**, *150*, 28–43. [[CrossRef](#)] [[PubMed](#)]
58. Coddou, C.; Lorca, R.; Acuña-Castillo, C.; Grauso, M.; Rassendren, F.; Huidobro-Toro, J. Heavy metals modulate the activity of the purinergic P2X4 receptor. *Toxicol. Appl. Pharmacol.* **2005**, *202*, 121–131. [[CrossRef](#)]
59. Codocedo, J.F.; Rodríguez, F.E.; Huidobro-Toro, J.P. Neurosteroids differentially modulate P2X4ATP-gated channels through non-genomic interactions. *J. Neurochem.* **2009**, *110*, 734–744. [[CrossRef](#)]
60. Popova, M.; Asatryan, L.; Ostrovskaya, O.; Wyatt, L.; Li, K.; Alkana, R.L.; Davies, D.L. A point mutation in the ectodomain-transmembrane 2 interface eliminates the inhibitory effects of ethanol in P2X4 receptors. *J. Neurochem.* **2009**, *112*, 307–317. [[CrossRef](#)]
61. Davies, D.L.; Kochegarov, A.A.; Kuo, S.T.; Kulkarni, A.A.; Woodward, J.J.; King, B.F.; Alkana, R.L. Ethanol differentially affects ATP-gated P2X and P2X receptor subtypes expressed in oocytes. *Neuropharmacol.* **2005**, *49*, 243–253. [[CrossRef](#)] [[PubMed](#)]
62. Stoop, R.; Surprenant, A.; North, R.A. Different sensitivities to pH of ATP-induced currents at four cloned P2X receptors. *J. Neurophysiol.* **1997**, *78*, 1837–1840. [[CrossRef](#)] [[PubMed](#)]
63. Li, J.; Fountain, S.J. Fluvastatin suppresses native and recombinant human P2X4 receptor function. *Purinergic Signal*. **2012**, *8*, 311–316. [[CrossRef](#)] [[PubMed](#)]
64. Ilyaskin, A.V.; Sure, F.; Nesterov, V.; Haerteis, S.; Korbmacher, C. Bile acids inhibit human purinergic receptor P2X4 in a heterologous expression system. *J. Gen. Physiol.* **2019**, *151*, 820–833. [[CrossRef](#)]
65. Schmid, R.; Evans, R.J. ATP-Gated P2X Receptor Channels: Molecular Insights into Functional Roles. *Annu. Rev. Physiol.* **2019**, *81*, 43–62. [[CrossRef](#)]
66. Garcia-Guzman, M.; Soto, F.; Gomez-Hernandez, J.M.; Lund, P.E.; Stuhmer, W. Characterization of recombinant human P2X4 receptor reveals pharmacological differences to the rat homologue. *Mol. Pharmacol.* **1997**, *51*, 109–118. [[CrossRef](#)]
67. Coddou, C.; Stojilkovic, S.S.; Huidobro-Toro, J.P. Allosteric modulation of ATP-gated P2X receptor channels. *Rev. Neurosci.* **2011**, *22*, 335–354. [[CrossRef](#)]
68. Kasuya, G.; Fujiwara, Y.; Takemoto, M.; Dohmae, N.; Nakada-Nakura, Y.; Ishitani, R.; Hattori, M.; Nureki, O. Structural Insights into Divalent Cation Modulations of ATP-Gated P2X Receptor Channels. *Cell Rep.* **2016**, *14*, 932–944. [[CrossRef](#)]
69. Clarke, C.E.; Benham, C.D.; Bridges, A.; George, A.R.; Meadows, H.J. Mutation of histidine 286 of the human P2X4 purinoceptor removes extracellular pH sensitivity. *J. Physiol.* **2000**, *523*, 697–703. [[CrossRef](#)]
70. Khakh, B.S.; Proctor, W.R.; Dunwiddie, T.V.; Labarca, C.; Lester, H.A. Allosteric control of gating and kinetics at P2X(4) receptor channels. *J. Neurosci.* **1999**, *19*, 7289–7299. [[CrossRef](#)]
71. Brown, D.A.; Yule, D.I. Protein kinase A regulation of P2X(4) receptors: Requirement for a specific motif in the C-terminus. *Biochim. Biophys. Acta* **2010**, *1803*, 275–287. [[CrossRef](#)] [[PubMed](#)]
72. Bernier, L.P.; Ase, A.R.; Chevallier, S.; Blais, D.; Zhao, Q.; Boue-Grabot, E.; Logothetis, D.; Seguela, P. Phosphoinositides regulate P2X4 ATP-gated channels through direct interactions. *J. Neurosci.* **2008**, *28*, 12938–12945. [[CrossRef](#)] [[PubMed](#)]
73. Ohata, Y.; Ogata, S.; Nakanishi, K.; Kanazawa, F.; Uenoyama, M.; Hiroi, S.; Tominaga, S.; Kawai, T. Expression of P2X4R mRNA and protein in rats with hypobaric hypoxia-induced pulmonary hypertension. *Circ. J.* **2011**, *75*, 945–954. [[CrossRef](#)] [[PubMed](#)]

74. Li, F.; Wang, L.; Li, J.W.; Gong, M.; He, L.; Feng, R.; Dai, Z.; Li, S.Q. Hypoxia induced amoeboid microglial cell activation in postnatal rat brain is mediated by ATP receptor P2X4. *BMC Neurosci.* **2011**, *12*, 111. [[CrossRef](#)]
75. Tang, Y.; Matsuoka, I.; Ono, T.; Inoue, K.; Kimura, J. Selective up-regulation of P2X4-receptor gene expression by interferon-gamma in vascular endothelial cells. *J. Pharmacol. Sci.* **2008**, *107*, 419–427. [[CrossRef](#)]
76. Gu, B.J.; Sun, C.; Valova, V.A.; Skarratt, K.K.; Wiley, J.S. Identification of the promoter region of the P2RX4 gene. *Mol. Biol. Rep.* **2010**, *37*, 3369–3376. [[CrossRef](#)]
77. Fountain, S.J.; North, R.A. A C-terminal lysine that controls human P2X4 receptor desensitization. *J. Biol. Chem.* **2006**, *281*, 15044–15049. [[CrossRef](#)]
78. Murrell-Lagnado, R.D. A role for P2X4 receptors in lysosome function. *J. Gen. Physiol.* **2018**, *150*, 185–187. [[CrossRef](#)]
79. Xu, J.; Chai, H.; Ehinger, K.; Egan, T.M.; Srinivasan, R.; Frick, M.; Khakh, B.S. Imaging P2X4 receptor subcellular distribution, trafficking, and regulation using P2X4-pHluorin. *J. Gen. Physiol.* **2014**, *144*, 81–104. [[CrossRef](#)]
80. Erlinge, D.; Burnstock, G. P2 receptors in cardiovascular regulation and disease. *Purinergic Signal.* **2008**, *4*, 1–20. [[CrossRef](#)]
81. Burnstock, G. Purinergic Signaling in the Cardiovascular System. *Circ. Res.* **2017**, *120*, 207–228. [[CrossRef](#)] [[PubMed](#)]
82. Burnstock, G.; Pelleg, A. Cardiac purinergic signalling in health and disease. *Purinergic Signal.* **2015**, *11*, 1–46. [[CrossRef](#)]
83. Musa, H.; Tellez, J.O.; Chandler, N.J.; Greener, I.D.; Maczewski, M.; Mackiewicz, U.; Beresewicz, A.; Molenaar, P.; Boyett, M.R.; Dobrzynski, H. P2 purinergic receptor mRNA in rat and human sinoatrial node and other heart regions. *Naunyn Schmiedebergs Arch. Pharmacol.* **2009**, *379*, 541–549. [[CrossRef](#)] [[PubMed](#)]
84. Bogdanov, Y.; Rubino, A.; Burnstock, G. Characterisation of subtypes of the P2X and P2Y families of ATP receptors in the foetal human heart. *Life Sci.* **1998**, *62*, 697–703. [[CrossRef](#)]
85. Braganca, B.; Nogueira-Marques, S.; Ferreirinha, F.; Fontes-Sousa, A.P.; Correia-de-Sa, P. The Ionotropic P2X4 Receptor has Unique Properties in the Heart by Mediating the Negative Chronotropic Effect of ATP While Increasing the Ventricular Inotropy. *Front. Pharmacol.* **2019**, *10*, 1103. [[CrossRef](#)] [[PubMed](#)]
86. Ma, B.; Ruan, H.Z.; Burnstock, G.; Dunn, P.M. Differential expression of P2X receptors on neurons from different parasympathetic ganglia. *Neuropharmacology* **2005**, *48*, 766–777. [[CrossRef](#)]
87. Chen, J.-B.; Liu, W.-J.; Che, H.; Liu, J.; Sun, H.-Y.; Li, G.-R. Adenosine-5'-triphosphate up-regulates proliferation of human cardiac fibroblasts. *Br. J. Pharmacol.* **2012**, *166*, 1140–1150. [[CrossRef](#)] [[PubMed](#)]
88. Hu, B.; Mei, Q.B.; Yao, X.J.; Smith, E.; Barry, W.H.; Liang, B.T. A novel contractile phenotype with cardiac transgenic expression of the human P2X4 receptor. *FASEB J.* **2001**, *15*, 2739–2741. [[CrossRef](#)]
89. Yang, A.; Sonin, D.; Jones, L.; Barry, W.H.; Liang, B.T. A beneficial role of cardiac P2X4 receptors in heart failure: Rescue of the caldesmon overexpression model of cardiomyopathy. *Am. J. Physiol. Heart Circ. Physiol.* **2004**, *287*, H1096–H1103. [[CrossRef](#)]
90. Sonin, D.; Zhou, S.Y.; Cronin, C.; Sonina, T.; Wu, J.; Jacobson, K.A.; Pappano, A.; Liang, B.T. Role of P2X purinergic receptors in the rescue of ischemic heart failure. *Am. J. Physiol. Heart Circ. Physiol.* **2008**, *295*, H1191–H1197. [[CrossRef](#)]
91. Shen, J.B.; Shutt, R.; Agosto, M.; Pappano, A.; Liang, B.T. Reversal of cardiac myocyte dysfunction as a unique mechanism of rescue by P2X4 receptors in cardiomyopathy. *Am. J. Physiol. Heart Circ. Physiol.* **2009**, *296*, H1089–H1095. [[CrossRef](#)] [[PubMed](#)]
92. Yang, T.; Shen, J.B.; Yang, R.; Redden, J.; Dodge-Kafka, K.; Grady, J.; Jacobson, K.A.; Liang, B.T. Novel protective role of endogenous cardiac myocyte P2X4 receptors in heart failure. *Circ. Heart Fail.* **2014**, *7*, 510–518. [[CrossRef](#)] [[PubMed](#)]
93. Eisner, D.A.; Caldwell, J.L.; Kistamas, K.; Trafford, A.W. Calcium and Excitation-Contraction Coupling in the Heart. *Circ. Res.* **2017**, *121*, 181–195. [[CrossRef](#)]
94. Shen, J.B.; Yang, R.; Pappano, A.; Liang, B.T. Cardiac P2X purinergic receptors as a new pathway for increasing Na(+) entry in cardiac myocytes. *Am. J. Physiol. Heart Circ. Physiol.* **2014**, *307*, H1469–H1477. [[CrossRef](#)] [[PubMed](#)]
95. Lakatta, E.G.; Maltsev, V.A.; Vinogradova, T.M. A coupled SYSTEM of intracellular Ca²⁺ clocks and surface membrane voltage clocks controls the timekeeping mechanism of the heart's pacemaker. *Circ. Res.* **2010**, *106*, 659–673. [[CrossRef](#)] [[PubMed](#)]

96. Belz, G.G.; Breithaupt-Grogler, K.; Osowski, U. Treatment of congestive heart failure—current status of use of digitoxin. *Eur. J. Clin. Investig.* **2001**, *31*, 10–17. [[CrossRef](#)]
97. Whayne, T.F., Jr. Clinical Use of Digitalis: A State of the Art Review. *Am. J. Cardiovasc. Drugs* **2018**, *18*, 427–440. [[CrossRef](#)]
98. Kirchhof, P.; Benussi, S.; Kotecha, D.; Ahlsson, A.; Atar, D.; Casadei, B.; Castella, M.; Diener, H.C.; Heidbuchel, H.; Hendriks, J.; et al. 2016 ESC Guidelines for the management of atrial fibrillation developed in collaboration with EACTS. *Eur. Heart J.* **2016**, *37*, 2893–2962. [[CrossRef](#)]
99. Ponikowski, P.; Voors, A.A.; Anker, S.D.; Bueno, H.; Cleland, J.G.F.; Coats, A.J.S.; Falk, V.; Gonzalez-Juanatey, J.R.; Harjola, V.P.; Jankowska, E.A.; et al. 2016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: The Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC) Developed with the special contribution of the Heart Failure Association (HFA) of the ESC. *Eur. Heart J.* **2016**, *37*, 2129–2200. [[CrossRef](#)]
100. Bossu, A.; Kostense, A.; Beekman, H.D.M.; Houtman, M.J.C.; van der Heyden, M.A.G.; Vos, M.A. Istaroxime, a positive inotropic agent devoid of proarrhythmic properties in sensitive chronic atrioventricular block dogs. *Pharmacol. Res.* **2018**, *133*, 132–140. [[CrossRef](#)]
101. Carubelli, V.; Zhang, Y.; Metra, M.; Lombardi, C.; Felker, G.M.; Filippatos, G.; O'Connor, C.M.; Teerlink, J.R.; Simmons, P.; Segal, R.; et al. Treatment with 24 hour istaroxime infusion in patients hospitalised for acute heart failure: A randomised, placebo-controlled trial. *Eur. J. Heart Fail.* **2020**. [[CrossRef](#)] [[PubMed](#)]
102. Chioncel, O.; Collins, S.P.; Butler, J. Istaroxime in acute heart failure: The holy grail is at HORIZON? *Eur. J. Heart Fail.* **2020**. [[CrossRef](#)] [[PubMed](#)]
103. Shah, S.J.; Blair, J.E.; Filippatos, G.S.; Macarie, C.; Ruzyllo, W.; Korewicki, J.; Bubenek-Turconi, S.I.; Ceracchi, M.; Bianchetti, M.; Carminati, P.; et al. Effects of istaroxime on diastolic stiffness in acute heart failure syndromes: Results from the Hemodynamic, Echocardiographic, and Neurohormonal Effects of Istaroxime, a Novel Intravenous Inotropic and Lusitropic Agent: A Randomized Controlled Trial in Patients Hospitalized with Heart Failure (HORIZON-HF) trial. *Am. Heart J.* **2009**, *157*, 1035–1041. [[CrossRef](#)] [[PubMed](#)]
104. Eisner, D.A.; Caldwell, J.L.; Trafford, A.W.; Hutchings, D.C. The Control of Diastolic Calcium in the Heart: Basic Mechanisms and Functional Implications. *Circ. Res.* **2020**, *126*, 395–412. [[CrossRef](#)]
105. Belhassen, B.; Pelleg, A.; Shoshani, D.; Laniado, S. Atrial fibrillation induced by adenosine triphosphate. *Am. J. Cardiol.* **1984**, *53*, 1405–1406. [[CrossRef](#)]
106. Tao, S.; Yamauchi, Y.; Maeda, S.; Okada, H.; Yamaguchi, T.; Hara, N.; Konishi, Y.; Umemoto, T.; Miyamoto, T.; Obayashi, T.; et al. Adenosine triphosphate-induced atrial fibrillation: The clinical significance and relevance to spontaneous atrial fibrillation. *J. Interv. Card. Electrophysiol.* **2014**, *39*, 103–109. [[CrossRef](#)]
107. Gilbert, G.; Demydenko, K.; Dries, E.; Puertas, R.D.; Jin, X.; Sipido, K.; Roderick, H.L. Calcium Signaling in Cardiomyocyte Function. *Cold Spring Harb. Perspect. Biol.* **2020**, *12*, 12. [[CrossRef](#)]
108. Farah, C.; Michel, L.Y.M.; Balligand, J.L. Nitric oxide signalling in cardiovascular health and disease. *Nat. Rev. Cardiol.* **2018**, *15*, 292–316. [[CrossRef](#)]
109. Vasan, R.S.; Sullivan, L.M.; Wilson, P.W.; Sempos, C.T.; Sundstrom, J.; Kannel, W.B.; Levy, D.; D'Agostino, R.B. Relative importance of borderline and elevated levels of coronary heart disease risk factors. *Ann. Intern. Med.* **2005**, *142*, 393–402. [[CrossRef](#)]
110. Group, S.R.; Wright, J.T., Jr.; Williamson, J.D.; Whelton, P.K.; Snyder, J.K.; Sink, K.M.; Rocco, M.V.; Reboussin, D.M.; Rahman, M.; Oparil, S.; et al. A Randomized Trial of Intensive versus Standard Blood-Pressure Control. *N. Engl. J. Med.* **2015**, *373*, 2103–2116. [[CrossRef](#)]
111. Rawshani, A.; Rawshani, A.; Franzén, S.; Sattar, N.; Eliasson, B.; Svensson, A.-M.; Zethelius, B.; Miftaraj, M.; McGuire, D.K.; Rosengren, A.; et al. Risk Factors, Mortality, and Cardiovascular Outcomes in Patients with Type 2 Diabetes. *N. Engl. J. Med.* **2018**, *379*, 633–644. [[CrossRef](#)]
112. Dzau, V.J.; Antman, E.M.; Black, H.R.; Hayes, D.L.; Manson, J.E.; Plutzky, J.; Popma, J.J.; Stevenson, W. The Cardiovascular Disease Continuum Validated: Clinical Evidence of Improved Patient Outcomes. *Circulation* **2006**, *114*, 2850–2870. [[CrossRef](#)]
113. Burnstock, G.; Novak, I. Purinergic signalling and diabetes. *Purinergic Signal.* **2013**, *9*, 307–324. [[CrossRef](#)] [[PubMed](#)]
114. Sparks, D.L.; Chatterjee, C. Purinergic signaling, dyslipidemia and inflammatory disease. *Cell Physiol. Biochem.* **2012**, *30*, 1333–1339. [[CrossRef](#)] [[PubMed](#)]

115. Novak, I.; Solini, A. P2X receptor-ion channels in the inflammatory response in adipose tissue and pancreas-potential triggers in onset of type 2 diabetes? *Curr. Opin. Immunol.* **2018**, *52*, 1–7. [[CrossRef](#)]
116. Ren, L.M.; Nakane, T.; Chiba, S. Muscarinic receptor subtypes mediating vasodilation and vasoconstriction in isolated, perfused simian coronary arteries. *J. Cardiovasc. Pharmacol.* **1993**, *22*, 841–846. [[CrossRef](#)] [[PubMed](#)]
117. Brock, J.A.; Van Helden, D.F. Enhanced excitatory junction potentials in mesenteric arteries from spontaneously hypertensive rats. *Pflugers Arch.* **1995**, *430*, 901–908. [[CrossRef](#)]
118. Vial, C.; Evans, R.J. P2X(1) receptor-deficient mice establish the native P2X receptor and a P2Y6-like receptor in arteries. *Mol. Pharmacol.* **2002**, *62*, 1438–1445. [[CrossRef](#)]
119. Lewis, C.J.; Evans, R.J. P2X receptor immunoreactivity in different arteries from the femoral, pulmonary, cerebral, coronary and renal circulations. *J. Vasc. Res.* **2001**, *38*, 332–340. [[CrossRef](#)]
120. Harhun, M.I.; Povstyan, O.V.; Albert, A.P.; Nichols, C.M. ATP-evoked sustained vasoconstrictions mediated by heteromeric P2X1/4 receptors in cerebral arteries. *Stroke* **2014**, *45*, 2444–2450. [[CrossRef](#)]
121. Milner, P.; Bodin, P.; Loesch, A.; Burnstock, G. Rapid release of endothelin and ATP from isolated aortic endothelial cells exposed to increased flow. *Biochem. Biophys. Res. Commun.* **1990**, *170*, 649–656. [[CrossRef](#)]
122. Yamamoto, K.; Sokabe, T.; Matsumoto, T.; Yoshimura, K.; Shibata, M.; Ohura, N.; Fukuda, T.; Sato, T.; Sekine, K.; Kato, S.; et al. Impaired flow-dependent control of vascular tone and remodeling in P2X4-deficient mice. *Nat. Med.* **2006**, *12*, 133–137. [[CrossRef](#)] [[PubMed](#)]
123. Piskuric, N.A.; Nurse, C.A. Expanding role of ATP as a versatile messenger at carotid and aortic body chemoreceptors. *J. Physiol.* **2013**, *591*, 415–422. [[CrossRef](#)] [[PubMed](#)]
124. Campanucci, V.A.; Zhang, M.; Vollmer, C.; Nurse, C.A. Expression of multiple P2X receptors by glossopharyngeal neurons projecting to rat carotid body O₂-chemoreceptors: Role in nitric oxide-mediated efferent inhibition. *J. Neurosci.* **2006**, *26*, 9482–9493. [[CrossRef](#)] [[PubMed](#)]
125. Pijacka, W.; Moraes, D.J.; Ratcliffe, L.E.; Nightingale, A.K.; Hart, E.C.; da Silva, M.P.; Machado, B.H.; McBryde, F.D.; Abdala, A.P.; Ford, A.P.; et al. Purinergic receptors in the carotid body as a new drug target for controlling hypertension. *Nat. Med.* **2016**, *22*, 1151–1159. [[CrossRef](#)] [[PubMed](#)]
126. Conde, S.V.; Monteiro, E.C.; Sacramento, J.F. Purines and Carotid Body: New Roles in Pathological Conditions. *Front. Pharmacol.* **2017**, *8*, 913. [[CrossRef](#)]
127. Libby, P.; Buring, J.E.; Badimon, L.; Hansson, G.K.; Deanfield, J.; Bittencourt, M.S.; Tokgozoglu, L.; Lewis, E.F. Atherosclerosis. *Nat. Rev. Dis. Primers* **2019**, *5*, 56. [[CrossRef](#)]
128. Stachon, P.; Hergeth, S.H.; Anto Michel, N.M.; Dufner, B.D.; Rodriguez, A.R.; Hoppe, N.H.; Wolf, D.W.; Bode, C.B.; Idzko, M.I.; Zirlik, A.Z. Extracellular atp contributes to atherogenesis via purinergic receptors by inducing leukocyte recruitment in mice. *Eur. Hear. J.* **2013**, *34*. [[CrossRef](#)]
129. Zhao, T.V.; Li, Y.; Liu, X.; Xia, S.; Shi, P.; Li, L.; Chen, Z.; Yin, C.; Eriguchi, M.; Chen, Y.; et al. ATP release drives heightened immune responses associated with hypertension. *Sci. Immunol.* **2019**, *4*, eaau6426. [[CrossRef](#)]
130. Sathanoori, R.; Sward, K.; Olde, B.; Erlinge, D. The ATP Receptors P2X7 and P2X4 Modulate High Glucose and Palmitate-Induced Inflammatory Responses in Endothelial Cells. *PLoS ONE* **2015**, *10*, e0125111. [[CrossRef](#)]
131. Pulvirenti, T.J.; Yin, J.L.; Chaufour, X.; McLachlan, C.; Hambly, B.D.; Bennett, M.R.; Barden, J.A. P2X (purinergic) receptor redistribution in rabbit aorta following injury to endothelial cells and cholesterol feeding. *J. Neurocytol.* **2000**, *29*, 623–631. [[CrossRef](#)] [[PubMed](#)]
132. Layhadi, J.A.; Turner, J.; Crossman, D.; Fountain, S.J. ATP Evokes Ca(2+) Responses and CXCL5 Secretion via P2X4 Receptor Activation in Human Monocyte-Derived Macrophages. *J. Immunol.* **2018**, *200*, 1159–1168. [[CrossRef](#)] [[PubMed](#)]
133. Rousselle, A.; Qadri, F.; Leukel, L.; Yilmaz, R.; Fontaine, J.F.; Sihh, G.; Bader, M.; Ahluwalia, A.; Duchene, J. CXCL5 limits macrophage foam cell formation in atherosclerosis. *J. Clin. Investig.* **2013**, *123*, 1343–1347. [[CrossRef](#)] [[PubMed](#)]
134. Sathanoori, R.; Rosi, F.; Gu, B.J.; Wiley, J.S.; Muller, C.E.; Olde, B.; Erlinge, D. Shear stress modulates endothelial KLF2 through activation of P2X4. *Purinergic Signal.* **2015**, *11*, 139–153. [[CrossRef](#)] [[PubMed](#)]
135. Ozaki, T.; Muramatsu, R.; Sasai, M.; Yamamoto, M.; Kubota, Y.; Fujinaka, T.; Yoshimine, T.; Yamashita, T. The P2X4 receptor is required for neuroprotection via ischemic preconditioning. *Sci. Rep.* **2016**, *6*, 25893. [[CrossRef](#)]
136. Anavekar, N.S.; McMurray, J.J.V.; Velazquez, E.J.; Solomon, S.D.; Kober, L.; Rouleau, J.-L.; White, H.D.; Nordlander, R.; Maggioni, A.; Dickstein, K.; et al. Relation between Renal Dysfunction and Cardiovascular Outcomes after Myocardial Infarction. *N. Engl. J. Med.* **2004**, *351*, 1285–1295. [[CrossRef](#)]

137. McAlister, F.A.; Ezekowitz, J.; Tarantini, L.; Squire, I.; Komajda, M.; Bayes-Genis, A.; Gotsman, I.; Whalley, G.; Earle, N.; Poppe, K.K.; et al. Renal dysfunction in patients with heart failure with preserved versus reduced ejection fraction: Impact of the new Chronic Kidney Disease-Epidemiology Collaboration Group formula. *Circ. Heart Fail.* **2012**, *5*, 309–314. [[CrossRef](#)]
138. Legrand, M.; Rossignol, P. Cardiovascular Consequences of Acute Kidney Injury. *N. Engl. J. Med.* **2020**, *382*, 2238–2247. [[CrossRef](#)]
139. Rangaswami, J.; Bhalla, V.; Blair, J.E.A.; Chang, T.I.; Costa, S.; Lentine, K.L.; Lerma, E.V.; Mezue, K.; Molitch, M.; Mullens, W.; et al. Cardiorenal Syndrome: Classification, Pathophysiology, Diagnosis, and Treatment Strategies: A Scientific Statement From the American Heart Association. *Circulation* **2019**, *139*, e840–e878. [[CrossRef](#)]
140. Burnstock, G.; Evans, L.C.; Bailey, M.A. Purinergic signalling in the kidney in health and disease. *Purinergic Signal.* **2014**, *10*, 71–101. [[CrossRef](#)]
141. Carlstrom, M.; Wilcox, C.S.; Arendshorst, W.J. Renal autoregulation in health and disease. *Physiol. Rev.* **2015**, *95*, 405–511. [[CrossRef](#)]
142. Fernandez, O.; Wangenstein, R.; Osuna, A.; Vargas, F. Renal vascular reactivity to P(2)-purinoceptor activation in spontaneously hypertensive rats. *Pharmacology* **2000**, *60*, 47–50. [[CrossRef](#)] [[PubMed](#)]
143. Menzies, R.I.; Unwin, R.J.; Dash, R.K.; Beard, D.A.; Cowley, A.W., Jr.; Carlson, B.E.; Mullins, J.J.; Bailey, M.A. Effect of P2X4 and P2X7 receptor antagonism on the pressure diuresis relationship in rats. *Front. Physiol.* **2013**, *4*, 305. [[CrossRef](#)] [[PubMed](#)]
144. Menzies, R.I.; Unwin, R.J.; Bailey, M.A. Renal P2 receptors and hypertension. *Acta Physiol.* **2015**, *213*, 232–241. [[CrossRef](#)]
145. Kowalski, R.; Kreft, E.; Kasztan, M.; Jankowski, M.; Szczepanska-Konkel, M. Chronic renal denervation increases renal tubular response to P2X receptor agonists in rats: Implication for renal sympathetic nerve ablation. *Nephrol. Dial. Transplant.* **2012**, *27*, 3443–3448. [[CrossRef](#)]
146. Wildman, S.S.; Marks, J.; Churchill, L.J.; Peppiatt, C.M.; Chraibi, A.; Shirley, D.G.; Horisberger, J.D.; King, B.F.; Unwin, R.J. Regulatory interdependence of cloned epithelial Na⁺ channels and P2X receptors. *J. Am. Soc. Nephrol.* **2005**, *16*, 2586–2597. [[CrossRef](#)] [[PubMed](#)]
147. Wildman, S.S.; Marks, J.; Turner, C.M.; Yew-Booth, L.; Peppiatt-Wildman, C.M.; King, B.F.; Shirley, D.G.; Wang, W.; Unwin, R.J. Sodium-dependent regulation of renal amiloride-sensitive currents by apical P2 receptors. *J. Am. Soc. Nephrol.* **2008**, *19*, 731–742. [[CrossRef](#)]
148. Craigie, E.; Menzies, R.I.; Larsen, C.K.; Jacquillet, G.; Carrel, M.; Wildman, S.S.; Loffing, J.; Leipziger, J.; Shirley, D.G.; Bailey, M.A.; et al. The renal and blood pressure response to low sodium diet in P2X4 receptor knockout mice. *Physiol. Rep.* **2018**, *6*, e13899. [[CrossRef](#)]
149. Howarth, A.R.; Conway, B.R.; Bailey, M.A. Vascular and inflammatory actions of P2X receptors in renal injury. *Auton. Neurosci.* **2015**, *191*, 135–140. [[CrossRef](#)]
150. Han, S.J.; Lovaszi, M.; Kim, M.; D’Agati, V.; Hasko, G.; Lee, H.T. P2X4 receptor exacerbates ischemic AKI and induces renal proximal tubular NLRP3 inflammasome signaling. *FASEB J.* **2020**, *34*, 5465–5482. [[CrossRef](#)]
151. Kim, M.J.; Turner, C.M.; Hewitt, R.; Smith, J.; Bhangal, G.; Pusey, C.D.; Unwin, R.J.; Tam, F.W. Exaggerated renal fibrosis in P2X4 receptor-deficient mice following unilateral ureteric obstruction. *Nephrol. Dial. Transplant.* **2014**, *29*, 1350–1361. [[CrossRef](#)] [[PubMed](#)]
152. Cohn, J.N.; Quyyumi, A.A.; Hollenberg, N.K.; Jamerson, K.A. Surrogate markers for cardiovascular disease: Functional markers. *Circulation* **2004**, *109*, IV31–IV46. [[CrossRef](#)]
153. Forst, A.L.; Olteanu, V.S.; Mollet, G.; Wlodkowski, T.; Schaefer, F.; Dietrich, A.; Reiser, J.; Gudermann, T.; Mederos y Schnitzler, M.; Storch, U. Podocyte Purinergic P2X4 Channels Are Mechanotransducers That Mediate Cytoskeletal Disorganization. *J. Am. Soc. Nephrol.* **2016**, *27*, 848–862. [[CrossRef](#)]
154. Chen, K.; Zhang, J.; Zhang, W.; Zhang, J.; Yang, J.; Li, K.; He, Y. ATP-P2X4 signaling mediates NLRP3 inflammasome activation: A novel pathway of diabetic nephropathy. *Int. J. Biochem. Cell Biol.* **2013**, *45*, 932–943. [[CrossRef](#)]
155. Forfia, P.R.; Vaidya, A.; Wiegers, S.E. Pulmonary heart disease: The heart-lung interaction and its impact on patient phenotypes. *Pulm. Circ.* **2013**, *3*, 5–19. [[CrossRef](#)]

156. Galie, N.; Humbert, M.; Vachiery, J.L.; Gibbs, S.; Lang, I.; Torbicki, A.; Simonneau, G.; Peacock, A.; Vonk Noordegraaf, A.; Beghetti, M.; et al. 2015 ESC/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension: The Joint Task Force for the Diagnosis and Treatment of Pulmonary Hypertension of the European Society of Cardiology (ESC) and the European Respiratory Society (ERS): Endorsed by: Association for European Paediatric and Congenital Cardiology (AEPC), International Society for Heart and Lung Transplantation (ISHLT). *Eur. Heart J.* **2016**, *37*, 67–119. [[CrossRef](#)]
157. Guibert, C.; Pacaud, P.; Loirand, G.; Marthan, R.; Savineau, J.P. Effect of extracellular ATP on cytosolic Ca²⁺ concentration in rat pulmonary artery myocytes. *Am. J. Physiol.* **1996**, *271*, L450–L458. [[CrossRef](#)]
158. Neely, C.F.; Matot, I.; Batra, V.K.; Bo, X.; Burnstock, G. P2X purinoceptors in the feline pulmonary vascular bed: Distribution and selective in vivo pharmacological probes. *Am. J. Physiol.* **1996**, *270*, L889–L897. [[CrossRef](#)]
159. Visovatti, S.H.; Hyman, M.C.; Goonewardena, S.N.; Anyanwu, A.C.; Kanthi, Y.; Robichaud, P.; Wang, J.; Petrovic-Djergovic, D.; Rattan, R.; Burant, C.F.; et al. Purinergic dysregulation in pulmonary hypertension. *Am. J. Physiol. Heart Circ. Physiol.* **2016**, *311*, H286–H298. [[CrossRef](#)] [[PubMed](#)]
160. Syed, N.I.; Tengah, A.; Paul, A.; Kennedy, C. Characterisation of P2X receptors expressed in rat pulmonary arteries. *Eur. J. Pharmacol.* **2010**, *649*, 342–348. [[CrossRef](#)] [[PubMed](#)]
161. Qasabian, R.A.; Schyvens, C.; Owe-Young, R.; Killen, J.P.; Macdonald, P.S.; Conigrave, A.D.; Williamson, D.J. Characterization of the P2 receptors in rabbit pulmonary artery. *Br. J. Pharmacol.* **1997**, *120*, 553–558. [[CrossRef](#)] [[PubMed](#)]
162. Yamamoto, K.; Sokabe, T.; Ohura, N.; Nakatsuka, H.; Kamiya, A.; Ando, J. Endogenously released ATP mediates shear stress-induced Ca²⁺ influx into pulmonary artery endothelial cells. *Am. J. Physiol. Heart Circ. Physiol.* **2003**, *285*, H793–H803. [[CrossRef](#)] [[PubMed](#)]
163. Yamamoto, K.; Imamura, H.; Ando, J. Shear stress augments mitochondrial ATP generation that triggers ATP release and Ca(2+) signaling in vascular endothelial cells. *Am. J. Physiol. Heart Circ. Physiol.* **2018**, *315*, H1477–H1485. [[CrossRef](#)]
164. Gao, Z.G.; Jacobson, K.A. Purinergic Signaling in Mast Cell Degranulation and Asthma. *Front. Pharmacol.* **2017**, *8*, 947. [[CrossRef](#)] [[PubMed](#)]
165. Burnstock, G.; Brouns, I.; Adriaensen, D.; Timmermans, J.P. Purinergic signaling in the airways. *Pharmacol. Rev.* **2012**, *64*, 834–868. [[CrossRef](#)]
166. Antonioli, L.; Blandizzi, C.; Pacher, P.; Hasko, G. The Purinergic System as a Pharmacological Target for the Treatment of Immune-Mediated Inflammatory Diseases. *Pharmacol. Rev.* **2019**, *71*, 345–382. [[CrossRef](#)]
167. Le, T.T.; Berg, N.K.; Harting, M.T.; Li, X.; Eltzschig, H.K.; Yuan, X. Purinergic Signaling in Pulmonary Inflammation. *Front. Immunol.* **2019**, *10*, 1633. [[CrossRef](#)]
168. Celutkienė, J.; Balciunas, M.; Kablucko, D.; Vaitkeviciute, L.; Blasciuk, J.; Danila, E. Challenges of Treating Acute Heart Failure in Patients with Chronic Obstructive Pulmonary Disease. *Card Fail. Rev.* **2017**, *3*, 56–61. [[CrossRef](#)]
169. Eltzschig, H.K.; Thompson, L.F.; Karhausen, J.; Cotta, R.J.; Ibla, J.C.; Robson, S.C.; Colgan, S.P. Endogenous adenosine produced during hypoxia attenuates neutrophil accumulation: Coordination by extracellular nucleotide metabolism. *Blood* **2004**, *104*, 3986–3992. [[CrossRef](#)] [[PubMed](#)]
170. Winkelmann, V.E.; Thompson, K.E.; Neuland, K.; Jaramillo, A.M.; Fois, G.; Schmidt, H.; Wittekindt, O.H.; Han, W.; Tuvim, M.J.; Dickey, B.F.; et al. Inflammation-induced upregulation of P2X4 expression augments mucin secretion in airway epithelia. *Am. J. Physiol. Lung Cell Mol. Physiol.* **2019**, *316*, L58–L70. [[CrossRef](#)]
171. Burnstock, G. P2X ion channel receptors and inflammation. *Purinergic Signal.* **2016**, *12*, 59–67. [[CrossRef](#)] [[PubMed](#)]
172. Zabala, A.; Vazquez-Villoldo, N.; Rissiek, B.; Gejo, J.; Martin, A.; Palomino, A.; Perez-Samartin, A.; Pulagam, K.R.; Lukowiak, M.; Capetillo-Zarate, E.; et al. P2X4 receptor controls microglia activation and favors remyelination in autoimmune encephalitis. *EMBO Mol. Med.* **2018**, *10*. [[CrossRef](#)] [[PubMed](#)]
173. Zech, A.; Wiesler, B.; Ayata, C.K.; Schlaich, T.; Durk, T.; Hossfeld, M.; Ehrat, N.; Cicko, S.; Idzko, M. P2rx4 deficiency in mice alleviates allergen-induced airway inflammation. *Oncotarget* **2016**, *7*, 80288–80297. [[CrossRef](#)]
174. Csoka, B.; Nemeth, Z.H.; Szabo, I.; Davies, D.L.; Varga, Z.V.; Paloczi, J.; Falzoni, S.; Di Virgilio, F.; Muramatsu, R.; Yamashita, T.; et al. Macrophage P2X4 receptors augment bacterial killing and protect against sepsis. *JCI Insight* **2018**, *3*, 3. [[CrossRef](#)] [[PubMed](#)]

175. Chen, H.; Xia, Q.; Feng, X.; Cao, F.; Yu, H.; Song, Y.; Ni, X. Effect of P2X4R on airway inflammation and airway remodeling in allergic airway challenge in mice. *Mol. Med. Rep.* **2016**, *13*, 697–704. [[CrossRef](#)] [[PubMed](#)]
176. Ledderose, C.; Liu, K.; Kondo, Y.; Slubowski, C.J.; Dertnig, T.; Denicolo, S.; Arbab, M.; Hubner, J.; Konrad, K.; Fakhari, M.; et al. Purinergic P2X4 receptors and mitochondrial ATP production regulate T cell migration. *J. Clin. Investig.* **2018**, *128*, 3583–3594. [[CrossRef](#)] [[PubMed](#)]
177. Paalme, V.; Rump, A.; Mado, K.; Teras, M.; Truumees, B.; Aitai, H.; Ratas, K.; Bourge, M.; Chiang, C.S.; Ghalali, A.; et al. Human Peripheral Blood Eosinophils Express High Levels of the Purinergic Receptor P2X4. *Front. Immunol.* **2019**, *10*, 2074. [[CrossRef](#)] [[PubMed](#)]
178. Nagaoka, M.; Nara, M.; Tamada, T.; Kume, H.; Oguma, T.; Kikuchi, T.; Zaini, J.; Moriya, T.; Ichinose, M.; Tamura, G.; et al. Regulation of adenosine 5'-triphosphate (ATP)-gated P2X(4) receptors on tracheal smooth muscle cells. *Respir. Physiol. Neurobiol.* **2009**, *166*, 61–67. [[CrossRef](#)]
179. Mishra, A.; Chintagari, N.R.; Guo, Y.; Weng, T.; Su, L.; Liu, L. Purinergic P2X7 receptor regulates lung surfactant secretion in a paracrine manner. *J. Cell Sci.* **2011**, *124*, 657–668. [[CrossRef](#)]
180. Patel, A.S.; Reigada, D.; Mitchell, C.H.; Bates, S.R.; Margulies, S.S.; Koval, M. Paracrine stimulation of surfactant secretion by extracellular ATP in response to mechanical deformation. *Am. J. Physiol. Lung Cell Mol. Physiol.* **2005**, *289*, L489–L496. [[CrossRef](#)]
181. Han, S.; Mallampalli, R.K. The Role of Surfactant in Lung Disease and Host Defense against Pulmonary Infections. *Ann. Am. Thorac Soc.* **2015**, *12*, 765–774. [[CrossRef](#)]
182. Miklavc, P.; Mair, N.; Wittekindt, O.H.; Haller, T.; Diel, P.; Felder, E.; Timmler, M.; Frick, M. Fusion-activated Ca²⁺ entry via vesicular P2X4 receptors promotes fusion pore opening and exocytotic content release in pneumocytes. *Proc. Natl. Acad. Sci. USA* **2011**, *108*, 14503–14508. [[CrossRef](#)] [[PubMed](#)]
183. Thompson, K.E.; Korbmayer, J.P.; Hecht, E.; Hobi, N.; Wittekindt, O.H.; Diel, P.; Kranz, C.; Frick, M. Fusion-activated cation entry (FACE) via P2X(4) couples surfactant secretion and alveolar fluid transport. *FASEB J.* **2013**, *27*, 1772–1783. [[CrossRef](#)] [[PubMed](#)]
184. Fois, G.; Winkelmann, V.E.; Bareis, L.; Staudenmaier, L.; Hecht, E.; Ziller, C.; Ehinger, K.; Schymeinsky, J.; Kranz, C.; Frick, M. ATP is stored in lamellar bodies to activate vesicular P2X4 in an autocrine fashion upon exocytosis. *J. Gen. Physiol.* **2018**, *150*, 277–291. [[CrossRef](#)]
185. Hasan, D.; Satalin, J.; van der Zee, P.; Kollisch-Singule, M.; Blankman, P.; Shono, A.; Somhorst, P.; den Uil, C.; Meeder, H.; Kotani, T.; et al. Excessive Extracellular ATP Desensitizes P2Y2 and P2X4 ATP Receptors Provoking Surfactant Impairment Ending in Ventilation-Induced Lung Injury. *Int. J. Mol. Sci.* **2018**, *19*, 1185. [[CrossRef](#)] [[PubMed](#)]
186. Rajagopal, K.; Keller, S.P.; Akkanti, B.; Bime, C.; Loyalka, P.; Cheema, F.H.; Zwischenberger, J.B.; El Banayosy, A.; Pappalardo, F.; Slaughter, M.S.; et al. Advanced Pulmonary and Cardiac Support of COVID-19 Patients: Emerging Recommendations From ASAIO-a Living Working Document. *Circ. Heart Fail.* **2020**, *13*, e007175. [[CrossRef](#)]
187. Matthay, M.A.; Zemans, R.L.; Zimmerman, G.A.; Arabi, Y.M.; Beitler, J.R.; Mercat, A.; Herridge, M.; Randolph, A.G.; Calfee, C.S. Acute respiratory distress syndrome. *Nat. Rev. Dis. Primers* **2019**, *5*, 18. [[CrossRef](#)]
188. Albert, R.K. The role of ventilation-induced surfactant dysfunction and atelectasis in causing acute respiratory distress syndrome. *Am. J. Respir Crit Care Med.* **2012**, *185*, 702–708. [[CrossRef](#)]
189. Stokes, L.; Layhadi, J.A.; Bibic, L.; Dhuna, K.; Fountain, S.J. P2X4 Receptor Function in the Nervous System and Current Breakthroughs in Pharmacology. *Front. Pharmacol.* **2017**, *8*, 291. [[CrossRef](#)]
190. Zhang, W.J.; Zhu, Z.M.; Liu, Z.X. The role of P2X4 receptor in neuropathic pain and its pharmacological properties. *Pharmacol. Res.* **2020**, *158*, 104875. [[CrossRef](#)]
191. Tsuda, M.; Shigemoto-Mogami, Y.; Koizumi, S.; Mizokoshi, A.; Kohsaka, S.; Salter, M.W.; Inoue, K. P2X4 receptors induced in spinal microglia gate tactile allodynia after nerve injury. *Nature* **2003**, *424*, 778–783. [[CrossRef](#)] [[PubMed](#)]
192. Ulmann, L.; Hatcher, J.P.; Hughes, J.P.; Chaumont, S.; Green, P.J.; Conquet, F.; Buell, G.N.; Reeve, A.J.; Chessell, I.P.; Rassendren, F. Up-regulation of P2X4 receptors in spinal microglia after peripheral nerve injury mediates BDNF release and neuropathic pain. *J. Neurosci.* **2008**, *28*, 11263–11268. [[CrossRef](#)] [[PubMed](#)]
193. Ulmann, L.; Hirbec, H.; Rassendren, F. P2X4 receptors mediate PGE2 release by tissue-resident macrophages and initiate inflammatory pain. *EMBO J.* **2010**, *29*, 2290–2300. [[CrossRef](#)]

194. Long, T.; He, W.; Pan, Q.; Zhang, S.; Zhang, D.; Qin, G.; Chen, L.; Zhou, J. Microglia P2X4R-BDNF signalling contributes to central sensitization in a recurrent nitroglycerin-induced chronic migraine model. *J. Headache Pain* **2020**, *21*, 4. [[CrossRef](#)]
195. Beggs, S.; Trang, T.; Salter, M.W. P2X4R+ microglia drive neuropathic pain. *Nat. Neurosci.* **2012**, *15*, 1068–1073. [[CrossRef](#)]
196. Sim, J.A.; Chaumont, S.; Jo, J.; Ulmann, L.; Young, M.T.; Cho, K.; Buell, G.; North, R.A.; Rassendren, F. Altered hippocampal synaptic potentiation in P2X4 knock-out mice. *J. Neurosci.* **2006**, *26*, 9006–9009. [[CrossRef](#)]
197. Baxter, A.W.; Choi, S.J.; Sim, J.A.; North, R.A. Role of P2X4 receptors in synaptic strengthening in mouse CA1 hippocampal neurons. *Eur. J. Neurosci.* **2011**, *34*, 213–220. [[CrossRef](#)] [[PubMed](#)]
198. de Rivero Vaccari, J.P.; Bastien, D.; Yurcisin, G.; Pineau, I.; Dietrich, W.D.; De Koninck, Y.; Keane, R.W.; Lacroix, S. P2X4 receptors influence inflammasome activation after spinal cord injury. *J. Neurosci.* **2012**, *32*, 3058–3066. [[CrossRef](#)] [[PubMed](#)]
199. Ulmann, L.; Levavasseur, F.; Avignone, E.; Peyroutou, R.; Hirbec, H.; Audinat, E.; Rassendren, F. Involvement of P2X4 receptors in hippocampal microglial activation after status epilepticus. *Glia* **2013**, *61*, 1306–1319. [[CrossRef](#)] [[PubMed](#)]
200. Khoja, S.; Huynh, N.; Warnecke, A.M.P.; Asatryan, L.; Jakowec, M.W.; Davies, D.L. Preclinical evaluation of avermectins as novel therapeutic agents for alcohol use disorders. *Psychopharmacology* **2018**, *235*, 1697–1709. [[CrossRef](#)] [[PubMed](#)]
201. Franklin, K.M.; Asatryan, L.; Jakowec, M.W.; Trudell, J.R.; Bell, R.L.; Davies, D.L. P2X4 receptors (P2X4Rs) represent a novel target for the development of drugs to prevent and/or treat alcohol use disorders. *Front. Neurosci.* **2014**, *8*, 176. [[CrossRef](#)] [[PubMed](#)]
202. Li, F.; Guo, N.; Ma, Y.; Ning, B.; Wang, Y.; Kou, L. Inhibition of P2X4 suppresses joint inflammation and damage in collagen-induced arthritis. *Inflammation* **2014**, *37*, 146–153. [[CrossRef](#)]



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).