

Resonance of Continued Fractions Related to ${}_2\psi_2$ Basic Bilateral Hypergeometric Series

PANKAJ SRIVASTAVA

*Department of Mathematics, Motilal Nehru National Institute of Technology,
Allahabad-211004, India*

e-mail: drpankaj23@gmail.com; pankajs23@rediffmail.com

ABSTRACT. In this paper, making use of transformation due to S. N. Singh [21], an attempt has been made to establish certain results involving basic bilateral hypergeometric series and continued fractions.

1. Introduction

Continued fractions offer a useful means of expressing numbers and functions. The Indian mathematician Arya Bhatt(475-550AD) used a continued fraction to solve a linear equation[15]. Continued fraction were used only in specific examples for more than one thousand years. In the beginning of the 20th century, the theory had advanced due to the Indian genius S. Ramanujan[17, 18]. Generalized hypergeometric series both ordinary and basic have been a very significant tool in the derivation of continued fraction representation. In the chapter 16 of Ramanujan's second notebook, first seventeen sections are devoted mainly to basic hypergeometric functions and chapter 12 of his second notebook is entirely devoted to the study of continued fractions. Ramanujan has made significant contribution to the theory of continued fractions making use of basic hypergeometric function as significant tool and provided a new dimension to this interesting branch of analysis. During the end of the 20th century continued fraction had been center of attraction for the mathematicians working in the field of q -series namely R. P. Agarwal [1], George E. Andrews [2], Bruce C. Berndt[7], S. Bhargava and C. S., Adiga[9], R. Y. Denis [10, 11, 12, 13], C. S. Adiga and D. D. Somashekara [3, 4], S. N. Singh [19, 20], Pankaj Srivastava [22], Maheshwar Pathak and Pankaj Srivastava [16], Bhaskar Srivastava [23], L. C. Zhang[24] etc. To investigate continued fraction representation for the ratio of two hypergeometric function ${}_2F_1$, basic hypergeometric series ${}_2\phi_1$, and ${}_3\phi_2$, several mathematicians namely R. P. Agarwal [1], R. Y. Denis [10, 11, 12, 13], S. N. Singh [19, 20], N. A. Bhagirathi [5, 6], S. Bhargava, C. S. Adiga

Received August 17, 2010; accepted April 26, 2011.

2010 Mathematics Subject Classification: 33D15, 11A55.

Key words and phrases: Basic hypergeometric series, basic bilateral series, continued fractions.

and D. D. Somashekara [8] established a number of results. In the present paper, transformation due to S. N. Singh [21] has been used to establish new continued fraction representation for the ratio of basic bilateral hypergeometric series ${}_2\psi_2$. S. N. Singh [21] establish the following general transformation for basic bilateral series into basic series:

$$(1.1) \quad {}_{r+3}\psi_{r+3} \left[\begin{matrix} \frac{a}{b}, & cq, & dq, & \frac{b_1}{b}q^{m_1}, & \dots, & \frac{b_r}{b}q^{m_r} \\ \frac{q}{b}, & c, & d, & \frac{b_1}{b}, & \dots, & \frac{b_r}{b} \end{matrix} ; q; z \right] =$$

$$\frac{\left(q, \frac{bq}{az}, \frac{az}{b}, \frac{q}{a}; q \right)_{\infty}}{\left(\frac{q}{b}, \frac{q}{az}, az, \frac{bq}{a}; q \right)_{\infty}} \times \frac{(1-bc)(1-bd)(b_1; q)_{m_1} \dots (b_r; q)_{m_r}}{(1-c)(1-d)(b_1/b; q)_{m_1} \dots (b_r/b; q)_{m_r}}$$

$$\times {}_{r+3}\phi_{r+2} \left[\begin{matrix} a, & bcq, & bdq, & b_1q^{m_1}, & \dots, & b_rq^{m_r} \\ bc, & bd, & b_1, & \dots, & b_r \end{matrix} ; q; z \right]$$

2. Definitions and notations

A basic hypergeometric series is defined as:

$$(2.1) \quad {}_r\phi_s \left[\begin{matrix} a_1, & \dots, & a_r \\ b_1, & \dots, & b_s \end{matrix} ; q; z \right] = \sum_{n=0}^{\infty} \frac{(a_1)_n (a_2)_n \dots (a_r)_n}{(b_1)_n (b_2)_n \dots (b_s)_n} \frac{z^n}{(q)_n}$$

subject to appropriate convergence series conditions, where $|q| < 1$.

$$(2.2) \quad (a)_n = (a; q)_n = \begin{cases} (1-a)(1-aq)\dots(1-aq^{n-1}), & n \geq 1 \\ 1, & n = 0 \end{cases}$$

A basic bilateral hypergeometric series is defined as:

$$(2.3) \quad {}_r\psi_r \left[\begin{matrix} a_1, & \dots, & a_r \\ b_1, & \dots, & b_r \end{matrix} ; q; z \right] = \sum_{n=-\infty}^{\infty} \frac{(a_1, a_2, \dots, a_r; q)_n}{(b_1, b_2, \dots, b_r; q)_n} z^n$$

where $|b_1 b_2 \dots b_r / a_1 a_2 \dots a_r| < |z| < 1$ for convergence. An expression of the form

$$(2.4) \quad \frac{a_1}{b_1 +} \frac{a_2}{b_2 +} \frac{a_3}{b_3 +} \dots \frac{a_n}{b_n}$$

is said to be a terminating continued fraction and as $n \rightarrow \infty$, it is said to be an infinite continued fraction.

We shall make use of following results in our analysis:

$$(2.5) \quad \frac{{}_2\phi_1 \left[\begin{matrix} \alpha, & \beta \\ & \gamma \end{matrix} ; xq \right]}{{}_2\phi_1 \left[\begin{matrix} \alpha, & \beta \\ & \gamma \end{matrix} ; x \right]} = \frac{1}{1+} \frac{(1-\alpha)(1-\beta)x}{(1-x)+} \frac{\alpha\beta xq - \gamma}{1+} \frac{(1-\alpha q)(1-\beta q)x}{(1-x)+}$$

$$+ \frac{\alpha\beta xq^3 - \gamma q}{1+} \frac{(1-\alpha q^2)(1-\beta q^2)x}{(1-x)+ \dots}$$

[Denis, [10]]

$$(2.6) \quad \frac{{}_2\phi_1 [a, b ; cq; x]}{{}_2\phi_1 [a, b ; c; x]} = \frac{1}{1+} \frac{xc(1-a)(1-b)/(1-c)(1-cq)}{(1-abx/cq)+}$$

$$+ \frac{x(a-cq)(b-cq)/cq(1-cq)(1-cq^2)}{1+} \frac{cxq^2(1-aq)(1-bq)/(1-cq^2)(1-cq^3)}{(1-abx/cq)+ \dots}$$

[R. P. Agarwal, [1]]

$$(2.7) \quad \frac{{}_2\phi_1 [a, bq ; c; x]}{{}_2\phi_1 [a, b ; c; xq]} = 1 + \frac{x(1-a)}{(1-x)(1-b)+} \frac{(b-c)/(1-axq)}{1+}$$

$$+ \frac{x(1-aq)}{(1-x)(1-b)+} \frac{(b-cq)(1-axq^2)}{1+ \dots}$$

[R. P. Agarwal, [1]]

$$(2.8) \quad \frac{{}_2\phi_1 [a, b ; c; xq]}{{}_2\phi_1 [aq, bq ; cq; x]} = \frac{(1-x)}{(1-c)-} \frac{c(1-abxq/c)}{(1-c)+} \frac{x(1-aq)(1-bq)}{(1-x)/(1-c)-}$$

$$- \frac{cq(1-abxq^2/c)}{(1-c)+} \frac{x(1-aq^2)(1-bq^2)}{(1-x)/(1-c)-} \frac{cq^2(1-abxq^3/c)}{(1-c)+ \dots}$$

[R. P. Agarwal, [1]]

$$(2.9) \quad \frac{{}_2\phi_1 [\alpha, \beta ; \gamma; z]}{{}_2\phi_1 [\alpha, \beta q ; \gamma; z]} = 1 - \frac{z\beta(1-\alpha)}{(1-\gamma/\beta q)+} \frac{\gamma(1-\beta q)(1-\alpha\beta zq/\gamma)/\beta q}{1-}$$

$$- \frac{z\beta q(1-\alpha q)}{(1-\gamma/\beta q)+} \frac{\gamma(1-\beta q^2)(1-\alpha\beta zq^2/\gamma)/\beta q}{1-} \frac{z\beta q^2(1-\alpha q^2)}{(1-\gamma/\beta q)+ \dots}$$

[S. N. Singh, [19]]

$$(2.10) \quad \frac{{}_2\phi_1 [\alpha, \beta ; \gamma; z]}{{}_2\phi_1 [\alpha, \beta ; \gamma q; z]} = 1 + \frac{z\gamma(1-\alpha)(1-\beta)/(1-\gamma)(1-\gamma q)}{(1-\alpha\beta z/\gamma q)+}$$

$$+ \frac{\mu(\alpha - \gamma q)/(\beta - \gamma q)/(1 - \gamma q)(1 - \gamma q^2)}{1 + \dots}$$

[S. N. Singh, [19]]

where $\mu = \frac{z}{\gamma q}$

$$(2.11) \quad \frac{{}_2\phi_1 \left[\begin{matrix} \alpha, & \beta q & ; \gamma; & x \end{matrix} \right]}{{}_2\phi_1 \left[\begin{matrix} \alpha, & \beta & ; \gamma; & xq \end{matrix} \right]} = 1 + \frac{x(1 - \alpha)}{(1 - x)(1 - \beta) +} \frac{(\beta - \gamma)/(1 - \alpha x q)}{1 +}$$

$$+ \frac{x(1 - \alpha q)}{(1 - x)(1 - \beta) +} \frac{(\beta - \gamma q)/(1 - \alpha x q^2)}{1 +} \frac{x(1 - \alpha q^2)}{(1 - x)(1 - \beta) + \dots}$$

[S. N. Singh, [20]]

$$(2.12) \quad \frac{(1 - cq^i) {}_2\phi_1 \left[\begin{matrix} aq^i, & bq^i & ; x \end{matrix} \right]}{(1 - c) {}_2\phi_1 \left[\begin{matrix} aq^{i+1}, & bq^{i+1} & ; x/q \end{matrix} \right]} = \frac{(1 - x/q)}{(1 - c)} - \frac{(1 - abxq^i/c)cq^i}{(1 - c) +}$$

$$+ \frac{x(1 - aq^{i+1})(1 - bq^{i+1})/b}{(1 - x/q)(1 - c) -} \frac{(1 - abxq^{i+1}/c)cq^{i+1}}{(1 - c) +} \frac{x(1 - aq^{i+2})(1 - bq^{i+2})/q}{(1 - x/q)/(1 - c) - \dots}$$

[Bhagirathi, [5]]

$$(2.13) \quad \frac{(1 - c) {}_2\phi_1 \left[\begin{matrix} aq^{i+1}, & bq^{i+1} & ; x/q \end{matrix} \right]}{{}_2\phi_1 \left[\begin{matrix} aq^{i+1}, & bq^{i+1} & ; x \end{matrix} \right]} =$$

$$(1 - c) + \frac{(1 - aq^{i+1})(1 - bq^{i+1})/q}{(1 - x/q)(1 - c) -} \frac{(1 - abxq^{i+1}/c)cq^{i+1}}{(1 - c) +}$$

$$+ \frac{x(1 - aq^{i+2})(1 - bq^{i+2})/q}{(1 - x/q)/(1 - c) -} \frac{(1 - abxq^{i+2}/c)cq^{i+2}}{(1 - c) +} \frac{x(1 - aq^{i+3})(1 - bq^{i+3})/q}{(1 - x/q)/(1 - c) - \dots}$$

[Bhagirathi, [5]]

$$(2.14) \quad \frac{\frac{(1 - cq^i)}{(1 - c)} {}_2\phi_1 \left[\begin{matrix} aq^i, & bq^i & ; x \end{matrix} \right]}{{}_2\phi_1 \left[\begin{matrix} aq^{i+1}, & bq^i & ; x/q \end{matrix} \right]}} = \frac{(1 - c/b)}{(1 - c) +} \frac{c(1 - bq^i)(1 - abxq^i/c)/b}{(1 - c) -}$$

$$\frac{(1 - aq^{i+1})xbq^i}{(1 - c/b)/(1 - c)+} \frac{c(1 - bq^{i+1})(1 - abxq^{i+1}/c)/b}{(1 - c)-} \frac{(1 - aq^{i+2})xbq^{i+1}}{(1 - c/b)/(1 - c) + \dots}$$

[Bhagirathi, [5]]

3. Main results

The following results are to be established in this section:

(3.1)
$$\frac{{}_2\psi_2 \left[\begin{matrix} \frac{a}{b}, & \frac{b_1}{b}q^{m_1} \\ \frac{q}{b}, & \frac{b_1}{b}q \end{matrix} ; q; z \right]}{{}_2\psi_2 \left[\begin{matrix} \frac{a}{b}, & \frac{b_1}{b}q^{m_1} \\ \frac{q}{b}, & \frac{b_1}{b} \end{matrix} ; q; z \right]} = \frac{b}{1+} \frac{(1 - a)(1 - b_1q^{m_1})z}{(1 - z)+} \frac{b_1(azq^{m_1+1} - 1)}{1+}$$

$$+ \frac{(1 - aq)(1 - b_1q^{m_1+1})z}{(1 - z)+} \frac{b_1q(zaq^{m_1+3} - 1)}{1+} \frac{(1 - aq^2)(1 - b_1q^{m_1+2})z}{(1 - z) + \dots}$$

(3.2)
$$\frac{{}_2\psi_2 \left[\begin{matrix} \frac{a}{b}, & \frac{b_1}{b}q^{m_1+1} \\ \frac{q}{b}, & \frac{b_1}{b}q \end{matrix} ; q; z \right]}{{}_2\psi_2 \left[\begin{matrix} \frac{a}{b}, & \frac{b_1}{b}q^{m_1} \\ \frac{q}{b}, & \frac{b_1}{b} \end{matrix} ; q; z \right]} =$$

$$\frac{(b - b_1)/b(1 - b_1)zb_1(1 - a)(1 - b_1q^{m_1})/(1 - b_1)(1 - b_1q)z(a - b_1q)(q^{m_1-1} - 1)/(1 - b_1q)(1 - b_1q^2)}{1+} \frac{(1 - azq^{m_1-1})+}{1+}$$

$$+ \frac{b_1zq^2(1 - aq)(1 - b_1q^{m_1+1}/b)/(1 - b_1q^2)(1 - b_1q^3)}{(1 - azq^{m_1-1}) + \dots}$$

(3.3)
$$\frac{{}_2\psi_2 \left[\begin{matrix} \frac{a}{b}, & \frac{b_1}{b}q^{m_1+1} \\ \frac{q}{b}, & \frac{b_1}{b}q \end{matrix} ; q; z \right]}{{}_2\psi_2 \left[\begin{matrix} \frac{a}{b}, & \frac{b_1}{b}q^{m_1} \\ \frac{q}{b}, & \frac{b_1}{b} \end{matrix} ; q; zq \right]} = \frac{(b - b_1)}{b^2(1 - b_1)} \left[1 + \frac{z(1 - a)}{(1 - z)(1 - b_1q^{m_1})+} \right.$$

$$\left. + \frac{b_1(q^{m_1} - 1)/(1 - azq)}{1+} \frac{z(1 - aq)}{(1 - z)(1 - b_1q^{m_1})+} \frac{b_1q(q^{m_1-1} - 1)(1 - azq^2)}{1 + \dots} \right]$$

$$(3.4) \quad \frac{{}_2\psi_2 \left[\begin{matrix} \frac{a}{b}, & \frac{b_1}{b} q^{m_1} \\ \frac{q}{b}, & \frac{b_1}{b} \end{matrix} ; q; zq \right]}{{}_2\psi_2 \left[\begin{matrix} \frac{aq}{b}, & \frac{b_1}{b} q^{m_1+1} \\ \frac{q}{b}, & \frac{b_1}{b} q \end{matrix} ; q; z \right]} = \frac{b(1-b_1)(a-b)}{(b-b_1)(a-1)} \left[\frac{(1-z)}{(1-b_1)-} \frac{b_1(1-azq^{m_1+1})}{(1-b_1)+} \frac{z(1-aq)(1-b_1q^{m_1+1})}{(1-z)/(1-b_1)-} - \frac{b_1q(1-azq^{m_1+2})}{(1-b_1)+} \frac{z(1-aq^2)(1-b_1q^{m_1+2})}{(1-z)/(1-b_1)-} \frac{b_1q^2(1-azq^{m_1+3})}{(1-b_1)+ \dots} \right].$$

$$(3.5) \quad \frac{{}_2\psi_2 \left[\begin{matrix} \frac{a}{b}, & \frac{b_1}{b} q^{m_1} \\ \frac{q}{b}, & \frac{b_1}{b} \end{matrix} ; q; z \right]}{{}_2\psi_2 \left[\begin{matrix} \frac{a}{b}, & \frac{b_1}{b} q^{m_1+1} \\ \frac{q}{b}, & \frac{b_1}{b} \end{matrix} ; q; z \right]} = 1 - \frac{zb_1q^{m_1}(1-a)}{(1-q^{-m_1-1})+} \frac{b_1(1-b_1q^{m_1+1})(1-azq^{m_1+1})/b_1q^{m_1+1}}{1-} - \frac{zb_1q^{m_1+1}(1-aq)}{(1-q^{-m_1-1})+} \frac{b_1(1-b_1q^{m_1+2})(1-zab_1q^{m_1+1})/b_1q^{m_1+1}}{1- \dots}.$$

$$(3.6) \quad \frac{{}_2\psi_2 \left[\begin{matrix} \frac{a}{b}, & \frac{b_1}{b} q^{m_1} \\ \frac{q}{b}, & \frac{b_1}{b} \end{matrix} ; q; z \right]}{{}_2\psi_2 \left[\begin{matrix} \frac{a}{b}, & \frac{b_1}{b} q^{m_1+1} \\ \frac{q}{b}, & \frac{b_1}{b} q \end{matrix} ; q; z \right]} = \frac{b(1-b_1)}{(b-b_1)} \left[1 + \frac{zb_1(1-a)(1-b_1q^{m_1})/(1-b_1)(1-b_1q)}{(1-azq^{m_1-1})+} + \frac{z/b_1q \times (a-b_1q)/[b_1(q^{m_1}-q)/(1-b_1q)(1-bq^2)]}{1+ \dots} \right].$$

$$(3.7) \quad \frac{{}_2\psi_2 \left[\begin{matrix} \frac{a}{b}, & \frac{b_1}{b} q^{m_1+1} \\ \frac{q}{b}, & \frac{b_1}{b} \end{matrix} ; q; z \right]}{{}_2\psi_2 \left[\begin{matrix} \frac{a}{b}, & \frac{b_1}{b} q^{m_1} \\ \frac{q}{b}, & \frac{b_1}{b} \end{matrix} ; q; zq \right]} =$$

$$\begin{aligned}
 & \frac{1}{b} \left[1 + \frac{z(1-a)}{(1-z)(1-b_1q^{m_1})+} \frac{b_1(q^{m_1}-1)/(1-azq)}{1+} \right. \\
 & \left. + \frac{z(1-aq)}{(1-z)(1-bq^{m_1})+} \frac{b_1q(q^{m_1-1}-1)(1-azq^2)}{1+} \frac{z(1-aq^2)}{(1-z)(1-b_1q^{m_1})+ \dots} \right]. \\
 (3.8) \quad & \frac{{}_2\psi_2 \left[\begin{matrix} \frac{a}{b}q^i, & \frac{b_1}{b}q^{m_1+i} \\ \frac{q}{b}, & \frac{b_1}{b}q^i \end{matrix} ; q, z \right]}{{}_2\psi_2 \left[\begin{matrix} \frac{a}{b}q^{i+1}, & \frac{b_1}{b}q^{m_1+i+1} \\ \frac{q}{b}, & \frac{b_1}{b}q^{i+1} \end{matrix} ; q, z/q \right]} = \frac{b(aq^i-b)(1-b_1q^i)(1-z/q)}{(aq^i-1)(b-b_1q^i)(1-b_1)} \\
 & - \frac{(1-azq^{m_1+i})b_1q^i}{(1-b_1)+} \frac{z(1-aq^{i+1})(1-b_1q^{m_1+i+1})/b_1q^{m_1}}{(1-z/q)/(1-b_1)-} \\
 & - \frac{(1-azq^{m_1+i+1})b_1q^{i+1}}{(1-b_1)+} \frac{z(1-aq^{i+2})(1-b_1q^{m_1+i+2})/q}{(1-z/q)/(1-b_1)- \dots} \\
 (3.9) \quad & \frac{{}_2\psi_2 \left[\begin{matrix} aq^{i+1}, & \frac{b_1}{b}q^{m_1+i+1} \\ \frac{q}{b}, & \frac{b_1}{b}q^{i+1} \end{matrix} ; q, z/q \right]}{{}_2\psi_2 \left[\begin{matrix} \frac{a}{b}q^{i+1}, & \frac{b_1}{b}q^{m_1+i+1} \\ \frac{q}{b}, & \frac{b_1}{b}q^{i+1} \end{matrix} ; q, z \right]} = \frac{1}{b(1-b_1)} \left[(1-b_1)+ \right. \\
 & \frac{(1-aq^{i+1})(1-b_1q^{m_1+i+1})/q}{(1-z/q)/(1-b_1)-} \frac{(1-azq^{m_1+i+1})b_1q^{i+1}}{(1-b_1)+} \frac{z(1-aq^{i+2})(1-b_1q^{m_1+i+2})/q}{(1-z/q)/(1-b_1)-} \\
 & \left. - \frac{(1-azq^{m_1+i+2})b_1q^{i+2}}{(1-b_1)+} \frac{z(1-aq^{i+3})(1-b_1q^{m_1+i+3})/q}{(1-z/q)/(1-b_1)- \dots} \right]. \\
 (3.10) \quad & \frac{{}_2\psi_2 \left[\begin{matrix} \frac{a}{b}q^i, & \frac{b_1}{b}q^{m_1+i} \\ \frac{q}{b}, & \frac{b_1}{b}q^i \end{matrix} ; q, z \right]}{{}_2\psi_2 \left[\begin{matrix} \frac{a}{b}q^{i+1}, & \frac{b_1}{b}q^{m_1+i} \\ \frac{q}{b}, & \frac{b_1}{b}q^{i+1} \end{matrix} ; q, z/q \right]} = \\
 & \frac{b(1-b_1)(aq^i-b)}{(b-b_1q^i)(aq^i-1)} \left[\frac{(1-1/q^{m_1})}{(1-b_1)+} \frac{b_1(1-b_1q^{m_1+i})(1-azq^{m_1+i})/b_1q^{m_1}}{(1-b_1)-} \frac{(1-aq^{i+1})zb_1q^{m_1+i}}{(1-1/q^{m_1})/(1-b_1)+} \right. \\
 & \left. + \frac{b_1(1-b_1q^{m_1+i+1})(1-azq^{m_1+i+1})/b_1q^{m_1}}{(1-b_1)-} \frac{(1-aq^{i+2})zb_1q^{m_1+i+1}}{(1-1/q^{m_1})/(1-b_1)+ \dots} \right].
 \end{aligned}$$

for $i = 0, 1, 2, 3, \dots$

Proof of (3.1)-(3.10).

As an illustration, we shall prove (3.1).

Taking $c=d=0$ and $r=1$ in (1.1), we have

$$(3.11) \quad {}_2\psi_2 \left[\begin{matrix} \frac{a}{b}, & \frac{b_1}{b} q^{m_1} \\ \frac{q}{b}, & \frac{b_1}{b} \end{matrix} ; q; z \right] = \frac{(b_1; q)_{m_1}}{(b_1/b; q)_{m_1}} \\ \times \frac{\left(q, \frac{bq}{az}, \frac{az}{b}, \frac{q}{a}; q \right)_{\infty}}{\left(\frac{q}{b}, \frac{q}{az}, az, \frac{bq}{a}; q \right)_{\infty}} {}_2\phi_1 \left[\begin{matrix} a, & b_1 q^{m_1} \\ b_1, & \end{matrix} ; q; z \right]$$

Now, replacing z by zq in (3.11) and finally taking the ratio of these two, we have

$$\frac{{}_2\psi_2 \left[\begin{matrix} \frac{a}{b}, & \frac{b_1}{b} q^{m_1} \\ \frac{q}{b}, & \frac{b_1}{b} \end{matrix} ; q; zq \right]}{{}_2\psi_2 \left[\begin{matrix} \frac{a}{b}, & \frac{b_1}{b} q^{m_1} \\ \frac{q}{b}, & \frac{b_1}{b} \end{matrix} ; q; z \right]} = b \frac{{}_2\phi_1 \left[\begin{matrix} a, & b_1 q^{m_1} \\ b_1, & \end{matrix} ; q; zq \right]}{{}_2\phi_1 \left[\begin{matrix} a, & b_1 q^{m_1} \\ b_1, & \end{matrix} ; q; z \right]}$$

Now, using the result due to Denis[10], (3.1) can be obtained easily.

Proceeding in the same way and using the results due to R. P. Agarwal [[1], (3.23), (3.21), (3.18) pages[69-71]], S. N. Singh [19, 20] and N. Bhagirathi [5], one can easily establish the results (3.2) to (3.10), respectively. \square

Note: It is important to mention that a number of results for the basic hypergeometric series ${}_2\phi_1$ can be deduced from the results established in §3.

References

- [1] R. P. Agarwal, *Resonance of Ramanujan's mathematics Vol. III*, New Age International Pvt. Ltd. Publishers, New Delhi, 1998.
- [2] G. E. Andrews, *Ramanujan's 'Lost' Note Book III, The Rogers-Ramanujan's continued fraction*, Adv. Math., 2, **41**(1981), 186-208.
- [3] C. Adiga, D. D. Somashekara, *On some Rogers-Ramanujan type continued fraction identities*, Mathematica Balkamica, New Series 1-2, **12**(1998), 37-45.
- [4] C. Adiga, D. D. Somashekara, *Rogers-Ramanujan Identities, continued fractions and their generalizations*, Selected Topics in Special Functions, Allied publisher limited, 2001, 208-226.

- [5] N. A. Bhagirathi, *On certain investigations in q -series and continued fractions*, Math. Student, 1-4, **56**(1988), 158-170.
- [6] N. A. Bhagirathi, *On basic bilateral hypergeometric series and continued fractions*, Math. Student, 1-4, **56**(1988), 135-141.
- [7] B. C. Berndt, *Ramanujan's Notebook part III*, Springer-Verlag, 1991.
- [8] S. Bhargava, C. Adiga, D. D. Somashekara, *On certain continued fractions related to ${}_3\phi_2$ basic hypergeometric functions*, J. Math. Phys. Sci., **21**(1987), 613-629.
- [9] S. Bhargava, C. Adiga, *On some continued fraction of Srinivas Ramanujan*, Proc. Amer. Math. Soc., **92**(1984), 13-18.
- [10] R. Y. Denis, *On basic hypergeometric function and continued fractions*, Math. Student, 1-4, **52**(1984), 129-136.
- [11] R. Y. Denis, *On certain q -series and continued fractions*, Math. Student, **44**(1983), 70-76.
- [12] R. Y. Denis, *On certain continued fraction of Ramanujan*, Jour. Math. Phys. Sci., **3**, **24**(1990), 193-205.
- [13] R. Y. Denis, *On generalization of Euler's continued fraction*, Indian J. Pure Appl. Math., **21**(1)(1990), 78-81.
- [14] M. D. Hirschhorn, *A continued fraction of Ramanujan*, J. Australian Math. Soc., **29**(1980), 80-86.
- [15] Kline, Morris, *Mathematical thought from ancient to modern time*, Oxford Univ. Press, 1972, 28-32.
- [16] Maheshwar Pathak and Pankaj Srivastava, *A note on continued fractions and ${}_3\psi_3$ series*, Italian J. Pure Appl. Math., **27**(2010), 191-200.
- [17] S. Ramanujan, *Note book of Ramanujan Vol. I & II*, T. I. F. R. Bombay, 1957.
- [18] S. Ramanujan, *The lost note book and other unpublished papers*, Narosa, New Delhi, 1988.
- [19] S. N. Singh, *On q -hypergeometric function and continued fractions*, Math. Student, 1-4, **56**(1988), 81-84.
- [20] S. N. Singh, *Basic hypergeometric series and continued fractions*, Math. Student, 1-4, **56**(1988), 91-96.
- [21] S. N. Singh, *Certain transformation formulae for basic and bibasic hypergeometric series*, Proc. Nat. Acad. Sci. India, **65**(A), **III**(1995), 319-329.
- [22] Pankaj Srivastava, *Certain continued fractions for quotients of two ${}_3\psi_3$ series*, Proc. Nat. Acad. Sci. India, **78**(A), **IV**(2008), 327-330.
- [23] Bhaskar Srivastava, *A continued fraction and some identities associated with Ramanujan*, Riv. Math. Univ. Parma., **17**(1991), 199-206.
- [24] L. C. Zhang, *Ramanujan's continued fraction for product of gamma functions*, J. Math. Anal. Appl., **174**(1993), 22-25.