


Accepted Article

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version record](#). Please cite this article as [doi:10.1002/ana.24594](https://doi.org/10.1002/ana.24594).



170x152mm (300 x 300 DPI)

Acce

Response to Treatment in Infantile Spasms

Title: **Response To Treatment In A Prospective National Infantile Spasms Cohort**

Running Head: **Response to treatment in Infantile Spasms**

Authors: Kelly G Knupp MD¹, Jason Coryell MD², Katherine C Nickels MD³, Nicole Ryan MD⁴, Erin

Leister⁵, [Tobias Loddenkemper MD⁶](#), [Zachary Grinspan MD, MS⁷](#), [Adam L Hartman MD⁸](#), [Eric H Kossoff MD⁸](#), [William D Gaillard MD⁹](#), [John R Mytinger MD¹⁰](#), [Sucheta Joshi MD¹¹](#), [Renée A Shellhaas MD, MS¹¹](#), [Joseph Sullivan MD¹²](#), [Dennis Dlugos MD⁴](#), [Lorie Hamikawa MD¹³](#), [Anne T Berg PhD¹⁴](#), [John Millichap MD¹⁴](#), [Douglas R Nordli Jr MD¹⁴](#), [Elaine Wirrell MD³](#) and the Pediatric Epilepsy Research Consortium

¹ Department of Pediatrics and Neurology, School of Medicine, University of Colorado Anschutz Medical Campus

² Departments of Pediatrics and Neurology, School of Medicine, Oregon Health & Sciences University

³ Departments of Neurology and Pediatrics, Mayo Clinic Rochester

⁴ Division of Neurology, The Children's Hospital of Philadelphia and Perelman School of Medicine at the University of Pennsylvania, Philadelphia

⁵ Colorado School of Public Health, Department of Biostatistics and Informatics, University of Colorado Anschutz Medical Campus

⁶ [Division of Epilepsy and Clinical Neurophysiology, Department of Neurology, Boston Children's Hospital, Boston, MA, USA](#)

⁷ [Weill Cornell Medical Center, New York, NY](#)

⁸ [Departments of Neurology and Pediatrics, Johns Hopkins Hospital, Baltimore Maryland](#)

⁹ [Center For Neuroscience, Children's National Health System](#)

¹⁰ [The Ohio State University, Nationwide Children's Hospital, Department of Pediatrics, Division of Pediatric Neurology, Columbus, Ohio](#)

¹¹ [Department of Pediatrics & Communicable Diseases \(Division of Pediatric Neurology\), University of Michigan](#)

¹² ~~Department~~[Departments of Pediatrics and Neurology, University of San Francisco](#)

¹³ [University of Washington](#)

- Formatted: Font: Font color: Black, Superscript
- Formatted: Line spacing: Double
- Formatted: Superscript
- Formatted: Superscript
- Formatted: Superscript
- Formatted: Font: Border: : (No border)
- Formatted: Superscript
- Formatted: Font: Border: : (No border)
- Formatted: Superscript
- Formatted: Line spacing: Double
- Formatted: Superscript
- Formatted: Superscript
- Formatted: Font: Border: : (No border)
- Formatted: Superscript
- Formatted: Line spacing: Double
- Formatted: Superscript

Response to Treatment in Infantile Spasms

¹⁴ [Ann & Robert H. Lurie Children's Hospital of Chicago; and Departments of Pediatrics and Neurology,
Northwestern University Feinberg School of Medicine](#)

Formatted: Superscript

Corresponding author:

Kelly G Knupp MD
13123 East 16th Ave
Aurora, CO 80045
Ph: 720 777 6871
Fax: 720 777 7285
Email: kelly.knupp@childrenscolorado.org

Characters:

Title: 65
Running head: 41

Words:

Abstract: 243
Introduction: 502
Discussion: 1234
Body of manuscript: 3480

Number of figures: 1

Number of tables: 5

Accel

Response to Treatment in Infantile Spasms

Abstract

Objective: Infantile spasms are seizures associated with a severe epileptic encephalopathy presenting in the first 2 years of life, and optimal treatment continues to be debated. This study evaluates early and sustained response to initial treatments and addresses both clinical remission and electrographic resolution of hypsarhythmia.

Secondarily, it assesses whether response to treatment differs by etiology or developmental status.

Methods: The National Infantile Spasms Consortium established a multi-center, prospective database enrolling infants with new diagnosis of infantile spasms. Children were considered responders if there was clinical remission and resolution of hypsarhythmia that was sustained at three months after first treatment initiation.

Standard treatments of ACTH, oral corticosteroids, and vigabatrin were considered individually, and all other non-standard therapies were analyzed collectively.

Developmental status and etiology were assessed. We compared response rates by treatment group using Chi-square tests and multivariable logistic regression models.

Results: Two hundred and thirty infants were enrolled from 22 centers. Overall, 46% of children receiving standard therapy responded compared to only 9% who responded to non-standard therapy ($p < 0.001$). 55% of infants receiving ACTH as initial treatment responded, compared to 39% for oral corticosteroids, 36% for vigabatrin, and 9% for other ($p < 0.001$). Neither etiology nor development significantly modified the response pattern by treatment group.

Response to Treatment in Infantile Spasms

Interpretation: Response rate varies by treatment choice. Standard therapies should be considered as initial treatment for infantile spasms, including those with impaired development or known structural or genetic/metabolic etiology. ACTH appeared to be more effective than other standard therapies.

Accepted Article

Response to Treatment in Infantile Spasms

Response To Treatment In A Prospective National Infantile Spasms Cohort

Introduction

West syndrome is an infantile epileptic encephalopathy, typically occurring within the first two years of life with an incidence of 2-5 per 10,000 live births.¹⁻⁴ Infantile spasms, a subtype of epileptic spasms are the pathognomonic seizure type and are frequently associated with hypsarhythmia on electroencephalogram (EEG) and developmental plateau or regression. The literature on preferred treatment is inconsistent and at times conflicting, suggesting multiple treatment options and dosing regimens. The most accepted treatments are adrenocorticotrophic hormone (ACTH), oral corticosteroids (OCS), and vigabatrin, although other therapies have been used.⁵ Despite consensus statements, recent surveys of child neurologists showed little agreement regarding best initial therapy, preferred dose, route of administration, and adjunctive medications.^{6,7}

The effective use of ACTH and OCS was first reported in 1958.^{8,9} The Food and Drug Administration dosing guidelines for ACTH recommend initial high dose therapy (150 international units [IU]/m²/Day divided twice daily)¹⁰. Studies have reported response rates of 86-93% with high dose ACTH,^{11,12} while a randomized study indicated that high dose ACTH was not superior to low dose (20-30 IU/ day).¹³ Some trials comparing ACTH and OCS utilized low doses (20-30 IU) of ACTH, and failed to observe significant

Response to Treatment in Infantile Spasms

differences in response rates.^{14, 15} The United Kingdom Infantile Spasms Study (UKISS) observed similar response rates between OCS and synthetic ACTH (tetracosactide) (70% and 76%), with both superior to vigabatrin (54%).¹⁶ UKISS was not powered to compare hormone subgroups and electroencephalogram (EEG) data was not used in the assessment of the primary outcome measure. Higher dosing of prednisolone in UKISS (40-60 mg/day in three to four divided doses) than in previous clinical trials, may account for the higher response rate. Finally, case series utilizing “high dose” OCS (initial 40-45 mg/day, increased to max 60 mg/day)¹⁷ and “very high dose” OCS (8 mg/kg/day, max 60 mg/day)¹⁸ reported two week response rates of 67% and 63%. Trials of vigabatrin (100-148 mg/kg/day in two divided doses) reported a 36% remission rate.¹⁹ The response rate to vigabatrin was higher in those with tuberous sclerosis compared to other conditions (52% and 16%)¹⁹. Additional trials comparing low dose ACTH to vigabatrin as first line therapy found better response to ACTH (74%) than vigabatrin (48%).²⁰

Aside from those including children with tuberous sclerosis, few prospective studies have assessed etiology and development as predictors of outcome. The largest demonstrated higher IS resolution in cryptogenic cases with ACTH compared to OCS.²¹ A retrospective study showed no difference in response between ACTH and vigabatrin (88% and 80%) among children with unknown etiology (normal early development and MRI brain).²²

Response to Treatment in Infantile Spasms

Multi-center collaboration has been recommended to increase understanding of effective treatments.⁶ This study presents data from a large multi-center consortium evaluating early and sustained response to initial treatments and addresses both clinical remission and electrographic resolution of hypsarhythmia. The study aims were to observe response rates by primary treatment type, and to ascertain if children with varying etiologies or developmental status had differential responses to treatment in a prospective manner.

Methods

In 2012, The Pediatric Epilepsy Research Consortium (PERC) developed the National Infantile Spasms Consortium (NISC) database, which is a multi-center database enrolling children prospectively. Children with new onset infantile spasms between two months and two years of age were eligible for enrollment. Clinical information was collected at time of diagnosis and three months after diagnosis. Medication dosing recommendations for ACTH, oral corticosteroids and vigabatrin is provided to all NISC centers (see table 1 for dosing recommendations) to improve homogeneity for analysis, although compliance with these recommendations was not necessary for inclusion. Treatment decisions for individual children were made by the treating clinician.

The parents or guardians provided consent for all children according to center-specific IRB requirements. The study was approved by all participating site IRBs. Data collected from June 2012 to July 2014 were used for this study. Children with an early

Response to Treatment in Infantile Spasms

infantile epileptic encephalopathy (Ohtahara syndrome/ Early Myoclonic Encephalopathy) and/or missing treatment or response data due to lack of follow up or incomplete data entry were excluded from the analysis.

Data collected for each child included age at onset of IS, gestational age at birth, sex, presence of seizures prior to spasms, etiology, height, weight, MRI, genetic and metabolic testing, developmental assessment, presence of hypsarhythmia at onset, IS medication, and dosage. Hypsarhythmia was assessed at individual institutions and defined as multifocal spikes, disorganization, and >200 microvolts [peak-to-peak] in any epoch on a bipolar longitudinal montage.²³ At three months, etiology, new MRI findings, new genetic and metabolic testing, developmental assessment, response to medication(s), and EEG findings were collected. Response to medications was recorded as response at 2 weeks from initiation of medication and response at 3 months from enrollment into study.

Standard therapy was defined as ACTH, oral corticosteroids or vigabatrin. All other medications were considered nonstandard therapy. Children initiated on simultaneous standard and non-standard therapy (e.g. ACTH and levetiracetam), had response attributed to the standard medication. Body surface area was calculated for each child using the Haycock formula.²⁴ Dosing for ACTH was considered high dose if greater than 140 IU/m²/day was used at initiation of medication. All others were categorized as low/intermediate dose. For analysis, ACTH was combined with all doses due to small number of children receiving low/intermediate dose ACTH.

Response to Treatment in Infantile Spasms

Response to initial medication was classified into one of four response categories based on clinical infantile spasm response rate and resolution of hypsarhythmia on EEG (when hypsarhythmia was present at onset): early responders, late responders, relapse and non-responders. Early responders had resolution of clinical spasms documented in the medical record by two weeks and ongoing remission three months after the start of treatment with an EEG to confirm remission of hypsarhythmia. Various EEGs were used including routine, video and prolonged EEG. Late responders had clinical remission starting after two weeks of treatment and had an EEG to confirm remission of hypsarhythmia. Relapse included children who initially met response criteria and then had return of either clinical spasms or hypsarhythmia. All others were considered non-responders. In our primary analyses, we defined two response categories as follows: responders included children who responded both early and late, and the non-responder group included non-responders and those who relapsed. In order to compare our results to other studies, in separate analyses, we defined 2 week responders as those who responded early and those who responded early but then relapsed.

Development was recorded as the clinician's perception of overall development, motor and cognitive status, with each defined as normal, mild or equivocal delay, or definite abnormality. These three domains were then used to create an overall assessment of development categorized as normal, mild, moderate and severe. Children with no domain marked as abnormal were classified as having normal development. If one domain was marked as mild, the child was included in the mild developmental delay group. The moderate developmental delay group consisted of children with two or more domains marked as mild or one domain marked as a definite

Response to Treatment in Infantile Spasms

abnormality. Severe developmental delay included children with two or more domains were marked as definite abnormality.

Etiology was classified into five primary etiologic classifications: genetic/metabolic, malformation of cortical development, prior acquired injury, other structural, and unknown. Tuberous sclerosis was classified as other structural according to International League Against Epilepsy (ILAE) guidelines²⁵ and was not removed from analysis due to small numbers. For data analysis, those with unknown etiology were further categorized into normal and abnormal development. Unknown etiology with normal development was analyzed as a separate category, while genetic/metabolic was combined with unknown etiology and abnormal development. The latter group likely represents presumed genetic causes, but without a determined etiology in the 3 month follow-up period (either due to late diagnosis, decreased utilization of testing, or genetic influences that are non-Mendelian). Additionally, malformations of cortical development, prior injury, and other structural were categorized together as a structural cause of epilepsy.

Statistical analysis

We compared demographic and clinical characteristics by treatment group (ACTH, oral corticosteroid, VGB, or other) using Chi-square tests for categorical covariates and Kruskal-Wallis tests for continuous covariates. To understand the association of demographic and clinical covariates with treatment response, we used Chi-square tests to compare the proportions of responders in each group. Next, we fit multivariable

Response to Treatment in Infantile Spasms

logistic regression models to estimate crude and adjusted relative risk of responding to treatment.²⁶ We also estimated the adjusted predicted probability of treatment response for each treatment and etiologic or developmental category of interest. Prior to selecting final models we explored the possibility of interaction between treatment and etiology or development. All analyses were performed using SAS version 9.4 (SAS Institute, Cary, NC).

Results

Entrance criteria were met by two hundred eighty-two subjects. 52 were excluded due to: 33 lack of adequate data, eight lost to follow up, six deceased and five with early infantile epileptic encephalopathy (Ohtahara syndrome/ Early Myoclonic Encephalopathy). Two hundred thirty subjects had complete follow-up information including clinical and EEG response to individual treatments for a three month period. Twenty-two centers contributed data with 1-43 children enrolled per center. The median (Q1, Q3) age of infantile spasm onset was 6.5 months (4.5, 8.5), and the time from spasm infantile onset to initiation of treatment was 15 days (6, 37) (Table 2).

The first treatment choice for infantile spasms was ACTH in 42%, oral corticosteroids (OC) in 23%, vigabatrin in 20%, while 14% were prescribed non-standard therapies (levetiracetam, topiramate, clobazam, valproic acid, zonisamide, oxcarbazepine, phenobarbital, clonazepam and ketogenic diet). Clinicians followed NISC dosing recommendations in 80/96 (83%) of ACTH treated children, 34/50 (68%) of OC treated children and 32/41 (78%) of those treated with vigabatrin. We did not observe any

Response to Treatment in Infantile Spasms

significant differences in the distribution of treatment choice based on sex, race, ethnicity, gestational age, age at infantile spasm onset, time to diagnosis, or time to treatment. However, we did observe differences in choice of treatment by other clinical characteristics. Children with a history of prior seizures or anti-seizure medication use were less likely to be treated with ACTH and more likely to be treated with a non-standard therapy. (Table 2) The etiology of infantile spasms was also associated treatment choice, with vigabatrin prescribed more frequently to those with a structural cause. There was prescribing variation based on development at the time of presentation; children prescribed ACTH had the highest percentage of mild or no developmental issues while children prescribed non-standard therapy were most likely to have severe development issues ($p < 0.001$). (Table 2)

Ninety-one out of 198 (46%) children receiving standard therapy were responders with early or late clinical remission and resolution of hypsarhythmia by EEG, while only three out of 32 (9%) responded to non-standard therapy ($p < 0.001$). Fifty-three out of 97 (55%) infants receiving ACTH as initial treatment responded, compared to 21/54 (39%) for oral corticosteroids, 17/47 (36%) for vigabatrin, and 3/32 (9%) for other (overall $p < 0.001$). When we compared response rate between pairs of treatments, we observed significant differences (all $P < 0.01$ between each standard treatment and other, non-standard treatment. The response rate in the ACTH group was significantly higher than the vigabatrin group ($p = 0.038$) and marginally higher than the oral corticosteroid group ($p = 0.06$). We did not observe a significant difference in response rate between oral corticosteroids and vigabatrin. In 11 children with tuberous sclerosis, eight were

Response to Treatment in Infantile Spasms

treated with VGB and five responded (62%); those treated with ACTH (1) and non-standard therapy (2) did not respond. Of the 80 children who received high dose ACTH, 46 (58%) responded, compared to 6 (38%) of the 16 who received low/intermediate dose ACTH ($p=0.14$).

Detailed two week and three month response by treatment is shown in Figure 1. We observed the greatest relapse rate in the oral corticosteroids group (24%) compared to the other three treatment groups (18% overall), but this difference was not significant ($p=0.21$). At two weeks 66/97 (68%) on ACTH and 30/54 (56%) on oral corticosteroids had responded ($p=0.13$). The response rate to ACTH at two weeks was significantly higher than the rate in those on vigabatrin (23/47, 49%, $p=0.027$) or non-standard treatment (7/32, 22%, $p<0.001$).

Response rates were also significantly higher for those without a history of prior seizures or anti-seizure medication use. (Table 3) In unadjusted analysis, children with normal development or mild developmental delay had a higher response rate than children with moderate or severe delay ($p=0.025$). Children with unknown etiology and normal development had a higher response rate (20/34, 59%) than children with genetic (40/106, 38%) or structural (34/90, 38%) etiologies though this was not statistically significant. ($p=0.07$).

Our final multivariate logistic regression analyses consisted of two models, one including treatment and developmental status as covariates, the other including

Response to Treatment in Infantile Spasms

treatment and etiology. Model ~~results are~~ results are shown in Tables 4 and 5. Crude and adjusted relative risks of response between treatment groups and development and etiology categories are shown in Table 4. These analyses showed that after adjusting for development and etiology, choice of treatment was still as significant predictor of response. Children on standard treatments had more than three times greater probability of responding than those on non-standard treatments, with children on ACTH over five times as likely to respond. In models adjusting for treatment, the effects of etiology and development on response were weaker than the effects seen in the unadjusted models. The predicted probability of response to specific treatments is presented for each etiology and developmental status group (Table 5). We did not observe modification of the overall treatment effect by either etiology or development; the predicted treatment response patterns were similar across the etiologic and developmental sub-groups.

Discussion

The National Infantile Spasms Consortium (NISC) database provides a multicenter, prospective cohort of children with newly diagnosed infantile spasms. The large prospective nature of the study allows for assessment of selection bias of treatment for a number of factors including baseline development and etiology. The size of this multicenter study also enables greater power to distinguish outcomes. Assessing outcome at three months allows for more a clinically appropriate analysis of response to medication accounting for relapse rates.

Response to Treatment in Infantile Spasms

The determination of efficacy by both immediate and sustained response at three months, in our study, allows for a more complete clinical picture regarding response to treatment. The majority of studies have assessed outcome at two weeks.^{11, 12, 16-20, 27} Similar to other studies, we found that ACTH (all doses combined) was associated with higher early response rate than vigabatrin or oral corticosteroids, although this did not reach statistical significance for oral corticosteroids.^{11, 16, 20, 22} At 3 months, after taking into account relapse rates, the sustained response rate in those treated with ACTH was still significantly higher compared to those treated with vigabatrin and marginally higher than those treated with oral corticosteroids. The difference in outcomes particularly between ACTH and oral corticosteroids may be due to the increased relapse rate that was present in the oral corticosteroid group. Previous studies with shorter follow up often did not account for relapse rates and the only long-term follow-up studies were smaller (24 and 97 subjects each)^{14, 15} and used variable dosing of ACTH and oral corticosteroids. This may have led to an overestimation of response rate to oral corticosteroids. In the current study, relapses occurred with all treatments, but were more frequent with oral corticosteroids. Future study designs should account for relapse rates as it may vary by treatment. Our data also demonstrate higher response rate to ACTH in children with unknown etiology which differs from a previously published study demonstrating vigabatrin as equally effective to ACTH;²⁰ this may be due to the larger size of our study and its prospective design.

Children were more likely to respond to standard therapies than non-standard therapies. Our findings suggest the importance of using standard therapeutic agents for treatment

Response to Treatment in Infantile Spasms

of infantile spasms. These differences were seen for early response, 3 month response, and relapse rates. The large cohort relative to other studies and observation nature of this study allows for evaluation of a sizeable group of children on non-standard therapy (therapies other than ACTH, oral corticosteroids, and vigabatrin). This was a heterogeneous group in regards to treatment. Currently available information regarding non-standard treatments for infantile spasms is limited to small retrospective cohorts and limited case studies. A recent study demonstrated improved rates of spasm remission after a standard treatment protocol was implemented (78.8% post-standardization compared to 30.6% pre-standardization). Treatment was standardized to ACTH, oral corticosteroids, or vigabatrin and other anti-seizure medications were no longer used, providing indirect evidence of poor outcomes with non-standard medications.²⁸ Our findings provide additional evidence of poor response to non-standard medications. The most recent guidelines on treatments of infantile spasms from AAN/CNS state there is insufficient evidence to determine whether other (non-standard) therapies, including valproic acid, pyridoxine, and ketogenic diet are effective treatments for spasms.

The etiology of spasms and the presence of moderate-severe delays at onset were not as strongly associated with treatment response as the choice of treatment itself. There were differences in prescriber choice of initial therapy based on patient development and etiology. Children with mild or no developmental issues were more often prescribed ACTH while those with severe impairments were more likely to be prescribed a non-standard treatment. This suggests some of the poor response to non-standard

Response to Treatment in Infantile Spasms

treatments may have been due to selection bias and underlying etiology. However, adjusted models demonstrate that response was not significantly affected by etiology or development once adjusted for treatment, though we had limited power to detect response differences between etiology and development groups. Based on these findings, there is not good evidence to alter treatment choice due to existence of a structural abnormality or pre-existing developmental delay. However, there is still likely a role for treatment-specific therapies, such as vigabatrin for those with tuberous sclerosis.

This study showed a trend for higher response rates with high dose ACTH compared to low/intermediate dose ACTH. However, this study was not sufficiently powered to determine differences in outcome between high dose and low/intermediate dose ACTH response. Due to the NISC dosing guidelines, there were fewer subjects on low/intermediate-dose ACTH. The most effective dose of ACTH is variably reported in the literature and several reports suggest low/intermediate dose ACTH is as effective as high dose ACTH.^{5, 13, 27, 29} The most recent evidence-based guidelines on the medical treatment of infantile spasms from the American Academy of Neurology and the Child Neurology Society suggests consideration for “low-dose” ACTH as an alternative to high dose ACTH,⁵ although the majority of surveyed child neurologists use higher doses of ACTH.⁶ While the efficacy of other therapies is often compared to the efficacy of ACTH, the dose, type and duration of treatment of ACTH is inconsistent among studies, making interpretation of the available literature challenging. This merits further study.

Response to Treatment in Infantile Spasms

There were some limitations to this study. First, the therapies were not randomized and providers were allowed to choose the medications, including those that were not considered “standard.” There is evidence of medication selection bias based on underlying etiology and developmental status, as well as variability in doses. However, in multivariable analysis we did not observe significant differences in overall treatment response patterns based on underlying etiology or development. Although this suggests that the lack of randomization did not contribute to the results of this study, we cannot exclude unknown confounders given the observational nature of study design. Developmental assessment was based on clinical exam and not on formal assessment tools, although recent studies suggest that tests such as the Bayley Scales of Infant Development (BSID) do not always correspond to long term cognitive outcome and are associated with reporting errors.^{30, 31} The use of broad developmental classifications in this study to determine the effect of development at the onset of spasms on treatment bias and response rates, though crude, was likely sufficient to identify those with moderate to severe developmental delay. As improvement in developmental outcomes is the ultimate goal in the treatment of infantile spasms, further long term developmental assessments need to be performed. Finally, this study was not sufficiently powered to identify response to treatment based on each specific etiology and merging of etiological categories was necessary.

Using a multicenter prospective design, this study was able to incorporate larger numbers of children than previous studies, providing increasing insight about the most

Response to Treatment in Infantile Spasms

effective initial treatment for infantile spasms. Extended follow-up of a large cohort provides the most extensive data regarding relapse rates, and how relapse varies by treatment choice providing a more complete clinical picture. The larger enrollment enabled assessment of response differences among developmental and etiologic subgroups. ACTH is likely more effective for children with infantile spasms regardless of development or etiology (perhaps with the exception of tuberous sclerosis). Our data do not support the use of non-standard medications as initial therapy for infantile spasms. Additional prospective treatment trials are needed to determine optimal dosing of ACTH, further explore the implications of etiology, provide expanded developmental assessments and long term outcome of development and epilepsy.

Acknowledgments

Funding for this study was received from the American Epilepsy Society and Pediatric Epilepsy Research Foundation. [The authors would like to appreciate and thank the numerous research assistants at all the institutions who gathered and entered data for making this study possible.](#)~~The authors would like to appreciate and thank the numerous research assistants at all the institutions that entered and corrected data, making this study possible.~~ The authors appreciate the contribution of SAS code by Dr. Lawrence C. Kleinman and Professor Edward C. Norton to assist with the regression risk analysis.

Author Contributions

The named authors participated in the conception and design of the study, and in the data analysis and editing. ~~Please see table indicating additional author~~

Formatted: Normal

Formatted: Border: : (No border)

Response to Treatment in Infantile Spasms

contributions. The members of the Pediatric Epilepsy Research Consortium contributed data and participated in editing the manuscript. They and their academic affiliations are included in Table X6.

Potential Conflicts of Interest

Author SH received grant support from Lundbeck which owns rights to vigabatrin, and served on the scientific advisory board of Questcor/Mallinckrodt which owns rights to ACTH. No additional authors had conflicts of interest to disclose.

References

1. Riikonen R, Donner M. Incidence and aetiology of infantile spasms from 1960 to 1976: a population study in Finland. *Developmental medicine and child neurology*. 1979 Jun;21(3):333-43.
2. Cowan LD, Hudson LS. The epidemiology and natural history of infantile spasms. *Journal of child neurology*. 1991 Oct;6(4):355-64.
3. Luthvigsson P, Olafsson E, Sigurthardottir S, Hauser WA. Epidemiologic features of infantile spasms in Iceland. *Epilepsia*. 1994 Jul-Aug;35(4):802-5.

Response to Treatment in Infantile Spasms

4. Sidenvall R, Eeg-Olofsson O. Epidemiology of infantile spasms in Sweden. *Epilepsia*. 1995 Jun;36(6):572-4.
5. Go CY, Mackay MT, Weiss SK, et al. Evidence-based guideline update: Medical treatment of infantile spasms: Report of the Guideline Development Subcommittee of the American Academy of Neurology and the Practice Committee of the Child Neurology Society. *Neurology*. 2012 Jun 12;78(24):1974-80.
6. Mytinger JR, Joshi S. The current evaluation and treatment of infantile spasms among members of the child neurology society. *Journal of child neurology*. 2012 Oct;27(10):1289-94.
7. Wheless JW, Clarke DF, Carpenter D. Treatment of pediatric epilepsy: expert opinion, 2005. *Journal of child neurology*. 2005 Dec;20 Suppl 1:S1-56; quiz S9-60.
8. Low NL. Infantile spasms with mental retardation. II. Treatment with cortisone and adrenocorticotropin. *Pediatrics*. 1958 Dec;22(6):1165-9.
9. Sorel L, Dusaucy-Bauloye A. [Findings in 21 cases of Gibbs' hypsarrhythmia; spectacular effectiveness of ACTH]. *Acta Neurol Psychiatr Belg*. 1958 Feb;58(2):130-41.
10. H.P. Acthar Gel [package insert]. Questcor pharmaceuticals, Inc, Union City CA, 2010. Available from: <http://www.acthar.com/pdf/Acthar-PI.pdf>.
11. Baram TZ, Mitchell WG, Tournay A, Snead OC, Hanson RA, Horton EJ. High-dose corticotropin (ACTH) versus prednisone for infantile spasms: a prospective, randomized, blinded study. *Pediatrics*. 1996 Mar;97(3):375-9.

Response to Treatment in Infantile Spasms

12. Snead OC, 3rd, Benton JW, Jr., Hosey LC, et al. Treatment of infantile spasms with high-dose ACTH: efficacy and plasma levels of ACTH and cortisol. *Neurology*. 1989 Aug;39(8):1027-31.
13. Hrachovy RA, Frost JD, Jr., Glaze DG. High-dose, long-duration versus low-dose, short-duration corticotropin therapy for infantile spasms. *The Journal of pediatrics*. 1994 May;124(5 Pt 1):803-6.
14. Hrachovy RA, Frost JD, Jr., Kellaway P, Zion TE. Double-blind study of ACTH vs prednisone therapy in infantile spasms. *The Journal of pediatrics*. 1983 Oct;103(4):641-5.
15. Wanigasinghe J, Arambepola C, Sri Ranganathan S, Sumanasena S, Muhandiram EC. The efficacy of moderate-to-high dose oral prednisolone versus low-to-moderate dose intramuscular corticotropin for improvement of hypsarrhythmia in West syndrome: a randomized, single-blind, parallel clinical trial. *Pediatric neurology*. 2014 Jul;51(1):24-30.
16. Lux AL, Edwards SW, Hancock E, et al. The United Kingdom Infantile Spasms Study comparing vigabatrin with prednisolone or tetracosactide at 14 days: a multicentre, randomised controlled trial. *Lancet*. 2004 Nov 13-19;364(9447):1773-8.
17. Kossoff EH, Hartman AL, Rubenstein JE, Vining EP. High-dose oral prednisolone for infantile spasms: an effective and less expensive alternative to ACTH. *Epilepsy & behavior : E&B*. 2009 Apr;14(4):674-6.
18. Hussain SA, Shinnar S, Kwong G, et al. Treatment of infantile spasms with very high dose prednisolone before high dose adrenocorticotrophic hormone. *Epilepsia*. 2014 Jan;55(1):103-7.

Response to Treatment in Infantile Spasms

19. Elterman RD, Shields WD, Mansfield KA, Nakagawa J, Group USISVS. Randomized trial of vigabatrin in patients with infantile spasms. *Neurology*. 2001 Oct 23;57(8):1416-21.
20. Vigeveno F, Cilio MR. Vigabatrin versus ACTH as first-line treatment for infantile spasms: a randomized, prospective study. *Epilepsia*. 1997 Dec;38(12):1270-4.
21. Lombroso CT. A prospective study of infantile spasms: clinical and therapeutic correlations. *Epilepsia*. 1983 Apr;24(2):135-58.
22. Cohen-Sadan S, Kramer U, Ben-Zeev B, et al. Multicenter long-term follow-up of children with idiopathic West syndrome: ACTH versus vigabatrin. *European journal of neurology : the official journal of the European Federation of Neurological Societies*. 2009 Apr;16(4):482-7.
23. Lux AL, Osborne JP. A proposal for case definitions and outcome measures in studies of infantile spasms and West syndrome: consensus statement of the West Delphi group. *Epilepsia*. 2004 Nov;45(11):1416-28.
24. Haycock GB, Schwartz GJ, Wisotsky DH. Geometric method for measuring body surface area: a height-weight formula validated in infants, children, and adults. *The Journal of pediatrics*. 1978 Jul;93(1):62-6.
25. Berg AT, Berkovic SF, Brodie MJ, et al. Revised terminology and concepts for organization of seizures and epilepsies: report of the ILAE Commission on Classification and Terminology, 2005-2009. *Epilepsia*. 2010 Apr;51(4):676-85.
26. Kleinman LC, Norton EC. What's the Risk? A simple approach for estimating adjusted risk measures from nonlinear models including logistic regression. *Health Serv Res*. 2009 Feb;44(1):288-302.

Response to Treatment in Infantile Spasms

27. Yanagaki S, Oguni H, Hayashi K, et al. A comparative study of high-dose and low-dose ACTH therapy for West syndrome. *Brain & development*. 1999 Oct;21(7):461-7.
28. Fedak EM, Patel AD, Heyer GL, Wood EG, Mytinger JR. Optimizing Care With a Standardized Management Protocol for Patients With Infantile Spasms. *Journal of child neurology*. 2014 Dec 22.
29. Hrachovy RA, Frost JD, Jr., Kellaway P, Zion T. A controlled study of ACTH therapy in infantile spasms. *Epilepsia*. 1980 Dec;21(6):631-6.
30. Costantini L, D'Ilario J, Moddemann D, Penner K, Schmidt B. Accuracy of Bayley scores as outcome measures in trials of neonatal therapies. *JAMA Pediatr*. 2015 Feb;169(2):188-9.
31. Hack M, Taylor HG, Drotar D, et al. Poor predictive validity of the Bayley Scales of Infant Development for cognitive function of extremely low birth weight children at school age. *Pediatrics*. 2005 Aug;116(2):333-41.

[AuthorsTable 6](#)

Author	Institution	Role
Wendy G Mitchell, MD Tobias Loddenkemper, MD-MPH	Children's Hospital Los Angeles Keck School of Medicine, University of Southern California Department of NeurologyBoston Children's	Data collection, edit manuscript Data collection, idea conception, edit manuscript
Amy Brooks-Kayal, MD Elaine C. Wirrell, MD	Department of Pediatrics and Neurology, University of Colorado Divisions of Epilepsy and Child and Adolescent Neurology, Department of Neurology, Mayo Clinic, Rochester MN	Data collection, edit manuscript Data collection, idea conception, edit manuscript
Cynthia Keator, MD Wendy G	Jane and John Justin	Data collection, edit

Formatted: Font: Not Italic, Font color: Auto

Response to Treatment in Infantile Spasms

Mitchell, MD	Neurosciences Department, Cook Children's Hospital Children's Hospital Los Angeles Keck School of Medicine, University of Southern California Department of Neurology	manuscript Data collection, edit manuscript
Shilpi Kumar MD Amy Brooks-Kayal, MD	Department of Pediatrics Wright State University, Dayton, Ohio Department of Pediatrics and Neurology, University of Colorado	Data collection, edit manuscript Data collection, edit manuscript
Gogi Kumar MD Cynthia Keator, MD	Department of Pediatrics Wright State University, Dayton, Ohio Jane and John Justin Neurosciences Department, Cook Children's Hospital	Data collection, edit manuscript Data collection, edit manuscript
Shilpi Kumar MD	The Children's Medical Center of Dayton	Data collection, edit manuscript
Gogi Kumar MD	The Children's Medical Center of Dayton	Data collection, edit manuscript
Zachary Grinspan MD, MS	Weill Cornell Medical Center, New York, NY	Data collection, idea conception, edit manuscript
Sarah A Kelley MD Adam L. Hartman MD	Departments of Neurology and Pediatrics Johns Hopkins Hospital Departments of Neurology and Pediatrics Johns Hopkins Hospital	Data collection, edit manuscript Data collection, idea conception, edit manuscript
Laura A Jansen MD, PhD Eric H. W. Kossoff, M.D.	University of Virginia Departments of Neurology and Pediatrics Johns Hopkins Hospital	Data collection, edit manuscript Data collection, idea conception, edit manuscript
Elissa Yozawitz MD Sarah A Kelley MD	Department of Neurology and Pediatrics Montefiore Medical Center Albert Einstein College of Medicine Departments of Neurology and Pediatrics Johns Hopkins Hospital	Data collection, edit manuscript Data collection, edit manuscript
Laura A Jansen MD, PhD	University of Virginia	Data collection, edit manuscript
Elissa Yozawitz MD	Department of Neurology and Pediatrics Montefiore Medical Center Albert Einstein College of Medicine	Data collection, edit manuscript

Formatted: Indent: First line: 0"

Response to Treatment in Infantile Spasms

Charuta N Joshi MBBS William D Gaillard MD	Division of Pediatric Neurology, Children's Hospital- Iowa Center For Neuroscience, Children's National Health System,	Data collection, edit manuscript Data collection, idea conception, edit manuscript
Ignacio Valencia MD John R Mytinger MD	St Christopher's Hospital for Children, Drexel University College of Medicine The Ohio State University, Nationwide Children's Hospital, Department of Pediatrics, Division of Pediatric Neurology, Columbus, Ohio	Data collection, edit manuscript Data collection, idea conception, edit manuscript
Charuta N Joshi MBBS	Division of Pediatric Neurology, Children's Hospital- Iowa.	Data collection, edit manuscript
Ignacio Valencia MD	St Christopher's Hospital for Children, Drexel University College of Medicine	Data collection, edit manuscript
Sucheta Joshi MD	University of Michigan	Data collection, idea conception, edit manuscript
Renée A Shollhaas MD, MS	University of Michigan	Idea conception Data collection, edit manuscript,
Lawrence Brown MD Joseph Sullivan MD	Division of Neurology, The Children's Hospital of Philadelphia and Perelman School of Medicine at the University of Pennsylvania, Philadelphia Department of Pediatrics and Neurology, University of San Francisco,	Data collection, edit manuscript Data collection, idea conception, edit manuscript
Courtney J. Wusthoff, MD, MS Dennis Dlugos MD	Department of Neurology & Neurological Sciences and by courtesy, Pediatrics- Neonatal and Developmental Medicine Stanford Division of Child Neurology Division of Neurology, The Children's Hospital of Philadelphia and Perelman School of Medicine at the University of Pennsylvania, Philadelphia,	Data collection, edit manuscript Data collection, idea conception, edit manuscript
Edward J. Novotny, MD Lawrence Brown MD	Departments of Neurology, Pediatrics, Neurosurgery and Radiology University of Washington Seattle Children's Hospital	Data collection, edit manuscript Data collection, edit manuscript

Formatted: Font: Not Bold, Not Italic, Font color: Auto

Formatted: Font: Not Italic, Font color: Black

Formatted: Font: Not Italic, Font color: Auto

Response to Treatment in Infantile Spasms

	Division of Neurology, The Children's Hospital of Philadelphia and Perelman School of Medicine at the University of Pennsylvania, Philadelphia	
Russell P. Saneto DO, PhD Gourtney J. Wusthoff, MD MS	Seattle Children's/University of Washington Division of Pediatric Neurology Department of Neurology & Neurological Sciences and by courtesy, Pediatrics-Neonatal and Developmental Medicine Stanford Division of Child Neurology	Data collection, edit manuscript Data collection, edit manuscript
Edward Novotny MD	University of Washington	Data collection, edit manuscript
Shaun A Hussain MD, MS Russell P. Saneto DO, PhD	Division of Pediatric Neurology David Geffen School of Medicine Mattel Children's Hospital UCLA Los Angeles, CA, USA 9009 Seattle Children's/University of Washington Division of Pediatric Neurology	Data collection, edit manuscript Data collection, edit manuscript
Elizabeth Thelle MD, PhD Lorie Hamiwka MD	Massachusetts General Hospital University of Washington	Data collection, edit manuscript Data collection, idea conception, edit manuscript
Catherine Chu MD Shaun A Hussain MD MS	Massachusetts General Hospital Division of Pediatric Neurology David Geffen School of Medicine Mattel Children's Hospital UCLA Los Angeles, CA, USA 90095	Data collection, edit manuscript Data collection, edit manuscript
Elizabeth Thelle MD PhD	Massachusetts General Hospital	Data collection, edit manuscript
Catherine Chu MD	Massachusetts General Hospital	Data collection, edit manuscript
Anne T. Berg, PhD	Lurie Children's Hospital Ann & Robert H. Lurie Children's Hospital of Chicago; Northwestern University Feinberg School of Medicine	Data collection, idea conception, edit manuscript
Cynthia Stack MD John J Millichap MD	Ann & Robert H. Lurie Children's Hospital of Chicago; and Departments of Pediatrics and Neurology, Northwestern University Feinberg School of Medicine Ann & Robert H. Lurie	Data collection, edit manuscript Data collection, idea conception, edit manuscript

Formatted: None, Space Before: 0 pt, After: 12 pt, Don't keep with next, Don't keep lines together

Formatted: Font: Not Italic, Font color: Auto

Formatted: Left

Formatted: Left

Response to Treatment in Infantile Spasms

	Children's Hospital of Chicago; and Departments of Pediatrics and Neurology, Northwestern University Feinberg School of Medicine	
Douglas R Nordli, Jr., MD	Ann & Robert H. Lurie Children's Hospital of Chicago; and Departments of Pediatrics and Neurology, Northwestern University Feinberg School of Medicine	Date collection, idea conception, edit manuscript
Cynthia Stack MD	Ann & Robert H. Lurie Children's Hospital of Chicago; and Departments of Pediatrics and Neurology, Northwestern University Feinberg School of Medicine	Data collection, edit manuscript

Accepted Article

Page 1: [1] Formatted Knupp, Kelly 12/17/2015 2:12:00 PM

Font: Font color: Black, Superscript

Page 1: [2] Formatted Knupp, Kelly 12/17/2015 2:12:00 PM

Font: Font color: Black, Superscript

Page 1: [3] Formatted Knupp, Kelly 12/18/2015 9:27:00 AM

Font: (Default) Arial, 10 pt, Not Bold, Font color: Auto

Page 1: [4] Formatted Knupp, Kelly 12/17/2015 2:36:00 PM

Line spacing: Double, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between : (No border), Bar : (No border)

Page 1: [5] Formatted Knupp, Kelly 12/17/2015 2:36:00 PM

Line spacing: Double, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between : (No border), Bar : (No border)

Page 1: [6] Formatted Kelly Knupp 12/21/2015 11:09:00 AM

Font: (Default) Arial, 10 pt, Font color: Auto

Accepted

Table 1 Treatment Guidelines Recommended by PERC

Treatment with intramuscular (IM) ACTH

Days	Dose – ACTH
1-14	75 U/m ² twice daily
15-17	30 U/m ² in the morning
18-20	15 U/m ² in the morning
21-23	10 U/m ² in the morning
24-29	10 U/m ² every other morning (3 total doses)

#If there is no clinical response by day 14, consider alternative treatment.

Treatment with oral prednisolone

Days	Dose – prednisolone
1-14	10 mg oral 4 times daily*#
15-19	10 mg oral 3 times daily*
20-24	10 mg oral 2 times daily
25-29	10 mg oral daily

*If there is no clinical response after day 7 (i.e. no 24 hour period free of infantile spasms), the dose can be increased to 20 mg three times daily. If done, the taper schedule from day 15-19 would be 10 mg 4 times daily and then proceed as in the table beginning on day 20.

#If no clinical response by day 14, consider alternative treatment.

Treatment with oral vigabatrin

Days*#	Dose – Vigabatrin
1 - 3	50 mg/kg/day divided 2 times daily
4 - 6	100 mg/kg/day divided 2 times daily
>7	150 mg/kg/day divided 2 times daily

*Side effects (e.g. sedation, hypotonia) may necessitate slower titration

#If no clinical response by day 14, consider alternative treatment

Table 2: Baseline characteristics by type of spasm treatment

Characteristic		Spasm treatment				Total N=230	P- value*
		ACTH N=97	Oral cortico- steroid N=54	Vigabatrin N=47	Other N=32		
Sex	Male	54 (56)	26 (48)	24 (51)	18 (56)	122 (53)	0.80
Race	Black	12 (14)	8 (17)	4 (9)	7 (24)	31 (15)	0.27
	White	59 (66)	34 (74)	35 (80)	19 (66)	147 (71)	
	Other	18 (20)	4 (9)	5 (11)	3 (10)	30 (14)	
Ethnicity	Hispanic	17 (20)	2 (5)	8 (18)	4 (15)	31 (16)	0.21
Gestational age	Weeks	39 (37,40)	40 (38,40)	39 (37,40)	38 (33,40)	39 (37,40)	0.10
Age at spasm onset	Months	6.0 (4.5, 8.0)	7.0 (4.5,9.0)	6.0 (4.5, 7.7)	7.7 (6.0,10.0)	6.5 (4.5, 8.5)	0.03
First spasm to diagnosis	Days	14 (5,32)	10 (5,18)	14 (4,73)	18 (2,28)	13 (5,31)	0.26
First spasm to treatment start	Days	16 (6,37)	12 (6,21)	21 (6,74)	15 (3,47)	15 (6,37)	0.13
Diagnosis to treatment start	Days	1 (0,2)	0 (0,1)	2 (1,5)	0 (-1,9)	1 (0,2)	<0.001
Prior seizures		19 (20)	17 (32)	19 (40)	20 (63)	75 (33)	<0.001
History of AED use		21 (22)	22 (41)	21 (45)	19 (59)	83 (36)	<0.001
Etiology**	Genetic/metabolic	18 (19)	11 (20)	13 (28)	8 (25)	50 (22)	<0.001
	Prior brain injury	17 (18)	15 (28)	11 (23)	9 (28)	52 (23)	
	MCD/other structural	9 (9)	5 (9)	16 (34)	8 (25)	38 (17)	
	Unknown abnormal	29 (30)	16 (30)	6 (13)	5 (16)	56 (24)	
	Unknown normal	24 (25)	7 (13)	1 (2)	2 (6)	34 (15)	
Developmental issues	None	31 (32)	7 (13)	2 (4)	3 (10)	43 (19)	<0.001
	Mild	10 (10)	6 (11)	4 (9)	1 (3)	21 (9)	
	Moderate	15 (16)	9 (17)	12 (26)	2 (7)	38 (17)	
	Severe	40 (42)	31 (59)	29 (62)	25 (81)	125 (55)	

Values are N (column %) or median (Q1, Q3).

*Chi-square test for categorical variables, Kruskal-Wallis test for continuous variables.

**There were 11 participants with tuberous sclerosis (included in the MCD/other structural etiology group). Eight were on vigabatrin, 1 on ACTH, and 2 Other.

Missing values (N): race (22), ethnicity (35), gestational age (2), age at spasm onset (5), time between first spasm and diagnosis (2), time between first spasm and treatment start (3), time between diagnosis and treatment start (3), and development (3).

Table 3: Characteristics by response to first spasm treatment

Characteristic		Response to treatment				P-value*
		Non-response		Response		
		N=136		N=94		
Sex	Female	63	(58)	45	(42)	0.82
	Male	73	(60)	49	(40)	
Race	Black	20	(65)	11	(36)	0.91
	White	89	(61)	58	(40)	
	Other	18	(60)	12	(40)	
Ethnicity	Hispanic	17	(55)	14	(45)	0.44
	Non-Hispanic	102	(62)	18	(51)	
Gestational age	<37 weeks	30	(63)	18	(38)	0.60
	At least 37 weeks	105	(58)	75	(42)	
Age at spasm onset	<12 months	119	(59)	84	(41)	0.97
	At least 12 months	13	(59)	9	(41)	
First spasm to treatment start	Within 4 weeks	93	(61)	59	(39)	0.35
	> 4 weeks	41	(55)	34	(45)	
Prior seizures	Yes	55	(73)	20	(27)	0.002
	No	81	(52)	74	(48)	
History of AED use	Yes	58	(70)	25	(30)	0.013
	No	78	(53)	69	(47)	
Etiology	Genetic/metabolic/unknown abnormal	66	(62)	40	(38)	0.070
	Prior brain injury/MCD/other structural	56	(62)	34	(38)	
	Unknown normal	14	(41)	20	(59)	
Developmental issues	None/Mild	30	(47)	34	(53)	0.025
	Moderate/Severe	103	(63)	60	(37)	
Treatment	ACTH	44	(45)	53	(55)	<0.001
	Oral corticosteroid	33	(61)	21	(39)	
	Vigabatrin	30	(64)	17	(36)	
	Other	29	(91)	3	(9)	

Values are N (row %)

*Chi-square test

Missing values (N): race (22), ethnicity (35), gestational age (2), age at spasm onset (5), time between first spasm and diagnosis (2), time between first spasm and treatment start (3), time between diagnosis and treatment start (3), and development (3).

Table 4: Relative risk of response, by treatment, etiology, and development

Covariate		Crude Relative Risk (95% CI)	Adjusted* Relative Risk (95% CI)	Adjusted** Relative Risk (95% CI)
Treatment	ACTH	5.81 (4.10, 9.40)	5.50 (3.83, 9.18)	5.24 (3.59, 8.70)
	Oral corticosteroid	4.15 (2.78, 6.76)	4.11 (2.78, 6.52)	3.88 (2.63, 6.46)
	Vigabatrin	3.91 (2.60, 6.45)	3.92 (2.64, 6.50)	3.69 (2.55, 6.18)
	Other	Reference	Reference	Reference
Etiology	Genetic/metabolic/unknown abnormal	Reference	Reference	---
	Prior brain injury/MCD/other structural	1.00 (0.69, 1.43)	1.06 (0.98, 1.16)	---
	Unknown normal	1.56 (1.04, 2.22)	1.39 (0.89, 1.99)	---
Development	None/Mild	1.42 (1.04, 2.01)	---	1.23 (0.86, 1.72)
	Moderate/Severe	Reference	---	Reference

Relative risks, or relative probabilities, of treatment response between groups were estimated using the method of Kleinman and Norton applied logistic regression models; confidence intervals estimated via bootstrapping

*Model including treatment and etiology as covariates

**Model including treatment and development as covariates

Accepte

Table 5: Etiology, Development, and Treatment effect on Response

Characteristic	Treatment	Response to treatment				
		Non-response		Response		Predicted Probability of response* (95% CI)
		N=136		N=94		
Etiology						
Genetic/metabolic/unknown abnormal	ACTH	25	(53)	22	(47)	49% (37, 62)
	Oral corticosteroid	15	(56)	12	(44)	35% (23, 50)
	Vigabatrin	13	(68)	6	(34)	33% (20, 50)
	Other	13	(100)	0	(0)	8% (3, 24)
Prior brain injury/MCD/other structural	ACTH	10	(39)	16	(62)	55% (40, 68)
	Oral corticosteroid	15	(75)	5	(25)	40% (26, 56)
	Vigabatrin	16	(59)	11	(41)	38% (24, 54)
	Other	15	(88)	2	(12)	10% (3, 27)
Unknown normal	ACTH	9	(38)	15	(63)	65% (48, 80)
	Oral corticosteroid	3	(43)	4	(57)	51% (30, 71)
	Vigabatrin	1	(100)	0	(0)	49% (26, 72)
	Other	1	(50)	1	(50)	15% (4, 41)
Developmental issues						
None/Mild	ACTH	17	(42)	24	(59)	61% (47, 73)
	Oral corticosteroid	7	(54)	6	(46)	47% (30, 64)
	Vigabatrin	3	(50)	3	(50)	44% (26, 64)
	Other	3	(75)	1	(25)	13% (4, 35)
Moderate/Severe	ACTH	26	(47)	29	(53)	51% (39, 63)
	Oral corticosteroid	25	(63)	15	(38)	37% (25, 52)
	Vigabatrin	27	(66)	14	(34)	35% (23, 50)
	Other	25	(93)	2	(7)	9% (3, 25)

Values are N (row %)

*Predicted probabilities of response to treatment (expressed as a percentage) were estimated via two logistic regression models: (1) model containing etiology (overall p-value p=0.29) and treatment (p=0.0016) as covariates, (2) model containing development (p=0.24) and treatment (p=0.0023) as covariates. ~~Once adjusted for treatment, etiology (p=0.29) and development (p=0.24) were no longer significant predictors of treatment response.~~