

# RESTRICTED PERMUTATIONS, CONTINUED FRACTIONS, AND CHEBYSHEV POLYNOMIALS

TOUFIK MANSOUR\* AND ALEK VAINSHTEIN†

\* Department of Mathematics

† Department of Mathematics and Department of Computer Science  
University of Haifa, Haifa, Israel 31905  
tmansur@study.haifa.ac.il, alek@mathcs.haifa.ac.il

Submitted: December 21, 1999; Accepted: February 27, 2000.

ABSTRACT. Let  $f_n^r(k)$  be the number of 132-avoiding permutations on  $n$  letters that contain exactly  $r$  occurrences of  $12\dots k$ , and let  $F_r(x; k)$  and  $F(x, y; k)$  be the generating functions defined by  $F_r(x; k) = \sum_{n \geq 0} f_n^r(k)x^n$  and  $F(x, y; k) = \sum_{r \geq 0} F_r(x; k)y^r$ . We find an explicit expression for  $F(x, y; k)$  in the form of a continued fraction. This allows us to express  $F_r(x; k)$  for  $1 \leq r \leq k$  via Chebyshev polynomials of the second kind.

2000 MATHEMATICS SUBJECT CLASSIFICATION: Primary 05A05, 05A15; Secondary 30B70, 42C05

Typeset by  $\mathcal{A}\mathcal{M}\mathcal{S}$ - $\mathcal{T}\mathcal{E}\mathcal{X}$

## 1. INTRODUCTION

Let  $[p] = \{1, \dots, p\}$  denote a totally ordered alphabet on  $p$  letters, and let  $\alpha = (\alpha_1, \dots, \alpha_m) \in [p_1]^m$ ,  $\beta = (\beta_1, \dots, \beta_m) \in [p_2]^m$ . We say that  $\alpha$  is *order-isomorphic* to  $\beta$  if for all  $1 \leq i < j \leq m$  one has  $\alpha_i < \alpha_j$  if and only if  $\beta_i < \beta_j$ . For two permutations  $\pi \in \mathfrak{S}_n$  and  $\tau \in \mathfrak{S}_k$ , an *occurrence* of  $\tau$  in  $\pi$  is a subsequence  $1 \leq i_1 < i_2 < \dots < i_k \leq n$  such that  $(\pi_{i_1}, \dots, \pi_{i_k})$  is order-isomorphic to  $\tau$ ; in such a context  $\tau$  is usually called the *pattern*. We say that  $\pi$  *avoids*  $\tau$ , or is  $\tau$ -*avoiding*, if there is no occurrence of  $\tau$  in  $\pi$ . The set of all  $\tau$ -avoiding permutations of all possible sizes including the empty permutation is denoted  $\mathfrak{S}(\tau)$ . Pattern avoidance proved to be a useful language in a variety of seemingly unrelated problems, from stack sorting [5] to singularities of Schubert varieties [6]. A complete study of pattern avoidance for the case  $\tau \in \mathfrak{S}_3$  is carried out in [11]. For the case  $\tau \in \mathfrak{S}_4$  see [14, 11, 12, 1].

A natural generalization of pattern avoidance is the restricted pattern inclusion, when a prescribed number of occurrences of  $\tau$  in  $\pi$  is required. Papers [8] and [3] contain simple expressions for the number of permutations containing exactly one 123 and 132 patterns, respectively. The main result of [B2] is that the generating function for the number of permutations containing exactly  $r$  132 patterns is a rational function in variables  $x$  and  $\sqrt{1-4x}$ . This proves a particular case of the general conjecture of Noonan and Zeilberger [9] which is that for any set  $T$  of patterns, the sequence of numbers enumerating permutations having a prescribed number of occurrences of patterns in  $T$  is  $P$ -recursive. Recent paper [10] presents the generating function for the number of 132-avoiding permutations that contain a prescribed number of 123 patterns. The generating function is given in the form of a continued fraction. In the present note we generalize the argument of [10] to get the generating function for the number of 132-avoiding permutations that contain a prescribed number of  $12\dots k$  patterns for arbitrary  $k \geq 3$ . The study of the obtained continued fraction allows us to recover and to generalize the result of [4] that relates the number of 132-avoiding permutations that contain no  $12\dots k$  patterns to Chebyshev polynomials of the second kind.

The authors are grateful to C. Krattenthaler, H. Wilf, and anonymous referee for useful comments concerning Theorems 4.1 and 4.2.

## 2. CONTINUED FRACTIONS

Let  $f_n^r(k)$  stand for the number of 132-avoiding permutations on  $n$  letters that contain exactly  $r$  occurrences of  $12\dots k$ . We denote by  $F(x, y; k)$  the generating function of the sequence  $\{f_n^r(k)\}$ , that is,

$$F(x, y; k) = \sum_{n \geq 0} \sum_{r \geq 0} f_n^r(k) x^n y^r.$$

Our first result is a natural generalization of the main theorem of [10].

**Theorem 2.1.** *The generating function  $F(x, y; k)$  for  $k \geq 1$  is given by the continued fraction*

$$F(x, y; k) = \frac{1}{1 - \frac{xy^{d_1}}{1 - \frac{xy^{d_2}}{1 - \frac{xy^{d_3}}{\dots}}}}$$

where  $d_i = \binom{i-1}{k-1}$ , and  $\binom{a}{b}$  is assumed 0 whenever  $a < b$  or  $b < 0$ .

*Proof.* Following [10] we define  $\eta_j(\pi)$ ,  $j \geq 1$ , as the number of occurrences of  $12 \dots j$  in  $\pi$ . Define  $\eta_0(\pi) = 1$  for any  $\pi$ , which means that the empty pattern occurs exactly once in each permutation. The *weight* of a permutation  $\pi$  is a monomial in  $k$  independent variables  $q_1, \dots, q_k$  defined by

$$w_k(\pi) = \prod_{j=1}^k q_j^{\eta_j(\pi)}.$$

The *total weight* is a polynomial

$$W_k(q_1, \dots, q_k) = \sum_{\pi \in \mathfrak{S}(132)} w_k(\pi).$$

The following proposition is implied immediately by the definitions.

**Proposition 2.1.**  *$F(x, y; k) = W_k(x, 1, \dots, 1, y)$  for  $k \geq 2$ , and  $F(x, y; 1) = W_1(xy)$ .*

We now find a recurrence relation for the numbers  $\eta_j(\pi)$ . Let  $\pi \in \mathfrak{S}_n$ , so that  $\pi = (\pi', n, \pi'')$ .

**Proposition 2.2.** *For any  $j \geq 1$  and any nonempty  $\pi \in \mathfrak{S}(132)$*

$$\eta_j(\pi) = \eta_j(\pi') + \eta_j(\pi'') + \eta_{j-1}(\pi').$$

*Proof.* Let  $l = \pi^{-1}(n)$ . Since  $\pi$  avoids 132, each number in  $\pi'$  is greater than any of the numbers in  $\pi''$ . Therefore,  $\pi'$  is a 132-avoiding permutation of the numbers  $\{n-l+1, n-l+2, \dots, n-1\}$ , while  $\pi''$  is a 132-avoiding permutation of the numbers  $\{1, 2, \dots, n-l\}$ . On the other hand, if  $\pi'$  is an arbitrary 132-avoiding permutation of the numbers  $\{n-l+1, n-l+2, \dots, n-1\}$  and  $\pi''$  is an arbitrary 132-avoiding permutation of the numbers  $\{1, 2, \dots, n-l\}$ , then  $\pi = (\pi', n, \pi'')$  is 132-avoiding. Finally, if  $(i_1, \dots, i_j)$  is an occurrence of  $12 \dots j$  in  $\pi$  then either  $i_j < l$ , and so it is also an occurrence of  $12 \dots j$  in  $\pi'$ , or  $i_1 > l$ , and so it is also an occurrence of  $12 \dots j$  in  $\pi''$ , or  $i_j = l$ , and so  $(i_1, \dots, i_{j-1})$  is an occurrence of  $12 \dots j-1$  in  $\pi'$ . The result follows.  $\square$

Now we are able to find the recurrence relation for the total weight  $W$ . Indeed, by Proposition 2.2,

$$\begin{aligned} W_k(q_1, \dots, q_k) &= 1 + \sum_{\emptyset \neq \pi \in \mathfrak{S}(132)} \prod_{j=1}^k q_j^{\eta_j(\pi') + \eta_j(\pi'') + \eta_{j-1}(\pi')} \\ &= 1 + \sum_{\pi' \in \mathfrak{S}(132)} \sum_{\pi'' \in \mathfrak{S}(132)} \prod_{j=1}^k q_j^{\eta_j(\pi'')} \cdot q_1 \prod_{j=1}^{k-1} (q_j q_{j+1})^{\eta_j(\pi')} \cdot q_k^{\eta_k(\pi')} \\ &= 1 + q_1 W_k(q_1, \dots, q_k) W_k(q_1 q_2, q_2 q_3, \dots, q_{k-1} q_k, q_k). \end{aligned} \tag{1}$$

For any  $d \geq 0$  and  $1 \leq m \leq k$  define

$$\mathbf{q}^{d,m} = \prod_{j=1}^k q_j^{\binom{d}{j-m}};$$

recall that  $\binom{a}{b} = 0$  if  $a < b$  or  $b < 0$ . The following proposition is implied immediately by the well-known properties of binomial coefficients.

**Proposition 2.3.** *For any  $d \geq 0$  and  $1 \leq m \leq k$*

$$\mathbf{q}^{d,m} \mathbf{q}^{d,m+1} = \mathbf{q}^{d+1,m}.$$

Observe now that  $W_k(q_1, \dots, q_k) = W_k(\mathbf{q}^{0,1}, \dots, \mathbf{q}^{0,k})$  and that by (1) and Proposition 2.3

$$W_k(\mathbf{q}^{d,1}, \dots, \mathbf{q}^{d,k}) = 1 + \mathbf{q}^{d,1} W_k(\mathbf{q}^{d,1}, \dots, \mathbf{q}^{d,k}) W_k(\mathbf{q}^{d+1,1}, \dots, \mathbf{q}^{d+1,k}),$$

therefore

$$W_k(q_1, \dots, q_k) = \frac{1}{1 - \frac{\mathbf{q}^{0,1}}{1 - \frac{\mathbf{q}^{1,1}}{1 - \frac{\mathbf{q}^{2,1}}{\dots}}}}.$$

To obtain the continued fraction representation for  $F(x, y; k)$  it is enough to use Proposition 2.1 and to observe that

$$\mathbf{q}^{d,1} \Big|_{q_1=x, q_2=\dots=q_{k-1}=1, q_k=y} = xy^{\binom{d}{k-1}}. \quad \square$$

*Remark.* For  $k = 1$  one recovers from Theorem 2.1 the well-known generating function for the Catalan numbers,  $(1 - \sqrt{1 - 4z})/2z$ . This result also follows immediately from Proposition 2.1 and equation (1), which for  $k = 1$  is reduced to  $W_1(q) = 1 + qW_1^2(q)$ .

### 3. CHEBYSHEV POLYNOMIALS

Let us denote by  $F_r(x; k)$  the generating function of the sequence  $\{f_n^r(k)\}$  for a given  $r$ , that is,

$$F_r(x; k) = \sum_{n \geq 0} f_n^r(k) x^n.$$

Recall that  $F(x, y; k) = \sum_{r \geq 0} F_r(x; k) y^r$ . In this section we find explicit expressions for  $F_r(x; k)$  in the case  $0 \leq r \leq k$ .

Consider a recurrence relation

$$T_j = \frac{1}{1 - xT_{j-1}}, \quad j \geq 1. \tag{2}$$

The solution of (2) with the initial condition  $T_0 = 0$  is denoted by  $R_j(x)$ , and the solution of (2) with the initial condition

$$T_0 = G(x, y; k) = \frac{y}{1 - \frac{xy^{\binom{k}{1}}}{1 - \frac{xy^{\binom{k+1}{2}}}{1 - \frac{xy^{\binom{k+2}{3}}}{\dots}}}}$$

is denoted by  $S_j(x, y; k)$ , or just  $S_j$  when the value of  $k$  is clear from the context. Our interest in (2) is stipulated by the following relation, which is an easy consequence of Theorem 2.1:

$$F(x, y; k) = S_k(x, y; k). \tag{3}$$

First of all, we find an explicit formula for the functions  $R_j(x)$ . Let  $U_j(\cos \theta) = \sin(j + 1)\theta / \sin \theta$  be the Chebyshev polynomials of the second kind.

**Lemma 3.1.** *For any  $j \geq 1$*

$$R_j(x) = \frac{U_{j-1}\left(\frac{1}{2\sqrt{x}}\right)}{\sqrt{x}U_j\left(\frac{1}{2\sqrt{x}}\right)}. \tag{4}$$

*Proof.* Indeed, it follows immediately from (2) that  $R_j(x)$  is the  $j$ th approximant for the continued fraction

$$\frac{1}{1 - \frac{x}{1 - \frac{x}{1 - \frac{x}{\dots}}}}$$

Hence, by [7, Theorem 2, p. 194], for any  $j \geq 1$  one has  $R_j(x) = A_j(x)/A_{j+1}(x)$ , where

$$A_j(x) = \left(\frac{1 + \sqrt{1 - 4x}}{2}\right)^j - \left(\frac{1 - \sqrt{1 - 4x}}{2}\right)^j.$$

Using substitution  $x \rightarrow 1/4t^2$  one gets  $(2t)^j A_j(1/4t^2) = 2\sqrt{t^2 - 1}U_{j-1}(t)$ , which gives  $A_j(x) = \sqrt{1/x - 4}x^{j/2}U_{j-1}(1/2\sqrt{x})$ , and the result follows.  $\square$

Next, we find an explicit expression for  $S_j$  in terms of  $G$  and  $R_j$ .

**Lemma 3.2.** *For any  $j \geq 1$  and any  $k \geq 1$*

$$S_j(x, y; k) = R_j(x) \frac{1 - xR_{j-1}(x)G(x, y; k)}{1 - xR_j(x)G(x, y; k)}. \tag{5}$$

*Proof.* Indeed, from (2) and  $S_0 = G$  we get  $S_1 = 1/(1 - xG)$ . On the other hand,  $R_0 = 0, R_1 = 1$ , so (5) holds for  $j = 1$ . Now let  $j > 1$ , then by induction

$$S_j = \frac{1}{1 - xS_{j-1}} = \frac{1}{1 - xR_{j-1}} \cdot \frac{1 - xR_{j-1}G}{1 - \frac{x(1 - xR_{j-2})R_{j-1}G}{1 - xR_{j-1}}}.$$

Relation (2) for  $R_j$  and  $R_{j-1}$  yields  $(1 - xR_{j-2})R_{j-1} = (1 - xR_{j-1})R_j = 1$ , which together with the above formula gives (5).  $\square$

As a corollary from Lemma 3.2 and (3) we get the following expression for the generating function  $F(x, y; k)$ .

**Corollary.**

$$F(x, y; k) = R_k(x) + (R_k(x) - R_{k-1}(x)) \sum_{m \geq 1} (xR_k(x)G(x, y; k))^m.$$

Now we are ready to express the generating functions  $F_r(x; k), 0 \leq r \leq k$ , via Chebyshev polynomials.

**Theorem 3.1.** *For any  $k \geq 1, F_r(x; k)$  is a rational function given by*

$$F_r(x; k) = \frac{x^{\frac{r-1}{2}}U_{k-1}^{r-1}\left(\frac{1}{2\sqrt{x}}\right)}{U_k^{r+1}\left(\frac{1}{2\sqrt{x}}\right)}, \quad 1 \leq r \leq k,$$

$$F_0(x; k) = \frac{U_{k-1}\left(\frac{1}{2\sqrt{x}}\right)}{\sqrt{x}U_k\left(\frac{1}{2\sqrt{x}}\right)},$$

where  $U_j$  is the  $j$ th Chebyshev polynomial of the second kind.

*Proof.* Observe that  $G(x, y; k) = y + y^{k+1}P(x, y)$ , so from Corollary we get

$$F(x, y; k) = R_k(x) + (R_k(x) - R_{k-1}(x)) \sum_{m=1}^k (xR_k(x))^m y^m + y^{k+1}P'(x, y),$$

where  $P(x, y)$  and  $P'(x, y)$  are formal power series. To complete the proof, it suffices to use (4) together with the identity  $U_{n-1}^2(z) - U_n(z)U_{n-2}(z) = 1$ , which follows easily from the trigonometric identity  $\sin^2 n\theta - \sin^2 \theta = \sin(n+1)\theta \sin(n-1)\theta$ .  $\square$

For the case  $r = 0$  this result was proved by a different method in [4].

#### 4. FURTHER RESULTS

There are several ways to generalize the results of the previous sections. First, one can try to get exact formulas for  $F_r(x; k)$  in the case  $r > k$ . The method described in Section 3 allows, in principle, to obtain such formulas, though they become more and more complicated. For example, the following theorem gives an explicit expression for  $F_r(x; k)$  when  $r \leq k(k+3)/2$ .

**Theorem 4.1.** *For any  $k \geq 1$  and  $1 \leq r \leq k(k+3)/2$ ,  $F_r(x; k)$  is a rational function given by*

$$F_r(x; k) = \frac{x^{\frac{r-1}{2}} U_{k-1}^{r-1} \left( \frac{1}{2\sqrt{x}} \right)}{U_k^{r+1} \left( \frac{1}{2\sqrt{x}} \right)} \sum_{j=0}^{\lfloor (r-1)/k \rfloor} \binom{r - kj + j - 1}{j} \left( \frac{U_k \left( \frac{1}{2\sqrt{x}} \right)}{x^{\frac{k-2}{2k}} U_{k-1} \left( \frac{1}{2\sqrt{x}} \right)} \right)^{kj},$$

where  $U_j$  is the  $j$ th Chebyshev polynomial of the second kind.

*Proof.* Indeed, the explicit expression for  $G(x, y; k)$  gives

$$G(x, y; k) = y(1 + xy^k + \dots + x^s y^{ks}) + y^t P(x, y),$$

where  $s = \lceil (k+1)/2 \rceil$ ,  $t = 1 + k(k+3)/2$ , and  $P(x, y)$  is a formal power series. Hence, by Corollary,

$$\begin{aligned} \frac{F(x, y; k) - R_k(x)}{R_k(x) - R_{k-1}(x)} &= \sum_{m \geq 1} (xR_k(x))^m y^m (1 + xy^k + \dots + x^s y^{ks})^m + y^t P'(x, y) \\ &= \sum_{m \geq 1} (xR_k(x))^m y^m \sum_{j=0}^{ms} \binom{m+j-1}{j} x^j y^{kj} + y^t P'(x, y) \\ &= \sum_{r \geq 1} y^r (xR_k(x))^r \sum_{j=0}^{\lfloor (r-1)/k \rfloor} \frac{\binom{r-kj+j-1}{j} x^j}{(xR_k(x))^{kj}} + y^t P''(x, y), \end{aligned}$$

where  $P'(x, y)$  and  $P''(x, y)$  are formal power series. The rest of the proof follows the proof of Theorem 3.1.  $\square$

Another possibility is to analyze the case of permutations containing exactly one 132 pattern and  $r$  12... $k$  patterns. Introducing the modified total weight  $\Omega_k(q_1, \dots, q_k)$  as the sum of the weights  $w_k(\pi)$  over all permutations containing exactly one 132 pattern, we get the following equation:

$$\begin{aligned} \Omega_k(q_1, \dots, q_k) &= q_1 W_k(q_1 q_2, \dots, q_{k-1} q_k, q_k) \Omega_k(q_1, \dots, q_k) \\ &\quad + q_1 W_k(q_1, \dots, q_k) \Omega_k(q_1 q_2, \dots, q_{k-1} q_k, q_k) \\ &\quad + q_1^2 q_2^2 W_k(q_1 q_2, \dots, q_{k-1} q_k, q_k) (W_k(q_1, \dots, q_k) - 1); \end{aligned}$$

for the case  $k = 3$  see [10]. By (1) and Proposition 2.3 this is equivalent to

$$\begin{aligned} \Omega_k(\mathbf{q}^{d,1}, \dots, \mathbf{q}^{d,k}) &= \mathbf{q}^{d,1} (\mathbf{q}^{d,2})^2 (W_k(\mathbf{q}^{d,1}, \dots, \mathbf{q}^{d,k}) - 1)^2 \\ &\quad + \mathbf{q}^{d,1} W_k^2(\mathbf{q}^{d,1}, \dots, \mathbf{q}^{d,k}) \Omega_k(\mathbf{q}^{d+1,1}, \dots, \mathbf{q}^{d+1,k}). \end{aligned} \tag{6}$$

Let now  $\varphi_n^r(k)$  be the number of permutations on  $n$  letters that contain exactly one 132 pattern and  $r$  12... $k$  patterns, and  $\Phi_r(x; k)$  be the generating function of the sequence  $\{\varphi_n^r(k)\}$  for a given  $r$ . In general, equation (6) allows us to find explicit expressions for  $\Phi_r(x; k)$ . However, they are rather cumbersome, so we restrict ourselves to the case  $r = 0$ .

**Theorem 4.2.** *For any  $k \geq 3$ ,  $\Phi_0(x; k)$  is a rational function given by*

$$\begin{aligned} \Phi_0(x; k) &= \frac{x}{U_k^2\left(\frac{1}{2\sqrt{x}}\right)} \sum_{j=1}^{k-2} U_j^2\left(\frac{1}{2\sqrt{x}}\right) \\ &= \frac{1}{16 \sin^2(k+1)t \cos^2 t} \left( 2k - 5 + 4 \cos^2 t - \frac{\sin(2k-1)t}{\sin t} \right), \end{aligned}$$

where  $U_j$  is the  $j$ th Chebyshev polynomial of the second kind and  $\cos t = 1/2\sqrt{x}$ .



## REFERENCES

1. M. Bona, *Permutations avoiding certain patterns: the case of length 4 and some generalizations*, Discr. Math. **175** (1997), 55–67.
2. M. Bona, *The number of permutations with exactly  $r$  132-subsequences is  $P$ -recursive in the size!*, Adv. Appl. Math. **18** (1997), 510–522.
3. M. Bona, *Permutations with one or two 132-subsequences*, Discr. Math. **181** (1998), 267–274.
4. T. Chow and J. West, *Forbidden subsequences and Chebyshev polynomials*, Discr. Math. **204** (1999), 119–128.
5. D. Knuth, *The Art of Computer Programming*, vol. 3, Addison Wesley, Reading, MA, 1973.
6. V. Lakshmibai and B. Sandhya, *Criterion for smoothness of Schubert varieties in  $Sl(n)/B$* , Proc. Indian Acad. Sci. **100** (1990), no. 1, 45–52.
7. L. Lorentzen and H. Waadeland, *Continued fractions with applications*, North-Holland, 1992.
8. J. Noonan, *The number of permutations containing exactly one increasing subsequence of length three*, Discr. Math. **152** (1996), 307–313.
9. J. Noonan and D. Zeilberger, *The enumeration of permutations with a prescribed number of “forbidden” patterns*, Adv. Appl. Math. **17** (1996), 381–407.
10. A. Robertson, H. Wilf, and D. Zeilberger, *Permutation patterns and continuous fractions*, Elec. J. Comb. **6** (1999), no. 1, R38.
11. R. Simion and F. Schmidt, *Restricted permutations*, Europ. J. Comb. **6** (1985), 383–406.
12. Z. Stankova, *Classification of forbidden subsequences of length 4*, Eur. J. Comb. **17** (1996), 501–517.
13. Z. Stankova, *Forbidden subsequences*, Discr. Math. **132** (1994), 291–316.
14. J. West, *Generating trees and the Catalan and Schröder numbers*, Discr. Math. **146** (1995), 247–262.