



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

## Reverse-Transcriptase Inhibitors in the Aicardi–Goutières Syndrome

### Citation for published version:

Rice, GI, Meyzer, C, Bouazza, N, Hully, M, Boddaert, N, Semeraro, M, Zeef, LAH, Rozenberg, F, Bondet, V, Duffy, D, Llibre, A, Baek, J, Sambe, MN, Henry, E, Jolaine, V, Barnerias, C, Barth, M, Belot, A, Cances, C, Debray, F-G, Doummar, D, Frémond, M-L, Kitabayashi, N, Lepelley, A, Levrat, V, Melki, I, Meyer, P, Nougues, M-C, Renaldo, F, Rodero, MP, Rodriguez, D, Roubertie, A, Seabra, L, Ugenti, C, Abdoul, H, Treluyer, J-M, Desguerre, I, Blanche, S & Crow, YJ 2018, 'Reverse-Transcriptase Inhibitors in the Aicardi–Goutières Syndrome', *New England Journal of Medicine*, vol. 379, no. 23, pp. 2275-7. <https://doi.org/10.1056/NEJMc1810983>

### Digital Object Identifier (DOI):

[10.1056/NEJMc1810983](https://doi.org/10.1056/NEJMc1810983)

### Link:

[Link to publication record in Edinburgh Research Explorer](#)

### Document Version:

Publisher's PDF, also known as Version of record

### Published In:

New England Journal of Medicine

### General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

### Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



## CORRESPONDENCE



## Reverse-Transcriptase Inhibitors in the Aicardi–Goutières Syndrome


**TO THE EDITOR:** The Aicardi–Goutières syndrome is a genetic encephalopathy that is associated with childhood illness and death. The syndrome is hypothesized to be due to misidentification of self-derived nucleic acids as nonself and the subsequent induction of a type I interferon–mediated response that simulates an antiviral reaction.<sup>1</sup> Endogenous retroelements, mobile genetic elements that can be transcribed to RNA and then to DNA by reverse transcription, constitute 40% of the human genome and represent a potential source of immunostimulatory nucleic acid in patients with this syndrome.<sup>2</sup>

In a single-center, open-label, pilot study involving patients with the Aicardi–Goutières syndrome (ClinicalTrials.gov number, NCT02363452), we administered a combination of three nucleoside analogue reverse-transcriptase inhibitors — abacavir, lamivudine, and zidovudine — for 12 months, at doses used in children with human immunodeficiency virus type 1 (HIV-1) infection. The study protocol is available with the full text of this letter at NEJM.org. The primary aim was to determine the effect of treatment on the interferon score, calculated from the expression of six interferon-stimulated genes; higher scores indicate greater interferon signaling, and scores higher than 2.47 are considered to be abnormal.<sup>3</sup> Interferon status was also determined by measurement of interferon- $\alpha$  protein levels in serum, plasma, and cerebrospinal fluid (CSF); the antiviral protective capacity (interferon activity) of patient serum and CSF; and genomewide sequencing of RNA extracted from whole blood. Clinical features and cerebral blood flow (measured by means of arterial spin labeling magnetic resonance imaging) were secondary efficacy measures.

Eight of 11 patients who were recruited from a pool of 68 patients in France known to have the syndrome completed the study (Table S2 in the Supplementary Appendix, available at NEJM.org). Three patients withdrew owing to an inability to swallow the volume of the study medication. There was an effect of treatment on interferon signaling, with the median interferon score across all 8 patients falling from 9.66 (interquartile range, 6.51 to 13.23) to 5.33 (interquartile range, 2.76 to 10.90) ( $P < 0.001$ ) (Fig. 1A). Interferon- $\alpha$  protein levels in serum and plasma and interferon antiviral activity in CSF were also reduced with treatment (Table S3 in the Supplementary Appendix). This effect was greatest among the 4 patients with mutations in components of the RNase H2 complex (with the median score in these 4 patients falling from 8.16 [interquartile range, 5.41 to 11.94] to 3.51 [interquartile range, 2.49 to 5.46]). RNA sequencing indicated a reduction of global interferon-stimulated gene expression after 12 months of treatment and a return

### THIS WEEK'S LETTERS

- 2275** Reverse-Transcriptase Inhibitors in the Aicardi–Goutières Syndrome
- 2277** Labor Induction vs. Expectant Management of Low-Risk Pregnancy
- 2279** Thromboprophylaxis after Hospitalization for Medical Illness
- 2280** Aspirin-Exacerbated Respiratory Disease
- 2282** Sequencing Circulating cfDNA during Pregnancy



**Figure 1.** Measures of Interferon Status in Eight Patients Completing 12 Months of Therapy.

Panel A shows the interferon score in blood, with higher scores indicating greater interferon signaling. There were 18 recorded values before the screening period, 30 recorded values during the screening period (6 months up to and including day 0), 40 recorded values during the 12-month treatment period, and 8 recorded values at month 18 (6 months after the discontinuation of treatment). Red lines represent the median values according to time period. The dashed line indicates the mean interferon score of controls plus 2 SD; values above this line (i.e.,  $>2.47$ ) are considered to be abnormal. Panel B shows a comparison of interferon scores in blood in the 6 months before treatment, during the treatment period, and after treatment (at month 18), performed with the use of a nonlinear mixed-effects model. Panel C shows the interferon- $\alpha$  protein level in blood, Panel D the interferon score in blood, Panel E the interferon- $\alpha$  protein level in cerebrospinal fluid (CSF), and Panel F interferon activity in CSF at the indicated time points. Two patients (Patient 5 and Patient 10) did not undergo lumbar puncture. In two other patients (Patient 3 and Patient 4), an insufficient CSF sample was available at day 0, month 12, or both for assessment of the interferon- $\alpha$  protein level, interferon activity, or both. The correlation coefficient between the interferon- $\alpha$  protein level in blood and the interferon score in blood was 0.704 ( $P < 0.001$ ).

to pretreatment levels 6 months after discontinuation of therapy (Figs. S5 and S6 in the Supplementary Appendix). There was an increase in cerebral blood flow during the treatment period in 3 of 5 patients with data that could be interpreted (Fig. S8 and Table S7 in the Supplementary Appendix).

These results support the hypothesis that HIV-1 reverse-transcriptase therapy can reduce interferon signaling in patients with the Aicardi-Goutières syndrome by inhibition of reverse tran-

scription of endogenous retroelements. Changes in interferon signaling and cerebral blood flow suggest that treatment could have clinical value, perhaps in combination with other therapies (e.g., inhibitors of Janus kinase 1 and 2).<sup>4</sup> The open-label design of the study and small sample require that a larger group of patients be evaluated in a controlled clinical trial.

Gillian I. Rice, Ph.D.  
University of Manchester  
Manchester, United Kingdom

Candice Meyzer, M.D.

Naïm Bouazza, Ph.D.

Assistance Publique–Hôpitaux de Paris  
Paris, France

Marie Hully, M.D.

Nathalie Boddaert, M.D.

Hôpital Necker–Enfants Malades  
Paris, France

Michaela Semeraro, M.D., Ph.D.

Université Paris Descartes  
Paris, France

Leo A.H. Zeeff, Ph.D.

University of Manchester  
Manchester, United Kingdom

Flore Rozenberg, M.D., Ph.D.

Université Paris Descartes  
Paris, France

Vincent Bondet, Ph.D.

Darragh Duffy, Ph.D.

Alba Llibre, Ph.D.

Institut Pasteur  
Paris, France

Jinmi Baek, M.Sc.

Mame N. Sambe, B.Sc.

Elodie Henry, M.Sc.

Valerie Jolaine, M.Sc.

Assistance Publique–Hôpitaux de Paris  
Paris, France

Christine Barnerias, M.D.

Hôpital Necker–Enfants Malades  
Paris, France

Magalie Barth, M.D.

Centre Hospitalier Universitaire Angers  
Angers, France

Alexandre Belot, M.D., Ph.D.

University of Lyon  
Lyon, France

Claude Cances, M.D.

Centre Hospitalier Universitaire Toulouse  
Toulouse, France

François-Guillaume Debray, M.D., Ph.D.

University of Liège  
Liège, Belgium

Diane Doummar, M.D.

Hôpital Armand Trousseau  
Paris, France

Marie-Louise Frémond, M.D.

Naoki Kitabayashi, B.Sc.

Alice Lepelley, Ph.D.

Institut Imagine  
Paris, France

Virginie Levrat, M.D.

Centre Hospitalier Annecy Genevois  
Pringy, France

Isabelle Melki, M.D., Ph.D.

Hôpital Robert Debré  
Paris, France

Pierre Meyer, M.D.

Université de Montpellier  
Montpellier, France

Marie-Christine Nougues, M.D.

Hôpital Armand Trousseau  
Paris, France

Florence Renaldo, M.D.

Hôpital Robert Debré  
Paris, France

Mathieu P. Rodero, Ph.D.

Institut Imagine  
Paris, France

Diana Rodriguez, M.D., Ph.D.

Sorbonne Université  
Paris, France

Agathe Roubertie, M.D., Ph.D.

Centre Hospitalier Universitaire de Montpellier  
Montpellier, France

Luis Seabra, M.Sc.

Institut Imagine  
Paris, France

Carolina Uggenti, Ph.D.

University of Edinburgh  
Edinburgh, United Kingdom

Hendy Abdoul, M.D., Ph.D.

Jean-Marc Treluyer, M.D.

Assistance Publique–Hôpitaux de Paris  
Paris, France

Isabelle Desguerre, M.D.

Stéphane Blanche, M.D.

Hôpital Necker–Enfants Malades  
Paris, France

Yanick J. Crow, M.D., Ph.D.

University of Edinburgh  
Edinburgh, United Kingdom  
yanickcrow@mac.com

Supported by grants from the European Leukodystrophy Association (ELA 2012-00811), the European Research Council (GA 309449 and 786142-E-T1IFNs), ERA-NET Neuron (MR/M501803/1), and the French National Research Agency (ANR-10-IAHU-01 and CE17001002). Medications were provided by GlaxoSmithKline and Viiv Healthcare.

Disclosure forms provided by the authors are available with the full text of this letter at [NEJM.org](http://NEJM.org).

1. Crow YJ, Manel N. Aicardi-Goutières syndrome and the type I interferonopathies. *Nat Rev Immunol* 2015;15:429-40.
2. Volkman HE, Stetson DB. The enemy within: endogenous retroelements and autoimmune disease. *Nat Immunol* 2014;15:415-22.
3. Rice GI, Forte GM, Szykiewicz M, et al. Assessment of interferon-related biomarkers in Aicardi-Goutières syndrome associated with mutations in TREX1, RNASEH2A, RNASEH2B, RNASEH2C, SAMHD1, and ADAR: a case-control study. *Lancet Neurol* 2013;12:1159-69.
4. Kothur K, Bandodkar S, Chu S, et al. An open-label trial of JAK 1/2 blockade in progressive *IFIH1*-associated neuroinflammation. *Neurology* 2018;90:289-91.

DOI: 10.1056/NEJMc1810983