

# Revisiting the one leptoquark solution to the $R(D^{(*)})$ anomalies and its phenomenological implications

Xin-Qiang Li, Ya-Dong Yang and Xin Zhang

*Institute of Particle Physics and Key Laboratory of Quark and Lepton Physics (MOE),  
 Central China Normal University,  
 Wuhan, Hubei 430079, China*

*E-mail:* [xqli@mail.ccnu.edu.cn](mailto:xqli@mail.ccnu.edu.cn), [yangyd@mail.ccnu.edu.cn](mailto:yangyd@mail.ccnu.edu.cn),  
[zhangxin027@mails.ccnu.edu.cn](mailto:zhangxin027@mails.ccnu.edu.cn)

**ABSTRACT:** It has been shown recently that the anomalies observed in  $\bar{B} \rightarrow D^{(*)}\tau\bar{\nu}_\tau$  and  $\bar{B} \rightarrow \bar{K}\ell^+\ell^-$  decays could be resolved with just one scalar leptoquark. Fitting to the current data on  $R(D^{(*)})$  along with acceptable  $q^2$  distributions in  $\bar{B} \rightarrow D^{(*)}\tau\bar{\nu}_\tau$  decays, four best-fit solutions for the operator coefficients have been found. In this paper, we explore the possibilities of how to discriminate these four solutions. Firstly, we find that two of them are already excluded by the decay  $B_c^- \rightarrow \tau^-\bar{\nu}_\tau$ , because the predicted decay widths have already overshoot the total width  $\Gamma_{B_c}$ . It is then found that the remaining two solutions result in two effective Hamiltonians governing  $b \rightarrow c\tau\bar{\nu}_\tau$  transition, which differ by a sign and enhance the absolute value of the coefficient of  $\bar{c}_L\gamma_\mu b_L \bar{\tau}_L\gamma^\mu\nu_{\tau L}$  operator by about 12%. However, they give nearly the same predictions as in the SM for the  $D^*$  and  $\tau$  longitudinal polarizations as well as the lepton forward-backward asymmetries in  $\bar{B} \rightarrow D^{(*)}\tau\bar{\nu}_\tau$  decays. For the other observables like  $\mathcal{B}(B_c^- \rightarrow \tau^-\bar{\nu}_\tau)$ ,  $\mathcal{B}(B_c^- \rightarrow \gamma\tau^-\bar{\nu}_\tau)$ ,  $R_{D^{(*)}}(q^2)$ ,  $d\mathcal{B}(\bar{B} \rightarrow D^{(*)}\tau\bar{\nu}_\tau)/dq^2$  and  $\mathcal{B}(\bar{B} \rightarrow X_c\tau\bar{\nu}_\tau)$ , on the other hand, the two solutions give sizable enhancements relative to the SM predictions. With measurement of  $B_c^- \rightarrow \tau^-\bar{\nu}_\tau$  at LHCb and refined measurements of observables in  $\bar{B} \rightarrow D^{(*)}\tau\bar{\nu}_\tau$  at both LHCb and Belle-II, such a specific NP scenario could be further deciphered.

**KEYWORDS:** Beyond Standard Model, Heavy Quark Physics

**ARXIV EPRINT:** [1605.09308](https://arxiv.org/abs/1605.09308)

---

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>The one scalar leptoquark scenario</b>	<b>3</b>
<b>3</b>	<b>The effects of scalar leptoquark in <math>B_c</math> and <math>B</math> decays</b>	<b>5</b>
3.1	Purely leptonic decay $B_c^- \rightarrow \tau^- \bar{\nu}_\tau$	5
3.2	Comparison between the solutions $P_A$ and $P_C$	6
<b>4</b>	<b>Effects of solutions <math>P_A</math> and <math>P_C</math> in <math>B_c^- \rightarrow \gamma \tau^- \bar{\nu}_\tau</math>, <math>\bar{B} \rightarrow D^{(*)} \tau \bar{\nu}_\tau</math> and <math>B \rightarrow X_c \tau \bar{\nu}_\tau</math> decays</b>	<b>6</b>
4.1	$B_c^- \rightarrow \gamma \tau^- \bar{\nu}_\tau$	6
4.2	$\bar{B} \rightarrow D^{(*)} \tau \bar{\nu}_\tau$	8
4.3	$B \rightarrow X_c \tau^- \bar{\nu}_\tau$	11
<b>5</b>	<b>Conclusion</b>	<b>12</b>
<b>A</b>	<b>Analytic formulae of <math>B_c^- \rightarrow \gamma \tau^- \bar{\nu}_\tau</math>, <math>\bar{B} \rightarrow D^{(*)} \tau \bar{\nu}_\tau</math> and <math>B \rightarrow X_c \tau \bar{\nu}_\tau</math></b>	<b>13</b>
A.1	The radiative leptonic decay $B_c^- \rightarrow \gamma \tau^- \bar{\nu}_\tau$	13
A.2	Exclusive semi-leptonic decays $\bar{B} \rightarrow D^{(*)} \tau \bar{\nu}_\tau$	14
A.3	Inclusive semi-leptonic decay $B \rightarrow X_c \tau \bar{\nu}_\tau$	17
<b>B</b>	<b>Hadronic amplitudes in <math>\bar{B} \rightarrow D^{(*)} \tau \bar{\nu}_\tau</math> decays</b>	<b>17</b>

---

## 1 Introduction

With the discovery of heavy quark spin-flavor symmetry and the formulation of heavy quark effective theory (HQET) [1–5], it has become clear that the physical observables in semi-leptonic  $\bar{B} \rightarrow D^{(*)} \ell \bar{\nu}_\ell$  could be rather reliably predicted within the Standard Model (SM), especially at the zero recoil point, allowing therefore a reliable determination of the Cabibbo-Kobayashi-Maskawa (CKM) element  $V_{cb}$ . It is also believed that the effect of New Physics (NP) beyond the SM should be tiny since these decays are induced by the tree-level charged current.

However, the BaBar [6, 7], Belle [8, 9] and LHCb [10] collaborations have recently observed anomalies in the ratios

$$R(D^{(*)}) = \frac{\mathcal{B}(\bar{B} \rightarrow D^{(*)} \tau \bar{\nu}_\tau)}{\mathcal{B}(\bar{B} \rightarrow D^{(*)} \ell \bar{\nu}_\ell)}, \quad \ell = e/\mu. \quad (1.1)$$

The Heavy Flavor Average Group (HFAG) gives the average values [11]

$$\begin{aligned} R(D)_{\text{avg}} &= 0.397 \pm 0.040 \pm 0.028, \\ R(D^*)_{\text{avg}} &= 0.316 \pm 0.016 \pm 0.010, \end{aligned} \tag{1.2}$$

which exceed the SM predictions [12, 13]

$$\begin{aligned} R(D)_{\text{SM}} &= 0.300 \pm 0.008, \\ R(D^*)_{\text{SM}} &= 0.252 \pm 0.003, \end{aligned} \tag{1.3}$$

by  $1.9\sigma$  and  $3.3\sigma$ , respectively. Especially when the  $R(D)$ - $R(D^*)$  correlation of  $-0.21$  is taken into account, the tension with the SM predictions would be at  $4.0\sigma$  level [11]. Theoretically,  $R(D)$  and  $R(D^*)$  can be rather reliably calculated, because they are independent of the CKM element  $|V_{cb}|$  and, to a large extent, of the  $B \rightarrow D^{(*)}$  transition form factors.

The above anomalies have been investigated extensively both within model-independent frameworks [14–35], as well as in some specific NP models where the  $b \rightarrow c\tau\bar{\nu}_\tau$  transition is mediated by leptoquarks [14, 15, 36–44], charged Higgses [14, 45–55], charged vector bosons [14, 56–58], and sparticles [59–61]. It is also interesting to point out that, besides the branching ratios, the measured differential distributions  $d\Gamma(\bar{B} \rightarrow D^{(*)}\tau\bar{\nu})/dq^2$  by BaBar [7] and Belle [8, 9] provide very complementary information to distinguish NP from the SM as well as different NP models from each other [14, 39, 45].

With both the ratios  $R(D^{(*)})$  and the  $q^2$  spectra taken into account, Freytsis, Ligeti and Ruderman have identified viable models with leptoquark mediators, which are consistent with minimal flavor violation and could provide good fits to the current data; especially, four best-fit solutions are found for the operator coefficients induced by scalar leptoquarks [14]. With this observation, Bauer and Neubert have recently proposed a very simple NP model by extending the SM with a single TeV-scale scalar leptoquark  $\phi$  transforming as  $(\mathbf{3}, \mathbf{1}, -\frac{1}{3})$  under the SM gauge group, and shown that the anomalies observed in  $\bar{B} \rightarrow D^{(*)}\tau\bar{\nu}$ ,  $\bar{B} \rightarrow \bar{K}\ell^+\ell^-$  [62], as well as the anomalous magnetic moment of muon [63] can be explained in a natural way, while constraints from other precision measurements in the flavor sector are also satisfied without fine-tuning [40].

To further test such an interesting scenario, in this paper, we shall explore in detail the effect of the scalar leptoquark on the purely leptonic  $B_c^- \rightarrow \tau^-\bar{\nu}_\tau$ , the radiative leptonic  $B_c^- \rightarrow \gamma\tau^-\bar{\nu}_\tau$ , the exclusive semi-leptonic  $\bar{B} \rightarrow D^{(*)}\tau\bar{\nu}_\tau$ , and the inclusive semi-leptonic  $B \rightarrow X_c\tau\bar{\nu}_\tau$  decays. It is found that two of the four best-fit solutions obtained in ref. [14] are already excluded by the decay  $B_c^- \rightarrow \tau^-\bar{\nu}_\tau$ , because the predicted decay widths have already overshoot the total width  $\Gamma_{B_c}$ . The remaining two solutions result in two effective Hamiltonians that differ by a sign, but give almost the same predictions as in the SM for the  $D^*$  and  $\tau$  longitudinal polarizations as well as the lepton forward-backward asymmetries in  $\bar{B} \rightarrow D^{(*)}\tau\bar{\nu}_\tau$  decays. For the observables  $\mathcal{B}(B_c^- \rightarrow \tau^-\bar{\nu}_\tau)$ ,  $\mathcal{B}(B_c^- \rightarrow \gamma\tau^-\bar{\nu}_\tau)$ ,  $R_{D^{(*)}}(q^2)$ ,  $d\mathcal{B}(\bar{B} \rightarrow D^{(*)}\tau\bar{\nu}_\tau)/dq^2$  and  $\mathcal{B}(\bar{B} \rightarrow X_c\tau\bar{\nu}_\tau)$ , on the other hand, the two solutions give sizable enhancements relative to the SM predictions. With measurement of  $B_c^- \rightarrow \tau^-\bar{\nu}_\tau$  at LHCb and refined measurements of observables in  $\bar{B} \rightarrow D^{(*)}\tau\bar{\nu}_\tau$  at both LHCb and Belle-II, such a specific NP scenario could be further deciphered.

This paper is organized as follows: in section 2, we recapitulate the scenario with just one scalar leptoquark introduced in ref. [40]. In section 3, we consider the purely leptonic  $B_c^- \rightarrow \tau^- \bar{\nu}_\tau$  decay, from which two of the four best-fit solutions for the operator coefficients are found to be already excluded. The effects of the remaining two solutions on  $B_c^- \rightarrow \gamma \tau^- \bar{\nu}_\tau$ ,  $\bar{B} \rightarrow D^{(*)} \tau \bar{\nu}_\tau$  and  $B \rightarrow X_c \tau \bar{\nu}_\tau$  decays are then investigated in section 4. Our conclusions are finally made in section 5. Appendices A and B contain the formulae relevant to these decays.

## 2 The one scalar leptoquark scenario

In this section, we recapitulate the model proposed very recently by Bauer and Neubert [40], where a single TeV-scale leptoquark  $\phi$  is added to the SM to address the aforementioned anomalies in flavor physics. The new scalar  $\phi$  transforms as  $(\mathbf{3}, \mathbf{1}, -\frac{1}{3})$  under the SM gauge group, and its couplings to fermions are described by the Lagrangian [38, 40]

$$\mathcal{L}_{\text{int}}^\phi = \bar{Q}_L^c \boldsymbol{\lambda}^L i\tau_2 L\phi^* + \bar{u}_R^c \boldsymbol{\lambda}^R l_R\phi^* + \text{h.c.}, \quad (2.1)$$

where  $\boldsymbol{\lambda}^{L,R}$  are the Yukawa coupling matrices in flavor space,  $Q_L$ ,  $L$  denote the left-handed quark and lepton doublet,  $u_R$ ,  $l_R$  the right-handed up-type quark and lepton singlet respectively, and  $\psi^c = C\bar{\psi}^T$ ,  $\bar{\psi}^c = \psi^T C$  ( $C = i\gamma^2\gamma^0$ ) are the charge-conjugated spinors. Rotating the Lagrangian from the weak to the mass basis for quarks and charged leptons, the interaction terms take the form

$$\mathcal{L}_{\text{int}}^\phi = \bar{u}_L^c \boldsymbol{\lambda}_{ul}^L l_L\phi^* - \bar{d}_L^c \boldsymbol{\lambda}_{d\nu}^L \nu_L\phi^* + \bar{u}_R^c \boldsymbol{\lambda}_{ul}^R l_R\phi^* + \text{h.c.}, \quad (2.2)$$

where  $\boldsymbol{\lambda}_{ul}^L$ ,  $\boldsymbol{\lambda}_{d\nu}^L$  and  $\boldsymbol{\lambda}_{ul}^R$  are now the coupling matrices in mass basis, and describe the strength of  $\phi$  interactions with fermions.

Writing down the tree-level  $\phi$ -exchange amplitude for the process  $b \rightarrow c\tau\bar{\nu}_\tau$  in the leading order in  $k^2/M_\phi^2$  expansion, where  $k \sim \mathcal{O}(m_b)$  is the momentum flowing through the  $\phi$  propagator and  $M_\phi$  the leptoquark mass, and then performing the Fierz transformation of the resulting four-fermion operators, one can get the effective Hamiltonian

$$\begin{aligned} \mathcal{H}_{\text{eff}}^\phi = & -\frac{1}{2M_\phi^2} \left[ \lambda_{b\nu_\tau}^L \lambda_{c\tau}^{L*} \bar{b}^c \gamma_\mu P_R c^c \bar{\tau} \gamma^\mu P_L \nu_\tau \right. \\ & \left. + \lambda_{b\nu_\tau}^L \lambda_{c\tau}^{R*} \left( \bar{b}^c P_L c^c \bar{\tau} P_L \nu_\tau + \frac{1}{4} \bar{b}^c \sigma_{\mu\nu} P_L c^c \bar{\tau} \sigma^{\mu\nu} P_L \nu_\tau \right) \right], \end{aligned} \quad (2.3)$$

where  $P_{L,R} = (1 \mp \gamma_5)/2$  are the chirality projectors. Using the definitions  $\psi^c = C\bar{\psi}^T$ ,  $\bar{\psi}^c = \psi^T C$ , and the relations  $C\gamma_\mu = -\gamma_\mu^T C$ ,  $C\gamma_5 = \gamma_5^T C$ , one can easily arrive at the equations

$$\bar{b}^c \gamma_\mu P_R c^c = -\bar{c} \gamma_\mu P_L b, \quad (2.4)$$

$$\bar{b}^c P_L c^c = \bar{c} P_L b, \quad (2.5)$$

$$\bar{b}^c \sigma_{\mu\nu} P_L c^c = -\bar{c} \sigma_{\mu\nu} P_L b. \quad (2.6)$$

Plugging the above three equations into eq. (2.3) and including also the SM contribution, one obtains then the total effective Hamiltonian governing the  $b \rightarrow c\tau\bar{\nu}_\tau$  transition

$$\mathcal{H}_{\text{eff}} = \frac{4G_F}{\sqrt{2}}V_{cb} \left[ C_V(M_\phi) \bar{c}\gamma_\mu P_L b \bar{\tau}\gamma^\mu P_L \nu_\tau + C_S(M_\phi) \bar{c}P_L b \bar{\tau}P_L \nu_\tau - \frac{1}{4}C_T(M_\phi) \bar{c}\sigma_{\mu\nu} P_L b \bar{\tau}\sigma^{\mu\nu} P_L \nu_\tau \right], \quad (2.7)$$

where  $C_V, C_S, C_T$  are the Wilson coefficients of the corresponding operators at the matching scale  $\mu = M_\phi$ , and are given explicitly as

$$C_V(M_\phi) = 1 + \frac{\lambda_{b\nu_\tau}^L \lambda_{c\tau}^{L*}}{4\sqrt{2}G_F V_{cb} M_\phi^2},$$

$$C_S(M_\phi) = C_T(M_\phi) = -\frac{\lambda_{b\nu_\tau}^L \lambda_{c\tau}^{R*}}{4\sqrt{2}G_F V_{cb} M_\phi^2}. \quad (2.8)$$

In order to re-sum potentially large logarithmic effects and to make predictions for physical observables, the Wilson coefficients given by eq. (2.8) should be run down to the characteristic scale of the processes we are interested in, i.e.,  $\mu_b \sim m_b$ . While the vector current is conserved and needs not be renormalized, the evolutions at the leading logarithmic approximation of the scalar  $C_S$  and tensor  $C_T$  coefficients are given, respectively, by [64]

$$C_S(\mu_b) = \left[ \frac{\alpha_s(m_t)}{\alpha_s(\mu_b)} \right]^{\frac{\gamma_S}{2\beta_0^{(5)}}} \left[ \frac{\alpha_s(M_\phi)}{\alpha_s(m_t)} \right]^{\frac{\gamma_S}{2\beta_0^{(6)}}} C_S(M_\phi),$$

$$C_T(\mu_b) = \left[ \frac{\alpha_s(m_t)}{\alpha_s(\mu_b)} \right]^{\frac{\gamma_T}{2\beta_0^{(5)}}} \left[ \frac{\alpha_s(M_\phi)}{\alpha_s(m_t)} \right]^{\frac{\gamma_T}{2\beta_0^{(6)}}} C_T(M_\phi), \quad (2.9)$$

where  $\gamma_S = -8$  [65] and  $\gamma_T = 8/3$  [66] are the LO anomalous dimensions of QCD scalar and tensor currents respectively, and  $\beta_0^{(f)} = 11 - 2n_f/3$  the LO beta function coefficient, with  $n_f$  being the number of active quark flavors.

As detailed in ref. [14], such a scalar leptoquark scenario introduced above could provide good explanations to the  $R(D)$  and  $R(D^*)$  anomalies along with acceptable  $q^2$  spectra. Taking  $M_\phi = 1$  TeV as a benchmark and performing a two-dimensional  $\chi^2$  fit, they found four best-fit solutions with  $\chi_{\text{min}}^2 < 5$  for the operator coefficients, which are listed below and denoted, respectively, as  $P_A, P_B, P_C, P_D$ :

$$(\lambda_{b\nu_\tau}^L \lambda_{c\tau}^{L*}, \lambda_{b\nu_\tau}^L \lambda_{c\tau}^{R*}) = (C_{S_R}^{\prime\prime}, C_{S_L}^{\prime\prime}) = \begin{cases} (0.35, -0.03), P_A \\ (0.96, 2.41), P_B \\ (-5.74, 0.03), P_C \\ (-6.34, -2.39), P_D \end{cases}. \quad (2.10)$$

It is noticed that only the solution  $P_A$  is adopted by Bauer and Neubert [40], arguing that the other three require significantly larger couplings. It would be worth investigating whether the four best-fit solutions could be discriminated from each other using the processes mediated by the same effective operators given by eq. (2.7). To this end, in

addition to  $\bar{B} \rightarrow D^{(*)}\tau\bar{\nu}_\tau$ , we shall examine their effects on  $B_c^- \rightarrow \tau^-\bar{\nu}_\tau$ ,  $B_c^- \rightarrow \gamma\tau^-\bar{\nu}_\tau$  and  $B \rightarrow X_c\tau\bar{\nu}_\tau$  decays.

As a final comment, it should be noted that the interaction Lagrangian eq. (2.2) also gives rise to tree-level neutral quark and lepton currents; after integrating out the scalar leptoquark and performing the Fierz transformation, one encounters the operators  $(\bar{u}_i u_j)(\ell^+ \ell^-)$  and  $(\bar{d}_i d_j)(\bar{\nu} \nu)$ , the Wilson coefficients of which can be constrained, for example, by the rare decays  $D^0 \rightarrow \mu^+ \mu^-$  and  $D^+ \rightarrow \pi^+ \mu^+ \mu^-$ , as well as  $B \rightarrow X_s \nu \bar{\nu}$ ,  $B \rightarrow K^{(*)} \nu \bar{\nu}$  and  $K \rightarrow \pi \nu \bar{\nu}$ , respectively. For more information about the low-energy constraints on the model, we refer the reader to refs. [14, 38, 40].

### 3 The effects of scalar leptoquark in $B_c$ and $B$ decays

In this section, we explore the effects of the scalar leptoquark  $\phi$  with the four best-fit solutions in the  $B_c$  and  $B$ -meson decays.

#### 3.1 Purely leptonic decay $B_c^- \rightarrow \tau^-\bar{\nu}_\tau$

Firstly, we investigate the effect of  $\phi$  on the purely leptonic decay  $B_c^- \rightarrow \tau^-\bar{\nu}_\tau$ , the decay amplitude of which, including both the SM and NP contributions, can be written as

$$\mathcal{A}(B_c^- \rightarrow \tau^-\bar{\nu}_\tau) = i \frac{G_F V_{cb}}{\sqrt{2}} \left[ C_V \langle 0 | \bar{c} \gamma_\mu \gamma_5 b | B_c \rangle \bar{\tau} \gamma^\mu (1 - \gamma_5) \nu_\tau + C_S \langle 0 | \bar{c} \gamma_5 b | B_c \rangle \bar{\tau} (1 - \gamma_5) \nu_\tau \right]. \quad (3.1)$$

Together with the definition of the  $B_c$ -meson decay constant  $f_{B_c}$ ,  $\langle 0 | \bar{c} \gamma^\mu \gamma_5 b | B_c(p) \rangle = i f_{B_c} p^\mu$ , and using the equation of motion, one can express the matrix element of pseudoscalar current as  $\langle 0 | \bar{c} \gamma_5 b | B_c(p) \rangle = -\frac{i f_{B_c} m_{B_c}^2}{m_b(\mu_b) + m_c(\mu_b)}$ . The decay width for this process then reads

$$\Gamma(B_c^- \rightarrow \tau^-\bar{\nu}_\tau) = \frac{G_F^2}{8\pi} |V_{cb}|^2 f_{B_c}^2 m_{B_c}^3 \frac{m_\tau^2}{m_{B_c}^2} \left( 1 - \frac{m_\tau^2}{m_{B_c}^2} \right)^2 \left| C_V - C_S \frac{m_{B_c}^2}{m_\tau [m_b(\mu_b) + m_c(\mu_b)]} \right|^2, \quad (3.2)$$

where  $C_V$  and  $C_S$  are the Wilson coefficients of (axial)vector and (pseudo)scalar operators, and  $m_b$  and  $m_c$  the current quark masses, all being given at the scale  $\mu_b = m_b$ .

With the input parameters collected in table 1 and  $f_{B_c} = 0.434$  GeV [67], we get

$$\Gamma(B_c^- \rightarrow \tau^-\bar{\nu}_\tau) = \begin{cases} 2.22 \times 10^{-2} \Gamma_{B_c}, & \text{SM} \\ 2.45 \times 10^{-2} \Gamma_{B_c}, & P_A \\ 1.33 \Gamma_{B_c}, & P_B \\ 2.39 \times 10^{-2} \Gamma_{B_c}, & P_C \\ 1.31 \Gamma_{B_c}, & P_D \end{cases}, \quad (3.3)$$

which are normalized to the total decay width  $\Gamma_{B_c} = 1/\tau_{B_c} = 1/0.507$  ps<sup>-1</sup> [68]. The results labelled by  $P_{A,B,C,D}$  are obtained using the four best-fit solutions given by eq. (2.10) and with  $M_\phi = 1$  TeV. Clearly, one can see that two of the four solutions,  $P_B$  and  $P_D$ , are already excluded by the decay  $B_c^- \rightarrow \tau^-\bar{\nu}_\tau$ , because the predicted decay widths have already overshoot the total width  $\Gamma_{B_c}$ . Therefore, in the following, we need only consider the remaining two best-fit solutions  $P_A$  and  $P_C$ .

### 3.2 Comparison between the solutions $P_A$ and $P_C$

Before going to detail their effects on the other decays, we give firstly a comparison between the remaining two best-fit solutions  $P_A$  and  $P_C$ . Plugging into the effective Hamiltonian eq. (2.7) the fitted values of the effective couplings in  $P_A$  and  $P_C$  solutions, we get

$$\mathcal{H}_{\text{fit}} = \frac{4G_F}{\sqrt{2}} V_{cb} \left\{ \left[ 1 + \begin{pmatrix} 0.129 \text{ for } P_A \\ -2.117 \text{ for } P_C \end{pmatrix} \right] \bar{c}\gamma_\mu P_L b \bar{\tau}\gamma^\mu P_L \nu_\tau + \begin{pmatrix} 0.018 \text{ for } P_A \\ -0.018 \text{ for } P_C \end{pmatrix} \bar{c}P_L b \bar{\tau}P_L \nu_\tau \right. \\ \left. + \begin{pmatrix} -0.002 \text{ for } P_A \\ 0.002 \text{ for } P_C \end{pmatrix} \bar{c}\sigma_{\mu\nu} P_L b \bar{\tau}\sigma^{\mu\nu} P_L \nu_\tau \right\}, \quad (3.4)$$

which have been run down from the scale  $\mu = M_\phi = 1$  TeV to the scale  $\mu = m_b = 4.18$  GeV.

It is observed that the coefficients of the operator  $\bar{c}\gamma_\mu P_L b \bar{\tau}\gamma^\mu P_L \nu_\tau$ ,

$$C_V^{\text{fit}} = \begin{cases} 1.129, & \text{for } P_A \\ -1.117, & \text{for } P_C \end{cases}, \quad (3.5)$$

have nearly the same absolute values, both enhancing the SM result by  $\sim 12\%$ , but the sign of solution  $P_C$  is flipped relative to the SM. Furthermore, the  $\sim 1.2\%$  difference in  $|C_V^{\text{fit}}|$  would be too small to be discriminated from each other phenomenologically. For the other two operators, on the other hand, the two solutions  $P_A$  and  $P_C$  result in the same (tiny) values of coefficients but with opposite signs.

## 4 Effects of solutions $P_A$ and $P_C$ in $B_c^- \rightarrow \gamma\tau^-\bar{\nu}_\tau$ , $\bar{B} \rightarrow D^{(*)}\tau\bar{\nu}_\tau$ and $B \rightarrow X_c\tau\bar{\nu}_\tau$ decays

In this section, we study the effects of solutions  $P_A$  and  $P_C$  in  $B_c^- \rightarrow \gamma\tau^-\bar{\nu}_\tau$ ,  $\bar{B} \rightarrow D^{(*)}\tau\bar{\nu}_\tau$  and  $B \rightarrow X_c\tau\bar{\nu}_\tau$  decays. All the relevant analytic formulae for these decays are collected in appendices A and B. In table 1, we list the input parameters used in our numerical analyses, with all the other ones taken from the Particle Data Group [68].

We give our predictions for the observables in  $B_c^- \rightarrow \gamma\tau^-\bar{\nu}_\tau$ ,  $\bar{B} \rightarrow D^{(*)}\tau\bar{\nu}_\tau$  and  $B \rightarrow X_c\tau\bar{\nu}_\tau$  decays both within the SM as well as in the one scalar leptoquark scenario with the operator parameters taking the values labelled by  $P_A$  and  $P_C$  in eq. (2.10). The theoretical uncertainty for an observable is evaluated by varying each input parameter within its corresponding allowed ranges and then adding the individual uncertainties in quadrature [75–77].

### 4.1 $B_c^- \rightarrow \gamma\tau^-\bar{\nu}_\tau$

Firstly, we present our predictions for the branching ratio by just giving their center values, which are shown in table 2. In our calculation, we adopt the cut condition for the photon energy  $E_\gamma \geq 1$  GeV, and take the constituent quark mass values  $m_b = 4.8$  GeV,  $m_c = 1.5$  GeV [78, 79].

From table 2, one can see that the branching ratios are enhanced by about 27% in both the  $P_A$  and  $P_C$  cases, relative to our SM prediction, which is in agreement with the ones in the literatures [79–85].

Parameter	Value	Reference(s)
$\alpha_s(M_Z)$	$0.1185 \pm 0.0006$	[68]
$\alpha_e$	1/128	[68]
$m_t$	$(173.21 \pm 0.87)$ GeV	[68]
$m_b(m_b)$	$(4.18 \pm 0.03)$ GeV	[68]
$m_c(m_c)$	$(1.275 \pm 0.025)$ GeV	[68]
$\tau_{B^+}$	$(1.638 \pm 0.004)$ ps	[68]
$\tau_{B^0}$	$(1.520 \pm 0.004)$ ps	[68]
$ V_{cb} $	$(41.1 \pm 1.3) \times 10^{-3}$	[68]
$\rho_D^2$	$1.086 \pm 0.070$	[69–71]
$V_1(1)$	$0.908 \pm 0.017$	[72]
$\rho_{D^*}^2$	$1.207 \pm 0.026$	[73]
$h_{A_1}(1)$	$0.908 \pm 0.017$	[74]
$R_1(1)$	$1.406 \pm 0.033$	[73]
$R_2(1)$	$0.853 \pm 0.020$	[73]

**Table 1.** Input parameters used in our numerical analyses.


Observable	SM	$P_A$	$P_C$
$\mathcal{B}(B_c^- \rightarrow \gamma \tau^- \bar{\nu}_\tau) \times 10^5$	2.36	3.01	2.94

**Table 2.** Branching ratios of  $B_c^- \rightarrow \gamma \tau^- \bar{\nu}_\tau$  both within the SM and in the  $P_A$  and  $P_C$  cases.

In figure 1, we present the dependence of the differential branching ratios on the photon energy  $E_\gamma$ . One can see that the effects of solutions  $P_A$  and  $P_C$  are both significant in the region  $1 \text{ GeV} < E_\gamma < 1.5 \text{ GeV}$ , but become tiny near the end point  $E_\gamma = (m_{B_c}^2 - m_\tau^2)/(2m_{B_c})$ . While being enhanced both in the  $P_A$  and  $P_C$  cases, the predicted differential branching ratios coincide almost with each other and are therefore indistinguishable.

It is well-known that, while the width for a purely leptonic decay of a charged pseudoscalar meson is helicity suppressed by  $m_\ell^2/m_{P^+}^2$  ( $m_{P^+}$  is the meson mass), the corre-





**Figure 1.** The dependence of the differential branching ratios on the photon energy  $E_\gamma$ .

sponding radiative leptonic decay relieves the helicity suppression, and enhances the decay rate, especially for  $\ell = e, \mu$ , at expense of much larger theoretical uncertainties [86–89]. For the  $B_c^- \rightarrow (\gamma)\tau^- \bar{\nu}_\tau$  decays, however, this is not the case; since  $B_c^- \rightarrow \tau^- \bar{\nu}_\tau$  does not suffer so much from the helicity suppression, the photons radiated from heavy quarks and heavy  $\tau$  do not enhance the decay rate, and the resulting extra electromagnetic coupling  $\alpha_e$  will suppress  $B_c^- \rightarrow \gamma\tau^- \bar{\nu}_\tau$  [79–85].

Together with the lattice QCD calculation of the decay constant  $f_{B_c}$  [67],  $\mathcal{B}(B_c^- \rightarrow \tau^- \bar{\nu}_\tau)$  could be reliably predicted. To test the one scalar leptoquark scenario, the purely leptonic decay  $B_c^- \rightarrow \tau^- \bar{\nu}_\tau$  is very powerful, especially if LHCb could measure the branching ratio  $\mathcal{B}(B_c^- \rightarrow \tau^- \bar{\nu}_\tau)$  with a precision of 5%, since the remaining two best-fit solutions  $P_A$  and  $P_C$  just enhance it by 10% and 8%, respectively, as shown in eq. (3.3). However, unlike the measurements of  $B_u^- \rightarrow \tau^- \bar{\nu}_\tau$  at BaBar and Belle operated at the  $\Upsilon(4S)$  resonance with  $B_u^\pm$  produced in pairs, it would be extremely difficult to measure  $B_c^- \rightarrow \tau^- \bar{\nu}_\tau$  at LHCb, because the presence of at least two neutrinos per  $B_c^-$  decay and the inability to impose kinematic constraints on the center-of-mass energy make background rejection incredibly challenging. The radiative mode would face even more challenges from vetoing photons from excited  $B_c^* \rightarrow B_c \gamma$  decays. While being very challenging, it is worthy for LHCb to make delicately experimental studies of these decays thanks to the high luminosity and the large  $B_c^-$  production cross-section at the LHC [90].


#### 4.2 $\bar{B} \rightarrow D^{(*)} \tau \bar{\nu}_\tau$

In this subsection, we present firstly in table 3 our predictions for the ratios  $R(D^{(*)})$  and the branching fractions  $\mathcal{B}(\bar{B} \rightarrow D^{(*)} \tau \bar{\nu}_\tau)$ , both within the SM and in the  $P_A$  and  $P_C$  cases. One can see from the table that the values of  $R(D^{(*)})$  in both the  $P_A$  and  $P_C$  cases coincide very well with the experimental data, as it should be.


We now analyze in turn the  $q^2$  distributions of the differential branching fractions (shown in figure 2), the ratios  $R_{D^{(*)}}(q^2)$  (figure 3), the polarizations of  $\tau$  (figure 4) and  $D^*$  (figure 5 (a)), as well as the lepton forward-backward asymmetry defined as the relative difference between the partial decay rates where the angle  $\theta$  between  $D^{(*)}$  and  $\tau$  three-momenta in the  $\tau\text{-}\bar{\nu}_\tau$  center-of-mass frame is greater or smaller than  $\pi/2$  (cf. eq. (A.9)) (figures 5 (b) and 5 (c)) in  $\bar{B} \rightarrow D^{(*)} \tau \bar{\nu}_\tau$  decays. For simplicity, we plot only

Observable	SM	$P_A$	$P_C$	Exp
$R(D)$	$0.298 \pm 0.009$	$0.388 \pm 0.012$	$0.380 \pm 0.011$	$0.397 \pm 0.049$ [11]
$R(D^*)$	$0.253 \pm 0.002$	$0.325 \pm 0.002$	$0.318 \pm 0.002$	$0.316 \pm 0.019$ [11]
$\mathcal{B}(\bar{B} \rightarrow D\tau\nu)$	$0.72 \pm 0.06$	$0.94 \pm 0.07$	$0.92 \pm 0.07$	$1.07 \pm 0.18$ [68]
$\mathcal{B}(\bar{B} \rightarrow D^*\tau\nu)$	$1.30 \pm 0.04$	$1.67 \pm 0.05$	$1.64 \pm 0.04$	$1.64 \pm 0.15$ [68]


**Table 3.** Theoretical and experimental values of  $R(D^{(*)})$  and  $\mathcal{B}(\bar{B} \rightarrow D^{(*)}\tau\nu_\tau)$  (in unit of  $10^{-2}$ ), both within the SM and in the  $P_A$  and  $P_C$  cases. The statistical and systematic uncertainties have been added in quadrature for the experimental data.




**Figure 2.** The  $q^2$  distributions of the differential branching fractions for  $\bar{B} \rightarrow D\tau\bar{\nu}_\tau$  (a) and for  $\bar{B} \rightarrow D^*\tau\bar{\nu}_\tau$  (b) decays.



**Figure 3.** The  $q^2$  distributions of the ratios  $R_D(q^2)$  (a) and  $R_{D^*}(q^2)$  (b).



**Figure 4.** The  $q^2$  distributions of the  $\tau$  longitudinal polarization in  $\bar{B} \rightarrow D\tau\bar{\nu}_\tau$  (a) and in  $\bar{B} \rightarrow D^*\tau\bar{\nu}_\tau$  (b) decays.




**Figure 5.** The  $q^2$  distributions of the  $D^*$  longitudinal polarization in  $\bar{B} \rightarrow D^*\tau\bar{\nu}_\tau$  (a), and that of the lepton forward-backward asymmetries in  $\bar{B} \rightarrow D\tau\bar{\nu}_\tau$  (b) and in  $\bar{B} \rightarrow D^*\tau\bar{\nu}_\tau$  (c) decays.

the central values of these observables at each  $q^2$  point. From these figures, we make the following observations:

- As shown in figure 2, the differential branching ratio  $d\mathcal{B}(\bar{B} \rightarrow D\tau\bar{\nu}_\tau)/dq^2$  is largely enhanced around  $q^2 \sim 7 \text{ GeV}^2$ , while  $d\mathcal{B}(\bar{B} \rightarrow D^*\tau\bar{\nu}_\tau)/dq^2$  around  $8 \text{ GeV}^2$ . Furthermore, both  $P_A$  and  $P_C$  predict similar  $q^2$  behaviors as in the SM for these two observables. As the measured differential distributions by BaBar [7] and Belle [8, 9] are still quite uncertain, it is currently unable to discriminate the NP from the SM predictions. More precise measurements of these observables by LHCb and Belle-II are, therefore, very necessary.
- From figure 3, one can see that the scalar leptoquark effects provide overall enhancements for both  $R_D(q^2)$  and  $R_{D^*}(q^2)$  in the whole kinematic region. However, the enhancement is small for  $R_D(q^2)$ , but quite large for  $R_{D^*}(q^2)$  in the large  $q^2$  region. This could be tested at Belle-II in the near future.
- As shown in figures 4 and 5, for the  $\tau$  and  $D^*$  longitudinal polarizations, as well as the lepton forward-backward asymmetries in these decays, results obtained in the  $P_A$

Observable	SM	$P_A$	$P_C$
$R(X_c)$	0.230	0.297	0.290
$\mathcal{B}(B^- \rightarrow X_c \tau^- \bar{\nu}_\tau)$	2.51%	3.24%	3.17%

**Table 4.** Predictions for  $R(X_c)$  and  $\mathcal{B}(B^- \rightarrow X_c \tau^- \bar{\nu}_\tau)$  in the SM and in the  $P_A$  and  $P_C$  cases.



**Figure 6.** The  $\hat{q}^2$  distributions of the ratio  $R_{X_c}(\hat{q}^2)$  (a) and the differential branching fraction  $d\mathcal{B}(B^- \rightarrow X_c \tau^- \bar{\nu}_\tau)/d\hat{q}^2$  (b), as well as the  $\tau$ -energy spectrum  $d\mathcal{B}(B^- \rightarrow X_c \tau^- \bar{\nu}_\tau)/dy$  (c).

and  $P_C$  cases coincide not only with each other, but also with the corresponding SM predictions. This is naively what should be expected, because the scalar leptoquark effects appear both in the numerator and in the denominator of these observables and are cancelled to a large extent, making these observables almost independent of their contributions.

### 4.3 $B \rightarrow X_c \tau^- \bar{\nu}_\tau$

Finally, we consider the inclusive semi-leptonic  $B$ -meson decays. Similar to the case in exclusive decays, we can also define a ratio  $R(X_c)$  for inclusive decay rates,

$$R(X_c) = \frac{\mathcal{B}(\bar{B} \rightarrow X_c \tau^- \bar{\nu}_\tau)}{\mathcal{B}(\bar{B} \rightarrow X_c \ell^- \bar{\nu}_\ell)}, \quad \ell = e/\mu, \quad (4.1)$$

which can be calculated precisely with an operator product expansion [4, 91–94]. With the most recent world average  $\mathcal{B}(B^- \rightarrow X_c e^- \bar{\nu}_e) = (10.92 \pm 0.16)\%$  [73, 95], one can then get the prediction for  $\mathcal{B}(B^- \rightarrow X_c \tau^- \bar{\nu}_\tau)$ , free of the large uncertainty due to the factor  $m_b^5$ . Here we consider neither the  $\mathcal{O}(\alpha_s)$  QCD nor the  $\mathcal{O}(\Lambda_{\text{QCD}}/m_b)$  power corrections, and take the heavy quark on-shell masses with  $m_b = 4.6$  GeV and  $m_c = 1.15$  GeV [96].

Our numerical results of the ratio  $R(X_c)$  and the branching fraction  $\mathcal{B}(B^- \rightarrow X_c \tau^- \bar{\nu}_\tau)$  are given in table 4. From the table, one can see that both  $R(X_c)$  and  $\mathcal{B}(B^- \rightarrow X_c \tau^- \bar{\nu}_\tau)$  are enhanced by the scalar leptoquark, and our SM value of  $R(X_c)$  is roughly consistent with the recent update within the 1S short-distance mass scheme [14, 91],  $R(X_c) = 0.223 \pm 0.004$ , obtained with both the  $\mathcal{O}(\Lambda_{\text{QCD}}^2/m_b^2)$  and the two-loop QCD corrections included [97].

In figure 6, we display the  $\hat{q}^2$  distributions of the ratio  $R_{X_c}(\hat{q}^2)$ , the differential branching fraction  $d\mathcal{B}(B^- \rightarrow X_c \tau^- \bar{\nu}_\tau)/d\hat{q}^2$ , as well as the differential spectrum of the  $\tau$  energy

$d\mathcal{B}(B^- \rightarrow X_c \tau \bar{\nu}_\tau)/dy$ , where  $\hat{q}^2 = q^2/m_b^2$  and  $y = 2E_\tau/m_b$ .<sup>1</sup> One can see that the  $\hat{q}^2$  distributions of these two observables are similar to that of  $R_{D^*}(q^2)$  and  $d\mathcal{B}(\bar{B} \rightarrow D^* \tau \bar{\nu}_\tau)/dq^2$ , respectively, since both  $R_{X_c}(\hat{q}^2)$  and  $d\mathcal{B}(B^- \rightarrow X_c \tau \bar{\nu}_\tau)/d\hat{q}^2$  are also enhanced by the scalar leptoquark contributions in the whole kinematic region, except for near the origin and the end point regions of  $\hat{q}^2$  for the latter. This is due to the fact that these observables have similar relations with respect to the operator coefficients  $C_V$ ,  $C_S$  and  $C_T$ , which can be seen from eqs. (A.13) and (A.16). It is also found that the  $\tau$ -energy spectrum shows a different behavior than the differential branching fraction and provides complementary information compared to the latter.

## 5 Conclusion

The anomalies observed in  $\bar{B} \rightarrow D^{(*)} \tau \bar{\nu}_\tau$  and  $\bar{B} \rightarrow \bar{K} \ell^+ \ell^-$  decays could be resolved with just one scalar leptoquark [40]. Fitting to the current experimental data on the ratios  $R(D^{(*)})$  and the  $q^2$  spectra of  $\bar{B} \rightarrow D^{(*)} \tau \bar{\nu}_\tau$ , four best-fit solutions denoted by  $P_A$ ,  $P_B$ ,  $P_C$  and  $P_D$  are obtained [14]. In this paper, we have explored the possibilities of how to discriminate these four solutions. Firstly, we have shown that two of them,  $P_B$  and  $P_D$ , are already excluded by the purely leptonic decay  $B_c^- \rightarrow \tau^- \bar{\nu}_\tau$ , because the predicted decay widths by  $P_B$  and  $P_D$  have already overshoot the total width  $\Gamma_{B_c}$ . The remaining two solutions  $P_A$  and  $P_C$  would enhance  $\mathcal{B}(B_c^- \rightarrow \tau^- \bar{\nu}_\tau)$  by 10% and 8%, respectively. Together with the lattice QCD calculation of the decay constant  $f_{B_c}$  [67],  $\mathcal{B}(B_c^- \rightarrow \tau^- \bar{\nu}_\tau)$  could be reliably predicted. Given the branching ratio measured to a precision of a few percent at the LHCb, one could then test the interesting one scalar leptoquark model.

By comparing the effects of  $P_A$  and  $P_C$  at the scale  $\mu = m_b$ , we find that the two solutions lead to overall different sign of  $\mathcal{H}_{\text{fit}}$ , but with just 1.2% difference in the coefficient  $C_V^{\text{fit}}$  of  $(V - A) \otimes (V - A)$  operator, which is too small to be discriminated from each other phenomenologically. Furthermore, in  $\mathcal{H}_{\text{fit}}$ , the coefficient  $|C_V^{\text{fit}}|$  is much larger than  $|C_S^{\text{fit}}|$  and  $|C_T^{\text{fit}}|$ , the coefficients of the new scalar and tensor operators, respectively.

Combining these observations and our numerical results, we may draw the following conclusions. The one scalar leptoquark scenario gives nearly the same predictions as in the SM for the  $D^*$  and  $\tau$  longitudinal polarizations and the lepton forward-backward asymmetries in  $\bar{B} \rightarrow D^{(*)} \tau \bar{\nu}_\tau$  decays. Although precision measurements of these observables would be very challenging at LHCb and/or Belle-II, any significant deviation from the SM predictions would lead to another model for the  $R(D^{(*)})$  anomalies. Otherwise, the one scalar leptoquark scenario would be viable and good. For the other observables like  $\mathcal{B}(B_c^- \rightarrow \tau^- \bar{\nu}_\tau)$ ,  $\mathcal{B}(B_c^- \rightarrow \gamma \tau^- \bar{\nu}_\tau)$ ,  $R_{D^{(*)}}(q^2)$ ,  $d\mathcal{B}(\bar{B} \rightarrow D^{(*)} \tau \bar{\nu}_\tau)/dq^2$  and  $\mathcal{B}(\bar{B} \rightarrow X_c \tau \bar{\nu}_\tau)$ , on the other hand, the model could generally give sizable enhancements relative to the SM predictions.

---

<sup>1</sup>It should be noted that the lowest-order (parton-level) prediction for the inclusive spectrum receives substantial corrections from nonperturbative  $\Lambda_{\text{QCD}}^2/m_b^2$  power corrections and shape-function convolutions in the large  $q^2$  (kinematic endpoint) region, which is also the part of the distribution where any reported data will likely be cleanest; for a recently detailed study, see ref. [91].

With future measurement of  $B_c^- \rightarrow \tau^- \bar{\nu}_\tau$  at LHCb and refined measurements of observables in  $\bar{B} \rightarrow D^{(*)} \tau \bar{\nu}_\tau$  decays at both LHCb and Belle-II, one could further decipher the various NP models that provide so far good explanations of the  $R(D^{(*)})$  anomalies.

Finally, we would like to point out that, due to the half-integer-spin of  $\Lambda_b$  and  $\Lambda_c$  baryons, the semi-leptonic  $\Lambda_b \rightarrow \Lambda_c \ell \bar{\nu}_\ell$  decays, which are mediated by the same quark-level transition as in  $\bar{B} \rightarrow D^{(*)} \ell \bar{\nu}_\ell$  decays, can provide additional polarization observables through angular decay distribution, such as the hadron-side asymmetries in the decay  $\Lambda_c^+ \rightarrow \Lambda^0 \pi^+$  and azimuthal correlations between the two final-state decay planes [98]. While the  $\Lambda_b$  baryons are not produced at an  $e^+e^-$  B-factory, they account for about 20% of the  $b$ -hadrons produced at the LHC [99], making the experimental study of these decays very promising in the near future. It would be, therefore, very interesting to make a comprehensive analysis of the scalar leptoquark effect in these baryonic decays, which will be presented in a forthcoming work [100].

## Acknowledgments

The work is supported by the National Natural Science Foundation of China (NSFC) under contract Nos. 11005032, 11225523, 11221504 and 11435003. XL is also supported by the Scientific Research Foundation for the Returned Overseas Chinese Scholars, State Education Ministry, and by the self-determined research funds of CCNU from the colleges' basic research and operation of MOE (CCNU15A02037). XZ is supported by the CCNU-QLPL Innovation Fund (QLPL2015P01).

## A Analytic formulae of $B_c^- \rightarrow \gamma \tau^- \bar{\nu}_\tau$ , $\bar{B} \rightarrow D^{(*)} \tau \bar{\nu}_\tau$ and $B \rightarrow X_c \tau \bar{\nu}_\tau$

In this appendix, we give all the relevant formulae used to calculate the observables in  $B_c^- \rightarrow \gamma \tau^- \bar{\nu}_\tau$ ,  $\bar{B} \rightarrow D^{(*)} \tau \bar{\nu}_\tau$  and  $B \rightarrow X_c \tau \bar{\nu}_\tau$  decays.


### A.1 The radiative leptonic decay $B_c^- \rightarrow \gamma \tau^- \bar{\nu}_\tau$

Starting from the effective Hamiltonian eq. (2.7), we find that there are only three tree-level Feynman diagrams contributing to  $B_c^- \rightarrow \gamma \tau \bar{\nu}_\tau$ , which are shown in figure 7.

To calculate these Feynman diagrams, we adopt the peaking approximation for the  $B_c$ -meson wave functions,  $\phi_{B_c}(x_c) = \delta(x_c - \frac{m_c}{m_{B_c}})$ , with  $x_c$  being the momentum fraction of the  $c$  quark [101–105]. The spinor part of the  $B_c$ -meson projector is given by [106, 107]

$$-\frac{i}{4} f_{B_c} (\not{p}_{B_c} + m_{B_c}) \gamma_5. \tag{A.1}$$

Then we get the amplitudes for figures 7(a), 7(b) and 7(c) containing both the SM and the



**Figure 7.** The leading-order Feynman diagrams for  $B_c \rightarrow \gamma \tau \bar{\nu}_\tau$ , starting with the effective Hamiltonian eq. (2.7).

scalar leptoquark contributions

$$\begin{aligned} \mathcal{A}_{(a)} = & -\frac{i4G_F}{\sqrt{2}} V_{cb} \left(-\frac{ie}{3}\right) \left(-\frac{i}{4} f_{B_c}\right) \frac{i}{\bar{p}_b^2 - m_b^2} \left\{ \text{Tr}[C_V \gamma_\mu P_L (\not{p}_b + m_b) \gamma_\rho (\not{p}_{B_c} + m_{B_c}) \gamma_5] \epsilon^{\rho*} \bar{\tau} \gamma^\mu P_L \nu_\tau \right. \\ & + \text{Tr}[C_S P_L (\not{p}_b + m_b) \gamma_\rho (\not{p}_{B_c} + m_{B_c}) \gamma_5] \epsilon^{\rho*} \bar{\tau} P_L \nu_\tau \\ & \left. + \text{Tr}[C_T \sigma_{\mu\nu} P_L (\not{p}_b + m_b) \gamma_\rho (\not{p}_{B_c} + m_{B_c}) \gamma_5] \epsilon^{\rho*} \bar{\tau} \sigma^{\mu\nu} P_L \nu_\tau \right\}, \end{aligned} \quad (\text{A.2})$$

$$\begin{aligned} \mathcal{A}_{(b)} = & -\frac{i4G_F}{\sqrt{2}} V_{cb} \left(\frac{2ie}{3}\right) \left(-\frac{i}{4} f_{B_c}\right) \frac{i}{\bar{p}_c^2 - m_c^2} \left\{ \text{Tr}[C_V \gamma_\rho (\not{p}_c + m_c) \gamma_\mu P_L (\not{p}_{B_c} + m_{B_c}) \gamma_5] \epsilon^{\rho*} \bar{\tau} \gamma^\mu P_L \nu_\tau \right. \\ & + \text{Tr}[C_S \gamma_\rho (\not{p}_c + m_c) P_L (\not{p}_{B_c} + m_{B_c}) \gamma_5] \epsilon^{\rho*} \bar{\tau} P_L \nu_\tau \\ & \left. + \text{Tr}[C_T \gamma_\rho (\not{p}_c + m_c) \sigma_{\mu\nu} P_L (\not{p}_{B_c} + m_{B_c}) \gamma_5] \epsilon^{\rho*} \bar{\tau} \sigma^{\mu\nu} P_L \nu_\tau \right\}, \end{aligned} \quad (\text{A.3})$$

$$\begin{aligned} \mathcal{A}_{(c)} = & -\frac{i4G_F}{\sqrt{2}} V_{cb} (-ie) \frac{-i f_{B_c}}{2} \frac{i}{\bar{p}_\tau^2 - m_\tau^2} \left[ C_V p_{B_c}^\mu \bar{\tau} \gamma_\rho (\not{p}_\tau + m_\tau) \gamma_\mu P_L \nu_\tau \epsilon^{\rho*} \right. \\ & \left. - C_S m_{B_c} \bar{\tau} \gamma_\rho (\not{p}_\tau + m_\tau) P_L \nu_\tau \epsilon^{\rho*} \right], \end{aligned} \quad (\text{A.4})$$

where  $C_{V,S,T}$  are the Wilson coefficients of the corresponding operators at the scale  $\mu = m_b$ , and  $\bar{p}_b = p_b - k$ ,  $\bar{p}_c = k - p_c$  and  $\bar{p}_\tau = p_\tau + k$  are, respectively, the momentum of the  $b$ ,  $c$  and  $\tau$  propagators, with  $p_b = \frac{m_b}{m_{B_c}} p_{B_c}$ ,  $p_c = \frac{m_c}{m_{B_c}} p_{B_c}$  and  $k$  the momentum of the photon. The photon energy dependence of the differential branching ratio is given by

$$d\mathcal{B}(B_c^- \rightarrow \gamma \tau^- \bar{\nu}_\tau) / dE_\gamma = \int dE_{\nu_\tau} \frac{\tau_{B_c}}{2m_{B_c}} |\mathcal{A}_{(a)} + \mathcal{A}_{(b)} + \mathcal{A}_{(c)}|^2 d\Phi_3, \quad (\text{A.5})$$

where  $d\Phi_3 = \frac{1}{32\pi^3} dE_\gamma dE_{\nu_\tau}$  is the three-body phase space. Integrating over the photon energy  $E_\gamma$ , one can then obtain the total branching ratio.

## A.2 Exclusive semi-leptonic decays $\bar{B} \rightarrow D^{(*)} \tau \bar{\nu}_\tau$

For the exclusive semi-leptonic  $\bar{B} \rightarrow D \tau \bar{\nu}_\tau$  and  $\bar{B} \rightarrow D^* \tau \bar{\nu}_\tau$  decays, we follow the helicity amplitude formalism that is commonly used in the literatures [13, 17, 19, 38, 44]. For simplicity, we list below the relevant formulae without any detailed derivations.

(1)  $\bar{B} \rightarrow D\tau\bar{\nu}_\tau$

- The differential decay rates

$$\begin{aligned} \frac{d\Gamma^{\lambda_\tau=1/2}}{dq^2} &= \frac{G_F^2 |V_{cb}|^2}{192\pi^3 m_B^3} q^2 \sqrt{\lambda(q^2)} \left(1 - \frac{m_\tau^2}{q^2}\right)^2 \left\{ \frac{1}{2} |C_V|^2 \frac{m_\tau^2}{q^2} (H_0^s{}^2 + 3H_t^s{}^2) + \frac{3}{2} |C_S|^2 \right. \\ &\quad \left. H^s{}^2 + 8|C_T|^2 H_{+-}^s{}^2 + 3\Re[C_V C_S^*] \frac{m_\tau}{\sqrt{q^2}} H^s H_t^s - 4\Re[C_V C_T^*] \frac{m_\tau}{\sqrt{q^2}} H_{+-}^s - H_0^s \right\}, \\ \frac{d\Gamma^{\lambda_\tau=-1/2}}{dq^2} &= \frac{G_F^2 |V_{cb}|^2}{192\pi^3 m_B^3} q^2 \sqrt{\lambda(q^2)} \left(1 - \frac{m_\tau^2}{q^2}\right)^2 \left\{ |C_V|^2 H_0^s{}^2 + 16|C_T|^2 \frac{m_\tau^2}{q^2} H_{+-}^s{}^2 \right. \\ &\quad \left. - 8\Re[C_V C_T^*] \frac{m_\tau}{\sqrt{q^2}} H_{+-}^s - H_0^s \right\}, \\ \frac{d\Gamma}{dq^2} &= \frac{d\Gamma^{\lambda_\tau=1/2}}{dq^2} + \frac{d\Gamma^{\lambda_\tau=-1/2}}{dq^2}, \end{aligned} \quad (\text{A.6})$$

where  $\lambda^{(*)}(q^2) = \lambda(m_B^2, m_{D^{(*)}}^2, q^2)$  with  $\lambda(a, b, c) = a^2 + b^2 + c^2 - 2(ab + bc + ca)$ ,  $H_s$  are the hadronic amplitudes given in appendix B, and all the Wilson coefficients  $C_i$  ( $i = V, S, T$ ) are evaluated at the scale  $\mu_b = m_b$ .

- The  $q^2$  dependent ratio

$$R_D(q^2) = \frac{d\Gamma(\bar{B} \rightarrow D\tau\bar{\nu}_\tau)/dq^2}{d\Gamma(\bar{B} \rightarrow D\ell\bar{\nu}_\ell)/dq^2}, \quad (\text{A.7})$$

where  $\ell$  denotes the light lepton ( $e$  or  $\mu$ ).

- The longitudinal polarization of  $\tau$

$$P_\tau^D(q^2) = \frac{d\Gamma^{\lambda_\tau=1/2}/dq^2 - d\Gamma^{\lambda_\tau=-1/2}/dq^2}{d\Gamma^{\lambda_\tau=1/2}/dq^2 + d\Gamma^{\lambda_\tau=-1/2}/dq^2}. \quad (\text{A.8})$$

- The lepton forward-backward asymmetry

$$A_{\text{FB}}^D(q^2) = \frac{\int_0^1 d\cos\theta (d^2\Gamma/dq^2 d\cos\theta) - \int_{-1}^0 d\cos\theta (d^2\Gamma/dq^2 d\cos\theta)}{d\Gamma/dq^2}, \quad (\text{A.9})$$

where  $\theta$  is the angle between the three-momentum of  $\tau$  and that of the  $D$  meson in the  $\tau\text{-}\bar{\nu}_\tau$  center-of-mass frame. Writing the double-differential decay rates as [44]

$$\frac{d^2\Gamma}{dq^2 d\cos\theta} = a_\theta(q^2) + b_\theta^{D^{(*)}}(q^2) \cos\theta + c_\theta(q^2) \cos^2\theta, \quad (\text{A.10})$$

one can then see clearly that the coefficient  $b_\theta$  determines the lepton forward-backward asymmetry, with

$$\begin{aligned} A_{\text{FB}}^D(q^2) = b_\theta^D(q^2) &= \frac{G_F^2 |V_{cb}|^2}{128\pi^3 m_B^3} q^2 \sqrt{\lambda(q^2)} \left(1 - \frac{m_\tau^2}{q^2}\right)^2 \left\{ |C_V|^2 \frac{m_\tau^2}{q^2} H_0^s H_t^s + \Re[C_V C_S^*] \right. \\ &\quad \left. \frac{m_\tau}{\sqrt{q^2}} H^s H_0^s - 4\Re[C_V C_T^*] \frac{m_\tau}{\sqrt{q^2}} H_{+-}^s - H_t^s - 4\Re[C_S C_T^*] H_{+-}^s - H^s \right\}, \end{aligned} \quad (\text{A.11})$$



and

$$\begin{aligned}
 A_{\text{FB}}^{D^*}(q^2) &= b_{\theta}^{D^*}(q^2) = \frac{G_F^2 |V_{cb}|^2}{128\pi^3 m_B^3} q^2 \sqrt{\lambda^*(q^2)} \left(1 - \frac{m_\tau^2}{q^2}\right)^2 \left\{ |C_V|^2 \left[ \frac{1}{2} (H_+^{+2} - H_-^{-2}) \right. \right. \\
 &\quad \left. \left. + \frac{m_\tau^2}{q^2} H_0^0 H_t^0 \right] + 8|C_T|^2 \frac{m_\tau^2}{q^2} \cdot (H_{+t}^{02} - H_{-t}^{02}) - \Re[C_V C_S^*] \frac{m_\tau}{\sqrt{q^2}} \right. \\
 &\quad \left. H_0^0 H_0^0 - 4\Re[C_V C_T^*] \frac{m_\tau}{\sqrt{q^2}} (H_{+-}^0 - H_t^0 + H_{+t}^0 H_+^+ - H_{-t}^0 H_-^-) \right. \\
 &\quad \left. + 4\Re[C_S C_T^*] H_{+-}^0 - H_0^0 \right\}. \tag{A.12}
 \end{aligned}$$

(2)  $\bar{B} \rightarrow D^* \tau \bar{\nu}_\tau$

- The differential decay rates

$$\begin{aligned}
 \frac{d\Gamma^{\lambda_\tau=1/2}}{dq^2} &= \frac{G_F^2 |V_{cb}|^2}{192\pi^3 m_B^3} q^2 \sqrt{\lambda^*(q^2)} \left(1 - \frac{m_\tau^2}{q^2}\right)^2 \left\{ \frac{1}{2} |C_V|^2 \frac{m_\tau^2}{q^2} (H_+^{+2} + H_-^{-2} + H_0^{02} + H_t^{02}) \right. \\
 &\quad \left. + \frac{3}{2} |C_S|^2 H^{02} + 8|C_T|^2 (H_{+t}^{+2} + H_{-t}^{-2} + H_{+-}^{02}) - 3\Re[C_V C_S^*] \frac{m_\tau}{\sqrt{q^2}} H^0 H_t^0 \right. \\
 &\quad \left. - 4\Re[C_V C_T^*] \frac{m_\tau}{\sqrt{q^2}} (H_{+-}^0 - H_0^0 + H_{+t}^+ H_+^+ + H_{-t}^- H_-^-) \right\}, \\
 \frac{d\Gamma^{\lambda_\tau=-1/2}}{dq^2} &= \frac{G_F^2 |V_{cb}|^2}{192\pi^3 m_B^3} q^2 \sqrt{\lambda^*(q^2)} \left(1 - \frac{m_\tau^2}{q^2}\right)^2 \left\{ |C_V|^2 (H_+^{+2} + H_-^{-2} + H_0^{02}) + 16|C_T|^2 \frac{m_\tau^2}{q^2} \right. \\
 &\quad \left. (H_{+t}^{+2} + H_{-t}^{-2} + H_{+-}^{02}) - 8\Re[C_V C_T^*] \frac{m_\tau}{\sqrt{q^2}} (H_{+-}^0 - H_0^0 + H_{+t}^+ H_+^+ + H_{-t}^- H_-^-) \right\}, \\
 \frac{d\Gamma^{\lambda_{D^*}=\pm 1}}{dq^2} &= \frac{G_F^2 |V_{cb}|^2}{192\pi^3 m_B^3} q^2 \sqrt{\lambda^*(q^2)} \left(1 - \frac{m_\tau^2}{q^2}\right)^2 \left\{ \left(1 + \frac{m_\tau^2}{2q^2}\right) (|C_V|^2 H_{\pm\pm}^{\pm 2} + 8|C_T|^2 \right. \\
 &\quad \left. \left(1 + \frac{2m_\tau^2}{q^2}\right) H_{\pm t}^{\pm 2} \mp 12\Re[C_V C_T^*] \frac{m_\tau}{\sqrt{q^2}} H_{\pm t}^{\pm 2} H_{\pm\pm}^{\pm 2} \right\}, \\
 \frac{d\Gamma^{\lambda_{D^*}=0}}{dq^2} &= \frac{G_F^2 |V_{cb}|^2}{192\pi^3 m_B^3} q^2 \sqrt{\lambda^*(q^2)} \left(1 - \frac{m_\tau^2}{q^2}\right)^2 \left\{ |C_V|^2 \left[ \left(1 + \frac{m_\tau^2}{2q^2}\right) H_0^{02} + \frac{3}{2} \frac{m_\tau^2}{q^2} H_t^{02} \right] \right. \\
 &\quad \left. + \frac{3}{2} |C_S|^2 H^{02} + 8|C_T|^2 \left(1 + \frac{2m_\tau^2}{q^2}\right) H_{+-}^{02} - 3\Re[C_V C_S^*] \frac{m_\tau}{\sqrt{q^2}} H^0 H_t^0 \right. \\
 &\quad \left. - 12\Re[C_V C_T^*] \frac{m_\tau}{\sqrt{q^2}} H_{+-}^0 - H_0^0 \right\}. \tag{A.13}
 \end{aligned}$$

Besides the observables similar to that defined in  $B^- \rightarrow D\tau^-\bar{\nu}_\tau$ , there are another two observables in this process, i.e., the longitudinal and transverse polarizations of the  $D^*$  meson defined, respectively, by

$$P_L^{D^*}(q^2) = \frac{d\Gamma^{\lambda_{D^*}=0}/dq^2}{d\Gamma^{\lambda_{D^*}=0}/dq^2 + d\Gamma^{\lambda_{D^*}=1}/dq^2 + d\Gamma^{\lambda_{D^*}=-1}/dq^2}, \tag{A.14}$$

$$P_T^{D^*}(q^2) = 1 - P_L^{D^*}(q^2). \tag{A.15}$$

### A.3 Inclusive semi-leptonic decay $B \rightarrow X_c \tau \bar{\nu}_\tau$

In the heavy-quark limit  $m_b \gg \Lambda_{\text{QCD}}$ , the inclusive semi-leptonic decay rate is equivalent to the perturbative quark-level  $b$  decay rate [4, 92–94]. This makes it possible to get the inclusive decay rate by calculating directly the rate for the quark-level process  $b \rightarrow c \tau \bar{\nu}_\tau$ , with the result given by

$$\begin{aligned} \frac{d\Gamma}{d\hat{q}^2} = & \frac{G_F^2 V_{cb}^2 m_b^5}{192\pi^3} \left(1 - \frac{x_\tau}{\hat{q}^2}\right)^2 \lambda^{1/2}(1, x_c, \hat{q}^2) \left[ 3|C_S|^2(1+x_c-\hat{q}^2)\hat{q}^2 + 6\Re[C_V C_S^*]x_c^{1/2}x_\tau^{1/2}(1-x_c+\hat{q}^2) \right. \\ & + 16|C_T|^2(2x_c^2-x_c(\hat{q}^2+4)-\hat{q}^4-\hat{q}^2+2) \left(1 + \frac{2x_\tau}{\hat{q}^2}\right) - 72\Re[C_V C_T^*]x_c^{1/2}x_\tau^{1/2}(1-x_c+\hat{q}^2) \\ & \left. + 2|C_V|^2 \left( (1+x_c-x_\tau)\hat{q}^2 + (1-x_c)^2 - x_\tau(1+x_c) + 2(1-x_c)^2 \frac{x_\tau}{\hat{q}^2} - 2\hat{q}^4 \right) \right], \quad (\text{A.16}) \end{aligned}$$

where  $x_i = \frac{m_i^2}{m_b^2}$ , and  $\hat{q}^2 = \frac{q^2}{m_b^2}$  with  $\hat{q}^2$  varying from  $x_\tau$  to  $(1-\sqrt{x_c})^2$ . The  $\tau$ -energy spectrum of this process is given by

$$\begin{aligned} \frac{d\Gamma}{dy} = & \frac{G_F^2 V_{cb}^2 m_b^5}{192\pi^3} \frac{\sqrt{y^2 - 4x_\tau}(1+x_\tau-x_c-y)^2}{2(1+x_\tau-y)^3} \left[ (4|C_V|^2 + |C_S|^2) (2y^3 - (x_c + 5x_\tau + 5)y^2) \right. \\ & + (3x_c(x_\tau + 1) + (3x_\tau^2 + 10x_\tau + 3))y - 4x_\tau(2x_c + x_\tau + 1) + 16|C_T|^2 (14y^3 \\ & - (x_c + 29x_\tau + 29)y^2 + (3x_c(x_\tau + 1) + 15x_\tau^2 + 34x_\tau + 15)y - 4x_\tau(2x_c + x_\tau + 1)) \\ & - 12(\Re[C_V C_S^*] - 12\Re[C_V C_T^*])\sqrt{x_c}\sqrt{x_\tau}(1+x_\tau-y)(y-2) + 8\Re[C_S C_T^*] (4y^3 \\ & \left. + (x_c - 7x_\tau - 7)y^2 - (3x_c(x_\tau + 1) - 3x_\tau^2 - 2x_\tau - 3)y + 4x_\tau(2x_c + x_\tau + 1)) \right], \quad (\text{A.17}) \end{aligned}$$

where  $y = \frac{2E_\tau}{m_b}$ , with  $y$  varying from  $2\sqrt{x_\tau}$  to  $1 + x_\tau - x_c$ .

## B Hadronic amplitudes in $\bar{B} \rightarrow D^{(*)} \tau \bar{\nu}_\tau$ decays

The hadronic amplitudes in  $\bar{B} \rightarrow D^{(*)} \tau \bar{\nu}_\tau$  decays are given, respectively, as [13, 17, 38, 44]

- $\bar{B} \rightarrow D \tau \bar{\nu}_\tau$ .

$$\begin{aligned} H_0^s(q^2) &= \sqrt{\frac{\lambda_D(q^2)}{q^2}} F_1(q^2), & H_t^s(q^2) &= \frac{m_B^2 - m_D^2}{\sqrt{q^2}} F_0(q^2), \\ H^s(q^2) &= \frac{m_B^2 - m_D^2}{m_b - m_c} F_0(q^2), & H_{+-}^s(q^2) &= H_{0t}^s(q^2) = -\frac{\sqrt{\lambda_D(q^2)}}{m_B + m_D} F_T(q^2). \end{aligned} \quad (\text{B.1})$$

- $\bar{B} \rightarrow D^* \tau \bar{\nu}_\tau$ .

$$\begin{aligned}
 H_{\pm}^{\pm}(q^2) &= (m_B + m_{D^*})A_1(q^2) \mp \frac{\sqrt{\lambda_{D^*}(q^2)}}{m_B + m_{D^*}}V(q^2), \\
 H_0^0(q^2) &= \frac{m_B + m_{D^*}}{2m_{D^*}\sqrt{q^2}} \left[ -(m_B^2 - m_{D^*}^2 - q^2)A_1(q^2) + \frac{\lambda_{D^*}(q^2)}{(m_B + m_{D^*})^2}A_2(q^2) \right], \\
 H_t^0(q^2) &= -\sqrt{\frac{\lambda_{D^*}(q^2)}{q^2}}A_0(q^2), \quad H^0(q^2) = \frac{\sqrt{\lambda_{D^*}(q^2)}}{m_b + m_c}A_0(q^2), \\
 H_{\pm t}^{\pm}(q^2) &= \frac{1}{\sqrt{q^2}} \left[ (m_B^2 - m_{D^*}^2)T_2(q^2) \pm \sqrt{\lambda_{D^*}(q^2)}T_1(q^2) \right], \\
 H_{+-}^0(q^2) = H_{0t}^0(q^2) &= \frac{1}{2m_{D^*}} \left[ -(m_B^2 + 3m_{D^*}^2 - q^2)T_2(q^2) + \frac{\lambda_{D^*}(q^2)}{m_B^2 - m_{D^*}^2}T_3(q^2) \right], \quad (\text{B.2})
 \end{aligned}$$

where all the  $B \rightarrow D$  and  $B \rightarrow D^*$  form factors are taken from refs. [44, 108], except for  $S_1(w)$ , for which we use the form given by eq. (A6) in ref. [51].

**Open Access.** This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

## References

- [1] B. Grinstein, *The static quark effective theory*, *Nucl. Phys. B* **339** (1990) 253 [[INSPIRE](#)].
- [2] E. Eichten and B.R. Hill, *An effective field theory for the calculation of matrix elements involving heavy quarks*, *Phys. Lett. B* **234** (1990) 511 [[INSPIRE](#)].
- [3] H. Georgi, *An effective field theory for heavy quarks at low-energies*, *Phys. Lett. B* **240** (1990) 447 [[INSPIRE](#)].
- [4] A.V. Manohar and M.B. Wise, *Heavy quark physics*, *Camb. Monogr. Part. Phys. Nucl. Phys. Cosmol.* **10** (2000) 1 [[INSPIRE](#)].
- [5] M. Neubert, *Heavy quark symmetry*, *Phys. Rept.* **245** (1994) 259 [[hep-ph/9306320](#)] [[INSPIRE](#)].
- [6] BABAR collaboration, J.P. Lees et al., *Evidence for an excess of  $\bar{B} \rightarrow D^{(*)} \tau^- \bar{\nu}_\tau$  decays*, *Phys. Rev. Lett.* **109** (2012) 101802 [[arXiv:1205.5442](#)] [[INSPIRE](#)].
- [7] BABAR collaboration, J.P. Lees et al., *Measurement of an excess of  $\bar{B} \rightarrow D^{(*)} \tau^- \bar{\nu}_\tau$  decays and implications for charged Higgs bosons*, *Phys. Rev. D* **88** (2013) 072012 [[arXiv:1303.0571](#)] [[INSPIRE](#)].
- [8] BELLE collaboration, M. Huschle et al., *Measurement of the branching ratio of  $\bar{B} \rightarrow D^{(*)} \tau^- \bar{\nu}_\tau$  relative to  $\bar{B} \rightarrow D^{(*)} \ell^- \bar{\nu}_\ell$  decays with hadronic tagging at Belle*, *Phys. Rev. D* **92** (2015) 072014 [[arXiv:1507.03233](#)] [[INSPIRE](#)].
- [9] BELLE collaboration, A. Abdesselam et al., *Measurement of the branching ratio of  $\bar{B}^0 \rightarrow D^{*+} \tau^- \bar{\nu}_\tau$  relative to  $\bar{B}^0 \rightarrow D^{*+} \ell^- \bar{\nu}_\ell$  decays with a semileptonic tagging method*, [arXiv:1603.06711](#) [[INSPIRE](#)].

- [10] LHCb collaboration, *Measurement of the ratio of branching fractions*  $\mathcal{B}(\bar{B}^0 \rightarrow D^{*+}\tau^-\bar{\nu}_\tau)/\mathcal{B}(\bar{B}^0 \rightarrow D^{*+}\mu^-\bar{\nu}_\mu)$ , *Phys. Rev. Lett.* **115** (2015) 111803 [*Addendum ibid.* **115** (2015) 159901] [[arXiv:1506.08614](#)] [[INSPIRE](#)].
- [11] HEAVY FLAVOR AVERAGING GROUP (HFAG) collaboration, *Average of  $R(D)$  and  $R(D^*)$  for Winter 2016 webpage*  
<http://www.slac.stanford.edu/xorg/hfag/semi/winter16/winter16.dtaunu.html>.
- [12] HPQCD collaboration, H. Na, C.M. Bouchard, G.P. Lepage, C. Monahan and J. Shigemitsu,  *$B \rightarrow D\ell\nu$  form factors at nonzero recoil and extraction of  $|V_{cb}|$* , *Phys. Rev. D* **92** (2015) 054510 [[arXiv:1505.03925](#)] [[INSPIRE](#)].
- [13] S. Fajfer, J.F. Kamenik and I. Nisandzic, *On the  $B \rightarrow D^*\tau\bar{\nu}_\tau$  sensitivity to new physics*, *Phys. Rev. D* **85** (2012) 094025 [[arXiv:1203.2654](#)] [[INSPIRE](#)].
- [14] M. Freytsis, Z. Ligeti and J.T. Ruderman, *Flavor models for  $\bar{B} \rightarrow D^{(*)}\tau\bar{\nu}$* , *Phys. Rev. D* **92** (2015) 054018 [[arXiv:1506.08896](#)] [[INSPIRE](#)].
- [15] L. Calibbi, A. Crivellin and T. Ota, *Effective field theory approach to  $b \rightarrow s\ell\ell^{(\prime)}$ ,  $B \rightarrow K^{(*)}\nu\bar{\nu}$  and  $B \rightarrow D^{(*)}\tau\nu$  with third generation couplings*, *Phys. Rev. Lett.* **115** (2015) 181801 [[arXiv:1506.02661](#)] [[INSPIRE](#)].
- [16] R. Alonso, B. Grinstein and J. Martin Camalich, *Lepton universality violation and lepton flavor conservation in  $B$ -meson decays*, *JHEP* **10** (2015) 184 [[arXiv:1505.05164](#)] [[INSPIRE](#)].
- [17] M. Tanaka and R. Watanabe, *New physics in the weak interaction of  $\bar{B} \rightarrow D^{(*)}\tau\bar{\nu}$* , *Phys. Rev. D* **87** (2013) 034028 [[arXiv:1212.1878](#)] [[INSPIRE](#)].
- [18] S. Fajfer, J.F. Kamenik, I. Nisandzic and J. Zupan, *Implications of lepton flavor universality violations in  $B$  decays*, *Phys. Rev. Lett.* **109** (2012) 161801 [[arXiv:1206.1872](#)] [[INSPIRE](#)].
- [19] D. Becirevic, S. Fajfer, I. Nisandzic and A. Tayduganov, *Angular distributions of  $\bar{B} \rightarrow D^{(*)}\ell\bar{\nu}_\ell$  decays and search of new physics*, [arXiv:1602.03030](#) [[INSPIRE](#)].
- [20] S. Bhattacharya, S. Nandi and S.K. Patra, *Optimal-observable analysis of possible new physics in  $B \rightarrow D^{(*)}\tau\nu_\tau$* , *Phys. Rev. D* **93** (2016) 034011 [[arXiv:1509.07259](#)] [[INSPIRE](#)].
- [21] B. Bhattacharya, A. Datta, D. London and S. Shivashankara, *Simultaneous explanation of the  $R_K$  and  $R(D^{(*)})$  puzzles*, *Phys. Lett. B* **742** (2015) 370 [[arXiv:1412.7164](#)] [[INSPIRE](#)].
- [22] M. Duraisamy, P. Sharma and A. Datta, *Azimuthal  $B \rightarrow D^*\tau^-\bar{\nu}_\tau$  angular distribution with tensor operators*, *Phys. Rev. D* **90** (2014) 074013 [[arXiv:1405.3719](#)] [[INSPIRE](#)].
- [23] K. Hagiwara, M.M. Nojiri and Y. Sakaki,  *$CP$  violation in  $B \rightarrow D\tau\nu_\tau$  using multipion  $\tau$  decays*, *Phys. Rev. D* **89** (2014) 094009 [[arXiv:1403.5892](#)] [[INSPIRE](#)].
- [24] R. Dutta, A. Bhol and A.K. Giri, *Effective theory approach to new physics in  $b \rightarrow u$  and  $b \rightarrow c$  leptonic and semileptonic decays*, *Phys. Rev. D* **88** (2013) 114023 [[arXiv:1307.6653](#)] [[INSPIRE](#)].
- [25] M. Duraisamy and A. Datta, *The full  $B \rightarrow D^*\tau^-\bar{\nu}_\tau$  angular distribution and  $CP$ -violating triple products*, *JHEP* **09** (2013) 059 [[arXiv:1302.7031](#)] [[INSPIRE](#)].
- [26] P. Biancofiore, P. Colangelo and F. De Fazio, *On the anomalous enhancement observed in  $B \rightarrow D^{(*)}\tau\bar{\nu}_\tau$  decays*, *Phys. Rev. D* **87** (2013) 074010 [[arXiv:1302.1042](#)] [[INSPIRE](#)].
- [27] J.A. Bailey et al., *Refining new-physics searches in  $B \rightarrow D\tau\nu$  decay with lattice QCD*, *Phys. Rev. Lett.* **109** (2012) 071802 [[arXiv:1206.4992](#)] [[INSPIRE](#)].

- [28] D. Bečirević, N. Košnik and A. Tayduganov,  $\bar{B} \rightarrow D\tau\bar{\nu}_\tau$  vs.  $\bar{B} \rightarrow D\mu\bar{\nu}_\mu$ , *Phys. Lett. B* **716** (2012) 208 [[arXiv:1206.4977](#)] [[INSPIRE](#)].
- [29] A. Datta, M. Duraisamy and D. Ghosh, *Diagnosing new physics in  $b \rightarrow c\tau\nu_\tau$  decays in the light of the recent BaBar result*, *Phys. Rev. D* **86** (2012) 034027 [[arXiv:1206.3760](#)] [[INSPIRE](#)].
- [30] S. Faller, T. Mannel and S. Turczyk, *Limits on new physics from exclusive  $B \rightarrow D^{(*)}\ell\bar{\nu}$  decays*, *Phys. Rev. D* **84** (2011) 014022 [[arXiv:1105.3679](#)] [[INSPIRE](#)].
- [31] C.-H. Chen and C.-Q. Geng, *Lepton angular asymmetries in semileptonic charmful  $B$  decays*, *Phys. Rev. D* **71** (2005) 077501 [[hep-ph/0503123](#)] [[INSPIRE](#)].
- [32] Y.-Y. Fan, W.-F. Wang, S. Cheng and Z.-J. Xiao, *Semileptonic decays  $B \rightarrow D^{(*)}\ell\nu$  in the perturbative QCD factorization approach*, *Chin. Sci. Bull.* **59** (2014) 125 [[arXiv:1301.6246](#)] [[INSPIRE](#)].
- [33] Y.-Y. Fan, Z.-J. Xiao, R.-M. Wang and B.-Z. Li, *The  $B \rightarrow D^{(*)}\ell\nu_\ell$  decays in the pQCD approach with the lattice QCD input*, [arXiv:1505.07169](#) [[INSPIRE](#)].
- [34] A.K. Alok, D. Kumar, S. Kumbhakar and S.U. Sankar,  *$D^*$  polarization as a probe to discriminate new physics in  $B \rightarrow D^*\tau\bar{\nu}$* , [arXiv:1606.03164](#) [[INSPIRE](#)].
- [35] M.A. Ivanov, J.G. Körner and C.-T. Tran, *Analyzing new physics in the decays  $\bar{B}^0 \rightarrow D^{(*)}\tau^-\bar{\nu}_\tau$  with form factors obtained from the covariant quark model*, [arXiv:1607.02932](#) [[INSPIRE](#)].
- [36] F.F. Deppisch, S. Kulkarni, H. Päs and E. Schumacher, *Leptoquark patterns unifying neutrino masses, flavor anomalies and the diphoton excess*, *Phys. Rev. D* **94** (2016) 013003 [[arXiv:1603.07672](#)] [[INSPIRE](#)].
- [37] B. Dumont, K. Nishiwaki and R. Watanabe, *LHC constraints and prospects for  $S_1$  scalar leptoquark explaining the  $\bar{B} \rightarrow D^{(*)}\tau\bar{\nu}$  anomaly*, [arXiv:1603.05248](#) [[INSPIRE](#)].
- [38] I. Doršner, S. Fajfer, A. Greljo, J.F. Kamenik and N. Košnik, *Physics of leptoquarks in precision experiments and at particle colliders*, *Phys. Rept.* **641** (2016) 1 [[arXiv:1603.04993](#)] [[INSPIRE](#)].
- [39] Y. Sakaki, M. Tanaka, A. Tayduganov and R. Watanabe, *Probing new physics with  $q^2$  distributions in  $\bar{B} \rightarrow D^{(*)}\tau\bar{\nu}$* , *Phys. Rev. D* **91** (2015) 114028 [[arXiv:1412.3761](#)] [[INSPIRE](#)].
- [40] M. Bauer and M. Neubert, *Minimal leptoquark explanation for the  $R_{D^{(*)}}$ ,  $R_K$  and  $(g-2)_g$  anomalies*, *Phys. Rev. Lett.* **116** (2016) 141802 [[arXiv:1511.01900](#)] [[INSPIRE](#)].
- [41] S. Fajfer and N. Košnik, *Vector leptoquark resolution of  $R_K$  and  $R_{D^{(*)}}$  puzzles*, *Phys. Lett. B* **755** (2016) 270 [[arXiv:1511.06024](#)] [[INSPIRE](#)].
- [42] S. Sahoo and R. Mohanta, *Lepton flavor violating  $B$  meson decays via a scalar leptoquark*, *Phys. Rev. D* **93** (2016) 114001 [[arXiv:1512.04657](#)] [[INSPIRE](#)].
- [43] R. Barbieri, G. Isidori, A. Pattori and F. Senia, *Anomalies in  $B$ -decays and  $U(2)$  flavour symmetry*, *Eur. Phys. J. C* **76** (2016) 67 [[arXiv:1512.01560](#)] [[INSPIRE](#)].
- [44] Y. Sakaki, M. Tanaka, A. Tayduganov and R. Watanabe, *Testing leptoquark models in  $\bar{B} \rightarrow D^{(*)}\tau\bar{\nu}$* , *Phys. Rev. D* **88** (2013) 094012 [[arXiv:1309.0301](#)] [[INSPIRE](#)].
- [45] A. Celis, M. Jung, X.-Q. Li and A. Pich, *Sensitivity to charged scalars in  $B \rightarrow D^{(*)}\tau\nu_\tau$  and  $B \rightarrow \tau\nu_\tau$  decays*, *JHEP* **01** (2013) 054 [[arXiv:1210.8443](#)] [[INSPIRE](#)].

- [46] J.M. Cline, *Scalar doublet models confront  $\tau$  and  $b$  anomalies*, *Phys. Rev. D* **93** (2016) 075017 [[arXiv:1512.02210](#)] [[INSPIRE](#)].
- [47] C.S. Kim, Y.W. Yoon and X.-B. Yuan, *Exploring top quark FCNC within 2HDM type-III in association with flavor physics*, *JHEP* **12** (2015) 038 [[arXiv:1509.00491](#)] [[INSPIRE](#)].
- [48] A. Crivellin, J. Heeck and P. Stoffer, *A perturbed lepton-specific two-Higgs-doublet model facing experimental hints for physics beyond the Standard Model*, *Phys. Rev. Lett.* **116** (2016) 081801 [[arXiv:1507.07567](#)] [[INSPIRE](#)].
- [49] D.S. Hwang, *Transverse spin polarization of  $\tau^-$  in  $\bar{B}^0 \rightarrow D^+ \tau^- \bar{\nu}$  and charged Higgs boson*, [arXiv:1504.06933](#) [[INSPIRE](#)].
- [50] A. Crivellin, A. Kokulu and C. Greub, *Flavor-phenomenology of two-Higgs-doublet models with generic Yukawa structure*, *Phys. Rev. D* **87** (2013) 094031 [[arXiv:1303.5877](#)] [[INSPIRE](#)].
- [51] Y. Sakaki and H. Tanaka, *Constraints on the charged scalar effects using the forward-backward asymmetry on  $\bar{B} \rightarrow D^{(*)} \tau \bar{\nu}_\tau$* , *Phys. Rev. D* **87** (2013) 054002 [[arXiv:1205.4908](#)] [[INSPIRE](#)].
- [52] U. Nierste, S. Trine and S. Westhoff, *Charged-Higgs effects in a new  $B \rightarrow D \tau \nu$  differential decay distribution*, *Phys. Rev. D* **78** (2008) 015006 [[arXiv:0801.4938](#)] [[INSPIRE](#)].
- [53] K. Kiers and A. Soni, *Improving constraints on  $\tan \beta / m_H$  using  $B \rightarrow D \tau \bar{\nu}$* , *Phys. Rev. D* **56** (1997) 5786 [[hep-ph/9706337](#)] [[INSPIRE](#)].
- [54] M. Tanaka, *Charged Higgs effects on exclusive semitauonic  $B$  decays*, *Z. Phys. C* **67** (1995) 321 [[hep-ph/9411405](#)] [[INSPIRE](#)].
- [55] W.-S. Hou, *Enhanced charged Higgs boson effects in  $B^- \rightarrow \tau \bar{\nu}$ ,  $\mu \bar{\nu}$  and  $b \rightarrow \tau \bar{\nu} + X$* , *Phys. Rev. D* **48** (1993) 2342 [[INSPIRE](#)].
- [56] S.M. Boucenna, A. Celis, J. Fuentes-Martin, A. Vicente and J. Virto, *Non-Abelian gauge extensions for  $B$ -decay anomalies*, *Phys. Lett. B* **760** (2016) 214 [[arXiv:1604.03088](#)] [[INSPIRE](#)].
- [57] C. Hati, G. Kumar and N. Mahajan,  *$\bar{B} \rightarrow D^{(*)} \tau \bar{\nu}$  excesses in ALRSM constrained from  $B$ ,  $D$  decays and  $D^0$ - $\bar{D}^0$  mixing*, *JHEP* **01** (2016) 117 [[arXiv:1511.03290](#)] [[INSPIRE](#)].
- [58] A. Greljo, G. Isidori and D. Marzocca, *On the breaking of lepton flavor universality in  $B$  decays*, *JHEP* **07** (2015) 142 [[arXiv:1506.01705](#)] [[INSPIRE](#)].
- [59] D. Das, C. Hati, G. Kumar and N. Mahajan, *Towards a unified explanation of  $R_{D^{(*)}}$ ,  $R_K$  and  $(g-2)_\mu$  anomalies in a  $L$ - $R$  model*, [arXiv:1605.06313](#) [[INSPIRE](#)].
- [60] J. Zhu, H.-M. Gan, R.-M. Wang, Y.-Y. Fan, Q. Chang and Y.-G. Xu, *Probing the  $R$ -parity violating supersymmetric effects in the exclusive  $b \rightarrow c \ell^- \bar{\nu}_\ell$  decays*, *Phys. Rev. D* **93** (2016) 094023 [[arXiv:1602.06491](#)] [[INSPIRE](#)].
- [61] N.G. Deshpande and A. Menon, *Hints of  $R$ -parity violation in  $B$  decays into  $\tau \nu$* , *JHEP* **01** (2013) 025 [[arXiv:1208.4134](#)] [[INSPIRE](#)].
- [62] LHCb collaboration, *Test of lepton universality using  $B^+ \rightarrow K^+ \ell^+ \ell^-$  decays*, *Phys. Rev. Lett.* **113** (2014) 151601 [[arXiv:1406.6482](#)] [[INSPIRE](#)].
- [63] M. Davier, A. Hoecker, B. Malaescu and Z. Zhang, *Reevaluation of the hadronic contributions to the muon  $g-2$  and to  $\alpha(M_Z^2)$* , *Eur. Phys. J. C* **71** (2011) 1515 [[Erratum ibid. C 72 \(2012\) 1874](#)] [[arXiv:1010.4180](#)] [[INSPIRE](#)].

- [64] I. Doršner, S. Fajfer, N. Košnik and I. Nišandžić, *Minimally flavored colored scalar in  $\bar{B} \rightarrow D^{(*)}\tau\bar{\nu}$  and the mass matrices constraints*, *JHEP* **11** (2013) 084 [[arXiv:1306.6493](#)] [[INSPIRE](#)].
- [65] K.G. Chetyrkin, *Quark mass anomalous dimension to  $O(\alpha_s^4)$* , *Phys. Lett. B* **404** (1997) 161 [[hep-ph/9703278](#)] [[INSPIRE](#)].
- [66] J.A. Gracey, *Three loop  $\overline{MS}$ -bar tensor current anomalous dimension in QCD*, *Phys. Lett. B* **488** (2000) 175 [[hep-ph/0007171](#)] [[INSPIRE](#)].
- [67] HPQCD collaboration, B. Colquhoun et al., *B-meson decay constants: a more complete picture from full lattice QCD*, *Phys. Rev. D* **91** (2015) 114509 [[arXiv:1503.05762](#)] [[INSPIRE](#)].
- [68] PARTICLE DATA GROUP collaboration, K.A. Olive et al., *Review of particle physics*, *Chin. Phys. C* **38** (2014) 090001 [[INSPIRE](#)].
- [69] BABAR collaboration, B. Aubert et al., *Measurements of the semileptonic decays  $\bar{B} \rightarrow D\ell\bar{\nu}$  and  $\bar{B} \rightarrow D^*\ell\bar{\nu}$  using a global fit to  $DX\ell\bar{\nu}$  final states*, *Phys. Rev. D* **79** (2009) 012002 [[arXiv:0809.0828](#)] [[INSPIRE](#)].
- [70] BABAR collaboration, B. Aubert et al., *Measurement of  $|V_{cb}|$  and the form-factor slope in  $\bar{B} \rightarrow D\ell^-\bar{\nu}_\ell$  decays in events tagged by a fully reconstructed B meson*, *Phys. Rev. Lett.* **104** (2010) 011802 [[arXiv:0904.4063](#)] [[INSPIRE](#)].
- [71] BELLE collaboration, R. Glattauer, *Semileptonic and leptonic B and  $B_s$  decays at Belle*, talk at the *ICHEP 2014*, Valencia Spain July 2–9 2014.
- [72] M. Okamoto et al., *Semileptonic  $D \rightarrow \pi/K$  and  $B \rightarrow \pi/D$  decays in 2 + 1 flavor lattice QCD*, *Nucl. Phys. Proc. Suppl.* **140** (2005) 461 [[hep-lat/0409116](#)] [[INSPIRE](#)].
- [73] HEAVY FLAVOR AVERAGING GROUP (HFAG) collaboration, Y. Amhis et al., *Averages of b-hadron, c-hadron and  $\tau$ -lepton properties as of summer 2014*, [arXiv:1412.7515](#) [[INSPIRE](#)].
- [74] FERMILAB LATTICE, MILC collaboration, J.A. Bailey et al.,  *$B \rightarrow D^*\ell\nu$  at zero recoil: an update*, *PoS(LATTICE 2010)311* [[arXiv:1011.2166](#)] [[INSPIRE](#)].
- [75] A. Hocker, H. Lacker, S. Laplace and F. Le Diberder, *A new approach to a global fit of the CKM matrix*, *Eur. Phys. J. C* **21** (2001) 225 [[hep-ph/0104062](#)] [[INSPIRE](#)].
- [76] CKMFITTER GROUP collaboration, J. Charles et al., *CP violation and the CKM matrix: assessing the impact of the asymmetric B factories*, *Eur. Phys. J. C* **41** (2005) 1 [[hep-ph/0406184](#)] [[INSPIRE](#)].
- [77] X.-Q. Li, Y.-D. Yang and X.-B. Yuan, *Exclusive radiative B-meson decays within minimal flavor-violating two-Higgs-doublet models*, *Phys. Rev. D* **89** (2014) 054024 [[arXiv:1311.2786](#)] [[INSPIRE](#)].
- [78] C.-F. Qiao, P. Sun, D. Yang and R.-L. Zhu,  *$B_c$  exclusive decays to charmonium and a light meson at next-to-leading order accuracy*, *Phys. Rev. D* **89** (2014) 034008 [[arXiv:1209.5859](#)] [[INSPIRE](#)].
- [79] W. Wang and R.-L. Zhu, *Radiative leptonic  $B_c \rightarrow \gamma\ell\bar{\nu}$  decay in effective field theory beyond leading order*, *Eur. Phys. J. C* **75** (2015) 360 [[arXiv:1501.04493](#)] [[INSPIRE](#)].
- [80] C.-H. Chang, J.-P. Cheng and C.-D. Lu, *Radiative leptonic decays of  $B_c$  meson*, *Phys. Lett. B* **425** (1998) 166 [[hep-ph/9712325](#)] [[INSPIRE](#)].

- [81] G. Chiladze, A.F. Falk and A.A. Petrov, *Radiative leptonic  $B_c$  decays in effective field theory*, *Phys. Rev. D* **60** (1999) 034011 [[hep-ph/9811405](#)] [[INSPIRE](#)].
- [82] P. Colangelo and F. De Fazio, *Radiative leptonic  $B_c$  decays*, *Mod. Phys. Lett. A* **14** (1999) 2303 [[hep-ph/9904363](#)] [[INSPIRE](#)].
- [83] C.C. Lih, C.Q. Geng and W.-M. Zhang, *Study of  $B_{c^+} \rightarrow \ell^+ \nu, \gamma$  decays in the light front model*, *Phys. Rev. D* **59** (1999) 114002 [[INSPIRE](#)].
- [84] C.-H. Chang, C.-D. Lu, G.-L. Wang and H.-S. Zong, *The pure leptonic decays of  $b_c$  meson and their radiative corrections*, *Phys. Rev. D* **60** (1999) 114013 [[hep-ph/9904471](#)] [[INSPIRE](#)].
- [85] N. Barik, S. Naimuddin, P.C. Dash and S. Kar, *Radiative leptonic  $B_c$  decay in the relativistic independent quark model*, *Phys. Rev. D* **78** (2008) 114030 [[INSPIRE](#)].
- [86] V.M. Braun and A. Khodjamirian, *Soft contribution to  $B \rightarrow \gamma \ell \nu_\ell$  and the  $B$ -meson distribution amplitude*, *Phys. Lett. B* **718** (2013) 1014 [[arXiv:1210.4453](#)] [[INSPIRE](#)].
- [87] M. Beneke and J. Rohrwild,  *$B$  meson distribution amplitude from  $B \rightarrow \gamma \ell \nu$* , *Eur. Phys. J. C* **71** (2011) 1818 [[arXiv:1110.3228](#)] [[INSPIRE](#)].
- [88] S. Descotes-Genon and C.T. Sachrajda, *Factorization, the light cone distribution amplitude of the  $B$  meson and the radiative decay  $B \rightarrow \gamma \ell \nu_\ell$* , *Nucl. Phys. B* **650** (2003) 356 [[hep-ph/0209216](#)] [[INSPIRE](#)].
- [89] G.P. Korchemsky, D. Pirjol and T.-M. Yan, *Radiative leptonic decays of  $B$  mesons in QCD*, *Phys. Rev. D* **61** (2000) 114510 [[hep-ph/9911427](#)] [[INSPIRE](#)].
- [90] I.P. Gouz, V.V. Kiselev, A.K. Likhoded, V.I. Romanovsky and O.P. Yushchenko, *Prospects for the  $B_c$  studies at LHCb*, *Phys. Atom. Nucl.* **67** (2004) 1559 [*Yad. Fiz.* **67** (2004) 1581] [[hep-ph/0211432](#)] [[INSPIRE](#)].
- [91] Z. Ligeti and F.J. Tackmann, *Precise predictions for  $B \rightarrow X_c \tau \bar{\nu}$  decay distributions*, *Phys. Rev. D* **90** (2014) 034021 [[arXiv:1406.7013](#)] [[INSPIRE](#)].
- [92] A.F. Falk, Z. Ligeti, M. Neubert and Y. Nir, *Heavy quark expansion for the inclusive decay  $B \rightarrow \tau \bar{\nu} X$* , *Phys. Lett. B* **326** (1994) 145 [[hep-ph/9401226](#)] [[INSPIRE](#)].
- [93] J. Chay, H. Georgi and B. Grinstein, *Lepton energy distributions in heavy meson decays from QCD*, *Phys. Lett. B* **247** (1990) 399 [[INSPIRE](#)].
- [94] M.A. Shifman and M.B. Voloshin, *Preasymptotic effects in inclusive weak decays of charmed particles*, *Sov. J. Nucl. Phys.* **41** (1985) 120 [*Yad. Fiz.* **41** (1985) 187] [[INSPIRE](#)].
- [95] F.U. Bernlochner, Z. Ligeti and S. Turczyk, *A proposal to solve some puzzles in semileptonic  $B$  decays*, *Phys. Rev. D* **85** (2012) 094033 [[arXiv:1202.1834](#)] [[INSPIRE](#)].
- [96] P. Gambino, *Inclusive semileptonic  $B$  decays and  $|V_{cb}|$ : in memoriam Kolya Uraltsev*, *Int. J. Mod. Phys. A* **30** (2015) 1543002 [[arXiv:1501.00314](#)] [[INSPIRE](#)].
- [97] S. Biswas and K. Melnikov, *Second order QCD corrections to inclusive semileptonic  $b \rightarrow X_c \ell \bar{\nu}_\ell$  decays with massless and massive lepton*, *JHEP* **02** (2010) 089 [[arXiv:0911.4142](#)] [[INSPIRE](#)].
- [98] T. Gutsche, M.A. Ivanov, J.G. Körner, V.E. Lyubovitskij, P. Santorelli and N. Haby, *Semileptonic decay  $\Lambda_b \rightarrow \Lambda_c + \tau^- + \bar{\nu}_\tau$  in the covariant confined quark model*, *Phys. Rev. D* **91** (2015) 074001 [*Erratum ibid.* **91** (2015) 119907] [[arXiv:1502.04864](#)] [[INSPIRE](#)].
- [99] LHCb collaboration, *Measurement of  $b$ -hadron production fractions in 7 TeV  $pp$  collisions*, *Phys. Rev. D* **85** (2012) 032008 [[arXiv:1111.2357](#)] [[INSPIRE](#)].



- [100] X.Q. Li, Y.D. Yang and X. Zhang, *The semileptonic  $\Lambda_b \rightarrow \Lambda_c \ell \nu_\ell$  decays with a scalar or vector leptoquark*, in preparation.
- [101] S.J. Brodsky and C.-R. Ji, *Exclusive production of higher generation hadrons and form-factor zeros in quantum chromodynamics*, *Phys. Rev. Lett.* **55** (1985) 2257 [[INSPIRE](#)].
- [102] J.G. Korner and P. Kroll, *Heavy quark symmetry at large recoil*, *Phys. Lett. B* **293** (1992) 201 [[INSPIRE](#)].
- [103] S.Y. Choi and H.S. Song, *Exclusive heavy meson pair production by  $\gamma\gamma$  collision in heavy quark effective theory*, *Phys. Lett. B* **296** (1992) 420 [[hep-ph/9209264](#)] [[INSPIRE](#)].
- [104] C.E. Carlson and J. Milana, *Perturbative QCD calculations of heavy meson exclusive decays*, *Phys. Lett. B* **301** (1993) 237 [[INSPIRE](#)].
- [105] D.-S. Du, X.-L. Li and Y.-D. Yang, *A study on the rare radiative decay  $B_c \rightarrow D_s * \gamma$* , *Phys. Lett. B* **380** (1996) 193 [[hep-ph/9603291](#)] [[INSPIRE](#)].
- [106] A.G. Grozin and M. Neubert, *Asymptotics of heavy meson form-factors*, *Phys. Rev. D* **55** (1997) 272 [[hep-ph/9607366](#)] [[INSPIRE](#)].
- [107] M. Beneke, G. Buchalla, M. Neubert and C.T. Sachrajda, *QCD factorization for exclusive, nonleptonic B meson decays: general arguments and the case of heavy light final states*, *Nucl. Phys. B* **591** (2000) 313 [[hep-ph/0006124](#)] [[INSPIRE](#)].
- [108] I. Caprini, L. Lellouch and M. Neubert, *Dispersive bounds on the shape of  $\bar{B} \rightarrow D(*)\ell\bar{\nu}$  form-factors*, *Nucl. Phys. B* **530** (1998) 153 [[hep-ph/9712417](#)] [[INSPIRE](#)].