

RIEMANNIAN MANIFOLDS WITH DENSE ORBITS UNDER LIE GROUPS OF MOTIONS

SHÔBIN KASHIWABARA

(Received November 24, 1967)

Let M be an n -dimensional connected complete differentiable¹⁾ Riemannian manifold admitting an intransitive effective connected Lie group H of motions²⁾ on M . For any $p \in M$, the differentiable submanifold $H(p) = \{h(p) : h \in H\}$ is usually called the orbit of p under H or the H -orbit of p . On the other hand, the group H can be regarded as an analytic subgroup of the Lie group $I(M)$ of all motions on M and the closure (in $I(M)$) of H forms a subgroup which is a connected Lie group. Such a Lie group we denote by \bar{H} . The closure of an H -orbit consists of one point or has the structure of a regularly imbedded³⁾ connected differentiable submanifold. This follows from the fact that the closure $\overline{H(p)}$, $p \in M$, coincides with the \bar{H} -orbit of p , i.e., $\overline{H(p)} = \bar{H}(p)$. We shall call such a manifold $\overline{H(p)}$ the *closure manifold* of $H(p)$. For any $q \in \overline{H(p)}$, we can see $\overline{H(p)} = \bar{H}(q)$.

In the following, suppose an H -orbit of M is dense in M ⁴⁾. The object of this note is to give geometrical structures of such a manifold M .

Now, as is easily seen, the group \bar{H} acts on M transitively and effectively. It is shown that the group H is normal in \bar{H} . Hence, for any $g \in \bar{H}$ we have $g \cdot H(p) = H(g \cdot p)$, where $p \in M$. Thus the following theorem is obtained:

THEOREM 1. 1) *Every H -orbit is dense in M ,*
2) *any element of \bar{H} carries every H -orbit into an H -orbit, and*
3) *M has the structure of a foliated manifold⁵⁾ with H -orbits as its leaves.*

Goto proved the following theorem (see [1] or [2]), which plays an important role in this note: For a connected Lie group G and its analytic

1) The word "differentiable" means " C^∞ -differentiable".

2) By a motion, we mean an isometry as usual.

3) This means that the topology of the submanifold coincides with the relative one.

4) An orbit is sometimes regarded as a subset of M , as is here the case.

5) For the definition, see [4].

subgroup S which is not closed in G , there exists a 1-parameter subgroup of S whose closure (in G) is not contained in S . We shall call such a subgroup a *Goto's 1-parameter subgroup*.

Next, let G be a connected abelian Lie group and let N be a complete differentiable Riemannian manifold. Then the following facts are well-known :

1) Any invariant differentiable Riemannian metric on G reduces to a euclidean metric (see [3]).

2) If G is a group of motions on N , then the isotropy subgroup G_p at $p \in N$ leaves the orbit $G(p)$ pointwise invariant.

3) If G is a 1-parameter group of motions on N , then the closure manifold of a G -orbit consists of one point, or is homeomorphic to a straight line or a torus of dimension ≥ 1 and the metric induced from N becomes euclidean (see [3]).

Let \bar{H}_p be the isotropy subgroup of \bar{H} at $p \in M$. The set $H \cdot \bar{H}_p$ forms a Lie subgroup of \bar{H} which is also the minimal subgroup containing H and \bar{H}_p . We denote this group by $J_{(p)}$. Then $\bar{H} \supset J_{(p)} \supset H$ and $J_{(p)} = J_{(q)}$ for any $q \in H(p)$. $J_{(p)}$ consists of all the elements of \bar{H} which leaves $H(p)$ invariant. Let $J_{(p)}^0$ denote the identity component of $J_{(p)}$. Then $J_{(p)}^0 \supset H$. By referring to Theorem 1, we see that $J_{(p)}^0$ leaves not only $H(p)$ but also $H(x)$ for any $x \in M$ invariant. Hence $J_{(p)}^0 = J_{(x)}^0$. So, we denote $J_{(p)}^0$ by J^0 hereafter. The closure (in $I(M)$) of J^0 coincides with \bar{H} and J^0 is not closed in \bar{H} . Therefore we have a Goto's 1-parameter subgroup $\gamma \subset J^0$ such that the closure $\bar{\gamma}$ (in \bar{H}) of γ is not contained in J^0 . For any $x \in M$, $\gamma(x) \subset H(x)$ and the closure manifold $\bar{\gamma(x)}$ (in M) of $\gamma(x)$ is not contained in $H(x)$. For, if $\bar{\gamma(x)} \subset H(x)$ we have $\bar{\gamma(x)} = \bar{\gamma(x)} \subset H(x)$. So $\bar{\gamma} \subset J_{(x)}$. This implies $\bar{\gamma} \subset J^0$ which is a contradiction. We can further see that the closure (in $H(x)$) of $\gamma(x)$ coincides with $\gamma(x)$. Otherwise, this closure is homeomorphic to a torus of dimension > 1 , and so compact under the topology of $H(x)$. Hence it would be shown that $H(x)$ contains the closure manifold $\bar{\gamma(x)}$ (in M) of $\gamma(x)$. This is a contradiction as mentioned above. Thus, $H(x)$ has a structure of product bundle with $\gamma(y)$, $y \in H(x)$, as fibers. Summing up these facts, we have

THEOREM 2. *The group \bar{H} has a 1-parameter subgroup γ with the following properties: for any $x \in M$,*

1) $\gamma(x) \subset H(x)$, but the closure manifold $\bar{\gamma(x)}$ (in M) is not included in $H(x)$,

2) $\bar{\gamma(x)}$ is homeomorphic to a torus of dimension > 1 and a euclidean metric is induced from M , and

3) $H(x)$ has a structure of product bundle with $\gamma(y)$, $y \in H(x)$, as fibers.

Further we suppose the group H is abelian. Then so also is the group \bar{H} . The isotropy subgroup \bar{H}_p , $p \in M$, leaves M pointwise invariant and so consists of the identity only, \bar{H} being effective. Hence the group \bar{H} is diffeomorphic to M and the metric on M must be euclidean. Thus we may prove the following

THEOREM 3. *Suppose the group H is abelian. Then,*

- 1) *M is diffeomorphic to the abelian Lie group \bar{H} ,*
 - 2) *The metric on M reduces to euclidean one,*
 - 3) *every H -orbit is totally geodesic,*
- and further, with respect to the 1-parameter subgroup γ in Theorem 2,*
- 4) *M has a structure of fiber bundle with $\overline{\gamma(x)}$, $x \in M$, as fibers, where fibers have euclidean metrics as the induced ones and are isometric to one another, and*
 - 5) *every $\overline{\gamma(x)}$ is totally geodesic.*

BIBLIOGRAPHY

- [1] M. GOTÔ, Faithful representations of Lie groups I, Math. Japonicae, 1(1948), 107-119.
- [2] S. HELGASON, Differential Geometry and Symmetric Spaces, Academic Press, 1962.
- [3] S. KASHIWABARA, A fibering of Riemannian manifolds admitting 1-parameter groups of motions, Tôhoku Math. Journ., 17(1965), 266-270.
- [4] B. L. REINHART, Foliated manifolds with bundle-like metrics, Ann. of Math., 69(1959), 119-132.

DEPARTMENT OF MATHEMATICS
KANAZAWA UNIVERSITY
KANAZAWA, JAPAN