



RIGID ORBITS AND SHEETS IN REDUCTIVE LIE ALGEBRAS OVER FIELDS OF PRIME CHARACTERISTIC

DOI:

[10.1017/S1474748016000086](https://doi.org/10.1017/S1474748016000086)

Document Version

Accepted author manuscript

[Link to publication record in Manchester Research Explorer](#)

Citation for published version (APA):

Premet, A., & Stewart, D. I. (2016). RIGID ORBITS AND SHEETS IN REDUCTIVE LIE ALGEBRAS OVER FIELDS OF PRIME CHARACTERISTIC. *Journal of the Institute of Mathematics of Jussieu*, 17(3), 583-613. <https://doi.org/10.1017/S1474748016000086>

Published in:

Journal of the Institute of Mathematics of Jussieu

Citing this paper

Please note that where the full-text provided on Manchester Research Explorer is the Author Accepted Manuscript or Proof version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version.

General rights

Copyright and moral rights for the publications made accessible in the Research Explorer are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Takedown policy

If you believe that this document breaches copyright please refer to the University of Manchester's Takedown Procedures [<http://man.ac.uk/04Y6Bo>] or contact uml.scholarlycommunications@manchester.ac.uk providing relevant details, so we can investigate your claim.



RIGID ORBITS AND SHEETS IN REDUCTIVE LIE ALGEBRAS OVER FIELDS OF PRIME CHARACTERISTIC

ALEXANDER PREMETS¹ AND DAVID I. STEWART²

¹The University of Manchester, Oxford Road, M13 9PL, UK
(alexander.premet@manchester.ac.uk)

²University of Newcastle, NE1 7RU, UK (dis20@cantab.net)

(Received 26 August 2015; revised 22 February 2016; accepted 22 February 2016)

Abstract Let G be a simple simply connected algebraic group over an algebraically closed field k of characteristic $p > 0$ with $\mathfrak{g} = \text{Lie}(G)$. We discuss various properties of nilpotent orbits in \mathfrak{g} , which have previously only been considered over \mathbb{C} . Using computational methods, we extend to positive characteristic various calculations of de Graaf with nilpotent orbits in exceptional Lie algebras. In particular, we classify those orbits which are reachable as well as those which satisfy a certain related condition due to Panyushev, and determine the codimension of the derived subalgebra $[\mathfrak{g}_e, \mathfrak{g}_e]$ in the centraliser \mathfrak{g}_e of any nilpotent element $e \in \mathfrak{g}$. Some of these calculations are used to show that the list of rigid nilpotent orbits in \mathfrak{g} , the classification of sheets of \mathfrak{g} and the distribution of the nilpotent orbits amongst them are independent of good characteristic, remaining the same as in the characteristic zero case. We also give a comprehensive account of the theory of sheets in reductive Lie algebras over algebraically closed fields of good characteristic.

Keywords: reductive group; decomposition class; induced nilpotent orbit

2010 *Mathematics subject classification:* 17B45; 20G15

1. Introduction

Let G be a reductive algebraic group over an algebraically closed field k of characteristic $p \geq 0$ with $\mathfrak{g} = \text{Lie}(G)$. We consider various properties of a nilpotent element $e \in \mathfrak{g}$ and its centraliser \mathfrak{g}_e . The element e is called *reachable* if $e \in [\mathfrak{g}_e, \mathfrak{g}_e]$. It is called *strongly reachable* if $[\mathfrak{g}_e, \mathfrak{g}_e] = \mathfrak{g}_e$, i.e., the subalgebra \mathfrak{g}_e is perfect. If p is a good prime for G , it is said to satisfy *Panyushev property* if in the associated grading $\mathfrak{g}_e = \bigoplus_{i \geq 0} \mathfrak{g}_e(i)$, the Lie subalgebra $\bigoplus_{i \geq 1} \mathfrak{g}_e(i)$ is generated by $\mathfrak{g}_e(1)$. For the case $p = 0$, in [7] the author applies various routines in the SLA package of GAP in order to confirm and finish the classification of nilpotent elements which are reachable, strongly reachable or satisfy the Panyushev property. In particular, he confirms that in characteristic 0, all reachable elements are Panyushev. For G classical and $p = 0$ each of three properties above characterises the class of rigid (i.e., non-induced) nilpotent orbits in \mathfrak{g} . This was established by Yakimova in [34].

One of the goals of this article is to record extensions to de Graaf's results to deal with the case where $p > 0$. For G exceptional and simply connected (and for any prime p) we compute the number $c(e) := \dim \mathfrak{g}_e / [\mathfrak{g}_e, \mathfrak{g}_e]$ for any nilpotent element $e \in \mathfrak{g}$. This data is recorded in Tables 2 and 3. It turns out that $c(e)$ is independent of p provided that p is good for G . Our computations agree with those in [7] made in the characteristic zero case.¹ We also consider another property, which is relevant in the theory of finite W -algebras. Let us call a nilpotent element *almost reachable* if it is not reachable, but $\mathfrak{g}_e = ke \oplus [\mathfrak{g}_e, \mathfrak{g}_e]$.

If G is SO_n or Sp_n and $p \neq 2$ then [26, Theorem 3(i)] provides a general formula for $c(e)$ in terms of the partition of n attached to e . We mention for completeness that for $G = \mathrm{SL}_n$ or GL_n the value of $c(e)$ is also known for all $e \in \mathfrak{g}$ (and all p) thanks to the explicit description of \mathfrak{g}_e given in [34]; see [26, Remark 1] for more detail. So in our paper we mostly deal with the groups of exceptional types. The main results of the computational part of our paper are as follows:

Theorem 1.1. *Let G be a simple, simply connected algebraic group of exceptional type over an algebraically closed field of characteristic $p > 0$ and let e be a nilpotent element in $\mathfrak{g} = \mathrm{Lie}(G)$. Then e is reachable or almost reachable in characteristic p if and only if the orbit of e is listed in the first column of Table 1. If, moreover, the equality $\mathfrak{g}_e = [\mathfrak{g}_e, \mathfrak{g}_e]$ holds for e in characteristic p then this is indicated in the third column of Table 1.*

Table 1 show that many new nilpotent orbits become reachable in bad characteristic and on the three occasions this happens when p is good for G . In this case, the new orbits are as follows:

- (A) $p = 5$, G is of type E_7 and e is of type $A_3 + A_2 + A_1$;
- (B) $p = 7$, G is of type E_7 and e is of type $A_2 + 3A_1$;
- (C) $p = 7$, G is of type E_8 and e is of type $A_4 + A_2 + A_1$.

It is interesting that these three orbits are responsible for the existence of new maximal Lie subalgebras in $\mathfrak{g} = \mathrm{Lie}(G)$ which have no analogues in characteristic 0. This will be explained in detail in a forthcoming paper by the authors.

We have also checked the validity of Panyushev property in good characteristic; see § 4.5 for more detail on our computations. This property will play an important role in proving Humphreys' conjecture on the existence of $U_\chi(\mathfrak{g})$ -modules of dimension $p^{(\dim G \cdot \chi)/2}$ (here $U_\chi(\mathfrak{g})$ stands for the reduced enveloping algebra of \mathfrak{g} associated with a linear function $\chi \in \mathfrak{g}^*$). We denote by $G_{\mathbb{C}}$ the complex simple algebraic group of the same type as G and let $\mathfrak{g}_{\mathbb{C}} = \mathrm{Lie}(G_{\mathbb{C}})$. Recall that if p is good for G then the nilpotent orbits in \mathfrak{g} have the same labels as those in $\mathfrak{g}_{\mathbb{C}}$; see [6, pp. 401–407], for example.

Theorem 1.2. *Let G be as in Theorem 1.1 and suppose p is a good prime for G . Then an element e in a nilpotent orbit \mathcal{O} satisfies the Panyushev condition if and only if the nilpotent orbit in $\mathfrak{g}_{\mathbb{C}}$ which has the same label as \mathcal{O} consists of reachable elements. In*

¹According to the usual notation one should replace \tilde{A}_1 by A_1 in [7, Table 5].

\mathfrak{g}	e	p	Strong	Almost
G_2	$\tilde{A}_1^{(3)}$	3		
	\tilde{A}_1	2, 3	3	$p \geq 5$
	A_1	Any	$p \geq 5$	
F_4	F_4	3		
	C_3	2		
	$C_3(a_1)$	2		
	$A_1 + \tilde{A}_2$	2, 3		$p \geq 5$
	B_2			$p \geq 3$
	$A_2 + \tilde{A}_1$	$p \geq 3$	$p \geq 5$	
	\tilde{A}_2	2	2	$p \geq 3$
	A_2			$p \geq 3$
	$A_1 + \tilde{A}_1$	Any	$p \geq 3$	
	$\tilde{A}_1^{(2)}$	2		
	\tilde{A}_1	Any	Any	
	A_1	Any	$p \geq 3$	
E_6	E_6	3		
	A_5	2		
	$A_4 + A_1$	2, 3		
	$A_3 + A_1$	2		
	$2A_2 + A_1$	Any	$p \geq 5$	
	$A_2 + 2A_1$	Any		
	$2A_2$	2		
	$A_2 + A_1$	Any		
	A_2			Any
	$3A_1$	Any	$p \geq 3$	
	$2A_1$	Any		
	A_1	Any	Any	
E_7	E_7	3		
	E_6	3		
	A_6	2		
	$A_5 + A_1$	3		$p \geq 5$
	$(A_5)'$	2		$p \geq 3$
	$A_4 + A_2$	2, 3		$p \geq 5$
	$A_4 + A_1$	Any		
	$D_4 + A_1$			$p \geq 3$
	$A_3 + A_2 + A_1$	3, 5		$p \geq 7$
	$A_3 + A_2$	2		
	$A_3 + 2A_1$	3		$p \geq 5$
	$(A_3 + A_1)'$	2		$p \geq 3$
$2A_2 + A_1$	Any	$p \geq 5$		

Table 1: continued

\mathfrak{g}	e	p	Strong	Almost	
E_8	$A_2 + 3A_1$	7		$p \neq 2, 7$	
	$2A_2$	2		$p \geq 3$	
	A_3			$p \geq 3$	
	$A_2 + 2A_1$	Any	$p \geq 3$		
	$A_2 + A_1$	Any			
	$4A_1$	$p \geq 3$	$p \geq 3$		
	A_2			Any	
	$(3A_1)'$	Any	$p \geq 3$		
	$(3A_1)''$	3		$p \geq 5$	
	$2A_1$	Any	$p \geq 3$		
	A_1	Any	Any		
	E_8	E_8	3, 5		
	$E_8(a_1)$	$E_8(a_1)$	3		
	E_7	E_7	3		
	D_7	D_7	2		
	$E_6 + A_1$	$E_6 + A_1$	3		
	$D_7(a_1)$	$D_7(a_1)$	2		
	$E_8(b_6)$	$E_8(b_6)$	3		
	$(A_7)^{(3)}$	$(A_7)^{(3)}$	3		
	A_7	A_7	2, 3		$p \geq 5$
	$D_7(a_2)$	$D_7(a_2)$	2		
	E_6	E_6	3		
	$A_6 + A_1$	$A_6 + A_1$	3, 5		$p \geq 7$
	A_6	A_6	2		
	$D_6(a_2)$	$D_6(a_2)$	2		
	$D_5(a_1) + A_2$	$D_5(a_1) + A_2$	2		$p \geq 5$
	$A_5 + A_1$	$A_5 + A_1$	2, 3		$p \geq 5$
	$A_4 + A_3$	$A_4 + A_3$	Any	$p \geq 7$	
	$D_4 + A_2$	$D_4 + A_2$	2		
	$A_4 + A_2 + A_1$	$A_4 + A_2 + A_1$	7		$p \neq 2, 5, 7$
	$D_5(a_1) + A_1$	$D_5(a_1) + A_1$			$p \geq 3$
	A_5	A_5	2		$p \geq 3$
	$A_4 + A_2$	$A_4 + A_2$	2, 3		$p \geq 5$
	$A_4 + 2A_1$	$A_4 + 2A_1$	Any		
$2A_3$	$2A_3$	Any	$p \geq 3$		
$A_4 + A_1$	$A_4 + A_1$	Any			
$D_4(a_1) + A_2$	$D_4(a_1) + A_2$	2		$p \geq 5$	
$D_4 + A_1$	$D_4 + A_1$			$p \geq 3$	

Table 1: continued

\mathfrak{g}	e	p	Strong	Almost
	$A_3 + A_2 + A_1$	$p \geq 3$	$p \geq 3$	
	$A_3 + A_2$	2		
	$D_4(a_1) + A_1$	$p \geq 3$	$p \geq 3$	
	$A_3 + 2A_1$	Any	$p \geq 3$	
	$2A_2 + 2A_1$	Any	$p \geq 5$	
	$A_3 + A_1$	2		$p \geq 3$
	$2A_2 + A_1$	Any	$p \geq 5$	
	$2A_2$	2		$p \geq 3$
	$A_2 + 3A_1$	Any	$p \geq 3$	
	A_3			Any
	$A_2 + 2A_1$	Any	$p \geq 3$	
	$A_2 + A_1$	Any	Any	
	$4A_1$	Any	$p \geq 3$	
	A_2			Any
	$3A_1$	Any	$p \geq 3$	
	$2A_1$	Any	Any	
	A_1	Any	Any	

TABLE 1. Strongly reachable and almost reachable orbits for Strongly reachable and almost reachable orbits for G_2, F_4, E_6, E_7 and E_8 .

other words, e satisfies the Panyushev condition if and only if it is reachable and not listed in cases (A), (B) or (C) above. 73
74

It follows from Theorems 1.1 and 1.2 that the Panyushev and strong reachability conditions are independent of good characteristic. 75
76

A nilpotent element $e \in \mathfrak{g}$ is called *rigid* if it cannot be obtained by Lusztig–Spaltenstein induction from a proper Levi subalgebra of \mathfrak{g} . Arguing as in [25, 3.2] one can reduce proving Humphreys’ conjecture to the case where $\chi \in \mathfrak{g}^*$ corresponds to a rigid nilpotent element of \mathfrak{g} under a G -equivariant isomorphism $\mathfrak{g} \cong \mathfrak{g}^*$. In the characteristic zero case, all rigid nilpotent orbits in exceptional Lie algebras are classified by Elashvili [10] and his computations are recently double-checked in [8] by using GAP. Since the way this is done in [8] relies heavily on the characteristic 0 hypothesis, we present in § 2 a classification of rigid nilpotent orbits in exceptional Lie algebras valid over any algebraically closed field of good characteristic (at some point we have to rely on GAP as well). 77
78
79
80
81
82
83
84
85

Theorem 1.3. *Let G be as in Theorem 1.1 and suppose p is a good prime for G . Then a nilpotent orbit \mathcal{O} is rigid in \mathfrak{g} if and only if so is the nilpotent orbit in $\mathfrak{g}_{\mathbb{C}}$ which has the same label as \mathcal{O} .* 86
87
88

Given $m \in \mathbb{N}$ we let $\mathfrak{g}_{(m)}$ denote the set of all elements of \mathfrak{g} whose adjoint G -orbit has codimension m in \mathfrak{g} . A subset $\mathcal{S} \subset \mathfrak{g}$ is called a *sheet* of \mathfrak{g} if it coincides with an irreducible component of one of the quasi-affine varieties $\mathfrak{g}_{(m)}$. According to a classical 89
90

Orbit in E_8	$\dim(\mathfrak{g}_e/[\mathfrak{g}_e, \mathfrak{g}_e])$ $p = 2, 3, 5, \geq 7$	Orbit in E_8	$\dim(\mathfrak{g}_e/[\mathfrak{g}_e, \mathfrak{g}_e])$ $p = 2, 3, 5, \geq 7$
E_8	12, 6, 3, 8	$D_5(a_1) + A_2$	5, 3, 1, 1
$E_8(a_1)$	12, 6, 9, 7	$A_5 + A_1$	5, 2, 1, 1
$E_8(a_2)$	12, 10, 8, 6	$A_4 + A_3$	5, 2, 2, 0
$E_8(a_3)$	12, 5, 7, 7	D_5	4, 3, 3, 3
$E_8(a_4)$	14, 8, 6, 6	$E_6(a_3)$	4, 3, 3, 3
E_7	11, 3, 4, 4	$(D_4 + A_2)^{(2)}$	3, -, -, -
$E_8(b_4)$	11, 6, 5, 5	$D_4 + A_2$	9, 2, 2, 2
$E_8(a_5)$	11, 6, 5, 5	$A_4 + A_2 + A_1$	3, 1, 3, 1
$E_7(a_1)$	11, 5, 5, 5	$D_5(a_1) + A_1$	4, 1, 1, 1
$E_8(b_5)$	11, 7, 7, 7	A_5	3, 1, 1, 1
$(D_7)^{(2)}$	11, -, -, -	$A_4 + A_2$	4, 1, 1, 1
D_7	10, 2, 2, 2	$A_4 + 2A_1$	1, 1, 1, 1
$E_8(a_6)$	11, 6, 6, 6	$D_5(a_1)$	2, 2, 2, 2
$E_7(a_2)$	6, 6, 4, 4	$2A_3$	9, 0, 0, 0
$E_6 + A_1$	6, 5, 2, 2	$A_4 + A_1$	1, 1, 1, 1
$(D_7(a_1))^{(2)}$	6, -, -, -	$D_4(a_1) + A_2$	9, 2, 1, 1
$D_7(a_1)$	10, 4, 4, 4	$D_4 + A_1$	9, 1, 1, 1
$E_8(b_6)$	7, 5, 5, 5	$(A_3 + A_2)^{(2)}$	9, -, -, -
$E_7(a_3)$	8, 4, 4, 4	$A_3 + A_2 + A_1$	9, 0, 0, 0
$E_6(a_1) + A_1$	3, 7, 3, 3	A_4	2, 2, 2, 2
$(A_7)^{(3)}$	-, 6, -, -	$A_3 + A_2$	4, 2, 2, 2
A_7	10, 3, 1, 1	$D_4(a_1) + A_1$	9, 0, 0, 0
$D_7(a_2)$	11, 3, 3, 3	$A_3 + 2A_1$	4, 0, 0, 0
E_6	5, 3, 4, 4	$2A_2 + 2A_1$	4, 4, 0, 0
D_6	10, 2, 2, 2	D_4	3, 2, 2, 2
$(D_5 + A_2)^{(2)}$	10, -, -, -	$D_4(a_1)$	3, 3, 3, 3
$D_5 + A_2$	10, 3, 3, 3	$A_3 + A_1$	2, 1, 1, 1
$E_6(a_1)$	6, 5, 4, 4	$2A_2 + A_1$	2, 2, 0, 0
$E_7(a_4)$	10, 4, 3, 3	$2A_2$	1, 1, 1, 1
$A_6 + A_1$	10, 3, 1, 1	$A_2 + 3A_1$	2, 0, 0, 0
$D_6(a_1)$	10, 3, 3, 3	A_3	1, 1, 1, 1
$(A_6)^{(2)}$	10, -, -, -	$A_2 + 2A_1$	1, 0, 0, 0
A_6	5, 2, 2, 2	$A_2 + A_1$	0, 0, 0, 0
$E_8(a_7)$	10, 10, 10, 10	$4A_1$	8, 0, 0, 0
$D_5 + A_1$	5, 2, 2, 2	A_2	1, 1, 1, 1
$E_7(a_5)$	5, 6, 6, 6	$3A_1$	2, 0, 0, 0
$E_6(a_3) + A_1$	5, 3, 3, 3	$2A_1$	0, 0, 0, 0
$D_6(a_2)$	5, 3, 3, 3	A_1	0, 0, 0, 0

Table 2: continued

Orbit in E_7	$\dim(\mathfrak{g}_e/[\mathfrak{g}_e, \mathfrak{g}_e])$ $p = 2, 3, \geq 5$	Orbit in E_7	$\dim(\mathfrak{g}_e/[\mathfrak{g}_e, \mathfrak{g}_e])$ $p = 2, 3, \geq 5$
E_7	10, 4, 7	$(A_5)''$	4, 3, 3
$E_7(a_1)$	10, 8, 6	$A_3 + A_2 + A_1$	8, 1, 1
$E_7(a_2)$	11, 7, 5	A_4	5, 3, 3
$E_7(a_3)$	13, 6, 6	$(A_3 + A_2)^{(2)}$	8, —, —
E_6	11, 3, 4	$A_3 + A_2$	9, 2, 2
$E_6(a_1)$	13, 6, 5	$D_4(a_1) + A_1$	9, 2, 2
D_6	9, 3, 3	D_4	9, 2, 2
$E_7(a_4)$	9, 5, 4	$A_3 + 2A_1$	4, 1, 1
$D_6(a_1)$	9, 4, 4	$D_4(a_1)$	9, 3, 3
$D_5 + A_1$	10, 3, 3	$(A_3 + A_1)'$	4, 1, 1
$(A_6)^{(2)}$	9, —, —	$2A_2 + A_1$	4, 2, 0
A_6	10, 2, 2	$(A_3 + A_1)''$	3, 2, 2
$E_7(a_5)$	10, 6, 6	$A_2 + 3A_1$	2, 1, 1
D_5	10, 3, 3	$2A_2$	3, 1, 1
$E_6(a_3)$	10, 3, 3	A_3	3, 1, 1
$D_6(a_2)$	5, 3, 3	$A_2 + 2A_1$	3, 0, 0
$D_5(a_1) + A_1$	6, 2, 2	$A_2 + A_1$	1, 1, 1
$A_5 + A_1$	5, 3, 1	$4A_1$	7, 0, 0
$(A_5)'$	5, 1, 1	A_2	1, 1, 1
$A_4 + A_2$	6, 4, 1	$(3A_1)'$	8, 0, 0
$D_5(a_1)$	7, 3, 3	$(3A_1)''$	2, 1, 1
$A_4 + A_1$	2, 2, 2	$2A_1$	2, 0, 0
$D_4 + A_1$	8, 1, 1	A_1	0, 0, 0

 TABLE 2. Codimension of $[\mathfrak{g}_e, \mathfrak{g}_e]$ in \mathfrak{g}_e for E_7 and E_8 .

result of Borho [3] the sheets of $\mathfrak{g}_{\mathbb{C}}$ are parametrised by the $G_{\mathbb{C}}$ -conjugacy classes of pairs $(\mathfrak{l}_{\mathbb{C}}, \mathcal{O}_0)$ where $\mathfrak{l}_{\mathbb{C}}$ is a Levi subalgebra of $\mathfrak{g}_{\mathbb{C}}$ and \mathcal{O}_0 is a rigid nilpotent orbit in \mathfrak{l} . In § 2, we give a comprehensive account of the theory of sheets in reductive Lie algebras $\mathfrak{g} = \text{Lie}(G)$ which satisfy the standard hypotheses (when G is simple and not of type $A_{r,p-1}$ this is equivalent to saying that p is a good prime for G). In particular, we show that every sheet of \mathfrak{g} contains a unique nilpotent orbit and Borho's classification of sheets remains valid under our assumptions on G .

For any sheet \mathcal{S} of a complex exceptional Lie algebra $\mathfrak{g}_{\mathbb{C}} = \text{Lie}(G_{\mathbb{C}})$, de Graaf and Elashvili determine the weighted Dynkin diagram of the unique nilpotent orbit $\mathcal{N}(\mathfrak{g}_{\mathbb{C}}) \cap \mathcal{S}$ and give a very nice representative $e_{\Gamma} = \sum_{\gamma \in \Gamma} e_{\gamma}$ in $\mathcal{N}(\mathfrak{g}_{\mathbb{C}}) \cap \mathcal{S}$ compatible with the combinatorial data that defines \mathcal{S} . Here $\Gamma = \Gamma(\mathcal{S})$ is a subset in the root system Φ of $\mathfrak{g}_{\mathbb{C}}$ and e_{γ} is a root vector of $\mathfrak{g}_{\mathbb{C}}$ corresponding to $\gamma \in \Phi$. Each set Γ is linearly independent in the vector space $\mathbb{Q}\Phi$ and the $G_{\mathbb{C}}$ -orbit of e_{Γ} is independent of the choices of root vectors $e_{\gamma} \in (\mathfrak{g}_{\mathbb{C}})_{\gamma}$. There is a natural way to attach to Γ a graph $D(\Gamma)$, and it turned

Orbit in E_6	$\dim(\mathfrak{g}_e/[\mathfrak{g}_e, \mathfrak{g}_e])$ $p = 2, 3, \geq 5$	Orbit in E_6	$\dim(\mathfrak{g}_e/[\mathfrak{g}_e, \mathfrak{g}_e])$ $p = 2, 3, \geq 5$
E_6	5, 4, 6	$D_4(a_1)$	5, 5, 5
$E_6(a_1)$	7, 8, 5	$A_3 + A_1$	3, 4, 2
D_5	4, 8, 4	$2A_2 + A_1$	3, 3, 0
$E_6(a_3)$	6, 6, 5	A_3	2, 2, 2
$D_5(a_1)$	3, 5, 3	$A_2 + 2A_1$	1, 5, 1
A_5	4, 3, 2	$2A_2$	2, 3, 2
$A_4 + A_1$	5, 4, 2	$A_2 + A_1$	1, 3, 1
D_4	3, 2, 2	A_2	1, 1, 1
A_4	3, 3, 3	$3A_1$	2, 0, 0
		$2A_1$	1, 1, 1
		A_1	0, 0, 0
Orbit in F_4	$\dim(\mathfrak{g}_e/[\mathfrak{g}_e, \mathfrak{g}_e])$ $p = 2, 3, \geq 5$	Orbit in F_4	$\dim(\mathfrak{g}_e/[\mathfrak{g}_e, \mathfrak{g}_e])$ $p = 2, 3, \geq 5$
F_4	5, 3, 4	$\tilde{A}_2 + A_1$	4, 2, 1
$F_4(a_1)$	5, 5, 4	$(B_2)^{(2)}$	7, -, -
$F_4(a_2)$	7, 4, 3	B_2	6, 1, 1
$(C_3)^{(2)}$	8, -, -	$A_2 + \tilde{A}_1$	7, 2, 0
C_3	4, 2, 2	\tilde{A}_2	0, 1, 1
B_3	5, 2, 2	$(A_2)^{(2)}$	7, -, -
$F_4(a_3)$	7, 6, 6	A_2	7, 1, 1
$C_3(a_1)^{(2)}$	8, -, -	$A_1 + \tilde{A}_1$	4, 0, 0
$C_3(a_1)$	4, 3, 3	$(\tilde{A}_1)^{(2)}$	6, -, -
$(\tilde{A}_2 + A_1)^{(2)}$	8, -, -	\tilde{A}_1	0, 0, 0
		A_1	6, 0, 0
Orbit in G_2	$\dim(\mathfrak{g}_e/[\mathfrak{g}_e, \mathfrak{g}_e])$ $p = 2, 3, \geq 5$		
G_2	3, 3, 2		
$G_2(a_1)$	3, 3, 3		
$(\tilde{A}_1)^{(3)}$	-, 3, -		
\tilde{A}_1	2, 0, 1		
A_1	2, 2, 0		

TABLE 3. Codimension of $[\mathfrak{g}_e, \mathfrak{g}_e]$ in \mathfrak{g}_e for G_2 , F_4 and E_6 .

104 out (not surprisingly) that in many cases the graphs thus obtained are admissible in the
105 sense of Carter; cf. [5] and the last column of the tables in [8].

106 Since we have a natural analogue of e_Γ in $\mathfrak{g} = \text{Lie}(G)$, where G is a simple algebraic
107 k -group of the same type as $G_{\mathbb{C}}$, and the discussion above indicates that there is a natural
108 bijection between the sheets of \mathfrak{g} and $\mathfrak{g}_{\mathbb{C}}$, one wonders whether a k -analogue of e_Γ still
109 belongs to the sheet of \mathfrak{g} given by the same data as \mathcal{S} . We verify that for exceptional
110 groups this is indeed the case and from the validity of the representatives of [8] in good
111 characteristic we then deduce the following:

Theorem 1.4. *If G is an exceptional algebraic k -group and $p = \text{char}(k)$ is a good prime for G , then the distribution of nilpotent orbits amongst the sheets of \mathfrak{g} agrees with that of $\mathfrak{g}_{\mathbb{C}}$ and can be read off from the tables in [8].*

For many elements e_{Γ} our proof of Theorem 1.4 is computer-free. For example, this is the case when $D(\Gamma)$ is a disjoint union of Dynkin graphs. However, at the end of the day we do rely on GAP (as did de Graaf and Elashvili) and we have decided to run our programme on all elements e_{Γ} in order to obtain an independent confirmation of the computations in [8]. At some point in the proof we have to show that the adjoint orbits of $e_{\Gamma} \in \mathfrak{g}$ and of its counterpart $e_{\Gamma, \mathbb{C}} \in \mathfrak{g}_{\mathbb{C}}$ have the same Dynkin labels. We deduce this by asking GAP to determine the Jordan block structure of each element $\text{ad } e_{\Gamma}$ and a chosen representative of each nilpotent orbit in $\text{ad } \mathfrak{g}$. Since it turned out, for G exceptional, that the Jordan block decomposition of $\text{ad } e$ identifies the orbit of e almost uniquely (except when $p = 7$ where the Jordan block decompositions of nilpotent elements of type B_3 and C_3 in Lie algebras of type F_4 are the same), this enabled us to determine the Dynkin label of e_{Γ} and finish the proof of Theorem 1.4.

2. Sheets and induced nilpotent orbits in good characteristic

2.1. The standard hypotheses

Let G be a connected reductive group over an algebraically closed field k of characteristic $p > 0$ and $\mathfrak{g} = \text{Lie}(G)$. Being the Lie algebra of an algebraic k -group, \mathfrak{g} carries a canonical p th power map $x \mapsto x^{[p]}$ equivariant under the adjoint action of G . Given $x \in \mathfrak{g}$ we denote by \mathfrak{g}_x (respectively, G_x), the centraliser of x in \mathfrak{g} (respectively, G). In order to apply [23, Theorem A] to all Levi subalgebras of \mathfrak{g} we shall assume, unless otherwise specified, that p is a good prime for G , the derived subgroup of G is simply connected and \mathfrak{g} admits an $(\text{Ad } G)$ -invariant non-degenerate symmetric bilinear form. We fix such a form and call it κ .

The above conditions on G are often referred as the *standard hypotheses*. If they hold then for any $x \in G$ we have the equality $\mathfrak{g}_x = \text{Lie}(G_x)$ (the latter is sometimes expressed by saying that the scheme-theoretic centraliser of e in G is smooth). One also knows that if G satisfies the standard hypotheses then the centraliser of any semi-simple element of the restricted Lie algebra \mathfrak{g} is a Levi subalgebra of \mathfrak{g} . It is worth remarking that any simple, simply connected algebraic k -group of type other than A_{rp-1} , where $p = \text{char}(k)$, satisfies the standard hypotheses if and only if p is a good prime for G .

If G satisfies the standard hypotheses then so does any Levi subgroup L of G . It is well known that up to conjugacy any such subgroup is associated with a subset of a chosen basis of simple roots of G . Let Φ be root system of G with respect to a maximal torus T of G and let Π be a basis of simple roots in Φ . Let $\mathfrak{t} = \text{Lie}(T)$. If L is the standard Levi subgroup of G associated with a subset Π_0 of Π and $\mathfrak{l} = \text{Lie}(L)$ then $\mathfrak{t}_0 := [\mathfrak{l}, \mathfrak{l}] \cap \mathfrak{t}$ has dimension equal to $\dim \mathfrak{t} - |\Pi_0|$ (this follows from the fact that the derived subgroup of L is simply connected). On the other hand, the orthogonal complement $\mathfrak{t}_0^{\perp} := \{t \in \mathfrak{t} \mid \kappa(t, \mathfrak{t}_0) = 0\}$ contains $\mathfrak{z}(\mathfrak{l})$ which forces $\dim \mathfrak{z}(\mathfrak{l}) \leq \dim \mathfrak{t}_0^{\perp} = \dim \mathfrak{t} - |\Pi_0|$. Since $\text{Lie}(Z(L)) \subseteq \mathfrak{z}(\mathfrak{l})$ has dimension equal to $\dim \mathfrak{t} - |\Pi_0|$ this shows that $\mathfrak{z}(\mathfrak{l}) = \text{Lie}(Z(L))$ for any Levi subgroup L of G . Furthermore, if Π_0 has no components of type A_{rp-1} then the structure theory of reductive Lie algebras yields $\mathfrak{z}([\mathfrak{l}, \mathfrak{l}]) = 0$ implying $\mathfrak{l} = [\mathfrak{l}, \mathfrak{l}] \oplus \mathfrak{z}(\mathfrak{l})$.

Let G_1, \dots, G_s be the simple components of G and $\mathfrak{g}_i = \text{Lie}(G_i)$. Let $T_i = T \cap G_i$ and $\mathfrak{t}_i = \text{Lie}(T_i)$. If G_i is not of type A_{rp-1} then it is well known that \mathfrak{g}_i is a simple Lie algebra and the restriction of κ to \mathfrak{g}_i is non-degenerate. If G_l has type $A_{m(l)}$ with $p|(m(l)+1)$ then $\mathfrak{g}_l \cong \mathfrak{sl}_{m(l)+1}$ as restricted Lie algebras and the restriction of κ to \mathfrak{g}_l coincides with a non-zero scalar multiple of the trace form, κ' , of the standard (vector) representation of G_l . Let $\alpha_1, \dots, \alpha_{m(l)}$ be a basis of simple roots of the root system of type $A_{m(l)}$ in Bourbaki's numbering. Then \mathfrak{t}_l has basis $h_1, \dots, h_{m(l)}$ such that $\alpha_i(h_i) = 2 = \kappa'(h_i, h_i)$ for all i and we also have that $\kappa'(h_i, h_{i+1}) = -1$ for $1 \leq i \leq m(l) - 1$, $\kappa'(h_i, h_j) = 0$ for $j \notin \{i-1, j, i+1\}$, and $h_i = [e_{\alpha_i}, e_{-\alpha_i}]$ for some root vectors $e_{\pm\alpha_i} \in \mathfrak{g}_l$ with $\kappa'(e_{\alpha_i}, e_{-\alpha_i}) = 1$. Suppose $\kappa|_{\mathfrak{g}_l} = b \cdot \kappa'$ where $b = b(l) \in k^\times$. Since the restriction of κ to \mathfrak{t} is non-degenerate, there exists an element $h_0 \in \mathfrak{t}$ orthogonal to all \mathfrak{g}_i with $i \neq m(l)$ and such that $\kappa(h_0, h_1) = b$ and $\kappa(h_0, h_i) = 0$ for $2 \leq i \leq m(l)$. In general, h_0 is not unique and we only know that $\kappa(h_0, h_0) = a$ for some $a = a(l) \in k$.

Let C_n denote the Cartan matrix of the root system of type A_n with entries reduced modulo p , so that $\det(C_n) = n+1 \pmod{p}$. Then the restriction of κ to the k -span of $h_0, h_1, \dots, h_{m(l)}$ is represented by the matrix

$$A = \begin{bmatrix} a & b & 0 & \cdots & 0 & 0 \\ b & 2b & -b & \cdots & 0 & 0 \\ 0 & -b & 2b & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \cdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 2b & -b \\ 0 & 0 & 0 & \cdots & -b & 2b \end{bmatrix}$$

of order $m(l)+1$ which is non-singular because $\det(A) = ab^{m(l)} \det(C_{m(l)}) - b^{m(l)+1} \det(C_{m(l)-1}) = b^{m(l)+1} \neq 0$ by the first row Laplace expansion. In particular, this shows that $h_0, h_1, \dots, h_{m(l)}$ are linearly independent. The \mathfrak{g} -invariance of κ yields

$$\begin{aligned} b &= \kappa(h_0, h_1) = \kappa([h_0, e_{\alpha_1}], e_{-\alpha_1}) = (d\alpha_1)_e(h_0)\kappa(e_{\alpha_1}, e_{-\alpha_1}) = b \cdot (d\alpha_1)_e(h_0), \\ 0 &= \kappa(h_0, h_i) = \kappa([h_0, e_{\alpha_i}], e_{-\alpha_i}) = (d\alpha_i)_e(h_0)\kappa(e_{\alpha_i}, e_{-\alpha_i}), \quad 2 \leq i \leq m(l). \end{aligned}$$

It follows that $[h_0, e_{\pm\alpha_1}] = \pm e_{\alpha_1}$ and $[h_0, e_{\pm\alpha_i}] = 0$ for $2 \leq i \leq m(l)$.

We now set $\tilde{\mathfrak{g}}_l := kh_0 \oplus \mathfrak{g}_l$ if G_l is of type A_{rp-1} and $\tilde{\mathfrak{g}}_l := \mathfrak{g}_l$ otherwise. It is immediate from above discussion that if G_l is of type A_{rp-1} then $\tilde{\mathfrak{g}}_l$ is an $(\text{Ad } G)$ -stable ideal of \mathfrak{g} isomorphic to \mathfrak{gl}_{rp} as an abstract Lie algebra and the restriction of κ to $\tilde{\mathfrak{g}}_l$ is non-degenerate. Moreover, the adjoint action of G_l on $\tilde{\mathfrak{g}}_l$ is induced by that of GL_{rp} . Since $\kappa|_{\tilde{\mathfrak{g}}_l}$ is non-degenerate for all l we can attach h_0 's to different components of type A_{rp-1} in such a way that they form an orthogonal set. Note also that the condition $\kappa(h_0, \mathfrak{g}_i) = 0$ forces $[h_0, \mathfrak{g}_i] = 0$ due to the \mathfrak{g} -invariance of κ . We thus deduce that $\mathfrak{g} = \tilde{\mathfrak{g}}_1 \oplus \cdots \oplus \tilde{\mathfrak{g}}_s \oplus \mathfrak{z}$ where \mathfrak{z} is a central subalgebra of \mathfrak{g} contained in \mathfrak{t} and all direct summands in this decomposition are $(\text{Ad } G)$ -stable (in [13, 2.9] this result is mentioned without a proof).

Of course, the above has bearing on the description of the sheets of \mathfrak{g} which turns out to be the same as in the characteristic 0 case. Since we shall require some rather detailed results on sheets of \mathfrak{g} and they are hard to find in the literature under our assumption on G , some short proofs will be given below.

2.2. Sheets and decomposition classes

Given $m \in \mathbb{Z}_{\geq 0}$ we denote by $\mathfrak{g}_{(m)}$ the set of all $x \in \mathfrak{g}$ for which $\dim G_x = m$. Each subset $\mathfrak{g}_{(m)}$ is locally closed in the Zariski topology of \mathfrak{g} and the irreducible components of the $\mathfrak{g}_{(m)}$'s are called the *sheets* of \mathfrak{g} .

Remark 2.1. In prime characteristic there are two meaningful ways to define sheets of \mathfrak{g} and \mathfrak{g}^* . Given $m \in \mathbb{Z}_{\geq 0}$ we let $\mathfrak{g}_{[m]}$ (respectively, $\mathfrak{g}_{[m]}^*$) denote the subset of \mathfrak{g} (respectively, \mathfrak{g}^*) consisting of all elements whose stabiliser in \mathfrak{g} has dimension m . We call the irreducible components of the quasi-affine varieties $\mathfrak{g}_{[m]}$ (respectively, $\mathfrak{g}_{[m]}^*$) the *infinitesimal sheets* of \mathfrak{g} (respectively, \mathfrak{g}^*). All infinitesimal sheets are G -stable, quasi-affine varieties. If G satisfies the standard hypotheses all these notions coincide, but in general there will be some subtle differences well worth investigating. We call a G -orbit \mathfrak{g} (respectively, \mathfrak{g}^*) *infinitesimally isolated* if it forms a single infinitesimal sheet in \mathfrak{g} (respectively, \mathfrak{g}^*). The problem of classifying all infinitesimally isolated nilpotent orbits in reductive Lie algebras is interesting and wide open at the moment. As an example, if k has characteristic p and $\mathfrak{g} = \mathfrak{pgl}_p$ then the regular nilpotent orbit \mathcal{O}_{reg} in \mathfrak{g} coincides with $\mathfrak{g}_{[p]}$ and therefore forms a single infinitesimal sheet of \mathfrak{g} . This can be deduced from the following elementary result of linear algebra: if $X, Y \in \mathfrak{gl}_p$ and $[X, Y] = I_p$ then both X and Y have a single Jordan block of size p and hence are mapped by the canonical homomorphism $\mathfrak{gl}_p \rightarrow \mathfrak{g}$ to a commuting pair of elements in \mathcal{O}_{reg} . So, in contrast with the classical case the regular nilpotent orbit in \mathfrak{pgl}_p is infinitesimally isolated. On the other hand, if $\mathfrak{g} = \mathfrak{sl}_p$ then $\mathfrak{g}_{[p]}$ consists of all traceless $p \times p$ matrices X for which k^p is an indecomposable $k[X]$ -module. From this it follows that $\mathfrak{g}_{[p]}$ is a single infinitesimal sheet of \mathfrak{g} , but $\mathcal{O}_{\text{reg}} \subset \mathfrak{g}_{[p]}$ is not infinitesimally isolated in \mathfrak{g} . In fact, there are no infinitesimally isolated orbits in $\mathfrak{g} = \mathfrak{sl}_p$ at all, because $I_p \in \mathfrak{g}$ and all sheets of \mathfrak{g} are invariant under the affine translations $x \mapsto x + \alpha I_p$ where $\alpha \in k$. Finally, we mention that the study of infinitesimal sheets in \mathfrak{g}^* finds interesting applications in the modular representation theory of reductive Lie algebras; see [24, Remark 5.6].

Remark 2.1 indicates that one cannot expect a uniform behaviour of sheets for all reductive Lie algebras over algebraically closed fields (even in good characteristic) and partially justifies the assumptions we imposed on \mathfrak{g} .

If \mathfrak{l} is a Levi subalgebra of \mathfrak{g} then its centre $\mathfrak{z}(\mathfrak{l})$ is a toral subalgebra of \mathfrak{g} and $\mathfrak{l} = \mathfrak{c}_{\mathfrak{g}}(\mathfrak{z}(\mathfrak{l}))$. We denote by $\mathfrak{z}(\mathfrak{l})_{\text{reg}}$ the set of all $t \in \mathfrak{z}(\mathfrak{l})$ such that $\mathfrak{l} = \mathfrak{c}_{\mathfrak{g}}(t)$. This is a non-empty Zariski open subset of $\mathfrak{z}(\mathfrak{l})$. Given a nilpotent element $e_0 \in \mathfrak{l}$ we set

$$\mathcal{D}(\mathfrak{l}, e_0) := (\text{Ad } G) \cdot (e_0 + \mathfrak{z}(\mathfrak{l})_{\text{reg}})$$

and we call $\mathcal{D}(\mathfrak{l}, e)$ the *decomposition class* of \mathfrak{g} associated with the pair (\mathfrak{l}, e_0) . If $x = x_s + x_n$ is the Jordan decomposition of $x \in \mathcal{D}(\mathfrak{l}, e_0)$ then x_s (respectively, x_n) is G -conjugate to an element of $\mathfrak{z}(\mathfrak{l})_{\text{reg}}$ (respectively, e_0). It follows that $\dim \mathfrak{g}_x = \dim \mathfrak{l}_{e_0}$ for every such x . Since there are finitely many nilpotent L -orbits in $\mathfrak{l} = \text{Lie}(L)$ and the number of G -conjugacy classes of Levi subalgebras of \mathfrak{g} is finite, \mathfrak{g} contains finitely many decomposition classes. Each of them is an irreducible, locally closed subset of \mathfrak{g} contained in one of the $\mathfrak{g}_{(r)}$'s and hence lies in a sheet of \mathfrak{g} . As the sheets are irreducible and their union is the whole of \mathfrak{g} , there can be only finitely many, and so every sheet contains a unique open decomposition class of \mathfrak{g} .

234 *Remark 2.2.* Since the number of nilpotent orbits is finite in all characteristics, the above
 235 discussion is essentially valid for all reductive Lie algebras over algebraically closed fields.
 236 However, in bad characteristic the notion of a decomposition class in \mathfrak{g} will involve the
 237 centraliser in G of a semi-simple element of \mathfrak{g} which is not always a Levi subgroup of G .
 238 In general, the connected component of that group is G -conjugate to a subgroup of G
 239 obtained by base change from some standard *pseudo-Levi* subgroup of $G_{\mathbb{C}}$.

240 2.3. The Zariski closure of a decomposition class

241 Borho's classification of sheets of \mathfrak{g} is based on an important result of Borho–Kraft which
 242 provides a description of the Zariski closure of a decomposition class in \mathfrak{g} ; see [1, Theorem
 243 5.4]. Since a modular version of this result valid under our assumptions on G seems to
 244 be missing in the literature, a short proof will be given below.

245 Let $\mathcal{D}(\mathfrak{l}, e_0)$ be a decomposition class of \mathfrak{g} and let $\lambda \in X_*(L)$ be an optimal cocharacter
 246 for $e_0 \in \mathfrak{l}$ as defined in [23, 2.2]. We adopt the notation of [23] and write $\mathfrak{l}(\lambda, i)$ for
 247 the i -weight space of the 1-dimensional torus $\lambda(k^\times)$ acting on \mathfrak{l} . By construction, $e_0 \in$
 248 $\mathfrak{l}(\lambda, 2)$. Given $k \in \mathbb{Z}$ we set $\mathfrak{l}(\lambda, \geq k) := \bigoplus_{i \geq k} \mathfrak{l}(\lambda, i)$. Then $\mathfrak{p}_\mathfrak{l}(\lambda) := \mathfrak{l}(\lambda, \geq 0)$ is the optimal
 249 parabolic subalgebra of $e_0 \in \mathfrak{l}$. We write $P_L(\lambda)$ for the corresponding parabolic subgroup
 250 of L . Note that $P_L(\lambda) = Z_L(\lambda)R_u(P_L)$ where $\text{Lie}(Z_L(\lambda)) = \mathfrak{l}(\lambda, 0)$ and $\text{Lie}(R_u(P_L(\lambda))) =$
 251 $\mathfrak{l}(\lambda, \geq 1)$. As explained in [23, p. 346], there exists a non-zero regular function φ on $\mathfrak{l}(\lambda, 2)$
 252 semi-invariant under the adjoint action of $Z_L(\lambda)$ and such that the orbit $(\text{Ad } Z_L(\lambda)) \cdot e_0$
 253 coincides with the principal open subset $\mathfrak{l}(\lambda, 2)_\varphi = \{x \in \mathfrak{l}(\lambda, 2) \mid \varphi(x) \neq 0\}$ of $\mathfrak{l}(\lambda, 2)$.

254 Let $\mathfrak{g} = \mathfrak{n}_- \oplus \mathfrak{l} \oplus \mathfrak{n}_+$ be a parabolic decomposition of \mathfrak{g} associated with the Levi
 255 subalgebra \mathfrak{l} . Then there exists a parabolic subgroup $P = LN_+$ such that $N_+ = R_u(P)$
 256 and $\mathfrak{n}_+ = \text{Lie}(N_+)$. We set $P(\mathfrak{l}, e_0) := P_L(\lambda)N_+$, a parabolic subgroup of G contained in
 257 P . It is easy to see that $\text{Lie}(P(\mathfrak{l}, e_0)) = \mathfrak{p}_\mathfrak{l}(\lambda) \oplus \mathfrak{n}_+$ and

$$258 \quad \mathfrak{r}(\mathfrak{l}, e_0) := \mathfrak{z}(\mathfrak{l}) \oplus \mathfrak{l}(\lambda, \geq 2) \oplus \mathfrak{n}_+$$

259 is a solvable ideal of $\text{Lie}(P(\mathfrak{l}, e_0))$ invariant under the adjoint action of $P(\mathfrak{l}, e_0)$. Identifying
 260 direct sums with direct products we put

$$261 \quad \mathfrak{r}(\mathfrak{l}, e_0)_{\text{reg}} := \mathfrak{z}(\mathfrak{l})_{\text{reg}} \oplus \mathfrak{l}(\lambda, 2)_\varphi \oplus \mathfrak{l}(\lambda, \geq 3) \oplus \mathfrak{n}_+.$$

262 By construction, $\mathfrak{r}(\mathfrak{l}, e_0)_{\text{reg}}$ is a Zariski open subset of $\mathfrak{r}(\mathfrak{l}, e_0)$ invariant under the adjoint
 263 action of $P(\mathfrak{l}, e_0)$. We mention for further references that $\mathfrak{z}(\mathfrak{l}) = \text{Lie}(Z_G(L))$ and there
 264 exists a connected algebraic subgroup $R(\mathfrak{l}, e_0)$ of G with a maximal torus $Z_G(L)^\circ$ such
 265 that $\mathfrak{r}(\mathfrak{l}, e_0) = \text{Lie}(R(\mathfrak{l}, e_0))$.

266 **Theorem 2.3.** *In the above notation, $\mathcal{D}(\mathfrak{l}, e_0) = (\text{Ad } G) \cdot \mathfrak{r}(\mathfrak{l}, e_0)_{\text{reg}}$ and the Zariski closure*
 267 *of $\mathcal{D}(\mathfrak{l}, e_0)$ in \mathfrak{g} coincides with $(\text{Ad } G) \cdot \mathfrak{r}(\mathfrak{l}, e_0)$.*

268 **Proof.** Let $z \in \mathfrak{z}(\mathfrak{l})_{\text{reg}}$. Then $(\text{Ad } P_L(\lambda)) \cdot (z + e_0) = z + (\text{Ad } P_L(\lambda)) \cdot e_0$. Since $[\mathfrak{p}_\mathfrak{l}(\lambda), e_0] =$
 269 $\mathfrak{l}(\lambda, \geq 2)$ by [23, Theorem 2.3(iv)] and the orbit $(\text{Ad } R_u(P_L(\lambda))) \cdot e_0 \subseteq e_0 + \mathfrak{l}(\lambda, \geq 3)$ is
 270 Zariski closed by Rosenlicht's theorem [27, Theorem 2], the equality $(\text{Ad } R_u(P_L(\lambda))) \cdot e_0 =$
 271 $e_0 + \mathfrak{l}(\lambda, \geq 3)$ must hold. It implies that

$$272 \quad (\text{Ad } P_L(\lambda)) \cdot (z + e_0) = z + (\text{Ad } Z_L(\lambda)) \cdot (e_0 + \mathfrak{l}(\lambda, \geq 3)) = z + \mathfrak{l}(\lambda, 2)_\varphi + \mathfrak{l}(\lambda, \geq 3).$$

As $z \in \mathfrak{z}(\mathfrak{l})_{\text{reg}}$ and e_0 is a nilpotent element of \mathfrak{l} , it must be the endomorphism $\text{ad}(z + e_0)$ acts invertibly on \mathfrak{n}_+ . As $(\text{Ad } N_+) \cdot (z + e_0) \subseteq z + e_0 + \mathfrak{n}_+$, another application of Rosenlicht's theorem yields $(\text{Ad } N_+) \cdot (z + e_0) = z + e_0 + \mathfrak{n}_+$. It follows that

$$\begin{aligned} (\text{Ad } P(\mathfrak{l}, e_0)) \cdot (z + e_0) &= (\text{Ad } P_L(\lambda)) \cdot ((\text{Ad } N_+) \cdot (z + e_0)) \\ &= (\text{Ad } P_L(\lambda)) \cdot (z + e_0 + \mathfrak{n}_+) = \mathfrak{n}_+ + (\text{Ad } P_L(\lambda)) \cdot (z + e_0) \\ &= z + \mathfrak{l}(\lambda, 2)_\varphi + \mathfrak{l}(\lambda, \geq 3) + \mathfrak{n}_+. \end{aligned} \tag{1}$$

As this holds for all $z \in \mathfrak{z}(\mathfrak{l})_{\text{reg}}$ we now deduce that $(\text{Ad } P(\mathfrak{l}, e_0)) \cdot (\mathfrak{z}(\mathfrak{l})_{\text{reg}} + e_0) = \mathfrak{r}(\mathfrak{l}, e_0)_{\text{reg}}$ and $\mathcal{D}(\mathfrak{l}, e_0) = (\text{Ad } G) \cdot \mathfrak{r}(\mathfrak{l}, e_0)_{\text{reg}}$. As $\mathfrak{r}(\mathfrak{l}, e_0)_{\text{reg}}$ is Zariski open in $\mathfrak{r}(\mathfrak{l}, e_0)$, the Zariski closure of $\mathcal{D}(\mathfrak{l}, e_0)$ contains $(\text{Ad } G) \cdot \mathfrak{r}(\mathfrak{l}, e_0)$. Since the latter set is the image of the action morphism

$$G \times^{P(\mathfrak{l}, e_0)} \mathfrak{r}(\mathfrak{l}, e_0) \longrightarrow \mathfrak{g}, \quad (g, r) \mapsto (\text{Ad } g) \cdot r,$$

and $P(\mathfrak{l}, e_0)$ is a parabolic subgroup of G , the set $(\text{Ad } G) \cdot \mathfrak{r}(\mathfrak{l}, e_0)$ is Zariski closed in \mathfrak{g} ; see [13, Lemma 8.7(c)], for example. Since $(\text{Ad } G) \cdot \mathfrak{r}(\mathfrak{l}, e_0)_{\text{reg}}$ is Zariski dense in $(\text{Ad } G) \cdot \mathfrak{r}(\mathfrak{l}, e_0)$ we conclude that $\overline{\mathcal{D}(\mathfrak{l}, e_0)} = (\text{Ad } G) \cdot \mathfrak{r}(\mathfrak{l}, e_0)$, thereby completing the proof. \square

Remark 2.4. Since $\lambda \in X_*(T)$ is an optimal cocharacter for $e_0 \in \mathcal{N}(\mathfrak{l})$ in the sense of the Kempf–Rousseau theory, it follows from the proof of Theorem 2.3 that for any $x \in \mathfrak{r}(\mathfrak{l}, e_0)_{\text{reg}}$ the inclusion $G_x \subset P(\mathfrak{l}, e_0)$ holds. In the case where $\text{char}(k) = 0$, this was first observed in [1, Zusatz 5.5(e)].

2.4. Restricting the adjoint quotient map to a decomposition class

Let $\mathfrak{t} = \text{Lie}(T)$ be a maximal toral subalgebra of \mathfrak{g} containing $\mathfrak{z}(\mathfrak{l})$ and let $W = N_G(T)/T$ be the Weyl group of G . Let $n = \dim G$ and $\ell = \text{rk } G$. Under our assumptions on G the invariant ring $k[\mathfrak{g}]^G$ is freely generated by ℓ homogeneous regular functions f_1, \dots, f_ℓ and the restriction map $k[\mathfrak{g}] \rightarrow k[\mathfrak{t}]$ induces a natural isomorphism of invariant rings $j: k[\mathfrak{g}]^G \rightarrow k[\mathfrak{t}]^W$. Furthermore, the categorical quotient morphism $F: \mathfrak{g} \rightarrow \mathfrak{g}/G \cong \mathfrak{t}/W$ sending any $x \in \mathfrak{g}$ to $(f_1(x), \dots, f_\ell(x)) \in \mathbb{A}^\ell$ is flat, surjective, and all its fibres are irreducible complete intersections of dimension $n - \ell$ in \mathfrak{g} consisting of finitely many G -orbits. If $x \in \mathfrak{g}$ then $f(x) = f(x_s)$ for every $f \in k[\mathfrak{g}]^G$, and for every $\xi \in \mathbb{A}^\ell$ there is a unique orbit $W \cdot t \subset \mathfrak{t}$ such that the fibre $F^{-1}(\xi)$ consists of all $x \in \mathfrak{g}$ such that $(\text{Ad } g) \cdot x_s = t$ for some $g \in G$. The special fibre $F^{-1}(0)$ coincides with the nilpotent cone $\mathcal{N}(\mathfrak{g})$ of \mathfrak{g} . All these results are well known and can be found in [13, § 7], for example.

In this subsection we are concerned with the restriction of the adjoint quotient morphism F to the Zariski closure of a decomposition class.

Proposition 2.5. *Let \bar{F} denote the restriction of the adjoint quotient morphism $F: \mathfrak{g} \rightarrow \mathfrak{t}/W$ to $\overline{\mathcal{D}(\mathfrak{l}, e_0)}$. Then all irreducible components of the fibres of \bar{F} have dimension $n - \dim \mathfrak{l}_{e_0}$, the special fibre $\bar{F}^{-1}(0) = \overline{\mathcal{D}(\mathfrak{l}, e_0)} \cap \mathcal{N}(\mathfrak{g})$ is irreducible, and $\dim \mathcal{D}(\mathfrak{l}, e_0) = n - \dim \mathfrak{l}_{e_0} + \dim \mathfrak{z}(\mathfrak{l})$.*

Proof. To ease notation we put $n(\mathfrak{l}, e_0) = \dim \mathcal{D}(\mathfrak{l}, e_0)$. Let X be a fibre of \bar{F} and $x \in X$. By Theorem 2.3, we may assume without loss that $x \in \mathfrak{r}(\mathfrak{l}, e_0)$. Since $\mathfrak{r}(\mathfrak{l}, e_0) = \text{Lie}(R(\mathfrak{l}, e_0))$ and $Z_G(L)^\circ$ is a maximal torus of $R(\mathfrak{l}, e_0)$, the standard fact in the theory

of algebraic groups that maximal tori are conjugate yields that the semi-simple part of x is $R(\mathfrak{l}, e_0)$ -conjugate to an element of $\mathfrak{z}(\mathfrak{l}) = \text{Lie}(Z_G(L)^\circ)$; see [2, 11.8]. In view of the discussion above this implies that

$$X = \bigcup_{t \in (W \cdot x_s) \cap \mathfrak{z}(\mathfrak{l})} (\text{Ad } G) \cdot (t + \mathfrak{l}(\lambda, \geq 2) + \mathfrak{n}_+ \cap \mathfrak{g}_t). \quad (2)$$

(As $t \in \mathfrak{z}(\mathfrak{l})$ we have $\mathfrak{l}(\lambda, \geq 2) \subset \mathfrak{g}_t$.) As $\mathfrak{z}(\mathfrak{l}) \subset \overline{\mathcal{D}(\mathfrak{l}, e_0)}$ this implies that \bar{F} maps $\overline{\mathcal{D}(\mathfrak{l}, e_0)}$ onto $\mathfrak{z}(\mathfrak{l})/W$, the image of $\mathfrak{z}(\mathfrak{l})$ in \mathfrak{t}/W . Since W is a finite group the quotient morphism $\mathfrak{t} \rightarrow \mathfrak{t}/W$ is finite, hence closed and has finite fibres. So $\mathfrak{z}(\mathfrak{l})/W$ is a closed subset of \mathfrak{t}/W of dimension $d := \dim \mathfrak{z}(\mathfrak{l})$. It follows that the minimal dimension of the fibres of \bar{F} equals $n(\mathfrak{l}, e_0) - d$.

On the other hand, it follows from [9, Theorem 14.8(a)] that the union of those irreducible components of the fibres of \bar{F} that have dimension $> n(\mathfrak{l}, e_0) - d$ is Zariski closed in $\overline{\mathcal{D}(\mathfrak{l}, e_0)}$. That union cannot coincide with $\overline{\mathcal{D}(\mathfrak{l}, e_0)}$ by the preceding remark. Since all irreducible components of the fibres of \bar{F} contain open G -orbits and $\mathcal{D}(\mathfrak{l}, e_0)$ is dense in $\overline{\mathcal{D}(\mathfrak{l}, e_0)}$, there exists $z \in \mathfrak{z}(\mathfrak{l})_{\text{reg}}$ such that $n - \dim \mathfrak{g}_{z+e_0} = n(\mathfrak{l}, e_0) - d$. Since z is the semi-simple part of $z + e_0$ we have the equality $\mathfrak{g}_{z+e_0} = \mathfrak{l}_{e_0}$. So $\dim \mathfrak{g}_{z+e_0} = \dim \mathfrak{l}_{e_0}$ implying $n(\mathfrak{l}, e_0) = n - \dim \mathfrak{l}_{e_0} + d$. This proves the last statement of the proposition.

If one of the fibres of \bar{F} has an irreducible component of dimension $> n(\mathfrak{l}, e_0) - d = n - \dim \mathfrak{l}_{e_0}$, then by the same token it contains an open G -orbit of the same dimension and hence an element $v \in \overline{\mathcal{D}(\mathfrak{l}, e_0)}$ such that $\dim \mathfrak{g}_v < \dim \mathfrak{l}_{e_0}$. Since the set $\{x \in \overline{\mathcal{D}(\mathfrak{l}, e_0)} \mid \dim \mathfrak{g}_x \geq \dim \mathfrak{l}_{e_0}\}$ is Zariski closed by Chevalley's semi-continuity theorem and $\dim \mathfrak{g}_x = \dim \mathfrak{l}_{e_0}$ for all $x \in \mathcal{D}(\mathfrak{l}, e_0)$ it follows that $\dim \mathfrak{g}_x \geq \dim \mathfrak{l}_{e_0}$ for all $x \in \overline{\mathcal{D}(\mathfrak{l}, e_0)}$. This contradiction shows that all irreducible components of the fibres of \bar{F} have the same dimension.

If X is the special fibre of \bar{F} , then $x_s = 0$ and (2) gives

$$\overline{\mathcal{D}(\mathfrak{l}, e_0)} \cap \mathcal{N}(\mathfrak{g}) = (\text{Ad } G) \cdot (\mathfrak{l}(\lambda, \geq 2) + \mathfrak{n}_+). \quad (3)$$

Therefore, the variety $\overline{\mathcal{D}(\mathfrak{l}, e_0)} \cap \mathcal{N}(\mathfrak{g})$ is irreducible which completes the proof. \square

Remark 2.6. If \mathcal{S} is a sheet of \mathfrak{g} whose open decomposition class equals $\mathcal{D}(\mathfrak{l}, e_0)$ then by maximality \mathcal{S} must coincide with the set of all G -orbits of maximal dimension contained in $\overline{\mathcal{D}(\mathfrak{l}, e_0)}$. Therefore, Proposition 2.5 implies that every sheet of \mathfrak{g} contains a unique nilpotent orbit. Since W is a finite group, the quotient morphism $\mathfrak{t} \rightarrow \mathfrak{t}/W$ is open and closed. As $\mathfrak{z}(\mathfrak{l})_{\text{reg}}$ is open in $\mathfrak{z}(\mathfrak{l}) \subseteq \mathfrak{t}$, the set $\mathfrak{z}(\mathfrak{l})_{\text{reg}}/W$ is open in $\mathfrak{z}(\mathfrak{l})/W$, a closed subset of \mathfrak{t}/W . So the proof of Proposition 2.5 also shows that any decomposition class $\mathcal{D}(\mathfrak{l}, e_0) = \bar{F}^{-1}(\mathfrak{z}(\mathfrak{l})_{\text{reg}}/W)$ of \mathfrak{g} is Zariski open in its closure. It is worth mentioning that all G -stable pieces in the union (2) are Zariski closed in \mathfrak{g} . This follows from [13, Lemma 8.7(c)] (see the end of the proof of Theorem 2.3 for a similar argument). If $\mathcal{D}(\mathfrak{l}, e_0)$ is the open decomposition class of a sheet \mathcal{S} and \mathcal{O} is the nilpotent orbit contained in \mathcal{S} then it follows from Proposition 2.5 that $\dim \mathfrak{z}(\mathfrak{l}) = \dim \mathcal{S} - \dim \mathcal{O}$. This number is called the *rank* of \mathcal{S} and denoted $\text{rk}(\mathcal{S})$. It is immediate from Proposition 2.5 that under our assumptions on G a sheet \mathcal{S} of \mathfrak{g} is a single nilpotent G -orbit if and only if $\text{rk}(\mathcal{S}) = 0$.

2.5. Induced nilpotent orbits and sheets

Given $x \in \mathfrak{g}$ we denote by $\mathcal{O}(x)$ the adjoint G -orbit of x . If x lies in a Levi subalgebra $\mathfrak{l} = \text{Lie}(L)$ of \mathfrak{g} we set $\mathcal{O}_L(x) := (\text{Ad } L) \cdot x$.

Combining Theorem 2.3 with Proposition 2.5, we observe that the unique open G -orbit $\mathcal{O}(e)$ in $\overline{\mathcal{D}(\mathfrak{l}, e_0)} \cap \mathcal{N}(\mathfrak{g})$ intersects densely with $\mathfrak{l}(\lambda, 2)_\varphi \oplus \mathfrak{l}(\lambda, \geq 3) \oplus \mathfrak{n}_+$ and has the property that $\dim \mathfrak{g}_e = \dim \mathfrak{l}_{e_0}$. As $\mathfrak{l}(\lambda, 2)_\varphi \oplus \mathfrak{l}(\lambda, \geq 3) = (\text{Ad } P_L(\lambda)) \cdot e_0$ by our choice of λ we see that $\mathcal{O}(e)$ intersects densely with $\mathcal{O}_L(e_0) + \mathfrak{n}_+$. In other words, the orbit $\mathcal{O}(e)$ coincides with $\text{Ind}_\mathfrak{l}^\mathfrak{g} \mathcal{O}_L(e_0)$, the nilpotent G -orbit obtained from $\mathcal{O}_L(e_0)$ by Lusztig–Spaltenstein induction.

As in the characteristic zero case, induction of nilpotent orbits is transitive in the following sense: if L, L' are Levi subgroups of G with $L \subset L'$ and e_0 is a nilpotent elements of $\mathfrak{l} = \text{Lie}(L)$ then

$$\text{Ind}_\mathfrak{l}^\mathfrak{g} \mathcal{O}_L(e_0) = \text{Ind}_{\mathfrak{l}'}^\mathfrak{g} (\text{Ind}_{\mathfrak{l}}^{\mathfrak{l}'} \mathcal{O}_L(e_0)) \tag{4}$$

where $\mathfrak{l}' = \text{Lie}(L')$. In order to see this it suffices to note that both orbits in (4) have the same dimension, the Zariski closure of the decomposition class $(\text{Ad } L') \cdot (e_0 + \mathfrak{z}(\mathfrak{l})_{\text{reg}})$ in \mathfrak{l}' contains $\mathfrak{z}(\mathfrak{l}')_{\text{reg}} + e$ for some $e \in \text{Ind}_{\mathfrak{l}}^{\mathfrak{l}'} \mathcal{O}_L(e_0)$, and $\mathfrak{z}(\mathfrak{l}') \subset \mathfrak{z}(\mathfrak{l})$. The statement then follows from Proposition 2.5.

Remark 2.7. Since the decomposition class $\mathcal{D}(\mathfrak{l}, e_0)$ is evidently independent of the choice of a triangular decomposition of \mathfrak{g} containing \mathfrak{l} , the above discussion implies that induction of orbits is independent of the choice of a parabolic subalgebra of \mathfrak{g} containing \mathfrak{l} . This was first observed by Lusztig–Spaltenstein for unipotent classes in reductive groups over algebraically closed fields of arbitrary characteristic; see [17, Theorem 2.2]. The argument in [17] relied on the theory of complex representations of finite groups of Lie type and some results of Mizuno. More recently, Lusztig returned to this subject and found another proof of the fact that induction is independent of the choice of a parabolic. His argument is similar to ours but still works in the setting of a reductive group, possibly disconnected; see [18, Lemma 10.3(a)]. In principle, one could use a Bardsley–Richardson projection $\pi : G \rightarrow \mathfrak{g}$ from [4] to obtain a Lie algebra analogue of [17, Theorem 2.2]. Indeed, in our situation π provides a nice G -equivariant isomorphism between the unipotent variety of G and $\mathcal{N}(\mathfrak{g})$. However, to the best of our knowledge this line of reasoning was never carried out in the literature under our assumption on G and we felt that arguing in the spirit of [1] was more relevant for our purposes (see e.g., Remark 2.4).

A nilpotent element e_0 in a Levi subalgebra $\mathfrak{l} = \text{Lie}(L)$ of \mathfrak{g} is called *rigid* if the orbit $\mathcal{O}_L(e_0)$ cannot be obtained by Lusztig–Spaltenstein induction from a proper Levi subalgebra of \mathfrak{l} .

Theorem 2.8. *If \mathcal{S} is a sheet of \mathfrak{g} and $\mathcal{D}(\mathfrak{l}, e_0)$ is the open decomposition class of \mathcal{S} , then e_0 is rigid in \mathfrak{l} and the unique nilpotent orbit in \mathcal{S} has the form $\mathcal{O} = \text{Ind}_\mathfrak{l}^\mathfrak{g} \mathcal{O}_L(e_0)$. Moreover, $\dim \mathfrak{g}_x = \dim \mathfrak{l}_{e_0}$ for any $x \in \mathcal{O}$. Conversely, for any pair (\mathfrak{l}, e_0) , where \mathfrak{l} is a Levi subalgebra of \mathfrak{g} and e_0 is a rigid nilpotent element of \mathfrak{l} , there is a unique sheet of \mathfrak{g} whose open decomposition class equals $\mathcal{D}(\mathfrak{l}, e_0)$.*

Proof. Let $\mathcal{D}(\mathfrak{l}', e'_0)$ be the open decomposition class of \mathcal{S} and suppose for a contradiction that $\mathcal{O}_{L'}(e'_0) = \text{Ind}_{\mathfrak{l}'}^{\mathfrak{g}} \mathcal{O}_L(e_0)$ for some proper Levi subalgebra \mathfrak{l}' of \mathfrak{g} (here L, L' are Levi subgroups of G with $L \subset L'$ such that $\mathfrak{l}' = \text{Lie}(L')$ and $\mathfrak{l} = \text{Lie}(L)$). Note that $\mathfrak{z}(\mathfrak{l}') \subset \mathfrak{z}(\mathfrak{l})$ and $(\text{Ad } L') \cdot (e_0 + \mathfrak{z}(\mathfrak{l})_{\text{reg}})$ lies in $\mathcal{D}(\mathfrak{l}, e_0)$. Let $\mathfrak{l}' = \mathfrak{n}'_- \oplus \mathfrak{l} \oplus \mathfrak{n}'_+$ be a triangular decomposition of \mathfrak{l}' . Replacing G and \mathfrak{g} , respectively, by L' and \mathfrak{l}' in the proof of Theorem 2.3 and applying (1) to any $z \in \mathfrak{z}(\mathfrak{l})_{\text{reg}}$, it is straightforward to see that $\mathcal{D}(\mathfrak{l}, e_0)$ contains $\mathfrak{z}(\mathfrak{l})_{\text{reg}} + \mathcal{O}_L(e_0) + \mathfrak{n}'_+$ and hence $\mathfrak{z}(\mathfrak{l})_{\text{reg}} + e'_0$. But then $\overline{\mathcal{D}(\mathfrak{l}, e_0)}$ contains $\mathfrak{z}(\mathfrak{l}) + e'_0$. As $\mathfrak{z}(\mathfrak{l}')_{\text{reg}} \subset \mathfrak{z}(\mathfrak{l})$ it follows that $\overline{\mathcal{S}} = \overline{\mathcal{D}(\mathfrak{l}', e'_0)}$ lies in $\overline{\mathcal{D}(\mathfrak{l}, e_0)}$. Combining Proposition 2.5 with (4) and our discussion at the beginning of this subsection we see that $\overline{\mathcal{D}(\mathfrak{l}', e'_0)}$ and $\overline{\mathcal{D}(\mathfrak{l}, e_0)}$ share the same nilpotent orbit $\mathcal{O} = \text{Ind}_{\mathfrak{l}'}^{\mathfrak{g}} \mathcal{O}_L(e_0)$. On the other hand, Proposition 2.5 gives

$$\begin{aligned} \dim \mathcal{S} &= \dim \mathcal{D}(\mathfrak{l}', e'_0) = n - \dim \mathfrak{l}'_{e'_0} + \dim \mathfrak{z}(\mathfrak{l}') = n - \dim \mathfrak{l}_{e_0} + \dim \mathfrak{z}(\mathfrak{l}') \\ &< n - \dim \mathfrak{l}_{e_0} + \dim \mathfrak{z}(\mathfrak{l}) = \dim \mathcal{D}(\mathfrak{l}, e_0) \end{aligned}$$

contrary to the fact that \mathcal{S} is an irreducible component of $\mathfrak{g}_{(m)}$ where $m = n - \dim \mathfrak{l}_{e_0}$. This contradiction shows that $\mathfrak{l}' = \mathfrak{l}$ and $e'_0 = e_0$ is rigid in \mathfrak{l} .

If \mathcal{O} is the nilpotent orbit of \mathcal{S} then $\dim \mathcal{O} = n - \dim \mathfrak{l}_{e_0}$ by Proposition 2.5. Then $\dim \mathfrak{g}_x = \dim \mathfrak{l}_{e_0}$ for all $x \in \mathcal{O}$ because $\mathfrak{g}_x = \text{Lie}(G_x)$.

Finally, suppose e'_0 is a rigid nilpotent element in a Levi subalgebra \mathfrak{l}' of \mathfrak{g} and put $s = \dim \mathfrak{l}'_{e'_0}$. Let \mathcal{S}' be the union of all adjoint G -orbits of dimension $n - s$ contained in $\overline{\mathcal{D}(\mathfrak{l}', e'_0)}$. It follows from Chevalley's semi-continuity theorem that \mathcal{S}' is a Zariski open subset of $\overline{\mathcal{D}(\mathfrak{l}', e'_0)}$ (see the end of the proof of Proposition 2.5 for a similar argument). Since $\overline{\mathcal{D}(\mathfrak{l}', e'_0)}$ is irreducible, so is the quasi-affine variety \mathcal{S}' . We claim that \mathcal{S}' is a sheet of \mathfrak{g} . Indeed, suppose the contrary. Then $\mathcal{S}' \subsetneq \tilde{\mathcal{S}}$ for some sheet $\tilde{\mathcal{S}}$ of \mathfrak{g} contained in $\mathfrak{g}_{(n-s)}$. Let $\mathcal{D}(\mathfrak{l}, e_0)$ be the open decomposition class of $\tilde{\mathcal{S}}$. We may assume without loss of generality that both \mathfrak{l}' and \mathfrak{l} are standard Levi subalgebras of \mathfrak{g} , so that the maximal toral subalgebra $\mathfrak{t} = \text{Lie}(T) \subseteq \mathfrak{l} \cap \mathfrak{l}'$ of \mathfrak{g} contains both $\mathfrak{z}(\mathfrak{l})$ and $\mathfrak{z}(\mathfrak{l}')$.

Let $z \in \mathfrak{z}(\mathfrak{l}')_{\text{reg}}$. Since $\tilde{\mathcal{S}}$ contains $\mathcal{D}(\mathfrak{l}', e'_0)$ and the Zariski closure of $\tilde{\mathcal{S}}$ coincides with $\overline{\mathcal{D}(\mathfrak{l}, e_0)}$ which, in turn, lies in $\overline{\mathfrak{g}_{(n-s)}}$, the G -orbit $\mathcal{O}(z + e'_0)$ is open in one of the fibres of the morphism $\tilde{F}: \overline{\mathcal{D}(\mathfrak{l}, e_0)} \rightarrow \mathfrak{t}/W$. In view of (2) it coincides with $(\text{Ad } G) \cdot (t + V)$ for some $t \in (W \cdot z) \cap \mathfrak{z}(\mathfrak{l})$ and some large subset V of $\mathfrak{l}(\lambda, \geq 2) + \mathfrak{n}_+ \cap \mathfrak{g}_{\mathfrak{t}}$ which we are now going to describe. Since both \mathfrak{l} and $\mathfrak{n}_+ \cap \mathfrak{g}_{\mathfrak{t}}$ are centralised by t , combining (2) with the inclusion $\overline{\mathcal{D}(\mathfrak{l}, e_0)} \subseteq \overline{\mathfrak{g}_{(n-s)}}$ one observes that V consists of all elements in $\mathfrak{l}(\lambda, \geq 2) + \mathfrak{n}_+ \cap \mathfrak{g}_{\mathfrak{t}}$ whose centraliser in $\mathfrak{g}_{\mathfrak{t}}$ has dimension s . Furthermore, V is Zariski open in $\mathfrak{l}(\lambda, \geq 2) + \mathfrak{n}_+ \cap \mathfrak{g}_{\mathfrak{t}}$.

Write $z + e'_0 = (\text{Ad } g) \cdot (t + v)$ where $g \in G$ and $v \in V$ and choose $n_w \in N_G(T)$ such that $(\text{Ad } n_w) \cdot z = t$. Since $[t, v] = 0 = [z, e'_0]$ we have that $(\text{Ad } g) \cdot t = z$ and $(\text{Ad } g) \cdot v = e'_0$. Then $gn_w \in G_z$. Since $z \in \mathfrak{z}(\mathfrak{l}')_{\text{reg}}$, and G satisfies the standard hypotheses, applying [31, 3.19, 4.2] yields $G_z = L'$. Set $v' := (\text{Ad } n_w^{-1}) \cdot v$. Since $e'_0 = (\text{Ad } gn_w) \cdot v'$, we now deduce that $v' \in \mathcal{O}_{L'}(e'_0)$. On the other hand, the concluding remark of the preceding paragraph in conjunction with (2) shows that v lies in the $G_{\mathfrak{t}}$ -orbit which intersects densely with

$\mathfrak{l}(\lambda, \geq 2) + \mathfrak{n}_+ \cap \mathfrak{g}_t \subset \mathfrak{g}_t$. Since $L' = n_w^{-1}G_t n_w$, it follows that $\mathcal{O}_{L'}(e'_0)$ intersects densely with $(\text{Ad } n_w^{-1}) \cdot (\mathfrak{l}(\lambda, \geq 2) + \mathfrak{n}_+ \cap \mathfrak{g}_t) \subset \mathfrak{g}_z = \mathfrak{l}'$.

Since the orbit $\mathcal{O}_{L'}(e'_0)$ is rigid in \mathfrak{l}' this yields $\mathfrak{n}_+ \cap \mathfrak{g}_t = 0$ (otherwise $\mathcal{O}_{L'}(e'_0)$ would be induced from $(\text{Ad } n_w^{-1})(\mathfrak{l})$, a proper Levi subalgebra of \mathfrak{l}'). Since $t \in \mathfrak{z}(\mathfrak{l})$ and $\mathfrak{g}_t = (\mathfrak{n}_- \cap \mathfrak{g}_t) \oplus \mathfrak{l} \oplus (\mathfrak{n}_+ \cap \mathfrak{g}_t)$ we now obtain that $t \in \mathfrak{z}(\mathfrak{l})_{\text{reg}}$. Since $t \in W \cdot z$ and $G_z = L'$ it follows that $\dim \mathfrak{z}(\mathfrak{l}) = \dim \mathfrak{z}(\mathfrak{l}')$ and

$$\dim \mathcal{S}' = \dim \mathcal{D}(\mathfrak{l}', e'_0) = n - s + \dim \mathfrak{z}(\mathfrak{l}') = n - s + \dim \mathfrak{z}(\mathfrak{l}) = \dim \mathcal{D}(\mathfrak{l}, e_0) = \dim \tilde{\mathcal{S}}$$

by Proposition 2.5. This contradiction shows that $\mathcal{S}' = \tilde{\mathcal{S}}$ is a sheet of \mathfrak{g} thereby completing the proof. \square

Remark 2.9. Theorem 2.8 implies that the sheets of \mathfrak{g} are in 1–1 correspondence with the G -conjugacy classes of pairs (L, \mathcal{O}_L) where L is a Levi subgroup of G and \mathcal{O}_L is a rigid nilpotent L -orbit in $\text{Lie}(L)$. In the characteristic zero case, this is a well-known result of Borho; see [3].

3. Classifying the rigid orbits in exceptional Lie algebras

3.1. Some historical remarks

In order to complete the picture outlined in §2 one needs to classify all rigid orbits in the reductive Lie algebras $\mathfrak{g} = \text{Lie}(G)$. Of course, the general case reduces quickly to the case where the group G is simple, and here the problem has a long history. All rigid nilpotent orbits in Lie algebras of classical groups are described by Kraft [15] in type A and by Kempken [14] and Spaltenstein [28] in types B, C, D. If $\mathfrak{g} = \mathfrak{sl}_n$ then $\{0\}$ is the only rigid orbit in \mathfrak{g} , whilst for $\mathfrak{g} = \mathfrak{so}_n$ or \mathfrak{sp}_n the classification is given in terms of partitions of n . More precisely, it is proved in [15] that a nilpotent element in \mathfrak{gl}_n corresponding to a partition λ of n is Richardson in a parabolic subalgebra of \mathfrak{gl}_n associated with the dual partition λ' . The arguments in [14] rely on Kraft's result in a crucial way.

Although the main theorems in [14] are stated under the assumption that the base field has characteristic 0, the proofs are based on linear algebra and the same combinatorial description is valid in any good characteristic, i.e., for $p \neq 2$. In characteristic 2, nilpotent orbits and sheets in \mathfrak{so}_n and \mathfrak{sp}_n were studied by Hesselink [12] and by Spaltenstein [29] who conjectured that every sheet in a reductive Lie algebra over an algebraically closed field of arbitrary characteristic should contain a *unique* nilpotent orbit. Of course, confirming this would imply that the number of nilpotent orbits in \mathfrak{g} is finite. The main results of [29] show that the conjecture holds for G classical, but the general case remains open in bad characteristic. We mention that Spaltenstein's conjecture makes sense in the context of infinitesimal sheets and their coadjoint analogues (see Remark 2.1) and confirming it would enable one to prove a natural analogue of the Kac–Weisfeiler conjecture in arbitrary characteristic; see [24, 5.6] for detail.

For G exceptional, the rigid orbits in $\mathcal{N}(\mathfrak{g})$ were first classified by Elashvili [10] under the assumption that $\text{char}(k) = 0$. Furthermore, for any orbit $\mathcal{O} \subset \mathcal{N}(\mathfrak{g})$, Elashvili lists all sheets of \mathfrak{g} containing \mathcal{O} . This important result is presented in [10] in the form of tables which were recently double-checked in [8] by using computational methods.

3.2. Odd nilpotent elements

It was first observed by Lusztig–Spaltenstein that if $\text{char}(k)$ is 0 or large for G then many nilpotent orbits in \mathfrak{g} can be induced in a very natural way; see [17, Proposition 1.9]. The goal of this subsection is to show that this result is still valid under the assumption that G satisfies the standard hypotheses.

Let Φ be the root system of G with respect to a maximal torus T of G and let Π be a basis of simple roots in Φ . Each orbit $\mathcal{O} \subset \mathcal{N}(\mathfrak{g})$ is uniquely labelled by its weighted Dynkin diagram D which assigns to each $\alpha \in \Pi$ an integer $D(\alpha) \in \{0, 1, 2\}$. According to [23, 2.3, 2.4], each such D gives rise to a cocharacter $\lambda = \lambda_D \in X_*(T \cap \mathcal{D}G)$ which is optimal in the sense of the Kempf–Rousseau theory for every element of a principal Zariski open subset of $\mathfrak{g}(\lambda, 2)$. The subset, denoted earlier by $\mathfrak{g}(\lambda, 2)_\varphi$, coincides with the intersection of \mathcal{O} with $\mathfrak{g}(\lambda, 2)$. A nilpotent orbit $\mathcal{O} = \mathcal{O}(D)$ is said to be *odd* if $D(\alpha) \in \{0, 1\}$ for all $\alpha \in \Pi$.

Given a weighted Dynkin diagram D we let $\Pi' = \Pi'(D)$ be the set of all $\alpha \in \Pi$ such that $D(\alpha) \in \{0, 1\}$ and denote by $L = L(D)$ the standard Levi subgroup of G generated by T and all unipotent root subgroups U_α with $\alpha \in \Pi'$. Given a non-empty subset $\Pi_0 \subseteq \Pi$ and a weighted Dynkin diagram D_0 of the standard Levi subalgebra of \mathfrak{g} associated with Π_0 , we define $\tilde{D}_0: \Pi \rightarrow \{0, 1, 2\}$ by setting $\tilde{D}_0(\alpha) = D_0(\alpha)$ for all $\alpha \in \Pi_0$ and $\tilde{D}_0(\alpha) = 2$ for all $\alpha \in \Pi \setminus \Pi_0$.

Proposition 3.1. *The following are true:*

(i) *If the orbit $\mathcal{O} = \mathcal{O}(D)$ is not odd then $\mathfrak{l} = \text{Lie}(L(D))$ is a proper Levi subalgebra of \mathfrak{g} and $\mathcal{O}(D) = \text{Ind}_\mathfrak{l}^\mathfrak{g} \mathcal{O}_L(e_0)$ for some nilpotent element $e_0 \in \mathfrak{l}$.*

(ii) *Let L' be the standard Levi subgroup of G associated with a non-empty subset Π_0 of Π and let $D_0: \Pi_0 \rightarrow \{0, 1, 2\}$ be the weighted Dynkin diagram of a nilpotent orbit in $\mathfrak{l}' = \text{Lie}(L')$. If \tilde{D}_0 is the weighted Dynkin diagram of a nilpotent orbit in \mathfrak{g} then $\mathcal{O}(\tilde{D}_0) = \text{Ind}_\mathfrak{l}'^\mathfrak{g} \mathcal{O}_{L'}(D_0)$.*

Proof. It is clear that if $D(\alpha) = 2$ for some $\alpha \in \Pi$ then \mathfrak{l} is a proper Levi subalgebra of \mathfrak{g} . Set $\Pi(j) := \{\alpha \in \Pi \mid D(\alpha) = j\}$ where $j \in \{0, 1, 2\}$. If $\Pi = \{\alpha_1, \dots, \alpha_\ell\}$ and $\beta = \sum_{i=1}^\ell m_i \alpha_i$, where $m_i \in \mathbb{Z}$, then we put $v_i(\beta) := m_i$ for all $1 \leq i \leq \ell$. From the definition of $\lambda = \lambda_D$ it follows that $\mathfrak{g}(\lambda, i) = \mathfrak{l}(\lambda, i)$ for $i = 0, 1$ and $\mathfrak{g}(\lambda, \pm 2) = \mathfrak{l}(\lambda, \pm 2) \oplus \mathfrak{m}(\lambda, \pm 2)$ where $\mathfrak{m}(\lambda, \pm 2)$ is the k -span of all root vectors $e_\beta \in \text{Lie}(U_\beta)$ with $v_i(\beta) = 1$ for $i \in \Pi(2)$ and $v(\beta) = 0$ for $i \in \Pi(1)$.

Let $e \in \mathfrak{g}(2, \lambda)_\varphi$ and write $e = e_0 + e_1$ with $e_0 \in \mathfrak{l}(\lambda, 2)$ and $e_1 \in \mathfrak{m}(\lambda, 2)$. Since $\mathfrak{g}_e \subset \mathfrak{g}(\lambda, \geq 0)$ and $[e, \mathfrak{g}(\lambda, \geq 0)] = \mathfrak{g}(\lambda, \geq 2)$ by [23, Theorem 2.3], we have that

$$\dim \mathfrak{g}_e = \dim \mathfrak{g}(\lambda, 0) + \dim \mathfrak{g}(\lambda, 1) = \dim \mathfrak{l}(\lambda, 0) + \dim \mathfrak{l}(\lambda, 1).$$

Therefore, in order to prove the first part of the proposition it suffices to show that $\dim \mathfrak{l}_{e_0} = \dim \mathfrak{l}(\lambda, 0) + \dim \mathfrak{l}(\lambda, 1)$. Indeed, in that case the orbit $\text{Ind}_\mathfrak{l}^\mathfrak{g} \mathcal{O}_L(e_0)$ would intersect densely with $\mathfrak{g}(\lambda, \geq 2)$ and hence coincide with $\mathcal{O}(D)$.

Since $\Phi = \Phi_+(\Pi) \sqcup \Phi_-(\Pi)$ it is straightforward to see that $[\mathfrak{l}(\lambda, -i), \mathfrak{m}(\lambda, 2)] = 0$ for all $i \in \mathbb{Z}_{>0}$. It follows that the restrictions of $\text{ad } e$ and $\text{ad } e_0$ to $\mathfrak{l}(\lambda, -i)$ coincide for all such i . Since $\mathfrak{g}_e \subset \mathfrak{g}(\lambda, \geq 0)$ by [23, Theorem A] this entails $\mathfrak{l}_{e_0} \subseteq \mathfrak{l}(\lambda, \geq 0)$.

It is well known (and easily seen) that the restriction of κ to \mathfrak{l} is non-degenerate and the subspaces $\mathfrak{l}(\lambda, \pm i)$ are dual to each other with respect to κ . Since \mathfrak{l}_{e_0} coincides with the orthogonal complement of $[e_0, \mathfrak{l}]$ with respect to the restriction of κ to \mathfrak{l} and $\mathfrak{l}_{e_0} \subseteq \mathfrak{l}(\lambda, \geq 0)$ by the above, the map $\text{ad } e_0: \mathfrak{l}(\lambda, i) \rightarrow \mathfrak{l}(\lambda, i+2)$ must be surjective for all $i \in \mathbb{Z}_{\geq 0}$. From this it is immediate that $\dim \mathfrak{l}_{e_0} = \dim \mathfrak{l}(\lambda, 0) + \dim \mathfrak{l}(\lambda, 1)$ proving (i).

To prove (ii), we let λ' and λ be the cocharacters in $X_*(T)$ attached to D_0 and \tilde{D}_0 at the beginning of this subsection. It is immediate from the definition of \tilde{D}_0 that $\mathfrak{l}'(\lambda', 2) \subseteq \mathfrak{g}(\lambda, 2)$ and $\mathfrak{g}(\lambda, i) \cap \mathfrak{l}' = \mathfrak{l}'(\lambda', i)$ for $i = 1, 2$. Let V be the T -stable subspace of $\mathfrak{g}(\lambda, 2)$ complementary to $\mathfrak{l}'(\lambda', 2)$. Being an open map the first projection $\text{pr}_1: \mathfrak{g}(\lambda, 2) = \mathfrak{l}'(\lambda', 2) \oplus V \rightarrow \mathfrak{l}'(\lambda', 2)$ takes the Zariski open subset $\mathfrak{g}(\lambda, 2)_\varphi$ of $\mathfrak{g}(\lambda, 2)$ onto a dense subset of $\mathfrak{l}'(\lambda', 2)$. In view of [23, Theorem 2.3], this implies that there exists $e' \in \mathcal{O}_{L'}(D_0) \cap \mathfrak{l}'(\lambda', 2)$ such that $e' + v \in \mathfrak{g}(\lambda, 2)_\varphi$ for some $v \in V$. As \tilde{D}_0 is a weighted Dynkin diagram for \mathfrak{g} , [23] also shows that the cocharacter λ is optimal for $e := e' + v$ and

$$\dim \mathfrak{g}_e = \dim \mathfrak{g}(\lambda, 0) + \dim \mathfrak{g}(\lambda, 1) = \dim \mathfrak{l}'(\lambda', 0) + \dim \mathfrak{l}'(\lambda', 1) = \dim \mathfrak{l}'_{e'}.$$

Since it is straightforward to see that there exists a standard parabolic subgroup $P = L'R_u(P)$ of G such that $V \subseteq \text{Lie}(R_u(P))$, we now conclude that $e = e' + v \in \text{Ind}_V^{\mathfrak{g}} \mathcal{O}_{L'}(D_0)$. This finishes the proof. \square

Remark 3.2. For G exceptional, the second part of Proposition 3.1 is an immediate consequence of [20, Theorem 3(i)] which was proved by computational methods.

3.3. A sufficient condition for rigidity

Given a nilpotent element $e \in \mathfrak{g}$ we denote by \mathfrak{c}_e the factor space $\mathfrak{g}_e / [\mathfrak{g}_e, \mathfrak{g}_e]$ and set $c(\mathfrak{g}_e) := \dim \mathfrak{c}_e$. Our next result is a generalisation of [34, Proposition 11].

Proposition 3.3. *Assume, as before, that G satisfies the standard hypotheses and let $\mathfrak{l} = \text{Lie}(L)$ be a Levi subalgebra of \mathfrak{g} . If $e \in \text{Ind}_V^{\mathfrak{g}} \mathcal{O}_L(e_0)$ for some $e_0 \in \mathcal{N}(\mathfrak{l})$ then $c(\mathfrak{g}_e) \geq c(\mathfrak{l}_{e_0})$ and $\dim \mathfrak{z}(\mathfrak{g}_e) \geq \dim \mathfrak{z}(\mathfrak{l}_{e_0})$. Furthermore, if L has no simple components isomorphic to SL_{rp} , where $p = \text{char}(k)$, then $c(\mathfrak{g}_e) \geq c([\mathfrak{l}, \mathfrak{l}_{e_0}] + \dim \mathfrak{z}(\mathfrak{l}))$ and $\dim \mathfrak{z}(\mathfrak{g}_e) \geq \dim \mathfrak{z}(\mathfrak{l}) + \dim \mathfrak{z}([\mathfrak{l}, \mathfrak{l}_{e_0}])$.*

Proof. Put $\mathfrak{g}[t] := \mathfrak{g} \otimes_k k[t]$ and $\mathfrak{g}(t) := \mathfrak{g} \otimes_k k(t)$, where t is an indeterminate, and let $\pi: \mathfrak{g}[t] \rightarrow \mathfrak{g}$ denote the canonical projection induced by the augmentation map $k[t] \rightarrow k$. Let $P = L \cdot R_u(P)$ be a parabolic subgroup of G containing L and $\mathfrak{n}_+ := \text{Lie}(R_u(P))$. By our assumption, $\dim \mathfrak{g}_e = \dim \mathfrak{l}_{e_0}$ and we may assume without loss of generality that $e = e_0 + e_1$ for some $e_1 \in \mathfrak{n}_+$.

Let $h \in \mathfrak{z}(\mathfrak{l})_{\text{reg}}$ and consider $th + e$, an element of $\mathfrak{g}[t]$. Let $G(t)$ be the group of $k(t)$ -rational points of G . The argument used in the proof of Theorem 2.3 then shows that $th + e$ is $G(t)$ -conjugate to $th + e_0$ (in fact, a conjugating element can be found already in the group of $k(t)$ -rational points of $R_u(P)$). From this it is immediate that $\mathfrak{g}(t)_{th+e} \cong \mathfrak{l}_{e_0} \otimes_k k(t)$ as Lie algebras over $k(t)$. It follows that

$$\dim_{k(t)}[\mathfrak{g}(t)_{th+e}, \mathfrak{g}(t)_{th+e}] = \dim[\mathfrak{l}_{e_0}, \mathfrak{l}_{e_0}],$$

$$\dim_{k(t)} \mathfrak{z}(\mathfrak{g}(t)_{th+e}) = \dim \mathfrak{z}(\mathfrak{l}_{e_0}).$$

549 Furthermore, if \mathfrak{l} has no components of type A_{rp-1} , our discussion in § 2.1 implies that
 550 $\dim[\mathfrak{l}_{e_0}, \mathfrak{l}_{e_0}] = \dim[\mathfrak{l}, \mathfrak{l}]_{e_0}$ and $\dim \mathfrak{z}(\mathfrak{l}_{e_0}) = \dim \mathfrak{z}(\mathfrak{l}) + \dim \mathfrak{z}([\mathfrak{l}, \mathfrak{l}]_{e_0})$.

551 Put $M_1 := \mathfrak{g}(t)_{th+e} \cap \mathfrak{g}[t]$, $M_2 := [\mathfrak{g}(t)_{th+e}, \mathfrak{g}(t)_{th+e}] \cap \mathfrak{g}[t]$ and $M_3 := \mathfrak{z}(\mathfrak{g}(t)_{th+e}) \cap \mathfrak{g}[t]$.
 552 Each of these is a $k[t]$ -submodule of the free $k[t]$ -module $\mathfrak{g}[t]$, and it is not hard to check
 553 that all factor modules $\mathfrak{g}[t]/M_i$ are torsion-free. Since $k[t]$ is a principal ideal domain
 554 and the $k[t]$ -module $\mathfrak{g}[t]$ is finitely generated, it follows that all $\mathfrak{g}[t]/M_i$ are free over $k[t]$.
 555 As a consequence, there exist free $k[t]$ -submodules N_i of $\mathfrak{g}[t]$ such that $\mathfrak{g}[t] = N_i \oplus M_i$
 556 as $k[t]$ -modules, where $i = 1, 2, 3$. This, in turn, shows that each M_i is free over $k[t]$ and
 557 $\mathrm{rk}_{k[t]} M_i = \dim_{k(t)} M_i \otimes_{k[t]} k(t)$.

558 It is straightforward to see that $\pi: \mathfrak{g}[t] \rightarrow \mathfrak{g}$ maps M_1 into \mathfrak{g}_e . Since $e \in \mathrm{Ind}_1^{\mathfrak{g}} \mathcal{O}_L(e_0)$
 559 we have that

$$560 \quad \dim \mathfrak{g}_e = \dim \mathfrak{l}_{e_0} = \dim_{k(t)} \mathfrak{g}(t)_{th+e} = \dim_{k(t)} M_1 \otimes_{k[t]} k(t) = \mathrm{rk}_{k[t]} M_1.$$

561 From this it follows that M_1 has a free basis $v_1(t), \dots, v_r(t)$ such that
 562 $\pi(v_1(t)), \dots, \pi(v_r(t))$ form a k -basis of \mathfrak{g}_e . For $1 \leq i \leq r$ put $v_i := \pi(v_i(t))$ and observe
 563 that

$$564 \quad \mathrm{rk}_{k[t]} M_2 = \dim M_2 \otimes_{k[t]} k(t) = \dim_{k(t)} [\mathfrak{g}(t)_{th+e}, \mathfrak{g}(t)_{th+e}] = \dim[\mathfrak{l}_{e_0}, \mathfrak{l}_{e_0}].$$

565 Since $[v_i(t), v_j(t)] \in M_2$ and $\pi([v_i(t), v_j(t)]) = [v_i, v_j]$ for $1 \leq i, j \leq r$,
 566 this gives $\dim[\mathfrak{g}_e, \mathfrak{g}_e] \leq \dim[\mathfrak{l}_{e_0}, \mathfrak{l}_{e_0}]$. Therefore, $c(\mathfrak{g}_e) = \dim \mathfrak{g}_e - \dim[\mathfrak{g}_e, \mathfrak{g}_e] \geq \dim \mathfrak{l}_{e_0} -$
 567 $\dim[\mathfrak{l}_{e_0}, \mathfrak{l}_{e_0}] = c(\mathfrak{l}_{e_0})$.

568 Finally, we observe that $\pi(M_3) \subseteq \mathfrak{z}(\mathfrak{g}_e)$ and

$$569 \quad \mathrm{rk}_{k[t]} M_3 = \dim_{k(t)} M_3 \otimes_{k[t]} k(t) = \dim_{k(t)} \mathfrak{z}(\mathfrak{g}(t)_{th+e}) = \dim \mathfrak{z}(\mathfrak{l}_{e_0}).$$

570 Consequently, $\dim \mathfrak{z}(\mathfrak{g}_e) \geq \dim \mathfrak{z}(\mathfrak{l}_{e_0})$. The second part of the proposition now follows from
 571 our remarks earlier in the proof. \square

572 Proposition 3.3 gives us a useful sufficient condition for rigidity of nilpotent elements
 573 in exceptional Lie algebras.

574 **Corollary 3.4.** *Suppose G is simple and p is good for G . If $e \in \mathcal{N}(\mathfrak{g})$ is such that $\mathfrak{g}_e =$
 575 $[\mathfrak{g}_e, \mathfrak{g}_e]$ then e is rigid in \mathfrak{g} .*

576 **Proof.** We may assume without loss that G is simply connected. For G classical
 577 the statement follows by comparing [26, Theorem 3] with the Kempken–Spaltenstein
 578 description of rigid nilpotent orbits in \mathfrak{g} . (Of course, in the characteristic zero
 579 case one can apply [34, Proposition 11] directly.)

580 So suppose from now that G is exceptional and p is a good prime for G . Since the Killing
 581 form of \mathfrak{g} is non-degenerate, G satisfies the standard hypotheses. By rank considerations,
 582 there are no Levi subgroups in G with components of type A_{rp-1} for $r \geq 2$.

583 Suppose $e \in \mathrm{Ind}_1^{\mathfrak{g}} \mathcal{O}_L(e_0)$ where L is a proper Levi subgroup of G and $e_0 \in \mathcal{N}(\mathfrak{l})$. Since
 584 $c(\mathfrak{g}_e) = 0$ in the present case, Proposition 3.3 yields that L has a component of type
 585 A_{p-1} . Then all components of L have type A. Let T be a maximal torus of L and
 586 $\mathfrak{t} = \mathrm{Lie}(T)$. We may assume that L is associated with a subset Π_0 of a basis of simple
 587 roots Π of the root system $\Phi(G, T)$. Since all components of L have type A we may

also assume without loss that $e_0 = \sum_{\alpha \in \Pi_1} e_\alpha$ for some subset Π_1 of Π_0 (possibly empty). Let $\mathfrak{t}_0 := \mathfrak{t} \cap [\mathfrak{l}, \mathfrak{l}]$ and $\mathfrak{t}_1 := \bigcap_{\alpha \in \Pi_1} \ker(d\alpha)_e$. Then $\mathfrak{t}_0 \cap \mathfrak{t}_1$ consists of all $x \in \mathfrak{t}_0$ such that $(d\alpha)_e(x) = 0$ for all $\alpha \in \Pi_1$. Since all simple components of L are simply connected, looking at the p -ranks of the Cartan matrices associated with Π and Π_0 one observes that $\dim \mathfrak{t}_0 = |\Pi_0|$, $\dim \mathfrak{t}_1 = \dim \mathfrak{t} - |\Pi_1|$ and $\dim(\mathfrak{t}_0 \cap \mathfrak{t}_1) = |\Pi_0| - |\Pi_1|$ unless one of the components of Π_1 has type A_{p-1} in which case $\dim(\mathfrak{t}_0 \cap \mathfrak{t}_1) = |\Pi_0| - |\Pi_1| + 1$.

If Π_1 does not contain components of type A_{p-1} the above shows that $\dim \mathfrak{t}_1 = |\Pi| - |\Pi_1| > |\Pi_0| - |\Pi_1| = \dim \mathfrak{t}_0 \cap \mathfrak{t}_1$. Hence, there exists $h \in \mathfrak{t}$ such that $h \notin [\mathfrak{l}, \mathfrak{l}]$ and $[h, e_0] = 0$. But then $\mathfrak{l}_{e_0} \supsetneq [\mathfrak{l}_{e_0}, \mathfrak{l}_{e_0}]$, so that $c(\mathfrak{l}_{e_0}) \geq 1$. Since this contradicts Proposition 3.3, we see that Π_1 must have a component of type A_{p-1} . Since Π_0 has such a component too, they must contain the same component of type A_{p-1} . Let L_1 be the simple component of L generated by the root subgroups $U_{\pm\alpha}$ with $\alpha \in \Pi_1$. Then $L_1 \cong \mathrm{SL}_p$ as algebraic groups. The Lie algebra \mathfrak{l} acts on its ideal $\mathfrak{l}_1 := \mathrm{Lie}(L_1)$ by derivations, giving a natural Lie algebra homomorphism $\psi: \mathfrak{l} \rightarrow \mathrm{Der}(\mathfrak{l}_1)$. Since $\mathfrak{l}_1 \cong \mathfrak{sl}_p$ we may identify $\mathrm{Der}(\mathfrak{l}_1)$ with \mathfrak{pgl}_p . Our earlier remarks now imply that $\psi(e_0)$ is a regular nilpotent element in \mathfrak{pgl}_p and hence has an abelian centraliser in \mathfrak{pgl}_p . As ψ maps \mathfrak{l}_{e_0} onto a non-zero subalgebra of the centraliser of $\psi(e_0)$ in \mathfrak{pgl}_p , the Lie algebra \mathfrak{l}_{e_0} has a non-trivial abelian quotient. But then \mathfrak{l}_{e_0} is not perfect, i.e., $c(\mathfrak{l}_{e_0}) \geq 1$. Since this contradicts Proposition 3.3, the element e is rigid in \mathfrak{g} . \square

3.4. Optimal cocharacters of nilpotent elements contained in regular subalgebras

In this subsection, we assume that G is an arbitrary connected reductive group defined over an algebraically closed field of good characteristic. A connected reductive subgroup K of G is called *regular* if it contains a maximal torus of G . We denote by \mathfrak{k} the Lie algebra of K . If T is a maximal torus of G contained in K then the root system Φ_0 of K with respect to T identifies with a root subsystem of the root system $\Phi = \Phi(G, T)$ and $\mathfrak{k} = \mathfrak{t} \oplus \sum_{\alpha \in \Phi_0} \mathfrak{g}_\alpha$ where $\mathfrak{t} = \mathrm{Lie}(T)$. Since \mathfrak{k} is a restricted Lie subalgebra of \mathfrak{g} any nilpotent element of \mathfrak{k} is contained in $\mathcal{N}(\mathfrak{g})$, the nilpotent cone of \mathfrak{g} .

If $\mathrm{char}(k) = 0$ then a standard argument involving \mathfrak{sl}_2 -triples shows that any non-zero nilpotent element $e \in \mathfrak{k}$ admits a rational cocharacter $\lambda: k^\times \rightarrow G$ which is optimal for e in the sense of the Kempf–Rousseau theory and has the property that $\lambda(k^\times) \subset K$. Our next goal is to show that this result still holds under our assumption on k . Given a Zariski closed subgroup H of G we write $X_*(H)$ for the set of all rational cocharacters $\lambda: k^\times \rightarrow H$ and we denote by $P_H(\lambda)$ the parabolic subgroup of H associated with $\lambda \in X_*(H)$. If e is a non-zero nilpotent element of \mathfrak{k} then we denote by $\hat{\Lambda}_K(e)$ (respectively, $\hat{\Lambda}_G(e)$) the set of all $\lambda \in X_*(K)$ (respectively, $\lambda \in X_*(G)$) which are optimal for e regarded as a K -unstable element of \mathfrak{k} (respectively, a G -unstable element of \mathfrak{g}).

Proposition 3.5. *If K is a regular reductive subgroup of G and $p = \mathrm{char}(k)$ is a good prime for G then for any non-zero nilpotent element of $e \in \mathfrak{k}$ we have the inclusion $\hat{\Lambda}_K(e) \subseteq \hat{\Lambda}_G(e)$.*

Proof. In proving this proposition we may assume without loss of generality that K is a maximal connected reductive subgroup of G . Thanks to [22, Lemma 14] we may also assume that G is a semi-simple group of adjoint type. Let Φ be the root system of G with respect to a maximal torus T contained in K and let Φ_0 be the set of roots of K with respect to T . The maximality of K then shows that Φ_0 is a maximal root subsystem of Φ . Choose a basis of simple roots Π in Φ and let $\tilde{\alpha}_1, \dots, \tilde{\alpha}_s$ be all highest roots of Φ with respect to Π (here s is the number of the simple components of G). For $1 \leq i \leq s$ write $\tilde{\alpha}_i = \sum_{\alpha \in \Pi} n(\alpha, i)\alpha$. In view of the Borel–de Siebenthal theorem, no generality will be lost by assuming that either K is a maximal standard Levi subgroup of G or there exist $\alpha_0 \in \Pi$ and a prime number q dividing $n(\alpha_0, i_0)$, where i_0 is the unique index $j \leq s$ for which $n(\alpha_0, j) \neq 0$, such that $\Phi_0 = \{\gamma = \sum_{\alpha \in \Pi} r_\gamma(\alpha)\alpha \in \Phi \mid q \text{ divides } r_\gamma(\alpha_0)\}$.

It is well known that there is a canonical duality $\langle \cdot, \cdot \rangle$ between $X_*(T)$ and the lattice of rational characters $X^*(T)$. Let $\{\varpi_\alpha^\vee \mid \alpha \in \Pi\} \subset X_*(T)$ be the system of fundamental coweights corresponding to $\Pi \subset X^*(T)$. It has the property that $\langle \varpi_\alpha^\vee, \beta \rangle = \delta_{\alpha, \beta}$ for all $\alpha, \beta \in \Pi$. Since G is a group of adjoint type, for every $\alpha \in \Pi$ there is a 1-parameter subgroup $\varpi_\alpha^\vee(k^\times)$ in T such that

$$(\text{Ad } \varpi_\alpha^\vee(t))(e_\gamma) = t^{r_\gamma(\alpha)} e_\gamma \quad (\forall \gamma \in \Phi, \forall e_\gamma \in \mathfrak{g}_\gamma, \forall t \in k^\times). \quad (5)$$

Since p is a good prime for G we have that $q \neq \text{char}(k)$. Hence, k^\times contains a q th primitive root of 1. Thanks to (5) and the preceding discussion this implies that T contains an element σ such that \mathfrak{k} coincides with \mathfrak{g}^σ , the Lie algebra of the fixed point group G^σ .

Let $\lambda_0 \in \hat{\Lambda}_K(e)$ and $\lambda \in \hat{\Lambda}_G(e)$. As $\mathfrak{k} = \mathfrak{g}^\sigma$ and $G_e \subset P_G(\lambda)$ by the optimality of $P_G(\lambda)$ it must be that $\sigma \in P_G(\lambda)$. As $T \subset K$, the element σ lies in the centre of K . As K is connected, the latter is contained in any parabolic subgroup of K . So $\sigma \in P_K(\lambda_0) \subset P_G(\lambda_0)$. But then σ belongs to the connected group $P_G(\lambda) \cap P_G(\lambda_0)$ and therefore lies in a maximal torus of $P_G(\lambda) \cap P_G(\lambda_0)$; we call it T' . Note that $T' \subset K$ because T' is a connected group commuting with σ .

Since T' is a maximal torus of $P_G(\lambda)$, the set $X_*(T') \cap \hat{\Lambda}_G(e)$ is non-empty; see [23, Theorem 2.1(iii)], for example. We claim that $X_*(T') \cap \hat{\Lambda}_G(e) \subset \hat{\Lambda}_K(e)$. To prove this we adopt the notation and conventions of [23, 2.2] and recall that there exists $\lambda' \in X_*(T') \cap \hat{\Lambda}_G(e)$ for which $m(\lambda', e) = 2$; see [23, Theorems 2.3(i) and 2.7]. Since $\lambda' \in \hat{\Lambda}_G(e)$ we have that

$$\frac{m(\lambda', e)}{\|\lambda'\|} \geq \frac{m(\mu, e)}{\|\mu\|} \quad (\forall \mu \in \hat{\Lambda}_K(e)).$$

But it also follows from [23] that there is a $\mu' \in \hat{\Lambda}_K(e)$ for which $m(\mu', e) = 2$ (here we regard e as an element of \mathfrak{k}). From this it is immediate that $\lambda' \in \hat{\Lambda}_K(e)$. Since the quantity $m(\mu, e)/\|\mu\|$ is independent of the choice of $\mu \in \hat{\Lambda}_K(e)$ the claim follows and completes the proof. \square

Working over \mathbb{C} , de Graaf and Elashvili determine the weighted Dynkin diagram of the unique nilpotent orbit contained in a sheet \mathcal{S} of an exceptional Lie algebra $\mathfrak{g}_{\mathbb{C}}$ and give a nice representative $e_{\Gamma, \mathbb{C}} = \sum_{\gamma \in \Gamma} e_{\gamma, \mathbb{C}}$ in that orbit. Here $\Gamma = \Gamma(\mathcal{S})$ is a subset of roots of the root system Φ of $\mathfrak{g}_{\mathbb{C}}$ and $e_{\gamma, \mathbb{C}}$ is a root vector of $\mathfrak{g}_{\mathbb{C}}$ corresponding to $\gamma \in \Phi$. Since each set Γ consists of linearly independent roots, the $G_{\mathbb{C}}$ -orbit of $e_{\Gamma, \mathbb{C}}$ is independent of the choices of root vectors $e_{\gamma, \mathbb{C}}$ (in the sense that each of them can be rescaled). In [8],

the roots in each set Γ are labelled by their positions in the ordering of positive roots used in GAP4.²

To each set Γ de Graaf and Elashvili assign a diagram $D(\Gamma)$. Specifically, the number of nodes of $D(\Gamma)$ equals the cardinality of Γ and the nodes depicting distinct $\gamma, \gamma' \in \Gamma$ are linked by $\langle \gamma, \gamma' \rangle \cdot \langle \gamma', \gamma \rangle$ edges. The edges are solid (respectively dotted) if $\langle \gamma, \gamma' \rangle$ is negative (respectively positive). If Φ has roots of two different lengths then the nodes of $D(\Gamma)$ corresponding to long roots of Φ are coloured in black.

Since $\mathfrak{g} = \text{Lie}(G)$ contains a natural analogue e_Γ of $e_{\Gamma, \mathbb{C}}$ we wonder whether the nilpotent orbits of e_Γ and $e_{\Gamma, \mathbb{C}}$ have the same weighted Dynkin diagram. In the next section, we give a positive answer to this question by computational methods, but our next result indicates that in many cases such computations can be avoided.

Corollary 3.6. *Suppose that p is a good prime for G , all roots in Φ have the same length, and Γ is such that $\gamma - \gamma' \notin \Phi$ for all distinct $\gamma, \gamma' \in \Gamma$. If $D(\Gamma)$ is a disjoint union of Dynkin graphs then the nilpotent orbits in \mathfrak{g} and $\mathfrak{g}_{\mathbb{C}}$ containing e_Γ and $e_{\Gamma, \mathbb{C}}$, respectively, have the same labels and dimensions.*

Proof. We may assume that Φ is the root system of G with respect to a maximal torus T of G and Γ is contained in the positive system $\Phi_+(\Pi)$ associated with a basis of simple roots Π of Φ . Let W_0 be the subgroup of the Weyl group $W(\Phi)$ generated by all reflections s_γ with $\gamma \in \Gamma$ and put $\Phi(\Gamma) := \{w(\gamma) \mid w \in W_0, \gamma \in \Gamma\}$. Using our assumptions on Γ it is straightforward to see that $\Phi(\Gamma) = \Phi_+(\Gamma) \sqcup -\Phi_+(\Gamma)$ where $\Phi_+(\Gamma)$ consists of roots in Φ which can be presented as linear combinations of elements in Γ with coefficients in $\mathbb{Z}_{\geq 0}$. It follows that $\Phi(\Gamma)$ is a root system in the \mathbb{Q} -span of Γ and Γ is a basis of simple roots in $\Phi(\Gamma)$.

Let K be the connected subgroup of G generated by T and all root subgroups $U_{\pm\gamma}$ with $\gamma \in \Phi(\Gamma)$. The preceding discussion shows that the quadruple $(X^*(T), \Phi(\Gamma), \Phi(\Gamma)^\vee, X_*(T))$ is a root datum for K . In particular, this implies that $\Phi(\Gamma)$ is the root system of K with respect to T . Since Γ is a basis of simple roots of $\Phi(\Gamma)$, we now see that e_Γ is a regular nilpotent element of the Lie algebra $\mathfrak{k} = \text{Lie}(K)$. Proposition 3.5 shows that the set $\hat{\Lambda}_K(e_\Gamma)$ consists of optimal cocharacters of $e_\Gamma \in \mathfrak{g}$. On the other hand, it is well known that $\hat{\Lambda}_K(e_\Gamma)$ contains a unique element $\lambda = \sum_{\gamma \in \Gamma} a_\gamma \gamma^\vee \in X_*(T)$ with $a_\gamma \in \mathbb{Z}$ which satisfies the conditions $\langle \lambda, \gamma \rangle = 2$ for all $\gamma \in \Gamma$. Moreover, the coefficients a_γ are independent of p and λ lies in $\hat{\Lambda}_{G_{\mathbb{C}}}(e_{\Gamma, \mathbb{C}})$ when regarded as an element of $X_*(G_{\mathbb{C}})$. In view of [23, 2.4] this yields that the nilpotent orbits $\mathcal{O}(e_\Gamma) \subset \mathfrak{g}$ and $\mathcal{O}(e_{\Gamma, \mathbb{C}}) \subset \mathfrak{g}_{\mathbb{C}}$ have the same labels and dimensions. \square

3.5. Comparing rigid orbits in \mathfrak{g} and $\mathfrak{g}_{\mathbb{C}}$

From now on we assume that G is a simply connected algebraic group of type G_2, F_4, E_6, E_7 or E_8 and denote by $G_{\mathbb{C}}$ the complex counterpart of G . We identify the root system Φ of G with that of $G_{\mathbb{C}}$ and let Π be a basis of simple roots. Since p is good for G , the nilpotent orbits in \mathfrak{g} and $\mathfrak{g}_{\mathbb{C}} = \text{Lie}(G_{\mathbb{C}})$ are parametrised by their weighted Dynkin

²We are grateful to Simon Goodwin for bringing this to our attention.

709 diagrams $D: \Pi \rightarrow \{0, 1, 2\}$. We let $\mathcal{O}_{\mathbb{C}}(D)$ be the nilpotent orbit in $\mathfrak{g}_{\mathbb{C}}$ corresponding to
 710 D . It follows from [23, Theorem 2.3], for example, that $\dim_k \mathcal{O}(D) = \dim_{\mathbb{C}} \mathcal{O}_{\mathbb{C}}(D)$. We
 711 shall denote this number by $N(D)$.

712 Combining [7] and [8] with the results of the previous subsection we obtain the
 713 following:

714 **Lemma 3.7.** *Let $D: \Pi \rightarrow \{0, 1, 2\}$ be a weighted Dynkin diagram. If the orbit $\mathcal{O}_{\mathbb{C}}(D)$ is*
 715 *rigid in $\mathfrak{g}_{\mathbb{C}}$ then the orbit $\mathcal{O}(D)$ is rigid in \mathfrak{g} .*

716 **Proof.** Suppose $\mathcal{O}_{\mathbb{C}}(D)$ is rigid in $\mathfrak{g}_{\mathbb{C}}$ and pick $e' \in \mathcal{O}_{\mathbb{C}}(D)$ and $e \in \mathcal{O}(D)$. It follows from
 717 [7] that either the centraliser of e' in $\mathfrak{g}_{\mathbb{C}}$ is perfect or $\mathcal{O}_{\mathbb{C}}(D)$ has one of the following
 718 types:

- 719 (a) \tilde{A}_1 in type G_2 ;
- 720 (b) $\tilde{A}_2 + A_1$ in type F_4 ;
- 721 (c) $(A_3 + A_1)'$ in type E_7 ;
- 722 (d) $A_3 + A_1, A_5 + A_1$ or $D_5(a_1) + A_2$ in type E_8 .

723 If the centraliser of e' in $\mathfrak{g}_{\mathbb{C}}$ is perfect then it follows from Table 1 that the same holds
 724 for the centraliser of e in \mathfrak{g} . In that case the orbit $\mathcal{O}(D)$ is rigid in \mathfrak{g} by Corollary 3.4.

725 If $\mathcal{O}(D)$ is listed in parts (a), (c), (d) then analysing the tables in [6, pp. 401–407] one
 726 finds that either $\mathcal{O}(D)$ is the only orbit of dimension $N(D)$ in \mathfrak{g} or G is of type E_8 and
 727 $\mathcal{O}(D)$ is one of $A_5 + A_1$ or $D_5(a_1) + A_2$. If $\mathcal{O}(D)$ is the only orbit of dimension $N(D)$ in
 728 \mathfrak{g} then it cannot be induced from a proper Levi subalgebra, because otherwise the same
 729 would be true for $\mathcal{O}_{\mathbb{C}}(D)$ by Theorem 2.8. In type E_8 , there are two nilpotent orbits of
 730 dimension 202 and they have types $A_5 + A_1$ and $D_5(a_1) + A_2$. Moreover, both are rigid
 731 in characteristic 0 by [8]. Therefore, their counterparts in $\mathcal{N}(\mathfrak{g})$ cannot be induced from
 732 proper Levi subalgebras by Theorem 2.8.

733 Finally, let $\mathcal{O}(D)$ be as in part (b). Then $N(D) = 36$ and $c(\mathfrak{g}_e) = 1$ by Table 3. Suppose
 734 for a contradiction that the orbit $\mathcal{O}(D)$ is induced from a nilpotent orbit \mathcal{O}_0 in a proper
 735 Levi subalgebra $\mathfrak{l} = \text{Lie}(L)$ of \mathfrak{g} . Since in the present case G has type F_4 and p is good for
 736 G , the Levi subgroup L cannot have components of type A_{rp-1} . Applying Proposition
 737 3.3 we then get $c(\mathfrak{l}, \mathfrak{l}_{e_0}) + \dim \mathfrak{z}(\mathfrak{l}) \leq 1$ which implies that $\dim \mathfrak{z}(\mathfrak{l}) = 1$ and $c(\mathfrak{l}_{e_0}, \mathfrak{l}_{e_0}) = 0$
 738 for any $e_0 \in \mathcal{O}_0$. Since L has no components of exceptional types, the orbit \mathcal{O}_0 must
 739 be rigid in \mathfrak{l} by [26, Theorem 3(i)]. Combining Theorem 2.8 and Proposition 2.5 we
 740 conclude that $\mathcal{O}(D)$ lies in a sheet of \mathfrak{g} of dimension $N(D) + \dim \mathfrak{z}(\mathfrak{l}) = 37$. But Theorem
 741 2.8 also shows that there is a bijection $\mathcal{S} \rightarrow \mathcal{S}_{\mathbb{C}}$ between the sheets of \mathfrak{g} and $\mathfrak{g}_{\mathbb{C}}$ such that
 742 $\dim_k \mathcal{S} = \dim_{\mathbb{C}} \mathcal{S}_{\mathbb{C}}$ for all sheets \mathcal{S} of \mathfrak{g} . By [8, Table 10], there exists only one sheet of
 743 dimension 37 in $\mathfrak{g}_{\mathbb{C}}$ and the nilpotent orbit contained in it has type B_2 . Using Proposition
 744 3.1(ii) it is not hard to observe that the orbit of that type in \mathfrak{g} is induced from the minimal
 745 nilpotent orbit in a Levi subalgebra of type C_3 . In view of Remark 2.6 this implies that
 746 $\mathcal{O}(D)$ cannot lie in the unique sheet of dimension 37 in \mathfrak{g} . By contradiction the result
 747 follows. \square

748 We are now in a position to prove one of the main results of this paper:

Theorem 3.8. *The orbit $\mathcal{O}(D)$ is rigid in \mathfrak{g} if and only if the orbit $\mathcal{O}_{\mathbb{C}}(D)$ is rigid in $\mathfrak{g}_{\mathbb{C}}$.* 749
750

Proof. Let $D: \Pi \rightarrow \{0, 1, 2\}$ be a weighted Dynkin diagram such that the orbit $\mathcal{O}(D)$ is rigid in \mathfrak{g} . In view of Lemma 3.7 we need to show that the orbit $\mathcal{O}_{\mathbb{C}}(D)$ is rigid in $\mathfrak{g}_{\mathbb{C}}$. So suppose $\mathcal{O}_{\mathbb{C}}(D)$ is induced from a proper Levi subalgebra of $\mathfrak{g}_{\mathbb{C}}$. Then $\mathcal{O}(D) \neq \{0\}$ and thanks to Proposition 3.1(i) we may assume that $D(\Pi) = \{0, 1\}$. If $\mathcal{O}_{\mathbb{C}}(D)$ is the only orbit of dimension $N(D)$ in $\mathfrak{g}_{\mathbb{C}}$ then Theorem 2.8 implies that the orbit $\mathcal{O}(D)$ must be induced in \mathfrak{g} . 751
752
753
754
755
756

Thus, we may assume from now that the orbit $\mathcal{O}_{\mathbb{C}}(D)$ is odd, induced, and there are at least two nilpotent orbits of dimension $N(D)$ in $\mathfrak{g}_{\mathbb{C}}$. Looking at the tables in [6, pp. 401, 402] one finds out that this never happens in types G_2, F_4 and E_6 . In type E_7 , there exist only two orbits fulfilling these conditions, namely, $A_3 + 2A_1$ and $D_4(a_1) + A_1$. In type E_8 we have to examine more closely the following orbits: $A_4 + A_1, A_4 + A_2 + A_1, D_6(a_2), E_6(a_3) + A_1, A_6 + A_1, A_7$ and $D_7(a_2)$. 757
758
759
760
761
762

If G is of type E_8 and $\mathcal{O}(D)$ has type $A_4 + A_1$ then there are two nilpotent orbits in \mathfrak{g} (and $\mathfrak{g}_{\mathbb{C}}$) of dimension $N(D) = 188$. One of these orbits is rigid in $\mathfrak{g}_{\mathbb{C}}$ and hence in \mathfrak{g} by Lemma 3.7 (it has type $2A_3$). On the other hand, combining Theorem 2.8 with [8, Table 8] one observes that \mathfrak{g} must have an *induced* orbit of dimension 188. This rules out the case where $\mathcal{O}(D)$ has that type. Looking at the diagrams $D(\Gamma)$ attached in [8] to the orbits labelled $A_3 + 2A_1$ in type E_7 and $A_4 + A_2 + A_1, E_6(a_3) + A_1, A_6 + A_1$ and A_7 in type E_8 one finds out that they satisfy the conditions of Corollary 3.6. In view of Proposition 3.5 [8] this means that these orbits cannot be rigid in \mathfrak{g} . 763
764
765
766
767
768
769
770

Since there was no obvious way to deal with remaining three cases in types E_7 and E_8 these have been checked through GAP. As our calculations in GAP in fact prove the stronger result of Theorem 1.4 without the need to use the present result, we leave the details of this calculation until § 4.2. \square 771
772
773
774
775

4. Further computations with nilpotent orbits 775

In this section, we describe routines performed in GAP which extend those of [7] and [8] to positive characteristic, in particular, justifying the tables at the end of the document. The basic problem in obtaining results in arbitrary characteristic is to reduce to a finite list the prime characteristics which must be considered separately. For this, the idea is always to produce an integral matrix such that it encodes this list via the prime divisors of its elementary divisors. 776
777
778
779
780
781

4.1. Nilpotent orbit representatives 782

For our calculations we require (i) a list of nilpotent orbit representatives for bad primes; (ii) a list of orbit representatives for good primes with associated cocharacters. 783
784

Those in (i) are provided by [19] or [30] (for F_4). Regarding notation, particularly that which appears in the tables at the end of the paper, the labels for the nilpotent elements 785

used are now quite standard and consistent with [19]. A nilpotent orbit has type X_r if it is regular in a Levi subalgebra whose root system has type X_r , using $+$ to indicate a union of irreducible root systems. Where there are short roots in the root system of \mathfrak{g} , we use $\tilde{}$ to indicate a Levi subalgebra with such roots. If a nilpotent orbit is distinguished in a Levi subalgebra of the same type it is labelled $X_r(a_i)$ or $X_r(b_i)$ for some value i . Occasionally in E_7 , there are two conjugacy classes of Levi subalgebras giving rise to two regular nilpotent orbits and these are denoted, for example, as $(A_5)'$ and $(A_5)''$. Finally, there are extra orbits which appear over fields of characteristics 2 and 3 and have no analogue over \mathbb{C} . These are closely related to certain orbits of type X_r , for some X and r , and are denoted $X_r^{(p)}$ for p being 2 or 3.

(ii) For this we use [20] which provides all the data we require.

Since there are a number of errors in the literature related to nilpotent orbits, we have made reasonable attempts to check the validity of the orbit representatives we use. For example, we checked that they had the same Jordan blocks as found in [32, 33] (with certain exceptions³) and [16] (in the group case). For the exceptional groups considered by [32], the paper only deals with the elements of the *restricted* nullcone $\mathcal{N}_p(\mathfrak{g}) = \{x \in \mathfrak{g} \mid x^{[p]} = 0\}$, but in [33] the authors refer to an unpublished result of Lawther which treats the case of an arbitrary nilpotent element of \mathfrak{g} . Therefore, our computations provide an independent verification of the Jordan block structure of those nilpotent elements $x \in \mathfrak{g}$ for which $x^{[p]} \neq 0$ (of course, such elements exist only when p is less than or equal to the Coxeter number of G). It turned out that as in the group case (investigated in [16]) there are no coincidences in Jordan block decompositions of representatives of different nilpotent orbits in \mathfrak{g} except when $p = 7$ where the Jordan block decompositions of nilpotent elements of type B_3 and C_3 in Lie algebras of type F_4 are the same. As this result is likely to be useful in future work, we state it formally.

Theorem 4.1 (McNinch, Lawther). *Let G be a connected reductive k -group satisfying the standard hypotheses and $\mathfrak{g} = \text{Lie}(G)$. Then the Jordan block structure of a nilpotent element of \mathfrak{g} on the adjoint module coincides with that of a unipotent element of G with the same Dynkin label.*

Proof. Since G satisfies the standard hypotheses, p is a good prime for G . Our discussion in §2.1 shows that no generality will be lost by assuming that either G is a simple exceptional algebraic group or G is a classical group of type other than $A_{r,p-1}$ or $G = GL_{rp}$. In the first case the theorem follows from Lawther's and our computer-aided calculations whilst in the other two cases it was deduced by McNinch [21] from earlier results of Fossum on formal group laws; see [11]. \square

It would be interesting to find a non-computational and case-free proof of Theorem 4.1. Finally, all our results recover those of De Graaf for generic prime characteristic.

³There are a number of errors in the representatives and associated Jordan blocks listed in [33]. Some of these relate to the transcription of representatives from MAGMA into LaTeX; in addition, representatives of type $D_r(a_i)$ and $E_8(b_6)$ are used which break down in certain low characteristics. This means that there are a number of mistakes in the calculation of adjoint Jordan blocks of *op. cit.* for types E_6 , E_7 and E_8 for $p = 2$ and $p = 3$. Corrected Jordan blocks for these low characteristics are expected to appear in a forthcoming publication by the second author.

4.2. Induced nilpotent representatives

Here we show that the nilpotent orbit representatives given in the tables of [8] are still valid in good characteristic and use the results of the previous section to prove this. Let $e \in \mathfrak{g}_{\mathbb{Z}}$ be a nilpotent orbit representative as given in the tables of [8]. We proceed by having GAP compute the adjoint matrix of each representative over \mathbb{Q} according to a Chevalley basis of $\mathfrak{g}_{\mathbb{Z}}$. Then a routine computes the integral divisors of successive powers of the matrix. For any prime dividing the integral divisors we know that the Jordan block structure of $\text{ad } e$ for $e \in \mathfrak{g} = \mathfrak{g}_{\mathbb{Z}} \otimes_{\mathbb{Z}} k$ differs from that in characteristic zero. Let this finite list of primes be S . If $p \notin S$ then since the Jordan block structure of $\text{ad } e$ determines it in good characteristic by [16], we are done in this case. For each good prime of S we have GAP compute the Jordan block structure separately. We find in all cases that the Jordan block structure is the same as that of a nilpotent element in $\mathfrak{g}_{\mathbb{C}} = \mathfrak{g}_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathbb{C}$ with the same label, as listed in [16].⁴

Proof of Theorem 1.4. Each nilpotent element of [8] is of the form $e_{\Gamma} = e_0 + e_1$ where $e_0 \in \mathfrak{l}$ is a rigid nilpotent element of \mathfrak{l} and $e_1 \in \mathfrak{n}$ for \mathfrak{n} the nilradical of a parabolic subalgebra $\mathfrak{p} = \mathfrak{l} + \mathfrak{n}$ of \mathfrak{g} . The orbit dimension $\dim \mathcal{O}_L(e_0) = \dim[\mathfrak{l}, e_0]$ is given by taking the rank of the matrix formed from the coefficients of the images under $\text{ad } e_0$ of a basis of \mathfrak{l} contained in $\mathfrak{g}_{\mathbb{Z}}$. Hence, after reduction modulo p , the orbit dimension can only go down. Since by 2.8 the dimension of the orbit $\text{Ind}_{\Gamma}^{\mathfrak{g}} \mathcal{O}_L(e_0)$ is given by the formula $\dim \mathfrak{g} - \dim \mathfrak{g}_e = \dim \mathfrak{g} - \dim \mathfrak{l}_{e_0}$ we know that if $\dim \mathfrak{g}_e$ is independent of good characteristic then e is indeed in the nilpotent orbit which intersects densely with $\mathcal{O}_L(e_0) + \mathfrak{n}$ in good characteristic. But our calculation of the Jordan blocks of $\text{ad } e_{\Gamma}$ (described in §4.1) shows that this is the case for all Γ 's listed in [8] and, moreover, the orbit $\mathcal{O}(e_{\Gamma})$ always has the same label as its counterpart $\mathcal{O}(e_{\Gamma, \mathbb{C}}) \subset \mathfrak{g}_{\mathbb{C}}$. The exceptional case where $p = 7$ and \mathfrak{g} is of type F_4 does not cause us serious problems because \mathfrak{g} contains only two sheets of dimension 44 both of which have rank 2 (this is immediate from [8, Table 10] and Remark 2.9). Proposition 3.1(ii) implies that one of them contains the orbit of type B_3 . On the other hand, a closer look at [8, Table 10] reveals that the sheets of dimension $\neq 44$ cannot contain nilpotent orbits of dimension 42. This yields that the other sheet of dimension 44 contains the orbit of type C_3 (as in the characteristic 0 case).

Next we observe that $e_0 = e_{\Gamma_0}$ and $e_1 = e_{\Gamma_1}$ for some full subgraphs Γ_0 and Γ_1 of Γ . Looking through the tables in [8] one finds out that if G is a group of type E then in all cases except two Γ_0 is a disjoint union of Dynkin graphs (possibly empty). Specifically, the two exceptional cases occur when G has type E_8 and e has type $E_7(a_5)$ or $D_5(a_1) + A_1$. In both cases \mathfrak{l} contains a *unique* nilpotent orbit whose dimension equals that of $\mathcal{O}_L(e_0)$. A close look at the tables in [8] shows the same holds for groups of type G_2 and F_4 in all cases of interest. Thanks to Corollary 3.6 this implies that the L -orbit of $e_0 = e_{\Gamma_0}$ has the same label as the $L_{\mathbb{C}}$ -orbit of $e_{\Gamma_0, \mathbb{C}}$.

⁴Our calculation threw up a misprint in [8]. Specifically, the first nilpotent representative for the induced nilpotent orbit $E_7(a_4)$ in E_8 has 27 over two different nodes of the Dynkin diagram. de Graaf kindly redid the calculation and found that the right-most 27 should be a 44, and the 44 (just below the wrong 27) should be a 14.

Given a nilpotent orbit $\mathcal{O} \subset \mathcal{N}(\mathfrak{g})$ we define

$$d(\mathcal{O}) := \text{Card} \{ \mathcal{S} \mid \mathcal{S} \text{ is a sheet of } \mathfrak{g} \text{ containing } \mathcal{O} \}$$

and denote by $d(\mathcal{O}_{\mathbb{C}})$ the number of sheets of $\mathfrak{g}_{\mathbb{C}}$ containing the nilpotent orbit having the same Dynkin diagram as \mathcal{O} . The preceding discussion in conjunction with [8] implies that $d(\mathcal{O}) \geq d(\mathcal{O}_{\mathbb{C}})$ for any orbit $\mathcal{O} \subset \mathcal{N}(\mathfrak{g})$. Since every sheet of \mathfrak{g} contains a unique nilpotent orbit (Remark 2.6) the total number of sheets of \mathfrak{g} equals $\sum_{\mathcal{O} \subset \mathcal{N}(\mathfrak{g})} d(\mathcal{O})$. Since Theorem 2.8 established a bijection between the sheets of \mathfrak{g} and $\mathfrak{g}_{\mathbb{C}}$, we now deduce that $d(\mathcal{O}) = d(\mathcal{O}_{\mathbb{C}})$ for any orbit $\mathcal{O} \subset \mathcal{N}(\mathfrak{g})$. This completes the proof of Theorem 1.4. \square

4.3. (Strong) reachability

We will reduce the problem of classifying the (strongly) reachable elements to a finite calculation which can be performed in GAP. For this we will (i) exhibit a bound on the number of characteristics where the result may differ from the situation in characteristic zero established by de Graaf, then (ii) give a method to deal with any specific prime.

For (ii) we reduce to a calculation over the prime field \mathbb{F}_p . Observe that the Lie algebras we deal with are defined over \mathbb{Z} . That is, there is a Lie algebra $\mathfrak{g}_{\mathbb{Z}}$ over the integers with a basis \mathcal{B} , and associated structure constants, such that $\mathfrak{g}_p := \mathfrak{g}_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathbb{F}_p$ and $\mathfrak{g} = \mathfrak{g}_p \otimes_{\mathbb{F}_p} k$ with \mathfrak{g} and \mathfrak{g}_p obtaining a basis and associated structure constants by taking \mathcal{B} together with its structure constants reduced modulo p .

For this, \mathcal{B} can be taken as a Chevalley basis $\{x_{\alpha} \mid \alpha \in \Phi\} \cup \{h_{\alpha} \mid \alpha \in \Pi\}$ of $\mathfrak{g}_{\mathbb{Z}}$. Then we have give a complete set of nilpotent representatives over k in terms of linear combinations of elements \mathcal{B} . Since the coefficients are all over \mathbb{Z} , these nilpotent elements are also elements of \mathfrak{g}_p by reduction mod p . Now since \mathfrak{g}_e is defined over \mathbb{F}_p , we may write $\mathfrak{g}_e = (\mathfrak{g}_p)_e \otimes_{\mathbb{F}_p} k$. Then in order to establish reachability of a nilpotent element e , it suffices to check whether we have e contained in the derived subalgebra of $(\mathfrak{g}_p)_e$.

This calculation can then be performed by GAP.

For (i), we assume that p is a good prime for \mathfrak{g} , any bad primes being dealt with in case (ii). We work over the integers, i.e., with the admissible \mathbb{Z} -form $\mathfrak{g}_{\mathbb{Z}}$ (here we use our assumption that the group G is simply connected). Fix a nilpotent element e . First, we ask GAP to calculate a basis \mathcal{B}_e of $(\mathfrak{g}_{\mathbb{C}})_e$ contained in $\mathfrak{g}_{\mathbb{Z}}$ in terms of linear combinations of elements of \mathcal{B} . By the theory of nilpotent elements it is possible to take this basis such that reduction of the coefficients modulo p also gives a basis of $(\mathfrak{g}_p)_e$.

Next we take the product of each pair of elements of \mathcal{B}_e and form a matrix M of the \mathcal{B} -coefficients of the resulting vectors. Then the dimension of $[(\mathfrak{g}_p)_e, (\mathfrak{g}_p)_e]$ is the rank of the reduction modulo p of the matrix M . Thus, if we take p bigger than any of the elementary divisors of M the result will agree with that over \mathbb{Z} . Similar remarks apply to the subalgebra $\mathbb{F}_p e + [(\mathfrak{g}_p)_e, (\mathfrak{g}_p)_e]$ of $(\mathfrak{g}_p)_e$ whose dimension can be determined by calculating the rank of the matrix M' which is formed by adjoining the row of coefficients of e itself to M . Thus, for p bigger than any of the elementary divisors of M and M' , we have that e is reachable over k if and only if it is reachable over a field of characteristic 0. More precisely, we establish that the only exceptional primes are less than or equal to 7.

4.4. Almost perfect centralisers and $c(\mathfrak{g}_e) = \dim(\mathfrak{g}_e/[\mathfrak{g}_e, \mathfrak{g}_e])$

For a given prime, identifying the orbits which have almost perfect centralisers (i.e., $\mathfrak{g}_e = [\mathfrak{g}_e, \mathfrak{g}_e] + ke$) is a straightforward calculation in $\mathfrak{g}_p = \mathfrak{g}_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathbb{F}_p$ with GAP: we simply ask GAP to list those nilpotent elements e for which

$$\dim(\mathfrak{g}_p)_e = \dim([\mathfrak{g}_p]_e, (\mathfrak{g}_p)_e) + 1$$

and

$$\dim(\mathbb{F}_p e + [\mathfrak{g}_e, \mathfrak{g}_e]) = \dim(\mathfrak{g}_p)_e.$$

This process deals in particular with the bad primes. The other possible exceptional primes were already calculated: if $\dim([\mathfrak{g}_p]_e, (\mathfrak{g}_p)_e)$ or $\dim(\mathbb{F}_p e + [(\mathfrak{g}_p)_e, (\mathfrak{g}_p)_e])$ differs from the analogous dimension over characteristic zero, then we established in § 4.3 that $p \leq 7$.

Much the same applies to calculating $c(\mathfrak{g}_e)$. Again one need only look in characteristics at most 7, where the calculation becomes finite, hence easily performed in GAP.

4.5. Panyushev property

Let $\mathfrak{g}_{\mathbb{Q}} = \mathfrak{g}_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathbb{Q}$, a \mathbb{Q} -form of $\mathfrak{g}_{\mathbb{C}}$. A nilpotent elements $e \in \mathcal{O}(D)$ is said to satisfy *Panyushev's property* if the nilradical $\mathfrak{g}_e(\lambda_D, \geq 1)$ of \mathfrak{g}_e is generated by $\mathfrak{g}_e(\lambda_D, 1)$ as a Lie algebra. Since this definition relies on the so-called associated cocharacters which do not exist for all nilpotent elements in bad characteristic, Panyushev's property is particularly interesting under our assumptions on G .

We assume that G is an exceptional group and p is a good prime for G . From [20] we take a nilpotent element $e \in \mathfrak{g}$ and associated cocharacter $\tau \in X_*(G)$. For $e \in \mathcal{O}(D)$ this cocharacter coincides with λ_D and it is optimal for e in the sense of the Kempf–Rousseau theory by one of the main results of [23]. Let Φ be the root system of G associated with a maximal torus containing $\tau(k^{\times})$. We start by working over rationals and ask GAP to compute the root vectors in $\mathfrak{g}_{\mathbb{Q}}(1) := \mathfrak{g}_{\mathbb{Q}}(\tau, 1)$. Let \mathcal{B} be a basis of $\mathfrak{g}_{\mathbb{Q}}$ containing a set of root vectors $\{e_{\alpha} \mid \alpha \in \Phi\}$. Now we form a $(\dim \mathfrak{g}_{\mathbb{Q}}(\tau, 1) \times \dim \mathfrak{g}_{\mathbb{Q}})$ -matrix M of the \mathcal{B} -coefficients of $[b, e]$ with $b \in \mathcal{B} \cap \mathfrak{g}_{\mathbb{Q}}(1)$. Then $(\mathfrak{g}_{\mathbb{Q}})_e(1)$ coincides with the kernel of $\text{ad } e: \mathfrak{g}_{\mathbb{Q}}(1) \rightarrow \mathfrak{g}_{\mathbb{Q}}$ which for GAP is $\{v \in \mathbb{Q}^{\dim \mathfrak{g}_{\mathbb{Q}}(\tau, 1)} \mid v \cdot M = 0\}$. Since p is good for G , we have that $(\mathfrak{g}_p)_e = (\mathfrak{g}_{\mathbb{Z}})_e \otimes_{\mathbb{Z}} \mathbb{F}_p$ and $(\mathfrak{g}_p)_e(1) = (\mathfrak{g}_{\mathbb{Z}})_e(1) \otimes_{\mathbb{Z}} \mathbb{F}_p$. In particular, $(\mathfrak{g}_p)_e(1)$ is characteristic independent.

Next we generate a Lie subring \mathfrak{m} of $\mathfrak{g}_{\mathbb{Z}}$ from a \mathbb{Q} -basis of $(\mathfrak{g}_{\mathbb{Q}})_e(1)$ contained in $\mathfrak{g}_{\mathbb{Z}}$ and take a basis $\mathcal{B}_{\mathfrak{m}}$ of the \mathbb{Q} -span of \mathfrak{m} that lies in $\mathfrak{g}_{\mathbb{Z}}$. We then form the matrix of \mathcal{B} -coefficients of the elements of $\mathcal{B}_{\mathfrak{m}}$ and take elementary divisors again. If p is bigger than any of these elementary divisors then the rank of this matrix will give the dimension of the Lie subalgebra of \mathfrak{g}_p generated by $(\mathfrak{g}_p)_e(1)$. Since it turned out that all elementary divisors appearing in these calculations involve bad primes only, we conclude that the Panyushev property is independent of good characteristic.

Remark 4.2. There are six rigid nilpotent orbits in exceptional Lie algebras which do not satisfy Panyushev's property in good characteristic. These orbits are \tilde{A}_1 in type G_2 , $\tilde{A}_2 + A_1$ in type F_4 , $(A_3 + A_1)'$ in type E_7 , and $A_3 + A_1$, $A_5 + A_1$, $D_5(a_1) + A_2$ in type E_8 . The above routine was applied to these orbits too in order to determine the smallest

number r for which $\sum_{i=1}^r \mathfrak{g}_e(\tau, i)$ generates the Lie algebra $\mathfrak{g}_e(\tau, \geq 1)$. It turned out that $r = 2$ in type G_2 , $r = 4$ in type E_8 when e has type $A_5 + A_1$ and $r = 3$ in the other four cases. All elementary divisors that we encountered in the process turned out to be divisible by 2 and 3 only.

Acknowledgements. We would like to thank Michaël Bulois, Jim Humphreys, Simon Goodwin, Willem de Graaf, George Lusztig and Dan Nakano for very useful discussions and email correspondence on the subject of this paper.

References

1. W. BORHO AND H. KRAFT, Über Bahnen und deren Deformationen bei linearen Aktionen reductiver Gruppen, *Comment. Math. Helv.* **54**(1) (1979), 61–104.
2. A. BOREL, *Linear Algebraic Groups*, second edition, Graduate Texts in Mathematics, Volume 126 (Springer, New York, 1991).
3. W. BORHO, Über Schichten halbeinfacher Lie-Algebren, *Invent. Math.* **65**(2) (1981/82), 283–317.
4. P. BARDSLEY AND R. W. RICHARDSON, Étale slices for algebraic transformation groups in characteristic p , *Proc. Lond. Math. Soc.* (3) **51**(2) (1985), 295–317.
5. R. W. CARTER, Conjugacy classes in the Weyl group, *Compos. Math.* **25** (1972), 1–59.
6. R. W. CARTER, *Finite Groups of Lie Type: Conjugacy Classes and Complex Characters* (Wiley Classics Library. John Wiley & Sons Ltd., Chichester, 1993). Reprint of the 1985 Wiley–Interscience Publication.
7. W. A. DE GRAAF, Computations with nilpotent orbits in SLA. Preprint, 2013.
8. W. A. DE GRAAF AND A. ELASHVILI, Induced nilpotent orbits of the simple Lie algebras of exceptional type, *Georgian Math. J.* **16**(2) (2009), 257–278.
9. D. EISENBUD, *Commutative Algebra with a View Toward Algebraic Geometry*, Graduate Texts in Mathematics, Volume 150 (Springer, New York, 1995).
10. A. ELASHVILI, Sheets of the exceptional Lie algebras, *Stud. Algebra* **171** (1984).
11. R. M. FOSSUM, One-dimensional formal group actions, in *Invariant Theory (Denton, TX, 1986)*, Contemporary Mathematics, Volume 88, pp. 227–397 (American Mathematical Society, Providence, RI, 1989).
12. W. H. HESSELINK, Nilpotency in classical groups over a field of characteristic 2, *Math. Z.* **166**(2) (1979), 165–181.
13. J. C. JANTZEN, Nilpotent orbits in representation theory, in *Lie Theory*, Progressive Mathematics, Volume 228, pp. 1–211 (Birkhäuser Boston, Boston, MA, 2004).
14. G. KEMPKEN, Induced conjugacy classes in classical Lie algebras, *Abh. Math. Sem. Univ. Hamburg* **53** (1983), 53–83.
15. H. KRAFT, Parametrisierung von Konjugationsklassen in \mathfrak{sl}_n , *Math. Ann.* **234**(3) (1978), 209–220.
16. R. LAWTHER, Jordan block sizes of unipotent elements in exceptional algebraic groups, *Comm. Algebra* **23**(11) (1995), 4125–4156.
17. G. LUSZTIG AND N. SPALTENSTEIN, Induced unipotent classes, *J. Lond. Math. Soc.* (2) **19**(1) (1979), 41–52.
18. G. LUSZTIG, Character sheaves on disconnected groups, II, *Represent. Theory* **8** (2004), 72–124.
19. M. W. LIEBECK AND G. M. SEITZ, *Unipotent and Nilpotent Classes in Simple Algebraic Groups and Lie Algebras*, Mathematical Surveys and Monographs, Volume 180 (American Mathematical Society, Providence, RI, 2012).

20. R. LAWTHER AND D. M. TESTERMAN, Centres of centralizers of unipotent elements in simple algebraic groups, *Mem. Amer. Math. Soc.* **210**(988) (2011), vi+188. 989
21. G. J. MCNINCH, Adjoint Jordan blocks. Preprint, 2002. 990
22. G. J. MCNINCH, Nilpotent orbits over ground fields of good characteristic, *Math. Ann.* **329**(1) (2004), 49–85. 991
23. A. PREMET, Nilpotent orbits in good characteristic and the Kempf–Rousseau theory, *J. Algebra* **260**(1) (2003), 338–366. Special issue celebrating the 80th birthday of Robert Steinberg. 992
24. A. PREMET AND S. SKRYABIN, Representations of restricted Lie algebras and families of associative \mathcal{L} -algebras, *J. Reine Angew. Math.* **507** (1999), 189–218. 993
25. A. PREMET, Commutative quotients of finite W -algebras, *Adv. Math.* **225** (2010), 269–306. 994
26. A. PREMET AND L. TOPLEY, Derived subalgebras of centralisers and finite W -algebras, *Compos. Math.* **150**(9) (2014), 1485–1548. 995
27. M. ROSENLICHT, On quotient varieties and the affine embedding of certain homogeneous spaces, *Trans. Amer. Math. Soc.* **101** (1961), 211–223. 996
28. N. SPALTENSTEIN, *Classes Unipotentes et Sous-groupes de Borel*, Lecture Notes in Mathematics, Volume 946 (Springer, Berlin–New York, 1982). 997
29. N. SPALTENSTEIN, Nilpotent classes and sheets of Lie algebras in bad characteristic, *Math. Z.* **181**(1) (1982), 31–48. 998
30. N. SPALTENSTEIN, Nilpotent classes in Lie algebras of type F_4 over fields of characteristic 2, *J. Fac. Sci. Univ. Tokyo Sect. IA Math.* **30**(3) (1984), 517–524. 999
31. T. A. SPRINGER AND R. STEINBERG, Conjugacy classes, in *Seminar on Algebraic Groups and Related Finite Groups (The Institute for Advanced Study, Princeton, NJ, 1968/69)*, Lecture Notes in Mathematics, Volume 131, pp. 167–266 (Springer, Berlin, 1970). 1000
32. UNIVERSITY OF GEORGIA VIGRE ALGEBRA GROUP, Varieties of nilpotent elements for simple Lie algebras I: good primes, *J. Algebra* **280** (2004), 719–737. The University of Georgia VIGRE Algebra Group: D. J. Benson, P. Bergonio, B. D. Boe, L. Chastkofsky, D. Cooper, J. Jungster, G. M. Guy, J. J. Hyun, G. Matthews, N. Mazza, D. K. Nakano, K. J. Platt. 1001
33. UNIVERSITY OF GEORGIA VIGRE ALGEBRA GROUP, Varieties of nilpotent elements for simple Lie algebras: II. Bad primes, *J. Algebra* **292** (2005), 65–99. The University of Georgia VIGRE Algebra Group: D. J. Benson, P. Bergonio, B. D. Boe, L. Chastkofsky, B. Cooper, G. M. Guy, J. Hower, M. Hunziker, J. J. Hyun, J. Kujawa, G. Matthews, N. Mazza, D. K. Nakano, K. J. Platt and C. Wright. 1002
34. O. YAKIMOVA, On the derived algebra of a centraliser, *Bull. Sci. Math.* **134**(6) (2010), 579–587. 1003