

Ringel duality for certain strongly quasi-hereditary algebras

Martin Kalck¹ · Joseph Karmazyn²

Received: 26 November 2017 / Revised: 5 April 2018 / Accepted: 8 April 2018 /
Published online: 2 May 2018
© The Author(s) 2018

Abstract We study quasi-hereditary endomorphism algebras defined over a new class of finite dimensional monomial algebras with a special ideal structure. The main result is a uniform formula describing the Ringel duals of these quasi-hereditary algebras. As special cases, we obtain a Ringel duality formula for a family of strongly quasi-hereditary algebras arising from a type A configuration of projective lines in a rational, projective surface as recently introduced by Hille and Ploog, for certain Auslander–Dlab–Ringel algebras, and for Eiriksson and Sauter’s nilpotent quiver algebras when the quiver has no sinks and no sources. We also recover Tan’s result that the Auslander algebras of self-injective Nakayama algebras are Ringel self-dual.

Keywords Quasi-hereditary algebras · Ringel duality · Monomial algebras · Knörrer invariant algebras

Mathematics Subject Classification 16S50 · 16G10

The first author was supported by EPSRC Grant EP/L017962/1. The second author was supported by EPSRC Grant EP/M017516/2.

✉ Joseph Karmazyn
j.h.karmazyn@sheffield.ac.uk
Martin Kalck
martin.maths@posteo.de

¹ Freiburg, Germany

² School of Mathematics and Statistics, University of Sheffield, Hicks Building, Hounsfield Road, Sheffield S3 7RH, UK

1 Introduction

Quasi-hereditary algebras form an important class of finite dimensional algebras with relations to Lie theory (this was the original motivation [10]) and exceptional sequences in algebraic geometry (see e.g. [9,23]). Examples of quasi-hereditary algebras include blocks of category \mathcal{O} and Schur algebras.

Ringel duality [34] is a fundamental phenomenon in the theory of quasi-hereditary algebras, see for example [8,13–15,19,22,26,28,33] for (recent) work on this topic. For any quasi-hereditary algebra A there exists a quasi-hereditary algebra $\mathfrak{R}(A)$, the *Ringel dual* of A , such that

$$A\text{-mod} \cong \mathfrak{R}(\mathfrak{R}(A))\text{-mod}.$$

However, computing the Ringel dual of a quasi-hereditary algebra explicitly may not be straightforward. In this paper we introduce a new class of quasi-hereditary algebras that admit a uniform description of their Ringel duals, see Theorem 1.2.

Let us make this more precise. Let k be an algebraically closed field, and R be a finite dimensional monomial k -algebra, i.e. $R = kQ/I$, where I is a two-sided ideal generated by paths in Q . For example $R = k\langle x_1, \dots, x_l \rangle / I$, where I is a two-sided ideal generated by monomials in $k\langle x_1, \dots, x_l \rangle$.

Definition 1.1 We call R *ideally ordered*, if for every primitive idempotent $e \in R$ and every pair of monomials $m, n \in eR$ there exists an epimorphism $Rm \rightarrow Rn$ or an epimorphism $Rn \rightarrow Rm$.

For an algebra R we consider the additive subcategory of all torsionless R -modules

$$\text{sub}(R) := \text{add}\{U \mid U \subseteq R^{\oplus n}\} \subseteq R\text{-mod},$$

define $\text{SUB}(R) := \bigoplus_{U \in \text{ind}(\text{sub}(R))} U$ to be the direct sum of all indecomposable modules in $\text{sub}(R)$ up to isomorphism, and set

$$E_R := \text{End}_R(\text{SUB}(R)).$$

For submodules $\Lambda \subset R$ we define the layer function $l(\Lambda) := \dim_k R - \dim_k \Lambda$ and we call l the *ideal layer function*. For an ideally ordered algebra R the isomorphism classes of submodules $\Lambda \subset R$ label the simple modules $S(\Lambda)$ of E_R and so the ideal layer function induces a partial ordering on the simple E_R -modules: $S(\Lambda_1) \leq S(\Lambda_2) \Leftrightarrow l(\Lambda_1) \leq l(\Lambda_2)$. We call this the *ideal layer ordering*.

The following is the main result of this paper and calculates the Ringel dual for algebras of the form E_R . See Theorem 5.1 for a more detailed version.

Theorem 1.2 *Let R be a finite dimensional ideally ordered monomial algebra. Then E_R is quasi-hereditary with respect to the ideal layer ordering, has global dimension ≤ 2 , and has Ringel dual $E_{R^{\text{op}}}^{\text{op}}$:*

$$\mathfrak{R}(E_R) \cong E_{R^{\text{op}}}^{\text{op}}.$$

Remark 1.3 As we were preparing to post this paper on the arXiv we became aware of the very recent paper [14] of Coulembier that had just appeared. This paper introduces a more general version of the Auslander–Dlab–Ringel construction and proves a Ringel duality formula in this setting. In particular, this generalises the Ringel duality formula of Conde and Erdmann [13] that we discuss below.

Our construction appears to be a special case that fits into this more general framework which, in particular, implies the Ringel duality formula of Theorem 1.2. However, the approach and proof in Coulembier’s work is different to the one in this paper. The work of Coulembier also seems to answer the questions we raise in Remark 5.3 (1) and at the end of Sect. 6.3 regarding the possibility of finding a more general framework in which a Ringel duality formula holds.

In light of this, the results of this paper can be thought of as providing a very explicit example of Coulembier’s Ringel duality formula, linking to several geometrically inspired examples such as Knörrer invariant algebras, and proving further properties that hold in our special case of the algebras E_R such as being simultaneous left and right strongly quasi-hereditary for the same quasi-hereditary order and being left ultra strongly quasi-hereditary.

The class of ideally ordered monomial algebras includes many well known examples, and in many of these examples the endomorphism algebras E_R are also well understood.

Example 1.4 The following families of finite dimensional monomial algebras are ideally ordered.

(0) Hereditary algebras.

(1) The algebras $R = k\langle x_1, \dots, x_l \rangle / (x_1, \dots, x_l)^m$ for positive integers l, m .

(2) More generally, for Q a finite quiver, $J \subseteq kQ$ the two-sided ideal generated by all arrows in Q , and $m \geq 0$ the algebra $R := kQ/J^m$ is ideally ordered.

To prove this, consider a monomial $p \in eR$. There is a surjection $Re \rightarrow Rp$ given by $g \mapsto gp$ with kernel

$$\{g \in Re \mid gp = 0\} \cong \{g \in kQ \mid gp \in J^m\} \cong J^{m-i}e$$

where i is minimal such that $p \in J^i$. Hence for any monomial $p \in eR$ there is an isomorphism $Rp \cong Re/J^l e$ for some $l \in \{1, \dots, m\}$. As a result, for any pair of monomials $p, q \in eR$ the monomial ideals Rp, Rq are isomorphic to some pair of quotient modules occurring in the chain of surjections

$$Re \cong Re/J^m e \rightarrow Re/J^{m-1} e \rightarrow \dots \rightarrow Re/J^1 e.$$

Hence there is a surjection $Rp \rightarrow Rq$ or $Rq \rightarrow Rp$.

(3) For every pair $0 < a < r$ of coprime integers the finite dimensional monomial Knörrer invariant algebra $K_{r,a}$ is defined in [27, Definition 4.6], and the results of [27, Section 6.4] describe its monomial ideals and imply that it is ideally ordered. The definition of these algebras is recapped in Sect. 6.1.

(4) Nakayama algebras, introduced in [31], are ideally ordered.

We give two constructions that can be used to produce ideally ordered monomial algebras.

- (5) Let R and K be ideally ordered monomial algebras and let ${}_R M_K$ be an R - K -bimodule which is projective as R -module and as K -module. Then

$$T := \begin{pmatrix} R & {}_R M_K \\ 0 & K \end{pmatrix}$$

is an ideally ordered monomial algebra. Example 2.8(a) shows that T need not be ideally ordered if we weaken the assumptions on ${}_R M_K$.

- (6) If R is ideally ordered and $e \in R$ is an arbitrary idempotent, then eRe is ideally ordered.

Suppose that $f \in eRe$ is a primitive idempotent and $p, q \in feRe = fRe$ are monomials. Then f is a primitive idempotent in R , $p, q \in fR$ are monomials, and as R is ideally ordered there is a surjection between Rp and Rq . Applying

$$eR \otimes_R (-) \cong \text{Hom}_R(Re, -): R\text{-mod} \rightarrow eRe\text{-mod}$$

to this surjection of R -modules will produce the required surjection of eRe -modules between eRp and eRq since $eR \otimes_R (-)$ is exact. This shows eRe is ideally ordered.

We finish by exhibiting a local commutative monomial algebra which is not ideally ordered.


- (7) The algebra $R = k[x, y]/(x^3, xy, y^3)$ is not ideally ordered. To see this consider the ideals Rx and Ry .

We briefly discuss how these examples of ideally ordered monomial algebras R , and the algebras $E_R := \text{End}_R(\text{SUB}(R))$ they define, relate to algebras and results in the literature.

Hille and Ploog’s algebras

The Ringel duality formula of Theorem 1.2, the definition of ideally ordered monomial algebras, and the construction of the algebras E_R in this paper are all geometrically inspired. They were first observed in our previous work [27] for a class of quasi-hereditary algebras Λ_α constructed by Hille and Ploog [24].

In more detail, the algebras Λ_α arise from an exceptional collection of line bundles associated to a type A configuration of intersecting rational curves C_i in a rational, projective surface as illustrated in the picture below.



The construction of Λ_α (recapped in Sect. 6.1) depends on the order of the curves C_i . Reversing the order of these curves, Hille and Ploog's construction yields an algebra Λ_{α^\vee} .

It is natural to ask how the algebras Λ_α and Λ_{α^\vee} are related from a representation theoretic perspective. Our answer below is phrased in terms of Ringel duality.

Proposition 1.5 *There is an isomorphism of algebras*

$$\mathfrak{R}(\Lambda_\alpha) \cong \Lambda_{\alpha^\vee}^{\text{op}}. \quad (1)$$

In order to see that (1) is a special case of our main Theorem 1.2, we recall that there are isomorphisms of algebras

$$\Lambda_\alpha \cong E_{K_{r,a}} \quad \text{and} \quad \Lambda_{\alpha^\vee} \cong E_{K_{r,a}^{\text{op}}}$$

described in [27, Section 6]. This is recalled in Proposition 6.7 and the discussion immediately beneath it. Here, $K_{r,a}$ denotes a Knörrer invariant algebra, which is the ideally ordered monomial in Example 1.4(3), and $0 < a < r$ are a pair of coprime integers depending on α .

We remark that in this setting the Ringel duality formula (1) also has an alternative proof, which is more geometric, see Proposition 1.5.

The aim of this paper was to find a more general representation theoretic framework extending the Ringel duality formula (1) to a larger class of (ultra) strongly quasi-hereditary algebras. In particular, the Knörrer invariant algebras are the original motivation for the ideally ordered condition.

Remark 1.6 The algebras $\Lambda_\alpha \cong E_{K_{r,a}}$ and $K_{r,a}$ were used to show a noncommutative version of Knörrer periodicity for cyclic quotient surface singularities in [27]. More precisely, it was proved there that the singularity category of a cyclic quotient surface singularity is equivalent to the singularity category of a corresponding Knörrer invariant algebra, generalising classical Knörrer's periodicity for the polynomials x^n and $x^n + y^2 + z^2$. The proof uses noncommutative resolutions and $\Lambda_\alpha \cong E_{K_{r,a}}$ plays the role of a noncommutative resolution for $K_{r,a}$.

Auslander–Dlab–Ringel and nilpotent quiver algebras

From a more representation theoretic viewpoint, a Ringel duality formula that looks similar to that of Theorem 1.2 was proved for Auslander–Dlab–Ringel algebras E_R^{ADR} by Conde and Erdmann [13, Theorem A]. We define these algebras, recall Conde and Erdmann's Ringel duality formula, and discuss the relationship between this result and the results of this paper in Sect. 6.3.

In particular, for the class of algebras $R := kQ/J^m$ in Example 1.4(2) the corresponding algebras E_R and E_R^{ADR} coincide if Q has no sources.

Proposition 6.12 *If $R := kQ/J^m$ for Q a finite quiver without sources and J the two-sided ideal generated by all arrows in Q , then there is an isomorphism of quasi-hereditary algebras $E_R^{\text{ADR}} \cong E_R$.*

We also prove that when Q has no sinks the ADR algebra coincides with the quiver nilpotent algebra $N_m(Q)$ introduced by Eiriksson and Sauter [20], which is motivated via a quiver graded version of Richardson orbits and is recapped in Sect. 6.4.

Proposition 6.15 *If $R := kQ/J^m$ for Q a finite quiver without sinks and J the two-sided ideal generated by all arrows in Q , then there is an isomorphism of quasi-hereditary algebras $N_m(Q) \cong E_{kQ/J^m}^{\text{ADR}}$.*

In particular, if $R = kQ/J^m$ (as in Example 1.4(2)) for a quiver with no sinks or sources, then $E_R \cong E_R^{\text{ADR}} \cong N_m(Q)$ and so Theorem 1.2 provides a Ringel duality formula for such nilpotent quiver algebras; see Corollary 6.17.

Nakayama and Auslander algebras

Several of the examples of ideally ordered monomial algebras above can be thought of as geometrically inspired by resolutions of singularities. Indeed, Examples 1.4(1)–(4) can be thought of as different generalisations of the algebra $k[x]/x^n$.

Work of Dlab and Ringel [17] shows that every finite dimensional algebra admits a noncommutative ‘resolution’ by a quasi-hereditary algebra, and a generalisation of this result led to Iyama’s proof of the finiteness of Auslander’s representation dimension [25].

Such a resolution for finite dimensional algebras of finite representation type is provided by the Auslander algebra. This also occurs in more geometric contexts; the categorical resolutions considered by Kuznetsov and Lunts [30] use a construction motivated by Auslander algebras to resolve non-reduced schemes.

For R a finite dimensional algebra of finite representation type let E_R^{AUS} denote the Auslander algebra of R , which we recall in Sect. 6.5.

Proposition 6.18 *If R is an ideally ordered monomial algebra, then $E_R^{\text{AUS}} \cong E_R$ if and only if R is self-injective.*

A particular example of a class of ideally ordered, monomial algebras of finite representation type are the Nakayama algebras (listed as Example 1.4(4)).

Corollary 6.19 *If R is self-injective Nakayama algebra, then $E_R^{\text{AUS}} \cong E_R$.*

In this setting Theorem 1.2 also generalises several known results in the literature, e.g. that the Auslander algebras of self-injective Nakayama algebras are Ringel self-dual, see [37].

Corollary 6.20 *If R is a self-injective Nakayama algebra then $\mathfrak{R}(E_R) \cong E_R$.*

For other related results see work by Baur et al. [6], Crawley-Boevey and Sauter [16] and Nguyen et al. [32].

Left and right strongly quasi-hereditary structure

A further special property of the quasi-hereditary algebras $E_R = \text{End}_R(\text{SUB}(R))$ is that the ideal layer function simultaneously realises both a left and right strongly quasi-hereditary structure on the algebras.

Since $\text{add SUB}(R)$ is closed under kernels $E_R = \text{End}_R(\text{SUB}(R))$ has global dimension 2, and it was recently shown by Tsukamoto [38] that this implies E_R admits both a left strongly quasi-hereditary structure and a right strongly quasi-hereditary structure (for a possibly different order), building on earlier work of Dlab and Ringel, and Iyama.

In general the left and right strongly quasi-hereditary structures cannot be realised using the same order. Indeed, Tsukamoto shows that for Auslander algebras of representation-finite algebras (which all have global dimension 2) this is possible precisely if the underlying algebra is a Nakayama algebra.

As seen in the examples above, the class of quasi-hereditary algebras E_R constructed from ideally ordered monomial algebras provides a larger class of such algebras.

Conventions

Throughout this paper k will denote an algebraically closed field. For paths $p, q \in kQ$ in the path algebra of a quiver Q the composition pq will denote the path q followed by the path p . For R a Noetherian ring $R\text{-mod}$ will denote the category of finitely generated left R -modules, and for $S \subset R\text{-mod}$ we will define $\text{add } S$ to be the additive subcategory generated by S : i.e. the smallest full subcategory of $R\text{-mod}$ containing S and closed under isomorphism, direct sums, and direct summands. In particular, the category of finitely generated projective R modules $\text{proj-}R$ is equivalent to $\text{add } R$.

We recall the category of torsionless R -modules $\text{sub}(R)$ from the introduction, and now give a more general definition: for an R -module M we define the following subcategory

$$\text{sub}(M) := \text{add}\{U \mid U \subseteq M^{\oplus n}\} \subseteq R\text{-mod}$$

with corresponding module $\text{SUB}(M) := \bigoplus_{U \in \text{ind}(\text{sub}(M))} U$. Moreover, for an R -module M , we set

$$\text{fac}(M) := \text{add}\{Q \mid M^{\oplus n} \rightarrow Q \rightarrow 0\} \subseteq R\text{-mod}$$

and let $\text{FAC}(M) := \bigoplus_{Q \in \text{ind}(\text{fac}(M))} Q$ denote the direct sum of all indecomposable objects in $\text{fac}(M)$ up to isomorphism.

We let \dagger denote the standard k -duality $\text{Hom}_k(-, k)$. For the injective cogenerator $I := R_R^\dagger$ we define the category of *divisible* R -modules

$$\text{fac}(I) := \text{add}\{Q \mid I^{\oplus n} \rightarrow Q \rightarrow 0\} \subseteq R\text{-mod}$$

and let $\text{FAC}(I) := \bigoplus_{Q \in \text{ind}(\text{fac}(I))} Q$ denote the direct sum of all indecomposable objects in $\text{fac}(I)$ up to isomorphism.

2 Strongly quasi-hereditary algebras

In this section, we will give necessary and sufficient conditions for certain endomorphism algebras over ideally ordered monomial algebras to be left or right strongly quasi-hereditary.

We first recall the definition of a quasi-hereditary algebra. This needs some preparation. For a finite dimensional k -algebra A choose a labelling $i \in I$ of the simple A -modules S_i up to isomorphism. A partial order \leq on the set I is called *adapted* if for each $M \in A\text{-mod}$ with top S_i and socle S_j incomparable there exists some $k > i$ or $k > j$ such that S_k is a composition factor of M . In particular, total orderings are adapted. We denote the projective cover and injective envelope of the simple S_i by P_i and Q_i respectively.

Definition 2.1 Given a partial ordering \leq on the index set I , for $i \in I$ the *standard* module Δ_i is the maximal factor module of P_i whose composition series consists only of simple modules S_j such that $j \leq i$. Similarly, the *costandard* module ∇_i is the maximal submodule of Q_i whose composition series consists only of simple modules S_j such that $j \leq i$.

The k -algebra A is *quasi-hereditary* with respect to an adapted partial ordering \leq if:

- (1) $\text{End}_A(\Delta_i) \cong k$ for each $i \in I$ and
- (2) A can be filtered by the standard modules under this ordering; i.e. there exists a series of A -modules $0 = M_n \subset M_{n-1} \subset \dots \subset M_1 \subset M_0 = A$ such that each quotient M_{i-1}/M_i is isomorphic to a direct sum of standard modules.

The following terminology is due to Ringel [35]. We refer to the references and discussions in [35] for earlier work.

Definition 2.2 A quasi-hereditary algebra A is called *left strongly quasi-hereditary* if all standard modules have projective dimension at most 1. It is called *right strongly quasi-hereditary* if all costandard A -modules have injective dimension at most 1.

This is an equivalent characterisation of left/right strongly quasi-hereditary condition given in [35, Appendix A1]. The original definition, introduced in [35, Section 4], is in terms of a layer function.

Definition 2.3 A k -algebra A is *left strongly quasi-hereditary* with n layers if there is a layer function $L: \{\text{simple } A\text{-modules}\}/\cong \rightarrow \{1, \dots, n\}$ such that for any simple module s with projective cover $P(s)$ there is an exact sequence

$$0 \rightarrow R(s) \rightarrow P(s) \rightarrow \Delta(s) \rightarrow 0$$

such that

- (a) The module $R(s)$ is the direct sum of projective covers $P(s')$ of simple modules s' such that $L(s') > L(s)$.
- (b) All simple factors s' of $\text{rad } \Delta(s)$ satisfy $L(s') < L(s)$.

The layer function induces an ordering on the simple A -modules and the modules $\Delta(s)$ are the standard modules for this strongly quasi-hereditary structure. Right strongly quasi-hereditary algebras are defined dually.

After some preparation, we introduce the class of endomorphism algebras which we are interested in. For the rest of this section we let R be a finite dimensional k -algebra. A submodule of the form $Rp \subset R$ is a *principal left ideal* if $p \in eR$ with $e \in R$ a primitive idempotent. We introduce the additive subcategory

$$\text{pi}(R) := \text{add}\{Rp \mid p \in eR, e \text{ primitive idempotent}\} \subset R\text{-mod},$$

and we let $\text{PI}(R) := \bigoplus_{Rp \in \text{ind}(\text{pi}(R))} Rp$ denote the direct sum of all principal left ideals up to isomorphism. In this section we assume that $\text{PI}(R)$ is finitely generated and define $E_R^{\text{PI}} := \text{End}_R(\text{PI}(R))$.

The assumption on $\text{PI}(R)$ is satisfied for ideally ordered monomial algebras R due to Lemma 7.3 but does not hold for all finite dimensional algebras; e.g. if $R = \mathbb{C}[x, y]/(x^2, y^2)$, then the ideals $I_\lambda := R(x + \lambda y)$ for $\lambda \in \mathbb{C}$ give a \mathbb{C} -indexed set of ideals that are pairwise non-isomorphic as left modules.

Throughout the rest of the paper we will label the simple and projective E_R^{PI} -modules by the principal ideals of R , as we now explain. To do this we use the additive anti-equivalence

$$\text{add PI}(R) \xrightarrow{\text{Hom}_R(-, \text{PI}(R))} E_R^{\text{PI}}\text{-proj}. \tag{2}$$

It is clear that $\text{Hom}_R(-, \text{PI}(R))$ is a contravariant functor, and one can show that it is an additive anti-equivalence using that it maps the additive generator $\text{PI}(R)$ of $\text{add PI}(R)$ to the additive generator E_R^{PI} of $E_R^{\text{PI}}\text{-proj}$. Under this anti-equivalence the indecomposable summands Λ of $\text{PI}(R)$ are in 1-to-1 correspondence with indecomposable projective E_R^{PI} -modules, which we denote by $P(\Lambda)$. The indecomposable projective modules $P(\Lambda)$ are in 1-to-1 correspondence with simple E_R^{PI} -modules $S(\Lambda)$ that occur as their heads (i.e, so that $P(\Lambda) \rightarrow S(\Lambda)$ is a projective cover). Hence the principal ideals $\Lambda \subset R$ index the simple modules $S(\Lambda)$ of E_R^{PI} . When given a partial ordering on the principal ideals, we use similar notation to label standard $\Delta(\Lambda)$ and costandard $\nabla(\Lambda)$ objects. This labelling allows to define the following layer function for the algebra E_R^{PI} .

Definition 2.4 Let R be a finite dimensional algebra. For principal left R -ideals Λ , we define $l(S(\Lambda)) := l(\Lambda) := \dim_k R - \dim_k \Lambda$ and we call l the *ideal layer function*. It induces a partial ordering on the principal left R -ideals, which we call the *ideal ordering*.

We will now determine when the ideal layer function induces a left or right strongly quasi-hereditary structure on E_R^{PI} by considering left and right minimal approximations with respect to the ideal ordering.

The notion of minimal approximation is common in representation theory; see [29] for a survey. A morphism $\alpha: \Gamma \rightarrow \Lambda$ is a left approximation for a class of modules \mathcal{C} if $\Lambda \in \mathcal{C}$ and the induced morphism $\text{Hom}_R(\Lambda, C) \rightarrow \text{Hom}_R(\Gamma, C)$ is surjective for all $C \in \mathcal{C}$. A morphism $\Gamma \xrightarrow{\alpha} \Lambda$ is left minimal if any endomorphism ϕ of Λ satisfying $\phi \circ \alpha = \alpha$ is an isomorphism. In particular, left minimal approximations are unique up to isomorphism.

Denote by $\text{pi}(R)_{>i} \subseteq \text{pi}(R)$ the full subcategory of direct sums of principal left R -ideals Λ with $l(\Lambda) > i$.

Lemma 2.5 *Let Γ be a principal left ideal of layer γ . There is a minimal left $\text{pi}(R)_{>\gamma}$ approximation $\alpha_\Gamma: \Gamma \rightarrow \Gamma_{>\gamma}$ of Γ .*

Proof It is well-known that Γ admits a left $\text{pi}(R)_{>\gamma}$ approximation $\Phi: \Gamma \rightarrow \Lambda$. Indeed, this follows since there are only finitely many indecomposable objects in $\text{pi}(R)_{>\gamma} \subseteq R\text{-mod}$ and since R is finite dimensional, see e.g. [5]. For the convenience of the reader, we recall the argument. We consider the module

$$\Lambda := \bigoplus_{M \in \text{pi}(R)_{>\gamma}} M^{\oplus \dim \text{Hom}_R(\Gamma, M)}$$

where the sum is taken over all indecomposable objects M in $\text{pi}(R)_{>\gamma}$ (up to isomorphism). Then $\Lambda \in \text{pi}(R)_{>\gamma}$ as each $\text{Hom}_R(\Gamma, M)$ is finite dimensional, $\text{PI}(R)$ is assumed to be finitely generated, and $\text{pi}(R)_{>\gamma}$ is closed under finite direct sums.

Choosing a basis $(\phi_i)_{i \in I}$ of

$$\bigoplus_{M \in \text{pi}(R)_{>\gamma}} \text{Hom}_R(\Gamma, M)$$

determines a morphism $\Phi: \Gamma \rightarrow \Lambda$ as the direct sum $\Phi = \bigoplus_{i \in I} \phi_i$. One can check that Φ is a left $\text{pi}(R)_{>\gamma}$ approximation.

The existence of a left approximation with a finite length target implies the existence of a minimal left approximation by, for example [4, Theorem I.2.4], which shows such a minimal approximation can be constructed from an approximation by projection onto a summand. Hence the existence of the approximation $\Phi: \Gamma \rightarrow \Lambda$ ensures that a minimal left $\text{pi}(R)_{>\gamma}$ approximation $\alpha_\Gamma: \Gamma \rightarrow \Gamma_{>\gamma}$ exists. □

Definition 2.6 We say that $\text{PI}(R)$ has *good left approximations* if

$$\text{Hom}_R(\text{coker } \alpha_\Gamma, \text{PI}(R)) = 0$$

for all principal left R -ideals Γ .

Lemma 2.7 *If R is an ideally ordered monomial algebra, then for a principal ideal Γ of layer γ the minimal left $\text{pi}(R)_{>\gamma}$ approximation is surjective. Hence when R is ideally ordered $\text{PI}(R)$ has good left approximations.*

Proof Since R is ideally ordered, we can use Lemma 7.3 to replace any principal R -ideal by an isomorphic monomial ideal wherever needed. In particular, without loss of generality let $\Gamma = Rg$ (with $g \in eR$ a monomial) be a principal left R -ideal of layer γ .

A surjection from Γ to a principal ideal exists, $\Gamma \rightarrow 0$ as 0 is a principal ideal. Using that R is finite dimensional there is a surjection to a principal ideal $\Gamma_{>\gamma}$ which has maximal dimension among all principal ideals that admit surjections from Γ . The existence of the surjection implies that Γ and $\Gamma_{>\gamma}$ have the same head. In particular, we can assume that $\Gamma_{>\gamma} = Rn$ for a monomial $n \in eR$. Using Lemma 7.1, the assignment $g \mapsto n$ defines an R -linear surjection $\alpha_\Gamma : \Gamma \rightarrow \Gamma_{>\gamma}$.

We now claim that α_Γ is an approximation. To prove this we consider a principal ideal Λ and will show that the induced map $\text{Hom}_R(\Gamma_{>\gamma}, \Lambda) \rightarrow \text{Hom}_R(\Gamma, \Lambda)$ is a surjection. Take a morphism $\beta \in \text{Hom}_R(\Gamma, \Lambda)$. We aim to show that β factors through α_Γ and hence is the image of some morphism in $\text{Hom}_R(\Gamma_{>\gamma}, \Lambda)$.

To see this, take the induced surjection $\beta : \Gamma \rightarrow \text{im } \beta$ and, as the image of a principal ideal in a principal ideal, $\text{im } \beta \cong Rm$ (with a monomial $m \in eR$) is a principal left R -ideal. Using the ideally ordered condition on R there is a surjection in at least one direction between $\text{im } \beta$ and $\Gamma_{>\gamma}$. As $\Gamma_{>\gamma}$ is a principal ideal of maximal dimension with a surjection from Γ , it follows that $\dim \Gamma_{>\gamma} \geq \dim \text{im } \beta$ and hence there is a surjection $\sigma : \Gamma_{>\gamma} \rightarrow \text{im } \beta$. Using Lemma 7.1, we can assume that σ is given by $n \mapsto m$. Hence, the composition $\pi := \sigma \circ \alpha_\Gamma$ is a surjection defined by $g \mapsto m$. Now Lemma 7.2 shows that the surjection $\beta : \Gamma \rightarrow \text{im } \beta$ factors over π . In particular, β factors over α_Γ . So α_Γ is an approximation.

Finally, we claim that this approximation is minimal. To see this consider an endomorphism $\phi : \Gamma_{>\gamma} \rightarrow \Gamma_{>\gamma}$ such that $\phi \circ \alpha_\Gamma \cong \alpha_\Gamma$. Then as α_Γ is a surjection it follows that ϕ is a surjection, and hence an isomorphism.

By construction, $\text{coker } \alpha_\Gamma = 0$ for all Γ so $\text{PI}(R)$ has good left approximations. \square


We give examples showing that our results above apply beyond the class of ideally ordered monomial algebras.

Example 2.8 (a) Consider the monomial algebra $R = kQ/I$, where

$$Q := \begin{array}{c} & & a & & \\ & & \curvearrowright & & \\ & y & & & \\ & \curvearrowleft & 1 & \curvearrowright & b \\ & & & & \\ & x & & & \\ & \curvearrowright & & & \\ & & c & & \end{array} \quad \text{and} \quad I = (a, b, c)^2 + (yb, xc).$$

This is not ideally ordered since there are no surjections between Rb and Rc , however $\text{PI}(R)$ still has good left approximations. It is a short exercise to find the five isomorphism classes of indecomposable principal ideals and calculate their minimal left approximations. All but one of these minimal approximations are surjective, and the one which is not surjective has cokernel S_1 , the simple at vertex 1. There are no morphisms from S_1 to any principal ideal, and hence $\text{PI}(R)$ has good left approximations.

(b) Let $n > 0$ be an integer. Consider the non-monomial algebras $R_n = kQ/I_n$ where



Again, $\text{PI}(R_n)$ has good left approximations; it is a short exercise to find the $n + 3$ principal ideals and calculate that the minimal left approximation for each one is surjective.

Proposition 2.9 *The algebra $E_R^{\text{PI}} = \text{End}_R(\text{PI}(R))$ is left strongly quasi-hereditary with respect to the ideal layer function l if and only if $\text{PI}(R)$ has good left approximations with respect to l .*

Proof Assume $\text{PI}(R)$ has good left approximations $\alpha_\Gamma : \Gamma \rightarrow \Gamma_{>\gamma}$. Using the condition on $\text{coker } \alpha_\Gamma$ and applying $\text{Hom}_R(-, \text{PI}(R))$ yields a short exact sequence

$$0 \rightarrow P(\Gamma_{>\gamma}) \xrightarrow{\iota(\Gamma)} P(\Gamma) \rightarrow \Delta(\Gamma) \rightarrow 0, \tag{3}$$

where $\iota(\Gamma) = \text{Hom}_R(\alpha_\Gamma, \text{PI}(R))$ and $\Delta(\Gamma)$ denotes the cokernel of $\iota(\Gamma)$. We claim that the ideal layer function defines a left strongly quasi-hereditary structure on E_R^{PI} such that the $\Delta(\Gamma)$ are standard modules. To see this we have to show that (3) satisfies conditions (a) and (b) outlined in Definition 2.3. Since all direct summands of $P(\Gamma_{>\gamma})$ are of the form $P(\Lambda)$ with $l(\Lambda) > \gamma$ condition (a) is satisfied by construction. Using the anti-equivalence $\text{Hom}_R(-, \text{PI}(R)) : \text{add PI}(R) \rightarrow \text{proj-}E_R^{\text{PI}}$ condition (b) translates to: every R -linear non-isomorphism $v : \Gamma \rightarrow \Lambda$ with $\Lambda \in \text{pi}(R)_{\geq \gamma}$ factors over α_Γ . By definition of α_Γ this holds for $\Lambda \in \text{pi}(R)_{>\gamma}$. If $\Lambda \in \text{pi}(R)_{=\gamma}$, then v cannot be surjective for otherwise it is an isomorphism since $\dim_k \Lambda = \dim_k \Gamma$. Therefore, $\text{im } v \subsetneq \Lambda$ is a principal left R -ideal with $l(\text{im } v) > l(\Lambda) = \gamma$. So v factors over α_Γ .

To see the converse direction, assume $\text{PI}(R)$ does not have good left approximations. Then there exists a principal left R -ideal Γ such that $\text{Hom}_R(\text{coker } \alpha_\Gamma, \text{PI}(R)) \neq 0$. Assume that E_R^{PI} is quasi-hereditary with respect to the ideal layer function l and let $\Delta(\Gamma)$ be the standard module corresponding to Γ . Since α_Γ is a minimal left $\text{pi}(R)_{>\gamma}$ approximation

$$P(\Gamma_{>\gamma}) \xrightarrow{\text{Hom}_R(\alpha_\Gamma, \text{PI}(R))} P(\Gamma) \rightarrow \Delta(\Gamma) \rightarrow 0,$$

is the start of a minimal projective resolution of $\Delta(\Gamma)$. By our choice of Γ the morphism $\text{Hom}_R(\alpha_\Gamma, \text{PI}(R))$ is not injective. Hence $\Delta(\Gamma)$ has projective dimension greater than 1 and, using Definition 2.2, A is not left strongly quasi-hereditary with respect to l in this case. □

Remark 2.10 Assume that E_R^{PI} is quasi-hereditary with respect to the ideal layer function. One can show that as a set the standard module $\Delta(\Gamma)$ is given by all (residue

classes of) monomorphisms starting in Γ . Indeed if $\nu: \Gamma \rightarrow \Lambda$ is not a monomorphism then an argument along the lines of the proof of the proposition shows that ν factors over α_Γ and therefore corresponds to the zero element in $\Delta(\Gamma)$.

Proposition 2.9 is related to [35, Theorem 5] by Ringel. He shows that for an R -module M there exists an R -module N such that $\text{End}_R(M \oplus N)$ is left strongly quasi-hereditary and all the indecomposable summands N are submodules of M . In particular, if M is an R -module such that all submodules are isomorphic to direct summands of M , then $\text{End}_R(M)$ is left strongly quasi-hereditary. We will see in Theorem 5.1 that $\text{PI}(R)$ has this property if R is ideally ordered monomial. However, our proof of Theorem 5.1 uses Proposition 2.9, so we cannot apply Ringel’s result in our approach.

Now we look at the ‘dual’ side. First we ‘dualise’ Definition 2.6 using the same notation.

Definition 2.11 For every principal left ideal Γ there is a minimal right $\text{pi}(R)_{>\gamma}$ approximation $\rho_\Gamma: \Gamma_{>\gamma} \rightarrow \Gamma$ with $\Gamma_{>\gamma} \in \text{pi}(R)_{>\gamma}$. We say that $\text{PI}(R)$ has *good right approximations* if

$$\text{Hom}_R(\text{PI}(R), \ker \rho_\Gamma) = 0.$$

Since $\text{PI}(R)$ contains R as a direct summand this is equivalent to $\ker \rho_\Gamma = 0$ for all principal left R -ideals Γ .

Example 2.12 (a) Let R be a finite dimensional monomial algebra. Then $\text{PI}(R)$ has good right approximations. Indeed, let Γ be a principal left R ideal. Since R is monomial, $\text{rad } \Gamma$ is a direct sum of principal left ideals in $\text{pi}(R)_{>\gamma}$ and the natural inclusion $\text{rad } \Gamma \rightarrow \Gamma$ gives the desired minimal right approximation ρ_Γ .

(b) The algebra in Example 2.8(b) does not have good right approximations: the minimal right approximation of the projective module P_1 is $P_2 \oplus P_3 \rightarrow P_1$ and this has kernel S_4 .

The following result is proved dually to Proposition 2.9

Proposition 2.13 $E_R^{\text{PI}} = \text{End}_R(\text{PI}(R))$ is right strongly quasi-hereditary with respect to the ideal layer function l if and only if $\text{PI}(R)$ has good right approximations. For example, this holds if R is finite dimensional monomial.

Combining Propositions 2.9 and 2.13 with Lemma 2.7 and Example 2.12(a) yields the following theorem.

Theorem 2.14 If R is an ideally ordered monomial algebra, then E_R^{PI} is both left and right strongly quasi-hereditary with respect to the ordering induced by the ideal layer function.

We let $\mathcal{Filt}(\Delta)$ and $\mathcal{Filt}(\nabla)$ denote the full subcategories of E_R^{PI} -mod of objects filtered by standard and costandard modules respectively.

Remark 2.15 Assume that E_R^{PI} is quasi-hereditary with respect to the ideal layer function. Similarly to the case above, one can show that as a set a costandard module $\nabla(\Lambda)$ is given by all surjections ending in Λ . In particular, each costandard module has head $S(\Pi)$ for some indecomposable projective R -module Π and $\mathcal{Filt}(\nabla) \subseteq \text{fac}(P(R))$.

Corollary 2.16 *If $\text{PI}(R)$ has good right and left approximations, then $\mathcal{Filt}(\Delta)$ is closed under submodules and $\mathcal{Filt}(\nabla)$ is closed under quotients.*

Proof If $\text{PI}(R)$ has good left approximations, then E_R^{PI} is left strongly quasi-hereditary by Proposition 2.9, and hence all standard objects have projective dimension 1. By [35, Proposition A.1], all standard modules having projective dimension 1 is equivalent to $\mathcal{Filt}(\nabla)$ being closed under quotients.

The analogous dual statement, using Proposition 2.13, shows that when $\text{PI}(R)$ has good right approximations then $\mathcal{Filt}(\Delta)$ is closed under submodules. □

3 The characteristic tilting module and Ringel duality

In the following section we first recall the characteristic tilting module T associated to a quasi-hereditary algebra. Then we show that our algebras E_R^{PI} are ultra strongly quasi-hereditary in the sense of Conde [12] and use this to determine a subcategory of the additive hull $\text{add}(T)$ of T (Corollary 3.6). In the proof of our main Theorem 5.1 we show that these categories coincide for ideally ordered monomial algebras R and as a consequence establish our Ringel duality formula in this setup.

The following proposition can be found in Ringel [34], which is based on work of Auslander and Reiten [3] and Auslander and Buchweitz [2].

Proposition 3.1 *Let A be a quasi-hereditary algebra. Then there exists a tilting module $T \in A\text{-mod}$ such that*

$$\text{add}(T) = \mathcal{Filt}(\Delta) \cap \mathcal{Filt}(\nabla),$$

where $\mathcal{Filt}(\Delta) \cap \mathcal{Filt}(\nabla)$ is the full subcategory of A -modules with filtrations by both standard and costandard modules.

Definition 3.2 A tilting module T occurring in Proposition 3.1 is called a *characteristic tilting module*. The *Ringel dual* $\mathfrak{R}(A)$ of an algebra A is defined by

$$\mathfrak{R}(A) := \text{End}_A(T)^{\text{op}}$$

for T the basic characteristic tilting module consisting of one copy of each indecomposable module in $\mathcal{Filt}(\Delta) \cap \mathcal{Filt}(\nabla)$ up to isomorphism: i.e. we assume $\mathfrak{R}(A)$ is a basic algebra.

The notion of an ultra strongly quasi-hereditary algebras was introduced by Conde, see [12, Section 2.2.2].

Definition 3.3 A quasi-hereditary algebra A is *left ultra strongly quasi-hereditary* if a projective module P_i is filtered by costandard modules whenever the corresponding costandard module ∇_i is simple.

Let $e_0 \in E_R^{\text{PI}} = \text{End}_R(\text{PI}(R))$ be the idempotent corresponding to the direct summand R of $\text{PI}(R)$. Note that e_0 is primitive if and only if R is local. We have the following.

Proposition 3.4 *Let R be a finite dimensional algebra. Assume that $\text{Pl}(R)$ has good left approximations, so that E_R^{Pl} is left strongly quasi-hereditary with respect to the ideal layer function l . Then the following conditions are equivalent:*

- (a) $E_R^{\text{Pl}}e_0$ is filtered by costandard objects.
- (b) $\alpha_\Gamma: \Gamma \rightarrow \Gamma_{>\gamma}$ is surjective for all principal R -ideals Γ .
- (c) E_R^{Pl} is left ultra strongly quasi-hereditary.

If R is monomial then these conditions are equivalent to

- (d) R is ideally ordered.

Proof We first show that (b) implies (a). By [34, Theorem 4], it suffices to show that $\text{Ext}_{E_R^{\text{Pl}}}^1(\Delta(\Gamma), P(Re_i)) = 0$ for all principal left R ideals Γ and all primitive idempotents $e_i \in R$. We can assume that $\Delta(\Gamma)$ is not projective. Then applying $\text{Hom}_R(-, \text{Pl}(R))$ to α_Γ produces the projective resolution

$$0 \rightarrow P(\Gamma_{>\gamma}) \xrightarrow{\iota(\Gamma)} P(\Gamma) \rightarrow \Delta(\Gamma) \rightarrow 0,$$

and we have to show that every morphism $P(\Gamma_{>\gamma}) \rightarrow P(Re_i)$ factors over $\iota(\Gamma)$. Applying the anti-equivalence given in equation (2) translates this statement to: every morphism $\varphi: Re_i \rightarrow \Gamma_{>\gamma}$ factors over $\alpha_\Gamma: \Gamma \rightarrow \Gamma_{>\gamma}$. This holds since Re_i is projective and α_Γ is surjective by assumption.

Conversely, if α_Γ is not surjective for some principal ideal Γ then there exists $x \in \Gamma_{>\gamma} \setminus \text{im } \alpha_\Gamma$. Since R is free there is an R -linear map $R \rightarrow \Gamma_{>\gamma}$, $1 \mapsto x$, which by construction does not factor over α_Γ . In combination with the anti-equivalence and projective resolution above this shows $\text{Ext}_{E_R^{\text{Pl}}}^1(\Delta(\Gamma), P(R)) \neq 0$ and [34, Theorem 4] completes the proof that (a) implies (b).

That (a) is equivalent to (c) follows from the fact that $\nabla(\Lambda)$ is simple if and only if Λ is projective, see Remark 2.15, and hence $\nabla(\Lambda)$ simple implies $P(\Lambda)$ is a direct summand of $E_R^{\text{Pl}}e_0$.

Let R be monomial. The implication (d) \Rightarrow (b) follows from Lemma 2.7. We now assume (b) and prove the converse.

Firstly, for any indecomposable principal ideal Γ the minimal left approximation $\alpha_\Gamma: \Gamma \rightarrow \Gamma_{>\gamma}$ is surjective by assumption (b), and we claim that $\Gamma_{>\gamma}$ is indecomposable.

To show this take $p \in eR$ for e a primitive idempotent and consider the principal ideal $\Gamma \cong Rp$. Now suppose that there is a decomposition $\Gamma_{>\gamma} \cong \bigoplus Rq_i$ for some principal ideals Rq_i . As α_Γ is surjective, after relabelling we can assume that the image of p is (q_1, \dots, q_n) and $q_1 \neq 0$. As the morphism α_Γ is surjective there must exist some $r \in Re$ such that $\alpha_\Gamma(rp) = (q_1, 0, \dots, 0)$; i.e. $rq_1 = q_1$ and $rq_j = 0$ for $j \geq 2$. As R is monomial, by considering the monomial of lowest degree occurring in q_1 and $rq_1 = q_1$ we can see that the degree 0 primitive idempotent e must occur in r . Then we can rewrite $r = e + r'$ where all monomials occurring in r' have degree greater than 0. As a result, q_j must be zero as $0 = rq_j = q_j + r'q_j$ so there can be no non-zero monomial of lowest degree occurring in q_j . Hence $q_j = 0$ for $j \geq 2$,

the decomposition is a trivial decomposition $Rq_1 \cong Rq_1 \oplus 0 \oplus \dots \oplus 0$, and $\Gamma_{>y}$ is indecomposable.

This allows the successive construction of left $\text{pi}_{>k}$ approximations starting with the indecomposable principal ideal Re

$$Re \xrightarrow{\alpha_{Re}} Re_{>i_1} \xrightarrow{\alpha_{Re_{>i_1}}} Re_{>i_2} \xrightarrow{\alpha_{Re_{>i_2}}} \dots \xrightarrow{\alpha_{Re_{>i_{n-1}}}} Re_{>i_n}$$

where $i_1 = l(Re)$, $i_{j+1} = l(Re_{>i_j})$, and $\alpha_{Re_{>i_j}} : Re_{>i_j} \rightarrow Re_{>i_{j+1}}$ is the minimal left $\text{pi}_{>i_{j+1}}$ approximation. Each $Re_{>i_j}$ is indecomposable, and the composition $\alpha_k : Re \rightarrow Re_{>i_k}$ of the left approximations is again a left approximation.

We claim that any indecomposable principal ideal Rx with $x \in eR$ is isomorphic to one of these successive approximations. To see this choose k to be maximal such that $l(Rx) > i_k$. Then there is a surjection $\pi : Re \rightarrow Rx$, and as $Rx \in \text{pi}(R)_{>i_k}$ this must factor through the left approximation $\alpha_k : Re \rightarrow Re_{>i_k}$ by a surjection $\phi : Re_{>i_k} \rightarrow Rx$. In particular, $\dim Re_{>i_k} \geq \dim Rx$ so $l(Re_{>i_k}) \leq l(Rx)$. But, by the definition of k , it is true that $i_{k+1} = l(Re_{>i_k}) \geq l(Rx)$, hence it must be the case that $l(Re_{>i_k}) = l(Rx)$ so $\dim Re_{>i_k} = \dim Rx$ and hence the surjective morphism ϕ is an isomorphism $Re_{>i_k} \cong Rx$.

Finally, any pair Rx and Ry of principal ideals with $x, y \in eR$ occur (up to isomorphism) in the successive approximation sequence, in which every morphism is surjective by assumption (b), and hence there is a surjection between them. This proves that the ideally ordered condition holds. □

Example 3.5 The non-monomial algebra in Example 2.8(b) satisfies the equivalent conditions (a), (b) and (c) of the theorem.

Corollary 3.6 *Suppose that $\text{Pl}(R)$ has both good left and right approximations. Then $\text{sub}(E_R^{\text{Pl}}e_0) \cap \text{fac}(E_R^{\text{Pl}}e_0) \subseteq \text{Filt}(\Delta) \cap \text{Filt}(\nabla) = \text{add}(T)$.*

Proof By the definition of a quasi-hereditary algebra every projective module is filtered by standard modules. Therefore, $(E_R^{\text{Pl}}e_0)^{\oplus n} \in \text{Filt}(\Delta)$ and by Proposition 3.4(a), we also have $(E_R^{\text{Pl}}e_0)^{\oplus n} \in \text{Filt}(\nabla)$. Now Corollary 2.16 yields $\text{sub}(E_R^{\text{Pl}}e_0) \subseteq \text{Filt}(\Delta)$ and $\text{fac}(E_R^{\text{Pl}}e_0) \subseteq \text{Filt}(\nabla)$. This implies the claim. □

Remark 3.7 In combination with Remark 2.15, we see that when $\text{Pl}(R)$ has both good left and right approximations $\text{fac}(E_R^{\text{Pl}}e_0) = \text{Filt}(\nabla)$. For ideally ordered monomial algebras R , Theorem 5.1 (e) shows that $\text{sub}(E_R^{\text{Pl}}e_0) = \text{Filt}(\Delta)$ holds as well.

Remark 3.8 Let $R = R_2$ be the non-monomial algebra from Example 2.8(b). The algebra $E_R^{\text{Pl}} = \text{End}_R(\text{Pl}(R))$ is left ultra strongly quasi-hereditary with respect to the ideal layer function (in particular, $E_R^{\text{Pl}}e_0$ is filtered by costandard modules) but not right strongly quasi-hereditary, so $\text{Filt}(\Delta)$ is not closed under subobjects. It turns out that there is precisely one indecomposable subobject of $E_R^{\text{Pl}}e_0$ which is not filtered by standard modules. This module is also a quotient of $E_R^{\text{Pl}}e_0$ and therefore $\text{sub}(E_R^{\text{Pl}}e_0) \cap \text{fac}(E_R^{\text{Pl}}e_0) \not\subseteq \text{Filt}(\Delta) \cap \text{Filt}(\nabla) = \text{add}(T)$. Restricting to the local submodules of $E_R^{\text{Pl}}e_0$ yields the desired inclusion into $\text{add}(T)$ in this case and can be used to show a

version of the Ringel duality formula (10) in this example. Unfortunately, we do not know how to fit this example into a larger framework.

4 An equivalence from idempotents

In this section, we show that there is an equivalence of categories

$$\text{Hom}_A(Ae_0, -) : \text{sub}(Ae_0) \cap \text{fac}(Ae_0) \rightarrow \text{sub}(e_0Ae_0).$$

where $A = E_R^{\text{Pl}}$ for a finite dimensional algebra R with $\text{Pl}(R)$ finitely generated and $e_0 \in A$ is the idempotent corresponding to the projection onto R .

To show this we recall several well-known lemmas.

Lemma 4.1 *Let \mathcal{A} be an abelian category with Serre subcategory \mathcal{S} and let $q : \mathcal{A} \rightarrow \mathcal{A}/\mathcal{S}$ be the quotient functor. Then the restriction of q ,*

$${}^\perp\mathcal{S} \cap \mathcal{S}^\perp \rightarrow \mathcal{A}/\mathcal{S},$$

is fully faithful. Here

$$\begin{aligned} {}^\perp\mathcal{S} &:= \{X \in \mathcal{A} \mid \text{Hom}_{\mathcal{A}}(X, S) = 0 \text{ for all } S \text{ in } \mathcal{S}\}, \text{ and} \\ \mathcal{S}^\perp &:= \{Y \in \mathcal{A} \mid \text{Hom}_{\mathcal{A}}(S, Y) = 0 \text{ for all } S \text{ in } \mathcal{S}\}. \end{aligned}$$

Proof This follows from the description of homomorphism spaces in the quotient category as colimits. Indeed for $X, Y \in {}^\perp\mathcal{S} \cap \mathcal{S}^\perp$ the colimit describing $\text{Hom}_{\mathcal{A}/\mathcal{S}}(X, Y)$ is taken over the single pair of subobjects $(X, 0)$ and the quotient functor sends a morphism $f : X \rightarrow Y$ to f . □

The following lemma can be found in [21, Proposition 5.3 (b)]

Lemma 4.2 *Let B be a noetherian ring and let $e \in B$ be an idempotent. Then*

$$F = \text{Hom}_B(Be, -) : B\text{-mod} \rightarrow eBe\text{-mod}$$

is an exact quotient functor with kernel $B/BeB\text{-mod}$. In particular, $B/BeB\text{-mod}$ is a Serre-subcategory in $B\text{-mod}$.

Corollary 4.3 *In the notation of Lemma 4.2, we have $\text{fac}(Be) \subseteq^\perp (B/BeB\text{-mod})$.*

Proof Consider $N \in \text{fac}(Be)$ and $M \in B/BeB\text{-mod}$. Applying the right exact functor $\text{Hom}_B(-, M)$ to the surjection $Be \rightarrow N \rightarrow 0$ yields the injection $0 \rightarrow \text{Hom}_B(N, M) \rightarrow \text{Hom}_B(Be, M)$. As $B/BeB\text{-mod}$ is the kernel of $\text{Hom}_B(Be, -)$ and $M \in B/BeB\text{-mod}$ it follows that $\text{Hom}_B(Be, M) = 0$ and hence $\text{Hom}_B(N, M) = 0$. □

From now on let $A = E_R^{\text{Pl}}$ for some finite dimensional algebra R , such that $\text{Pl}(R)$ is finitely generated.

Lemma 4.4 *In the notation of Sect. 3, we have $\text{soc } Ae_0 \subseteq S_0^{\oplus n}$ for some natural number n . Here $S_0 = Ae_0/\text{rad } Ae_0$ is the semi-simple head of Ae_0 .*

Proof Indeed Ae_0 consists of all R -homomorphisms $R \rightarrow \text{PI}(R)$. Let Λ be a principal left R -ideal. If $R \rightarrow \Lambda$ is non-zero, then the composition with the canonical inclusion $R \rightarrow \Lambda \rightarrow R$ is non-zero. Therefore every maximal sequence of non-zero morphisms starting in R ends in R , proving the claim. \square

Corollary 4.5 $\text{sub}(Ae_0) \subseteq (A/Ae_0A\text{-mod})^\perp$.

Proof Assume that $f: X \rightarrow U$ is a non-zero map, where U in $\text{sub}(Ae_0)$ and X in $A/Ae_0A\text{-mod}$. Lemma 4.4 implies that $\text{im } f$ contains a non-zero direct summand of S_0 . But $\text{im } f \in A/Ae_0A\text{-mod}$ since X is contained in $A/Ae_0A\text{-mod}$. It follows that $\text{im } f$ has no submodule which is a direct summand of S_0 . A contradiction. So there is no non-zero morphism $f: X \rightarrow U$. \square

The following statement is the main result of this section.

Proposition 4.6 *The exact functor $F = \text{Hom}_A(Ae_0, -)$ restricts to an additive equivalence*

$$\text{sub}(Ae_0) \cap \text{fac}(Ae_0) \rightarrow \text{sub}(e_0Ae_0) \cap \text{fac}(e_0Ae_0) = \text{sub}(e_0Ae_0).$$

Proof The equality on the right follows from the fact that $\text{fac}(e_0Ae_0) = e_0Ae_0\text{-mod}$. Since F is exact and maps an A -module M to e_0M , the restriction is well-defined. We can apply Lemma 4.1 to $q = F$ to deduce that F is fully faithful. Indeed, by Lemma 4.2, F is a quotient functor corresponding to the Serre subcategory $A/Ae_0A\text{-mod}$ and Corollaries 4.3 and 4.5 show that the required orthogonality conditions are satisfied.

It remains to show that F is essentially surjective. Let $U \subseteq (e_0Ae_0)^{\oplus n}$ be generated by $u_1, \dots, u_n \in (e_0Ae_0)^n$. The u_i are elements of $(Ae_0)^n$. Let $V \subseteq (Ae_0)^{\oplus n}$ be the A -submodule generated by the u_i . One can check that $F(V) = U$ and since $e_0u_i = u_i$ for all i V is a factor module of $(Ae_0)^{\oplus m}$ for some m . This shows that V is contained in $\text{sub}(Ae_0) \cap \text{fac}(Ae_0)$ and completes the proof. \square

5 Proof of Ringel duality formula

In this section we prove the following main result of this paper, which is an extended version of Theorem 1.2 stated in the introduction.

Theorem 5.1 *Let R be a finite dimensional ideally ordered monomial algebra and $E_R = \text{End}_R(\text{SUB}(R))$. Then E_R is quasi-hereditary and the Ringel duality formula*

$$\mathfrak{R}(E_R) \cong (E_{R^{\text{op}}})^{\text{op}}$$

holds. More explicitly, where \dagger denotes the standard k -duality,

$$\mathfrak{R}(\text{End}_R(\text{SUB}(R))) \cong \text{End}_R(\text{FAC}(R_R^\dagger)) \cong \text{End}_{R^{\text{op}}}(\text{SUB}(R^{\text{op}}))^{\text{op}}, \tag{4}$$

and if we consider $\text{sub}(R) := \text{addSUB}(R)$ and $\text{fac}(R^\dagger) := \text{addFAC}(R_R^\dagger)$ as exact categories with split exact structures then this Ringel duality induces the derived equivalence

$$D^b(\text{sub}(R)) \cong D^b(\text{fac}(R^\dagger)).$$

Moreover:

- (a) Every indecomposable submodule of R^n is isomorphic to a principal left ideal, every principal left ideal is isomorphic to a monomial ideal, and hence $\text{sub}(R) \cong \text{pi}(R)$ so $E_R^{\text{Pl}} \cong E_R$.
- (b) The algebra E_R is left and right strongly quasi-hereditary with respect to the ideal layer function. In particular, E_R has global dimension at most 2. Moreover, it is left ultra strongly quasi-hereditary in the sense of Conde [12].
- (c) The ideal order is the unique order defining a quasi-hereditary structure on E_R if R is local and satisfies the following condition: if there exists a surjection $\Lambda \rightarrow \Gamma$ between principal left ideals, then there is an inclusion $\Gamma \rightarrow \Lambda$.
- (d) Let T be the characteristic tilting module of E_R and $e_0 \in E_R$ be the idempotent corresponding to R . Then there is an equality of subcategories $\text{add}(T) = \text{sub}(E_R e_0) \cap \text{fac}(E_R e_0)$. In other words, the indecomposable direct summands T_i of T are precisely those indecomposable E_R -modules which are both quotients and submodules of the projective module $E_R e_0$.
- (e) We can describe the subcategories $\text{Filt}(\Delta)$ and $\text{Filt}(\nabla)$ of $E_R\text{-mod}$ as follows:

$$\text{Filt}(\Delta) = \text{sub}(T) = \text{sub}(E_R e_0) = \text{sub}(E_R), \text{ and} \tag{5}$$

$$\text{Filt}(\nabla) = \text{fac}(T) = \text{fac}(E_R e_0). \tag{6}$$

Proof We first prove the main Ringel duality formula, and in the process also prove (a) and (d). Let $E_R^{\text{Pl}} = \text{End}_R(\text{Pl}(R))$ and let $e_0 \in E_R^{\text{Pl}}$ be the idempotent corresponding to R . By Corollary 3.6, we have an inclusion

$$\text{sub}(E_R^{\text{Pl}} e_0) \cap \text{fac}(E_R^{\text{Pl}} e_0) \subseteq \text{add}(T) \tag{7}$$

where T is the characteristic tilting module for E_R^{Pl} . In combination with Proposition 4.6, we get an inclusion

$$\text{sub}(R^{\text{op}}) \rightarrow \text{add}(T), \tag{8}$$

since $e_0 E_R^{\text{Pl}} e_0 \cong \text{End}_R(R) \cong R^{\text{op}}$. Let p (respectively, p^{op}) be the number of indecomposable direct summands of $\text{Pl}(R)$ (respectively, $\text{Pl}(R^{\text{op}})$) By definition of E_R^{Pl} , the number p also equals the number of simple E_R^{Pl} -modules. Which in turn equals the number of indecomposable summands of T since T is tilting. Let s (respectively, s^{op}) be the number of indecomposable direct summands of $\text{SUB}(R)$ (respectively, $\text{SUB}(R^{\text{op}})$). By (8), $s^{\text{op}} \leq p$ (in particular, s^{op} is finite). Moreover, $\text{Pl}(R) \subseteq \text{SUB}(R)$

implies $p \leq s$. It follows from [36, Theorem 1.1] that $s = s^{\text{op}}$. Summing up, we have that $s^{\text{op}} = p = s$. In particular, this yields equivalences $\text{pi}(R) \cong \text{sub}(R)$, and therefore $E_R \cong E_R^{\text{Pl}}$ so proves (a). Moreover, using $s^{\text{op}} = p$ and Proposition 4.6 the inclusions (7) and (8) are equivalences

$$\text{sub}(R^{\text{op}}) \cong \text{add}(T) = \text{sub}(E_R^{\text{Pl}} e_0) \cap \text{fac}(E_R^{\text{Pl}} e_0). \tag{9}$$

In particular, this shows part (d).

By definition, the Ringel dual of E_R is $\mathfrak{R}(E_R) = \text{End}_{E_R}(T)^{\text{op}}$. Using $\text{sub}(R^{\text{op}}) \cong \text{add}(T)$ we obtain $\text{End}_{E_R}(T)^{\text{op}} \cong \text{End}_{R^{\text{op}}}(\text{SUB}(R^{\text{op}}))^{\text{op}}$. Under the standard k -duality the latter identifies with $\text{End}_R(\text{FAC}(R_R^\dagger))$. This completes the proof of the main Ringel duality statement as given in formula (4). As a consequence we get the equivalence $D^b(\text{sub}(R)) \cong D^b(\text{fac}(R^\dagger))$.

We now consider part (b). By part (a) we know $E_R \cong E_R^{\text{Pl}}$, and as R is ideally ordered Theorem 2.14 implies that E_R^{Pl} is both left and right strongly quasi-hereditary with respect to the ideal layer function. An algebra which is left and right strongly quasi-hereditary with respect to the same ideal layer function has global dimension at most two by [35, first Proposition in A.2]. Proposition 3.4 shows that $E_R \cong E_R^{\text{Pl}}$ is also left ultra strongly quasi-hereditary, and so completes the proof of statement (b).

We now prove (c). Let $[M : S]$ denote the number of simple E_R -modules S that occur in a Jordan Hölder filtration of an E_R -module M . If a partial ordering on I induces a quasi-hereditary structure, then $[\Delta_i, S_i] = 1$ for all $i \in I$; as k is algebraically closed this is equivalent to $\text{End}_{E_R}(\Delta_i) \cong k$, see [18, Lemma 1.6].

Using the additional assumption in (c) that R is local, the ideally ordered condition produces a surjection between any two summands of $\text{Pl}(R)$ (as all principal ideals are monomial by Lemma 7.3). Hence the ideal layer function induces an ordering on the summands of $\text{Pl}(R)$ of the form $\Lambda_0 < \Lambda_1 < \dots < \Lambda_t$. Now consider another partial order that also produces a quasi-hereditary ordering.

We first prove that both orderings have the same maximal element. If Λ_i is maximal with respect to the new order, then the projective module $P_i := P(\Lambda_i)$ is also a standard module in this order. If the new order gives rise to a quasi-hereditary structure then, as P_i is standard in this ordering, $[P_i : S_i] = 1$. As P_i is projective $[P_i, S_i] = \dim \text{Hom}_{E_R}(P_i, P_i)$. Under the anti-equivalence $\text{Hom}_R(-, \text{Pl}(R))$, described in formula (2), this implies $\dim \text{End}_R(\Lambda_i) = 1$. Hence the identity morphism must equal socle projection so Λ_i is the simple R -module, which is unique as R is assumed to be local. The simple R -module is the largest summand Λ_t of $\text{Pl}(R)$ under the ideal layer function ordering, and hence $i = t$.

Secondly, we assume that the orderings match for $k, k + 1, \dots, t$, let $\Lambda_j < \Lambda_k$ be an immediate predecessor of Λ_k under the new order, and aim to show that $j = k - 1$. As R is ideally ordered there is a surjection between Λ_j and Λ_{j+1} (where Λ_{j+1} exists as $j < k \leq t$). As they are labelled by the ideal layer function $\dim \Lambda_j > \dim \Lambda_{j+1}$ and there is a surjection $\Lambda_j \rightarrow \Lambda_{j+1}$. By the condition assumed in (c), the existence of this surjection implies an inclusion $\Lambda_{j+1} \rightarrow \Lambda_j$. Together these produce a non-trivial endomorphism $\Lambda_j \rightarrow \Lambda_{j+1} \rightarrow \Lambda_j$ which does not factor over Λ_i for $i > j + 1$. Using the anti equivalence $\text{Hom}_R(-, \text{Pl}(R))$ again, this translates into a non-trivial endomorphism of P_j that does not factor over P_i for $i > j + 1$. In particular, the

standard object under the new order Δ_j is the cokernel of a morphism $P \rightarrow P_j$ where the summands of P are projective modules P_i such that $i > j$ under the new ordering, see [18, Lemma 1.1']. If $k \neq j + 1$, then both the trivial endomorphism and the non-trivial endomorphism constructed above do not factor via P and hence $\dim \text{Hom}_{E_R}(P_j, \Delta_j) \geq 2$. By considering the images of these morphisms we see $[\Delta_j : S_j] \geq 2$. This would imply that the new ordering does not give a quasi-hereditary structure. Therefore $j = k - 1$.

Finally, by proceeding in this way we recover the ideal order and conclude that there is only one quasi-hereditary structure.

We show part (e). To prove (5), we explain the following chain of subcategories

$$\text{Filt}(\Delta) = \text{sub}(T) \subseteq \text{sub}(E_R e_0) \subseteq \text{sub}(E_R) \subseteq \text{Filt}(\Delta).$$

By part (b), E_R is right strongly quasi-hereditary. The first equality holds for all right strongly quasi-hereditary algebras, for example by a dual version of [35, Proposition A.1]. Using (9) and part (a), we see that $T \in \text{sub}(E_R e_0)$ so $\text{sub}(T) \subseteq \text{sub}(E_R e_0)$. The next inclusion follows from $E_R e_0 \subseteq E_R$. The last inclusion holds for any right strongly quasi-hereditary algebra using that $E_R \in \text{Filt}(\Delta)$, which is closed under submodules as noted in Corollary 2.16. Using (9) and the fact that E_R is left ultra strongly quasi-hereditary by part (b), dual arguments establish the following chain

$$\text{Filt}(\nabla) = \text{fac}(T) \subseteq \text{fac}(E_R e_0) \subseteq \text{Filt}(\nabla)$$

(the last inclusion was also shown in the proof of Corollary 3.6). This implies (6) and completes the proof of part (e). □

For a monomial algebra R there is an equivalence of additive categories $\langle \text{rad}^i R \mid i = 1, \dots, m \text{ for } \text{rad}^m R \cong 0 \rangle \cong \text{pi}(R)$, and so E_R^{PI} is Morita equivalent to

$$\text{End}_R\left(\bigoplus_{i=1}^m \text{rad}^i R\right).$$

This construction is considered in the general context of pre-radicals in Conde’s thesis. An additional special feature of the ideally ordered algebras is that $\text{pi}(R) \cong \text{sub}(R)$, and this property does not hold for general monomial algebras. For example, consider the following example that was communicated to us by Xiao-Wu Chen.

Example 5.2 Let R be the path algebra of the following quiver with monomial relations.

$$1 \begin{matrix} \xleftarrow{x_1} \\ \xrightarrow{x_2} \\ \xrightarrow{x_3} \end{matrix} 2 \begin{matrix} \xleftarrow{y_1} \\ \xrightarrow{y_2} \\ \xrightarrow{y_3} \end{matrix} 3 \quad y_j x_i = 0 \text{ for } i \neq j.$$

Then the left ideal $I = (x_1 + x_2, x_2 + x_3)$ is indecomposable but not principal.

Remark 5.3 We give several further remarks on this result.

(1) For the non-monomial algebra $R = R_2$ in Example 2.8(b), formula (4) from the theorem fails but the following Ringel duality formula holds:

$$\mathfrak{R}(\text{End}_R(\text{Pl}(R))) \cong \text{End}_{R^{\text{op}}}(\text{Pl}(R^{\text{op}}))^{\text{op}}. \tag{10}$$

For ideally ordered monomial algebras this formula coincides with formula (4) above. Unfortunately, we were not able to find a more general setup where the formula (10) works.

Knörrer invariant algebras [27, Section 6.4.], see Example 1.4(3) and Sect. 6.1, and truncated free algebras $R = k\langle x_1, \dots, x_l \rangle / (x_1, \dots, x_l)^m$ satisfy the additional condition imposed in (c).

(2) The statement that $D^b(\text{sub}(R)) \cong D^b(\text{fac}(R^\dagger))$ is related to Ringel’s [36, Remark before Corollary 2.2]. It would be interesting to see in what generality this equivalence holds.

We observe that it holds for $k\langle x, y, z \rangle / (p, zx, xy, zy, yz, z^2)$ where p runs over all paths of length 3, which is not ideally ordered but in which every principal left ideal is isomorphic to a monomial ideal. Indeed, in this case the equivalence is given by a tilting module which is obtained by mutating the characteristic tilting module (for the quasi-hereditary algebra structure defined by the ideal layer function) once.

(3) Consider $R = k\langle x, y \rangle / (x^3, y^3, y^2x, yx^2, xy)$, which is an ideally ordered finite dimensional local monomial algebra. Then there is a surjection $Rx \rightarrow Ry$ but Ry does not include into Rx . One can check that the order $R < Ry < Rx < Rx^2$ on indecomposable submodules of R defines a (left but not right strongly) quasi-hereditary structure on $E_R := \text{End}_R(\text{SUB}(R))$. In particular, in this case the ideal order is not the unique quasi-hereditary order.

(4) Part (c) can fail if R is not local (even if all the other conditions are satisfied). Indeed consider for example the algebra $R = kQ/J^2$ where

$$Q := \begin{array}{c} \curvearrowright \\ 1 \longrightarrow 2 \end{array}$$

and J is the ideal generated by all arrows. Then R is ideally ordered and for every surjection between principal left ideals $\Gamma \rightarrow \Lambda$ there is an inclusion $\Lambda \rightarrow \Gamma$. The order $P_2 < P_1 < S_1$ defines a quasi-hereditary structure on $E_R^{\text{Pl}} = \text{End}_R(\text{Pl}(R))$ which is not left strongly quasi-hereditary. Hence, it differs from the quasi-hereditary structure defined by the ideal layer function (where $P_2 = P_1 < S_1$), and there is no unique quasi-hereditary structure in this case.

(5) It is true that R is ideally ordered iff R^{op} is ideally ordered, and using this fact one can also prove the theorem without relying on Ringel’s result [36, Theorem 1.1].

6 Applications and examples

We discuss some relationships between Theorem 5.1 and several classes of algebras that have been studied in separate work.

6.1 Hille and Ploog’s algebras

The results of this paper were originally motivated by an investigation in [27] of a class of geometrically inspired quasi-hereditary algebras introduced by Hille and Ploog [24] for which the Ringel duality formula has a geometric interpretation, and we briefly recall this geometric setup and these algebras below.

As the geometric background, consider a type A_n configuration of intersecting rational curves C_1, \dots, C_n in a smooth, rational, projective surface X with negative self-intersection numbers $C_i \cdot C_i =: -\alpha_i \leq -2$. Starting with this data, Hille and Ploog consider the full triangulated subcategory

$$\langle \mathcal{O}_X(-C_1 - \dots - C_n), \mathcal{O}_X(-C_2 - \dots - C_n), \dots, \mathcal{O}_X(-C_n), \mathcal{O}_X \rangle \subset D^b(\text{Coh}(X)),$$

where we recall that $\mathcal{O}_X(-D)$ denotes the line bundle occurring as the ideal sheaf of an effective divisor $D \subset X$. Hille and Ploog show that this subcategory carries an (exact) tilting object Λ . To do this they make use of universal (co)extensions, see [18] and also [23] for the special case of vector bundles on a rational surface. We briefly recall the definition in this setting.

Definition 6.1 Consider an ordered pair of vector bundles $\mathcal{E}_1, \mathcal{E}_2$ on a smooth projective rational surface X . Their *universal (co)extension* is defined to be the vector bundle occurring in the middle of the short exact sequence

$$\mathcal{E}_2 \otimes \text{Ext}_X^1(\mathcal{E}_1, \mathcal{E}_2)^\dagger \rightarrow \mathcal{F} \rightarrow \mathcal{E}_1 \tag{extension}$$

$$\mathcal{E}_2 \rightarrow \mathcal{F} \rightarrow \mathcal{E}_1 \otimes \text{Ext}_X^1(\mathcal{E}_1, \mathcal{E}_2) \tag{coextension}$$

where both sequences are determined by the identity element in $\text{End}(\text{Ext}^1(\mathcal{E}_1, \mathcal{E}_2)) \cong \text{Ext}_X(\mathcal{E}_1, \mathcal{E}_2) \otimes \text{Ext}_X(\mathcal{E}_1, \mathcal{E}_2)^\dagger$.

Hille and Ploog show that

$$\mathbb{E} := (\mathcal{O}_X(-C_1 - \dots - C_n), \mathcal{O}_X(-C_2 - \dots - C_n), \dots, \mathcal{O}_X(-C_n), \mathcal{O}_X)$$

is an exceptional sequence of line bundles and that iterated universal extension along this sequence produces a tilting bundle Λ , see [24, Section 2]. This defines a corresponding algebra

$$\Lambda_{[\alpha_1, \dots, \alpha_n]} := \text{End}_X(\Lambda)^{\text{op}},$$

where we assume that Λ is taken to be a basic representative of the tilting object. These algebras are quasi-hereditary by construction.

We note that the algebra depends on the choice of consecutive ordering for the labelling of the curves and that there are two choices, C_1, \dots, C_n or C_n, \dots, C_1 , for the same geometric set up that produce two different algebras $\Lambda_{[\alpha_1, \dots, \alpha_n]}$ and $\Lambda_{[\alpha_n, \dots, \alpha_1]}$. This phenomenon is explained by the following result.

Proposition 6.2 *There is an isomorphism of algebras $\mathfrak{R}(\Lambda_{[\alpha_1, \dots, \alpha_n]}) \cong \Lambda_{[\alpha_n, \dots, \alpha_1]}^{\text{op}}$.*

Remark 6.3 The algebra $\Lambda_{[\alpha_1, \dots, \alpha_n]}$ can in fact be realised in the form E_R where R is an ideally ordered monomial Knörrer invariant algebra, as we describe below. Then Proposition 6.2 is an consequence of Theorem 5.1. However, the following alternative, short, geometric proof was explained to us by Agnieszka Bodzenta; indeed it was the existence of a Ringel duality formula in this special case that inspired the representation-theoretic generalisation in this paper. Work of Bodzenta and Bondal also realises a Ringel duality associated to birational morphisms of smooth surfaces by gluing t-structures with reversed orderings, see [7].

Proof Let X be a smooth, rational, projective surface containing a type A_n configuration of rational curves with self-intersection numbers $\alpha := [\alpha_1, \dots, \alpha_n]$. Consider the exceptional sequence \mathbb{E} in the Hom-finite abelian category $\text{Coh}(X)$. By definition, $\Lambda_\alpha := \text{End}_X(\Lambda)^{\text{op}}$, where $\Lambda \in \text{Coh}(X)$ is obtained from \mathbb{E} by taking iterated universal extensions and by passing to a basic representative, see [27, Section 2.3]. On the other hand, taking iterated universal coextensions of \mathbb{E} yields $T \in \text{Coh}(X)$ (again we replace this by a basic version if necessary) and it follows from [18, paragraph above Proposition 3.1.] that there is an algebra isomorphism

$$\mathfrak{R}(\text{End}_X(\Lambda)) \cong \text{End}_X(T)^{\text{op}} \tag{11}$$

where $\mathfrak{R}(\text{End}_X(\Lambda))$ denotes the Ringel dual of $\text{End}_X(\Lambda)$. More precisely, since \mathbb{E} is standardisable, Dlab and Ringel [18, Theorem 2] show that

$$\text{Hom}_X(\Lambda, -) : \mathit{Filt}(\mathbb{E}) \rightarrow \mathit{Filt}(\Delta_{\Lambda_\alpha})$$

defines an exact equivalence sending \mathbb{E} to the sequence of standard modules Δ_{Λ_α} . By Ringel [34, p.217 and Proposition 2], the characteristic tilting module $T_{\Lambda_\alpha} \in \text{mod-}\Lambda_\alpha$ is obtained from Δ_{Λ_α} by iterated universal coextensions (and passing to a basic module if necessary). In particular, the exact equivalence $\text{Hom}_X(\Lambda, -)$ sends T to T_{Λ_α} . Combining this with definition of the Ringel dual we see

$$\mathfrak{R}(\Lambda_\alpha) := \text{End}_{\Lambda_\alpha}(T_{\Lambda_\alpha})^{\text{op}} \cong \text{End}_X(T)^{\text{op}}$$

as claimed.

Now consider the duality

$$\begin{aligned} \ddagger : D(\text{QCoh}(X)) &\rightarrow D(\text{QCoh}(X)) \\ \mathcal{E}^\ddagger &:= R\mathcal{H}om_X(\mathcal{E} \otimes_X \mathcal{O}(-C_1 - C_2 - \dots - C_n), \mathcal{O}). \end{aligned}$$

Then $\mathbb{E}^\ddagger = (\mathcal{O}(-C_1 - C_2 - \dots - C_n), \mathcal{O}(-C_1 - C_2 - \dots - C_{n-1}), \dots, \mathcal{O}(-C_1), \mathcal{O})$ and T^\ddagger is obtained from this sequence by iterated universal extensions. By definition, $\Lambda_{[\alpha_n, \dots, \alpha_1]} \cong \text{End}_X(T^\ddagger)^{\text{op}}$. Since \ddagger is a duality, $\Lambda_{[\alpha_n, \dots, \alpha_1]} \cong \text{End}_X(T^\ddagger)^{\text{op}} \cong \text{End}_X(T)$. In combination with (11) this completes the proof. \square

Remark 6.4 We note that there is a change in conventions for compositions of morphisms between this paper and [27]. This corresponds to exchanging algebras with their opposite algebras, or left modules with right modules. The effect this has on the quasi-hereditary structure and Ringel duality is as follows: if A is a quasi-hereditary algebra with defining layer function L and characteristic tilting module T , then T^\ddagger is the characteristic tilting module for A^{op} where $\ddagger: A\text{-mod} \rightarrow A^{\text{op}}\text{-mod}$ denotes the standard k -duality and the layer function on A^{op} on is L^\ddagger defined by $L^\ddagger(S^\ddagger) := L(S)$. In particular, $\mathfrak{R}(A^{\text{op}}) \cong \mathfrak{R}(A)^{\text{op}}$.

We briefly recap how the algebras Λ defined by Hille and Ploog fit into the general setup of Theorem 5.1. To do so we recall the definition of the Hirzebruch–Jung continued fraction expansion, the Knörrer invariant algebras $K_{r,a}$, and a description of the form $\Lambda_\alpha \cong E_{K_{r,a}}$.

Definition 6.5 For coprime integers $0 < a < r$ the Hirzebruch–Jung continued fraction $[\alpha_1, \dots, \alpha_n]$ is the collection of integers $\alpha_i \geq 2$ defined by

$$\frac{r}{a} = \alpha_1 - \frac{1}{\alpha_2 - \frac{1}{\dots - \frac{1}{\alpha_n}}}$$

Definition 6.6 ([27, Definitions 4.6, 6.20 and Corollary 6.27]) For coprime integers $0 < a < r$ the Knörrer invariant algebra $K_{r,a}$ is defined to be

$$K_{r,a} := \frac{\mathbb{C}\langle z_1, \dots, z_l \rangle}{\left\langle \begin{array}{l} z_i(z_i^{\beta_i-2})(z_{i+1}^{\beta_{i+1}-2}) \cdots (z_{j-1}^{\beta_{j-1}-2})(z_j^{\beta_j-2})z_j = 0 \text{ for } j \leq i \\ z_i z_j = 0 \text{ if } i < j \end{array} \right\rangle}$$

where the parameters $l \geq 1$ and $\beta_i \geq 2$ are defined by the Hirzebruch–Jung continued fraction expansion $[\beta_1, \dots, \beta_l]$ for the fraction $r/(r - a)$.

We recall that the results of [27, Section 6.4] describe the monomial ideal structure on $K_{r,a}$, and in particular combining [27, Theorem 6.26] and [27, Propositions 6.22 and 6.24] yields the following result.

Proposition 6.7 *The Knörrer invariant algebra $K_{r,a}$ is an ideally order monomial algebra and there is an isomorphism of quasi-hereditary algebras $\Lambda_{[\alpha_1, \dots, \alpha_n]} \cong E_{K_{r,a}}$ where $[\alpha_1, \dots, \alpha_n]$ is defined by the Hirzebruch–Jung continued fraction expansion of r/a .*


Suppose that $aa' \equiv 1 \pmod{r}$. If $r/a = [\alpha_1, \dots, \alpha_n]$, then $r/a' = [\alpha_1, \dots, \alpha_n]$. Similarly, if $r/(r - a) = [\beta_1, \dots, \beta_l]$, then $r/(r - a') = [\beta_1, \dots, \beta_l]$. Using this

result it can be seen from the explicit definition of $K_{r,a}$ that $K_{r,a'} \cong K_{r,a}^{\text{op}}$. As a result $\Lambda_{[\alpha_n, \dots, \alpha_1]} \cong E_{K_{r,a}^{\text{op}}}$ by Proposition 6.7, and hence Theorem 5.1 is a generalisation of Proposition 6.2.


6.2 Example of an application of the Ringel duality formula

In this section we consider as an example the pair of algebras $\Lambda_{[3,2]}$ and $\Lambda_{[2,3]}$. After giving explicit presentations, we discuss their relationship via Ringel duality, their construction from related Knörrer invariant algebras, and explicitly list the distinguished modules in their quasi-hereditary structures in order to verify the Ringel duality formula.

Firstly, by [27, Proposition 6.18] the algebras $\Lambda_{[3,2]}$ and $\Lambda_{[2,3]}$ can respectively be presented as the path algebra of the following quivers with relations:



and




Secondly, the Ringel duality formula of Proposition 6.2 states that

$$\mathfrak{R}(\Lambda_{[3,2]}) \cong (\Lambda_{[2,3]})^{\text{op}}.$$

Thirdly, by Proposition 6.7 the corresponding Knörrer invariant algebras are


$$K_{[3,2]} := K_{5,2} \cong \frac{\mathbb{C}\langle z_1, z_2 \rangle}{(z_1^2, z_2^3, z_1 z_2, z_2^2 z_1)} \quad \text{and} \quad K_{[2,3]} := K_{5,3} \cong \frac{\mathbb{C}\langle z_1, z_2 \rangle}{(z_1^3, z_2^2, z_1 z_2, z_2 z_1^2)}$$

and these can be presented via the following monomial diagrams:



where the nodes represent the monomial basis of $K_{r,a}$ with the root of the tree representing 1 and the arrows labelled i representing left multiplication by z_i of the node at the source equalling the node at the target. Using these monomial diagrams one can show that $K_{[3,2]} \cong K_{[2,3]}^{\text{op}}$ and to calculate all the left monomial ideals. The left

monomial ideals for $K_{[3,2]}$ are $M_0 \cong (1)$, $M_1 \cong (z_1)$ and $M_2 \cong (z_2z_1)$ and the left monomial ideals for $K_{[2,3]}$ are $N_0 \cong (1)$, $N_1 \cong (z_1)$ and $N_2 \cong (z_1^2)$. These can be represented pictorially as subsets of the monomial diagrams by



It is explicit that $\Lambda_{[3,2]} \cong E_{K_{[3,2]}} := \text{End}_{K_{[3,2]}}(\oplus M_i)$ and $\Lambda_{[2,3]} \cong E_{K_{[2,3]}} := \text{End}_{K_{[2,3]}}(\oplus N_i)$.

In order to explicitly verify the Ringel duality formula in this case we first describe the quasi-hereditary structure by calculating the projective P_i , injective I_i , standard Δ_i , costandard ∇_i , and characteristic tilting T_i objects for each algebra. We list these modules in the table below in terms of the simples, S_i notated by i , occurring in their composition series with the heads written at the top.

$\Lambda_{[3,2]}$	P_i	I_i	Δ_i	∇_i	T_i	
$i = 0$	$\begin{array}{c} 0 \\ c_1 \mid \\ 1 \\ a_1 / c_2' \setminus k_2 \\ 0 \ 2 \ 0 \\ a_2 \setminus / c_1 \\ 1 \\ a_1 / \setminus k_2 \\ 0 \ 0 \end{array}$	$\begin{array}{c} 0 \\ c_1 \mid \\ 1 \\ c_2 / \setminus k_2 \\ 2 \ 0 \\ a_2 \setminus / c_1 \\ 1 \\ k_2 \setminus \ 0 \end{array}$	$\begin{array}{c} 0 \\ c_1 \mid \\ 1 \\ c_2 / \setminus k_2 \\ 2 \ 0 \\ a_2 \setminus / c_1 \\ 1 \\ 0 \end{array}$	$\begin{array}{c} 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{array}$	$\begin{array}{c} 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{array}$	$\begin{array}{c} 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{array}$
$i = 1$	$\begin{array}{c} 1 \\ a_1 / c_2' \setminus k_2 \\ 0 \ 2 \ 0 \\ a_2 \setminus / c_1 \\ 1 \\ a_1 / \setminus k_2 \\ 0 \ 0 \end{array}$	$\begin{array}{c} 0 \\ c_1 \mid \\ 1 \\ c_2 / \setminus k_2 \\ 2 \ 0 \\ a_2 \setminus / c_1 \\ 1 \end{array}$	$\begin{array}{c} 1 \\ a_1 / \setminus k_2 \\ 0 \ 0 \end{array}$	$\begin{array}{c} 0 \\ c_1 \mid \\ 1 \\ 1 \\ 2 \end{array}$	$\begin{array}{c} 0 \\ c_1 \\ 1 \\ a_1 / \setminus k_2 \\ 0 \ 0 \end{array}$	$\begin{array}{c} 0 \\ c_1 \\ 1 \\ a_1 / \setminus k_2 \\ 0 \ 0 \end{array}$
$i = 2$	$\begin{array}{c} 2 \\ a_2 \mid \\ 1 \\ a_1 / \setminus k_2 \\ 0 \ 0 \end{array}$	$\begin{array}{c} 0 \\ c_1 \mid \\ 1 \\ c_2 \mid \\ 2 \end{array}$	$\begin{array}{c} 2 \\ a_2 \mid \\ 1 \\ a_1 / \setminus k_2 \\ 0 \ 0 \end{array}$	$\begin{array}{c} 0 \\ c_1 \mid \\ 1 \\ c_2 \mid \\ 2 \end{array}$	$\begin{array}{c} 0 \\ c_1 \mid \\ 1 \\ a_1 / c_2' \setminus k_2 \\ 0 \ 2 \ 0 \\ a_2 \setminus / c_1 \\ 1 \\ a_1 / \setminus k_2 \\ 0 \ 0 \end{array}$	$\begin{array}{c} 0 \\ c_1 \mid \\ 1 \\ a_1 / c_2' \setminus k_2 \\ 0 \ 2 \ 0 \\ a_2 \setminus / c_1 \\ 1 \\ a_1 / \setminus k_2 \\ 0 \ 0 \end{array}$

Remark 6.8 We observe some further properties of, and relations between, the modules in the tables above. These are all special cases of the general theory developed above.

- (1) If $i \leq j$ in the partial order, then there is an inclusion $P_j \subseteq P_i$ (and a projection $I_j \twoheadrightarrow I_i$). This holds for all left (respectively right) strongly quasi-hereditary algebras. In other words, in this situation it is a consequence of Theorem 2.14.
- (2) Every submodule of a standard module Δ_i or a projective module P_i is filtered by standard modules. This is a consequence of Corollary 2.16. Dually, quotients of costandard modules ∇_i or injective modules I_i are filtered by costandard modules, again by Corollary 2.16.
- (3) For both algebras the only simple costandard module is ∇_0 . One can check that the corresponding projective modules P_0 are filtered by costandard modules. This illustrates Proposition 3.4 in these cases.
- (3^{op}) For both algebras the only simple standard module is Δ_0 . The corresponding injective hulls I_0 are not filtered by standard modules. In other words, the algebras $\Lambda_{[3,2]}$ and $\Lambda_{[2,3]}$ are not right ultra strongly quasi-hereditary.
- (4) The summands T_i of the characteristic tilting module are precisely those indecomposable modules which are both quotients and submodules of the projective module P_0 , see Theorem 5.1 (d). In particular, they have head S_0 and a socle in $\text{add}(S_0)$.

6.3 Auslander–Dlab–Ringel algebras

Recent results of Conde–Erdmann [13], and work in Conde’s thesis, produce a Ringel duality formula similar to that of Theorem 5.1 for the class of Auslander–Dlab–Ringel (ADR) algebras.

Definition 6.9 Let R be a finite dimensional algebra of Loewy length L_R . Define the additive subcategory

$$\text{adr}(R) := \text{add} \{ Re/\text{rad}^i Re \mid e \text{ a primitive idempotent and } i = 1, \dots, L_R \}$$

and let $\text{ADR}(R) := \bigoplus_{M \in \text{ind}(\text{adr}(R))} M$ be the direct sum of indecomposable elements of the additive category $\text{adr}(R)$ up to isomorphism. Then the associated *ADR algebra* is defined to be

$$E_R^{\text{ADR}} := \text{End}_R(\text{ADR}(R)).$$

This is the basic algebra Morita equivalent to $\text{End}_R(\bigoplus_{i=1}^{L_R} R/\text{rad}^i R)$. In particular, the indecomposable modules in $\text{adr}(R)$ are exactly those of the form $Re/\text{rad}^i Re$ for e a primitive idempotent and $1 \leq i \leq L_{Re}$ where L_{Re} is the Loewy length of Re .

Remark 6.10 We remark that the ADR algebra defined here is the opposite algebra of the ADR algebra defined by Conde and Erdmann in [13], however the effect on the quasi-hereditary structure is straightforward as is explained in Remark 6.4.

The ADR algebra E_R^{ADR} is quasi-hereditary for the layer function $l(Re_i/\text{rad}^l Re_i) := L_R - l$; this induces the partial ordering

$$Re_i/\text{rad}^l Re_i < Re_j/\text{rad}^k Re_j \iff l > k$$

on indecomposable modules in $\text{adr}(R)$. Indeed it is left ultra strongly quasi-hereditary (see [11, Section 5]), and Conde and Erdmann obtain the following Ringel duality formula for ADR algebras satisfying a regularity condition; we recall that a module is *rigid* if its radical and socle series coincide.

Theorem 6.11 *Let R be an Artin algebra with Loewy length L . If all projective and injective indecomposable R -modules are rigid with Loewy length L , then*

$$\mathfrak{R}(E_R^{\text{ADR}}) \cong (E_{R^{\text{op}}}^{\text{ADR}})^{\text{op}}.$$

That is, the Ringel dual of E_R^{ADR} is isomorphic to the opposite algebra of $E_{R^{\text{op}}}^{\text{ADR}}$.

This formula looks very similar to the formula in Theorem 5.1 of this paper. However, in general $E_R \not\cong E_R^{\text{ADR}}$ and there does not appear to be any reason to think the overlap is large.

For example, ADR algebras are not left and right strongly hereditary in general and so not all ADR algebras are in the E_R algebra class. Moreover, it can be seen that Hille and Ploog’s algebras are not always ADR algebras. Indeed, in the example of Sect. 6.2 the modules $R/\text{rad}^i R$ are straightforward to calculate from the monomial diagrams, and the additive category generated by such objects can be seen to coincide with the additive category $\text{sub}(R) \cong \text{pi}(R)$ for $R = K_{[3,2]}$ so $E_R \cong E_R^{\text{ADR}}$ but not for $R^{\text{op}} \cong K_{[2,3]}$ where $E_R \not\cong E_R^{\text{ADR}}$.

Indeed, the results of Conde and Erdmann also only describe the Ringel dual of an ADR algebra when the dual is also an ADR algebra. However, as can be seen in the example of Sect. 6.2, there are examples of ADR algebras of the form E_R whose dual is not an ADR algebra but whose Ringel dual can still be described by Theorem 5.1: for $R = K_{[3,2]}$ and $R^{\text{op}} \cong K_{[2,3]}$

$$\mathfrak{R}(E_R^{\text{ADR}}) \cong \mathfrak{R}(E_R) \cong E_{R^{\text{op}}}^{\text{op}} \not\cong E_{R^{\text{op}}}^{\text{ADR}}.$$

Indeed, it is also straightforward to calculate the socle and radical filtrations in this example and hence clear to see that $K_{[3,2]}$ is rigid whereas $K_{[2,3]}$ is not.

Whilst these classes of algebras may not be related in general, there are cases which fall into both classes of algebras. Recall the monomial algebras $R := kQ/J^m$ of Example 1.4(2) which are ideally ordered and for which $\text{sub}(R) \cong \text{pi}(R) \subset \text{adr}(R)$. In particular, in this case E_R is a corner algebra of E_R^{ADR} : i.e. there is an idempotent $e \in E_R^{\text{ADR}}$ such that $E_R \cong eE_R^{\text{ADR}}e$.

Proposition 6.12 *Let Q be a finite quiver without sources and J be the two-sided ideal generated by all arrows in Q . Then $R := kQ/J^m$ is an ideally ordered monomial algebra and there is an isomorphism of quasi-hereditary algebras $E_R^{\text{ADR}} \cong E_R$.*

Proof The algebra R has Loewy length m and, as noted in Example 1.4(2), any monomial ideal is isomorphic to $Re/\text{rad}^l Re$ for some $l = 1, \dots, m$ and some primitive idempotent $e \in R$, hence R is ideally ordered and $\text{pi}(R) \subset \text{adr}(R)$. As R is ideally ordered $\text{sub}(R) \cong \text{pi}(R)$ by Theorem 5.1 (a), and hence to show that $E_R \cong E_R^{\text{ADR}}$ it is sufficient to show that $\text{adr}(R) \subset \text{sub}(R)$.

To show this consider an indecomposable object of $\text{adr}(R)$. This is necessarily of the form $Re_i/\text{rad}^l Re_i$ for some primitive idempotent e_i corresponding to a vertex $i \in Q$ and integer $l = 1, \dots, m$. As Q has no sources it follows that there exists a series of arrows $j_{m-l} \xrightarrow{a_{m-l}} \dots \xrightarrow{a_3} j_2 \xrightarrow{a_2} j_1 \xrightarrow{a_1} i$ such that the path $a := a_{m-l} \dots a_1$ induces a homomorphism $Re_i \xrightarrow{a} Re_{j_{m-l}}$ of indecomposable projective R -modules. By construction this has kernel $\text{rad}^l Re_i$, and hence there is an inclusion $Re_i/\text{rad}^l Re_i \rightarrow Re_{j_{m-l}}$. In particular $Re_i/\text{rad}^l Re_i \in \text{sub}(R)$, and hence $\text{adr}(R) \subset \text{sub}(R)$. Hence $E_R \cong E_R^{\text{ADR}}$.

Whilst the layer functions defining the quasi-hereditary structures on E_R and E_R^{ADR} are not identical in general, we claim that the corresponding orderings do induce the same standard modules and hence the same quasi-hereditary structure on $E_R^{\text{ADR}} \cong E_R$. To show this we let $P_{i,l}$ denote the projective $E_R \cong E_R^{\text{ADR}}$ -module $P(Re_i/\text{rad}^l Re_i)$ and $S_{i,l}$ denote its simple quotient. We recall the order for E_R^{ADR} is defined by $S_{i,l} < S_{j,k} \Leftrightarrow l > k$ and the order for E_R is defined by $S_{i,l} < S_{j,k} \Leftrightarrow \dim Re_i/\text{rad}^l Re_i > \dim Re_j/\text{rad}^k Re_j$. In particular, both orderings induce strongly quasi-hereditary structures, and hence for both orderings there are short exact sequences defining the respective standard modules

$$0 \rightarrow \bigoplus P_{j,k} \rightarrow P_{i,l} \rightarrow \Delta(Re_i/\text{rad}^l Re_i) \rightarrow 0 \tag{12}$$

for each projective module $P_{i,l}$, see Definition 2.3. Hence to show that the two orderings induce the same quasi-hereditary structure it is sufficient to show that the projective submodules $P_{j,k}$ of $P_{i,l}$ appearing in (12) are the same for both orderings. For this we note that under the additive anti-equivalence

$$\text{Hom}_R(-, \text{PI}(R)): \text{pi}(R) \rightarrow E_R\text{-proj}$$

an E_R -module $P_{j,k}$ is a proper submodule of $P_{i,l}$ if and only if the corresponding R -module $Re_j/\text{rad}^k Re_j$ is a proper quotient of $Re_i/\text{rad}^l Re_i$. This in turn is equivalent to $\dim Re_j/\text{rad}^k Re_j < \dim Re_i/\text{rad}^l Re_i$ and is also equivalent to $i = j$ and $k < l$. This shows that the two orderings induce the same quasi-hereditary structure. \square

It is a natural question whether it is possible to find an expanded class of algebras with a more general Ringel duality formula that encompasses both Theorems 5.1 and 6.11.

6.4 Nilpotent quiver algebras

The nilpotent quiver algebras introduced by Eiriksson and Sauter [20, Section 3] are a class of quasi-hereditary algebras.

Definition 6.13 Let $Q = (Q_0, Q_1)$ be a finite quiver. For $s \in \mathbb{Z}_{>0}$ the nilpotent quiver algebra is defined to be

$$N_s(Q) := kQ^{(s)}/J$$

where $Q^{(s)}$ is the staircase quiver $Q^{(s)}$ defined by having vertices i_l for $i \in Q_0$ and $l \in \{1, \dots, s\}$ and arrows

$$\begin{aligned} b(i_l) : i_{l+1} &\rightarrow i_l && \text{for } i \in Q_0 \text{ and } l \in \{1, \dots, s-1\}, \\ a_l : h(a)_{l-1} &\rightarrow t(a)_l && \text{for } a \in Q_1 \text{ and } l \in \{2, \dots, s\}, \end{aligned}$$

and where $J \subset kQ^{(s)}$ is the two-sided ideal generated by the relations

$$\begin{aligned} b(t(a)_l)a_{l+1} &= a_l b(h(a)_{l-1}) && \text{for all } a \in Q_1 \text{ and } l \in \{2, \dots, s\}, \\ b(t(a)_1)a_2 &= 0 && \text{for all } a \in Q_1. \end{aligned}$$

Remark 6.14 We remark again that the nilpotent quiver algebra defined here is the opposite algebra of the nilpotent quiver algebra defined by Eiriksson and Sauter in [20], however the effect on the quasi-hereditary structure is straightforward as is explained in Remark 6.4.

It follows from [20, Proposition 3.15] that all nilpotent quiver algebras $N_s(Q)$ are right strongly quasi-hereditary and left ultra strongly quasi-hereditary for the quasi-hereditary structure determined by the layer function $L(i_l) = s - l$.

In particular, for $R = kQ/J^m$ the ADR and nilpotent quiver algebras are related as follows.

Proposition 6.15 *Let Q be a finite quiver, J the two-sided ideal generated by all arrows in Q , and m a positive integer. Then there is an isomorphism of quasi-hereditary algebras*

$$N_m(Q) \cong E_{kQ/J^m}^{\text{ADR}}$$

if and only if all projective kQ/J^m -modules have Loewy length m : i.e. Q contains no sinks and m is arbitrary or $m = 1$ and Q is arbitrary.

Proof Let $R = kQ/J^m$, and let $e_i \in R$ for $i \in Q_0$ denote the primitive idempotents corresponding to vertices of Q . Up to isomorphism, the indecomposable modules in $\text{adr}(R)$ are exactly $Re_i/\text{rad}^l Re_i$ for $1 \leq l \leq L(Re_i)$ and $i \in Q_0$, where $L(Re_i)$ is the Loewy length of the projective Re_i .

In particular, the maximal Loewy length of a projective module in $R := kQ/J^m$ is m and so the maximum possible number of non-isomorphic indecomposables in $\text{adr}(R)$ is $m|Q_0|$. But $|Q_0^{(m)}| = m|Q_0|$, so for E_R^{ADR} to be isomorphic to $N_m(Q)$ it is necessary that all projective R -modules have Loewy length m .

Now suppose that all projective R -modules do have Loewy length m and consider the algebra $E_R^{\text{ADR}} := \text{End}_R(\text{ADR})$. We start by labelling the indecomposable module


in $\text{adr}(R)$ corresponding to $Re_i/\text{rad}^l Re_i$ by i_l and hence label the corresponding primitive idempotent by e_{i_l} . There are indecomposable modules i_l for $i \in Q_0$ and $l \in \{1, \dots, m\}$, matching the definition of the vertices in the staircase quiver $Q^{(m)}$.

We now want to produce a morphism $N_m(Q) \rightarrow E_R^{\text{ADR}}$, and to do this we consider the morphisms between the indecomposable modules in $\text{adr}(R)$. Firstly, there are surjections $Re_i/\text{rad}^{l+1} Re_i \rightarrow Re_i/\text{rad}^l Re_i$ which we label by arrows $\beta(i_l): i_{l+1} \rightarrow i_l$ for $i \in Q_0$ and $l \in \{1, \dots, m-1\}$.

Secondly, an arrow $a: i \rightarrow j \in Q_1$ corresponds to a morphism of projectives $a: Re_j \rightarrow Re_i$ and for each l this induces a morphism $Re_j \rightarrow Re_i/\text{rad}^l Re_i$ with kernel $\text{rad}^{l-1} Re_j$ which in turn induces an injective morphism

$$Re_j/\text{rad}^{l-1} Re_j \rightarrow Re_i/\text{rad}^l Re_i$$

for each $l \in \{2, \dots, m\}$. We label these morphisms by $\rho(a)_l: h(a)_{l-1} \rightarrow t(a)_l$ for $a \in Q_1$ and $l \in \{2, \dots, m\}$. In particular, the morphisms described here match the arrows of the staircase quiver $Q^{(m)}$ under the identification $a_l = \rho(a)_l$ and $b(i_l) = \beta(i_l)$. In particular, an arrow $a: i \rightarrow j$ in Q corresponds to a morphism $Re_j \rightarrow Re_i$ which induces morphisms



where the relations $\beta(i_l)\rho(a)_{l+1} \cong \rho(a)_l\beta(j_{l-1})$ and $\beta(i_1)\rho(a)_2 \cong 0$ hold.

This allows us to define a morphism from the path algebra of the staircase algebra $kQ^{(m)}$ to E_R^{ADR} by

$$e_i \mapsto e_i, \quad b(i_l) \mapsto \beta(i_l), \quad \text{and} \quad a_l \mapsto \rho(a)_l$$

and, as the relations imposed on $kQ^{(m)}$ by $N_m(Q)$ are mapped to 0, this induces a morphism

$$\Phi: N_m(Q) \rightarrow E_R^{\text{ADR}}.$$

We will now show that Φ is surjective, and then calculate the dimensions of $N_m(Q)$ and E_R^{ADR} to show that it is an isomorphism.

Suppose that $f \in E_R^{\text{ADR}} := \text{End}_R(\text{ADR})$ is a morphism

$$f: Re_i/\text{rad}^l Re_i \rightarrow Re_j/\text{rad}^k Re_j$$

for some $i, j \in Q_0$ and $l, k \in \{1, \dots, m\}$. There is a surjection $\pi_{i,l}: Re_i \rightarrow Re_i/\text{rad}^l Re_i$ and so f gives a morphism $f \circ \pi_{i,l}: Re_i \rightarrow Re_j/\text{rad}^k Re_j$. There is

also a surjection $\pi_{j,k}: Re_j \rightarrow Re_j/\text{rad}^k Re_j$, and as Re_i is projective this induces a uniquely defined morphism $g: Re_i \rightarrow Re_j$ such that

$$\pi_{j,k} \circ g \cong f \circ \pi_{i,l}.$$

As a morphism between projective modules, the morphism $g: Re_i \rightarrow Re_j$ corresponds to an element $g \in e_i Re_j \subset R \cong \text{End}_R(\bigoplus Re_i)^{\text{op}}$. In particular, $g = \sum_p \lambda_p p \in kQ/J^m = R$ for scalars λ_p and homogeneous paths p from j to i in $kQ/J^m = R$ corresponding to morphisms $p: Re_i \rightarrow Re_j$.

We now work with one indecomposable path p , corresponding to a morphism $p: Re_i \rightarrow Re_j$, and suppose that p consists of $n := |p|$ arrows $p = a_n a_{n-1} \dots a_1$ for $a_i \in Q_1$. We define a corresponding path in $N_m(Q)$ from i_{l-n} to j_l by

$$(p)_l := (a_1)_l \dots (a_{n-1})_{l+1} (a_n)_{l-n+1}: i_{l-n} \rightarrow j_l$$

for $l \in \{n + 1, \dots, m\}$. Similarly, we define the path in $N_m(Q)$

$$(\pi_{i,l}) := b(i_l) b(i_{l+1}) \dots b(i_{m-1}): i_m \rightarrow i_l$$

for $i \in Q_0$ and $l \in \{1, \dots, m\}$ such that $\Phi((\pi_{i,l})) \cong \pi_{i,l}$. Then the morphism p factors over its kernel, which is $\text{rad}^{m-n} Re_i$, so

$$p \cong \Phi((p)_m (\pi_{i,m-n})).$$

Hence

$$\pi_{j,k} \circ p \cong \Phi((\pi_{j,k}) (p)_m (\pi_{i,m-n})).$$

Using the relations in $N_m(Q)$ we can rearrange this expression as

$$\begin{aligned} (\pi_{j,k}) (p)_m (\pi_{i,m-n}) &= b(j_k) \dots b(j_{m-1}) (p)_m (\pi_{i,m-n}) \\ &= (p)_k b(i_{k-n}) \dots b(i_{m-n-1}) (\pi_{i,m-n}) = (p)_k (\pi_{i,k-n}) \end{aligned}$$

where we note that if g is non-zero then $k - n \leq l$ and hence

$$(\pi_{j,k}) (p)_m (\pi_{i,m-n}) = (p)_k (\pi_{i,k-n}) = (p)_k b(i_{k-n}) \dots b(i_{l-1}) (\pi_{i,l}),$$

and hence

$$\Phi((p)_k b(i_{k-n}) \dots b(i_{l-1}) (\pi_{i,l})) \cong \pi_{j,k} \circ p.$$

Returning to the morphism $g = \sum \lambda_p p$ we see that

$$\sum_p \lambda_p \pi_{j,k} \circ p \cong \pi_{j,k} \circ \sum_p \lambda_p p \cong \pi_{j,k} \circ g \cong f \circ \pi_{i,l}$$

and we can now conclude that

$$\begin{aligned} \sum_p \lambda_p \Phi((p)_k b(i_{k-|p|}) \cdots b(i_{l-1})(\pi_{i,l})) \\ \cong \Phi\left(\sum_p \lambda_p (p)_k b(i_{k-|p|}) \cdots b(i_{l-1})\right) \circ \pi_{i,l} \cong f \circ \pi_{i,l} \end{aligned}$$

where $|p|$ is the length of the path p , but $\pi_{i,l}$ is surjective and hence

$$f \cong \Phi\left(\sum_p \lambda_p (p)_k b(i_{k-|p|}) \cdots b(i_{l-1})\right).$$

We conclude that Φ is a surjection, and we now show that this surjective morphism is in fact an isomorphism by calculating the dimensions of $N_m(Q)$ and E_R^{ADR} .

We first calculate the dimension of E_R^{ADR} by calculating the dimension of the morphisms between any two indecomposables in $\text{adr}(R)$. As shown above, a morphism in E_R^{ADR} of the form $f : Re_i / \text{rad}^l Re_i \rightarrow Re_j / \text{rad}^k Re_j$ is induced by a particular element in kQ/J^m corresponding to a morphism of projective modules $Re_i \rightarrow Re_j$. Such elements are spanned by the paths, and we now calculate the morphisms in E_R^{ADR} that are induced by such path in $R = kQ/J^m$. These will give a basis for the morphisms $Re_i / \text{rad}^l Re_i \rightarrow Re_j / \text{rad}^k Re_j$. A path $p : j \rightarrow i \in kQ/J^m = R$ of length $|p|$ (under the length grading on Q) induces the morphism $p : Re_i \rightarrow Re_j$ which composes to give a non-zero morphism $Re_i \rightarrow Re_j / \text{rad}^k Re_j$ if and only if $|p| < k$. In turn, this descends to give a non-zero morphism $Re_i / \text{rad}^l Re_i \rightarrow Re_j / \text{rad}^k Re_j$ if and only if $\text{rad}^l Re_i \subset \ker(p) = \text{rad}^{k-|p|} Re_i$, which occurs if and only if $l \geq k - |p|$. As such there are isomorphisms of vector spaces

$$\begin{aligned} e_{jk} E_R^{\text{ADR}} e_{il} &= \left\langle \begin{array}{c} \text{elements of } \text{Hom}_R(Re_i, Re_j) \text{ that factor through} \\ \text{Hom}_R(Re_i / \text{rad}^l Re_i, Re_j / \text{rad}^k Re_j) \end{array} \right\rangle \\ &= \left\langle \begin{array}{c} \text{paths } p \in e_i Re_j \\ \text{such that } k - l \leq |p| < k \end{array} \right\rangle. \end{aligned}$$

We then calculate the dimension of $N_m(Q)$ by counting the number of paths between any two vertices. Using the explicit description of $N_m(Q)$ above, any path in $N_m(Q)$ corresponds to the composition of arrows of type a_l and arrows of type $b(i_t)$, these commute $b(t(a)_1)a_{l+1} = a_l b(h(a)_{l-1})$, and $b(t(a)_1)a_2 = 0$. Using these relations any non-zero path can be rearranged such that all the $b(i_t)$ type arrows occur in the path before the a_l type arrows. That is: a path from i_l to j_k in $N_m(Q)$ exactly corresponds to the path $(a_1)_k \cdots (a_n)_{k-|p|+1}$ in $N_m(Q)$ induced by a path $p = a_{|p|} \cdots a_1$ from j to i in Q of length $|p|$ pre-composed with $l - k + |p|$ arrows of $b(i_t)$ type

$$(a_1)_k \cdots (a_n)_{k-|p|+1} b(i_{k-|p|}) \cdots b(i_{l-1}) : i_l \rightarrow j_k$$

so that the induced path is from i_l to j_k . However, the path is non-zero if and only if the number of type $b(i_t)$ arrows is greater than or equal to 0 and strictly less than l , and it follows that

$$e_{jk} N_m(Q) e_{il} = \left\langle \begin{array}{l} \text{paths } p \in e_i R e_j \\ \text{such that } 0 \leq l - k + |p| < l \end{array} \right\rangle.$$

Hence

$$\dim e_{jk} N_m(Q) e_{il} = \dim \{p \in e_i R e_j \mid k - l \leq |p| < k\} = \dim e_{jk} E_R^{\text{ADR}} e_{il}.$$

It follows that the surjective homomorphism $\Phi: N_m(Q) \rightarrow E_R^{\text{ADR}}$ is in fact an isomorphism as $\dim N_m(Q) = \dim E_R^{\text{ADR}}$. Hence $E_R^{\text{ADR}} \cong N_m(Q)$.

Further, under this isomorphism the layer functions defining the quasi-hereditary structures on $N_m(Q)$ and E_R^{ADR} are identified and hence this is an isomorphism of quasi-hereditary algebras. \square

Example 6.16 We give a brief example of Proposition 6.15. Consider the quiver

$$Q := \begin{array}{ccc} & \circlearrowleft x & \\ & 1 & \xleftarrow{a} 2 \end{array}$$

and let J denote the two-sided ideal generated by all arrows. Define $R := \mathbb{C}Q/J^3$, and then we present the two algebras $N_3(Q)$ and E_R^{ADR} .

Firstly, the algebra $N_3(Q)$ is defined to be the path algebra of the quiver with relations

$$Q^{(3)} := \begin{array}{ccc} 1_3 & & 2_3 \\ b(1_2) \downarrow & \nearrow x_3 & \downarrow b(2_2) \\ 1_2 & & 2_2 \\ b(1_1) \downarrow & \nearrow x_2 & \downarrow b(2_1) \\ 1_1 & & 2_1 \end{array} \quad \begin{array}{l} b(1_1)x_2 = 0, \\ b(2_1)a_2 = 0, \\ b(1_2)x_3 = x_2b(1_1), \text{ and} \\ b(2_2)a_3 = a_2b(1_1). \end{array}$$

Secondly, we consider the indecomposable modules in $\text{adr}(R)$. There are six classes and we list them and a basis for all injective or surjective maps between them below.

$$\begin{array}{ll} 1_3 := Re_1 = \langle e_1, x, x^2 \rangle, & 2_3 := Re_2 = \langle e_2, a, xa \rangle, \\ 1_2 := Re_1/\text{rad}^2 Re_1 = \langle e_1, x \rangle, & 2_2 := Re_2/\text{rad}^2 Re_2 = \langle e_2, a \rangle, \\ 1_1 := Re_1/\text{rad}^1 Re_1 = \langle e_1 \rangle, & 2_1 := Re_2/\text{rad}^1 Re_2 = \langle e_2 \rangle. \end{array} \quad \begin{array}{ccc} 1_3 & & 2_3 \\ \downarrow x & \nearrow a & \downarrow \\ 1_2 & & 2_2 \\ \downarrow x & \nearrow a & \downarrow \\ 1_1 & & 2_1 \end{array}$$

This describes $E_R^{\text{ADR}} := \text{End}_R(\text{ADR})$ and matches the path algebra with relations description of $N_3(Q)$ above.

Combining Proposition 6.15 with Proposition 6.12 and Theorem 5.1 (or Theorem 6.11) instantly gives the following corollary.

Corollary 6.17 *If Q is a finite quiver without sinks or sources and m is a positive integer, then there are isomorphisms of quasi-hereditary algebras*

$$E_{kQ/J^m} \cong E_{kQ/J^m}^{\text{ADR}} \cong N_m(Q).$$

In particular, the Ringel dual for a nilpotent quiver algebra without sinks or sources is determined by the formula

$$\mathfrak{R}(N_m(Q)) \cong N_m(Q^{\text{op}})^{\text{op}}.$$

We note that if Q is a finite quiver with no sinks but with sources then $E_{kQ/J^m}^{\text{ADR}} \cong N_m(Q)$ but $E_{kQ^{\text{op}}/J^m}^{\text{ADR}} \not\cong N_m(Q^{\text{op}})^{\text{op}}$ (if $m > 1$) as kQ^{op} contains sinks. In particular, Proposition 6.15 and Theorem 6.11 cannot be used to strengthen the Ringel duality formula of Corollary 6.17 to all quivers with no sources.

6.5 Auslander and Nakayama algebras

For a finite dimensional algebra R of finite representation type we define $\text{AUS} := \bigoplus_{M \in \text{ind}(R\text{-mod})} M$, where the sum is taken over all indecomposable $M \in R\text{-mod}$ up to isomorphism, and the Auslander algebra is defined to be

$$E_R^{\text{AUS}} := \text{End}_R(\text{AUS}).$$

Proposition 6.18 *If R is an ideally ordered monomial algebra, then $E_R^{\text{AUS}} \cong E_R$ if and only if R is self-injective.*

Proof If $E_R^{\text{AUS}} \cong E_R$ then $R\text{-mod} \cong \text{sub}(R)$, hence every injective R -module I embeds into R^n . Therefore, I is a direct summand of R^n , hence projective, and hence R is self-injective.

Conversely, if R is self-injective, then every injective R -module embeds into R^n for some n and hence every injective module is also a projective module. Then every object in $R\text{-mod}$ is a submodule of an injective R -module, hence of a projective R -module, hence $R\text{-mod} \cong \text{sub}(R)$ and $E_R^{\text{AUS}} \cong E_R$. □

The Nakayama algebras, introduced in [31], are a well known class of finite dimensional algebras with finite representation type; see e.g. [4, Theorem VI.2.1]. Recall that a self-injective Nakayama algebra is of the form kC_n/J^m where C_n is an oriented cycle with n vertices and J is the ideal generated by all arrows, see e.g. [1, Theorem 32.4] for the a description of the underlying quiver of a general Nakayama algebra. In particular, the self-injective Nakayama algebras are ideally ordered monomial algebras.

Corollary 6.19 *If R is a self-injective Nakayama algebra, then $E_R^{\text{AUS}} \cong E_R$.*

It follows from the explicit description $R = kC_n/J^m$ that $E_R = N_m(C_n)$ by Corollary 6.17 and so this corollary recovers the well known explicit description of the Auslander

algebras of self-injective Nakayama algebra E_R in terms of quivers with relations. As $(kC_n/J^m)^{op} \cong kC_n/J^m$ and $N_m(C_n) \cong N_m(C_n^{op})^{op}$ the Ringel duality formula recovers the result of [37] that the Auslander algebras of self-injective Nakayama algebras are Ringel self-dual for the ideal layer function.

Corollary 6.20 *For a self-injective Nakayama algebra R , $\mathfrak{R}(E_R) \cong E_R$.*

Remark 6.21 In order to give another perspective on Proposition 6.18, and Corollaries 6.19 and 6.20, we recall that self-injective finite dimensional monomial algebras R are Nakayama algebras. To see this, we have to show that the quiver Q underlying R is a union of oriented lines and oriented cycles. In other words, at every vertex of Q there is at most one incoming and at most one outgoing arrow. Assume that there is a vertex i with more than one outgoing arrow. Then, as R is monomial, the corresponding indecomposable projective R -module P_i does not have a simple socle - in particular, P_i is not injective contradicting our assumption that R is self-injective. A dual argument shows that Q does not have vertices with more than one incoming arrow.

7 Appendix: Results on finite dimensional monomial algebras

In this section we collect some technical results on finite dimensional monomial algebras $R = kQ/I$ (where I is generated by a collection of paths in Q). We will use the term ‘monomial’ to mean a monomial expression in the generators (i.e. arrows and lazy paths) of such an algebra.

Lemma 7.1 *Let R be a monomial algebra and $n, m \in R$ monomials. If there exists a surjection $\phi: Rm \rightarrow Rn$, then the map $Rm \rightarrow Rn$ defined by $m \mapsto n$ is R -linear.*

Proof It suffices to show that $\text{ann}_R(m)$ is contained in $\text{ann}_R(n)$. Take $r \in R$ with $rm = 0$, and we aim to show that $rn = 0$. We write $r = \sum \lambda_i r_i$ with monomials r_i and non-zero scalars λ_i . Since R is monomial, it follows that $r_i m = 0$ for all i . The existence of a surjection $\phi: Rm \rightarrow Rn$ implies $m, n \in eR$ for some primitive idempotent $e \in R$ and that there exist $s, t \in R$ such that $\phi(tm) = n$ and $\phi(m) = sn$. In particular, $tsn = n$ and so $s = \mu_0 e + \sum_{i=1}^t \mu_i s_i$ for some non-zero scalars μ_i and distinct monomials $s_i \neq e$. Therefore $r_i sn = \phi(r_i m) = \phi(0) = 0$, and so as R is monomial it follows that all monomials that make up $r_i sn$ are 0. In particular, $r_i \mu_0 en = \mu_0 r_i n = 0$. This implies that $r_i n = 0$ for all i , and hence $rn = 0$ so $\text{ann}_R(m) \subset \text{ann}_R(n)$ finishing the proof. □

Lemma 7.2 *Let $m, n \in R$ be monomials. If R is ideally ordered, then every surjection $Rm \rightarrow Rn$ factors over $\pi: Rm \rightarrow Rn, m \mapsto n$.*

Proof Let $\psi: Rm \rightarrow Rn$ be a surjection. In particular, $m, n \in eR$ for some primitive idempotent $e \in R$ and there exist $s, t \in R$ such that $\psi(m) = sn$ and $\psi(tm) = n$. It follows that $tsn = n$, so $s = \lambda_0 e + \sum \lambda_i s_i \in eRe$ for non-zero scalars λ_i and distinct monomials $s_i \neq e$. Hence $sn = \lambda_0 n + \sum \lambda_i s_i n$. In particular, $Rs_i n \subsetneq Rn$, and since R is ideally ordered there exists surjections $Rn \rightarrow Rs_i n$ which, using Lemma 7.1,

we can assume are defined by $n \mapsto s_i n$. Denote the composition of such a surjection with the inclusion $Rs_i n \subseteq Rn$ by φ_i and define $\varphi: Rn \rightarrow Rn$ as $\varphi = \lambda_0 \text{id} + \sum \lambda_i \varphi_i$. Then $\varphi(n) = sn$ and therefore $\psi = \varphi\pi$ factors as claimed. \square

Lemma 7.3 *Let $p \in eR$ for a primitive idempotent $e \in R$. If R is ideally ordered, then the principal left ideal Rp is isomorphic to a principal ideal Rm , for a monomial $m \in eR$.*

Proof Since R is monomial, we may write p as linear combination of monomials $p = \sum_{i=1}^t \lambda_i p_i$ with λ_i non-zero scalars and $p_i \in eR$ monomials. Since R is ideally ordered we may assume that the p_i are labelled in such a way that $Rp_1 \rightarrow Rp_2 \rightarrow \dots \rightarrow Rp_t$ are surjections.

We now wish to rewrite p so that none of the p_i can be expressed in the form np_1 for a monomial n . To do this, let I index the p_i such that there is a monomial r_i with $p_i = r_i p_1$ for $i \in I$. Then we define $s = \lambda_1 e + \sum_{i \in I} \lambda_i r_i$ and $p = sp_1 + \sum_{i \notin I} \lambda_i p_i$. As $r := \sum_{i \in I} \lambda_i r_i \in \text{rad } R \cap eRe$ it follows that $s = \lambda_1 e + r$ is a unit in eRe and there exists $t \in eRe$ such that $st = e$. In particular, $Rtp = Rp$. Then we rewrite $tp = tsp_1 + \sum_{i \notin I} \lambda_i t p_i = p_1 + \sum_{j=2}^{t'} \mu_j q_j$ for some non-zero scalars μ_j and monomials $q_j \neq e$. For each q_j there is some p_i such that $Rq_j \subset Rp_i$ by their definition, and hence there are surjections $Rp_1 \rightarrow Rq_j$ for all j . As $Rtp \cong Rp$ we now work with tp rather than p and tp has the property that there are no q_j with $np_1 = q_j$ for a monomial n .

We claim that $Rtp \cong Rp_1$, hence $Rp \cong Rp_1$. As there are surjections $Rp_1 \rightarrow Rq_j$ there are surjections $Rp_1 \rightarrow Rq_j, p_1 \mapsto q_j$ by Lemma 7.1. Let φ_j be the composition of such a surjection with the canonical inclusion $Rq_j \rightarrow R$ and let $\iota: Rp_1 \rightarrow R$ be the canonical inclusion. Define $\psi: Rp_1 \rightarrow R$ by $\psi = \iota + \sum_{j=2}^{t'} \mu_j \varphi_j$. Then $\psi(p_1) = p_1 + \sum_{j=2}^{t'} \mu_j q_j = tp$ so $\text{im } \psi = Rtp$. Hence ψ defines a surjective morphism $\phi: Rp_1 \rightarrow Rtp$.

We must now check that this morphism is also injective. If $\psi(rp_1) = 0$, then $rp_1 + r \sum_{j=2}^{t'} \mu_j q_j = 0$. As R is monomial if rp_1 is non-zero there must exist monomials $n, m \in R$ such that $np_1 = mq_j$ for some j , and if this occurs either $p_1 = m'q_j$ or $n'p_1 = q_j$ for submonomials m' and n' neither equal e . The first case cannot occur as this implies $Rp_1 \subsetneq Rq_j$ which contradicts the existence of a surjection $Rp_1 \rightarrow Rq_j$. The second situation also cannot occur as the construction of the q_j above ensured none were of this form. Hence $rp_1 = 0$ so the morphism is also injective and $Rp_1 \cong Rtp \cong Rp$. \square

Acknowledgements We thank Teresa Conde for interesting discussions about this work and about relations to her thesis. We are grateful to Karin Erdmann for pointing us to Ringel’s paper which simplifies the proof of our main result and adds another perspective to this work. We also thank Agnieszka Bodzenta who, in particular, explained to us the proof of Proposition 6.2 and Xiao-Wu Chen who shared with us Example 5.2. We would also like to thank Ögmundur Eiriksson, Julian Külshammer, Daiva Pučinskaitė, Špela Špenko, and Michael Wemyss for interesting and helpful discussions and David Ploog for pointing out misprints in an earlier version. We would also particularly like to thank the anonymous referee, whose many useful comments have greatly improved the paper.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution,

and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

1. Anderson, F.W., Fuller, K.R.: Rings and Categories of Modules. Graduate Texts in Mathematics, vol. 13, 2nd edn. Springer, New York (1992)
2. Auslander, M., Buchweitz, R.-O.: The homological theory of maximal Cohen–Macaulay approximations. *Mém. Soc. Math. France (N.S.)* **38**, 5–37 (1989)
3. Auslander, M., Reiten, I.: Applications of contravariantly finite subcategories. *Adv. Math.* **86**(1), 111–152 (1991)
4. Auslander, M., Reiten, I., Smalø, S.O.: Representation Theory of Artin Algebras. Cambridge Studies in Advanced Mathematics, vol. 36. Cambridge University Press, Cambridge (1997). Corrected reprint of the 1995 original
5. Auslander, M., Smalø, S.O.: Preprojective modules over Artin algebras. *J. Algebra* **66**(1), 61–122 (1980)
6. Baur, K., Erdmann, K., Parker, A.: Δ -filtered modules and nilpotent orbits of a parabolic subgroup in O_N . *J. Pure Appl. Algebra* **215**(5), 885–901 (2011)
7. Bodzenta, A., Bondal, A.: Derived categories of smooth surface contractions (in preparation)
8. Bodzenta, A., Külshammer, J.: Ringel duality as an instance of Koszul duality (2017). [arXiv:1701.06222](https://arxiv.org/abs/1701.06222)
9. Buchweitz, R.-O., Leuschke, G.J., Van den Bergh, M.: On the derived category of Grassmannians in arbitrary characteristic. *Compositio Math.* **151**(7), 1242–1264 (2015)
10. Cline, E., Parshall, B., Scott, L.: Algebraic stratification in representation categories. *J. Algebra* **117**(2), 504–521 (1988)
11. Conde, T.: The quasihereditary structure of the Auslander–Dlab–Ringel algebra. *J. Algebra* **460**, 181–202 (2016)
12. Conde, T.: Δ -filtrations and projective resolutions for the Auslander–Dlab–Ringel algebra. *Algebr. Represent. Theory* (2017). <https://doi.org/10.1007/s10468-017-9730-z>
13. Conde, T., Erdmann, K.: The Ringel dual of the Auslander–Dlab–Ringel algebra (2017). [arXiv:1708.05766](https://arxiv.org/abs/1708.05766)
14. Coulembier, K.: Ringel duality and Auslander–Dlab–Ringel algebras (2017). [arXiv:1710.08019](https://arxiv.org/abs/1710.08019)
15. Coulembier, K., Mazorchuk, V.: Dualities and derived equivalences for category \mathcal{O} . *Israel J. Math.* **219**(2), 661–706 (2017)
16. Crawley-Boevey, W., Sauter, J.: On quiver Grassmannians and orbit closures for representation-finite algebras. *Math. Z.* **285**(1–2), 367–395 (2017)
17. Dlab, V., Ringel, C.M.: Every semiprimary ring is the endomorphism ring of a projective module over a quasihereditary ring. *Proc. Amer. Math. Soc.* **107**(1), 1–5 (1989)
18. Dlab, V., Ringel, C.M.: The module theoretical approach to quasi-hereditary algebras. In: Tachikawa, H., Brenner, S. (eds.) *Representations of Algebras and Related Topics*. London Mathematical Society Lecture Note Series, vol. 168, pp. 200–224. Cambridge University Press, Cambridge (1992)
19. Erdmann, K., Parker, A.E.: On the global and ∇ -filtration dimensions of quasi-hereditary algebras. *J. Pure Appl. Algebra* **194**(1–2), 95–111 (2004)
20. Eiríksson, Ö., Sauter, J.: Quiver-graded Richardson orbits (2017). [arXiv:1707.03244](https://arxiv.org/abs/1707.03244)
21. Geigle, W., Lenzing, H.: Perpendicular categories with applications to representations and sheaves. *J. Algebra* **144**(2), 273–343 (1991)
22. Ginzburg, V., Guay, N., Opdam, E., Rouquier, R.: On the category \mathcal{O} for rational Cherednik algebras. *Invent. Math.* **154**(3), 617–651 (2003)
23. Hille, L., Perling, M.: Tilting bundles on rational surfaces and quasi-hereditary algebras. *Ann. Inst. Fourier (Grenoble)* **64**(2), 625–644 (2014)
24. Hille, L., Ploog, D.: Tilting chains of negative curves on rational surfaces. *Nagoya Math. J.* (2017). <https://doi.org/10.1017/nmj.2017.40>
25. Iyama, O.: Finiteness of representation dimension. *Proc. Amer. Math. Soc.* **131**(4), 1011–1014 (2003)
26. Iyama, O., Reiten, I.: 2-Auslander algebras associated with reduced words in Coxeter groups. *Int. Math. Res. Not. IMRN* **2011**(8), 1782–1803 (2011)

27. Kalck, M., Karmazyn, J.: Noncommutative Knörrer type equivalences via noncommutative resolutions of singularities (2017). [arXiv:1707.02836](https://arxiv.org/abs/1707.02836)
28. Krause, H.: Koszul, Ringel and Serre duality for strict polynomial functors. *Compositio Math.* **149**(6), 996–1018 (2013)
29. Krause, H., Saorín, M.: On minimal approximations of modules. In: Green, E.L., Huisgen-Zimmermann, B. (eds.) *Trends in the Representation Theory of Finite-Dimensional Algebras. Contemporary Mathematics*, vol. 229, pp. 227–236. American Mathematical Society, Providence (1998)
30. Kuznetsov, A., Lunts, V.A.: Categorical resolutions of irrational singularities. *Int. Math. Res. Not. IMRN* **2015**(13), 4536–4625 (2015)
31. Nakayama, T.: Note on uni-serial and generalized uni-serial rings. *Proc. Imp. Acad. Tokyo* **16**, 285–289 (1940)
32. Nguyen, V.C., Reiten, I., Todorov, G., Zhu, S.: Dominant dimension and tilting modules (2017). [arXiv:1706.00475](https://arxiv.org/abs/1706.00475)
33. Pučinskaitė, D.: Quasi-hereditary algebras via generator-cogenerators of local self-injective algebras and transfer of Ringel duality. *Math. Z.* **279**(3–4), 641–668 (2015)
34. Ringel, C.M.: The category of modules with good filtrations over a quasi-hereditary algebra has almost split sequences. *Math. Z.* **208**(2), 209–223 (1991)
35. Ringel, C.M.: Iyama’s finiteness theorem via strongly quasi-hereditary algebras. *J. Pure Appl. Algebra* **214**(9), 1687–1692 (2010)
36. Ringel, C.M.: On the representation dimension of Artin algebras. *Bull. Inst. Math. Acad. Sin. (N.S.)* **7**(1), 33–70 (2012)
37. Tan, R.: Auslander algebras of self-injective Nakayama algebras. *Pure Mathematical Sciences* **2**(2), 89–108 (2013)
38. Tsukamoto, M.: Strongly quasi-hereditary algebras and rejective subcategories. *Nagoya Math. J.* (2018). <https://doi.org/10.1017/nmj.2018.9>