

 Open access • Journal Article • DOI:10.1097/00001573-200111000-00005

Risk and benefit of treatment of isolated systolic hypertension in the elderly: evidence from the Systolic Hypertension in Europe Trial — [Source link](#)

Hilde Celis, Robert Fagard, Jan A. Staessen, Lutgarde Thijs

Institutions: Katholieke Universiteit Leuven

Published on: 01 Nov 2001 - Current Opinion in Cardiology (Curr Opin Cardiol)

Topics: Systolic hypertension, Ambulatory blood pressure, Blood pressure, Pulse pressure and Nitrendipine

Related papers:

- [Treatment of isolated systolic hypertension in the elderly: evidence from three clinical trials](#)
- [Overview of the outcome trials in older patients with isolated systolic hypertension](#)
- [Subgroup and Per-Protocol Analysis of the Randomized European Trial on Isolated Systolic Hypertension in the Elderly](#)
- [Effects of hypertension control on stroke incidence and fatality: report from Syst.-China and post-stroke antihypertensive treatment.](#)
- [The role of diastolic blood pressure when treating isolated systolic hypertension.](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/risk-and-benefit-of-treatment-of-isolated-systolic-1bijmkqq6t>

Risk and benefit of treatment of isolated systolic hypertension in the elderly: evidence from the Systolic Hypertension in Europe Trial

Hilde Celis, MD, Robert H. Fagard, MD, PhD, Jan A. Staessen, MD, PhD, and Lutgarde Thijs, BSc

The Syst-Eur trial investigated whether active treatment starting with the dihydropyridine calcium channel blocker (CCB) nitrendipine, could reduce the cardiovascular complications of isolated systolic hypertension (ISH) in the elderly. The intention-to-treat analysis showed that active treatment improved outcome. The per-protocol analysis largely confirmed these results. The effect of treatment on total and cardiovascular mortality might be attenuated in very old patients. Further analysis also suggested benefit in those patients who remained on nitrendipine monotherapy. Active treatment was more beneficial in patients with diabetes as compared with those without diabetes at entry and reduced the incidence of dementia by 50%. Analyses of data from the Ambulatory Blood Pressure Monitoring (ABPM) Side Project suggested that most of the benefit of treatment was seen in patients with a daytime systolic BP ≥ 160 mm Hg. Finally, a meta-analysis partly based on Syst-Eur data showed that in older hypertensive patients pulse pressure and not mean pressure is the major determinant of cardiovascular risk. *Curr Opin Cardiol* 2001, 16:342–348 © 2001 Lippincott Williams & Wilkins, Inc.

This review was written for the Systolic Hypertension in Europe Trial Investigators, Hypertension and Cardiovascular Rehabilitation Unit, Department of Molecular and Cardiovascular Research, Faculty of Medicine, University of Leuven K.U. Leuven, Leuven, Belgium. A complete list of the investigators appears in reference [1].

Correspondence to Hilde Celis, MD, Studietoördinatiecentrum, Laboratorium Hypertensie, Campus Gasthuisberg, Herestraat 49, B-3000 Leuven, Belgium; E-mail: hilde.celis@med.kuleuven.ac.be

Current Opinion in Cardiology 2001, 16:342–348

The presence of isolated systolic hypertension (ISH) rises curvilinearly with age. Among septagenarians it averages 8% and beyond 80 years it rises to more than 25% [2]. Isolated systolic hypertension thus affects a large proportion of the elderly population. Against this background, the European Working Party on High Blood Pressure in the Elderly started the placebo-controlled double-blind Syst-Eur (Systolic Hypertension in Europe) trial [3]. In 1991 the Systolic Hypertension in the Elderly (SHEP) trial showed that diuretic-based treatment prevented stroke, myocardial infarction, and congestive heart failure [4]. However, because of the remaining uncertainties about the treatment of ISH in the elderly [5], the Syst-Eur trial continued after the publication of the SHEP results [4]. Furthermore, the controversy on the role of calcium channel blockers (CCBs) as first-line antihypertensive agents [6–9], highlighted the lack of evidence regarding the reduction of cardiovascular risk by these agents. The Syst-Eur trial stopped on February 14th, 1997, after the second of four planned interim analyses, because the primary endpoint of a significant benefit for stroke was reached [3].

Patients and methods

The protocol of the Syst-Eur trial, described in detail elsewhere [3], was approved by the Ethics Committees of all participating centers. Eligible patients had to be 60 years or older and have a sitting systolic BP of 160 to 219 mm Hg with sitting diastolic BP below 95 mm Hg during a run-in phase on single-blind placebo. After stratification by center, sex, and previous cardiovascular complications, patients were randomly assigned double-blind treatment with active treatment or placebo. Active treatment was initiated with nitrendipine (10–40 mg per day), if necessary combined or replaced by enalapril (5–20 mg per day) or hydrochlorothiazide (12.5–25 mg per day) or both drugs. In the control group matching placebos were used. The study medications were stepwise titrated and combined to reduce the systolic BP by 20 mm Hg or more to below 150 mm Hg. Patients withdrawing from double-blind treatment remained in open follow-up.

Overview of previously published Syst-Eur results

Of 4695 randomized patients, 2398 were randomly assigned to active treatment [1,10•,11,12,13••,14,15••,16•].

In the intention-to-treat analysis [1], the between-group difference in BP was 10.1 mm Hg systolic and 4.5 mm Hg diastolic at the median follow-up of 2.0 years. Cardiovascular mortality was slightly lower on active treatment (-26% , $P = 0.08$), but all-cause mortality was not significantly changed (-13% , $P = 0.28$) [10]. Active treatment reduced the incidence of total stroke by 42% ($P = 0.002$) and of nonfatal stroke by 45% ($P = 0.004$) [10]. All fatal and nonfatal cardiac endpoints decreased by 25% ($P = 0.03$) and all fatal and nonfatal cardiovascular endpoints by 30% ($P < 0.001$) [10]. The incidence of fatal and nonfatal cancer (-12% , $P = 0.42$) and of bleeding (excluding cerebral and retinal hemorrhages; -9% , $P = 0.75$) was not different in the two treatment groups [10]. In terms of absolute benefit, treating 1000 elderly patients with ISH for 5 years could prevent 29 strokes or 53 major cardiovascular events.


Subgroup analysis [11] showed that male sex, previous cardiovascular complications, older age, higher systolic BP, and smoking at randomization were positively and independently correlated with cardiovascular risk. The relative benefit of active treatment was different according to sex or to the presence of cardiovascular complications at entry. Furthermore, for total ($P = 0.009$) and cardiovascular ($P = 0.09$) mortality, the benefit of antihypertensive treatment weakened with advancing age (Fig. 1). For total mortality ($P = 0.05$), it increased with higher systolic BP at entry (Fig. 1) and for fatal and nonfatal stroke ($P = 0.01$), it was most evident in non-smokers (92.5% of all patients).

The results of the intention-to-treat [1,10] and per-protocol [11] (including only the endpoints occurring during the double-blind phase) analysis were largely similar. Active treatment reduced total mortality by 26% ($P = 0.05$); similar but nonsignificant trends were observed for cardiac and cerebrovascular mortality. Cardiovascular, cardiac, and cerebrovascular events declined by respectively, 32% ($P < 0.001$), 26% ($P = 0.05$), and 44% ($P = 0.004$). In terms of absolute benefit, the per-protocol analysis suggested that treating 1000 patients for 5 years would prevent 24 deaths, 29 strokes, 25 cardiac endpoints, or 54 major cardiovascular events.

The relative benefits of antihypertensive treatment in the Syst-Eur trial were largely similar to those of other trials in older patients with combined systolic and diastolic hypertension [17–22] or with isolated systolic hypertension [4,23,24].

In view of the persistent controversy about the possible adverse effects of CCBs [8,9,25–28], the question was raised whether treatment with nitrendipine alone would also influence outcome. Further analyses [12] also suggested similar benefit in the patients who remained on nitrendipine monotherapy. In these analyses, 1327 patients who remained on single nitrendipine treatment

Figure 1. Risk according to age and initial systolic blood pressure



Adjusted relative hazard rates of total and cardiovascular mortality according to age and initial systolic blood pressure. The hazard rates (placebo/active treatment), calculated by intention-to-treat, are presented as continuous risk functions with 95% confidence intervals. P values refer to the interaction terms between treatment and independent predictor variable. Published with permission [11].

throughout the whole trial, were matched by sex, age, previous cardiovascular complications, and systolic BP at entry, with an equal number of placebo patients from the control group. Compared with this matched control group, treatment with nitrendipine reduced cardiovascular mortality by 41% ($P = 0.05$), all cardiovascular endpoints by 33% ($P = 0.01$), fatal and nonfatal cardiac endpoints by 33% ($P = 0.05$) and fatal and nonfatal heart failure by 48% ($P = 0.05$).

These data from the Syst-Eur trial invalidate circumstantial evidence based on case-controlled and observational studies [6–9,29–33], which according to the investigators' interpretation left a margin of uncertainty regarding the occurrence of potentially dangerous side effects. Several studies investigated the effects of dihydropyridine CCBs in Chinese hypertensive patients [23,24,34–36]. Although some of these trials used unorthodox designs [34,35], they also demonstrated a positive influence on outcome.

At randomization, 492 patients (10.5%) had diabetes mellitus [13]. After adjustment for possible confounders, active treatment reduced all-cause mortality by 55%, car-

diovascular mortality by 76%, all cardiovascular endpoints by 69%, fatal and nonfatal stroke by 73% and all cardiac endpoints by 63% in the group of diabetic patients. In the nondiabetic patients, all cardiovascular endpoints were reduced by 26% and all fatal and nonfatal strokes by 38%. On active treatment, the reductions in total mortality ($P = 0.04$), cardiovascular mortality ($P = 0.02$) and all cardiovascular endpoints ($P = 0.01$) were significantly larger in diabetic patients than in nondiabetic patients.

These data from the Syst-Eur trial [13,37] were the first to prove that antihypertensive drug treatment starting with a dihydropyridine CCB was particularly beneficial in diabetic patients. These results contradicted the suggestion that some (second-generation) dihydropyridine CCBs might be harmful, particularly in hypertensive patients with diabetes mellitus [29–33,38,39].

Systolic hypertension is associated with an increased risk of dementia in elderly people. The primary hypothesis of the Syst-Eur Vascular Dementia Substudy, was that a reduction in BP would protect against vascular dementia [14]. Two thousand four hundred and eighteen patients were enrolled [15]. At the median follow-up of 2.0 years, active treatment reduced the incidence of dementia by 50% ($P = 0.05$) from 7.7 to 3.8 cases per 1000 patient-years. Active treatment prevented mainly Alzheimer's dementia (8 versus 15 cases), but also vascular (0 versus 2) and mixed (3 versus 4) dementia. According to the intention-to-treat analysis, treating 1000 elderly patients with ISH for 5 years might prevent 19 cases of dementia.

The prevention of degenerative dementia was somewhat unexpected, although recent studies indicate that vascular factors, in particular hypertension, might also play a role in the development of degenerative dementia [40]. An alternative explanation, although still unproven, could be that lipophilic CCBs, which cross the blood-brain barrier and bind to brain receptors located in areas affected by Alzheimer's disease, might confer a specific neuroprotective effect [41,42].

Overview of recent Syst-Eur results **Response to antihypertensive therapy in patients with sustained and nonsustained isolated systolic hypertension (ISH)**

Patients with nonsustained hypertension (also called white coat or isolated clinic hypertension) have normal ambulatory blood pressure (ABP), but elevated clinic blood pressure (CBP) [43–45]. Target organ damage [46–50] and cardiovascular risk [49,51,52] is reported to be lower in patients with nonsustained as compared with patients with sustained (ABP and CBP both elevated) hypertension, but data on nonsustained hypertension in elderly patients with ISH are generally lacking.


Therefore, the data from the Ambulatory Blood Pressure Monitoring (ABPM) Substudy [53,54] of the Syst-Eur trial were analyzed. The objectives of this analysis were to evaluate the consequences of nonsustained systolic hypertension as well as the impact of antihypertensive treatment. Patients enrolled in the ABPM Substudy were classified according to baseline daytime systolic ABP in 1 of 3 subgroups: nonsustained ISH (< 140 mm Hg), mild sustained ISH (140 to 159 mm Hg), and moderate sustained ISH (≥ 160 mm Hg).

At baseline, diastolic daytime and systolic and diastolic nighttime and clinic BPs were higher in sustained than in nonsustained hypertensives. Patients with nonsustained ISH had smaller ECG voltages than patients with mild and moderate sustained ISH. The differences in ECG voltages between these three groups remained significant after controlling for systolic CBP, but significance was lost after adjusting for daytime systolic ABP.

During follow-up, active treatment reduced daytime and nighttime ABP significantly in the patients with sustained ISH but not in patients with nonsustained ISH. CBP, however, decreased significantly in the three subgroups. Active treatment reduced ABP and CBP more than placebo in patients with mild and moderate sustained ISH. By contrast in patients with nonsustained ISH, the changes in ABP between the treatment groups were not significantly different (except for daytime SBP), whereas the changes in CBP were also more pronounced on active treatment as compared with placebo.

Patients with nonsustained ISH had a lower incidence of stroke ($P < 0.05$) and of cardiovascular events ($P = 0.01$) during follow-up than patients with sustained ISH. Active treatment significantly reduced ($P < 0.05$) the ECG voltages in patients with sustained ISH, but not in these with nonsustained ISH. The influence of active treatment on the incidence of stroke ($P < 0.05$) and cardiovascular events ($P = 0.06$) (Fig. 2) and on ECG voltages was significantly more beneficial than that of placebo only in patients with moderate sustained ISH.

These results confirm previous studies in other hypertensive populations [49,51,52,55,56], which suggested that white coat hypertension is associated with a better outcome [49,51,52] or that the predictive value of ABP persists after controlling for CBP [55,56]. A previous analysis of data from the ABPM Substudy, in which BP was treated as a continuous variable, showed that the systolic ABP was a significant predictor of cardiovascular complications over and beyond CBP [57]. The night/day ratio had a predictive value independent of the 24 hour ABP [57]. In this analysis diastolic ABP and CBP were not related to outcome [57]. The previously reported findings on surrogate (ECG voltages) and hard (outcome) endpoints [43] allow us to conclude that sustained

Figure 2. Risk according to hypertension status and treatment group

Number of strokes and cardiovascular events per 1000-patient-years and absolute number of events during follow-up in patients with Non-SH (nonsustained hypertension), MiSH (mild sustained hypertension), and MoSH (moderate sustained hypertension), divided according to treatment group (open columns, placebo treatment; filled columns, active treatment). Results are from intention-to-treat analysis. P value refers to comparison of rates between 2 treatment groups within each subgroup according to daytime systolic BP. Published with permission [43].

ISH is more harmful than nonsustained ISH, in particular when the daytime systolic ABP ≥ 160 mm Hg. Moreover, the benefit of antihypertensive therapy is also mainly seen in the latter patients, and becomes less evident when daytime systolic ABP is less than 160 mmHg.


Pulse pressure as cardiovascular risk factor in elderly hypertensives

For the moment, guidelines for the management of hypertension are mainly based on the measurement of systolic and diastolic BP and these are usually considered as being two isolated variables [44,45,58,59]. However, BP is more correctly described as consisting of a pulsatile (pulse pressure [PP]) and a steady (mean pressure [MP]) component [50]. Ventricular ejection, arterial stiffness and the timing of the wave reflections are the major determinants of PP, while MP mainly depends on peripheral vascular resistance and cardiac output. Several observational studies suggested that in elderly people PP might be a better predictor of cardiovascular risk than MP [60–69]. Since PP widens with advancing age [2], the results of outcome trials in older hypertensives might provide an ideal database to test this hypothesis. Therefore the data of three placebo-controlled trials in older patients with hypertension (EWPHE [17], Syst-Eur [1],

Syst-China [23]) were pooled to assess the independent roles of PP and MP as determinants of cardiovascular risk with sufficient statistical power. The meta-analysis was based on individual patient data. Relative hazard rates associated with PP and MP were calculated using Cox regression analysis and adjustments were made for sex, age, previous cardiovascular complications, smoking, and active treatment. Furthermore the relative hazard rates for PP were also adjusted for MP and *vice versa*.

In the overall Cox regression analysis with stratification for the three trials and with adjustment for the previously mentioned covariates, a wider PP at baseline increased the risk of major cardiovascular complications. The increase in risk associated with a 10 mm Hg wider PP, ranged from about 13% for all coronary end points ($P = 0.02$) to nearly 20% for cardiovascular mortality ($P = 0.001$). In a similar analysis, MP could only be identified as a significant predictor of risk, after removal of PP as an explanatory variable from the model. The 2-year probability of a major cardiovascular endpoint increased with higher systolic BP ($P < 0.001$) (Fig. 3). Furthermore, at any given level of systolic BP, the risk also increased with lower diastolic BP ($P = 0.001$). This observation suggests that the wider PP was driving the risk of major complications.

The role of PP as a significant predictor of cardiovascular risk had already been identified by several investigators and in different groups of patients [60,61,63–67,69–71]. Moreover, a recent meta-analysis [45] including data of more than 15 000 patients with ISH from 8 different outcome trials also confirmed the role of PP as a risk

Figure 3. Risk estimates for all cardiovascular endpoints based on three trials

Risk associated with increasing systolic blood pressure at fixed levels of diastolic blood pressure. The 2-year probability of a cardiovascular endpoint was adjusted for active treatment, sex, age, previous cardiovascular complications, and smoking by Cox multiple regression with stratification for trial (EWPHE [17], Syst-Eur [1], Syst-China [23]). Published with permission [44].

factor. Furthermore, the authors also reported that active antihypertensive treatment was particularly beneficial in men, in patients aged 70 years or more, and in those with previous cardiovascular complications or wider PP at baseline. If PP at baseline was 90 mm Hg or greater, 63 patients had to be treated to prevent one cardiovascular death, whereas 119 patients had to be treated if pulse pressure was less than 90 mm Hg.

From the present data it can be concluded that in older hypertensives, PP, but not MP is the major determinant of cardiovascular risk. These findings must however be interpreted with caution. The positive and independent association between PP and the incidence of cardiovascular complications does not automatically imply a causal or reversible relationship. Whether these data in the elderly can be extrapolated to younger or middle-aged patients also remains to be proven. Furthermore, the present findings may also have important clinical implications [72]. They suggest that the prediction of cardiovascular complications might be improved by accounting for both the pulsatile (PP) and the steady (MP) component of BP. This should however be further investigated in randomized clinical trials in which the pulsatile component of BP is differently affected by antihypertensive drug treatment.

Conclusion

From the previous data some important conclusions emerge. First, that stepwise antihypertensive treatment starting with the CCB nitrendipine improves prognosis in elderly patients with ISH and may particularly be indicated in diabetic patients with ISH or in those at risk of dementia. Second, that most of the benefit of treatment is seen in patients with a daytime systolic ABP ≥ 160 mm Hg. And finally, that in elderly hypertensive patients pulse pressure and not mean pressure is the major determinant of cardiovascular risk.

Acknowledgment

The Syst-Eur trial, initiated by Antoon Amery, MD (died on November 2, 1994), was a concerted action of the BIOMED Research Program sponsored by the European Union. The trial was carried out in consultation with the World Health Organization, the International Society of Hypertension, the European Society of Hypertension, and the World Hypertension League. The trial was sponsored by Bayer AG (Wuppertal, Germany). The National Fund for Scientific Research (Brussels, Belgium) provided additional support. Study medication was donated by Bayer AG and Merck Sharp and Dohme Inc (West Point, PA, USA).

References and recommended reading

Papers of particular interest, published within the annual period of review, have been highlighted as:

- Of special interest
- Of outstanding interest

- 1 Staessen JA, Fagard R, Thijs L, et al., for the Systolic Hypertension in Europe (Syst-Eur) Trial Investigators: Randomised double-blind comparison of placebo and active treatment for older patients with isolated systolic hypertension. *Lancet* 1997, 350:757–764.
- 2 Staessen JA, Amery A, Fagard R: Editorial review. Isolated systolic hypertension in the elderly. *J Hypertens* 1990, 8:393–405.
- 3 Amery A, Birkenhager W, Bulpitt CJ, et al.: Syst-Eur. A multicentre trial on the

treatment of isolated systolic hypertension in the elderly: objectives, protocol, and organization. *Aging* 1991, 3:287–302.

- 4 SHEP Cooperative Research Group: Prevention of stroke by antihypertensive drug treatment in older persons with isolated systolic hypertension. Final results of the Systolic Hypertension in the Elderly Program (SHEP). *JAMA* 1991, 265:3255–3264.
- 5 Staessen JA, Fagard R, Amery A: Isolated systolic hypertension in the elderly: implications of SHEP for clinical practice and for the ongoing trials. *J Hum Hypertens* 1991, 5:469–474.
- 6 Pahor M, Guralnik JM, Salive ME, et al.: Do calcium channel blockers increase the risk of cancer? *Am J Hypertens* 1996, 9:695–699.
- 7 Pahor M, Guralnik JM, Furberg CD, et al.: Risk of gastrointestinal haemorrhage with calcium antagonists in hypertensive persons over 67 years old. *Lancet* 1996, 347:1061–1065.
- 8 Furberg CD, Psaty BM, Meyer JV: Nifedipine - dose-related increase in mortality in patients with coronary heart disease. *Circulation* 1995, 92:1326–1331.
- 9 Psaty BM, Heckbert SR, Koepsell TD, et al.: The risk of myocardial infarction associated with antihypertensive drug therapies. *JAMA* 1995, 274 :620–625.
- 10 Staessen JA, Thijs L, Birkenhager W, et al., on behalf of the Syst-Eur investigators: Update on the Systolic Hypertension in Europe (Syst-Eur) Trial. *Hypertension* 1999, 33:1476–1477.
- This short paper provides updated outcome results based on the final Syst-Eur database. Although a slightly greater number of endpoints was available, the conclusions of the initial Syst-Eur report were not affected.
- 11 Staessen JA, Fagard R, Thijs L, et al., for the Systolic Hypertension in Europe Trial Investigators: Subgroup and per-protocol analysis of the randomized European trial on isolated systolic hypertension in the elderly. *Arch Intern Med* 1998, 158:1681–1691.
- 12 Staessen JA, Thijs L, Fagard R, et al., for the Systolic Hypertension in Europe (Syst-Eur) Trial Investigators: Calcium channel blockade and cardiovascular prognosis in the European Trial on Isolated Systolic Hypertension. *Hypertension* 1998, 32:410–416.
- 13 Tuomilehto J, Rastenyte D, Birkenhager W, et al.: Effects of calcium-channel blockade in older patients with diabetes and systolic hypertension. *N Engl J Med* 1999, 340:677–684.
- Ten percent of all 4695 Syst-Eur patients had diabetes mellitus. This post-hoc analysis compared outcome in the diabetic and nondiabetic patients. It demonstrated that nitrendipine-based antihypertensive treatment is particularly beneficial in older diabetic patients with ISH and refuted the hypothesis that the use of CCB might be harmful in diabetic patients.
- 14 Forette F, Amery A, Staessen JA, et al.: Is prevention of vascular dementia possible? The Syst-Eur Vascular Dementia Project. *Aging Clin Exp Res* 1991, 3:373–382.
- 15 Forette F, Seux ML, Staessen JA, et al., on behalf of the Syst-Eur investigators: Prevention of dementia in randomised double-blind placebo-controlled Systolic Hypertension in Europe (Syst-Eur) trial. *Lancet* 1998, 352:1347–1351.
- Analysis of data from the Vascular Dementia Side Project of the Syst-Eur Trial showed a reduction in the incidence of dementia of 50%. Treatment mainly prevented Alzheimer's dementia, a somewhat unexpected finding. The authors suggest that specific neuroprotection by CCBs might partly explain this result.
- 16 Staessen JA, Wang JG, Thijs L, et al., for the Systolic Hypertension in Europe Trial Investigators: Use of dihydropyridines for antihypertensive treatment in older patients: evidence from the Systolic Hypertension in Europe Trial. *J Clin Basic Cardiol* 2000, 3:15–21.

This paper reviews the main Syst-Eur results published before 2000. These results show that stepwise antihypertensive drug treatment starting with the CCB nitrendipine, improves prognosis in older patients with ISH.

- 17 Amery A, Brixio P, Clement D, et al.: Mortality and morbidity results from the European Working Party on High Blood Pressure in the Elderly Trial. *Lancet* 1985, 1349–1354.
- 18 Coope J, Warrender TS: Randomised trial of treatment of hypertension in elderly patients in primary care. *BMJ* 1986, 293:1145–1151.
- 19 The Management Committee of the National Blood Pressure Study: Treatment of mild hypertension in the elderly. A study initiated and administered by the National Heart Foundation of Australia. *Med J Aust* 1981, 2:398–402.
- 20 Kuramoto K, Matsushita S, Kuwajima I, Murakami M: Prospective study on the treatment of mild hypertension in the aged. *Jpn Heart J* 1981, 22:75–85.
- 21 Dahlof B, Lindholm LH, Hansson L, et al.: Morbidity and mortality in the Swedish Trial in Old Patients with Hypertension (STOP-Hypertension). *Lancet* 1991, 338:1281–1285.

- 22 MRC Working Party: Medical Research Council trial of treatment of hypertension in older adults: principal results. *BMJ* 1992, 304:405-412.
 - 23 Liu L, Wang JG, Gong L, et al., for the Systolic Hypertension in China (Syst-China) Collaborative Group: Comparison of active treatment and placebo for older patients with isolated systolic hypertension. *J Hypertens* 1998, 16:1829.
- In older Chinese patients with ISH, antihypertensive treatment prevents cardiovascular complications, in particular strokes. The Syst-China results suggest that approximately 60% of all stroke-related deaths might be prevented by nitrendipine-based treatment.
- 24 Wang JG, Staessen JA, Gong L, et al., for the Systolic Hypertension in China Investigators: Subgroup analysis of the placebo-controlled Chinese trial on isolated systolic hypertension in the elderly. *Arch Intern Med* 2000, 160:211-220.
- This subgroup analysis of the Syst-China Trial showed that the benefit of treating older Chinese patients with ISH was particularly evident in diabetics and for cardiovascular endpoints tended to be larger in nonsmokers. Other baseline patient characteristics did not significantly influence the benefit of treatment.
- 25 Furberg CD, Psaty BM: Calcium antagonists: not appropriate as first line antihypertensive agents. *Am J Hypertens* 1996, 9:122-125.
 - 26 Furberg CD, Psaty BM: JNC VI: timing is everything. *Lancet* 1997, 350:1413.
 - 27 Heckbert SR, Longstreth WT, Psaty BM, et al.: The association of antihypertensive agents with MRI white matter findings and modified mini-mental state examination in older adults. *J Am Geriatric Soc* 1997, 45:1423-1433.
 - 28 Psaty BM, Furberg CD: Clinical implications of the World Health Organization - International Society of Hypertension statement on calcium antagonists. *J Hypertens* 1997, 15:1197-1200.
 - 29 Estacio RO, Jeffers BW, Hiatt WR, et al.: The effect of nisoldipine as compared with enalapril on cardiovascular outcomes in patients with non-insulin-dependent diabetes and hypertension. *N Engl J Med* 1998, 338:645-652.
 - 30 Tatti P, Pahor M, Byington RB, et al.: Outcome results of the Fosinopril versus Amlodipine Cardiovascular Events Randomized Trial (FACET) in patients with hypertension and NIDDM. *Diabetes Care* 1998, 21:597-603.
 - 31 Alderman MH, Madhavan S, Cohen H: Calcium antagonists and cardiovascular events in patients with hypertension and diabetes. *Lancet* 1998, 351:216-217.
 - 32 Byington RB, Craven TE, Furberg CD, et al.: Isradipine, raised glycosylated haemoglobin, and risk of cardiovascular events. *Lancet* 1998, 350:1075-1076.
 - 33 Pahor M, Kritchewsky SB, Zuccala G, et al.: Diabetes and risk of adverse events with calcium antagonists. *Diabetes Care* 1998, 21:193-194.
 - 34 Hamet P, Gong L: Chinese contribution to trials on antihypertensive therapy. *J Hypertens* 1996, 14:S123-S128.
 - 35 Gong L, Zhang W, Zhu Y, et al.: Shanghai trial of nifedipine in the elderly (STONE). *J Hypertens* 1996, 14:1237-1245.
 - 36 Wang JG, Liu G, Wang X, et al., on behalf of the Syst-China Investigators: Long-term blood pressure control in older Chinese patients with isolated systolic hypertension: a progress report on the Syst-China trial. *J Hum Hypertens* 1996, 10:735-742.
 - 37 Staessen JA, Thijs L, Gasowski J, et al., for the Systolic Hypertension in Europe (Syst-Eur) Trial Investigators: Treatment of isolated systolic hypertension in the elderly: further evidence from the systolic hypertension in Europe (Syst-Eur) trial. *Am J Cardiol* 1998, 82:20R-22R.
 - 38 Pahor M, Psaty BM, Furberg CD: Treatment of hypertensive patients with diabetes. *Lancet* 1998, 351:689-690.
 - 39 Schrier RW, Estacio RO: Additional follow-up from the ABCD Trial in patients with type 2 diabetes and hypertension [Editorial]. *N Eng J Med* 2000, 343:1969-1969.
- This update of the ABCD trial reports on additional cases of myocardial infarction, congestive heart failure, and cardiovascular death, discovered after the publication of the main results in 1998. Based on these new findings, the authors report that the relative risk of fatal and nonfatal myocardial infarction in the nisoldipine group has decreased to 4.2 (as compared to 7.0 according to the data published in 1998). This does not however alter the conclusion of the study.
- 40 Hofman A, Ott H, Breteler MMB, et al.: Atherosclerosis, apolipoprotein E, and prevalence of dementia and Alzheimer's disease in the Rotterdam Study. *Lancet* 1997, 349:151-154.
 - 41 Morich FJ, Bieber F, Lewis JM, et al., for the North American Nimodipine Study Group: Nimodipine in the treatment of probable Alzheimer's disease: result of two multicentre trials. *Clin Drug Invest* 1996, 11:185-195.
 - 42 Parneti L, Senin U, Mecocci P: Cognitive enhancement therapy for Alzheimer's disease. The way forward. *Drugs* 1997, 53:752-768.
 - 43 Fagard R, Staessen JA, Thijs L, et al., for the Systolic Hypertension in Europe (Syst-Eur) Trial Investigators: Response to antihypertensive therapy in older patients with sustained and nonsustained systolic hypertension. *Circulation* 2000, 102:1139-1144.
- This analysis of data from the Ambulatory Blood Pressure Monitoring Side Project of the Syst-Eur Trial showed that in elderly patients, sustained ISH is more harmful than nonsustained or white coat hypertension and that the favorable effects of active treatment were most evident in those with daytime systolic ambulatory BP ≥ 160 mm Hg.
- 44 Blacher J, Staessen JA, Gierd X, et al.: Pulse pressure not mean pressure determines cardiovascular risk in older hypertensive patients. *Arch Intern Med* 2000, 160:1085-1089.
- This meta-analysis, based on individual patient data of three outcome trials in elderly hypertensives, indicated that pulse pressure, and not mean pressure is the major determinant of cardiovascular risk.
- 45 Staessen JA, Gasowski J, Wang JG, et al.: Risk of untreated and treated isolated systolic hypertension in the elderly: meta-analysis of outcome trials. *Lancet* 2000, 355:865-872.
- This meta-analysis includes data of over 15,000 patients from 8 trials. In untreated patients systolic BP was a better predictor of outcome than diastolic BP. The authors conclude that antihypertensive drug treatment improved outcome in older patients with systolic BP ≥ 160 mm Hg. Furthermore, the absolute benefit was greater in men and in patients with cardiovascular complications and increased with older age and higher pulse pressure.
- 46 Cavallini MC, Roman MJ, Pickering TG, et al.: Is white coat hypertension associated with arterial disease or left ventricular hypertrophy? *Hypertension* 1995, 26:413-419.
 - 47 Cuspidi C, Marabini M, Lonati L, et al.: Cardiac and carotid structure in patients with established hypertension and white coat hypertension. *J Hypertens* 1995, 13:1707-1711.
 - 48 Verdecchia P, Schillaci G, Borgioni C, et al.: White coat hypertension and white coat effect: similarities and differences. *Am J Hypertens* 1995, 8:790-798.
 - 49 Khattar RS, Senior R, Lahiri A: Cardiovascular outcome in white-coat versus sustained mild hypertension - a 10-year follow-up study. *Circulation* 1998, 98:1892-1897.
 - 50 Nichols WW, Rourke MF: Properties of the Arterial wall. In McDonald's Blood Flow in Arteries: Theoretical, Experimental and Clinical Principles, 4th ed. Edited by Edward Arnold. London: 1998.
 - 51 Verdecchia P, Porcellati C, Schillaci G, et al.: Ambulatory blood pressure: an independent predictor of prognosis in essential hypertension. *Hypertension* 1994, 24:793-801.
 - 52 Redon J, Campos C, Narciso ML, et al.: Prognostic value of ambulatory blood pressure monitoring in refractory hypertension: a prospective study. *Hypertension* 1998, 31:712-718.
 - 53 Staessen JA, Amery A, Clement D, et al.: Twenty-four hour blood pressure monitoring in the Syst-Eur trial. *Aging Clin Exp Res* 1992, 4:85-91.
 - 54 Emelianov D, Thijs L, Staessen JA, et al., on behalf of the Syst-Eur investigators: Conventional and ambulatory measurements of blood pressure in old patients with isolated systolic hypertension: baseline observations in the Syst-Eur trial. *Blood Press Monit* 1998, 3:173-180.
 - 55 Perloff D, Sokolow M, Cowan RM, et al.: Prognostic value of ambulatory blood pressure measurements: further analysis. *J Hypertens* 1989, 7(Suppl):3-10.
 - 56 Ohkubo T, Imai Y, Tsuji I, et al.: Prediction of mortality by ambulatory blood pressure monitoring versus screening blood pressure measurements: a pilot study in Ohasama. *J Hypertens* 1997, 15:357-364.
 - 57 Staessen JA, Thijs L, Fagard R, et al., for the Systolic Hypertension in Europe Trial Investigators: Predicting cardiovascular risk using conventional vs ambulatory pressure in older patients with systolic hypertension. *JAMA* 1999, 282:539-546.
 - 58 The Joint National Committee on Detection, Evaluation and Treatment of High Blood Pressure: The sixth report of the Joint National Committee on Detection, Evaluation and Treatment of High Blood Pressure (JNC-VI). *Arch Intern Med* 1997, 157:2413-2446.
 - 59 Guidelines Subcommittee: 1999 World Health Organization - International Society of Hypertension Guidelines for the Management of Hypertension. *J Hypertens* 1999, 17:151-183.
 - 60 Darné B, Gierd X, Safar M, et al.: Pulsatile versus steady component of blood pressure: a cross-sectional analysis and a prospective analysis on cardiovascular mortality. *Hypertension* 1989, 13:392-400.
 - 61 Rutan GH, Kuller LH, Neaton JD, et al.: Mortality associated with diastolic hypertension and isolated systolic hypertension among men screened for the Multiple Risk Factor Intervention Trial. *Circulation* 1988, 77:504-514.

- 62 Wilking SVB, Belanger AJ, Kannel WB, et al.: Determinants of isolated systolic hypertension. *JAMA* 1988, 260:3451–3455.
- 63 Millar JA, Lever AF: Pulse pressure predicts coronary but not cerebrovascular events in placebo-treated male subjects of the MRC trial [abstract]. *J Hypertens* 1996, 14(Suppl):194.
- 64 Benetos A, Safar M, Rudnicki A, et al.: Pulse pressure: a predictor of long-term cardiovascular mortality in a French male population. *Hypertension* 1997, 30:1410–1415.
- 65 Madhavan S, Ooi WL, Cohen H, et al.: Relation of pulse pressure and blood pressure reduction to the incidence of myocardial infarction. *Hypertension* 1994, 23:395–401.
- 66 Fang J, Madhavan S, Cohen H, Alderman MH: Measures of blood pressure and myocardial infarction in treated hypertensive patients. *J Hypertens* 1995, 13:413–419.
- 67 Mitchell GF, Moyé LA, Braunwald E, et al.: Sphygmomanometrically determined pulse pressure is a powerful independent predictor of recurrent events after myocardial infarction in patients with impaired left ventricular function. *Circulation* 1997, 96:4254–4260.
- 68 Franklin SS, Gustin WIV, Wong ND, et al.: Hemodynamic patterns of age-related changes in blood pressure: the Framingham Heart Study. *Circulation* 1997, 96:308–315.
- 69 Neaton JD, Wentworth D, for the Multiple Risk Factor Intervention Trial Research Group: Serum cholesterol, blood pressure, cigarette smoking, and death from coronary heart disease: overall findings and differences by age for 316 099 white men. *Arch Intern Med* 1992, 152:56–64.
- 70 Abernethy J, Borhani NO, Hawkins CM, et al.: Systolic blood pressure as an independent predictor of mortality in the hypertension detection and follow-up program. *Am J Prev Med* 1986, 2:123–132.
- 71 Kannel WB, Gordon T, Schwartz MJ: Systolic versus diastolic blood pressure and risk of coronary heart disease: the Framingham Study. *Am J Cardiol* 1971, 27:335–346.
- 72 Safar M: Roles of systolic and pulse pressure. *Hypertension* 1997, 30:146–147.