

Risk Assessment for the Population of Kyiv, Ukraine as a Result of Atmospheric Air Pollution

Oleksandr Popov,¹ 
 Andrii Iatsyshyn,¹ 
 Valeriia Kovach,¹ 
 Volodymyr Artemchuk,² 
 Iryna Kameneva,² 
 Dmytro Taraduda,³ 
 Vitaliy Sobyna,³ 
 Dmitry Sokolov,³ 
 Maksim Dement,³ 
 Teodoziia Yatsyshyn,⁴ 

1 State Institution, Institute of Environmental Geochemistry of National Academy of Sciences of Ukraine, Kiev, Ukraine

2 Pukhov Institute for Modelling in Energy Engineering of National Academy of Sciences of Ukraine, Kiev, Ukraine

3 National University of Civil Defense of Ukraine, Kharkiv, Ukraine

4 Ivano-Frankivsk National Technical University of Oil and Gas, Ivano-Frankivsk, Ukraine

Corresponding author:
 Valeriia Kovach
 Tel. +380967815496
 valeriia.kovach@gmail.com

Introduction

Air pollution is a large problem in many countries around the world. The main sources of atmospheric air pollution are transportation, power companies, chemically hazardous enterprises, metallurgy, machine-building, metal-processing enterprises and industries, building materials industries, and lumber operations, among others.¹ Atmospheric pollution is considered to be the most dangerous form of pollution by volume, as

Background. According to the World Health Organization, 92% of the world's population lives in places where air quality levels exceed recommended limits. Recently, Ukraine had the most deaths per every 100,000 people (out of 120 countries) attributed to atmospheric air pollution. High levels of atmospheric air pollution have been observed not only in typically industrial regions, but in Ukraine's capital, Kyiv, as well.

Objectives. The aim of the present study was to establish the state of air pollution in Kyiv and perform a risk assessment of associated human health effects.

Methods. Using official statistics and state monitoring data, the study aimed to identify and analyze risks to the health of Kyiv's population associated with air pollution. The following methods were used: systematic, functional and comparative analysis, risk theory, mathematical modeling, probability theory and mathematical statistics, as well as geographic information system technologies for digital map design and objective-oriented methodology for software design systems.

Results. The risk values across different areas of the city varied significantly, indicating that atmospheric air quality remains unstable. Areas with the highest and lowest risk values were identified.

Conclusions. The environmental state of atmospheric air in Kyiv requires greater attention and additional research to identify the causes of air pollution, along with implementation of measures to improve air quality.

Competing Interests. The authors declare no competing financial interests.

Keywords. air pollution, air quality, risk assessment, population health.

Received July 8, 2019. Accepted November 22, 2019.

J Health Pollution 25: (200303) 2020

© Pure Earth

pollution of industrially developed cities by toxic agents becomes irreversible and has negative impacts on the health of the population.

According to the World Health Organization (WHO), 92% of world population breathes air polluted beyond the acceptable limits.² Ukraine is no exception and recently (in September, 2017) ranked first (out of 120 world countries) in deaths caused by atmosphere air pollution, per 100,000 people. Air pollution in Ukraine causes four times more deaths per capita than the five cleanest countries altogether, according to the WHO.³

The main sources of pollution in Kyiv are motor transport and energy enterprises (thermal stations), enterprises of the construction industry, mechanical engineering, chemical-pharmaceutical, and the food industry. In 2015, the total amount of pollutants in the atmosphere from stationary and mobile sources in Kyiv amounted to 144.3 thousand tons. More detailed data on emissions from stationary technogenic facilities in Kiev are presented in Table 1.⁴

The air monitoring system in Ukraine is currently ineffective and has significant problems. There is no data collection and analysis center and

monitoring results are stored in paper form. Most atmospheric scientific research results with some elements of statistical and comparative analysis do not consider the environmental risks to public health. Therefore, a risk assessment of the health of Kyiv's population due to air pollution supported by data and analysis is urgently needed. The aim of the present study was to establish the state of air pollution in Kyiv and perform a risk assessment of associated human health effects.

Methods

Kyiv, the capital of Ukraine, is one of the biggest and oldest cities in Europe. Kyiv is situated in the south of Ukraine, on the border of Polissya. The Dnipro River runs through the city and there is forest steppe on either side of the river. Kyiv has an area of 836 km². The climate is moderately continental, with mild winters and warm summers. The average monthly temperature in January and July is -3.5°C and +20.5°C, respectively, with an average annual precipitation of 649 mm. Forests, parks and gardens make up more than half of the almost 80 000 hectares. The population in Kyiv was 2.9 million in December 2017, according to data of the General Statistics Department.⁵ The present study was conducted from September 2016 till September 2018.

In Ukraine, programs monitoring the quality of atmospheric air are aimed at the control of primary pollutants at stationary and mobile stations. In Kyiv, the Central Geophysical Observatory systematically monitors the content of pollutants in atmospheric air at 16 stationary (but manual) pollution observation stations (POS) with samples taken every 6 days, 3-4 times a day. These stations collect data on suspended substances, sulfur dioxide, carbonic oxide, hydrogen

Abbreviations			
MPC	Maximum permissible concentration	PrJSC	Private joint stock company
PJSC	Public joint stock company	RCI	Risk of chronic intoxication
POS	Pollution observation stations	RITE	Risk of immediate toxic effects
		SMC	Software-modeling complex

sulfide, phenol, hydrogen fluoride, ammonia, formaldehyde, and eight heavy metals. Their composition is defined depending on the emissions of pollutants in atmospheric air from enterprises located in the area of the observation stations. The distribution of POS in Kyiv is shown in Figure 1.

In the countries of the former USSR, health standards for permissible pollutants burden consists of maximum permissible concentration (MPC), relatively safe impact levels and different complex values. In Ukraine, and most post-Soviet

states, there are approved MPCs for pollutants used for the assessment of atmosphere air. Maximum permissible concentrations are set for each substance and for certain combinations of pollutants in atmospheric air. Maximum permissible concentrations of pollutants in the atmospheric air of populated areas are regulated as daily averages and maximum single MPC.

When the concentration values are taken, they are compared to maximum single values of those concentrations (thus the amount of excess is defined).

Stationary sources	2010	2013	2014	2015	2016
Total air emissions from stationary pollution sources, (thousand tons)	28.6	31.9	31.4	26.7	34.3
Air emissions from stationary pollution sources per km ² , (tons)	34.3	38.1	37.6	31.9	41.1
Air emissions from stationary pollution sources per capita, (kg)	10.3	11.2	10.9	9.2	11.8
Air emissions of particulate matters from stationary pollution sources, (thousand tons)	4.0	4.8	4.9	4.7	5.8
Air emissions of sulfur dioxide from stationary pollution sources, (thousand tons)	7.6	11.3	9.9	6.5	12.3
Air emissions of nitrogen dioxide from stationary pollution sources, (thousand tons)	9.8	9.1	7.8	6.8	7.5
Air emissions of carbon monoxide from stationary pollution sources, (thousand tons)	2.8	2.5	2.1	2.1	2.2
Air emissions of non-methane volatile organic compounds from stationary pollution sources, (thousand tons)	3.5	3.4	5.9	5.8	5.7

Table 1 — Dynamics of Emissions from Stationary Sources




Figure 1 — Primary enterprises impacting air quality in Kyiv (blue triangles, see Table 6) and atmospheric air pollution observation stations (red triangles)

The average monthly or yearly values are likewise compared with MPCs of daily average values.

In order to assess atmospheric air pollution levels, the air pollution index was used, following Equation 1.

Equation 1

$$API_i = \left(\frac{C_i}{MPC_{a.d.i}} \right)^{\beta_i}$$

where, API is the air pollution index; C_i is the average concentration of i -additive in time (per month or per year) calculated for the station, city or a group of cities; $MPC_{a.d.i}$ is the

maximum permissible daily average concentration of a pollutant in the air; and β_i is the dimensionless constant of reducing the damage of i -substance to the damage of sulfur dioxide which accepts the values of 1.7; 1.3; 1.0; 0.9 for 1, 2, 3, 4 classes of hazard of pollutants, respectively.

The above-mentioned health standards characterize only the level of chemical air pollution of a territory under investigation and do not define the impact of pollution on human health. Therefore, a risk assessment was conducted to determine the probability of negative effects on human health for individuals or groups of people affected by a particular chemical

reagent. It is characterized by the value that lies within (0...1) interval, where 0 indicates the absence of effect and 1 indicates the greatest effect.

In order to assess the negative impact of anthropogenic air pollution on the health of a population which may manifest itself in immediate or chronic toxic effects (carcinogenic and immunotoxic), two groups of models have been introduced: threshold and non-threshold.⁶ Acute toxicity (immediate toxic manifestations) has a definite threshold character. In order to assess the risk of immediate toxic effects (RITE), the model of personal impact thresholds is used. In terms of air pollution, this model is defined by

the formula in Equation 2:

Equation 2

$$RITE = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{a+b \lg(C/MPC_{m.s.})} \exp(-\tau^2/2) d\tau$$

where, a, b are the parameters depending on toxicological properties of the substance; C is the toxicant concentration in the air (mg/m^3); and $MPC_{m.s.}$ is the maximum permissible single concentration of a pollutant in the air. To find RITE, it is necessary to calculate a defined exponential function integral: $\exp(-\tau^2/2) = e^{-\tau^2/2}$ (number $e = 2,71$), where, τ is an integration variable that takes a value from $-\infty$ to $a + b \lg(C/MPC_{m.s.})$ and $d\tau$ is a differential variable τ .

The risk of chronic intoxication (RCI) is defined for a certain time period using Equation 3.

Equation 3

$$RCI = 1 - \exp \left[-0,174 \left(\frac{C}{MPC_{a.d.} \cdot K} \right)^\beta \cdot t \right]$$

where, β is the coefficient that takes into account toxic properties of the substance; K is the parameter depending on the class of hazard of the substance; C is the pollutant concentration in the air over time (t) (mg/m^3); and $MPC_{a.d.}$ is the maximum permissible daily average concentration.

The values of $a, b, \beta,$ and K depend on the substance hazard classification and are given in Table 2.

In order to define the level of air pollution attributed to a certain substance, average concentration values along with the highest values of concentration at a certain period of time are identified.

Substance hazard class	Substance characteristics	a	b	β	K
1 st	Extremely hazardous	-9.15	11.66	2.40	7.5
2 nd	Highly hazardous	-5.51	7.49	1.31	6.0
3 rd	Moderately hazardous	-2.35	3.73	1.00	4.5
4 th	Low hazard	-1.41	2.33	0.86	3.0

Table 2 — Coefficient values of a, b, β and K

The level of integral risk from simultaneous impact of all investigated substances was calculated with the formula in Equation 4:

Equation 4

$$R_\Sigma = 1 - \prod_{i=1}^n (1 - R_i)$$

where, R_Σ is the level of integral risk from simultaneous impact of all investigated substances; R_i is the health risk value as a result of the impact of i -substance; and n is the number of substances under investigation.

In Ukraine, according to applicable instructional guidelines of the Ministry of Health of Ukraine, the individual carcinogenic risk, hazard coefficient for exposure to i -substance, level of noncarcinogenic risks and hazard index are defined.⁷ The carcinogenic risk was calculated using Equation 5:

Equation 5

$$CR = LADD \cdot SF$$

where, CR is the carcinogenic risk; $LADD$ is the average daily dose over a lifetime ($mg/(kg \cdot day)$); and SF is the slope factor ($mg/(kg \cdot day)^{-1}$).

The average daily dose over a lifetime is defined by Equation 6:

Equation 6

$$LADD = [(C_a \cdot T_{out} \cdot V_{out}) + (C_h \cdot T_{in} \cdot V_{in})] \cdot EF \cdot ED / (BW \cdot AT \cdot 365)$$

where, C_a is the concentration of the substance in the ambient air (mg/m^3); C_h is the concentration of the substance in the room air (mg/m^3); T_{out} is the time spent outdoors (h/day); T_{in} is the time spent indoors (h/day); V_{out} is the breathing rate outdoors (m^3/hr); V_{in} is the rate of breathing indoors (m^3/hr); EF is the frequency of exposure (days/year); ED is the duration of exposure (years); BW is body weight (kg); and AT is the exposure average period (years).

The hazard coefficient is defined using Equation 7:

Equation 7

$$HQ_i = C_i / RfC_i$$

where, HQ_i is the hazard coefficient for exposure of i -substance; C_i is the level of exposure to i -substance (mg/m^3); and RfC_i is the safe exposure level of i -substance (mg/m^3).

The hazard index is defined using Equation 8:




Figure 2 — Module structure of risk assessment system

level), or were set in accordance with study conditions.⁸

The authors of the present study have designed a software-modeling complex (SMC) to solve problems of environmental pollution in technogenic loaded territories.⁸⁻¹⁰ The essential advantages of this SMC above other models are the wide range of investigated environmental problems, the presence of a universal database monitoring the sources of pollution and pollutants, POS data, different environmental health values (demographics, population morbidity, etc.) and the existence of a module of environmental risks calculation, which is thought to be the most objective hazard assessment. The SMC database obtained data from different monitoring sources in Ukraine from 2005 to 2017.

Risk level	Hazard coefficient of noncarcinogenic effects for separate combinations	Hazard index of noncarcinogenic effects for a group of unidirectional action	Individual lifetime carcinogenic risk
High	>3	>6	>10 ⁻³
Alert	1.1-3	3.1-6	1.1 · 10 ⁻⁴ -1.0 · 10 ⁻³
Acceptable	0.11-1.0	1.1-3.0	1.1 · 10 ⁻⁶ -1.0 · 10 ⁻⁴
Minimal (target)	0.1 and less	1.0 and less	10 ⁻⁶ and less

Table 3 — Risk Level Classification

The designed SMC makes it possible to observe the spatial distribution of atmospheric air pollution levels and individual risk distributions for a population (risk maps). The risk assessment completes the process of modeling anthropogenic pollution loads on a city's population and defines the criteria for managerial decisions.

Figure 2 shows the module structure of the risk assessment system that is a part of the SMC. The input modeling data are POS data, pollutant characteristics, sources of pollution parameters, and metrological parameters.

The designed SMC realizes an expert approach to the tasks of risk assessment for public health, corresponding with the current legislation in Ukraine, taking into account regulations of the US Environmental Protection Agency, and grounded by the results of the analysis and appropriate interpretation of ecological monitoring data.^{11,12} The

Equation 8

$$HI = \sum_{i=1}^n HQ_i$$

where, *HI* is the hazard index; *HQ_i* is the hazard coefficients for individual components of the solution of chemical reagents that produce an effect.

The main parameter determining

the types of risk considered in this analysis is the concentration of pollutants in atmospheric air, which is measured by the monitoring system or in cases where measurements are absent, modeling is based on data on emissions and meteorological conditions. All other parameters listed in the above formulas are either selected from the relevant reference literature tables, for example Popov *et al.*, for each substance respectively (MPC, slope factor, safe exposure

given approach includes the following stages:

1. Hazard identification: defining informative parameters (dangerous substances, hazard indices) on the basis of monitoring and expert data.¹³
2. Assessment of the impact of anthropogenic pollution loads as a result of data analysis and spatially distributed modeling: defining and visualizing territories with excess critical concentration of hazardous substances.
3. Assessment dependence, “dose – reaction”: defining with the help of mathematic models the probable risk assessment for carcinogenic and noncarcinogenic diseases.
4. Characterizing risk distribution: risk ranking and taking into account the approved methodology and other means of visual presentation of the obtained results.⁷

The framework of the given approach uses the US Environmental Protection Agency classification of human health risks and reduces uncertainties during the assessment of total noncarcinogenic risk of substances of unidirectional action (Table 3).¹¹

Results

According to the air pollution index, air pollution in the summer months of 2017 was generally characterized as very high. It differed across regions: at POS 7 it was characterized as very high (greater than 14); at POS 5 as high (7-14); at POS 2 as elevated (5-7); and at POS 1 as low (less than 5). In general, the average monthly concentrations of four pollutants in the city exceeded the maximum permissible daily average concentration: formaldehyde by 4.3




Figure 3 — Risk of chronic intoxication (a) and risk of immediate toxic effects (b) as a result of air pollution in Kyiv in January-December 2017

Location	2005		2008		2011		2017	
	RCI	RITE	RCI	RITE	RCI	RITE	RCI	RITE
Hydropark	0.085	0.137	0.063	0.140	0.068	0.025	0.077	0.026
National complex "Ekspocentr Ukraina"	0.061	0.051	0.053	0.120	0.071	0.032	0.087	0.067
Moskovska Square	0.192	0.406	0.060	0.089	0.144	0.157	0.178	0.160
Peremohy Square	0.170	0.458	0.136	0.298	0.210	0.306	0.219	0.226
Bessarabska Square	0.193	0.693	0.275	0.472	0.255	0.321	0.296	0.211
Maidan Nezalezhnosti	0.173	0.514	0.202	0.390	0.212	0.264	0.241	0.202
Darnytska Square	0.163	0.386	0.151	0.358	0.174	0.215	0.182	0.187

Table 4 — Dynamics of Risk Values for the Health of the Population in Kyiv




Figure 4a — Risk of chronic intoxication in Kyiv territory according to averaged risk values from 2011-2017

times, nitrogen dioxide by 4 times, nitrogen oxide by 1.7 times, phenol by 1.1 times, and suspended substances by 1.3 times. The contents of other substances in the air did not exceed health-based exposure limits.

Since 2007, significant seasonal changes in formaldehyde concentrations have been observed.

Thus, in the winter months, this figure does not exceed the MPC, and in the summer the concentration increases sharply and exceeds the MPC by 5-7 times, which is quite dangerous for the most sensitive categories of the city population. These fluctuations may be caused by the intensification of construction and repair work, which is common in the city within each

district. However, further investigation is needed to identify all the causes of this situation.

In terms of spatial distribution, the most polluted areas in June were the districts of Bessarabska Square, Oleksandr Dovzhenko street (near Shuliavka subway station), Demiiivska Square, Kaunasska street, Obolonsky Avenue, and Peremohy Square (the region of Sviatoshyn subway station), where the pollution level was characterized as very high. Somewhat less polluted, but still within the "high" level were the regions of Semena Skliarenka street, Akademika Strazheska (intercrossing with Vatslava Havela Boulevard), Lesi Ukraiinky Boulevard and Popudrenka street. Pollution levels were lower at Inzhenera Borodina street (area of Darnytsia carriage repair plant) and the Hydropark (area near a subway bridge and highway). A low pollution level was observed at Nauky Avenue (green zone, with high concentrations of parks and forest).

Figure 3 shows map examples of integral RITE and RCI as the result of air pollution in Kyiv during 2017, drawn by the interpolation method of weighted distances. Interpolation was conducted according to the multiplicity of exceeding MPC. The obtained risk values are presented in Figure 3 as outlines in different shades according to the values shown in the map legend.

The risk values for the population of Kyiv proved to be very high. The worst environmental conditions were identified in the northwest where the RCI exceeded 0.2, indicating that more than 20% of the population of the given districts are at risk for chronic diseases. The RITE is also highest for the population in the northwestern part of the city.

The designed SMC makes it possible to obtain the dynamics of risk values for the region under investigation (Table 4).

According to the data, the risk values differ across districts of the city, indicating unstable ecological conditions of atmospheric air in Kyiv. Thus, the lowest risk values were found at the territory of the Hydropark (from 0.025 to 0.137) which is a park complex with minimum impact from traffic and industry.

The greatest risk values are found in Bessarabska Square and Maidan Nezalezhnosti (0.693 and 0.514, respectively, for immediate toxic effects in 2005). In general, the level of RITE in Kyiv is very high, although for some districts it has somewhat lessened compared to previous years and the values of RCI are slowly increasing.

Figure 4a shows the distribution of Kyiv's territory with the risk levels of chronic intoxication according to the averaged risk data that were obtained from 2011 to 2017. Each bar of Figure 4a represents the percentage of Kyiv's territory (Y axis), which is characterised by the correspondent risk levels (X axis). Figure 4b shows the distribution of Kyiv's territory with the risk of immediate toxic effects according to the averaged risk values for the same period of time (each bar represents part of Kyiv's territory in percentage with the defined risk levels).

The analysis showed that the majority of Kyiv territory is characterized by 15-17% RCI values, 30-50% RITE values, a hazard index value of 4-6, a hazard coefficient value (for nitrogen dioxide) from 2-2.5, and a carcinogenic risk (for formaldehyde) from $2 \cdot 10^{-5}$ - $5 \cdot 10^{-5}$ (Table 5).




Figure 4b — Risk of immediate toxic effects in Kyiv territory according to averaged risk values from 2011-2017

Value (risk type and pollutant)	Year			
	2005	2008	2011	2017
Hazard coefficient				
Nitrogen dioxide 2-2.5	41%	67%	60%	57%
Carcinogenic risk				
Formaldehyde $2 \cdot 10^{-5} - 2.5 \cdot 10^{-5}$	55%			
Formaldehyde $3 \cdot 10^{-5} - 4 \cdot 10^{-5}$		64%		
Formaldehyde $4 \cdot 10^{-5} - 5 \cdot 10^{-5}$			53%	51%

Table 5 — Analysis of Kyiv Area Distribution According to Hazard Coefficient and Carcinogenic Risk Values

In order to obtain a more precise environmental risk assessment, it is necessary to modernize the existing monitoring system, increase the number of observation stations and measurements of harmful pollutants. Risk value deviations depend on high concentrations of formaldehyde, which is not measured at some observation stations. The absence of values for some pollutants and

the lack of statistical data affect the accuracy of the obtained results and limit the possibilities of their current interpretation.

Discussion

There have been many studies of the impact of air pollution on population health. Min *et al.* and Mate *et al.* have reported on the risk of mortality

from air pollution by solid particles with a diameter of $\leq 10 \mu\text{m}$, nitrogen dioxide and sulfur dioxide.^{14,15} In Buoli *et al.*, Italian scientists investigated the effect of particulate matter and nitrogen oxides on the mental health of the population, in particular the

risk of new symptoms of depression.¹⁶ In work by Sanidas *et al.*, American scientists stated that air pollution is a significant risk factor for hypertension in all age groups.¹⁷ Studies have shown that the presence of particulate matter of different sizes in city air

increases the risk of cardiovascular disease and significantly increases the likelihood of adverse health effects such as chronic cough, asthma, renal failure and infertility.^{18,19} In a study by Yang *et al.*, Chinese and American scientists investigated the association between the increase in the number of stillbirths with increasing average daily concentrations of particulate matter with a diameter of ≤ 2.5 and $\leq 10 \mu\text{m}$ at different stages of pregnancy.²⁰

The results of the present analysis are generally in agreement with the conclusions of previous investigations. For example, an analysis of macro scale observations from 1997-2013 reported continuous pollution of carcinogenic substances and the variability of their concentration in separate periods in Kyiv's atmosphere air was found to depend on the character of industrial activities.²¹ In addition, the results indicated stable pollutants slightly above the MPC for benz/a/pyrene, formaldehyde and lead. All of the benz/a/pyrene samples exceeded the MPC by 2.3-3.3 times, almost all granules (particles) were 1.1-1.5 times bigger than their normal size.

The above-mentioned study calculated the total carcinogenic risk for human health in Kyiv under

Downloaded from http://meridian.allenpress.com/jhp/article-pdf/10/25/200303/2649919/2156-9614-10_25_200303.pdf by India user on 16 August 2022

Number on the map	Object	Number on the map	Object
1, 2	TCPS-5 (pipes 1 and 2)	16	PrJSC "Kyiv Electric Car Repair Plant"
3	TCPS-4 (Darnytska)	17	SE, "State-Run Enterprise of Special Tool Engineering, Arsenal"
4	Korchevatsky building materials mill	18	Chipboards plant "Avers"
5	Plant "Energiya"	19	LLC "Kyivguma"
6	Branch "Darnytsia Carriage-Repair Plant" owned by PJSC "Ukrzaliznytsya"	20,21,22	PrJSC Asphalt and concrete plant "Stolychnyi" (pipes 1, 2, 3)
7	Bortnyky aeration station	23	Locomotive depot "Darnytsya"
8	Open joint stock company "Ukrplastyk"	24	Plant "Radykal"
9	TCPS-6 (pipe 1)	25	PrJSC "Kyiv Glass-Container Plant"
10	TCPS-6 (pipe 2)	26	Banknote-printing and minting works of "National Bank of Ukraine"
11	PJSC "Plant Mayak"	27	State joint stock holding company "Artem"
12	SE "Plant 410 of Civil Aviation"	28	PJSC "Pharmak"
13	PJSC Scientific and production center "Borshchahivskiy Chemical and Pharmaceutical Plant"	29	SE "Generator Plant"
14	Specialized utility company "Kyiv Crematorium"	30	PrJSC Plant "Smithy at Rybalskyi"
15	Joint stock company "First Kyiv Machine-Building Plant"		

Abbreviations: TCPS, thermal cogenerating power station; SE, state enterprise; LLC, limited liability company.

Table 6 — Largest Enterprises Influencing Atmospheric Air Quality in Kyiv (with Reference to Figure 1)

Year	Total thousand tons		Sources				In the calculation			
			Stationary polluting sources		Mobile polluting sources		Per capita, kg		Per 1 km ² of the region area, kg	
	Sulfur dioxide	Nitrogen dioxide	Sulfur dioxide	Nitrogen dioxide	Sulfur dioxide	Nitrogen dioxide	Sulfur dioxide	Nitrogen dioxide	Sulfur dioxide	Nitrogen dioxide
2005	5.9	29.5	4.7	13.1	1.2	16.4	2.2	11.1	7.1	35.3
2008	8	32.1	6	8.5	2	23.6	2.9	11.7	9.6	38.4
2011	11.9	31.9	9.9	10.7	2	21.2	4.3	11.4	14.2	38.2
2015	8	22.1	6.5	6.8	1.5	15.3	2.8	7.6	10	26.4
2016	12.3	7.5	12.3	7.5	no data	no data	4.2	2.6	14.8	8.9
2017	15	9.3	15	9.3	no data	no data	5.1	3.2	17.9	11.2

Table 7 — Dynamics of Dioxide and Nitrogen Dioxide (Converted to Nitrogen Dioxide) Emissions from Mobile Pollutions Sources

the impact of eight hazardous substances (benz/a/pyrene, formaldehyde, N-nitrosodiethylamine, nitrosodiethylamine, cadmium, lead, nickel and chrome VI) from 1997-2013. The total carcinogenic risk deviated within the limits of $1.5-1.9 \cdot 10^{-3}$ of the risk level. According to the US Environmental Protection Agency classification, these health risk levels are considered high ($>10^{-3}$) and require reduction measures.¹¹ Thus, air pollution levels in Kyiv are not safe for the health of its inhabitants according to an analysis of carcinogenic substances.

The risk distributions and risk dynamics obtained in the present study explain the annual morbidity growth in Kyiv for diseases of the respiratory system, oncology diseases and thyroid cancer.^{22,23} The associations between morbidity level attributed to these diseases in Kyiv and air pollution levels have also been confirmed in studies by Chernichenko *et al.* and the US Environmental Protection Agency.^{11,24}

The power industry is the largest enterprise contributing to the atmospheric air pollution in Kyiv by stationary sources. Since 2012, energy companies, the public joint stock company (PJSC) Kyivenergo and private joint stock company (PrJSC) "Ecostandart" increased their emissions up to 32.8 thousand tons (contributing 79.5% of the total amount of emissions produced by stationary pollution sources in the city).²⁵ Precise data concerning the volume of emissions of the largest stationary pollution sources over recent years is unavailable, as this information is not provided on the website of the Ministry of Environment and Natural Resources of Ukraine. However, since 2017, Darnytsya has used coal of higher quality with less ash content and

higher caloric content. Thus, the amount of particulate matter emissions in 2017 was reduced by 632.0 tons or by 27.8% compared to 2016.²⁶ The largest enterprises in Kyiv that influence the atmospheric air quality are given in Table 6 and their location is shown on the map (Figure 1).

According to data on mobile pollution sources by the General Statistics Department, the amount of polluting emissions in Kyiv has been declining (200.7 thousand tons in 2006, 236.4 thousand tons in 2010, 215.8 thousand tons in 2013, and 144.3 thousand tons in 2015). Unfortunately, in 2016 no calculation of polluting emissions by mobile polluting sources was conducted. In addition, emissions of sulfur dioxide (converted to nitrogen dioxide) increased between 2016 and 2017 (Table 7).²⁷

Conclusions

Analysis of the results of atmospheric air monitoring data demonstrate an unfavorable environmental health situation in Kyiv from 2016-2017. Significant excesses are noted for formaldehyde, nitrogen dioxide, nitrogen oxide, phenol and suspended substances, which significantly raise the risk of toxicity and associated chronic diseases.

Risk assessments, grounded on the basis of monitoring data, reveal that the majority of the city's population is exposed to pollution levels significantly surpassing acceptable limits. The results of the present analysis show that unhealthy atmospheric air conditions have contributed to the increasing morbidity rate in Kiev by 20-30%. The worst situation was observed in the northwestern part of the city.

Further studies are needed examining the status of atmospheric air in

Kyiv along with implementation of necessary measures aimed at air quality improvement. From a public health viewpoint, planning and execution of measures aimed at reducing toxic air levels are urgently needed, along with programs designed to inform the public about the influence of pollution on the health of the residents in Kyiv.

Acknowledgements

This study was funded as part of employment

Copyright Policy

This is an Open Access article distributed in accordance with Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0/>).

References

1. Sówka, I., Bezyk, Y., & Pachurka, L. Analysis and assessment of air quality in the cities area of Wrocław (Poland) and Lviv (Ukraine). *Sci Rev Eng and Env Sci*, 2015(2), 178-192. [cited 2020 Jan 8] Available from http://iiks_pn.sggw.pl/z68/ang7.pdf
2. Fairburn, J., Schüle, S. A., Dreger, S., Karla Hilz, L., Bolte, G. (2019). Social Inequalities in Exposure to Ambient Air Pollution: A Systematic Review in the WHO European Region. *J Environ Res Public Health*, 2019,16(17), 3127. [cited 2020 Jan 8] Available from: <https://www.mdpi.com/1660-4601/16/17/3127>
3. Kubatko, O., Kubatko, O. (2019). Economic estimations of air pollution health nexus. *Environment, Development and Sustainability*, 2019, 21(3), 1507-1517 [cited 2020 Jan 8] Available from: <https://link.springer.com/article/10.1007/s10668-018-0252-6>. (Subscription required to view)
4. Статистичний збірник: довкілля укр аїни за 2016 рік [Statistical yearbook: environment of Ukraine, 2016] [Internet]. Kyiv, Ukraine: State Statistics Service of Ukraine; 2017 [cited 2018 May 30]. 226 p. Available from: http://www.ukrstat.gov.ua/druk/publicat/kat_u/2017/zb/11/zb_du2016pdf.zip Ukrainian.
5. Statistical yearbook of Ukraine for 2017

[Internet]. Kyiv, Ukraine: State Statistics Service of Ukraine; 2018 [cited 2020 Jan 7]. 544 p. Available from: http://www.ukrstat.gov.ua/druk/publicat/kat_u/2018/zb/11/zb_seu2017_e.pdf

6. **Alymov V, Tarasova N.** Tekhnogennyi risk. Analiz i otsenka [Technogenic risk: analysis and assessment]. Moscow, Russia: Akademkniga; 2004. 118 p. Russian.

7. Про затвердження методичних рекомендацій: оцінка ризику для здоров'я населення від забруднення атмосферного повітря [On approval of the methodological recommendations: risk assessment for public health from air pollution] [Internet]. Kyiv, Ukraine: Ministry of Health of Ukraine; 2007 Apr 13 [cited 2018 May 1]. Order No.: 184. Available from: <http://mozdocs.kiev.ua/view.php?id=6902> Ukrainian.

8. **Popov O, Iatsyshyn A, Kovach V, Artemchuk V, Taraduda D, Sobyna V, Sokolov D, Dement M, Yatsyshyn T.** Концептуальні підходи до створення інформаційно-аналітичної експертної системи для оцінки впливу АЕС на довкілля [Conceptual approaches for development of informational and analytical expert system for assessing the NPP impact on the environment]. Nucl Radiat Saf [Internet]. 2018 August 28 [cited 2019 Nov 12];3(79):56-65. Available from: [https://doi.org/10.32918/nrs.2018.3\(79\).09](https://doi.org/10.32918/nrs.2018.3(79).09) Ukrainian.

9. **Popov O, Yatsyshyn A.** Mathematical tools to assess soil contamination by deposition of technogenic emissions. In: Dent D, Dmytruk Y, editor. Soil science working for a living. Cham, Switzerland: Springer; 2017. p. 127-37.

10. **Shkitsa L, Yatsyshyn T, Popov A, Artemchuk V.** Математические средства для улучшения экологического контроля атмосферы на территории буровой установки [The development of mathematical tools for ecological safe of atmosphere on the drilling well area]. [Oil Industry] [Internet]. 2013 Nov [cited 2019 Nov 12]:136-40. Available from: https://www.oil-industry.net/en/Journal/archive_detail.php?ID=9852&art=220821 Subscription required to view. Russian.

11. Supplementary guidance for conducting health risk assessment of chemical mixtures. Washington, DC: US Environmental Protection Agency; 2000 Aug. 209 p. Report No.: EPA/630/R-00/002.

12. **Iatsyshyn AV, Popov OO, Artemchuk VO, Kovach VO, Zinovieva IS.** Автоматизовані інформаційні системи підтримки прийняття управлінських рішень у галузі екологічної безпеки [Automated and information decision support systems for environmental safety]. Inf Technol Learn Tools [Internet]. 2019 [cited 2019 Nov 12];72(4):286-305.

Available from: <https://journal.iitta.gov.ua/index.php/ilt/article/view/2993> Ukrainian.

13. **Popov O, Iatsyshyn A, Kovach V, Artemchuk V, Taraduda D, Sobyna V, Sokolov D, Dement M, Yatsyshyn T, Matvieieva I.** Аналіз можливих причин виникнення надзвичайних ситуацій на аес з метою мінімізації ризику їх виникнення [Analysis of possible causes of NPP emergencies to minimize risk of their occurrence]. Nucl Radiat Saf [Internet]. 2019 March 21 [cited 2019 Nov 12];1(81):75-80. Available from: [https://doi.org/10.32918/nrs.2019.1\(81\).13](https://doi.org/10.32918/nrs.2019.1(81).13) Ukrainian.

14. **Min JY, Kim HJ, Min KB.** Long-term exposure to air pollution and the risk of suicide death: a population-based cohort study. Sci Total Environ [Internet]. 2018 Jul 1 [cited 2019 Nov 12];628-629:573-9. Available from: <https://doi.org/10.1016/j.scitotenv.2018.02.011> Subscription required to view.

15. **Mate T, Guaita R, Pichiule M, Linares C, Diaz J.** Short-term effect of fine particulate matter (PM_{2.5}) on daily mortality due to diseases of the circulatory system in Madrid (Spain). Sci Total Environ [Internet]. 2010 Nov 1 [cited 2019 Nov 12];408(23):5750-7. Available from: <https://doi.org/10.1016/j.scitotenv.2010.07.083> Subscription required to view.

16. **Buoli M, Grassi S, Caldiroli A, Carnevali GS, Mucci F, Iodice S, Cantone L, Pergoli L, Bollati V.** Is there a link between air pollution and mental disorders? Environ Int [Internet]. 2018 Sep [cited 2019 Nov 12];118:154-68. Available from: <https://doi.org/10.1016/j.envint.2018.05.044> Subscription required to view.

17. **Sanidas E, Papadopoulos DP, Grassos H, Velliou M, Tsioufis K, Barbetseas J, Papademetriou V.** Air pollution and arterial hypertension. A new risk factor is in the air. J Am Soc Hypertens [Internet]. 2017 Nov [cited 2019 Nov 12];11(11):709-15. Available from: <https://doi.org/10.1016/j.jash.2017.09.008> Subscription required to view.

18. **Polichetti G, Cocco S, Spinali A, Trimarco V, Nunziata A.** Effects of particulate matter (PM₁₀, PM_{2.5} and PM₁) on the cardiovascular system. Toxicol [Internet]. 2009 June 30 [cited 2019 Nov 12];261(1-2):1-8. Available from: <https://doi.org/10.1016/j.tox.2009.04.035> Subscription required to view.

19. Health aspects of air pollution with particulate matter, ozone and nitrogen dioxide [Internet]. Report on a WHO Working Group; 2003 Jan 13-15; Bonn, Germany. Geneva, Switzerland: World Health Organization; 2003 [cited 2019 Nov 12]. 98 p. Report No.: EUR/03/5042688. Available from: http://www.euro.who.int/__data/assets/pdf_file/0005/112199/E79097.pdf

20. **Yang S, Tan Y, Mei H, Wang F, Li N, Zhao J, Zhang Y, Qian Z, Chang JJ, Syberg KM, Peng A, Mei H, Zhang D, Zhang Y, Xu S, Li Y, Zheng T, Zhang B.** Ambient air pollution the risk of stillbirth: a prospective birth cohort study in Wuhan, China. Int J Hyg Environ Health [Internet]. 2018 Apr [cited 2019 No 12];221(3):502-9. Available from: <https://doi.org/10.1016/j.ijheh.2018.01.014>

21. **Chernychenko IO, Lytvychenko OM, Sovertkova LS, Tsybaliuk SM.** Оцінка канцерогенного ризику для населення промислових міст України [Cancer risk assessment for the population of the industrial cities of Ukraine] Environ Health [Internet]. 2017 May 31 [cited 2019 Nov 12];2(82):17-22. Available from: <http://oaji.net/articles/2017/1827-1498199269.pdf> Ukrainian.

22. Захворюваність населення [Population morbidity] [Internet]. Kyiv, Ukraine: State Statistics Service of Ukraine; [cited 2018 May 30]. [about 1 screen]. Available from: <http://www.kiev.ukrstat.gov.ua/p.php3?c=510&lang=1> Ukrainian.

23. Національний канцер-реєстр України [National Cancer Registry of Ukraine] [Internet]. Kyiv, Ukraine: [The National Cancer Institute]; [cited 2018 May 30]. Available from: <http://www.ncru.inf.ua/publications/index.htm> Ukrainian.

24. **Chernichenko I, Litvichenko O, Tsybalyuk S, Schwager O, Sovetkova L, Balenko N.** Хімічні канцерогени атмосферного повітря: вплив на захворюваність населення на рак щитоподібної залози [Chemical carcinogens of ambient air: influence on the morbidity of the population on thyroid cancer]. [Hygiene of Settlements] [Internet]. 2015 [cited 2019 Nov 12];66:104-10. Available from: http://nbuv.gov.ua/UJRN/gnm_2015_66_17 Ukrainian.

25. Екологічний паспорт міста Київ за 2012 [Kyiv ecological passport for 2012]. Kyiv, Ukraine: Ministry of Ecology and Natural Resources of Ukraine; 2013. Ukrainian.

26. Міфи про Дарницький золовідвал [Myths about Darnytskyi ash dump] [Internet]. Kyiv, Ukraine: Euro Reconstruction; c2019 [cited 2018 May 30]. [about 3 screens] Available from: <http://tec4.kiev.ua/mify-pro-darnytskyj-zolovidval> Ukrainian.

27. Викиди забруднюючих речовин та діоксиду вуглецю у атмосферне повітря (1990-2018 роки) [Emissions of pollutants and carbon dioxide into the atmosphere (1990-2018)] [Internet]. Kyiv, Ukraine: State Statistics Service of Ukraine; [cited 2018 May 30]. [about 2 screens]. Available from: <http://kiev.ukrstat.gov.ua/p.php3?c=1730&lang=1> Ukrainian.