

Risk factors for internal anal sphincter dysfunction in Japanese adults

Tatsuya Abe^{1*}, Toru Kono^{2,3}, Yoshikazu Hachiro¹, Masao Kunimoto¹, Hiroyuki Furukawa²

¹Department of Proctology, Kunimoto Hospital, Asahikawa, Japan

²Division of Gastroenterologic and General Surgery, Department of Surgery, Asahikawa Medical University, Asahikawa, Japan

³Advanced Surgery Center, Sapporo Higashi Tokushukai Hospital, Sapporo, Japan

Email: *t-abe@cf6.so-net.ne.jp

Received 13 December 2012; revised 13 January 2013; accepted 20 January 2013

ABSTRACT

Purpose: The internal anal sphincter provides most of the resting anal tone and is the main muscle responsible for continence. This study was designed to estimate the prevalence of, and identify risk factors associated with, internal anal sphincter dysfunction in Japanese adults. **Methods:** Anorectal manometry was performed in 1193 women and 1124 men aged 20 years or older. The maximal resting pressure, measured by a rapid pull-through technique, was defined as the highest resting pressure recorded. Internal anal sphincter dysfunction was defined as a maximal resting pressure less than 30 mmHg. Potential risk factors were assessed through self-reports, interviews, physical examinations, and medical record reviews. Multivariate logistic regression analysis was used to identify independent risk factors for internal anal sphincter dysfunction. **Results:** Significant differences in maximal resting pressure were seen between women (58.1 ± 24.9 mmHg) and men (68.8 ± 23.5 mmHg, $P < 0.001$). Maximal resting pressure decreased significantly with increasing age in both sexes. The prevalence of internal anal sphincter dysfunction was 10.4% (15.5% in women, 5.1% in men). In a multivariate logistic regression model, age, mental disease, pelvic organ prolapse repair, and fecal incontinence were independently associated with a greater risk of internal anal sphincter dysfunction in women and men. **Conclusions:** Internal anal sphincter dysfunction is a common problem for women and men. Several of the identified risk factors are preventable or modifiable, and may direct future research in fecal incontinence therapy.

Keywords: Fecal Incontinence; Internal Anal Sphincter; Anorectal Manometry; Maximal Resting Pressure

*Corresponding author.

1. INTRODUCTION

Fecal incontinence (FI) is a common disorder with a significant impact on quality of life. The prevalence of FI in the general population increases with age in both women and men [1]. Although the cause of FI is often multifactorial, FI is frequently caused by anal sphincter insufficiency [2]. The anal sphincter consist of the circular internal (IAS) and external (EAS) anal sphincter muscles together with the sling-shaped puborectalis muscle. Although both sphincter muscles are important for the maintenance of continence, the IAS, composed of smooth muscle arranged in oblique bundles, provides most of the resting anal tone and is the main muscle responsible for preventing fecal leakage. The IAS contributes an estimated 55% - 85% to maximal resting pressure (MRP) [3,4]. Low MRP is the most important predictor of FI and correlates with the Fecal Incontinence Severity Index (FISI) [5].

Despite a number of studies on FI in western countries [6-8], there has been a paucity of research about risk factors associated with IAS dysfunction (IASD). Identifying preventable or modifiable risk factors for IASD may guide future research for the prevention or treatment of FI.

The objective of this study was to provide a comprehensive description of IASD in Japanese adults and to describe demographic and other risk factors associated with IASD after multivariate adjustments.

2. MATERIALS AND METHODS

Between January 2006 and December 2008, 3190 consecutive subjects aged 20 years or older referred to our hospital were enrolled.

Factors potentially associated with IASD or FI were assessed by questionnaires, interviews, medical record reviews, and physical examinations. Participants filled in a structured questionnaire, which included questions about demographic characteristics (age, sex, body mass

index (BMI), employment), stool consistency, bowel movements, presence of selected medical conditions, previous anorectal surgeries, previous cholecystectomy, current habits, laxative use, FI, and parity in women. All medical conditions and medication use were determined according to self-reports as well as in-person interviews during which written responses were clarified. Patient weight and height were measured, and BMI was categorized according to the classification proposed for Japanese adults. The validated Bristol Stool Scale was used to describe the participant's usual stool consistency [9]. This scale consists of 7 types of stool and includes pictures of each stool type to aid participants in classifying their stools. We combined stool type 1 and 2 (hard and lumpy), 3 - 5 (normal consistencies), and 6 and 7 (mushy and watery). Smoking status was categorized as non-smoker and current smoker (≥ 1 cigarettes per day). Alcohol intake was categorized as nondrinker and current drinker (more than once a week). For this study, FI was defined as any involuntary loss of mucus, liquid, or solid stool during the last 30 days; this definition of FI did not include gas.

Anorectal manometry (ARM) was performed by a single experienced examiner using a 5 mm diameter, 1-channel solid-state catheter with a microtipped transducer ARM system (P-31, Star Medical Co., Tokyo, Japan). All subjects were examined in the left lateral position with the hips flexed to 90°. The lubricated catheter was introduced into the rectum. The MRP measured by means of a rapid pull-through technique was defined as the highest resting pressure recorded. IASD was defined as "low MRP" in cases where MRP was less than 30 mmHg [10].

Statistical analysis was performed using JMP version 9.0.2 (SAS Institute Inc, Cary, NC). MRP values were expressed as means \pm standard deviation. The Mann-Whitney U test was used to analyze for sex differences in MRP, and regression analysis was used to search for associations with age. The chi-square test was used to test the association between low MRP and each risk factor individually. All risk factors found to be significantly associated with low MRP in univariate analyses were combined into separate multivariate logistic models for women and men. Adjusted odds ratios (OR) and 95% confidence intervals (CI) were used to describe associations between risk factors and low MRP in univariate analyses and in multivariate analyses. P values < 0.05 were considered to indicate statistical significance for all analyses.

This study was a retrospective review of existing clinical data prospectively collected on a hospital anal physiology unit computer database. This study was approved by the research and ethics committee of Kunitomo Hospital and informed consent for ARM and future

use of their data was provided by all participants in the study.

3. RESULTS

Data from 2317 subjects (1193 women and 1124 men) who gave complete responses were analyzed. Mean age was 56.0 ± 17.9 years in women and 53.2 ± 16.1 years in men. All participants were stratified into 10-year age groups. The general characteristics of the participants are shown in **Table 1**. The most common medical condition was hypertension (26.8%). FI was reported by 15.1% of women and 7.7% of men.

Significant differences in MRP were seen between women (58.1 ± 24.9 mmHg) and men (68.8 ± 23.5 mmHg; $P < 0.001$). MRP decreased significantly with increasing age in both sexes (**Figure 1**).

The prevalence of low MRP was 10.4% (15.5% in women, 5.1% in men). Among women, prevalence increased from 1.7% for 20 - 29 year-olds up to 52.4% for 80 years or older. For women and men combined, the most common medical conditions associated with low MRP were cognitive impairment and pelvic organ prolapse repair (**Tables 2 and 3**).

In univariate analyses, variables associated with low MRP in women and men were age, employment status, hypertension, stroke, cognitive impairment, mental disease, ischemic heart disease, any history of cancer, pelvic organ prolapse repair, laxative use, and FI. In addition, parity, dyslipidemia, spinal cord injury, anal fistula surgery, and hysterectomy were associated with low MRP in women but not in men, while BMI, usual stool consistency, diabetes mellitus, hemorrhoidectomy, and rectal surgery were associated with low MRP in men alone.

In a multivariate logistic regression model, age, mental disease, pelvic organ prolapse repair, and FI were independently associated with a greater risk of low MRP in women and men. In addition, ischemic heart disease was associated with low MRP in women alone. By contrast, smoking or alcohol intake and diabetes mellitus were significantly negative associated with low MRP in women (**Tables 4 and 5**).

4. DISCUSSION

The present study investigated the influence of age, sex, and other possible risk factors for low MRP in a large number of Japanese adults spanning a wide age range. The main results are the following: 1) women had significantly lower MRP than men; 2) MRP decreased with increasing age in both sexes; 3) independent risk factors for low MRP in both sexes include advancing age, FI, mental disease, and pelvic organ prolapse repair.

FI is defined as either the involuntary passage of or the inability to control the discharge of fecal matter through

Table 1. General characteristics of study subjects.

Characteristic		Overall n = 2317		Women n = 1193		Men n = 1124		
Age (year)	20 - 29	201	(8.7)	117	(9.8)	84	(7.5)	
	30 - 39	348	(15.0)	149	(12.5)	199	(17.7)	
	40 - 49	339	(14.6)	157	(13.2)	182	(16.2)	
	50 - 59	470	(20.3)	216	(18.1)	254	(22.6)	
	60 - 69	421	(18.2)	231	(19.4)	190	(16.9)	
	70 - 79	391	(16.9)	220	(18.4)	171	(15.2)	
	≥80	147	(6.3)	103	(8.6)	44	(3.9)	
BMI	Underweight (<18.5)	140	(6.0)	87	(7.3)	53	(4.7)	
	Normal (18.5 - 24.9)	1628	(70.3)	898	(75.3)	730	(64.9)	
	Overweight (≥25)	549	(23.7)	208	(17.4)	341	(30.3)	
Parity	At least one vaginal			924	(77.4)			
Employment status	Employed/student	1281	(55.3)	470	(39.4)	811	(72.2)	
	Unemployed/other	1036	(44.7)	723	(60.6)	313	(27.8)	
Usual stool consistency	Normal stools	1403	(60.6)	635	(53.2)	768	(68.3)	
	Hard, lumpy stools	688	(29.7)	466	(39.1)	222	(19.8)	
	Loose, watery stools	150	(6.5)	47	(3.9)	103	(9.2)	
	Other	76	(3.3)	45	(3.8)	31	(2.8)	
Frequency of bowel movements	>21 BM/week	113	(4.9)	49	(4.1)	64	(5.7)	
	3 - 21 BM/week	2001	(86.4)	1001	(83.9)	1000	(89.0)	
	<3 BM/week	203	(8.8)	143	(12.0)	60	(5.3)	
Comorbid conditions	Hypertension	620	(26.8)	317	(26.6)	303	(27.0)	
	Diabetes mellitus	155	(6.7)	75	(6.3)	80	(7.1)	
	Dyslipidemia	257	(11.1)	154	(12.9)	103	(9.2)	
	Stroke	93	(4.0)	44	(3.7)	49	(4.4)	
	Cognitive impairment	29	(1.3)	17	(1.4)	12	(1.1)	
	Mental disease	102	(4.4)	60	(5.0)	42	(3.7)	
	Ischemic heart disease	64	(2.9)	27	(2.3)	37	(3.3)	
	Any history of cancer	108	(4.7)	58	(4.9)	50	(4.4)	
	Bronchial asthma	98	(4.2)	57	(4.8)	41	(3.6)	
	Spinal cord injury	46	(2.0)	22	(1.8)	24	(2.1)	
	Previous surgeries	Hemorrhoidectomy	314	(13.6)	156	(13.1)	158	(14.1)
		Anal fissure surgery	28	(1.2)	16	(1.3)	12	(1.1)
		Anal fistula surgery	82	(3.5)	16	(1.3)	66	(5.9)
Hysterectomy				97	(8.1)			
Pelvic organ prolapse repair		51	(2.2)	43	(3.6)	8	(0.7)	
Rectal surgery		16	(0.7)	7	(0.6)	9	(0.8)	
Cholecystectomy		43	(1.9)	17	(1.4)	26	(2.3)	
Current habits		Smoking	572	(24.7)	194	(16.3)	378	(33.6)
	Alcohol intake	979	(42.3)	293	(24.6)	686	(61.0)	
Laxative use		471	(20.3)	347	(29.1)	124	(11.0)	
Fecal incontinence		266	(11.5)	180	(15.1)	86	(7.7)	

BMI = body mass index. BM = bowel movements. Data are numbers of subjects with percentages in parentheses. Percentages may not sum to 100 because of rounding.




Figure 1. Regression analysis on the influence of age on maximal resting pressure (MRP) in women (a) and men (b). MRP decreased significantly with increasing age in both sexes (women, $r = -0.721$, $P < 0.001$; men, $r = -0.583$, $P < 0.001$).

the anal canal [1]. Clinically there are 3 subtypes: 1) passive incontinence—the involuntary discharge of rectal contents without awareness; 2) urge incontinence—the discharge of stool in spite of active attempts to retain bowel contents; and 3) combined incontinence—both passive and urge incontinence [2]. The majority of FI patients have passive incontinence [11]. Passive incontinence is generally associated with dysfunction of the smooth muscle of the IAS, whereas urge incontinence is related to dysfunction of the striated muscle of the EAS [1,2]. At rest, the IAS is in a tonically contracted state, and is innervated by the autonomic nervous system. The IAS plays a key role in maintaining continence, providing 55% - 85% of MRP [3,4].

Several studies have investigated the effect of sex on resting pressure of the anal canal [12-15]. The data are contradictory, with about half of the studies reporting comparable resting pressures in women and men, and the other half citing lower resting pressure in women. In most of these studies, the number of male and female subjects was rather low. Laurberg and Swash [12], who studied a larger cohort of subjects (102 women and 19 men), observed a significantly lower resting pressure in women. This is in accordance with the results of our study on more than 1000 people of both sexes.

Reports on the effects of aging on MRP are also conflicting. Some studies observed a significant lowering of MRP with age (more often in women than in men), while others found only a small decrease in MRP that did not reach statistical significance. Other studies did not see a decreased MRP in their older subjects [12-15]. In addition to the aforementioned problem of small sample size, most studies included patients of varying ages and grouped them differently. In the present study, we divided the

subjects into 7 age groups by decade, and used regression analysis to search for effects of age on MRP, and found that MRP significantly decreases with age in women as well as in men. This age-dependent decrease in MRP is thought to be due to fibrosis of the IAS in elderly patients [16,17].

Studies examining independent risk factors for FI have implicated age, obstetric and gynecologic factors, several medical conditions, and poor health. In particular, age, parity, vaginal delivery, diabetes, diarrhea, and neurologic conditions have notably been associated [6-8,18,19].

This study has several limitations that should be considered when interpreting the results. First, prevalence estimates were limited to non-institutionalized individuals, and because the prevalence of FI in the nursing home population is higher than in the community [1,2], the overall prevalence of IASD is likely underestimated, especially among older participants. Second, as in previous large epidemiological studies, comorbid conditions were determined according to self-reports instead of through clinician-derived methods.

Since women were found to have a higher prevalence of FI than men, in some prior studies, it was hypothesized that this sex difference is due to obstetrical injuries being a major risk factor for FI [18,19]. In this survey of the entire adult lifespan, there was a significant difference in the prevalence of low MRP between women and men. In univariate analysis, vaginal delivery was associated with a higher prevalence of low MRP in women, but this association was no longer significant after multivariate adjustment. Whitehead *et al.* [7] did not find a significant difference in the prevalence of FI between women and men comprising the entire adult lifespan. Other studies that included a broad range of ages have

Table 2. Baseline characteristics of the risks for low MRP of women.

Risk factor		Women n = 1193		Low MRP n = 185 (15.5)		P
Age (year)	20 - 29	117	(9.8)	2	(1.7)	<0.001
	30 - 39	149	(12.5)	4	(2.7)	
	40 - 49	157	(13.2)	5	(3.1)	
	50 - 59	216	(18.1)	9	(4.5)	
	60 - 69	231	(19.4)	42	(18.2)	
	70 - 79	220	(18.4)	69	(31.4)	
	≥80	103	(8.6)	54	(52.4)	
BMI	Underweight (<18.5)	87	(7.3)	20	(23.0)	0.13
	Normal (18.5 - 24.9)	898	(75.3)	135	(15.0)	
	Overweight (≥25)	208	(17.4)	30	(14.4)	
Parity	At least one vaginal	924	(77.4)	173	(18.7)	<0.001
	None	269	(22.6)	12	(4.5)	
Employment status	Employed/student	470	(39.4)	33	(7.0)	<0.001
	Unemployed/other	723	(60.6)	152	(21.0)	
Usual stool consistency	Normal stools	635	(53.2)	102	(16.1)	0.41
	Hard, lumpy stools	466	(39.1)	73	(15.7)	
	Loose, watery stools	47	(3.9)	7	(14.9)	
	Other	45	(3.8)	3	(6.7)	
Frequency of bowel movements	>21 BM/week	49	(4.1)	12	(24.5)	0.06
	3 - 21 BM/week	1001	(83.9)	145	(14.5)	
	<3 BM/week	143	(12.0)	28	(19.6)	
Comorbid conditions	Hypertension	317	(26.6)	81	(25.6)	<0.001
	Diabetes mellitus	75	(6.3)	16	(21.3)	0.15
	Dyslipidemia	154	(12.9)	34	(22.1)	0.01
	Stroke	44	(3.7)	13	(29.6)	0.009
	Cognitive impairment	17	(1.4)	10	(58.8)	<0.001
	Mental disease	60	(5.0)	17	(28.3)	0.005
	Ischemic heart disease	27	(2.3)	15	(55.6)	<0.001
	Any history of cancer	58	(4.9)	16	(27.6)	0.009
	Bronchial asthma	57	(4.8)	13	(22.8)	0.12
	Spinal cord injury	22	(1.8)	11	(50.0)	<0.001
Previous surgeries	Hemorrhoidectomy	156	(13.1)	29	(18.6)	0.25
	Anal fissure surgery	16	(1.3)	4	(25.0)	0.29
	Anal fistula surgery	16	(1.3)	6	(37.5)	0.01
	Hysterectomy	97	(8.1)	25	(25.8)	0.004
	Pelvic organ prolapse repair	43	(3.6)	25	(58.1)	<0.001
	Rectal surgery	7	(0.6)	3	(42.9)	0.08
	Cholecystectomy	17	(1.4)	5	(29.4)	0.11
Current habits	Smoking	194	(16.3)	5	(2.6)	<0.001
	Alcohol intake	293	(24.6)	7	(2.4)	<0.001
Laxative use		347	(29.1)	77	(22.2)	<0.001
Fecal incontinence		180	(15.1)	103	(57.2)	<0.001

MRP = maximal resting pressure. BMI = body mass index. BM = bowel movements. Data are numbers of subjects with percentages in parentheses. Percentages may not sum to 100 because of rounding.

Table 3. Baseline characteristics of the risks for low MRP of men.

Risk factor		Men n = 1124		Low MRP n = 57 (5.1)		P
Age (year)	20 - 29	84	(7.5)	1	(1.2)	<0.001
	30 - 39	199	(17.7)	2	(1.0)	
	40 - 49	182	(16.2)	1	(0.6)	
	50 - 59	254	(22.6)	3	(1.2)	
	60 - 69	190	(16.9)	11	(5.8)	
	70 - 79	171	(15.2)	26	(15.2)	
	≥80	44	(3.9)	13	(29.5)	
BMI	Underweight (<18.5)	53	(4.7)	5	(9.4)	0.01
	Normal (18.5 - 24.9)	730	(64.9)	44	(6.0)	
	Overweight (≥25)	341	(30.3)	8	(2.3)	
Employment status	Employed/student	811	(72.2)	17	(2.1)	<0.001
	Unemployed/other	313	(27.8)	40	(12.8)	
Usual stool consistency	Normal stools	768	(68.3)	27	(3.5)	0.005
	Hard, lumpy stools	222	(19.8)	20	(9.0)	
	Loose, watery stools	103	(9.2)	7	(6.8)	
	Other	31	(2.8)	3	(9.7)	
Frequency of bowel movements	>21 BM/week	64	(5.7)	6	(9.4)	0.27
	3 - 21 BM/week	1000	(89.0)	48	(4.8)	
	<3 BM/week	60	(5.3)	3	(5.0)	
Comorbid conditions	Hypertension	303	(27.0)	25	(8.6)	0.003
	Diabetes mellitus	80	(7.1)	8	(10.0)	0.04
	Dyslipidemia	103	(9.2)	6	(5.8)	0.71
	Stroke	49	(4.4)	6	(12.2)	0.02
	Cognitive impairment	12	(1.1)	7	(58.3)	<0.001
	Mental disease	42	(3.7)	8	(19.1)	<0.001
	Ischemic heart disease	37	(3.3)	5	(13.5)	0.02
	Any history of cancer	50	(4.4)	6	(12.0)	0.02
	Bronchial asthma	41	(3.6)	3	(7.3)	0.50
	Spinal cord injury	24	(2.0)	2	(8.3)	0.46
	Previous surgeries	Hemorrhoidectomy	158	(14.1)	19	(12.0)
Anal fissure surgery		12	(1.1)	1	(8.3)	0.61
Anal fistula surgery		66	(5.9)	6	(9.1)	0.13
Pelvic organ prolapse repair		8	(0.7)	4	(50.0)	<0.001
Rectal surgery		9	(0.8)	3	(33.3)	<0.001
Cholecystectomy		26	(2.3)	1	(3.8)	0.77
Current habits	Smoking	378	(33.6)	15	(4.0)	0.37
	Alcohol intake	686	(61.0)	25	(3.6)	0.007
Laxative use		124	(11.0)	19	(15.3)	<0.001
Fecal incontinence		86	(7.7)	26	(30.2)	<0.001

MRP = maximal resting pressure. BMI = body mass index. BM = bowel movements. Data are numbers of subjects with percentages in parentheses. Percentages may not sum to 100 because of rounding.

Table 4. Odds ratios for variables associated with low MRP of women.

Risk factor	Univariate analysis OR (95% CI)		P	Multivariate analysis OR (95% CI)		P
Age (10-year interval)	2.34	(2.04, 2.71)	<0.001	1.74	(1.44, 2.12)	<0.001
BMI (vs. normal)						
Underweight	1.69	(0.97, 2.82)	0.06	2.05	(0.92, 4.42)	0.08
Overweight	0.95	(0.61, 1.44)	0.82	0.70	(0.41, 1.16)	0.17
Parity (vs. none)	4.93	(2.82, 9.49)	<0.001	1.83	(0.87, 4.17)	0.11
Employment status (vs. employed/student)						
Unemployed/other	3.53	(2.40, 5.32)	<0.001	1.49	(0.86, 2.56)	0.15
Usual stool consistency (vs. normal stools)						
Hard, lumpy stools	0.97	(0.70, 1.34)	0.86	0.75	(0.46, 1.23)	0.26
Loose, watery stools	0.91	(0.37, 1.98)	0.83	0.93	(0.33, 2.83)	0.89
Other	0.37	(0.09, 1.05)	0.06	0.81	(0.17, 2.92)	0.77
Comorbid condition						
Hypertension	2.55	(1.84, 3.53)	<0.001	0.74	(0.46, 1.16)	0.19
Diabetes mellitus	1.52	(0.85, 2.71)	0.15	0.40	(0.18, 0.82)	0.01
Dyslipidemia	1.67	(1.10, 2.53)	0.01	1.41	(0.82, 2.38)	0.21
Stroke	2.38	(1.22, 4.64)	0.009	0.47	(0.19, 1.12)	0.09
Cognitive impairment	8.17	(3.07, 21.8)	<0.001	0.83	(0.24, 3.07)	0.78
Mental disease	2.27	(1.27, 4.08)	0.005	2.23	(1.00, 4.84)	0.049
Ischemic heart disease	7.32	(3.37, 15.9)	<0.001	3.14	(1.23, 8.22)	0.02
Any history of cancer	2.18	(1.20, 3.96)	0.009	1.11	(0.47, 2.50)	0.81
Spinal cord injury	5.73	(2.45, 13.4)	<0.001	2.51	(0.82, 7.58)	0.10
Previous surgeries						
Hemorrhoidectomy	1.29	(0.83, 2.00)	0.25	0.97	(0.55, 1.65)	0.91
Anal fistula surgery	3.35	(1.20, 9.32)	0.01	1.95	(0.48, 7.04)	0.33
Hysterectomy	2.03	(1.25, 3.30)	0.004	1.57	(0.82, 2.95)	0.17
Pelvic organ prolapse repair	8.59	(4.58, 16.1)	<0.001	2.48	(1.07, 5.84)	0.03
Rectal surgery	4.14	(0.80, 18.9)	0.08	1.11	(0.17, 6.75)	0.91
Current habits						
Smoking	0.12	(0.04, 0.27)	<0.001	0.25	(0.10, 0.56)	<0.001
Alcohol intake	0.10	(0.04, 0.20)	<0.001	0.27	(0.08, 0.72)	0.007
Laxative use	1.95	(1.41, 2.69)	<0.001	1.29	(0.81, 2.05)	0.29
Fecal incontinence	15.17	(10.5, 22.1)	<0.001	8.16	(5.20, 12.9)	<0.001

also failed to find a sex difference in FI prevalence or an effect of obstetrical injury [6,8,20]. This suggests that obstetrical injuries are not the most common cause of FI in women. Perineal tears are classified into 4 degrees. A fourth-degree tear is defined as an injury that extends through the fascia and musculature of the perineal body and involves the anal sphincter complex (EAS and IAS) and anal epithelium. Third-degree tears involve some or all of the fibers of the EAS, but do not necessarily involve the IAS. Faltin *et al.* [21] reported that clinically undetected tears of the anal sphincter were diagnosed by

anal endosonography in 42 of 150 women (28%). EAS injury alone was observed in 30 women (20%), IAS injury alone in 2 (1.3%), and both in 10 (7%). In a study of 62 women with FI related to obstetrical procedures, anal endosonography revealed EAS defects in 90% and IAS defects in 65% of the subjects [22]. Our data also suggests that EAS injury is the predominant cause of FI after vaginal delivery.

Previous studies have found diabetes mellitus to increase the risk of FI [8,23]. Ward and Tunuguntla [23] reported that 4% of diabetic patients suffered from FI.

Table 5. Odds ratios for variables associated with low MRP of men.

Risk factor	Univariate analysis OR (95% CI)		P	Multivariate analysis OR (95% CI)		P
Age (10-year interval)	2.54	(2.02, 3.29)	<0.001	2.36	(1.68, 3.41)	<0.001
BMI (vs. Normal)						
Underweight	1.62	(0.54, 3.94)	0.35	1.15	(0.36, 4.55)	0.82
Overweight	0.37	(0.16, 0.76)	0.006	0.49	(0.19, 1.13)	0.10
Employment status (vs. Employed/student)						
Unemployed/Other	6.84	(3.88, 12.6)	<0.001	1.10	(0.48, 2.49)	0.81
Usual stool consistency (vs. normal stools)						
Hard, lumpy stools	2.72	(1.48, 4.93)	0.002	1.94	(0.59, 5.68)	0.27
Loose, watery stools	2.00	(0.79, 4.48)	0.14	1.57	(0.68, 3.52)	0.29
Other	2.94	(0.67, 8.99)	0.13	2.30	(0.37, 10.3)	0.85
Comorbid condition						
Hypertension	2.22	(1.29, 3.81)	0.003	0.67	(0.31, 1.41)	0.30
Diabetes mellitus	2.26	(1.03, 4.95)	0.04	0.97	(0.33, 2.53)	0.96
Dyslipidemia	1.18	(0.49, 2.81)	0.71	0.86	(0.27, 2.29)	0.78
Stroke	2.80	(1.14, 6.89)	0.02	0.48	(0.13, 1.47)	0.21
Cognitive impairment	29.73	(9.11, 98.0)	<0.001	4.37	(0.97, 21.5)	0.06
Mental disease	4.96	(2.18, 11.3)	<0.001	7.29	(2.36, 21.2)	<0.001
Ischemic heart disease	3.11	(1.16, 8.31)	0.02	1.98	(0.53, 6.32)	0.29
Any history of cancer	2.74	(1.11, 6.72)	0.02	0.62	(0.16, 1.98)	0.44
Spinal cord injury	1.73	(0.40, 7.53)	0.46	1.40	(0.19, 5.98)	0.70
Previous surgeries						
Hemorrhoidectomy	3.34	(1.87, 5.96)	<0.001	1.55	(0.72, 3.21)	0.26
Anal fistula surgery	1.97	(0.81, 4.78)	0.13	2.02	(0.57, 6.02)	0.26
Pelvic organ prolapse repair	20.06	(4.88, 82.4)	<0.001	11.87	(1.71, 88.9)	0.01
Rectal surgery	9.82	(2.39, 40.3)	<0.001	5.94	(0.76, 43.9)	0.09
Current habits						
Smoking	0.69	(0.37, 1.24)	0.37	0.79	(0.65, 2.42)	0.50
Alcohol intake	0.47	(0.28, 0.82)	0.007	1.15	(0.53, 2.40)	0.72
Laxative use	4.58	(2.55, 8.23)	<0.001	1.57	(0.68, 3.51)	0.28
Fecal incontinence	14.08	(7.83, 25.23)	<0.001	5.13	(2.14, 12.12)	0.001

MRP = maximal resting pressure. OR = odds ratio. CI = confidence interval. BMI = body mass index. Only variables found to be significant in univariate analyses for women or men were included in the multivariate regressions and shown in the table. Multivariate odds ratios are adjusted for all other risk factors in the table.

The underlying pathophysiology of anorectal dysfunction in diabetic patients with FI is not well understood. Diabetes mellitus may contribute to FI through sphincter or pelvic floor weakness from anatomic defects, nerve damage, or microvascular complications. Schiller *et al.* [24] found IASD secondary to autonomic neuropathy responsible for FI in 16 diabetic patients. In contrast, Rogers [25] reported that EAS dysfunction might play an important role. Other reports suggested that both IAS and EAS dysfunction might be involved [26,27]. In the present study, the univariate association between diabetes mellitus and low MRP was significant for men, but after

multivariate adjustment, this relationship did not hold. On the contrary, in women low MRP was significantly negatively associated with diabetes mellitus after multivariate adjustment. The lumbar sympathetic innervation inhibits the colon and provides motor stimulation to the IAS. The pudendal nerves, on the other hand, are somatic nerves that innervate the EAS. Diabetic pudendal neuropathy leads to delay of pudendal nerve terminal motor latency, causes EAS damage, and may result in FI [28]. Our results suggest that somatic neuropathy plays a more important role in FI in diabetic patients.

FI was significantly associated with low MRP after

multivariate adjustment in both sexes. More treatment options for FI makes a proper pretreatment evaluation increasingly important, whether by means of subjective severity questionnaires, such as the FISFI, or by objective measurements, such as ARM, anal endosonography, electromyography, and defecography [1]. Bordeianou *et al.* [5] reported MRP was the only objective measurement that seems to correlate with both FISFI and the presence of sphincteric defects on anal endosonography, while maximal squeeze pressure (MSP) generated by the EAS was not correlated. Our results confirm a strong association between low MRP and FI.

Mental disease was found to be independently associated with low MRP. The IAS has spontaneous myogenic tone, but it also receives inputs from the sympathetic and parasympathetic nervous systems, both of which exert excitatory and inhibitory influences on the sphincteric smooth muscle [29]. Yamato and Rattan [30] demonstrated that stimulation of $\alpha 1$ -adrenoreceptors causes an increase in the resting pressure developed by the IAS. Multi-acting receptor targeted antipsychotics (MARTA) have a high affinity for a number of receptors including the dopaminergic D2, serotonergic 5-HT_{2A}, $\alpha 1$ -adren-ergic, and muscarinic receptors. Two cases of FI have been reported in patients taking MARTA [31,32]. The pathophysiology is thought to be related to its central mechanism of action and inhibitory effects on the anal sphincters. Siproudhis *et al.* [33] reported that a single oral administration of 2 types of antidepressants (amitriptyline and fluoxetine) reduced pressure in the upper anal canal in 10 healthy male volunteers. Both drugs are likely to decrease the contractile activity of smooth muscle by inhibiting calcium channels.

Pelvic organ prolapse repair was also an independent risk factor for low MRP. Most studies reported a greater than 10% prevalence of FI in women with genital prolapse, and more than a third of women and men with rectal prolapse reported FI [34]. Internal rectal prolapse is a circular infolding of the rectal wall that occurs on straining to defecate. It descends into the rectum and may reach into the anal canal. Harmston *et al.* [35] found a significant reduction in MRP with progressive increases in the grade of internal rectal prolapse without a demonstrable effect on MSP. By contrast, external rectal prolapse was associated with a significant reduction in both MRP and MSP [36]. The underlying mechanism has been postulated to result from nerve damage by stretching due to perineal descent or sphincter dilatation and inappropriate stimulation of the recto-anal inhibitory reflex by the prolapsing rectum [37]. The finding of decreased MRP without a significant change in MSP suggests that the predominant effect of internal rectal prolapse is on reducing IAS tone without an effect on EAS.

5. CONCLUSION

IASD has received little attention, and the results of this study indicate that it is a major problem for elderly people in Japan. In addition, the importance of IASD as a public health problem is likely to increase in the near future as the elderly population continues to grow. This cross-sectional study offers evidence that IASD is correlated with the presence of certain conditions such as age, sex, FI, mental disease, and pelvic organ prolapse. Several of these risk factors for IASD are potentially preventable or modifiable; for example, several studies have shown a recovery in MRP after rectal prolapse repair [35]. Improving our understanding of risk factors, particularly modifiable risk factors, is critical for developing future prevention guidelines and improving the specificity of FI treatment.

REFERENCES

- [1] Madoff, R.D., Parker, S.C., Varma, M.G., *et al.* (2004) Faecal incontinence in adults. *Lancet*, **364**, 621-632. [doi:10.1016/S0140-6736\(04\)16856-6](https://doi.org/10.1016/S0140-6736(04)16856-6)
- [2] Rao, S.S.C. (2004) Diagnosis and management of fecal incontinence. *American Journal of Gastroenterology*, **99**, 1585-1604. [doi:10.1111/j.1572-0241.2004.40105.x](https://doi.org/10.1111/j.1572-0241.2004.40105.x)
- [3] Frenckner, B. and Euler, C.V. (1975) Influence of pudendal block on the function of the anal sphincters. *Gut*, **16**, 482-489. [doi:10.1136/gut.16.6.482](https://doi.org/10.1136/gut.16.6.482)
- [4] Barnett, J.L., Hasler, W.L. and Camilleri, M. (1999) American gastroenterological association medical position statement on anorectal testing techniques. *Gastroenterology*, **116**, 732-760. [doi:10.1016/S0016-5085\(99\)70194-0](https://doi.org/10.1016/S0016-5085(99)70194-0)
- [5] Bordeianou, L., Lee, K.Y., Rockwood, T., *et al.* (2008) Anal resting pressures at manometry correlate with the fecal incontinence severity index and with presence of sphincter defects on ultrasound. *Disease of Colon and Rectum*, **51**, 1010-1014. [doi:10.1007/s10350-008-9230-7](https://doi.org/10.1007/s10350-008-9230-7)
- [6] Varma, M.G., Brown, J.S., Creasman, J.M., *et al.* (2006) Fecal incontinence in females older than aged 40 years: Who is at risk? *Disease of Colon and Rectum*, **49**, 841-851. [doi:10.1007/s10350-006-0535-0](https://doi.org/10.1007/s10350-006-0535-0)
- [7] Whitehead, W.E., Borrud, L., Goode, P.S., *et al.* (2009) Fecal incontinence in US adults: Epidemiology and risk factors. *Gastroenterology*, **137**, 512-517. [doi:10.1053/j.gastro.2009.04.054](https://doi.org/10.1053/j.gastro.2009.04.054)
- [8] Joh, H.K., Seong, M.K. and Oh, S.W. (2010) Fecal incontinence in elderly Koreans. *Journal of the American Geriatrics Society*, **58**, 116-121. [doi:10.1111/j.1532-5415.2009.02613.x](https://doi.org/10.1111/j.1532-5415.2009.02613.x)
- [9] Lewis, S.J. and Heaton, K.W. (1997) Stool form scale as a useful guide to intestinal transit time. *Scandinavian Journal of Gastroenterology*, **32**, 920-924. [doi:10.3109/00365529709011203](https://doi.org/10.3109/00365529709011203)
- [10] Rao, S.S. and Patel, R.S. (1997) How useful are manometric tests of anorectal function in the management of

- defecation disorders? *American Journal of Gastroenterology*, **92**, 469-475.
- [11] Abe, T., Kunimoto, M. and Hachiro, Y. (2008) Diagnosis and management of fecal incontinence at a specialty outpatient clinic. *Journal of Japan Society of Coloproctology*, **61**, 247-253. doi:10.3862/jcoloproctology.61.247
- [12] Laurberg, S. and Swash, M. (1989) Effects of aging on the anorectal sphincters and their innervation. *Disease of Colon and Rectum*, **32**, 737-742. doi:10.1007/BF02562120
- [13] Enck, P., Kuhlbusch, R., Lübke, H., *et al.* (1989) Age and sex and anorectal manometry in incontinence. *Disease of Colon and Rectum*, **32**, 1026-1030. doi:10.1007/BF02553874
- [14] Rao, S.S., Hatfield, R., Soffer, E., *et al.* (1999) Manometric tests in anorectal function in healthy adults. *American Journal of Gastroenterology*, **94**, 773-783. doi:10.1111/j.1572-0241.1999.00950.x
- [15] Gundling, F., Seidl, H., Scalercio, N., *et al.* (2009) Influence of gender and age on anorectal function: Normal values from anorectal manometry in a large Caucasian population. *Digestion*, **81**, 207-213. doi:10.1159/000258662
- [16] Speakman, C.T.M., Hoyle, C.H.V., Kamm, M.A., *et al.* (1995) Abnormal internal anal sphincter fibrosis and elasticity in fecal incontinence. *Disease of Colon and Rectum*, **38**, 407-410. doi:10.1007/BF02054231
- [17] Abe, T., Sato, Y., Kunimoto, M., *et al.* (2008) Effect of aging and gender on internal anal sphincter thickness. *Anti-Aging Medicine*, **5**, 46-48. doi:10.3793/jaam.5.46
- [18] MacLennan, A.H., Taylor, A.W., Wilson, D.H., *et al.* (2000) The prevalence of pelvic floor disorders and their relationship to gender, age, parity and mode of delivery. *British Journal of Obstetrics and Gynaecology*, **107**, 1460-1470. doi:10.1111/j.1471-0528.2000.tb11669.x
- [19] Nelson, R., Norton, N., Cautley, E., *et al.* (1995) Community-based prevalence of anal incontinence. *Journal of the American Medical Association*, **274**, 559-561. doi:10.1001/jama.274.7.559
- [20] Bharucha, A.E., Zinsmeister, A.R., Locke, G.R., *et al.* (2006) Risk factors for fecal incontinence: A population-based study in women. *American Journal of Gastroenterology*, **101**, 1305-1312. doi:10.1111/j.1572-0241.2006.00553.x
- [21] Faltin, D.L., Boulvain, M., Irion, O., *et al.* (2000) Diagnosis of anal sphincter tears by postpartum endosonography to predict fecal incontinence. *Obstetrics and Gynecology*, **95**, 643-647. doi:10.1016/S0029-7844(99)00631-6
- [22] Burnett, S.J., Spence-Jones, C., Speakman, C.T., *et al.* (1991) Unsuspected sphincter damage following childbirth revealed by anal endosonography. *British Journal of Radiology*, **64**, 225-227. doi:10.1259/0007-1285-64-759-225
- [23] Ward, A. and Tunuguntla, A.K. (1984) Anorectal sensorimotor dysfunction in fecal incontinence and diabetes mellitus. *New England Journal of Medicine*, **310**, 1282-1287. doi:10.1056/NEJM198405173102003
- [24] Schiller, L.R., Santa Ana, C.A., Schmulen, C., *et al.* (1982) Pathogenesis of fecal incontinence in diabetes mellitus: Evidence for internal anal sphincter dysfunction. *New England Journal of Medicine*, **307**, 1666-1671. doi:10.1056/NEJM198212303072702
- [25] Rogers, J., Levy, D.M., Henry, M.M., *et al.* (1988) Pelvic floor neuropathy: A comparative study of diabetes mellitus and idiopathic faecal incontinence. *Gut*, **29**, 756-761. doi:10.1136/gut.29.6.756
- [26] Pintor, M.P., Zara, G.P., Falletto, E., *et al.* (1994) Pudendal neuropathy in diabetic patients with faecal incontinence. *International Journal of Colorectal Disease*, **9**, 105-109. doi:10.1007/BF00699423
- [27] Erckenbrecht, J.F., Winter, H.J., Cicmir, I., *et al.* (1988) Faecal incontinence in diabetes mellitus: Is it correlated to diabetic autonomic or peripheral neuropathy? *Zeitschrift für Gastroenterologie*, **26**, 731-736.
- [28] Watanabe, M., Tsunoda, A., Kamiyama, G., *et al.* (2003) Pathophysiology in diabetic patients with fecal incontinence. *Showa University Journal of Medical Sciences*, **15**, 21-26.
- [29] Mills, K. and Chess-Williams, R. (2009) Pharmacology of the internal anal sphincter and its relevance to faecal incontinence. *Autonomic and Autacoid Pharmacology*, **29**, 85-95. doi:10.1111/j.1474-8673.2009.00437.x
- [30] Yamato, S. and Rattan, S. (1990) Role of alpha adrenoreceptors in opossum internal anal sphincter. *Journal of Clinical Investigation*, **86**, 424-429. doi:10.1172/JCI114728
- [31] Mendhekar, D.N., Srivastav, P.K., Sarin, S.K., *et al.* (2003) A case report of olanzapine-induced fecal incontinence. *Journal of Clinical Psychiatry*, **61**, 601-602.
- [32] Sagar, R., Varghese, S.T. and Balhara, Y.P. (2005) Olanzapine-induced double incontinence. *Indian Journal of Medical Sciences*, **59**, 163-164. doi:10.4103/0019-5359.16123
- [33] Siproudhis, L., Dinasquet, M., Sébille, V., *et al.* (2004) Differential effects of two types of antidepressants, amitriptyline and fluoxetine, on anorectal motility and visceral perception. *Alimentary Pharmacology and Therapeutics*, **20**, 689-695. doi:10.1111/j.1365-2036.2004.02151.x
- [34] Shamliyan, T., Wyman, J., Bliss, D.Z., *et al.* (2007) Prevention of urinary and fecal incontinence in adults. *Evidence Reports/Technology Assessments*, **161**, 1-379.
- [35] Harmston, C., Jones, O.M., Cunningham, C., *et al.* (2011) The relationship between internal rectal prolapse and internal anal sphincter function. *Colorectal Disease*, **13**, 791-795. doi:10.1111/j.1463-1318.2010.02266.x
- [36] Hiltunen, K.M., Matikainen, M., Auvinen, O., *et al.* (1986) Clinical and manometric evaluation of anal sphincter function in patients with rectal prolapse. *American Journal of Surgery*, **151**, 489-492. doi:10.1016/0002-9610(86)90110-8
- [37] Furouk, R., Duthie, G.S., MacGregor, A.B., *et al.* (1994) Recto-anal inhibition and incontinence in patients with rectal prolapse. *British Journal of Surgery*, **81**, 743-746. doi:10.1002/bjs.1800810542