



RNA-Seq Analysis in Fruit Science: A Review

Ozhan Simsek^{1,*}, Dicle Donmez², Yildiz Aka Kacar^{1,2}

¹Faculty of Agriculture, Department of Horticulture, Cukurova University, Adana, Turkey

²Institute of Natural and Applied Sciences, Department of Biotechnology, Cukurova University, Adana, Turkey

Email address:

ozhan12@gmail.com (O. Simsek), dicleonmez4@gmail.com (D. Donmez), ykacar@cu.edu.tr (Y. A. Kacar)

*Corresponding author

To cite this article:

Ozhan Simsek, Dicle Donmez, Yildiz Aka Kacar. RNA-Seq Analysis in Fruit Science: A Review. *American Journal of Plant Biology*. Special Issue: Plant Molecular Biology and Biotechnology. Vol. 2, No. 3-1, 2017, pp. 1-7. doi: 10.11648/j.ajpb.s.2017020301.11

Received: February 11, 2017; **Accepted:** February 14, 2017; **Published:** February 28, 2017

Abstract: Fruit breeding is an ancient technology with dynamic current techniques and an exciting future. There are a number of restraints to conventional fruit breeding which are especially limiting in tree fruits with their long juvenile period, large plant size, and which are represented by unique, highly-selected heterozygous genotypes. Biotechnology offers to minimize disadvantages of classical breeding techniques. In this sense, fruit breeding refers to the purposeful genetic improvement of fruit crops through various techniques including selection, hybridization, mutation induction, and molecular techniques. Among molecular techniques, sequencing technology have been used for many years and recently a new concept titled “RNA-Seq” have been started to performed to understand molecular mechanisms in fruits. RNA-Seq analysis is an effective tool to understand which genes involved and expressed in different mechanisms and organs/cells of a plant. Recently, many articles have been published using RNA-Seq in fruits. In the present review, we illustrated how to apply different RNA-Seq platforms in fruits with examples.

Keywords: Sequencing, Illumina, Gene, Breeding, Fruit

1. Introduction

1.1. Importance of Fruit Science

Among the horticultural plants; fruits are the primary source of essential minerals and vitamins. Fruits will continue to expand due to demand of better tasting, more varied and nutritious. Reproductive behavior in plant species is a crucial part of ecosystem functioning. Therefore, patterns of production of fruit have received much attention from a wide range of ecologists [1, 2]. How do we keep fruit production on par with the burgeoning population? Although conventional plant breeding techniques have made considerable progress in the development of improved varieties, they have not been able to keep pace with the increasing demand for vegetables and fruits in the developing countries. Therefore an immediate need is felt to integrate biotechnology to speed up the crop improvement programs. Especially modern biotechnological methods have been taken part in breeding strategies of fruit crops in addition to classical plant breeding strategies. Several biotechnological methods can be applied to plant to have better ones in the

process of fruit breeding [3].

RNA-Seq analysis is a magic method in order to understand which genes involved and expressed in different mechanisms and organs/cells of a plant. Recently, many article have been published related to RNA-Seq in fruits. In the present review, we illustrated how to apply RNA-Seq in fruits with examples.

1.2. RNA-Seq Technology

High-throughput sequencing of cDNA fragment populations, commonly known as RNA-Seq. RNA-Seq is a powerful tool for transcriptome analysis and uses deep-sequencing technologies to produce millions of short cDNA reads. The resulting reads are either aligned to a reference genome or reference transcripts, or assembled *de novo* (without the genomic sequence) to produce a genome-scale transcription map that consists of both the transcript structure and level of expression for each gene at any particular developmental stage [4, 5, 6]. RNA-Seq has been applied successfully in transcriptome profiling of species without genome sequencing data [7, 8].

The recent RNA-Seq based on NGS (next-generation

sequencing) enables studies to be carried out on species without corresponding sequenced genome information as a reference [9]. It has become widely applied to model as well as non-model organisms to obtain mass sequence data for molecular marker development, gene discovery and transcriptional analysis [9, 10]. Compared with traditional laboratory methods, RNA-Seq is a high throughput technology, overcoming the weakness of microarrays in exploring unknown genes. Furthermore, it has great advantages in examining transcriptome fine structure, such as detection of allele-specific expression and splice junction variation [11]. This powerful technology, available since only a couple of years, is already making substantial contributions towards the understanding of genome expression and regulation in living organisms [7, 12, 13]. This technology allows us to survey multiple levels of natural variation at unprecedented resolution [14] and can be very useful in the assessment of alternative splicing and the detection of novel gene structures [15]. Werner [16] indicated the great potential contribution of this technology to functional genomics with a special focus on gene regulation by transcription factor binding rates [17].

In principle, RNA-Seq allows analysis of all expressed transcripts, with three key goals: (i) annotating the structures of all transcribed genes including their 5' and 3' ends and all splice junctions [18, 19], (ii) quantifying expression of each transcript [20, 21] and (iii) measuring the extent of alternative splicing [5, 22, 23]. Many methods have been recently developed for strand-specific RNA-Seq, and they fall into two main classes. One class relies on attaching different adaptors in a known orientation relative to the 5' and 3' ends of the RNA transcript. These protocols generate a cDNA library flanked by two distinct adaptor sequences, marking the 5' end and the 3' end of the original mRNA. A second class of methods relies on marking one strand by chemical modification, either on the RNA itself by bisulfite treatment or during second-strand cDNA synthesis followed by degradation of the unmarked strand [23]. For RNA-Seq four high-throughput DNA sequencing technologies (Roche 454, Illumina, Helicos BioSciences and Life Technologies) can be used, and new technologies are in development by others. Among these methods, the most widely used system to generate RNA-Seq data is the Illumina system mainly due to the cheaper cost per base sequenced. Given the importance of sequencing capabilities, such as throughput, read length, error rate and ability to perform paired reads, for RNA-Seq as well as genomic studies, NGS companies are constantly improving their platforms to provide the best sequencing performance at the lowest cost [24, 25]. The GS instrument was introduced in 2005, developed by 454 Life Sciences, as the first next-generation system on the market. The method has recently increased the achieved reading length to the 400–500 base range, with paired-end reads, and as such is being applied to genome (bacterial, animal, human) sequencing. One spectacular application of the system was the identification of the culprit in the recent honey-bee disease epidemics. A relatively high cost of operation and

generally lower reading accuracy in homopolymer stretches of identical bases are mentioned presently as the few drawbacks of the method. The next upgrade 454 FLX Titanium will quintuple the data output from 100 Mb to about 500 Mb, and the new picotiter plate in the device uses smaller beads about 1 mm diameter. The device, schema of operation, its further developments and list of publications with applications can be found at <http://www.454.com/index.asp> [26, 27].

The Illumina Genome Analyser produces over 100 million short reads (35–76 bases, depending on the sequencing chemistry) leading to 3–6 gigabases of sequencing data in one run. A megabase costs about US\$4. We refer to this technology as a short read technology. It is based on solid-phase bridge PCR and uses a 'sequencing by synthesis' approach, with fluorescent dye-labelled terminator nucleotides. It uses fragmented double-stranded DNA as template. Fragments of up to 10 kb can be used for the construction of paired end sequencing libraries. The technology is also referred to as Solexa sequencing. The accuracy of the produced sequence data is greater 98.5% (<http://www.solexa.com>) [24, 28].

The Helicos Genetic Analysis System (<http://www.helicosbio.com>) was the first commercially available single-molecule sequencing system (SMS) on the market [29]. Helicos method produces average read lengths of 35 bp across 600 million to 1 billion reads, totaling 21 – 35 Gb per run at a rate of > 1 Gb/h. The Helicos platform lends itself well to multiplexing with up to 96 samples per channel or 4800 samples per run (<http://www.helicosbio.com>) [30].

The Applied Biosystems SOLiD system is based on emulsion PCR in combination with sequencing by ligation with dye-labelled oligonucleotides [31]. It produces up to one billion short reads (up to 50 bases) per run, leading to a total sequence output of up to 30 gigabases per single read run. As templates it uses fragmented double-stranded DNA. Fragment sizes for the construction of paired end sequencing libraries can be up to 10 kb. The sequences produced are 99.94% accurate (http://www3.appliedbiosystems.com/AB_Home) [24, 28].

RNA-Seq technology represents the latest and most powerful tool for characterizing transcriptomes [32]. RNA-Seq is its ability to provide information on transcripts that are expressed at very low levels, limited only by the total number of reads that are generated [33]. Because RNA-Seq is performed using tagged libraries of short cDNAs, prepared from fragmented or unfragmented RNA, it does not require prior knowledge of the sequences to be profiled [17].

2. Instances of RNA-Seq Analysis in Fruit Crops

Many RNA-Seq articles have been recently published in fruits to investigate genes expressed and involved in different mechanisms. RNA-Seq is also good tool to understand expression level of genes. Different RNA-Seq platforms have

been released for several years. These platforms were used for RNA sequencing in fruit species.

A comparative transcriptomic analysis between HW (Hot Water) treated and non-treated tomato fruit before and after cold storage was carried out. In the study massive sequencing were performed on a 5500 SOLID System (with Exact Call Chemistry module). RNA-Seq analysis detected a large number of differentially expressed genes that ranged from 2235 (heat shock) to 5433 (cold storage). Three clusters of genes were identified after 2 weeks of cold storage: the chilling-response included the down-regulation of genes involved in photo-synthesis, metabolism of cell wall, lipid and ethylene, as well as the up-regulation of genes for trehalose synthesis and transcription factors (DOF and MYB); the chilling-susceptibility was associated with the down-regulation of genes involved in carotenoid biosynthesis, which correlates with the main CI symptom of uneven ripening; meanwhile, the chilling-tolerance was related to the up-regulation of genes for heat stress (heat shock proteins and heat shock transcription factors) and detoxification (glutathione S-transferases) [34].

In another study, in order to understand the systemic effect of the fungal symbiosis on the tomato fruit, Zouari et al. [35] used RNA-Seq to perform global transcriptome profiling on Moneymaker tomato fruits at the turning ripening stage. In this study fruits were collected at 55 days after flowering, from plants colonized with *Funneliformis mosseae* and from control plants, which were fertilized to avoid responses related to nutrient deficiency. RNA-Seq analysis were performed using the Illumina Genome Analyzer (Solexa). Transcriptome analysis identified 712 genes that are differentially expressed in fruits from mycorrhizal and control plants. Gene Ontology (GO) enrichment analysis of these genes showed 81 overrepresented functional GO classes. Up-regulated GO classes include photosynthesis, stress response, transport, amino acid synthesis and carbohydrate metabolism functions, suggesting a general impact of fungal symbiosis on primary metabolisms and, particularly, on mineral nutrition. Down-regulated GO classes include cell wall, metabolism and ethylene response pathways.

De novo RNA-Seq of ripe blackberries grown under field conditions was performed using Illumina Hi-Seq 2000. Almost 9 billion nucleotide bases were sequenced in total. Following assembly, 42,062 consensus sequences were detected. For functional annotation, 33,040 (NR), 32,762 (NT), 21,932 (Swiss-Prot), 20,134 (KEGG), 13,676 (COG), 24,168 (GO) consensus sequences were annotated using different databases; in total 34,552 annotated sequences were identified [6]. Gupta et al. [36] performed high-throughput transcriptome sequencing (RNA-Seq) and differential gene expression analysis of five stages of berry development and ripening. RNA-Seq expression profiling showed that blueberry growth, maturation, and ripening involve dynamic gene expression changes, including coordinated up- and down-regulation of metabolic pathway enzymes and transcriptional regulators. Analysis of RNA-Seq alignments

identified developmentally regulated alternative splicing, promoter use, and 3' end formation. Kim et al. [37] optimized a procedure for high-quality RNA isolation from vegetative and reproductive tissues of climacteric and non-climacteric plum cultivars and conducted high-throughput transcriptomics. Researchers were identified 20 candidate reference genes from significantly non-differentially expressed transcripts of RNA-Seq data and verified their expression stability using qRT-PCR on a total of 141 plum samples which included flesh, peel, and leaf tissues of several cultivars collected from three locations over a 3-year period.

Fruit skin color is one of the most important traits for both the commercial and esthetic value of strawberry. Anthocyanins are the most prominent pigments in strawberry that bring red, pink, white, and yellow hues to the fruits in which they accumulate. Zhang et al. [8] conducted a *de novo* assembly of the fruit transcriptome of woodland strawberry and compared the gene expression profiles with yellow (Yellow Wonder, YW) and red (Ruegen, RG) fruits. The cDNA library was sequenced on the Illumina sequencing platform (Hi-Seq™ 2500). *De novo* assembly yielded 75,426 unigenes, 21.3% of which were longer than 1,000 bp. Among the high-quality unique sequences, 45,387 (60.2%) had at least one significant match to an existing gene model. A total of 595 genes, representing 0.79% of total unigenes, were differentially expressed in YW and RG. Among them, 224 genes were up-regulated and 371 genes were down-regulated in the fruit of YW.

Wang et al. [38] investigated to facilitate isolation of genes controlling important horticultural traits of peach, transcriptome sequencing was conducted. The cDNA libraries were sequenced using Illumina Hi-Seq2000 sequencer. A total of 133 million pair-end RNA-Seq reads were generated from leaf, flower, and fruit, and 90% of reads were mapped to the peach draft genome. Sequence assembly revealed 1,162 transcription factors and 2,140 novel transcribed regions (NTRs). Of these 2,140 NTRs, 723 contain an open reading frame, while the rest 1,417 are non-coding RNAs.

Liu et al. [39] were reported that Solexa sequencing has been used to discover small RNA populations and compare miRNAs on genome-wide scale in watermelon grafting system. A total of 11,458,476, 11,614,094 and 9,339,089 raw reads representing 2,957,751, 2,880,328 and 2,964,990 unique sequences were obtained from the scions of self-grafted watermelon and watermelon grafted on-to bottle gourd and squash at two true-leaf stage, respectively. 39 known miRNAs belonging to 30 miRNA families and 80 novel miRNAs were identified in our small RNA dataset. Compared with self-grafted watermelon, 20 (5 known miRNA families and 15 novel miRNAs) and 47 (17 known miRNA families and 30 novel miRNAs) miRNAs were expressed significantly different in watermelon grafted on to bottle gourd and squash, respectively. miRNAs expressed differentially when watermelon was grafted onto different rootstocks, suggesting that miRNAs might play an important

role in diverse biological and metabolic processes in watermelon and grafting may possibly by changing miRNAs expressions to regulate plant growth and development as well as adaptation to stresses. Feng *et al.* [40] were designed to obtain transcript sequence data and examine gene expression in bayberry developing fruit based on RNA-Seq and bioinformatic analysis, to provide a foundation for understanding the molecular mechanisms controlling fruit quality changes during ripening. Four RNA samples containing various tissues and fruit of different development and ripening stages were sequenced using the latest Illumina deep sequencing technique. RNA-Seq generated 1.92 G raw data, which was then *de novo* assembled into 41.239 UniGenes with a mean length of 531 bp. Approximately 80% of the UniGenes (32.805) were annotated against public protein databases, and coding sequences (CDS) of 31.665 UniGenes were determined. Over 3.600 UniGenes were differentially expressed during fruit ripening, with 826 up-regulated and 1.407 down-regulated. GO comparisons between the UniGenes of these two types and interactive pathways (Ipath) analysis found that energy-related

metabolism was enhanced, and catalytic activity was increased. All genes involved in anthocyanin biosynthesis were up-regulated during the fruit ripening processes, concurrent with color change.

Zhang *et al.* [41] investigated to identify novel as well as conserved miRNAs in citrus, deep sequencing of small RNA library combined with microarray was performed in precocious trifoliolate orange (an early flowering mutant of trifoliolate orange, *Poncirus trifoliata* L. Raf.). Illumina technology were used in sequencing analysis. A total of 114 conserved miRNAs belonging to 38 families and 155 novel miRNAs were determined. The miRNA star sequences of 39 conserved miRNAs and 27 novel miRNAs were also discovered among newly identified miRNAs, providing additional evidence for the existence of miRNAs.

RNA-Seq analysis could be also performed for molecular marker developments. There are some articles in fruits in order to develop molecular markers (Pomegranate [42], (Pomelo) [43], (Banana) [44].

Some recent examples by using RNA-Seq technologies, and mechanisms for fruits were presented Table 1.

Table 1. Some examples by using RNA-Seq technologies in fruits.

Species	Mechanism	RNA-Seq Technology	References
Sweet Orange (<i>Citrus sinensis</i>)	Fruit ripening	Illumina	[45]
Sweet Orange (<i>Citrus sinensis</i>)	Heterozygous	Solid Bioscope	[46]
Mango (<i>Mangifera indica</i> L.)	Characterization of transcriptome and chloroplast genome	Illumina Hi-Seq 2000	[47]
Pomegranate (<i>Punica granatum</i> L.)	Molecular Marker Development	454-GS-FLX Titanium	[42]
Mango (<i>Mangifera indica</i> Linn)	Fruit development and ripening	Illumina	[48]
Apple (<i>Malus communis</i>)	Transcriptome sequencing	Illumina Hi-Seq 2000	[49]
Trifoliolate orange <i>Poncirus trifoliata</i> (L.)	Phosphorus deficiency	Illumina Hi-Seq 2000	[50]
Apple (<i>Malus xiaojinensis</i>)	Iron tolerance	Illumina Hi-Seq 2000 and 454-GS-FLX Titanium	[51]
Cheery (<i>Prunus avium</i> L.)	Fruit development and exocarp	Illumina Hi-Seq 2000	[52]
Sweet Orange (<i>Citrus sinensis</i>)	Fruit ripening	Illumina Hi-Seq 2000	[53]
Sweet Orange (<i>Citrus sinensis</i>)	Fruit ripening	Illumina Hi-Seq 2000	[54]
Nectarine (<i>Prunus persica</i>)	Cold tolerance under storage conditions	Illumina Hi-Seq 2000	[55]
Xiangshui lemon (<i>Citrus limon</i> (L.) Burm. F)	Self-incompatibility	Illumina Hi-Seq 2000	[8]
Cherry (<i>Prunus avium</i> L.)	Biosynthesis of anthocyanin	Illumina/Solexa	[56]
Sweet Orange (<i>Citrus sinensis</i>)	Chitosan and salicylic acid applications	Illumina/Solexa	[57]
Apple (<i>Malus sieversii</i> v. niedzwetzkyana X <i>Malus domestica</i>)	Color of fruits	Illumina Hi-Seq 2000	[58]
Pummelo (<i>Citrus grandis</i>)	Pericarp colors and contents	Illumina	[59]
Apple (<i>Malus sieversii</i> v. niedzwetzkyana)	Biosynthesis of anthocyanin	Illumina Hi-Seq 2000	[60]
Pummelo (<i>Citrus grandis</i>)	Development of Molecular Markers	Illumina	[43]
Almond (<i>Prunus mongolica</i> maxim)	Drought stress	Illumina Hi-Seq 2000	[61]
Cherry (<i>Prunus pseudocerasus</i>)	Gene expression in seeds	Illumina	[62]
Trifoliolate orange (<i>Poncirus trifoliata</i> (L.) Raf.)	Cold stress	Illumina	[63]
Apple (<i>Malus x domestica</i>)	Russeting in exocarp	Illumina	[64]
Apple (<i>Malus domestica</i>)	Sorbitol synthesis	Illumina Hi-Seq 2000	[65]
Banana (<i>Musa balbisiana</i>)	Transcriptome sequencing, molecular marker development	Ion Torrent	[44]
Blackberry (<i>Rubus</i> sp. Var. Lochness)	Transcriptome sequencing	Illumina Hi-Seq 2000	[6]
Strawberry (<i>Fragaria vesca</i>)	Post-harvest auxin abscisic acid response	Illumina Hi-Seq 2000/2500	[66]
Grapevine (<i>Vitis vinifera</i> X <i>Vitis labrusca</i>)	Transcriptome sequencing	Illumina Hi-Seq 2500	[67]
Grape (<i>Vitis vinifera</i>)	Bud dormancy	Illumina Hi-Seq 2000	[68]

3. Conclusion

Many fruit crops are generated by hybridization and selection. However, crop hybridization breeding has

limitations that are difficult to overcome [3]. This requires enormous amounts of labor and land resources, although fast track breeding techniques and molecular technologies may accelerate breeding and selection processes. Finally advances in biotechnology must be attached to breeding efforts.

Recently different RNA-Seq platforms were released for researchers. It seems that RNA-Seq platforms and bioinformatics for data analysis will continue to have more significance for fruits.

References

- [1] K. Walter, and J. Knops, "The Mystery of Masting in Trees Some trees reproduce synchronously over large areas, with widespread ecological effects, but how and why?," *American Scientist*, vol. 93 (4), pp. 340-347, 2005.
- [2] M. Fernández-Martínez, S. Vicca, I. A. Janssens, J. M. Espelta, and J. Peñuelas, "The role of nutrients, productivity and climate in determining tree fruit production in European forests," *New Phytologist*, vol. 213 (2), pp. 669-679, 2017.
- [3] D. Dönmez, Ö. Şimşek, and Y. A. Kaçar, "Genetic Engineering Techniques in Fruit Science," *International Journal of Environmental & Agriculture Research (IJOEAR)*, vol. 2 (12), pp. 115-128, 2016.
- [4] S. A. Simon, J. Zhai, R. S. Nandety, K. P. McCormick, J. Zeng, D. Mejia, and B. C. Meyers, "Short-read sequencing technologies for transcriptional analyses," *Annu. Rev. Plant Biol*, vol. 60, pp. 305–33, 2009.
- [5] C. Trapnell, B. A. Williams, G. Pertea, A. Mortazavi, G. Kwan, M. J. van Baren, S. L. Salzberg, B. J. Wold, and L. Pachter, "Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts and isoform switching during cell differentiation," *Nature Biotechnol*, vol. 28 (5), pp. 511–515, 2010.
- [6] D. Garcia-Seco, Y. Zhang, F. J. Gutierrez-Mañero, C. Martin, and B. Ramos-Solano, "RNA-Seq analysis and transcriptome assembly for blackberry (*Rubus* sp. Var. Lochness) fruit," *BMC Genomics*, vol. 16 (5), 2015.
- [7] B. T. Wilhelm, and J. R. Landry, "RNA-Seq-quantitative measurement of expression through massively parallel RNA-sequencing," *Methods*, vol. 48, pp. 249–257, 2009.
- [8] S. Zhang, F. Ding, X. He, C. Luo, G. Huang, and Y. Hu, "Characterization of the 'Xiangshui' lemon transcriptome by de novo assembly to discover genes associated with self-incompatibility," *Molecular Genetics and Genomics*, vol. 290 (1), pp. 365-375, 2015.
- [9] A. Brautigam, T. Mullick, S. Schliesky, and A. P. M. Weber, "Critical assessment of assembly strategies for non-model species mRNA-Seq data and application of next-generation sequencing to the comparison of C(3) and C(4) species," *J. Exp. Bot*, vol. 62 (9), pp. 3093-3102, 2011.
- [10] J. Wu, Y. Zhang, H. Zhang, H. Huang, K. Folta, and J. Lu, "Whole genome wide expression profiles of *Vitisamurensis* grape responding to downy mildew by using Solexa sequencing technology," *BMC Plant Biol*, vol. 10 (1), pp. 234, 2010.
- [11] J. H. Malone, and B. Oliver, "Microarrays, deep sequencing and the true measure of the transcriptome," *BMC Biol*, vol. 9 (1), pp. 34, 2011.
- [12] D. Parkhomchuk, T. Borodina, V. Amstislavskiy, M. Banaru, L. Hallen, S. Krobtsch, H. Lehrach, and A. Soldatov, "Transcriptome analysis by strand-specific sequencing of complementary DNA," *Nucleic Acids Res*, vol. 37, pp. e123, 2009.
- [13] S. Marguerat, and J. Bahler, "RNA-Seq: from technology to biology," *Cell Mol. Life Sci*, vol. 67, pp. 569–579, 2010.
- [14] Y. Gilad, J. K. Pritchard, and K. Thornton, "Characterizing natural variation using next-generation sequencing technologies," *Trends Genet*, vol. 25, pp. 463–471, 2009.
- [15] S. B. Montgomery, and E. T. Dermitzakis, "The resolution of the genetics of gene expression," *Hum. Mol. Genet*, vol. 18, pp. R211–R215, 2009.
- [16] T. Werner, "Next generation sequencing in functional genomics," *Briefs Bioinform*, vol. 11, pp. 499–511, 2010.
- [17] P. Martínez-Gómez, C. H. Crisosto, C. Bonghi, "New approaches to Prunus transcriptome analysis," *Genetica*, vol. 139, pp. 755, 2011.
- [18] B. T. Wilhelm, S. Marguerat, S. Watt, F. Schubert, V. Wood, I. Goodhead, C. J. Penkett, J. Rogers, and J. Bähler, "Dynamic repertoire of a eukaryotic transcriptome surveyed at single-nucleotide resolution," *Nature*, vol. 453, pp. 1239–1243, 2008.
- [19] M. Yassour, T. Kaplan, H. B. Fraser, J. Z. Levin, J. Pfiffner, X. Adiconis, G. Schroth, S. Luo, I. Khrebtkova, A. Gnirke, C. Nusbaum, D. A. Thompson, N. Friedman, and A. Regev, "Ab initio construction of a eukaryotic transcriptome by massively parallel mRNA sequencing," *Proc. Natl. Acad. Sci. USA*, vol. 106, pp. 3264–3269, 2009.
- [20] J. C. Marioni, C. E. Mason, S. M. Mane, M. Stephens, and Y. Gilad, "RNA-seq: an assessment of technical reproducibility and comparison with gene expression arrays," *Genome Res*, vol. 18, pp. 1509–1517, 2008.
- [21] A. Mortazavi, B. A. Williams, K. McCue, L. Schaeffer, and B. Wold, "Mapping and quantifying mammalian transcriptomes by RNA-Seq," *Nat. Methods*, vol. 5, pp. 621–628, 2008.
- [22] Q. Pan, O. Shai, L. J. Lee, B. J. Frey, and B. J. Blencowe, "Deep surveying of alternative splicing complexity in the human transcriptome by high-throughput sequencing," *Nat. Genet*, vol. 40, pp. 1413–1415, 2008.
- [23] J. Z. Levin, M. Yassour, X. Adiconis, C. Nusbaum, D. A. Thompson, N. Friedman, A. Gnirke, and A. Regev, "Comprehensive comparative analysis of strand-specific RNRNA sequencing methods," *Nature Methods*, vol. 7 (9), pp. 709-715, 2010.
- [24] M. L. Metzker, "Sequencing technologies — the next generation," *Nature Rev. Genet*, vol. 11, pp. 31–46, 2010.
- [25] F. Ozsolak, and P. M. Milos, "RNA sequencing: advances, challenges and opportunities," *Nature Reviews/Genetics*, vol. 12, pp. 87-98, 2011.
- [26] S. C. Schuster, K. R. Chi, N. Rusk, V. Kiermer, B. Wold, and R. M. Myers, "Method of the year, next-generation DNA sequencing. Functional genomics and medical applications," *Nat. Methods*, vol. 5, pp. 11–21, 2008.
- [27] W. J. Ansorge, "Next-generation DNA sequencing techniques," *New Biotechnology*, vol. 25 (4), 2009.
- [28] A. Brautigam, and U. Gowik, "What can next generation sequencing do for you? Next generation sequencing as a valuable tool in plant research," *Plant Biology*, ISSN 1435-8603, 2010.

- [29] T. D. Harris, P. R. Buzby, H. Babcock, E. Beer, J. Bowers, I. Braslavsky, M. Causey, J. Colonell, J. Dimeo, J. W. Efcavitch, E. Giladi, J. Gill, J. Healy, M. Jarosz, D. Lapen, K. Moulton, S. R. Quake, K. Steinmann, E. Thayer, A. Tyurina, R. Ward, H. Weiss, and Z. Xie, "Single-molecule DNA sequencing of a viral genome," *Science*, vol. 320, pp. 106-109, 2008.
- [30] A. N. Egan, J. Schlueter, and D. M. Spooner, "Applications of next-generation sequencing in plant biology," *American Journal of Botany*, vol. 99 (2), pp. 175-185, 2012.
- [31] J. Shendure, G. J. Porreca, N. B. Reppas, X. X. Lin, J. P. McCutcheon, A. M. Rosenbaum, M. D. Wang, K. Zhang, R. D. Mitra, and G. M. Church, "Accurate multiplex polony sequencing of an evolved bacterial genome," *Science*, vol. 309, pp. 1728-1732, 2005.
- [32] Z. Wang, M. Gerstein, and M. Snyder, "RNA-Seq: a revolutionary tool for transcriptomics," *Nat. Rev. Genet.*, vol. 10, pp. 57-63, 2009.
- [33] L. Flintoft, "Transcriptomics: digging deep with RNA-Seq," *Nat. Rev. Genet.*, vol. 9, pp. 413, 2008.
- [34] A. Cruz-Mendivil, J. A. López-Valenzuela, C. L. Calderón-Vázquez, M. O. Vega-García, C. Reyes-Moreno, and A. Valdez-Ortiz, "Transcriptional changes associated with chilling tolerance and susceptibility in 'Micro-Tom' tomato fruit using RNA-Seq," *Postharvest Biology and Technology*, vol. 99, pp. 141-151, 2015.
- [35] I. Zouari, A. Salvioli, M. Chialva, M. Novero, L. Miozzi, G. C. Tenore, P. Bagnaresi, and P. Bonfante, "From root to fruit: RNA-Seq analysis shows that arbuscular mycorrhizal symbiosis may affect tomato fruit metabolism," *BMC Genomics*, vol. 15, pp. 221, 2014.
- [36] V. Gupta, A. D. Estrada, I. Blakley, R. Reid, K. Patel, M. D. Meyer, S. U. Andersen, A. F. Brown, M. A. Lila, and A. E. Loraine, "RNA-Seq analysis and annotation of a draft blueberry genome assembly identifies candidate genes involved in fruit ripening, biosynthesis of bioactive compounds, and stage-specific alternative splicing," *GigaScience*, vol. 4 (1), pp. 1, 2015.
- [37] H. Y. Kim, P. Saha, M. Faruh, B. Li, A. Sadka, and E. Blumwald, "RNA-Seq Analysis of Spatiotemporal Gene Expression Patterns During Fruit Development Revealed Reference Genes for Transcript Normalization in Plums," *Plant Mol. Biol. Rep.*, vol. 33, pp. 1634-1649, 2015.
- [38] L. Wang, S. Zhao, C. Gu, Y. Zhou, H. Zhou, J. Ma, J. Cheng, and Y. Han, "Deep RNA-Seq uncovers the peach transcriptome landscape," *Plant Mol. Biol.*, vol. 83, pp. 365-377, 2013.
- [39] N. Liu, J. Yang, S. Guo, Y. Xu, and M. Zhang, "Genome-wide identification and comparative analysis of conserved and novel microRNAs in grafted watermelon by high-throughput sequencing," *PloS one*, vol. 8 (2), pp. e57359, 2013.
- [40] C. Feng, M. Chen, C. J. Xu, L. Bai, X. Yin, X. Li, A. C. Allan, I. B. Ferguson, and K. S. Chen, "Transcriptomic analysis of Chinese bayberry (*Myricarubra*) fruit development and ripening using RNA-Seq," *BMC Genomics*, vol. 13, pp. 19, 2012.
- [41] J. Z. Zhang, X. Y. Ai, W. W. Guo, S. A. Peng, X. X. Deng, and C. G. Hu, "Identification of miRNAs and their target genes using deep sequencing and degradome analysis in trifoliate orange [*Poncirus trifoliata* (L.) Raf]," *Molecular biotechnology*, vol. 51 (1), pp. 44-57, 2012.
- [42] R. Ophir, A. Sherman, M. Rubinstein, R. Eshed, M. S. Schwager, R. Harel-Beja, I. B. Ya'akov, and D. Holland, "Singlenucleotide polymorphism markers from de-novo assembly of the pomegranate transcriptome reveal germplasm genetic diversity," *PloS One*, vol. 9 (2), 88998, 2014.
- [43] M. Liang, X. Yang, H. Li, S. Su, H. Yi, L. Chai, and X. Deng, X., "De novo transcriptome assembly of pummelo and molecular marker development," *PloS One*, vol. 10 (3), 0120615, 2015.
- [44] S. Backiyarani, S. Uma, M. S. Saraswathi, A. S. Saravanakumar, and A. Chandrasekar, "Transcriptome analysis of banana (*Musa balbisiana*) based on next-generation sequencing technology," *Turkish Journal of Agriculture and Forestry*, vol. 39 (5), pp. 705-717, 2015.
- [45] K. Yu, Q. Xu, X. Da, F. Guo, Y. Ding, and X. Deng, "Transcriptome changes during fruit development and ripening of sweet orange (*Citrus sinensis*)," *BMC Genomics*, vol. 13 (1), pp. 1, 2012.
- [46] W. B. Jiao, D. Huang, F. Xing, Y. Hu, X. X. Deng, Q. Xu, and L. L. Chen, "Genome-wide characterization and expression analysis of genetic variants in sweet orange," *The Plant Journal*, vol. 75 (6), pp. 954-964, 2013.
- [47] M. K. Azim, I. A. Khan, and Y. Zhang, "Characterization of mango (*Mangifera indica* L.) transcriptome and chloroplast genome," *Plant Molecular Biology*, vol. 85 (1-2), pp. 193-208, 2014.
- [48] H. X. Wu, X. W. Ma, S. B. Wang, Q. S. Yao, W. T. Xu, Y. G. Zhou, Z. S. Gao, and R. L. Zhan, "Transcriptome and proteomic analysis of mango (*Mangifera indica* Linn) fruits," *Journal of Proteomics*, vol. 105, pp. 19-30, 2014.
- [49] Y. Bai, L. Dougherty, and K. Xu, "Towards an improved apple reference transcriptome using RNA-Seq," *Molecular Genetics and Genomics*, vol. 289 (3), pp. 427-438, 2014.
- [50] F. Bai, C. Chen, J. An, S. Xiao, X. Deng, and Z. Pan, "Transcriptome responses to phosphate deficiency in *Poncirus trifoliata* (L.) Raf," *Acta Physiologiae Plantarum*, vol. 36 (12), pp. 3207-3215, 2014.
- [51] S. Wang, B. Lu, T. Wu, X. Zhang, X. Xu, Z. Han, and Y. Wang, "Transcriptomic analysis demonstrates the early responses of local ethylene and redox signaling to low iron stress in *Malus xiaojinensis*," *Tree Genetics & Genomes*, vol. 10 (3), pp. 573-584, 2014.
- [52] M. Alkio, U. Jonas, M. Declercq, S. Van Nocker, and M. Knoche, "Transcriptional dynamics of the developing sweet cherry (*Prunus avium* L.) fruit: sequencing, annotation and expression profiling of exocarp-associated genes," *Horticulture Research*, vol. 11, pp. 1-15, 2014.
- [53] J. Wu, Z. Xu, Y. Zhang, L. Chai, H. Yi, and X. Deng, "An integrative analysis of the transcriptome and proteome of the pulp of a spontaneous late-ripening sweet orange mutant and its wild type improves our understanding of fruit ripening in citrus," *Journal of Experimental Botany*, vol. 65 (6), pp. 1651-1671, 2014.
- [54] Y. J. Zhang, X. J. Wang, J. X. Wu, S. Y. Chen, H. Chen, L. J. Chai, and H. L. Yi, "Comparative Transcriptome Analyses between a Spontaneous Late Ripening Sweet Orange Mutant and Its Wild Type Suggest the Functions of ABA, Sucrose and JA during Citrus Fruit Ripening," *PloS One*, vol. 9 (12), pp. 116056, 2014.

- [55] D. Sanhueza, P. Vizoso, I. Balic, R. Campos-Vargas, and C. Meneses, "Transcriptomic analysis of fruit stored under cold conditions using controlled atmosphere in *Prunus persica* cv. 'Red Pearl', Frontiers in Plant Science, vol. 6, pp. 788, 2015.
- [56] H. Wei, X. Chen, X. Zong, H. Shu, D. Gao, and Q. Liu, "Comparative transcriptome analysis of genes involved in anthocyanin biosynthesis in the red and yellow fruits of sweet cherry (*Prunus avium* L.)," PloS One, vol. 10 (3), pp. 0121164, 2015.
- [57] D. S. O. Coqueiro, A. A. De Souza, M. A. Takita, C. M. Rodrigues, L. T. Kishi, and M. A. Machado, "Transcriptional profile of sweet orange in response to chitosan and salicylic acid," BMC Genomics, vol. 16 (1), pp. 1, 2015.
- [58] N. Wang, Y. Zheng, N. Duan, Z. Zhang, X. Ji, S. Jiang, S. Sun, L. Yang, Y. Bai, Z. Fei, and X. Chen, "Comparative Transcriptomes Analysis of Red-and White-Fleshed Apples in an F1 Population of *Malus sieversii* f. *niedzwetzkyana* Crossed with *M. domestica* 'Fuji', PloS One, vol. 10 (7), pp. e0133468, 2015.
- [59] F. Guo, H. Yu, Q. Xu, and X. Deng, X., "Transcriptomic analysis of differentially expressed genes in an orange-pericarp mutant and wild type in pummelo (*Citrus grandis*)," BMC Plant Biology, 15 (1):1, 2015.
- [60] X. Ji, H. Zhang, R. Wang, N. Yang, L. and X. S. Chen, "Transcriptome profiling reveals auxin suppressed anthocyanin biosynthesis in red-fleshed apple callus (*Malus sieversii* f. *niedzwetzkyana*)," Plant Cell, Tissue and Organ Culture (PCTOC), vol. 123 (2), pp. 389-404, 2015.
- [61] J. Wang, R. Zheng, S. Bai, X. Gao, M. Liu, and W. Yan, "Mongolian almond (*Prunus mongolica* Maxim): the morpho-physiological, biochemical and transcriptomic response to drought stress," PloS One, vol. 10 (4), pp. e0124442, 2015.
- [62] Y. Zhu, Y. Li, D. Xin, W. Chen, X. Shao, Y. Wang, and W. Guo, "RNA-Seq-based transcriptome analysis of dormant flower buds of Chinese cherry (*Prunus pseudocerasus*)," Gene, vol. 25; 555 (2), pp. 362-76, 2015.
- [63] M. Wang, X. Zhang, and J. H. Liu, "Deep sequencing-based characterization of transcriptome of trifoliolate orange (*Poncirus trifoliata* (L.) Raf.) in response to cold stress," BMC Genomics, vol. 16 (1), pp. 1, 2015.
- [64] S. Legay, G. Guerriero, A. Deleruelle, M. Lateur, D. Evers, C. M. André, and J. F. Hausman, "Apple russetting as seen through the RNA-Seq lens: strong alterations in the exocarp cell wall," Plant Molecular Biology, vol. 88 (1-2), pp. 21-40, 2015.
- [65] T. Wu, Y. Wang, Y. Zheng, Z. Fei, A. M. Dandekar, K. Xu, Z. Han, and L. Cheng, "Suppressing Sorbitol Synthesis Substantially Alters the Global Expression Profile of Stress Response Genes in Apple (*Malus domestica*) Leaves," Plant and Cell Physiology, pcv092, 2015.
- [66] J. Chen, L. Mao, W. Lu, T. Ying, and Z. Luo, "Transcriptome profiling of postharvest strawberry fruit in response to exogenous auxin and abscisic acid," Planta, vol. 243 (1), pp. 183-197, 2016.
- [67] L. Shangguan, Q. Mu, X. Fang, K. Zhang, H. Jia, X. Li, Y. Bao, and J. Fang, "RNA- sequencing reveals biological networks during table grapevine ('Fujiminori') fruit development," PloS one, vol. 12 (1), e0170571, 2017.
- [68] M. Khalil-Ur-Rehman, L. Sun, C. X. Li, M. Faheem, W. Wang, and J. M. Tao, "Comparative RNA-seq based transcriptomic analysis of bud dormancy in grape," BMC Plant Biology, vol. 17 (1), pp. 18, 2017.