

REVIEW

Open Access

# Rodent models of neuroinflammation for Alzheimer's disease

Amir Nazem<sup>1\*†</sup>, Roman Sankowski<sup>1†</sup>, Michael Bacher<sup>2</sup> and Yousef Al-Abed<sup>3\*</sup>

## Abstract

Alzheimer's disease remains incurable, and the failures of current disease-modifying strategies for Alzheimer's disease could be attributed to a lack of *in vivo* models that recapitulate the underlying etiology of late-onset Alzheimer's disease. The etiology of late-onset Alzheimer's disease is not based on mutations related to amyloid- $\beta$  (A $\beta$ ) or tau production which are currently the basis of *in vivo* models of Alzheimer's disease. It has recently been suggested that mechanisms like chronic neuroinflammation may occur prior to amyloid- $\beta$  and tau pathologies in late-onset Alzheimer's disease. The aim of this study is to analyze the characteristics of rodent models of neuroinflammation in late-onset Alzheimer's disease. Our search criteria were based on characteristics of an idealistic disease model that should recapitulate causes, symptoms, and lesions in a chronological order similar to the actual disease. Therefore, a model based on the inflammation hypothesis of late-onset Alzheimer's disease should include the following features: (i) primary chronic neuroinflammation, (ii) manifestations of memory and cognitive impairment, and (iii) late development of tau and A $\beta$  pathologies. The following models fit the pre-defined criteria: lipopolysaccharide- and PolyI:C-induced models of immune challenge; streptozotocin-, okadaic acid-, and colchicine neurotoxin-induced neuroinflammation models, as well as interleukin-1 $\beta$ , anti-nerve growth factor and p25 transgenic models. Among these models, streptozotocin, PolyI:C-induced, and p25 neuroinflammation models are compatible with the inflammation hypothesis of Alzheimer's disease.

**Keywords:** Alzheimer's disease, Neuroinflammation, Neurodegeneration, Animal models, Innate immunity, Amyloid- $\beta$ , Tau protein

## Introduction

After decades of research, Alzheimer's disease (AD) remains incurable and thus is considered a major human healthcare challenge [1]. A therapeutic intervention with the potential to cure AD should be a mechanistic disease-modifying agent that can slow or halt the neurodegenerative process; and ideally reverse it towards regeneration [2]. Recently, several potentially disease-modifying agents have been suggested for AD. Many of these suggested therapeutic agents have passed the efficacy testing in animal models. However, all of the ensuing phase three clinical trials have failed [3]. These failures question not only our accurate understanding of the disease [1], based on

which the therapeutic agents have been designed, but also the animal models in which both our understanding of the disease and drug discovery efforts are rooted [4].

Failure in AD drug discovery may in part be attributable to the so-called lesion seduction [1], a simplistic paradigm postulating that AD-related histopathological lesions are a direct reflection of its etiology [1]. Following this paradigm, the most commonly used animal models of AD are designed to recapitulate the lesions of AD [5], namely amyloid- $\beta$  (A $\beta$ ) plaques and neurofibrillary tangles through transgenic induction of mutations related to amyloid and tau production (amyloid precursor protein (APP), presenilin-1 (PS1) and PS2, or tau mutations [6,1,7]). However, since late-onset AD (LOAD) is not caused by such mutations [6], the results from these animal models cannot be reliably extrapolated to the human condition, further widening the gap between human AD pathology and its most commonly used models.

\* Correspondence: anazem@nshs.edu; yalabed@nshs.edu

<sup>†</sup>Equal contributors

<sup>1</sup>Elmezzi Graduate School of Molecular Medicine, The Feinstein Institute for Medical Research, 350 Community drive, Manhasset, NY 11030, USA

<sup>3</sup>Center for Molecular Innovation, The Feinstein Institute for Medical Research, 350 Community drive, Manhasset, NY 11030, USA

Full list of author information is available at the end of the article

In fact, there is growing evidence of very early involvement of other mechanisms which may commence even before the emergence of tau and A $\beta$  pathologies in LOAD pathogenesis [5,8]. These potentially triggering mechanisms include but are not limited to vascular pathology [9], mitochondrial dysfunction [10], oxidative stress [1], hypoxia [11], insulin resistance [12], and chronic neuroinflammation [13].

It has been proposed that a combination of chronic neuroinflammation and (pathological) aging, the so-called 'neuroinflammaging' state [14,15], plays a major role in the mechanism of neurodegenerative disorders, including AD [16,17]. Notably, genetic variants associated with the regulation of innate immunity and phagocytosis (for example, TREM2 [18,19] or CD33 [20]) have been identified as risk factors for LOAD [21,20]. Similarly, neuropathological studies have supported early involvement of neuroinflammation in AD through demonstrating the accumulation of activated microglia and inflammatory mediators in the cerebral neocortex at a low Braak stage for AD pathology [22].

In this study, we focused on characterizing the models suitable for studying the inflammation hypothesis of Alzheimer's disease, based on which neuroinflammation is considered as the driving force of AD pathology and starts early in the course of the disease, prior to tau hyperphosphorylation and amyloid plaque formation [23,13]. A thorough characterization of these models will enable future research to understand the possible transition from 'neuroinflammaging' state to neurodegeneration and also to test the efficacy of potential therapeutic agents to prevent such a transition.

### Neuroinflammation in Alzheimer's disease

Neuroinflammation is known as a key component in the neurodegenerative process of Alzheimer's disease [24]. Characteristics of neuroinflammation, like severity and duration, vary depending on the underlying causes. At one end of the spectrum, there are autoimmune disorders, such as multiple sclerosis, featuring chronic neuroinflammation mainly driven by Th1 cells (reviewed in [25]); At the other end of the spectrum, there is a less fulminant 'smoldering' form of chronic neuroinflammation driven by cells of the innate immunity ([26], reviewed in [27]). The latter is mainly due to age-related impairment of anti-inflammatory mechanisms that leads to the aforementioned 'inflammaging' state [18,20,28] and causes subtle clinical symptoms, as exemplified by neuroinflammation following traumatic brain injury, which may persists for years prior to clinical manifestation as AD [29].

The most recent perspective of the inflammation hypothesis of LOAD, proposed by Krstic et al. [13], has provided a comprehensive sequence of pathological events leading to AD pathology. Based on this hypothesis, the natural neuronal response to inflammatory stress includes

hyperphosphorylation of tau (hp-Tau) and mislocalization of hp-Tau towards the somatodendritic compartment as well as increased APP synthesis [13]. Under physiological conditions, the resulting APP aggregates are cleared by neuroprotective microglia [13]. However, in the setting of pathological aging, for example, midlife overweight and obesity [30], microglia become hyper-reactive with increased release of pro-inflammatory cytokines and dysfunctional phagocytosis [13]. This results in further exposure of neurons to a neurotoxic pro-inflammatory environment without the guard of neuroprotective microglia [13]. The consequent neuronal injury includes breakdown of the axonal cytoskeleton leading to the impairment of axonal transport, formation of axonal swellings of APP aggregates [13], and eventually dystrophic neurites that cannot be removed by hyper-reactive microglia [13]. Secondary to this neuronal degeneration, A $\beta$  plaques are formed from the intracellular APP aggregates [13]. These plaques will trigger further release of pro-inflammatory molecules leading to a vicious circle of neurotoxic pro-inflammatory response [13].

The pathological activation of microglia, which is the center of this proinflammatory response, is characterized by upregulation of MHC antigens and complement receptors [31], as well as release of various pro-inflammatory factors like tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ), interleukin-1 $\beta$  (IL-1 $\beta$ ), IL-6, and reactive nitrogen and oxygen species [31]. These pro-inflammatory factors are neurotoxic especially if accumulated during a chronic neuroinflammatory process [31].

Alternatively, some microglia deteriorate in the process of immune system senescence [32]. Histopathologically, this status manifests with microglial dystrophy, which is distinguished from cytoplasmic hypertrophy as seen in activated microglia. Dystrophic microglia are also associated with neurofibrillary degeneration in AD brain, especially in the temporal lobe [33]. Such observations have led to the hypothesis that senescence of microglia itself might be the initial trigger of Alzheimer's disease neuropathology; in this regard, Alzheimer's disease would be viewed as an immune senescent rather than neuroinflammatory condition [33,32]. However, growing evidence suggests that hyper-reactive microglia is involved in early stages of LOAD [34], but may more rapidly undergo the process of senescence, and thus become non-functional after the initial induction of an aberrant inflammatory response. It is noteworthy that in such dystrophic status, like hyper-reactive mode, microglia remain unable to fulfill their physiological roles of clearing the neurotoxic aggregates [35], like A $\beta$  oligomers, and producing neurotrophic factors, therefore, allowing the process of neurodegeneration to progress. Such paradigm may explain the early-stage responsiveness of the disease to NSAIDs compared to its NSAIDs-induced aggravation at later stages. Restraining

microglial activity in early stage will slow the disease; but in later stages, it will accelerate the disease process probably through restraining the residual neuroprotective and clearance function of dystrophic microglia [36].

#### Rodent models of neuroinflammation

In conventional transgenic animal models of AD, neuroinflammation is mainly known as a secondary response to sustained A $\beta$  overproduction and deposition. It includes microglial activation and variable involvement of the complement system and production of cytokines [13,17,37]. Altogether, in these models, the inflammatory response is incomplete and less severe compared to AD in humans [13]. Janelins and colleagues detected early activation of inflammatory processes in the entorhinal cortex (but not hippocampus) of the triple transgenic model (3xTg) of AD at 3 months of age [38]. Interestingly, the neuroinflammation process was concurrent with the production and accumulation of intracellular A $\beta$  but occurred prior to any significant extracellular A $\beta$  plaque deposition, which manifests at about 12 months of age in the 3xTg mice [38]. Of note, this neuroinflammatory process was characterized by a selective trend of increasing expression of TNF- $\alpha$  and monocyte chemoattractant protein-1 (MCP-1), which was not detected for 21 other cytokines tested [38]. Moreover, a substantial microgliosis was detectable at 6 months of age. Although this study provided valuable evidence for a contributory role of inflammatory factors like TNF- $\alpha$  and MCP-1 in AD pathology, the model system replicates the familial but not sporadic type of AD [38].

An ideal disease model should recapitulate causes, lesions, and symptoms in a chronological order similar to the actual disease [7]. A faithful model to the inflammation hypothesis of AD should be an aged animal that recapitulates early chronic neuroinflammation prior to hyperphosphorylation of tau and A $\beta$  plaque deposition. In rats, a neuroinflammatory process lasting more than 7 days is considered chronic neuroinflammation [39]; and rodents older than 22 months are considered senescent [40].

Here, we reviewed potential rodent models of AD that present early neuroinflammation in the disease process and are not genetically manipulated by mutations related to A $\beta$  or tau production (summarized in Table 1). In this regard, inflammatory responses in amyloid-injected models (reviewed in [41]) are beyond the theme of this article. We categorized models, based on the mechanism of their creation, to immune challenge-based, toxin-induced, and (non-AD) transgenic models. Current knowledge on the chronology of pathological events was analyzed for each model to discuss its potential compatibility with the inflammation hypothesis of AD (see Figure 1 for the compatible models).

#### Immune challenge-based models

##### *Lipopolysaccharide (LPS)-induced neuroinflammation*

A commonly studied model of neuroinflammation is LPS-induced neuroinflammation which represents the current standard paradigm to study neuroinflammation both *in vivo* [42,43] and *in vitro* [26,44]. LPS, also known as endotoxin, is a component of the outer membrane of gram-negative bacteria. LPS binds CD14 on microglia membranes. The LPS-CD14 complex then interacts with the toll-like receptor-4 (TLR-4) [26,45], which, in turn, activates microglia by initiating signal transduction cascades leading to rapid transcription and release of pro-inflammatory cytokines [46] (including IL-1, IL-6, IL-12, p40, and TNF- $\alpha$ ), chemokines (for example, CCL2, CCL5, and CXCL8), the complement system proteins (for example, C3, C3a, and C5a receptors) [46], as well as anti-inflammatory cytokines like IL-10 [47] and transforming growth factor- $\beta$  (TGF- $\beta$ ) [48].


Different paradigms of LPS-induced neuroinflammation exist with respect to the route of administration, duration of exposure and age of the animals [49]. While chronic central administration of LPS can induce memory and learning deficits analogous to AD cognitive decline [50], systemic LPS administration led to selective hippocampal impairment in context-object discrimination but not spatial memory [51]. Moreover, Bordou and colleagues recently investigated the role of duration of exposure to LPS as well as the age of exposed rats on the neuroinflammatory response to LPS. Male rats at three age groups of young (3 months), middle-aged (9 months), and aged (23 months) received continuous infusion of picomolar levels of LPS (or artificial CSF as control) into their fourth ventricle [49]. The duration of exposure was either 3 or 8 weeks. Among all cytokines, TNF- $\alpha$  increase in response to LPS infusion was similar in different age groups [49]. However, in contrast to young rats, IL-1 $\beta$  did not significantly increase after 3 weeks of infusion in middle-aged and aged rats. Instead, aged rats had significantly increased IFN- $\gamma$  compared with younger rats [49]. Among rats of the same age group, longer duration of exposure to LPS infusion significantly increased the elevations of IL-1- $\alpha$ , IL-2, IL-4, IL-5, IL-6, IL-12, IL-13, and GM-CSF levels [49]. This study provides an evidence of the influence of age and chronicity of infection on neuroinflammatory responses in certain regions of the brain, like locus coeruleus, which undergo significant cell loss in early stages of AD [52].

In similar studies performed by Wenk and coworkers [43,53-55], chronic neuroinflammation was modeled through continuous infusion of picomolar concentrations of LPS into the fourth ventricle of adult rats. A widespread activation of microglia was detected 2 days after the initiation of LPS infusion [55]. Within 2 weeks after the cessation of LPS infusion, microglial activation

**Table 1 Rodent models of neuroinflammation**

Models	Predisposing factors/causes	Time of appearance of lesions		Signs (time detectable)	SLC reading key	Reference
		hp-Tau	A $\beta$ depositions			
LPS	Peripheral immune challenge, chronic neuroinflammation	?	?	Fear memory (?) Spatial memory (?)	S1LOC1	[165]
PolyI:C	Peripheral immune challenge, chronic neuroinflammation	3m (PHF, but not NFTs)	12m (APP depositions)	Spatial memory (20 m)	S1L1C1	[5]
ICV-STZ	Disrupted insulin signaling, chronic neuroinflammation	6-7w	12w	Spatial memory Visual recognition memory (3w)	S1L1C1	[85]
ICV-OKA	Inhibition of serine/threonine phosphatases 1 and 2A	2w (PHF, but not NFTs)	6w (Non-fibrillar A $\beta$ deposits)	Spatial memory (?)	S1L1C0	[102] [104]
ICV-colchicine	Inhibition of tubulin formation/ microtubule breakdown	? (Tau dephosphorylation)	? (Amyloid plaque)	Spatial memory (14d to 21d)	S1LOC1	[113] [117]
p25 Tg	Upregulation of cPLA2, neuroinflammation	4w	8w	Contextual fear memory (6w)	S1L1C1	[145]
IL-1 $\beta$ Tg	Chronic neuroinflammation	?	? (Increased clearance of amyloid plaques)	Contextual fear memory (12w)	S1LOC0	[39]
Anti-NGF antibody Tg	Blockade of NGF signaling pathway	? (Neurofibrillary pathology)	? (Amyloid plaques)	Visual recognition memory (4 m); Spatial memory (9 m)	S1L1C0	[148] [149]

This table summarizes the suggested models of late-onset AD (LOAD) displaying neuroinflammation as one of the prominent pathological events. The SLC reading key is a scoring system that represents the compatibility of an animal model with the disease in humans with respect to signs (S), lesions (L), and causes (C) [7]. Compatibility is indicated by 1 and incompatibility by 0. Based on SLC reading key, p25 tg, PolyI:C-, and STZ-induced neuroinflammation models are compatible with the inflammation hypothesis of LOAD [13]. (Abbreviations: ? unavailable data; LPS lipopolysaccharide; PolyI:C polyriboinosinic-polyribocytidilic acid; p25 Tg p25 transgenic model; NGF nerve growth factor; IL-1 $\beta$  Tg interleukin-1 $\beta$  transgenic model; ICV intracerebroventricular; STZ streptozotocin; OKA okadaic acid; hp-Tau hyperphosphorylated tau; A $\beta$  amyloid  $\beta$ ; PHF paired helical filaments; NFT neurofibrillary tangles; cPLA2 cytosolic phospholipase 2; SLC Signs, Lesions, Causes; w week; m month).



decreased in most brain areas barring the hippocampus, and after the following 2 weeks, inflammation was mainly localized in the hippocampus [55]. Furthermore, MRI scans showed shrinkage of the temporal lobe and enlargement of the lateral ventricles. Of note, electron microscopic studies showed impaired protein synthesis in hippocampal neurons of LPS-injected animals [54]. Moreover, neuronal loss and impairment of long-term potentiation were reported in the entorhinal cortex [56] and the dentate gyrus of the hippocampus respectively [57], altogether explaining the decline in spatial memory [56]. In this model, LPS-induced neuroinflammation was time dependent (maximal within 4 weeks of infusion) as well as cell and region specific (microglia in hippocampus) [55].

Other groups have provided evidence of exacerbated AD-related protein pathology such as increased Aβ production

through enhanced β-secretase activity in APP sweTg [44] and tau hyperphosphorylation in 3xTg-AD mice following LPS injection [42]. However, wild-type animals injected with LPS showed no increased Aβ deposition in the time course of 3 months. The authors argue that this process might occur at a later time point due to severe neuronal dysfunction and neurodegeneration [44]. Another explanation for the lack of Aβ deposition in LPS-injected animals was proposed by DiCarlo et al. by showing a reduction of established Aβ plaques after intrahippocampal LPS injection through stimulation of Aβ clearance [58].

In line with neurodegenerative structural changes, LPS-induced neuroinflammation caused cognitive deficits leading to impaired performance in associative and spatial learning tasks [59,60]. Finally, a long-term characterization of LPS-induced changes with regard to the chronology of

histological and ultrastructural changes, as well as cognitive deficits, is required to assess the compatibility of LPS neuroinflammation model with the inflammation hypothesis of AD [13].

#### **PolyI:C-induced neuroinflammation**

It is well understood that peripheral infections influence the inflammatory state of the central nervous system [61,62]. The brain innate immune system reacts to systemic inflammation through activation of microglia (reviewed in [13,31]), which may become a potential trigger for neurodegenerative processes [31,61], especially when neuroinflammation becomes chronic in an aging brain [31]. Based on this evidence, a recent study proposed a non-transgenic mouse model of LOAD featuring chronic neuroinflammation after *in utero* systemic immune challenge [5]. Polyriboinosinic-polyribocytidilic acid (PolyI:C) is a synthetic double-stranded RNA that induces an innate immune response analogous to acute viral infections in treated animals. In the CNS, this immune response is mainly mediated by TLR-3-induced activation of microglia [63], followed by an NF- $\kappa$ B-dependent induction of proinflammatory cytokines including IL-1 $\beta$ , IL-6, IL-8, TNF- $\alpha$ , and type I and II interferons [64].

Exposure to PolyI:C leads to inflammation in the injected animal, as well as a chronic proinflammatory state in the fetus of pregnant animals [5,65,66]. Systemic injection of PolyI:C on gestational day 17 led to chronic neuroinflammation, as observed by significantly higher levels of IL-1 and IL-6 compared to the control group [5]. Increased brain cytokine levels were detectable as early as the age of 3 weeks and sustained throughout aging [5]. However, tau hyperphosphorylation started only after 3 months of age and, in spite of some fluctuation, was significantly higher than controls at 6 and 15 months of age [5]. Amyloidogenesis started as late as 12 months of age [5]. Later, at 20 months, animals in the immune-challenged group presented with a significant impairment of spatial recognition memory compared to the control group.

There is growing evidence that early-life infection may lead to abnormalities in cognitive aging [67], probably due to glial priming effect, leading to significantly enhanced glial reactivity to a subsequent immune challenge later in life [67]. The late-gestational PolyI:C induced neuroinflammation model is an informative approach to study the effects of life-long neuroinflammation on cognitive function. Notably, a single intraperitoneal injection of the pregnant mouse was sufficient to change the inflammatory state of the progeny throughout the entire life span, with two additional PolyI:C injections in adulthood exacerbating the pathology. Further work is required to more thoroughly assess the cognitive phenotype of the aged animals, especially since Bitanirwe et al. described neuropsychiatric changes in prenatally stimulated adults [68].

It is noteworthy that the APP deposition and paired helical filaments (PHF) in this model do not completely replicate the lesion in advanced human AD pathology, that is, A $\beta$  plaques and neurofibrillary tangles (NFT). However, the protein pathology is entirely endogenous and not relying on overexpression of human proteins in murine cells. Thus, the differences between PolyI:C-challenged mice and human AD patients might reflect characteristics of the respective species. Finally, the chronological order of pathological events and cognitive deficits in the PolyI:C model is compatible with the inflammation hypothesis of AD [13]; and therefore, it may be a suitable model for studying early stages of LOAD pathology from this perspective.

#### **Neurotoxin-induced models**

##### ***Streptozotocin-induced neuroinflammation***

Since Siegfried Hoyer [69,70] and Suzanne Craft [71] first described the causality between impaired insulin signaling and cognitive dysfunction, a growing body of evidence has emerged on impaired brain insulin signaling and glucose metabolism in LOAD (reviewed in [72]). Peripheral injection of the glucosamine-nitrosurea compound streptozotocin (STZ) selectively damages pancreatic  $\beta$ -cells after being taken up via the glucose transporter SLC2A2 [73,74]. Thus, repetitive intraperitoneal injection of STZ is an established animal model of diabetes mellitus [75]. Interestingly, after STZ-mediated induction of diabetes, rodents display impaired neuronal plasticity and learning deficits [76]. In a recent study, STZ-induced diabetic rats presented frontal lobe neurodegeneration (as evidenced by FJC staining) and hippocampal atrophy, accompanied by A $\beta$  aggregation, synapse loss, and the consequent cognitive decline 4 months after receiving IV STZ [77]. These deficits are mainly induced by peripheral effects of STZ as the molecule does not cross the blood-brain barrier [78,79].

Acutely, STZ causes oxidative stress through the generation of hydrogen peroxide [80] and NO [81]. Furthermore, it leads to DNA damage by alkylation and methylation leading to apoptosis [82]. Notably, since PARP knockout animals are resistant to STZ-induced diabetes, cell death is likely caused by depletion of reduction equivalents followed by ATP depletion [83]. In the long term, STZ induces a state of metabolic imbalance characterized by impaired insulin secretion (reviewed in [78]) and inflammation [84]. Intracerebroventricular (ICV) [85-87] or intracerebral [88] STZ injection induces impaired brain insulin signaling in rodents. Regardless of the route of administration, the animals develop neuroinflammation and cognitive deficits. The following discussion focuses on the ICV injection of STZ.

A single ICV injection of 1 or 3 mg/ml STZ in rats has been shown to cause chronic neuroinflammation,

dilation of the ventricles, and atrophy of the septum with reduction of neuronal cell counts [89]. Both STZ concentrations cause these effects; however, they are more pronounced at 3 mg/ml [89].

When administered to transgenic models of AD, ICV-STZ was shown to exacerbate neuroinflammation, cognitive deficits, plaque pathology, and tau hyperphosphorylation [90,91], indicating that STZ renders the brain more susceptible to the pathological hallmarks of AD. Similar exacerbating effects were observed when STZ was administered intraperitoneally to AD transgenic mice [92-94].

Chen and coworkers have recently compared ICV-STZ wild-type and the widely used 3xTg AD mouse models [85]. Immunohistochemical studies showed early and extensive neuroinflammation in the ICV-STZ mice, characterized by increased astroglial and microglial activation especially in hippocampal CA1, CA3, and dentate gyrus [85]. STZ-induced neuroinflammation was more pronounced as compared to neuroinflammation in 3xTg mice in the same regions [85].

In the ICV-STZ wild-type model, neuroinflammation was detectable 1 week after low-dose ICV-STZ injection while paraventricular A $\beta$  depositions and hippocampal hyperphosphorylated tau appeared within 3 months of ICV-STZ injection [89,85]. With high-dose ICV-STZ injection, however, tau hyperphosphorylation was detectable as early as 4 weeks in rats [95] and 6 weeks in mice [85]. Within 3 weeks of STZ injection in wild-type mice, spatial and short-term memory deficits developed, indistinguishable from cognitive deficits in 3xTg mice.

All in all, the ICV-STZ model not only displays neuroinflammation but also reproduces tau [85] and amyloid [89] pathologies as well as AD-like cognitive deficits [96,97] with a chronology compatible with the inflammation hypothesis of AD [13].

#### ***Okadaic acid-induced neuroinflammation***

A comparable model to STZ-induced neuroinflammation is okadaic acid (OKA)-induced neuroinflammation. OKA is a major polyether toxin that selectively inhibits serine/threonine phosphatases 1 and 2A [98]. The decreased activity of protein phosphatase 2A (PP2A) has been observed in the pathology of AD [99] and was proposed to be involved in hyperphosphorylation of tau [100].

In line with the abovementioned molecular link to AD pathology [101], (ICV) OKA injection develops memory impairment in rats [102,103], making it suitable for further characterization as a potential AD model [104]. In studies performed by Arendt and colleagues [105], ICV infusion of OKA (70 ng/day; for up to 4 months), could replicate some AD-associated pathologies including hyperphosphorylation of tau (at Ser-202/Thr-205) and apoptotic cell death within 2 weeks, as well as cortical deposition of non-fibrillar A $\beta$  within 6 weeks of infusion.

Interestingly, Lee and colleagues later confirmed the formation of paired helical filaments of tau following intra-hippocampal injection of OKA (1 mM, 0.5 ml) [106]. It is noteworthy, however, that in this model, hyperphosphorylated tau aggregates do not develop into NFTs [105].

In addition to AD-like histopathological changes, in a recent study, memory impairment was reported in the Morris water maze test 15 days after ICV-OKA (200 ng) injection [107]. In contrast to control and vehicle groups, OKA 200 ng treated rats did not present significant decrease in latency time to reach the platform in the second and third sessions as compared to the first session [107].

Interestingly, OKA-induced memory impairment is found to be associated with neuroinflammation [108]. In OKA-injected rats, neuroinflammation was characterized by increased expression of proinflammatory cytokine TNF- $\alpha$  and IL-1 $\beta$  as well as total nitrite in both hippocampus and cortex [107].

However, the effect of antidementic (non-antiinflammatory) drugs on subsiding the neuroinflammation supports the reactive rather than the causative role of neuroinflammation (with regard to neurodegeneration) in this model [109,103]. Thirteen days of pretreatment of OKA-injected rat with anti-dementic drugs memantine (10 mg/kg) and donepezil (5.0 mg/kg) could protect not only the ICV-OKA-induced memory impairment but also the associated changes in TNF- $\alpha$ , IL- $\beta$ , and total nitrite levels as well as expressions of iNOS and nNOS [107,110]. In addition, recent studies have shown that OKA-induced oxidative stress [111,112] is linked to dysfunction of astrocytic neuroprotection [102]. Twelve days after intra-hippocampal injection of OKA (100 ng), rats developed spatial cognitive impairment, accompanied by hippocampal astrogliosis (as evident by, increased GFAP), and oxidative stress (for example, decreased glutamine synthetase and decrease in reduced glutathione content) [102]. Thus, the application of this model as an etiology-based model of neuroinflammation in LOAD requires further characterization of the sequence of different pathological events, including the possible precedence of oxidative stress.

#### ***Colchicine-induced neuroinflammation***

Similarly, ICV injection of colchicine in rats could induce AD-like pathology with consequential cognitive and behavioral alterations similar to AD [113]. Colchicine is a cytotoxic agent that irreversibly binds to tubulin molecules and thus halts the aggregation of tubulin dimers to the fast growing end, causing disruption of microtubule polymerization [114]. Blocking axoplasmic transport, colchicine severely damages hippocampal granule cells and mossy fibers, eventually leading to neuronal loss, which manifests with cognitive impairment and spontaneous motor activity [115]. Systemic and neurologic symptomatology

of rats in response to the central administration of high-dose colchicine is further detailed in [113].

Neuroinflammation plays a key role in the development of AD-like neurodegeneration in this model [113,116,117]. In a recent study, Ho and colleagues found both *in vitro* and *in vivo* evidence for the involvement of COX-2-mediated apoptotic mechanisms in colchicine-induced neurotoxicity [116]. After intra-hippocampal injection of colchicine in rats, a significant increase in COX-2 mRNA levels was found in dentate gyrus granule cells, followed by apoptotic morphological changes [116]. Similarly, Sil et al. demonstrated that colchicine-induced neurodegeneration is mediated by COX-induced neuroinflammation [117]. In this study, the TNF- $\alpha$  level of the hippocampus was significantly higher in the colchicine-injected (15  $\mu$ g, ICV) rats compared to control and sham-operated rats. Likewise, the nitrite level and the ROS level of hippocampus were also significantly higher in colchicine-induced neurodegeneration compared to controls and sham-operated rats [117]. This study also reported significant increase in amyloid plaques found in the hippocampus. It is noteworthy that naproxen administration (doses 5, 10, or 20 mg/kg) could prevent from TNF- $\alpha$  increase and reduce the amyloid plaque formation [117].

Of note, AD-like tau pathology, however, does not occur in colchicine model [118]. In fact, although colchicine leads to microtubule breakdown [118], as seen in Alzheimer's disease, its mechanism is based on tau dephosphorylation rather than hyperphosphorylation [119,120]. The chronological association of the pathological events with neuroinflammation is yet to be investigated in this model. Without chronological characterization, it is unclear if this model is compatible with the inflammation hypothesis of AD.

#### Genetically manipulated models (unrelated to mutations in familial AD)

##### *IL-1 $\beta$ overexpression model*

Interleukin-1 $\beta$  (IL-1 $\beta$ ), regulating acute and chronic neuroinflammatory responses (reviewed in [121,122]), is found elevated in AD patients [123,121,124,125]. Based on this notion, O'Banion and coworkers have developed an inducible IL-1 $\beta$  overexpression model of chronic neuroinflammation (IL-1 $\beta$  excisional activation transgenic (XAT) mouse model) [126]. Prolonged IL-1 $\beta$  elevation induces microgliosis and astrogliosis alongside with chronic elevation of the proinflammatory cytokines, IL-6 and TNF- $\alpha$  [126]. After activation of the inducible transgene in this model, neuroinflammation may last as long as 10 months [126].

IL-1 $\beta$  transgenic mice displayed a dualistic histopathological presentation with respect to the hallmarks of AD. On the one hand, APP production and processing are unaltered despite prolonged overexpression of the transgene [126]; and notably, the amyloid plaque burden was

even reduced in crossed IL-1 $\beta$  XAT and APP/PS1 Tg mice [126,127]. On the other hand, crossing IL-1 $\beta$  XAT and 3xTg AD mice led to significant exacerbation of tau hyperphosphorylation within 1 month of IL-1 $\beta$  overexpression [128]. It is conceivable that in this model, IL-1 $\beta$  per se induces activation of microglia without shifting it to a dysfunctional hyper-reactive state, therefore sparing its ability to readopt the phagocytic mode in the aid of clearance of A $\beta$  plaques [129].

It is likewise noteworthy that the IL-1 $\beta$  XAT mouse model lacks overt neuronal loss or apoptosis within 2 and 5 months of IL-1 $\beta$  overexpression, respectively [127,126]. However, despite the absence of neuronal loss, the animals manifested significant cognitive deficits, including contextual fear memory and spatial memory defects within 3 months of transgene induction [39].

These divergent effects of IL-1 $\beta$  overexpression on A $\beta$  and tau pathology can be explained in the context of inflammation hypothesis of AD proposed by Krstic and Knuesel [13]. Based on this hypothesis, tau hyperphosphorylation is an early neuronal response to neuroinflammatory stress, while A $\beta$  pathology emerges after microglial shift to the pro-inflammatory M1 phenotype as opposed to the phagocytic M2 phenotype [13]. This implies that in the IL-1 $\beta$  XAT mouse model, microglia maintain their physiological function [130] in contrast with immune challenge-based animal models (for example, LPS and PolyI:C injection) or neurotoxin (that is, STZ) models.

Altogether, in spite of corroborating the association between chronic neuroinflammation and cognitive deficits, the IL-1 $\beta$  model could not reproduce the main lesions of AD pathology; therefore it may not be a suitable model for sporadic AD.

##### *p25 transgenic model*

Neurons are considered as terminally differentiated non-dividing cells. However, the evidence of expression of cell cycle-specific proteins [131] and DNA replication in neurons prior to neurodegeneration in AD-prone brain regions [132-134] supports the association of AD neurodegeneration with cell cycle dysregulations. Regulation of the cell cycle is performed by cyclins and cyclin-dependent kinases (CDKs) (reviewed in [135]). CDK5, in particular, also plays an important role in the brain development by promoting neurite outgrowth in post-mitotic neurons [136,137]. p35 is a regulatory subunit of CDK5 [138], and cleavage of p25 by the calcium-dependent kinase calpain leads to neurotoxicity through aberrant CDK5 activation [139]. Interestingly, neurotoxicity itself drives p35 cleavage possibly creating a vicious cycle [140].

CDK5 activation and aberrant p25 accumulation was shown in AD patients [139]. Notably, mice overexpressing human p25 displayed AD-like pathology [141], suggesting



p25 overexpression as a potential mechanism to model AD [142]. Unlike its precursor p35, the p25-CDK5 complex was shown to induce tau hyperphosphorylation possibly explaining the corollary AD-like pathology [141]. In p25 Tg mice, neuroinflammation is detectable before other AD pathology hallmarks, including tau hyperphosphorylation [143]. Said neuroinflammation is characterized by astrogliosis and increased levels of pro-inflammatory cytokines, like TNF- $\alpha$ , IL-1 $\beta$ , and MIP-1 $\alpha$  [143]. Activation of microglia was also detectable 4 weeks after induction of p25 overexpression [143].

Interestingly, Sundaram and colleagues reported a relatively clear temporal sequence between neuroinflammation and secondary pathological hallmarks associated with AD [143]. While neuroinflammation occurred as early as first week, tau hyperphosphorylation and amyloidogenesis were detectable at 4 and 8 weeks after p25 induction, respectively [143]. Finally, cognitive deficit (contextual fear memory) was detectable within 6 weeks of p25 induction [144].

In summary, the p25 Tg mice resemble the histopathological hallmarks of AD with amyloid depositions [143], tau hyperphosphorylation [143], and neurodegeneration [145]. Additionally, these mice display cognitive deficits [144]. These pathological events occur in a chronological order compatible with the inflammation hypothesis of AD [13].

#### ***Anti-nerve growth factor (NGF) transgenic models***

The transgenic expression of anti-nerve growth factor (NGF) antibodies led to an overt neurodegenerative phenotype in aged mice [146,147]. This model, known as AD11 model, resembles the insidious cognitive decline of LOAD by manifesting with a significantly progressive deficit in visual recognition memory (as evaluated through an object recognition test) and spatial memory (as evaluated through an eight-arm radial maze), starting at 4 months and 9 months of age, respectively [148]. Of note, in this model, neurodegeneration was characterized by neuronal loss, cholinergic deficit, tau hyperphosphorylation (associated with neurofibrillary pathology) and A $\beta$  plaques [149].

The primary assumptions alluded to a link between chronic deprivation of NGF and abnormal processing of amyloid precursor protein, leading to A $\beta$  excessive formation and deposition [149,150]. However, growing evidence, including gene expression profiles [151], is showing early involvement of neuroinflammatory elements in AD11 model [152]. Interestingly, changes in the expression of inflammatory and immune response genes were the earliest and most significant [151]. Specifically, significant changes were found in the expression of genes encoding for proteins of the complement cascade and the major histocompatibility complex (MHC). D'Onofrio and colleagues demonstrated that in AD11 model, overexpression

of C3 mRNA was significantly high as early as P30 in the cortex and hippocampus and at P90 in all brain areas [151]. Also, C1qb was significantly upregulated in the cortex and hippocampus at P90. Similarly, a significant dysregulation of MHC class I gene expression was observed in AD11, as some mRNAs of Class I MHC (in particular H2-Q1) were severely reduced in some areas at P30 but significantly increased at P90 [151]. This dysregulated MHC class I expression may explain the disruption in dynamic synaptic strength and connectivity, processes involved in memory formation.

Beside the above-explained 'non-immune' involvement of inflammatory factors in AD11 models, there is growing evidence on their role in a concurrent neuroinflammation process, further aggravating neurodegeneration. For example, the gene expression profiles showed significant changes in the expression levels of other inflammatory-related mRNAs, such as Ccl5 (chemokine ligand 5)/RANTES, IL-1 $\beta$ , TNF- $\alpha$ , IFN- $\gamma$ -induced ATPase, Cd47, Ccl17, Cd300lf, Cd72, and Cox-2 [151]. Altogether, AD11 mice demonstrate a complex neurotrophic and inflammatory dysregulation in key brain regions with potential role in the outset of the neurodegeneration process described in this model [146]. However, in order to avoid probable autoimmune reactions and achieve a more brain-specific phenotype, an inducible NGF knockout might be a more relevant approach.

Similarly, inflammatory response gene expression was significantly activated in AD10 mice [153], a variant of AD11 model without the antibody heavy chain, leading to a similar neurodegenerative picture [154]. Interestingly, however, when AD10 mice were housed in a murine pathogen-free environment, the inflammatory gene response significantly subsided, and neither an overt neurodegeneration nor behavioral symptoms occurred [154].

Histopathological studies that characterize the microglial activation in this model are lacking. Of special interest would be the temporal relationship of neuroinflammation and other AD neurodegeneration processes, including tau pathology and amyloid deposition. Thus, the compatibility of this model with the neuroinflammation hypothesis of AD is yet to be investigated.

#### ***TGF- $\beta$ -deficient models***

TGF- $\beta$  is a cytokine involved in several opposing physiological functions in inflammation pathways and cell growth, depending on the target cell type, cell environment, as well as amount and duration of exposure to TGF- $\beta$  [155,156]. In the CNS, TGF- $\beta$  is produced by both neurons and glial cells [157]. Recent evidence from autopsied samples of AD patients showed elevated TGF- $\beta$  in brain microvessels, leading to the release of pro-inflammatory cytokines like TNF- $\beta$  and IL-1 $\beta$  from endothelial cells in the brain [158]. This is in line with

experimental findings on TGF- $\beta$  transgenic mice, showing that long-term increase in expression of TGF- $\beta$  is associated with increased perivascular amyloidogenesis [156]. On the other hand, short-term increase in TGF- $\beta$  expression was neuroprotective [156]. Notably, a modest increase of astroglial TGF- $\beta$ 1 expression enhanced A $\beta$  clearance in aged human APP transgenic mice [159], confirming the neuroprotective role of TGF- $\beta$ 1. Thus TGF- $\beta$  knockout models, completely lacking the mentioned neuroprotective effects, have demonstrated overt neurodegeneration (reviewed in [24]). TGF- $\beta$ -/- mice displayed neurodegeneration and neuroinflammation at P1 and P21 without a clear temporal sequence of the two events [160]. However, in order to create a model based on TGF- $\beta$ 1 deficiency to study the inflammation hypothesis of AD, the effects of TGF- $\beta$ 1 deficiency (and thus its neuroprotective function) on neuroinflammatory dysregulations and its potential effect on triggering AD-like neurodegeneration in aged rodents should be investigated.

## Discussion

The focus of this study was to recognize rodent models of LOAD that present a process of chronic neuroinflammation prior to tau and amyloid pathology, as described by the inflammation hypothesis of AD [13]. Among models of chronic neuroinflammation, here we found ICV-STZ and PolyI:C-induced neuroinflammation and p25 transgenic (p25 tg) models most compatible with the inflammation hypothesis of AD in terms of temporal ordering of pathological events. The evidence on temporal order of pathological events is more controversial in LPS- and OKA-induced neuroinflammation. The timeline of AD-related pathological events in colchicine-induced neuroinflammation as well as anti-NGF overexpression and TGF- $\beta$ 1 knockout models is yet to be investigated. Owing to the clear temporal order of pathological events, ICV-STZ, PolyI:C, and p25 tg models are more suited to study the effect of anti-inflammatory agents on different stages of LOAD; the results of such studies may improve the design of clinical trials for potential preventive or therapeutic agents.

Since most cells (for example, microglia) and molecules (for example, TNF- $\alpha$ , TGF- $\beta$ ) of the immune system are multifunctional [161] and sometimes even demonstrate completely opposite functions, depending on the context [161], neuropathological characterization of neuroinflammation provides little information on its actual role in AD pathogenesis [24]. In addition, while temporal precedence of neuroinflammation supports its causative role in neurodegeneration [17], it does not rule out a physiological neuroprotective role of an early neuroinflammatory process against an underlying pathological process [161]. Of course, a physiological role is much less probable in the case of chronic neuroinflammation which by definition implies the

inability of the immune system to completely remove or deactivate the injurious agent and then resolve [17].

The detrimental role of chronic neuroinflammation is more clear in those models based on direct stimulation of the immune system (that are, LPS, PolyI:C, and Il-1beta overexpression model) [161]. For models not based on direct stimulation of immune system, further evidence is required to analyze the actual role of neuroinflammation in respect to each stage of AD pathology. For instance, studies that demonstrated reduced AD-related symptoms and pathology in animals pretreated with anti-inflammatory drugs support the pathological role of neuroinflammation in those models. For example, Dhull and colleagues reported significant increase in survival of hippocampal neurons and improvement in memory performance (as tested in Morris water maze) in ICV STZ rats who received COX-1 and Cox-2 inhibitors [162]. Similarly, administration of naproxen (a nonspecific COX inhibitor) reduced the amyloid plaque formation in the ICV colchicine-induced neuroinflammation model in a dose-dependent manner [117].

Neuropathologically, AD is characterized by extracellular amyloid plaques and intracellular neurofibrillary tangles [17,163]. Although amyloid or tau pathology followed chronic neuroinflammation in the above-discussed models, formation of typical senile plaques and neurofibrillary tangles, as seen in human AD, was not observed in these studies. Instead, intermediate pathological species, like A $\beta$  aggregates [143] and hyper-phosphorylated tau [97,141], were considered as signs of amyloid and tau pathologies, respectively. Of note, the recapitulation of these pathological hallmarks is a great challenge in every rodent model of AD [164].

Moreover, among the discussed studies, different types of molecular evidence were considered as the sign of AD-like amyloid pathology, including A $\beta$  plaque-like structures in STZ-induced model [96], APP containing plaques in PolyI:C-induced model [5], increased levels of alpha-, beta-, and gamma-secretase activity in brain lysate, and intracellular A $\beta$  aggregates in LPS model [50]. Similarly, the evidence of neurodegeneration was also different among these models; for instance, LPS-induced model led to apoptotic neuronal loss [165], while P25 transgenic model underwent extensive non-apoptotic neuronal death [145]. Thus, from these perspectives, a direct comparison of these animal models would be challenging. Future experiments aimed to compare the efficacy of these models in recapitulating AD amyloid and tau pathologies would be illuminating.

Finally, models reviewed in this article were those without underlying genetic manipulations related to familial AD. There are potential models that their effects on neuroinflammation have not yet been evaluated on wild-type animals. For example, degeneration of the locus ceruleus

through alkylating agent N-(2-chloroethyl)-N-ethyl-bromobenzylamine (DSP4) in APP transgenic animals displayed increased neuroinflammation and exacerbation of plaque pathology and behavioral deficits [166]. Future studies on aged wild-type animals can provide evidence for application of this model in sporadic AD.

## Conclusion

Disease-specific animal models are indispensable for the understanding of possible disease mechanisms as well as for preclinical drug development. Undoubtedly, conventional transgenic models of AD are the basis of our today's in-depth understanding of several mechanisms that are probably involved in AD. However, since all potential Alzheimer's disease-modifying agents tested in these models have failed in phase-3 clinical trials, their application in drug discovery is under question.

Different strategies can be considered to bridge the gap between human AD pathology and rodent AD models. On one hand, major efforts should be undertaken to thoroughly characterize conventional animal models with newly available methods [167], to allow for more realistic translation of the results from animal models to human LOAD. On the other hand, etiology-based models should be established for LOAD [1]. Thus far, several hypotheses regarding the probable etiology of AD have been suggested, however, appropriate *in vivo* models to test these hypotheses are still lacking. In this review, we synthesized the current information about rodent models potentially compatible with the inflammation hypothesis of AD [13]. All in all, the choice of an animal model should be an informed decision on behalf of the investigator. Nevertheless, using etiology-based models of LOAD may create a breakthrough in understanding of the disease pathology, designing precise diagnostic modalities and discovery of effective therapeutic agents.

## Abbreviations

APP: amyloid precursor protein; A $\beta$ : amyloid- $\beta$ ; CDK: cyclin-dependent kinases; IFN: interferon; hpTau: hyperphosphorylation of tau; IL: interleukin; ICV: intracerebroventricular; LOAD: late-onset Alzheimer's disease; LPS: lipopolysaccharide; MHC: major histocompatibility complex; NGF: nerve growth factor; NFT: neurofibrillary tangles; OKA: okadaic acid; PHF: paired helical filaments; PolyI:C: polyriboinosinic-polyribocytidilic acid; PS1: presenilin-1; PP2A: protein phosphatase 2A; STZ: streptozotocin; TLR-4: toll-like receptor-4; TGF- $\beta$ : transforming growth factor- $\beta$ ; TNF- $\alpha$ : tumor necrosis factor- $\alpha$ .

## Competing interests

The authors declare that they have no competing interests.

## Authors' contributions

AN designed the study, carried out the literature review, and drafted the manuscript. RS carried out the literature review, and participated in drafting the manuscript. MB critically edited the manuscript. YA conceived of the study and participated in its design and coordination. All authors read and approved the final manuscript.

## Author details

<sup>1</sup>Elmezzi Graduate School of Molecular Medicine, The Feinstein Institute for Medical Research, 350 Community drive, Manhasset, NY 11030, USA. <sup>2</sup>Institute

of Immunology, Philipps University Marburg, Hans-Meerwein-Str., 35043 Marburg, Germany. <sup>3</sup>Center for Molecular Innovation, The Feinstein Institute for Medical Research, 350 Community drive, Manhasset, NY 11030, USA.

Received: 7 February 2015 Accepted: 27 March 2015

Published online: 17 April 2015

## References

- Castellani RJ, Perry G. Pathogenesis and disease-modifying therapy in Alzheimer's disease: the flat line of progress. *Arch Med Res.* 2012;43(8):694–8. doi:10.1016/j.arcmed.2012.09.009.
- Nazem A, Mansoori GA. Nanotechnology solutions for Alzheimer's disease: advances in research tools, diagnostic methods and therapeutic agents. *J Alzheimers Dis.* 2008;13(2):199–223.
- Mangialasche F, Solomon A, Winblad B, Mecocci P, Kivipelto M. Alzheimer's disease: clinical trials and drug development. *Lancet Neurol.* 2010;9(7):702–16. doi:10.1016/s1474-4422(10)70119-8.
- Sabbagh JJ, Kinney JW, Cummings JL. Animal systems in the development of treatments for Alzheimer's disease: challenges, methods, and implications. *Neurobiol Aging.* 2013;34(1):169–83. doi:10.1016/j.neurobiolaging.2012.02.027.
- Krstic D, Madhusudan A, Doehner J, Vogel P, Notter T, Imhof C, et al. Systemic immune challenges trigger and drive Alzheimer-like neuropathology in mice. *J Neuroinflammation.* 2012;9:151. doi:10.1186/1742-2094-9-151.
- Epis R, Gardoni F, Marcello E, Genazzani A, Canonico PL, Di Luca M. Searching for new animal models of Alzheimer's disease. *Eur J Pharmacol.* 2010;626(1):57–63. doi:10.1016/j.ejphar.2009.10.020.
- Duyckaerts C, Potier MC, Delatour B. Alzheimer disease models and human neuropathology: similarities and differences. *Acta Neuropathol.* 2008;115(1):5–38. doi:10.1007/s00401-007-0312-8.
- Chetelat G. Alzheimer disease: Abeta-independent processes-rethinking preclinical AD. *Nat Rev Neurol.* 2013;9(3):123–4. doi:10.1038/nrneuro.2013.21.
- Zlokovic BV. Neurovascular pathways to neurodegeneration in Alzheimer's disease and other disorders. *Nat Rev Neurosci.* 2011;12(12):723–38. doi:10.1038/nrn3114.
- Ankarcrona M, Mangialasche F, Winblad B. Rethinking Alzheimer's disease therapy: are mitochondria the key? *J Alzheimers Dis.* 2010;20 Suppl 2:S579–90. doi:10.3233/jad-2010-100327.
- Oresic M, Hyotylainen T, Herukka SK, Sysi-Aho M, Mattila I, Seppanan-Laakso T, et al. Metabolome in progression to Alzheimer's disease. *Translational Psychiatry.* 2011;1, e57. doi:10.1038/tp.2011.55.
- Kuusisto J, Koivisto K, Mykkanen L, Helkala EL, Vanhanen M, Hanninen T, et al. Association between features of the insulin resistance syndrome and Alzheimer's disease independently of apolipoprotein E4 phenotype: cross sectional population based study. *BMJ.* 1997;315(7115):1045–9.
- Krstic D, Knuesel I. Deciphering the mechanism underlying late-onset Alzheimer disease. *Nat Rev Neurol.* 2013;9(1):25–34. doi:10.1038/nrneuro.2012.236.
- Pizza V, Agresta A, D'Acunzio CW, Festa M, Capasso A. Neuroinflamm-aging and neurodegenerative diseases: an overview. *CNS Neurol Disord Drug Targets.* 2011;10(5):621–34.
- Franceschi C, Capri M, Monti D, Giunta S, Olivieri F, Sevini F, et al. Inflammaging and anti-inflammaging: a systemic perspective on aging and longevity emerged from studies in humans. *Mech Ageing Dev.* 2007;128(1):92–105. doi:10.1016/j.mad.2006.11.016.
- Frank-Cannon TC, Alto LT, McAlpine FE, Tansey MG. Does neuroinflammation fan the flame in neurodegenerative diseases? *Mol Neurodegener.* 2009;4:47. doi:10.1186/1750-1326-4-47.
- Wyss-Coray T. Inflammation in Alzheimer disease: driving force, bystander or beneficial response? *Nat Med.* 2006;12(9):1005–15. doi:10.1038/nm1484.
- Guerreiro R, Wojtas A, Bras J, Carrasquillo M, Rogaeva E, Majounie E, et al. TREM2 variants in Alzheimer's disease. *N Engl J Med.* 2013;368(2):117–27. doi:10.1056/NEJMoa1211851.
- Jonsson T, Stefansson H, Steinberg S, Jonsdottir I, Jonsson PV, Snaedal J, et al. Variant of TREM2 associated with the risk of Alzheimer's disease. *N Engl J Med.* 2013;368(2):107–16. doi:10.1056/NEJMoa1211103.
- Naj AC, Jun G, Beecham GW, Wang LS, Vardarajan BN, Burros J, et al. Common variants at MS4A4/MS4A6E, CD2AP, CD33 and EPHA1 are associated with late-onset Alzheimer's disease. *Nat Genet.* 2011;43(5):436–41. doi:10.1038/ng.801.
- Pugliese L, Tanzi RE, Kovacs DM. Alzheimer's disease: the cholesterol connection. *Nat Neurosci.* 2003;6(4):345–51. doi:10.1038/nn0403-345.

22. Eikelenboom P, van Exel E, Hoozemans JJ, Veerhuis R, Rozemuller AJ, van Gool WA. Neuroinflammation - an early event in both the history and pathogenesis of Alzheimer's disease. *Neurodegener Dis*. 2010;7(1-3):38–41. doi:10.1159/000283480.
23. Ferretti MT, Cuello AC. Does a pro-inflammatory process precede Alzheimer's disease and mild cognitive impairment? *Curr Alzheimer Res*. 2011;8(2):164–74.
24. Buckwalter MS, Wyss-Coray T. Modelling neuroinflammatory phenotypes in vivo. *J Neuroinflammation*. 2004;1(1):10. doi:10.1186/1742-2094-1-10.
25. Frohman EM, Racke MK, Raine CS. Multiple sclerosis - the plaque and its pathogenesis. *N Engl J Med*. 2006;354(9):942–55. doi:10.1056/NEJMr052130.
26. Lehnardt S, Massillon L, Follett P, Jensen FE, Ratan R, Rosenberg PA, et al. Activation of innate immunity in the CNS triggers neurodegeneration through a toll-like receptor 4-dependent pathway. *Proc Natl Acad Sci U S A*. 2003;100(14):8514–9. doi:10.1073/pnas.1432609100.
27. Akiyama H, Barger S, Barnum S, Bradt B, Bauer J, Cole GM, et al. Inflammation and Alzheimer's disease. *Neurobiol Aging*. 2000;21(3):383–421.
28. Walker DG, Dalsing-Hernandez JE, Campbell NA, Lue LF. Decreased expression of CD200 and CD200 receptor in Alzheimer's disease: a potential mechanism leading to chronic inflammation. *Exp Neurol*. 2009;215(1):5–19. doi:10.1016/j.expneurol.2008.09.003.
29. Johnson VE, Stewart JE, Begbie FD, Trojanowski JQ, Smith DH, Stewart W. Inflammation and white matter degeneration persist for years after a single traumatic brain injury. *Brain*. 2013;136(Pt 1):28–42. doi:10.1093/brain/aw322.
30. Xu WL, Atti AR, Gatz M, Pedersen NL, Johansson B, Fratiglioni L. Midlife overweight and obesity increase late-life dementia risk: a population-based twin study. *Neurology*. 2011;76(18):1568–74. doi:10.1212/WNL.0b013e3182190d09.
31. Block ML, Zecca L, Hong JS. Microglia-mediated neurotoxicity: uncovering the molecular mechanisms. *Nat Rev Neurosci*. 2007;8(1):57–69. doi:10.1038/nrn2038.
32. Streit WJ, Xue QS. Human CNS immune senescence and neurodegeneration. *Curr Opin Immunol*. 2014;29:93–6. doi:10.1016/j.coi.2014.05.005.
33. Streit WJ, Braak H, Xue QS, Bechmann I. Dystrophic (senescent) rather than activated microglial cells are associated with tau pathology and likely precede neurodegeneration in Alzheimer's disease. *Acta Neuropathol*. 2009;118(4):475–85. doi:10.1007/s00401-009-0556-6.
34. Wojtera M, Sobow T, Kloszewska I, Liberski PP, Brown DR, Sikorska B. Expression of immunohistochemical markers on microglia in Creutzfeldt-Jakob disease and Alzheimer's disease: morphometric study and review of the literature. *Folia neuropathologica / Association of Polish Neuropathologists and Medical Research Centre, Polish Academy of Sciences*. 2012;50(1):74–84.
35. Li W. Phagocyte dysfunction, tissue aging and degeneration. *Ageing Res Rev*. 2013;12(4):1005–12. doi:10.1016/j.arr.2013.05.006.
36. Caldeira C, Oliveira AF, Cunha C, Vaz AR, Falcao AS, Fernandes A, et al. Microglia change from a reactive to an age-like phenotype with the time in culture. *Front Cell Neurosci*. 2014;8:152. doi:10.3389/fncel.2014.00152.
37. Schwab C, Klegeris A, McGeer PL. Inflammation in transgenic mouse models of neurodegenerative disorders. *Biochim Biophys Acta*. 2010;1802(10):889–902. doi:10.1016/j.bbdis.2009.10.013.
38. Janelins MC, Mastrangelo MA, Oddo S, LaFerla FM, Federoff HJ, Bowers WJ. Early correlation of microglial activation with enhanced tumor necrosis factor- $\alpha$  and monocyte chemoattractant protein-1 expression specifically within the entorhinal cortex of triple transgenic Alzheimer's disease mice. *J Neuroinflammation*. 2005;2:23. doi:10.1186/1742-2094-2-23.
39. Moore AH, Wu M, Shaftel SS, Graham KA, O'Banion MK. Sustained expression of interleukin-1beta in mouse hippocampus impairs spatial memory. *Neuroscience*. 2009;164(4):1484–95. doi:10.1016/j.neuroscience.2009.08.073.
40. Burton M, Johnson R. Interleukin-6 trans-signaling in the senescent mouse brain is involved in infection-related deficits in contextual fear conditioning. *Brain Behav Immun*. 2012;26(5):732–8. doi:10.1016/j.bbi.2011.10.008.
41. McLarnon JG. Correlated inflammatory responses and neurodegeneration in peptide-injected animal models of Alzheimer's disease. *Biomed Res Int*. 2014;2014:923670. doi:10.1155/2014/923670.
42. Kitazawa M, Oddo S, Yamasaki TR, Green KN, LaFerla FM. Lipopolysaccharide-induced inflammation exacerbates tau pathology by a cyclin-dependent kinase 5-mediated pathway in a transgenic model of Alzheimer's disease. *J Neurosci*. 2005;25(39):8843–53. doi:10.1523/JNEUROSCI.2868-05.2005.
43. Hauss-Wegrzyniak B, Dobrzanski P, Stoehr JD, Wenk GL. Chronic neuroinflammation in rats reproduces components of the neurobiology of Alzheimer's disease. *Brain Res*. 1998;780(2):294–303.
44. Sheng JG, Bora SH, Xu G, Borchelt DR, Price DL, Koliatsos VE. Lipopolysaccharide-induced-neuroinflammation increases intracellular accumulation of amyloid precursor protein and amyloid beta peptide in APPswe transgenic mice. *Neurobiol Dis*. 2003;14(1):133–45.
45. Hailman E, Lichenstein HS, Wurfel MM, Miller DS, Johnson DA, Kelley M, et al. Lipopolysaccharide (LPS)-binding protein accelerates the binding of LPS to CD14. *J Exp Med*. 1994;179(1):269–77.
46. Rivest S. Regulation of innate immune responses in the brain. *Nat Rev Immunol*. 2009;9(6):429–39. doi:10.1038/nri2565.
47. Mizuno T, Sawada M, Marunouchi T, Suzumura A. Production of interleukin-10 by mouse glial cells in culture. *Biochem Biophys Res Commun*. 1994;205(3):1907–15. doi:10.1006/bbrc.1994.2893.
48. Welsler-Alves JV, Milner R. Microglia are the major source of TNF- $\alpha$  and TGF- $\beta$ 1 in postnatal glial cultures; regulation by cytokines, lipopolysaccharide, and vitronectin. *Neurochem Int*. 2013;63(1):47–53. doi:10.1016/j.neuint.2013.04.007.
49. Bardou I, Kaercher RM, Brothers HM, Hopp SC, Royer S, Wenk GL. Age and duration of inflammatory environment differentially affect the neuroimmune response and catecholaminergic neurons in the midbrain and brainstem. *Neurobiol Aging*. 2014;35(5):1065–73. doi:10.1016/j.neurobiolaging.2013.11.006.
50. Lee JW, Lee YK, Yuk DY, Choi DY, Ban SB, Oh KW, et al. Neuro-inflammation induced by lipopolysaccharide causes cognitive impairment through enhancement of beta-amyloid generation. *J Neuroinflammation*. 2008;5:37. doi:10.1186/1742-2094-5-37.
51. Czerniawski J, Miyashita T, Lewandowski G, Guzowski JF. Systemic lipopolysaccharide administration impairs retrieval of context-object discrimination, but not spatial, memory: evidence for selective disruption of specific hippocampus-dependent memory functions during acute neuroinflammation. *Brain Behav Immun*. 2015;44:159–66. doi:10.1016/j.bbi.2014.09.014.
52. Grudzien A, Shaw P, Weintraub S, Bigio E, Mash DC, Mesulam MM. Locus coeruleus neurofibrillary degeneration in aging, mild cognitive impairment and early Alzheimer's disease. *Neurobiol Aging*. 2007;28(3):327–35. doi:10.1016/j.neurobiolaging.2006.02.007.
53. Cerbai F, Lana D, Nosi D, Petkova-Kirova P, Zecchi S, Brothers H, et al. The neuron-astrocyte-microglia triad in normal brain ageing and in a model of neuroinflammation in the rat hippocampus. *PLoS One*. 2012;7(9), e45250.
54. Hauss-Wegrzyniak B, Vannucchi MG, Wenk GL. Behavioral and ultrastructural changes induced by chronic neuroinflammation in young rats. *Brain Res*. 2000;859(1):157–66.
55. Wenk GL, McGann K, Mencarelli A, Hauss-Wegrzyniak B, Del Soldato P, Fiorucci S. Mechanisms to prevent the toxicity of chronic neuroinflammation on forebrain cholinergic neurons. *Eur J Pharmacol*. 2000;402(1-2):77–85.
56. Hafting T, Fyhn M, Molden S, Moser MB, Moser EI. Microstructure of a spatial map in the entorhinal cortex. *Nature*. 2005;436(7052):801–6. doi:10.1038/nature03721.
57. Hauss-Wegrzyniak B, Lynch MA, Vraniak PD, Wenk GL. Chronic brain inflammation results in cell loss in the entorhinal cortex and impaired LTP in perforant path-granule cell synapses. *Exp Neurol*. 2002;176(2):336–41.
58. DiCarlo G, Wilcock D, Henderson D, Gordon M, Morgan D. Intrahippocampal LPS injections reduce Abeta load in APP + PS1 transgenic mice. *Neurobiol Aging*. 2001;22(6):1007–12.
59. Deng XH, Ai WM, Lei DL, Luo XG, Yan XX, Li Z. Lipopolysaccharide induces paired immunoglobulin-like receptor B (PirB) expression, synaptic alteration, and learning-memory deficit in rats. *Neuroscience*. 2012;209:161–70. doi:10.1016/j.neuroscience.2012.02.022.
60. Liu Y, Qin L, Wilson B, Wu X, Qian L, Granholm AC, et al. Endotoxin induces a delayed loss of TH-IR neurons in substantia nigra and motor behavioral deficits. *Neurotoxicology*. 2008;29(5):864–70. doi:10.1016/j.neuro.2008.02.014.
61. Perry VH, Nicoll JA, Holmes C. Microglia in neurodegenerative disease. *Nat Rev Neurol*. 2010;6(4):193–201. doi:10.1038/nrneuro.2010.17.
62. Perry VH. Contribution of systemic inflammation to chronic neurodegeneration. *Acta Neuropathol*. 2010;120(3):277–86. doi:10.1007/s00401-010-0722-x.
63. Town T, Jeng D, Alexopoulou L, Tan J, Flavell RA, Alexopoulou L, et al. Microglia recognize double-stranded RNA via TLR3 recognition of double-stranded RNA and activation of NF- $\kappa$ B by toll-like receptor 3. *J Immunol*. 2006;176(6):3804–12. doi:10.1038/35099560.
64. De Miranda J, Yaddanapudi K, Hornig M, Villar G, Serge R, Lipkin WI. Induction of toll-like receptor 3-mediated immunity during gestation inhibits cortical neurogenesis and causes behavioral disturbances. *MBio*. 2010;1(4). doi:10.1128/mBio.00176-10.
65. Kimura M, Toth LA, Agostini H, Cady AB, Majde JA, Krueger JM. Comparison of acute phase responses induced in rabbits by lipopolysaccharide and double-stranded RNA. *Am J Physiol*. 1994;267(6 Pt 2):R1596–605.

66. Meyer U, Nyffeler M, Engler A, Urwyler A, Schedlowski M, Knuesel I, et al. The time of prenatal immune challenge determines the specificity of inflammation-mediated brain and behavioral pathology. *J Neurosci*. 2006;26(18):4752–62. doi:10.1523/JNEUROSCI.0099-06.2006.
67. Bilbo SD, Smith SH, Schwarz JM. A lifespan approach to neuroinflammatory and cognitive disorders: a critical role for glia. *J Neuroimmune Pharmacol*. 2012;7(1):24–41. doi:10.1007/s11481-011-9299-y.
68. Bitanihirwe BK, Peleg-Raibstein D, Mouttet F, Feldon J, Meyer U. Late prenatal immune activation in mice leads to behavioral and neurochemical abnormalities relevant to the negative symptoms of schizophrenia. *Neuropsychopharmacology*. 2010;35(12):2462–78. doi:10.1038/npp.2010.129.
69. Hoyer S, Nitsch R. Cerebral excess release of neurotransmitter amino acids subsequent to reduced cerebral glucose metabolism in early-onset dementia of Alzheimer type. *J Neural Transm (Vienna, Austria)*. 1989;75(3):227–32.
70. Hoyer S, Nitsch R, Oesterreich K. Predominant abnormality in cerebral glucose utilization in late-onset dementia of the Alzheimer type: a cross-sectional comparison against advanced late-onset and incipient early-onset cases. *J Neural Transm Park Dis Dement Sect*. 1991;3(1):1–14.
71. Craft S, Zallen G, Baker LD. Glucose and memory in mild senile dementia of the Alzheimer type. *J Clin Exp Neuropsychol*. 1992;14(2):253–67. doi:10.1080/01688639208402827.
72. Sims-Robinson C, Kim B, Rosko A, Feldman EL. How does diabetes accelerate Alzheimer disease pathology? *Nat Rev Neurol*. 2010;6(10):551–9. doi:10.1038/nrneuro.2010.130.
73. Yamamoto H, Uchigata Y, Okamoto H. Streptozotocin and alloxan induce DNA strand breaks and poly(ADP-ribose) synthetase in pancreatic islets. *Nature*. 1981;294(5838):284–6.
74. Mansford KR, Opie L. Comparison of metabolic abnormalities in diabetes mellitus induced by streptozotocin or by alloxan. *Lancet*. 1968;1(7544):670–1.
75. Like AA, Rossini AA. Streptozotocin-induced pancreatic insulinitis: new model of diabetes mellitus. *Science*. 1976;193(4251):415–7.
76. Stranahan AM, Arumugam TV, Cutler RG, Lee K, Egan JM, Mattson MP. Diabetes impairs hippocampal function through glucocorticoid-mediated effects on new and mature neurons. *Nat Neurosci*. 2008;11(3):309–17. doi:10.1038/nn2055.
77. Wang JQ, Yin J, Song YF, Zhang L, Ren YX, Wang DG, et al. Brain aging and AD-like pathology in streptozotocin-induced diabetic rats. *J Diabetes Res*. 2014;2014:796840. doi:10.1155/2014/796840.
78. Lenzen S. The mechanisms of alloxan- and streptozotocin-induced diabetes. *Diabetologia*. 2008;51(2):216–26. doi:10.1007/s00125-007-0886-7.
79. Szkudelski T. The mechanism of alloxan and streptozotocin action in B cells of the rat pancreas. *Physiol Res*. 2001;50(6):537–46.
80. Takasu N, Komiya I, Asawa T, Nagasawa Y, Yamada T. Streptozotocin- and alloxan-induced H<sub>2</sub>O<sub>2</sub> generation and DNA fragmentation in pancreatic islets: H<sub>2</sub>O<sub>2</sub> as mediator for DNA fragmentation. *Diabetes*. 1991;40(9):1141–5.
81. Turk J, Corbett JA, Ramanadham S, Bohrer A, McDaniel ML. Biochemical evidence for nitric oxide formation from streptozotocin in isolated pancreatic islets. *Biochem Biophys Res Commun*. 1993;197(3):1458–64. doi:10.1006/bbrc.1993.2641.
82. Murata M, Takahashi A, Saito I, Kawanishi S. Site-specific DNA methylation and apoptosis: induction by diabetogenic streptozotocin. *Biochem Pharmacol*. 1999;57(8):881–7.
83. Pieper AA, Brat DJ, Krug DK, Watkins CC, Gupta A, Blackshaw S, et al. Poly (ADP-ribose) polymerase-deficient mice are protected from streptozotocin-induced diabetes. *Proc Natl Acad Sci U S A*. 1999;96(6):3059–64.
84. Mensah-Brown EP, Obineche EN, Galadari S, Chandranath E, Shahin A, Ahmed I, et al. Streptozotocin-induced diabetic nephropathy in rats: the role of inflammatory cytokines. *Cytokine*. 2005;31(3):180–90. doi:10.1016/j.cyt.2005.04.006.
85. Chen YLZ, Blanchard J, Dai CL, Sun S, Lee MH, Grundke-Iqbal I, et al. A non-transgenic mouse model (icv-STZ Mouse) of Alzheimer's disease: similarities to and differences from the transgenic model (3xTg-AD Mouse). *Mol Neurobiol*. 2013;47(2):711–25.
86. Nitsch R, Hoyer S. Local action of the diabetogenic drug, streptozotocin, on glucose and energy metabolism in rat brain cortex. *Neurosci Lett*. 1991;128(2):199–202.
87. Plaschke K, Hoyer S. Action of the diabetogenic drug streptozotocin on glycolytic and glycogenolytic metabolism in adult rat brain cortex and hippocampus. *Int J Dev Neurosci*. 1993;11(4):477–83.
88. Lester-Coll N, Rivera EJ, Soscia SJ, Doiron K, Wands JR, de la Monte SM. Intracerebral streptozotocin model of type 3 diabetes: relevance to sporadic Alzheimer's disease. *J Alzheimer's Disease: JAD*. 2006;9(1):13–33.
89. Kraska ASM, Dorieux O, Joseph-Mathurin N, Bourrin E, Petit F, Jan C, et al. In vivo cross-sectional characterization of cerebral alterations induced by intracerebroventricular administration of streptozotocin. *PLoS One*. 2012;7(9), e46196. doi:10.1371/journal.pone.0046196.
90. Chen Y, Liang Z, Tian Z, Blanchard J, Dai CL, Chalbot S, et al. Intracerebroventricular streptozotocin exacerbates Alzheimer-like changes of 3xTg-AD mice. *Mol Neurobiol*. 2014;49(1):547–62. doi:10.1007/s12035-013-8539-y.
91. Plaschke K, Kopitz J, Siegelin M, Schliebs R, Salkovic-Petrisic M, Riederer P, et al. Insulin-resistant brain state after intracerebroventricular streptozotocin injection exacerbates Alzheimer-like changes in Tg2576 AbetaPP-overexpressing mice. *J Alzheimers Dis*. 2010;19(2):691–704. doi:10.3233/jad-2010-1270.
92. Ke YD, Delerue F, Gladbach A, Götz J, Ittner LM. Experimental diabetes mellitus exacerbates tau pathology in a transgenic mouse model of Alzheimer's disease. *PLoS One*. 2009;4(11), e7917. doi:10.1371/journal.pone.0007917.
93. Devi L, Alldred MJ, Ginsberg SD, Ohno M. Mechanisms underlying insulin deficiency-induced acceleration of  $\beta$ -amyloidosis in a mouse model of Alzheimer's disease. *PLoS One*. 2012;7(3):e32792. doi:10.1371/journal.pone.0032792.
94. Jolival CG, Hurford R, Lee CA, Dumaop W, Rockenstein E, Masliah E. Type 1 diabetes exaggerates features of Alzheimer's disease in APP transgenic mice. *Exp Neurol*. 2010;223(2):422–31. doi:10.1016/j.expneurol.2009.11.005.
95. Liu P, Zou LB, Wang LH, Jiao Q, Chi TY, Ji XF, et al. Xanthoceraside attenuates tau hyperphosphorylation and cognitive deficits in intracerebroventricular-streptozotocin injected rats. *Psychopharmacology (Berl)*. 2013. doi:10.1007/s00213-013-3240-4.
96. Salkovic-Petrisic M, Osmanovic-Barilar J, Brückner M, Hoyer S, Arendt T, Riederer P. Cerebral amyloid angiopathy in streptozotocin rat model of sporadic Alzheimer's disease: a long-term follow up study. *J Neural Transm*. 2011;118(5):765–72.
97. Grünblatt ES-PM, Osmanovic J, Riederer P, Hoyer S. Brain insulin system dysfunction in streptozotocin intracerebroventricularly treated rats generates hyperphosphorylated tau protein. *J Neurochem*. 2007;101(3):757–70.
98. Tapia R, Pena F, Arias C. Neurotoxic and synaptic effects of okadaic acid, an inhibitor of protein phosphatases. *Neurochem Res*. 1999;24(11):1423–30.
99. Sontag JM, Sontag E. Protein phosphatase 2A dysfunction in Alzheimer's disease. *Front Mol Neurosci*. 2014;7:16. doi:10.3389/fnmol.2014.00016.
100. Cohen PT, Brewis ND, Hughes V, Mann DJ. Protein serine/threonine phosphatases: an expanding family. *FEBS Lett*. 1990;268(2):355–9.
101. Kamat PK, Tota S, Rai S, Shukla R, Ali S, Najmi AK, et al. Okadaic acid induced neurotoxicity leads to central cholinergic dysfunction in rats. *Eur J Pharmacol*. 2012;690(1-3):90–8. doi:10.1016/j.ejphar.2012.06.006.
102. Costa AP, Tramontina AC, Biasibetti R, Batassini C, Lopes MW, Wartchow KM, et al. Neuroglial alterations in rats submitted to the okadaic acid-induced model of dementia. *Behav Brain Res*. 2012;226(2):420–7. doi:10.1016/j.bbr.2011.09.035.
103. Kamat PK, Tota S, Saxena G, Shukla R, Nath C. Okadaic acid (ICV) induced memory impairment in rats: a suitable experimental model to test anti-dementia activity. *Brain Res*. 2010;1309:66–74. doi:10.1016/j.brainres.2009.10.064.
104. Kamat PK, Rai S, Swarnkar S, Shukla R, Nath C. Molecular and cellular mechanism of okadaic acid (OKA)-induced neurotoxicity: a novel tool for Alzheimer's disease therapeutic application. *Mol Neurobiol*. 2014;50(3):852–65. doi:10.1007/s12035-014-8699-4.
105. Arendt T, Holzer M, Fruth R, Bruckner MK, Gartner U. Phosphorylation of tau, Abeta-formation, and apoptosis after in vivo inhibition of PP-1 and PP-2A. *Neurobiol Aging*. 1998;19(1):3–13.
106. Lee J, Hong H, Im J, Byun H, Kim D. The formation of PHF-1 and SMI-31 positive dystrophic neurites in rat hippocampus following acute injection of okadaic acid. *Neurosci Lett*. 2000;282(1-2):49–52.
107. Kamat PK, Tota S, Rai S, Swarnkar S, Shukla R, Nath C. A study on neuroinflammatory marker in brain areas of okadaic acid (ICV) induced memory impaired rats. *Life Sci*. 2012;90(19-20):713–20. doi:10.1016/j.lfs.2012.03.012.
108. Rajasekar N, Dwivedi S, Tota SK, Kamat PK, Hanif K, Nath C, et al. Neuroprotective effect of curcumin on okadaic acid induced memory impairment in mice. *Eur J Pharmacol*. 2013;715(1-3):381–94. doi:10.1016/j.ejphar.2013.04.033.
109. Kamat PK, Rai S, Swarnkar S, Shukla R, Ali S, Najmi AK, et al. Okadaic acid-induced tau phosphorylation in rat brain: role of NMDA receptor. *Neuroscience*. 2013;238:97–113. doi:10.1016/j.neuroscience.2013.01.075.
110. Zimmer ER, Kalinina E, Haas CB, Torrez VR, Souza DO, Muller AP, et al. Pretreatment with memantine prevents Alzheimer-like alterations induced by intrahippocampal okadaic acid administration in rats. *Curr Alzheimer Res*. 2012;9(10):1182–90.

111. Kamat PK, Rai S, Swarnkar S, Shukla R, Nath C. Mechanism of synapse redox stress in okadaic acid (ICV) induced memory impairment: role of NMDA receptor. *Neurochem Int.* 2014;76:32–41. doi:10.1016/j.neuint.2014.06.012.
112. Valdiglesias V, Laffon B, Pasaro E, Cemeli E, Anderson D, Mendez J. Induction of oxidative DNA damage by the marine toxin okadaic acid depends on human cell type. *Toxicol.* 2011;57(6):882–8. doi:10.1016/j.toxicol.2011.03.005.
113. Kumar A, Seghal N, Naidu PS, Padi SS, Goyal R. Colchicines-induced neurotoxicity as an animal model of sporadic dementia of Alzheimer's type. *Pharmacol Rep.* 2007;59(3):274–83.
114. Nakayama T, Sawada T. Involvement of microtubule integrity in memory impairment caused by colchicine. *Pharmacol Biochem Behav.* 2002;71(1-2):119–38.
115. Tilson HA, Rogers BC, Grimes L, Harry GJ, Peterson NJ, Hong JS, et al. Time-dependent neurobiological effects of colchicine administered directly into the hippocampus of rats. *Brain Res.* 1987;408(1-2):163–72.
116. Ho L, Osaka H, Aisen PS, Pasinetti GM. Induction of cyclooxygenase (COX)-2 but not COX-1 gene expression in apoptotic cell death. *J Neuroimmunol.* 1998;89(1-2):142–9.
117. Sil S, Goswami AR, Dutta G, Ghosh T. Effects of naproxen on immune responses in a colchicine-induced rat model of Alzheimer's disease. *Neuroimmunomodulation.* 2014;21(6):304–21. doi:10.1159/000357735.
118. Geddes JW, Bondada V, Keller JN. Effects of intrahippocampal colchicine administration on the levels and localization of microtubule-associated proteins, tau and MAP2. *Brain Res.* 1994;633(1-2):1–8.
119. McMartin DN, Schedlbauer LM. Effect of experimental colchicine encephalopathy on brain protein synthesis and tubulin metabolism. *J Neurobiol.* 1978;9(6):453–63. doi:10.1002/neu.480090605.
120. Merrick SE, Demoise DC, Lee VM. Site-specific dephosphorylation of tau protein at Ser202/Thr205 in response to microtubule depolymerization in cultured human neurons involves protein phosphatase 2A. *J Biol Chem.* 1996;271(10):5589–94.
121. Shaftel SS, Griffin WS, O'Banion MK. The role of interleukin-1 in neuroinflammation and Alzheimer disease: an evolving perspective. *J Neuroinflammation.* 2008;5:7. doi:10.1186/1742-2094-5-7.
122. Alboni S, Cervia D, Sugama S, Conti B. Interleukin 18 in the CNS. *J Neuroinflammation.* 2010;7:9. doi:10.1186/1742-2094-7-9.
123. Griffin WS, Stanley LC, Ling C, White L, MacLeod V, Perrot LJ, et al. Brain interleukin 1 and S-100 immunoreactivity are elevated in Down syndrome and Alzheimer disease. *Proc Natl Acad Sci U S A.* 1989;86(19):7611–5.
124. Ojala J, Alafuzoff I, Herukka SK, van Groen T, Tanila H, Pirttilä T. Expression of interleukin-18 is increased in the brains of Alzheimer's disease patients. *Neurobiol Aging.* 2009;30(2):198–209. doi:10.1016/j.neurobiolaging.2007.06.006.
125. Malaguarnera L, Motta M, Di Rosa M, Anzaldi M, Malaguarnera M. Interleukin-18 and transforming growth factor-beta 1 plasma levels in Alzheimer's disease and vascular dementia. *Neuropathology.* 2006;26(4):307–12.
126. Shaftel SS, Kyrkanides S, Olschowka JA, Miller JN, Johnson RE, O'Banion MK. Sustained hippocampal IL-1 beta overexpression mediates chronic neuroinflammation and ameliorates Alzheimer plaque pathology. *J Clin Invest.* 2007;117(6):1595–604. doi:10.1172/jci31450.
127. Matousek SB, Ghosh S, Shaftel SS, Kyrkanides S, Olschowka JA, O'Banion MK. Chronic IL-1beta-mediated neuroinflammation mitigates amyloid pathology in a mouse model of Alzheimer's disease without inducing overt neurodegeneration. *J Neuroimmune Pharmacol.* 2012;7(1):156–64. doi:10.1007/s11481-011-9331-2.
128. Ghosh S, Wu MD, Shaftel SS, Kyrkanides S, Laferla FM, Olschowka JA, et al. Sustained interleukin-1beta overexpression exacerbates tau pathology despite reduced amyloid burden in an Alzheimer's mouse model. *J Neurosci.* 2013;33:5053–64.
129. Neumann H, Kotter MR, Franklin RJ. Debris clearance by microglia: an essential link between degeneration and regeneration. *Brain.* 2009;132(Pt 2):288–95. doi:10.1093/brain/awn109.
130. Norden DM, Muccigrosso MM, Godbout JP. Microglial Priming and Enhanced Reactivity to Secondary Insult in Aging, and Traumatic CNS Injury, and Neurodegenerative Disease. *Neuropharmacology.* 2014. pii: S0028-3908(14)00403-1. doi: 10.1016/j.neuropharm.2014.10.028.
131. Smith MZ, Nagy Z, Esiri MM. Cell cycle-related protein expression in vascular dementia and Alzheimer's disease. *Neurosci Lett.* 1999;271(1):45–8.
132. Yang Y, Geldmacher DS, Herrup K. DNA replication precedes neuronal cell death in Alzheimer's disease. *J Neurosci.* 2001;21(8):2661–8.
133. Smith TW, Lippa CF. Ki-67 immunoreactivity in Alzheimer's disease and other neurodegenerative disorders. *J Neuropathol Exp Neurol.* 1995;54(3):297–303.
134. Busser J, Geldmacher DS, Herrup K. Ectopic cell cycle proteins predict the sites of neuronal cell death in Alzheimer's disease brain. *J Neurosci.* 1998;18(8):2801–7.
135. Morgan DO. Cyclin-dependent kinases: engines, clocks, and microprocessors. *Annu Rev Cell Dev Biol.* 1997;13:261–91. doi: 10.1146/annurev.cellbio.13.1.261.
136. Nikolic M, Dudek H, Kwon YT, Ramos YF, Tsai LH. The cdk5/p35 kinase is essential for neurite outgrowth during neuronal differentiation. *Genes Dev.* 1996;10(7):816–25.
137. Hellmich MR, Pant HC, Wada E, Battey JF. Neuronal cdc2-like kinase: a cdc2-related protein kinase with predominantly neuronal expression. *Proc Natl Acad Sci U S A.* 1992;89(22):10867–71.
138. Tsai LH, Delalle I, Caviness VS, Chae T, Harlow E. p35 is a neural-specific regulatory subunit of cyclin-dependent kinase 5. *Nature.* 1994;371(6496):419–23. doi:10.1038/371419a0.
139. Patrick GN, Zukerberg L, Nikolic M, de la Monte S, Dikkes P, Tsai LH. Conversion of p35 to p25 deregulates Cdk5 activity and promotes neurodegeneration. *Nature.* 1999;402(6762):615–22. doi:10.1038/45159.
140. Lee MS, Kwon YT, Li M, Peng J, Friedlander RM, Tsai LH. Neurotoxicity induces cleavage of p35 to p25 by calpain. *Nature.* 2000;405(6784):360–4. doi:10.1038/35012636.
141. Ahijanian MK, Barrezueta NX, Williams RD, Jakowski A, Kowz KP, McCarthy S, et al. Hyperphosphorylated tau and neurofilament and cytoskeletal disruptions in mice overexpressing human p25, an activator of cdk5. *Proc Natl Acad Sci U S A.* 2000;97(6):2910–5.
142. Monaco EA. Recent evidence regarding a role for Cdk5 dysregulation in Alzheimer's disease. *Curr Alzheimer Res.* 2004;1(1):33–8.
143. Sundaram JR, Chan ES, Poore CP, Pareek TK, Cheong WF, Shui G, et al. Cdk5/p25-induced cytosolic PLA2-mediated lysophosphatidylcholine production regulates neuroinflammation and triggers neurodegeneration. *J Neurosci.* 2012;32(3):1020–34. doi: 10.523/JNEUROSCI.5177-11.2012.
144. Fischer A, Sananbenesi F, Pang PT, Lu B, Tsai LH. Opposing roles of transient and prolonged expression of p25 in synaptic plasticity and hippocampus-dependent memory. *Neuron.* 2005;48(5):825–38.
145. Muyliaert D, Terwel D, Kremer A, Sennvik K, Borghgraef P, Devijver H, et al. Neurodegeneration and neuroinflammation in cdk5/p25-inducible mice: a model for hippocampal sclerosis and neocortical degeneration. *Am J Pathol.* 2008;172(2):470–85. doi:10.2353/ajpath.008.070693. Epub 2008 Jan 17.
146. Houeland G, Romani A, Marchetti C, Amato G, Capsoni S, Cattaneo A, et al. Transgenic mice with chronic NGF deprivation and Alzheimer's disease-like pathology display hippocampal region-specific impairments in short- and long-term plasticities. *J Neurosci.* 2010;30(39):13089–94. doi:10.1523/jneurosci.0457-10.2010.
147. Capsoni S, Tiveron C, Vignone D, Amato G, Cattaneo A. Dissecting the involvement of tropomyosin-related kinase A and p75 neurotrophin receptor signaling in NGF deficit-induced neurodegeneration. *Proc Natl Acad Sci U S A.* 2010;107(27):12299–304. doi:10.1073/pnas.1007181107.
148. De Rosa R, Garcia AA, Braschi C, Capsoni S, Maffei L, Berardi N, et al. Intranasal administration of nerve growth factor (NGF) rescues recognition memory deficits in AD11 anti-NGF transgenic mice. *Proc Natl Acad Sci U S A.* 2005;102(10):3811–6. doi:10.1073/pnas.0500195102.
149. Capsoni S, Giannotta S, Cattaneo A. Beta-amyloid plaques in a model for sporadic Alzheimer's disease based on transgenic anti-nerve growth factor antibodies. *Mol Cell Neurosci.* 2002;21(1):15–28.
150. Capsoni S, Cattaneo A. On the molecular basis linking nerve growth factor (NGF) to Alzheimer's disease. *Cell Mol Neurobiol.* 2006;26(4-6):619–33. doi:10.1007/s10571-006-9112-2.
151. D'Onofrio M, Arisi I, Brandi R, Di Mambro A, Felsani A, Capsoni S, et al. Early inflammation and immune response mRNAs in the brain of AD11 anti-NGF mice. *Neurobiol Aging.* 2011;32(6):1007–22. doi:10.1016/j.neurobiolaging.2009.05.023.
152. Capsoni S, Brandi R, Arisi I, D'Onofrio M, Cattaneo A. A dual mechanism linking NGF/proNGF imbalance and early inflammation to Alzheimer's disease neurodegeneration in the AD11 anti-NGF mouse model. *CNS Neurol Disord Drug Targets.* 2011;10(5):635–47.
153. Capsoni S, Tiveron C, Amato G, Vignone D, Cattaneo A. Peripheral neutralization of nerve growth factor induces immunosympathectomy and central neurodegeneration in transgenic mice. *J Alzheimers Dis.* 2010;20(2):527–46. doi:10.3233/jad-2010-091357.
154. Tzanoulina S, Brandi R, Arisi I, D'Onofrio M, Urfer SM, Sandi C, et al. Pathogen-free husbandry conditions alleviate behavioral deficits and

- neurodegeneration in AD10 anti-NGF mice. *J Alzheimers Dis.* 2014;38(4):951–64. doi:10.3233/jad-131037.
155. Flanders KC, Ren RF, Lippa CF. Transforming growth factor-betas in neurodegenerative disease. *Prog Neurobiol.* 1998;54(1):71–85.
  156. Ueberham U, Ueberham E, Bruckner MK, Seeger G, Gartner U, Gruschka H, et al. Inducible neuronal expression of transgenic TGF-beta1 in vivo: dissection of short-term and long-term effects. *Eur J Neurosci.* 2005;22(1):50–64. doi:10.1111/j.1460-9568.2005.04189.x.
  157. Unsicker K, Kriegstein K. TGF-betas and their roles in the regulation of neuron survival. *Adv Exp Med Biol.* 2002;513:353–74.
  158. Grammas P, Ovase R. Cerebrovascular transforming growth factor-beta contributes to inflammation in the Alzheimer's disease brain. *Am J Pathol.* 2002;160(5):1583–7.
  159. Wyss-Coray T, Lin C, Yan F, Yu GQ, Rohde M, McConlogue L, et al. TGF-beta1 promotes microglial amyloid-beta clearance and reduces plaque burden in transgenic mice. *Nat Med.* 2001;7(5):612–8. doi:10.1038/87945.
  160. Brionne TC, Tesseur I, Masliah E, Wyss-Coray T. Loss of TGF-beta 1 leads to increased neuronal cell death and microgliosis in mouse brain. *Neuron.* 2003;40(6):1133–45.
  161. Wyss-Coray T, Mucke L. Inflammation in neurodegenerative disease - a double-edged sword. *Neuron.* 2002;35(3):419–32.
  162. Dhull DK, Jindal A, Dhull RK, Aggarwal S, Bhateja D, Padi SS. Neuroprotective effect of cyclooxygenase inhibitors in ICV-STZ induced sporadic Alzheimer's disease in rats. *J Mol Neurosci.* 2012;46(1):223–35. doi:10.1007/s12031-011-9583-6.
  163. Grundke-Iqbal I, Iqbal K, Tung YC, Quinlan M, Wisniewski HM, Binder LI. Abnormal phosphorylation of the microtubule-associated protein tau (tau) in Alzheimer cytoskeletal pathology. *Proc Natl Acad Sci U S A.* 1986;83(13):4913–7.
  164. Jaworski T, Dewachter I, Seymour CM, Borghgraef P, Devijver H, Kugler S, et al. Alzheimer's disease: old problem, new views from transgenic and viral models. *Biochim Biophys Acta.* 2010;1802(10):808–18. doi:10.1016/j.bbadis.2010.03.005.
  165. Pintado C, Gavilan MP, Gavilan E, Garcia-Cuervo L, Gutierrez A, Vitorica J, et al. Lipopolysaccharide-induced neuroinflammation leads to the accumulation of ubiquitinated proteins and increases susceptibility to neurodegeneration induced by proteasome inhibition in rat hippocampus. *J Neuroinflammation.* 2012;9(1):87. doi:10.1186/1742-2094-9-87.
  166. Heneka MT, Nadrigny F, Regen T, Martinez-Hernandez A, Dumitrescu-Ozimek L, Terwel D, et al. Locus ceruleus controls Alzheimer's disease pathology by modulating microglial functions through norepinephrine. *Proc Natl Acad Sci U S A.* 2010;107(13):6058–63. doi:10.1073/pnas.0909586107.
  167. Maia LF, Kaeser SA, Reichwald J, Hruscha M, Martus P, Staufenbiel M, et al. Changes in amyloid-beta and tau in the cerebrospinal fluid of transgenic mice overexpressing amyloid precursor protein. *Sci Transl Med.* 2013;5(194):194re2. doi: 10.1126/scitranslmed.3006446.

**Submit your next manuscript to BioMed Central  
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

