

Role is in the eye of the beholder—the multiple functions of the antibacterial compound tropodithietic acid produced by marine *Rhodobacteraceae*

Nathalie N. S. E. Henriksen [†], Laura L. Lindqvist [†], Mario Wibowo [†], Eva C. Sonnenschein [†], Mikkel Bentzon-Tilia [†], Lone Gram [†]

Department of Biotechnology and Biomedicine, Technical University of Denmark, Søtofts, Plads Bldg. 221, DK-2800 Kgs. Lyngby, Denmark

*Corresponding author: Department of Bioechnology and Biomedicine, Technical University of Denmark, Søtofts Plads Bldg. 221, DK-2800 Kgs. Lyngby, Denmark.

Tel: +45 23688295; E-mail: gram@bio.dtu.dk

One sentence summary: Review of the multiple roles and functions of the secondary metabolite, tropodithietic acid, produced by some marine *Rhodobacteraceae*.

[†]These authors contributed equally.

Editor: Miguel Camara

Abstract

Many microbial secondary metabolites have been studied for decades primarily because of their antimicrobial properties. However, several of these metabolites also possess nonantimicrobial functions, both influencing the physiology of the producer and their ecological neighbors. An example of a versatile bacterial secondary metabolite with multiple functions is the tropone derivative tropodithietic acid (TDA). TDA is a broad-spectrum antimicrobial compound produced by several members of the *Rhodobacteraceae* family, a major marine bacterial lineage, within the genera *Phaeobacter*, *Tritonibacter*, and *Pseudovibrio*. The production of TDA is governed by the mode of growth and influenced by the availability of nutrient sources. The antibacterial effect of TDA is caused by disruption of the proton motive force of target microorganisms and, potentially, by its iron-chelating properties. TDA also acts as a signaling molecule, affecting gene expression in other bacteria, and altering phenotypic traits such as motility, biofilm formation, and antibiotic production in the producer. In microbial communities, TDA-producing bacteria cause a reduction of the relative abundance of closely related species and some fast-growing heterotrophic bacteria. Here, we summarize the current understanding of the chemical ecology of TDA, including the environmental niches of TDA-producing bacteria, and the molecular mechanisms governing the function and regulation of TDA.

Keywords: antimicrobials, secondary metabolites, *Rhodobacteraceae*, tropodithietic acid, marine microbiomes

Introduction

Since the discovery of penicillin by Alexander Flemming in 1929 (Fleming 1929), humanity has benefited from its antimicrobial effects and an array of similar bioactive compounds produced by microorganisms. These molecules, commonly referred to as secondary metabolites, have been studied for decades because of their antimicrobial properties and their use as anti-infective agents in the clinic. It has been the perception that the antibiotic properties also define their predominant function and role in natural microbial niches—as weapons to kill off competitors (Traxler and Kolter 2015). However, sublethal concentrations of antibiotics can influence gene expression in exposed microorganisms and result in changes in phenotypes such as biofilm formation or motility (Goh *et al.* 2002; Linares *et al.* 2006; Straight, Willey and Kolter 2006; Liu *et al.* 2013). These effects have predominantly been observed in bacteria exposed to ‘external’ antimicrobials, but it is not known if these effects are caused indirectly by induction of a stress response or by a direct effect of the compounds on cellular targets (Romero *et al.* 2011; Foster and Bell 2012; Cornforth and Foster 2013; Yoon and Nodwell 2014; Dittmann *et al.* 2019a; Li *et al.* 2021). Some antimicrobial secondary metabolites can also modulate gene expression in the producer itself, acting as signaling molecules mediating quorum sensing (QS; Romero *et al.*

2011; Beyersmann *et al.* 2017). Other secondary metabolites have nonantibiotic functions serving as iron scavengers (siderophores), in predator defense, as antivirulence compounds, or as promoters of horizontal gene transfer (Mansson *et al.* 2011; Seyedsayamdost *et al.* 2011b; Briand *et al.* 2016; Zhang *et al.* 2016, 2021; Danevčič *et al.* 2021). Thus, this challenges the original perception of antibiotic secondary metabolites being predominantly involved in direct interference competition between microbes, whilst not being required for growth and metabolism (Linares *et al.* 2006; Yim, Huimi Wang and Davies 2007; Davies 2013; Pishchany and Kolter 2020). In contrast, it has been argued that the physiological effects at subinhibitory concentrations serve to prime the recipient for competition to come, supporting the natural role of these compounds in interference competition (Foster and Bell 2012; Cornforth and Foster 2013; Abrudan *et al.* 2015). Despite the different perceptions of the predominant role of antimicrobial secondary metabolites, if a single role exists, one concept may not overrule the other (Firn and Jones 2000). However, it is evident that the physiological and ecological function of microbial secondary metabolites should be re-examined in the producing organism and in natural systems in an ecological context. Fortunately, the ‘-omics’ era has facilitated global studies of complex microbial communities, also in the presence of a host organism. In addition, the development of

Received: September 24, 2021. Revised: January 24, 2022. Accepted: January 27, 2022

© The Author(s) 2022. Published by Oxford University Press on behalf of FEMS. This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

in situ chemical detection of metabolites, for instance by mass spectrometry imaging, allows direct, high resolution analyses of secondary metabolites not hampered by bulk extraction, and enabling analysis of spatial metabolomes (Moree et al. 2012; Geier et al. 2020).

An example of an antimicrobial secondary metabolite with multiple functions and roles is the redox active metabolite pyocyanin, produced by *Pseudomonas aeruginosa*. It serves as an antimicrobial, but also as a respiratory pigment and a quorum-related molecule (Hernandez and Newman 2001). Similarly, in *Streptomyces coelicolor*, the antimicrobial red pigment prodigiosin can induce programmed cell death in subpopulations, and thereby provide nutrients to the surviving kin, suggesting an additional ecological role besides being an antibiotic (Tenconi et al. 2018). Expanding our holistic understanding of secondary metabolites demands in-depth studies of their facets, from chemistry to ecology. Here, we present a case study of tropodithietic acid (TDA; Fig. 1), a sulfur-containing tropone derivative with several functions.

TDA can act as antimicrobial and is produced by marine members of the *Rhodobacteraceae* family (class of Alphaproteobacteria; Brinkhoff et al. 2004; Bruhn et al. 2005; Geng et al. 2008; Harrington et al. 2014; Sonnenschein et al. 2017a, 2018; Duan et al. 2020). In addition to the antimicrobial properties of TDA, the compound exhibits other activities as a signaling molecule, anti-cancer agent, and weak iron chelator (Geng and Belas 2010; Wichmann et al. 2015; D'Alvise et al. 2016; Wilson et al. 2016; Beyersmann et al. 2017). TDA-producing bacteria have been detected in oceanic metagenomic data sets (Segev et al. 2016; Sonnenschein et al. 2017a) and have also been isolated from marine aquaculture and oceanic environments (Hjelm et al. 2004a; Lauzon et al. 2008; Porsby, Nielsen and Gram 2008; Grotkjær et al. 2016b). They can inhibit, or kill, fish pathogenic bacteria when co-cultivated in aquaculture live feed (microalgae, *Artemia*, rotifers, and copepods) or fish larvae, sparking a commercial interest in TDA-producing bacteria as aquaculture probiotics (D'Alvise et al. 2012; Grotkjær et al. 2016a; Dittmann et al. 2017; Rasmussen et al. 2018; Sonnenschein et al. 2021).

The purpose of this review is to provide an overview of the different functions of TDA, thus also addressing its possible broader physiological and ecological roles, with perspectives to other secondary metabolites. We focus on TDA-producing bacteria belonging to the marine *Rhodobacteraceae* family, but it should be noted that TDA-producing bacteria outside of this group have been isolated (Kintaka et al. 1984; Tsubotani et al. 1984; Kawano et al. 1997). The term 'function' will cover direct molecular or chemical responses attributed to TDA, whilst 'roles' will comprise the more holistic possible ecological effects of the compound (Fig. 1).

Given the potential broader functions of antimicrobial secondary metabolites, their definition and terminology have been debated (Bérdy 2005; Price-Whelan, Dietrich and Newman 2006; Davies 2013; Chevrette et al. 2020). 'Secondary metabolites' was introduced by the Nobel Prize laureate Albrecht Kossel in 1891 (Kossel 1891), and adopted by the botanist Friedrich Czapek, who in the 1920s coined the term 'secondary modifications' in work on plant nitrogen metabolism (Czapek 1922), with the purpose of distinguishing the compounds from growth-related primary metabolites. More recently, the term 'specialized metabolites' has gained traction emphasizing functions broader than merely secondary (Price-Whelan, Dietrich and Newman 2006; Davies 2013). We will, however, use the term 'secondary metabolites' in this review as this remains the term most commonly used.

TDA-producing bacteria and their environmental niches

The tautomer of TDA, thiotropocin, was discovered in 1984 in a *Pseudomonas* species isolated from soil (Kintaka et al. 1984; Tsubotani et al. 1984), and TDA was later detected in a marine bacterium, *Roseobacter gallaeciensis* (now: *Phaeobacter inhibens*; Brinkhoff et al. 2004). Since then, TDA has only been detected in a subset of members belonging to the *Rhodobacteraceae* family. This includes strains belonging to the three genera *Phaeobacter* (formerly *Roseobacter*; Ruiz-Ponte et al. 1998; Brinkhoff et al. 2004; Martens et al. 2006; Porsby, Nielsen and Gram 2008; Geng and Belas 2010; Berger et al. 2011; Breider et al. 2014; Sonnenschein et al. 2017b), *Tritonibacter* (formerly *Epibacterium*, *Ruegeria*, or *Silicibacter*; Hjelm et al. 2004a; Bruhn, Gram and Belas 2007; Muramatsu et al. 2007; Geng et al. 2008), and *Pseudovibrio* (Enticknap et al. 2006; Geng and Belas 2010; Penesyan et al. 2011; Bondarev et al. 2013; Harrington et al. 2014).

TDA is produced by strains belonging to four of the six described *Phaeobacter* species (Sonnenschein et al. 2018), namely *Phaeobacter gallaeciensis* (Martens et al. 2006), *P. inhibens* (Martens et al. 2006), *Phaeobacter piscinae* (Sonnenschein et al. 2017b), and *Phaeobacter porticola* (Breider et al. 2017). Production of TDA has so far not been detected in the species *Phaeobacter italicus* (Wirth and Whitman 2018) and the proposed species *Phaeobacter marinintestinus* (Lee et al. 2015), nor has the biosynthetic gene cluster of TDA been detected through genome mining (NCBI accession numbers NZ_VOGO01000001.1 (*P. marinintestinus* UB-M7) and FOOZ00000000.1 (*P. italicus* DSM26436)) using antiSMASH 5.0 in these species (Blin et al. 2019). Several strains of *Tritonibacter mobilis* (formerly *Epibacterium mobilis*) produce TDA, while strains belonging to other species in the genera, e.g. *Tritonibacter scottomollicae* (formerly *Epibacterium scottomollicae*), do not (Geng et al. 2008; Wang and Seyedsayamdost 2017; Sonnenschein et al. 2017a). In the *Pseudovibrio* genus, TDA-producing strains (unclassified at species level) have repeatedly been isolated (Enticknap et al. 2006; Geng and Belas 2010; Penesyan et al. 2011; Bondarev et al. 2013; Harrington et al. 2014) with *Pseudovibrio ascidiacei* being the closest relative to the TDA-producing *Pseudovibrio* isolates (Penesyan et al. 2011). However, a number of *Pseudovibrio* strains do not harbor the *tda* genes, suggesting that TDA production is not a widely distributed trait within this genus (Crowley et al. 2014; Romano 2018).

Since TDA production is not a conserved trait within the three genera, and since *Phaeobacter*, *Tritonibacter*, and *Pseudovibrio* are not close phylogenetic neighbors within the *Rhodobacteraceae* family, this could suggest that TDA genes and the ability to produce the compound have been distributed by horizontal gene transfer (Sonnenschein et al. 2018). However, short-term, noncompetitive biofilm cultivation of *P. inhibens* 2.10 induced single nucleotide polymorphisms in genes responsible for TDA production, leading to TDA deficiency (Majzoub et al. 2021). This points toward strong selection for the loss of TDA production in *P. inhibens*, presenting another possible explanation for the nonconserved pattern of TDA genes observed within the genera. To this day, the evolutionary history of TDA production is not fully understood, but would be important for unravelling the ecological role(s) of TDA. For most secondary metabolites the evolutionary route responsible for the chemical diversification remains poorly understood. Horizontal gene transfer has been identified to be an integral driver of secondary metabolite evolution (Fischbach, Walsh and Clardy 2008; Medema et al. 2014), but it is evident that vertical inheritance also influences the evolution of secondary metabolite




Figure 1. Proposal for three functions of TDA and potential ecological roles. Bold text indicate functions. Nonbold text indicate potential ecological roles. Created with Biorender.com.

gene clusters (Lind et al. 2017; Adamek et al. 2018; Chase et al. 2021; Undabarrena et al. 2021). Genetic diversification of the salinosporamides A biosynthetic gene cluster, found in the marine *Salinispora* genus, had direct consequences on the secondary metabolite production (Chase et al. 2021). This highlights that long-term evolutionary processes can lead to genetic and chemical diversification of secondary metabolites within closely related species, potentially generating new chemical diversity.

The genera *Phaeobacter* and *Tritonibacter* belong to the *Roseobacter* group ('roseobacters'), which is a paraphyletic subgroup within the family *Rhodobacteraceae* (Simon et al. 2017). The *Roseobacter* group contains multiple branching clades (Newton et al. 2010; Simon et al. 2017), where clade 1 contains strains that produce TDA (Brinkhoff et al. 2004; Newton et al. 2010). In the environment, the abundance of roseobacters is highest near the surface of temperate coastal waters and polar oceans (Buchan, González and Moran

2005), globally averaging 3.8% of bacterial populations (Wietz et al. 2010). The abundance of roseobacters correlates positively with chlorophyll *a* concentrations (Wietz et al. 2010), and roseobacters can constitute as much as 30% of the bacterial population in microalgal blooms (González and Moran 1997; Gonzalez et al. 2000; West et al. 2008).

In global marine metagenomic data, the *Phaeobacter* genus represents approximately 0.03% of all bacterial sequences in surface waters (Sunagawa et al. 2015; Sonnenschein et al. 2021), and while *Phaeobacter* species have not been isolated from open ocean water (Gram, Melchiorson and Bruhn 2010), they have frequently been recovered from aquaculture facilities (Hjelm et al. 2004a; Porsby, Nielsen and Gram 2008; Grotkjær et al. 2016b) and solid surfaces in harbors (Ruiz-Ponte et al. 1998; Bernbom et al. 2011; Gram et al. 2015; Breider et al. 2017). This is in concordance with comparative genomic analyses of the *Phaeobacter* genus suggesting an

adaptation to a surface-associated lifestyle (Dang and Lovell 2000; Hjelm et al. 2004a; Rao, Webb and Kjelleberg 2006; Porsby, Nielsen and Gram 2008; Thole et al. 2012; Gram et al. 2015; Freese et al. 2017). Antagonistic interactions are believed to be more frequent in particle- and between surface-associated bacteria due to the high cell densities in structured microenvironments (Long and Azam 2001; Gram, Melchiorson and Bruhn 2010). This could indicate that production of TDA provides a competitive advantage for the producer in surface colonization. TDA-producing *Tritonibacter* strains have also been isolated from aquaculture environments (Buchan, González and Moran 2005; Porsby, Nielsen and Gram 2008; Alsmark et al. 2013), as well from open ocean waters (Gram, Melchiorson and Bruhn 2010; Sonnenschein et al. 2017a). The *Tritonibacter* genus represents 0.2% of the bacterial population in the surface ocean, and is thus more abundant in oceanic surface waters than *Phaeobacter* (Sunagawa et al. 2015). However, like *Phaeobacter*, *Tritonibacter* is adapted to a surface-attached lifestyle and occurs rather in the particle-associated than the free-living fraction of seawater (Sonnenschein et al. 2017a). Consequently, it has been proposed that *T. mobilis* could be the open-water equivalent to the coastal, macrosurface-attached *Phaeobacter*.

TDA-producing bacteria are also found in association with a range of marine eukaryotes, such as zooplankton (Freese et al.), sponges (Harrington et al. 2014), molluscs (Ruiz-Ponte et al. 1999; Prado et al. 2009), and algae (Rao, Webb and Kjelleberg 2006; Nappi, Soldi and Egan 2019) and it has been suggested that up to one-third of TDA-producing bacteria are host-associated (Nappi, Soldi and Egan 2019). This is specifically seen with TDA-producing *Pseudovibrio* strains that are present on, and genetically adapted to, a symbiotic lifestyle with marine invertebrates such as corals and macroalgae (Enticknap et al. 2006; Penesyan et al. 2011; Bondarev et al. 2013; Crowley et al. 2014; Raina et al. 2016; Romano 2018). *Pseudovibrio* represent 0.04% of all bacteria in oceanic surface waters (Sunagawa et al. 2015) and TDA-producing *Pseudovibrio* species harbor genes associated with a free-living lifestyle (Enticknap et al. 2006; Bondarev et al. 2013).

Like other members of the *Rhodobacteraceae* family, TDA-producing bacteria are characterized by a high versatility of metabolic pathways (Newton et al. 2010; Bondarev et al. 2013; Zech et al. 2013; Sonnenschein et al. 2017a). TDA-producing *Phaeobacter* and *Tritonibacter* can catabolize several algal osmolytes, such as dimethylsulfoniopropionate (DMSP; Miller and Belas 2004; Newton et al. 2010; Thole et al. 2012), which has been suggested to act as a chemo-attractant for TDA-producing bacteria (Miller and Belas 2004; Miller et al. 2004) capable of utilizing DMSP in their primary metabolism (Curson et al. 2011). Furthermore, DMSP may provide sulfur to be incorporated into TDA (Geng et al. 2008), and TDA-producing bacteria, thus benefit from living in association with microalgae and corals (Raina et al. 2009, 2016; Harrington et al. 2014; Segev et al. 2016). However, other studies have not been able to detect the incorporation of DMSP sulfur in TDA in either *P. inhibens* DSM17395 or in *P. gallaeciensis* DSM26640, and instead suggested that cysteine serves as the sulfur precursor (Dickschat et al. 2017).

Biosynthesis, tautomers, and analogues of TDA

TDA is a sulfur-containing tropone derivative (Fig. 2). It can tautomerize into two other known sulfur-containing tropones; thiotropocin and troposulfenin (Fig. 2). The structures of these compounds are characterized by an aromatic cyclohepta-2,4,6-trienone moiety. Their biosynthesis draws on the central carbon and sulfur metabolism, as well as a range of enzymes encoded by

a cluster of dedicated so-called *tda* genes (Fig. 3A; Duan et al. 2020). For an in-depth review of the chemistry underlying the biosynthesis of TDA as well as other tropones, we refer to the review by Duan et al. (2020).

The basic skeleton of TDA and other tropone natural products arise from phenylacetic acid (**1**, PAA) catabolism (Fig. 3A), where PaaK, PaaABC(D)E (*E. coli* nomenclature, also referred to as PaaGHI(J)K in *Pseudomonas putida*), PaaG, as well as PaaZ are required to form a highly reactive intermediate, 3-oxo-5,6-dehydrosuberil-CoA semialdehyde (**3**, Fig. 3B, Teufel et al. 2010, 2011; Berger et al. 2012; Brock, Nikolay and Dickschat 2014). Interestingly, in TDA-producers, a homologue of PaaZ, PaaZ2, is encoded on a megaplasmid along with the core enzymes of TDA biosynthesis, TdaABCDEF (Fig. 3B; Brock, Nikolay and Dickschat 2014). PaaZ contains two domains; a C-terminal enoyl-CoA hydratase (ECH) and a N-terminal aldehyde dehydrogenase (ALDH) domain (Brock, Nikolay and Dickschat 2014). In PaaZ2, only the C-terminal ECH is conserved (Brock, Nikolay and Dickschat 2014). In the absence of the ADHL domain, the intermediate **3** cyclizes to a seven-membered ring 2-hydroxycyclohepta-1,4,6-triene-1-formyl-CoA (**4**), the proposed universal tropone precursor, via a spontaneous intramolecular condensation (Teufel et al. 2011). A recent study pointed to the acyl-CoA dehydrogenase-like flavoenzyme TdaE being the linker between primary metabolism and TDA biosynthesis (Duan et al. 2021). TdaE converts the PAA catabolism shunt product **4** and carries out a series of reactions, including dehydrogenation, CoA-ester oxygenolysis, and ring epoxidation to form **7** (Duan et al. 2021), which then presumably to be converted by TdaF, TdaB, and PatB to form TDA (**5**; Duan et al. 2021). This biosynthetic pathway therefore serves as an example where the bacteria 'direct' primary metabolism toward secondary metabolism.

Different analogues of TDA have been detected from TDA-producing bacteria (Choudhary et al. 2018; Phippen et al. 2019). The first analogue was tentatively characterized as TDA-methyl ester, was isolated from a TDA-producing *Pseudovibrio* sp. (Choudhary et al. 2018; Fig. 2). Methyl-troposulfenin, an S-methylated congener of TDA, was identified as a natural analogue of TDA from a TDA-producing *P. inhibens* (Phippen et al. 2019; Fig. 2). Notably, the antimicrobial activity of methyl-troposulfenin is lower than that of TDA (Phippen et al. 2019). A comparison of the MS fragmentation patterns of methyl-troposulfenin (Phippen et al. 2019) and the proposed TDA-methyl ester (Choudhary et al. 2018) show a very high level of similarity (Choudhary et al. 2018; Phippen et al. 2019), and due to the lack of NMR experiments in the characterization of TDA-methyl ester, it cannot be ruled out that TDA-methyl ester is a misassignment of methyl-troposulfenin. For other secondary metabolites, the production of analogues may serve to broaden the chemical (antimicrobial) repertoire of the producing organism or may serve as a detoxification and self-protection of the producer (Li et al. 2013; Gallagher et al. 2017).

Total synthesis of TDA has not been reported thus far. However, owing to its bioactivity, a series of TDA inspired analogues have been synthesized to elucidate structure activity relationship (Rabe et al. 2014). One particular synthetic analogue, tropone-2-carboxylic acid (Fig. 2B), a nonsulfur variant of TDA, had a stronger antibacterial activity against *Staphylococcus aureus* and *Vibrio anguillarum* than TDA itself, suggesting that the sulfur atoms are not necessary for the antimicrobial effect of TDA (Rabe et al. 2014).




Figure 2. TDA and its tautomers (upper panel) and analogues (lower panel). Created with ChemDraw Professional (PerkinElmer Informatics).

Factors influencing TDA production

As previously mentioned, TDA biosynthesis draws on central carbon metabolism as well as sulfur metabolism (Geng et al. 2008). The cascading production of TDA across different metabolic processes may explain why TDA production is influenced by a multitude of factors including mode of growth, nutrients, and cell density.

Mode of growth

When *P. piscinae* 24–7 is cultured in Marine Broth (MB), stagnant as compared to shaken conditions increase brown pigmentation (Bruhn et al. 2005), which is a TDA–iron complex, and thus a proxy for TDA production (D’Alvise et al. 2016). Similarly, TDA production by *Tritonibacter* is facilitated by stagnant growth conditions, whereas several *Phaeobacter* strains also produce TDA in aerated cultures (Bruhn et al. 2005, 2007; Porsby, Nielsen and Gram 2008; Belas et al. 2009; Geng and Belas 2010; Berger et al. 2011; D’Alvise et al. 2014).

A distinct phenotype associated with stagnant growth of TDA-producing bacteria is the formation of a thick biofilm at the liquid–air interface (Bruhn, Gram and Belas 2007; Gram, Melchiorson and Bruhn 2010) and the appearance of the brown TDA–iron complex in this biofilm. This observation has prompted the hypothesis that TDA production and biofilm formation could be linked (Bruhn et al. 2005). In *Tritonibacter* sp. TM1040, deficiency of the swimming regulator *flaC* is associated with a shift toward the motile phase, a reduction in biofilm formation as well as decreased antibiotic activity (Belas et al. 2009). The secondary messenger cyclic dimeric guanosin-monophosphate (c-di-GMP), is likely involved in the interconnection of these phenotypic traits as increased production of c-di-GMP induces both biofilm formation and TDA production in *T. mobilis* F1926 (D’Alvise et al. 2014). However, attachment, biofilm formation, and TDA biosynthesis are not universally linked across TDA-producing genera (Prol García et al. 2014; Zhao et al. 2016; Majzoub et al. 2018). While stagnant growth conditions facilitate biofilm formation and TDA production in many *Tritonibacter* strains, this is not the case in most *Phaeobacter* species. Collectively, these observations support the notion that *Phaeobacter* and *Tritonibacter* occupy separate niches (Sonnenschein et al.

2017a), where *Tritonibacter* species predominantly reside in open waters, and requires a tighter regulation of TDA production as not to waste metabolic energy during dispersed planktonic growth.

Nutrients

Variation of carbon, nitrogen, sulfur, phosphorus, or iron sources affect production of TDA. In *P. inhibens* DSM17395, production of TDA increases dramatically when phenylalanine is used as the primary carbon source instead of glucose, and more TDA is produced in general, when aromatic compounds are utilized as a carbon source (Berger et al. 2012). As the backbone of TDA originates from the phenylacetic acid catabolon, this increase in TDA production is likely a result of increased precursor availability (Berger et al. 2012; Brock, Nikolay and Dickschat 2014).

TDA is a sulfur-containing compound and in *Tritonibacter* sp. TM1040, growth on DMSP is associated with an increase in TDA concentrations as compared to growth on sulfate-containing substrates (Geng and Belas 2010). DMSP is produced by phytoplankton and roseobacters preferentially metabolize DMSP over the more readily available sulfate (Kiene et al. 1999), likely pointing to a niche-specific adaptation. In *Pseudovibrio* sp. FO-BEG1, phosphate limitation induces TDA production but this is likely attributed to a global change in sulfur metabolism rather than a specific phosphate effect (Romano et al. 2015).

In laboratory cultures, production of bioactive TDA is dependent on iron in concentrations that far exceed those observed in natural marine systems (D’Alvise et al. 2016). Despite this, *Phaeobacter* exhibits TDA-dependent antagonism against vibrios in low-iron artificial seawater as well as in systems mimicking multitrophic level seawater systems (D’Alvise et al. 2010, 2012). A nonantibacterial (inactive) form of TDA (‘pre-TDA’) is produced when iron is not available, and this ‘pre-TDA’ can be converted to TDA by acidification (D’Alvise et al. 2016). Thus, the biosynthesis of TDA does not appear to be regulated by iron at the transcriptional level, yet bioactive TDA only forms in its presence. This is in contrast to other iron-chelating secondary metabolites, siderophores, which are typically upregulated in the absence of iron. The weak iron-chelating properties of TDA indicate that iron sequestering is not its main function, but potentially relate to its mode of action (Wilson et al. 2016), or to symbiosis where the TDA–iron complex




Figure 3. TDA biosynthesis and genes responsible for TDA production. **(A)** Biosynthetic genes involved in TDA biosynthesis in *P. inhibens* DSM17395. *paaABCDEIJK1K2* and *patB* (in green) are located on the chromosome, whilst *tdaABCDEF* and *paaZ2* (in blue) are located on a 262-kb megaplasmid. **(B)** TDA biosynthesis draws on primary metabolism for formation of the carbon backbone. Phenylacetic acid, **1**, is converted to **2** by PaaK, PaaABC(D)E, and PaaG. PaaZ-ECH or PaaZ2 catalyzes hydrolytic ring cleavage to form **3**, which is then either converted to **9** by the ALDH domain of PaaZ or spontaneously cyclized to **4**. TdaE then connects primary and secondary metabolism through a series of reactions: dehydrogenation to **5**, CoA-ester oxygenolysis to **6**, and ring epoxidation to form **7**. TdaBCDF subsequently forms TDA, **8**. Stippled lines indicate multiple reactions taking place. Created with ChemDraw Professional (PerkinElmer Informatics) and Biorender.com.

could serve as an iron reservoir (D'Alvise et al. 2016), similarly to what has been proposed for vibrioferrin produced by *Marinobacter* (Amin et al. 2009; Yarimizu et al. 2019).

Quorum sensing and autoinduction

TDA production is measurable in late exponential or early stationary growth phase (Geng and Belas 2010; Berger et al. 2011; Harrington et al. 2014; Romano et al. 2015), which is correlated with high cell densities. Since many roseobacters also produce the QS molecules acyl-homoserine lactones (AHLs), it has been suggested that TDA production is QS regulated (Gram et al. 2002; Bruhn et al. 2005, 2006, 2007; Martens et al. 2007; Berger et al. 2011; Zan et al. 2014). Transposon insertions in either the AHL synthase gene, *pgaI*, or the gene encoding the response regulator, *pgaR*, of *P. inhibens* DSM17395 results in a reduction in TDA production (Berger et al. 2011). However, the effect is temporary, indicating that TDA biosynthesis is not fully dependent on this QS system (Prol Garcia, D'Alvise and Gram 2013). Furthermore, the effect is reversible

through supplementation with phenylalanine, suggesting a hierarchical regulation of TDA biosynthesis (Berger et al. 2012). AHLs are not universally involved in the regulation of TDA production since *Tritonibacter* sp. TM1040 does not produce any known AHLs and no AHL synthase genes have been identified in their genomes (Sonnenschein et al. 2017a). Other compounds than AHLs may be involved in QS, and using a *tdaCp::lacZ* reporter plasmid in several TDA-deficient *Tritonibacter* sp. TM1040 strains, it was discovered that *tdaC* was not expressed in the absence of TDA (Geng and Belas 2010). Subsequently, *tdaC* expression was restored by cross-feeding with the wildtype, demonstrating that TDA acts as an autoinducer of its own biosynthesis in a manner similar to AHL signaling TDA (Geng and Belas 2010). Addition of both the AHL produced by *pgaI*, 3-OH-C10-homoserine lactone (HSL), and TDA restored TDA production in the *pgaI*⁻ mutant, but not in the *pgaR*⁻ mutant, indicating that TDA could potentially act as an autoinducer through the same response regulator as 3-OH-C10-HSL (Berger et al. 2011). This provides a possible link between autoin-

duction and AHL-mediated regulation of TDA biosynthesis (Berger et al. 2011).

QS regulation of TDA production has only been demonstrated in the strains TM1040 and DSM17395. Adding TDA to a final concentration of 1 μ M to the TDA-producing *Pseudovibrio* sp. W74 did not result in earlier onset or higher production of TDA, indicating that autoinduction did not take place (Harrington et al. 2014; D'Alvise et al. 2016). However, since W74 produces TDA, TDA would also be present in the control cultures and testing TDA addition to a TDA-deficient mutant of W74 would be required to determine any possible QS function.

The exact regulatory network controlling TDA biosynthesis has not been elucidated, however, using a *tdaCp::lacZ* reporter fusion, it has been demonstrated that TdaA is necessary for the transcriptional activation of *tdaC* expression in *Tritonibacter* sp. TM1040, independent of TDA (Geng and Belas 2011). TdaA also acts as a transcriptional activator in *P. inhibens* DSM17395, where expression of *tdaB*, *tdaE*, and *tdaF* is strongly downregulated in *tdaA*⁻ mutants (Berger et al. 2011). A putative binding site of TdaA near the *tdaC* promoter has been identified in *Tritonibacter* sp. TM1040, and binding of TdaA to the *tdaC* promoter has been confirmed using an electrophoretic mobility shift assay (Geng and Belas 2011). Understanding the regulatory switches governing TDA biosynthesis, and how these differ between genera, may provide an important clue as to the ecological role of this compound.

Effect of TDA on the producing bacteria

Deletion of the 262-kb megaplasmid encoding the last part of the TDA biosynthetic pathway leads to an increase in growth rate and yield, demonstrating a significant burden of this plasmid (Trautwein et al. 2016; Wünsch et al. 2020). Similarly, transposon insertions in any of the *tdaA*, *tdaB*, *tdaC*, or *tdaE* genes on the plasmid result in TDA-deficient mutants exhibiting increased growth rates and yields (Will et al. 2017). The negative effects of TDA on the producer is, however, not due to the metabolic cost associated with TDA biosynthesis as supplementation with TDA-containing supernatant reverts the growth of the TDA-deficient mutant to wildtype levels (Will et al. 2017) indicating an autoantibacterial activity. Thus, as TDA production impairs growth of the producer, the compound must represent a significant ecological advantage in natural marine systems.

The fact that TDA interacts with AHL response regulators in *Tritonibacter* sp. TM1040 has major implications for global gene expression in the producing organism (Berger et al. 2011; Beyersmann et al. 2017). With the exception of 15 genes, TDA regulates the same genetic circuitry as the AHL 3-OH-C10-HSL in *P. inhibens* DSM17395, including genes involved in chemotaxis, motility, attachment, and biofilm integrity (Beyersmann et al. 2017). In effect, the QS signaling molecule and TDA in conjunction likely facilitate a swim-and-stick lifestyle through the induction of biofilm dispersion. While similar mechanisms have been observed for *P. inhibens* 2.10 (Majzoub et al. 2018), the evident connection between TDA and the biofilm mode of growth is likely different among producers with different niche adaptations, and more studies are necessary to fully resolve the effect of TDA on the producing organism. Several other antimicrobial secondary metabolites are known to function as QS signals in the producing strain, e.g. surfactin in *Bacillus subtilis* and *Bacillus amyloliquefaciens* (López et al. 2009; Chen et al. 2020), and pyocyanine in *P. aeruginosa* (Dietrich et al. 2006). Similarly to TDA, these compounds also affect biofilm formation and motility (Das and Manefield 2012). These molecules vary widely in chemical structure and are produced by bacteria

that are taxonomically distant; hence, it seems plausible that this may be a somewhat common function of antimicrobial secondary metabolites which is likely facilitated by many different mechanisms.

TDA-mediated inhibition of other microorganisms

TDA can inhibit or kill a wide range of Gram-positive and Gram-negative bacteria, including both fish and human pathogens (Ruiz-Ponte et al. 1998, 1999; Hjelm et al. 2004a; Planas et al. 2006; Porsby, Nielsen and Gram 2008; Prado et al. 2009; Porsby et al. 2011; Porsby and Gram 2016; Zhao et al. 2016; Grotkjær et al. 2016b; Sonnenschein et al. 2021). Also eukaryotes such as the fungal pathogens *Rizoctonia solani*, *Candida albicans*, and some microalgal strains of the genera *Chlorella* and *Scenedesmus* are negatively affected by TDA (Kintaka et al. 1984; Liang 2003). Pure TDA can be lethal to some mammalian cell lines, including neuronal and cancer cells, and it has been suggested that TDA may lead to disruption of the mitochondrial membrane potential and activation of oxidative stress responses (Wichmann et al. 2015). The algal compound DMSP is to some extent protective against TDA-induced cytotoxicity, as preincubation with DMSP of mammalian neural cells exerted protective effects against TDA, potentially due to DMSP acting as an antioxidant (Wichmann et al. 2016). This could suggest that DMSP could have a role in the interplay between marine eukaryotes and TDA-producing bacteria, thus acting as a protectant against TDA, although this is highly speculative and requires further studies (Duan et al. 2020). Despite the negative effects of TDA on eukaryotic cells, no adverse effect of TDA producers has so far been observed on higher organisms (Sonnenschein et al. 2021) such as microalgae, *Artemia*, rotifers, or nauplii (D'Alvise et al. 2012; Prol García, D'Alvise and Gram 2013; Rasmussen et al. 2018, 2019; Sonnenschein et al. 2018). In fact, TDA producers may be important symbionts for microalgae and corals, and TDA has been proposed to act as algal and coral protectant (Seyedsayamdost et al. 2011b; Raina et al. 2016). Some TDA-producers, i.e. *P. inhibens*, *P. gallegiensis*, and *P. piscinae*, also produce the algicidal troponoids roseobacticides (Seyedsayamdost et al. 2011a; Sonnenschein et al. 2018). These secondary metabolites are synthesized in response to *p*-coumaric acid, a potential senescence signal produced by algae (Seyedsayamdost et al. 2011a). In *P. inhibens* DSM17395, TDA and roseobacticides share parts of the same biosynthetic pathway, and the metabolites probably share the same precursor (Wang, Gallant and Seyedsayamdost 2016). This, thus, challenges the one-cluster-one-compound paradigm. These findings highlight that secondary metabolites may also be involved in beneficial interkingdom cross-talk. A well-studied example of this is found in *Bacillus*-plant interactions, where production of the secondary metabolite surfactin is stimulated in response to plant host cues and in turn triggers plant immunity against pathogens in *Bacillus*-plant interactions (Hoff et al. 2021).

In the proposed mechanism of antibacterial action, TDA disrupts the proton motive force by binding extracellular protons to the carboxyl group and transporting them across the cell membrane to the cytosol (Wilson et al. 2016). Here, the proton is exchanged for a metal ion, i.e. transported back to the extracellular space (Wilson et al. 2016). This destroys the transmembrane proton gradient whilst maintaining the membrane potential, making TDA an electroneutral proton antiporter (Wilson et al. 2016). In concordance with this, exposure of a *Vibrio vulnificus* to sublethal concentrations of pure TDA lead to upregulation of genes involved

in iron-uptake, oxidative stress, and regeneration of the cell envelope (Dittmann *et al.* 2019a).

The antibacterial effect of TDA against fish pathogenic bacteria has been extensively studied, due to the interest in using TDA-producing bacteria as probiotics in marine aquaculture (Ruiz-Ponte *et al.* 1998, 1999; Bruhn, Gram and Belas 2007; D'Alvise *et al.* 2012; Porsby and Gram 2016; Zhao *et al.* 2016; Grotkjær *et al.* 2016b; Rasmussen *et al.* 2018, 2019; Dittmann *et al.* 2019a; Ringø 2020). TDA-producing *Phaeobacter* and *Tritonibacter* are antibacterial against *Vibrio* species, such as *V. anguillarum*, *V. vulnificus*, and *Vibrio coralliilyticus* (D'Alvise *et al.* 2010; Porsby and Gram 2016; Zhao *et al.* 2016). Extracts from TDA-producing *Pseudovibrio* sp. P12 strongly inhibited the growth of *V. coralliilyticus* and *Vibrio owenii*, two coral pathogens causing white syndrome in Scleractinian corals (Raina *et al.* 2016). Growth of *V. coralliilyticus* is not suppressed by common coral-associated bacteria and the bacterium exhibits antibiotic resistance to a wide range of commercial antibiotics, greater than that of other vibrios such as *V. vulnificus* (Shnit-Orland and Kushmaro 2009; Rypien, Ward and Azam 2010; Vizcaino *et al.* 2010), emphasizing the antibiotic potential of TDA. TDA also inhibits other fish pathogens such as *Aeromonas* and *Tenacibaculum* spp. (Porsby and Gram 2016; Grotkjær *et al.* 2016b; Tesdorpf *et al.* 2022).

The antagonistic effect of TDA-producing strains has primarily been determined using agar-based assays (Brinkhoff *et al.* 2004; Hjelm *et al.* 2004a; Bruhn *et al.* 2005; Rao *et al.* 2007; Porsby, Nielsen and Gram 2008; Prado *et al.* 2009) and different broth and/or biofilm-based co-culture setups (Hjelm *et al.* 2004b; Porsby, Nielsen and Gram 2008; Prado *et al.* 2009; Grotkjær *et al.* 2016b). Exposure of a *V. anguillarum* to surface-attached *Phaeobacter* or *Tritonibacter* resulted in significant reduction or complete elimination of *V. anguillarum* (D'Alvise *et al.* 2010). Furthermore, *P. inhibens* DSM17395 successfully inhibited *V. vulnificus* in co-culture experiments, keeping it at inoculum level, whereas monocultures of *V. vulnificus* were 1000-fold higher (Porsby and Gram 2016). TDA is most likely responsible for this inhibition, since TDA-negative mutants did not inhibit *V. anguillarum* (D'Alvise *et al.* 2010).

The activity of TDA-producing roseobacters against vibrios has also been assessed in more complex, nonaxenic, and microcosm experiments including marine organisms of multiple trophic levels. As pathogenic *Vibrio* strains may enter the aquaculture unit via live feed, many experiments have been conducted in cultures of microalgae, rotifers, brine shrimps, and copepods (D'Alvise *et al.* 2012; Prol García, D'Alvise and Gram 2013; Porsby and Gram 2016; Grotkjær *et al.* 2016a; Rasmussen *et al.* 2018). In these laboratory experiments, TDA-producing *Phaeobacter* can reduce the number of vibrios and other fast-growing heterotrophic bacteria (D'Alvise *et al.* 2012; Grotkjær *et al.* 2016a; Rasmussen *et al.* 2018, 2019).

Resistance and tolerance to TDA

Phaeobacter inhibens DSM17395 carries three genes, *tdaR123*, conferring its self-resistance to TDA (Wilson *et al.* 2016). *tdaR1* and *tdaR2* are predicted to encode transmembrane proteins, while *tdaR3* is predicted to encode a γ -glutamyl-cyclotransferase, which in *Escherichia coli* is involved in cation-proton exchange (Wilson *et al.* 2016). All three genes are co-located to the TDA gene cluster on the megaplasmid, which to this day has not proven transmissible or to encode transmission genes (Petersen *et al.* 2013). Sensitivity to TDA in *E. coli* is reduced when the *tdaR123* genes are transferred and heterologously expressed (Wilson *et al.* 2016). However, the *tdaR123* genes have not been found in bacteria not producing TDA, and it has not been possible to develop resistance to

TDA in target bacteria *in vitro* (Porsby *et al.* 2011; Rasmussen *et al.* 2016). Different *in vitro* approaches have been used to induce mutations or adaptations conferring TDA resistance in the non-TDA-producer species *P. aeruginosa*, *S. aureus*, *Salmonella typhimurium*, and *E. coli*, which are all species susceptible to TDA (Porsby *et al.* 2011). Using adaptive laboratory evolution experiments, *V. anguillarum* strains capable of tolerating two times the minimum inhibitory concentration were evolved, however, the tolerance was transient and vanished after one passage in medium free of TDA (Rasmussen *et al.* 2016). The difficulty in developing TDA resistance or tolerance could suggest that TDA has multiple additional targets beside the disruption of the proton motive force. Since only a few TDA resistant bacteria have been found, and resistance is difficult to develop, the use of TDA as an antibiotic and of TDA producers as probiotics will not add to the risk of antimicrobial resistance (Sonnenschein *et al.* 2021).

TDA tolerance has been observed in natural microbial communities containing indigenous TDA-producing *Pseudovibrio*. Here, 126 out of 136 isolated non-TDA producing bacteria were tolerant to TDA (Harrington *et al.* 2014). Among the TDA-tolerant isolates were *Psychrobacter*, *Alteromonas*, *Salinibacter*, *Alcanivorax*, *Flavobacterium*, and *Micrococcus* strains, whilst TDA-sensitive isolates included *Staphylococcus*, *Idiomarina*, and *Rhodococcus* strains (Harrington *et al.* 2014). The mechanisms enabling this tolerance are not known.

Influence of TDA or TDA producers on marine microbial communities

The potential use of TDA-producing bacteria as probiotics in marine aquaculture has prompted studies determining how TDA or TDA-producing bacteria affect natural marine microbiomes. The possible effect of TDA or bacteria producing TDA on taxonomic composition of natural microbiomes has been determined, typically by 16S rRNA gene amplicon sequencing (Table 1).

Adding pure TDA to the microalgae *Nannochloropsis salina* colonized by a seawater microbiome caused a decrease in relative abundance of bacteria belonging to *Rhodobacteraceae*, *Flavobacteriia*, and *Alteromonadaceae* after 24 h, while bacteria of unclassified families within the Alteromonadales order increased in relative abundance (Geng *et al.* 2016). The changes were more rapidly seen in communities exposed to high concentrations (500 nM as opposed to 21 nM) of TDA, indicating a dose-dependent effect of TDA. The addition of TDA accelerated the development of the microbial community that after 3 h had a composition similar to the one reached in the nontreated community after 24 h (Geng *et al.* 2016). It should be noted that the concentration of TDA found in natural systems is not known. Addition of pure TDA to the microalgae *Tetraselmis suecica* could be detected to a lower limit of 50 nM, but TDA was not detectable when TDA-producing bacteria were cultured in the system (D'Alvise *et al.* 2012). Thus, it remains uncertain if the concentrations used in the *N. salina* study (Geng *et al.* 2016) were representative of the TDA concentrations in natural communities. Subsequent studies have typically studied changes in the bacterial community in the presence of TDA-producing bacteria.

The TDA-producing *P. inhibens* DSM17395 was added to the microbiome of the marine microalgae *Emiliana huxleyi* in concentrations reflecting the *in situ* abundances of roseobacters during algal blooms (Amin *et al.* 2015; Segev *et al.* 2016; Sonnenschein *et al.* 2018; Dittmann *et al.* 2019b). The addition of 10⁶ CFU/ml of DSM17395 caused a decrease in the relative abundance of bacteria belonging to Rhodobacterales, with *Loktanella* and *Marivita*

Table 1. Model systems used to study the effect of TDA-producing bacteria and TDA on microbial communities.

In vivo model	TDA dose	Controls	Duration (days)	Bacteria that increase	Bacteria that decrease	Reference
<i>Nannochloropsis salina</i> (microalgae)	Pure TDA (31–500 nM)	Untreated (glucose)	0–1	Alteromonadales (Unclassified families)	Alteromonadaceae, Flavobacteriia, Rhodobacteraceae	Geng et al. (2016)
<i>Emiliania huxleyi</i> (microalgae)	<i>P. inhibens</i> DSM17395 (10 ⁴ and 10 ⁶ cells/ml)	Untreated (medium)	0–4	<i>Colwellia</i> sp., <i>Winogradskyella</i> sp., <i>Neptuniibacter</i> sp. (absent in controls)	<i>Vibrio</i> sp., <i>Pseudoalteromonas</i> sp., Alteromonadales	Dittmann et al. (2019b)
<i>Ostrea edulis</i> (oyster)	<i>P. inhibens</i> DSM17395 (10 ⁴ and 10 ⁶ cells/ml)	Untreated (medium)	0–4	<i>Mycoplasma</i> sp.	<i>Mariivita</i> , <i>Vibrionaceae</i> , <i>Mycoplasma</i> sp., <i>Pseudoalteromonas</i> sp., <i>Shewanella</i> sp.	Dittmann et al. (2019b)
<i>Tetraselmis suecica</i> (microalgae)	<i>P. inhibens</i> DSM17395 (10 ⁶ cells/ml)	Untreated (medium)	0–4			Dittmann et al. (2020)
<i>Acartia tonsa</i> (copepod)	<i>P. inhibens</i> DSM17395 (10 ⁶ cells/ml)	Untreated (medium)	0–4		Rhodobacteraceae: <i>Ruegeria</i> , <i>Celeribacter</i> , <i>Pseudophaeobacter</i>	Dittmann et al. (2020)
<i>Scophthalmus maximus</i> (turbot larvae)	<i>P. inhibens</i> DSM17395 (10 ⁶ cells/ml)	Untreated (medium)	0–4		Rhodobacteraceae: <i>Ruegeria</i> , <i>Celeribacter</i> , <i>Pseudophaeobacter</i>	Dittmann et al. (2020)
<i>Thalassiosira rotula</i> (microalgae)	<i>P. inhibens</i> 2.10	TDA-deficient mutant	0–8		Rhodobacteraceae: <i>Sulfitobacter</i> , <i>Phaeobacter</i> , <i>Pelagicola</i> , <i>Loktanella</i>	Majzoub et al. (2019)

being the most affected genera (Dittmann *et al.* 2019b). This is similar to the changes observed in the *N. salina* study, where pure TDA caused a decrease in relative abundance of *Rhodobacteraceae* (Geng *et al.* 2016). When adding DSM17395 to the European flat oyster, *Ostrea edulis*, the relative abundance of the Vibrionales decreased markedly as compared to untreated microbiomes. In the *E. huxleyi* microbiome, individual amplicon sequence variants (ASVs) assigned as *Vibrio* sp. also decreased when treated with DSM17395, although no changes were seen at the order level. Also, the relative abundance of Alteromonadales in the *O. edulis* microbiome was higher in the DSM17395-treated microbiomes (up to 70%) compared to the untreated microbiomes (up to 47%). However, specific ASVs assigned as *Pseudoalteromonas* sp. (belonging to the Alteromonadales) decreased upon the addition of DSM17395, which was also observed in the *E. huxleyi* microbiome treated with DSM17395. Once again, this is similar to the *N. salina* study using pure TDA, where both increases and decreases were found for bacteria belonging to the Alteromonadales when treated with TDA (Geng *et al.* 2016).

The addition of DSM17395 to three common aquaculture live-feeds—the microalgae (*T. suecica*), copepod nauplii (*Acartia tonsa*), and turbot eggs/larvae (*Scophthalmus maximus*)—caused a decrease in relative abundance of closely related taxa particularly of the Rhodobacterales in the microbiomes (Dittmann *et al.* 2020). Specifically, species of *Ruegeria*, *Celeribacter*, and *Pseudophaeobacter* decreased in relative abundance. However, in contrast to the *E. huxleyi* microbiome (Dittmann *et al.* 2019b), bacteria belonging to the Vibrionales and Alteromonadales were not affected by DSM17395 in any of the microbiomes, despite the *S. maximus* microbiome having a high relative abundance of ASVs assigned as *Vibrio* sp. This is surprising since several studies have demonstrated an anti-*Vibrio* effect of TDA-producing *Phaeobacter* compared to a TDA-deficient mutants (D'Alvise *et al.* 2010, 2012), indicating that the effect of DSM17395 toward vibrios depend on the commensal microbiome composition or TDA being species-specific. Future studies should include a TDA-deficient mutant to specifically address the role of TDA in the microbiome development. Such a comparison of the effect of TDA-producing *P. inhibens* 2.10 and its TDA-deficient mutant was conducted on the microbiome assembly of the microalgae *Thalassiosira rotula* (Majzoub *et al.* 2019). Strain 2.10 demonstrated strain-specific effects in the microbiome, in concordance with the previous microbiome studies (Geng *et al.* 2016; Majzoub *et al.* 2019; Dittmann *et al.* 2019b, 2020). Furthermore, closely related strains belonging to the *Sulfotobacter*, *Phaeobacter*, *Pelagicola*, and *Loktanella* genera were reduced or eliminated by the TDA-producing wildtype but not by the TDA-deficient mutant (Majzoub *et al.* 2019).

Overall, the addition of a TDA-producing *P. inhibens* has only minor effects on the taxonomic composition of marine microbial communities. Changes due to the presence of TDA-producing *P. inhibens* strains or TDA appear to be species, if not strain-specific, particularly decreasing the relative abundance of closely related taxa. This is in line with the competition-relatedness concept (Russel *et al.* 2017), where closely related species more often compete for the same metabolic and environmental niches. Niche competition has been suggested to be one of the evolutionary explanations for the selection of antimicrobial compounds—a concept known as competition sensing (Cornforth and Foster 2013). In several studies (Geng *et al.* 2016; Dittmann *et al.* 2019b), the relative abundance of bacteria belonging to the fast-growing heterotrophs of Vibrionales or Alteromonadales were affected by the addition of TDA or the TDA-producing strains, in particular the genus *Pseudoalteromonas*, which generally decreased in relative abundance

(Geng *et al.* 2016; Dittmann *et al.* 2019b). Since TDA can act as an antiporter (Wilson *et al.* 2016), the compound may be particularly effective in antagonizing fast-growing bacteria depending on a high metabolic turnover. Several species of *Pseudoalteromonas* are potent secondary metabolite producers (Paulsen *et al.* 2019), and it has also been suggested that TDA-producing bacteria antagonize specifically potent secondary metabolite producing bacteria found in the same ecological niches (Lutz *et al.* 2016; Dittmann *et al.* 2019b), potentially due to competition sensing (Cornforth and Foster 2013). In fact, the ability of *P. inhibens* to produce TDA has been suggested to be maintained by interspecies competition with *Pseudoalteromonas tunicata* in biofilms (Lutz *et al.* 2016; Majzoub *et al.* 2018). However, the specific mechanism driving this pattern between TDA and *Pseudoalteromonas* species is not fully explored.

Conclusions

TDA is a molecule with multiple functions: antibiotic (disruption of the proton motive force in target bacteria), QS signal, and iron chelation (Fig. 1). The (weak) extracellular iron chelation by TDA could indicate that TDA also can act as an iron provider (reservoir) for other organisms, similar to other weak iron chelators, such as vibrioferrin, suggested to promote bacterial–algal mutualism (Amin *et al.* 2009; Yarimizu *et al.* 2019).

In microbial communities, TDA has predominantly been studied as an antimicrobial compound. It reduces the relative abundance of bacteria closely related to the TDA producer, and sometimes fast-growing, potential secondary metabolite producers, such as vibrios and members of the *Pseudoalteromonas* genus. These observations point toward TDA playing a role in niche competition. The natural concentrations of TDA are not known and we speculate that most of the studies addressing the (antimicrobial) effect of TDA, or its producer are using concentrations of the compound or producer higher than the natural levels. Thus, there is a need for tractable model systems that reflect the natural environment in order to study secondary metabolism and community dynamics (Pessotti, Hansen and Traxler 2018; Gralka *et al.* 2020).

The effects of TDA on gene expression patterns in TDA-producing bacteria indicate that TDA can serve as a QS signal, affecting biofilm formation and motility in the TDA producer. A putative receptor, *pgaR*, is also present in *P. inhibens*, but more studies are necessary to fully understand the molecular mechanism by which TDA regulates gene expression. TDA production may be part of adaptation to a surface-associated lifestyle, possibly in association with eukaryotic host organisms. TDA production in *Tritonibacter* occurs primarily during stagnant growth, whilst in *Phaeobacter*, TDA is also produced in aerated cultures, perhaps reflecting a tighter regulation of TDA biosynthesis in the open water *Tritonibacter* than in its coastal relative *Phaeobacter*.

In conclusion, TDA is indeed an antimicrobial secondary metabolite and as such serves multiple ecological roles such as an algal or coral protectant. However, TDA has other less explored functions being involved in QS regulation. Other antimicrobial secondary metabolites, such as surfactin, also serve multiple functions and this may be the case for several other secondary metabolites. Most studies of antimicrobial secondary metabolites has predominantly been motivated by their antibacterial activity and their effects studied on pure cultures of bacteria, mainly pathogens, using concentrations that are likely higher than those found in natural settings. To fully unravel the roles of antimicrobial secondary metabolites can and may play in natural commu-

nities, we must study the producing organisms and the compound in situ in natural systems.

Acknowledgments

This work was supported by the Danish National Research Foundation to the Center for Microbial Secondary Metabolites (DNRF137) and The Independent Research Fund Denmark (grant number DFF-8048-00035B) to MBT. The authors would like to thank Roberto Kolter for input and discussion on the terminology of secondary metabolites. Lastly, we would like to thank Lars Jelsbak and Aaron J. C. Andersen for input and corrections.

Conflict of interest. The authors declare no conflict of interest.

References

- Abrudan MI, Smakman F, Grimbergen AJ et al. Socially mediated induction and suppression of antibiosis during bacterial coexistence. *Proc Natl Acad Sci* 2015;**112**:11054–9.
- Adamek M, Alanjary M, Sales-Ortells H et al. Comparative genomics reveals phylogenetic distribution patterns of secondary metabolites in *Amycolatopsis* species. *BMC Genomics* 2018;**19**:1–15.
- Alsmark C, Strese Å, Wedén C et al. Microbial diversity of *Alcyonium digitatum*. *Phytochem Rev* 2013;**12**:531–42.
- Amin SA, Green DH, Küpper FC et al. Vibrioferin, an unusual marine siderophore: iron binding, photochemistry, and biological implications. *Inorg Chem* 2009;**48**:11451–8.
- Amin SA, Hmelo LR, Van Tol HM et al. Interaction and signalling between a cosmopolitan phytoplankton and associated bacteria. *Nature* 2015;**522**:98–101.
- Belas R, Horikawa E, Aizawa SI et al. Genetic determinants of *Silicibacter* sp. TM1040 motility. *J Bacteriol* 2009;**191**:4502–12.
- Bérdy J. Bioactive microbial metabolites. *J Antibiot* 2005;**58**:1–26.
- Berger M, Brock NL, Liesegang H et al. Genetic analysis of the upper phenylacetate catabolic pathway in the production of tropodithietic acid by *Phaeobacter gallaeciensis*. *Appl Environ Microbiol* 2012;**78**:3539–51.
- Berger M, Neumann A, Schulz S et al. Tropodithietic acid production in *Phaeobacter gallaeciensis* is regulated by N-acyl homoserine lactone-mediated quorum sensing. *J Bacteriol* 2011;**193**:6576–85.
- Bernbom N, Ng YY, Kjelleberg S et al. Marine bacteria from danish coastal waters show antifouling activity against the marine fouling bacterium *Pseudalteromonas* sp. strain S91 and zoospores of the green alga *Ulva australis* independent of bacteriocidal activity. *Appl Environ Microbiol* 2011;**77**:8557–67.
- Beyersmann PG, Tomasch J, Son K et al. Dual function of tropodithietic acid as antibiotic and signaling molecule in global gene regulation of the probiotic bacterium *Phaeobacter inhibens*. *Sci Rep* 2017;**7**:0–9.
- Blin K, Shaw S, Steinke K et al. AntiSMASH 5.0: updates to the secondary metabolite genome mining pipeline. *Nucleic Acids Res* 2019;**47**:W81–7.
- Bondarev V, Richter M, Romano S et al. The genus *Pseudovibrio* contains metabolically versatile bacteria adapted for symbiosis. *Environ Microbiol* 2013;**15**:2095–113.
- Breider S, Freese HM, Spröer C et al. *Phaeobacter porticola* sp. nov., an antibiotic-producing bacterium isolated from a sea harbour. *Int J Syst Evol Microbiol* 2017;**67**:2153–9.
- Breider S, Scheuner C, Schumann P et al. Genome-scale data suggest reclassifications in the *Leisingera-Phaeobacter* cluster including proposals for *Sedimentitalea* gen. nov. and *Pseudophaeobacter* gen. nov. *Front Microbiol* 2014;**5**:1–13.
- Briand E, Bormans M, Gugger M et al. Changes in secondary metabolic profiles of *Microcystis aeruginosa* strains in response to intraspecific interactions. *Environ Microbiol* 2016;**18**:384–400.
- Brinkhoff T, Bach G, Heidorn T et al. Antibiotic production by a *Roseobacter* clade-affiliated species from the German Wadden Sea and its antagonistic effects on indigenous isolates. *Appl Environ Microbiol* 2004;**70**:2560–5.
- Brock NL, Nikolay A, Dickschat JS. Biosynthesis of the antibiotic tropodithietic acid by the marine bacterium *Phaeobacter inhibens*. *Chem Commun* 2014;**50**:5487–9.
- Bruhn JB, Gram L, Belas R. Production of antibacterial compounds and biofilm formation by *Roseobacter* species are influenced by culture conditions. *Appl Environ Microbiol* 2007;**73**:442–50.
- Bruhn JB, Haagensen JAJ, Bagge-Ravn D et al. Culture conditions of *roseobacter* strain 27-4 affect its attachment and biofilm formation as quantified by real-time PCR. *Appl Environ Microbiol* 2006;**72**:3011–5.
- Bruhn JB, Nielsen KF, Hjelm M et al. Ecology, inhibitory activity, and morphogenesis of a marine antagonistic bacterium belonging to the *Roseobacter* clade. *Appl Environ Microbiol* 2005;**71**:7263–70.
- Buchan A, González JM, Moran MA. Overview of the marine *Roseobacter* lineage. *Appl Environ Microbiol* 2005;**71**:5665–77.
- Chase AB, Sweeney D, Muskat MN et al. Vertical inheritance facilitates interspecies diversification in biosynthetic gene clusters and specialized metabolites. *MBio* 2021;**12**:e02700–21.
- Chen B, Wen J, Zhao X et al. Surfactin: a quorum-sensing signal molecule to relieve CCR in *Bacillus amyloliquefaciens*. *Front Microbiol* 2020;**11**:1–16.
- Chevrette MG, Gutiérrez-García K, Selem-Mojica N et al. Evolutionary dynamics of natural product biosynthesis in bacteria. *Nat Prod Rep* 2020;**37**:566–99.
- Choudhary A, Naughton LM, Dobson ADW et al. High-performance liquid chromatography/electrospray ionisation mass spectrometric characterisation of metabolites produced by *Pseudovibrio* sp. W64, a marine sponge derived bacterium isolated from Irish waters. *Rapid Commun Mass Spectrom* 2018;**32**:1737–45.
- Cornforth DM, Foster KR. Competition sensing: the social side of bacterial stress responses. *Nat Rev Microbiol* 2013;**11**:285–93.
- Crowley S, O’Gara F, O’Sullivan O et al. Marine *Pseudovibrio* sp. as a novel source of antimicrobials. *Mar Drugs* 2014;**12**:5916–29.
- Curson ARJ, Todd JD, Sullivan MJ et al. Catabolism of dimethylsulphoniopropionate: microorganisms, enzymes and genes. *Nat Rev Microbiol* 2011;**9**:849–59.
- Czapek F. *Biochemie Der Pflanzen*. 3rd edn. Jena: G. Fischer, 1922.
- D’Alvise PW, Lillebø S, Prol-Garcia MJ et al. *Phaeobacter gallaeciensis* reduces *Vibrio anguillarum* in cultures of microalgae and rotifers, and prevents vibriosis in cod larvae. *PLoS ONE* 2012;**7**:e43996.
- D’Alvise PW, Magdenoska O, Melchiorson J et al. Biofilm formation and antibiotic production in *Ruegeria mobilis* are influenced by intracellular concentrations of cyclic dimeric guanosinmonophosphate. *Environ Microbiol* 2014;**16**:1252–66.
- D’Alvise PW, Melchiorson J, Porsby CH et al. Inactivation of *Vibrio anguillarum* by attached and planktonic *roseobacter* cells. *Appl Environ Microbiol* 2010;**76**:2366–70.
- D’Alvise PW, Phippen CBW, Nielsen KF et al. Influence of iron on production of the antibacterial compound tropodithietic acid and its noninhibitory analog in *Phaeobacter inhibens*. *Appl Environ Microbiol* 2016;**82**:502–9.
- Danevčić T, Dragoš A, Spacapan M et al. Surfactin facilitates horizontal gene transfer in *Bacillus subtilis*. *Front Microbiol* 2021;**12**:1–8.
- Dang H, Lovell CR. Bacterial primary colonization and early succession on surfaces in marine waters as determined by ampli-

- fied rRNA gene restriction analysis and sequence analysis of 16S rRNA genes. *Appl Environ Microbiol* 2000;**66**:467–75.
- Das T, Manefield M. Pyocyanin promotes extracellular DNA release in *Pseudomonas aeruginosa*. *PLoS ONE* 2012;**7**. DOI: 10.1371/journal.pone.0046718.
- Davies J. Specialized microbial metabolites: functions and origins. *J Antibiot* 2013;**66**:361–4.
- Dickschat JS, Rinkel J, Klapschinski T et al. Characterisation of the l-cystine β -lyase PatB from *Phaeobacter inhibens*: an enzyme involved in the biosynthesis of the marine antibiotic tropodithietic acid. *ChemBioChem* 2017;**18**:2260–7.
- Dietrich LEP, Price-Whelan A, Petersen A et al. The phenazine pyocyanin is a terminal signalling factor in the quorum sensing network of *Pseudomonas aeruginosa*. *Mol Microbiol* 2006;**61**:1308–21.
- Dittmann KK, Porsby CH, Goncalves P et al. Tropodithietic acid induces oxidative stress response, cell envelope biogenesis and iron uptake in *Vibrio vulnificus*. *Environ Microbiol Rep* 2019a;**11**:581–8.
- Dittmann KK, Rasmussen BB, Castex M et al. The aquaculture microbiome at the centre of business creation. *Microb Biotechnol* 2017;**10**:1279–82.
- Dittmann KK, Rasmussen BB, Melchiorson J et al. Changes in the microbiome of mariculture feed organisms after treatment with a potentially probiotic strain of *Phaeobacter inhibens*. *Appl Environ Microbiol* 2020;**86**:1–17.
- Dittmann KK, Sonnenschein EC, Egan S et al. Impact of *Phaeobacter inhibens* on marine eukaryote-associated microbial communities. *Environ Microbiol Rep* 2019b;**11**:401–13.
- Duan Y, Petzold M, Saleem-Batcha R et al. Bacterial tropone natural products and derivatives: overview of their biosynthesis, bioactivities, ecological role and biotechnological potential. *ChemBioChem* 2020;**21**:2384–407.
- Duan Y, Toplak M, Hou A et al. A flavoprotein dioxygenase steers bacterial tropone biosynthesis via coenzyme A-ester oxygenolysis and ring epoxidation. *J Am Chem Soc* 2021;**143**:10413–21.
- Enticknap JJ, Kelly M, Peraud O et al. Characterization of a culturable alphaproteobacterial symbiont common to many marine sponges and evidence for vertical transmission via sponge larvae. *Appl Environ Microbiol* 2006;**72**:3724–32.
- Firn RD, Jones CG. The evolution of secondary metabolism - a unifying model. *Mol Microbiol* 2000;**37**:989–94.
- Fischbach MA, Walsh CT, Clardy J. The evolution of gene collectives: how natural selection drives chemical innovation. *Proc Natl Acad Sci* 2008;**105**:4601–8.
- Fleming A. On the antibacterial action of cultures of a *Penicillium*, with special reference to their use in the isolation of *Bacillus influenzae*. *Br J Exp Pathol* 1929;**10**:226.
- Foster KR, Bell T. Competition, not cooperation, dominates interactions among culturable microbial species. *Curr Biol* 2012;**22**:1845–50.
- Freese HM, Methner A, Overmann J. Adaptation of surface-associated bacteria to the open ocean: a genomically distinct subpopulation of *Phaeobacter gallaeciensis* colonizes pacific mesozooplankton. *Front Microbiol* 2017;**8**:1–12.
- Gallagher KA, Wanger G, Henderson J et al. Ecological implications of hypoxia-triggered shifts in secondary metabolism. *Environ Microbiol* 2017;**19**:2182–91.
- Geier B, Sogin EM, Michellod D et al. Spatial metabolomics of in situ host–microbe interactions at the micrometre scale. *Nat Microbiol* 2020;**5**:498–510.
- Geng H, Belas R. Expression of tropodithietic acid biosynthesis is controlled by a novel autoinducer. *J Bacteriol* 2010;**192**:4377–87.
- Geng H, Belas R. TdaA regulates tropodithietic acid synthesis by binding to the tdaC promoter region. *J Bacteriol* 2011;**193**:4002–5.
- Geng H, Bruhn JB, Nielsen KF et al. Genetic dissection of tropodithietic acid biosynthesis by marine roseobacters. *Appl Environ Microbiol* 2008;**74**:1535–45.
- Geng H, Tran-Gyamfi MB, Lane TW et al. Changes in the structure of the microbial community associated with *Nannochloropsis salina* following treatments with antibiotics and bioactive compounds. *Front Microbiol* 2016;**7**:1–13.
- Goh EB, Yim G, Tsui W et al. Transcriptional modulation of bacterial gene expression by subinhibitory concentrations of antibiotics. *Proc Natl Acad Sci* 2002;**99**:17025–30.
- González JM, Moran MA. Numerical dominance of a group of marine bacteria in the alpha-subclass of the class proteobacteria in coastal seawater. *Appl Environ Microbiol* 1997;**63**:4237–42.
- Gonzalez JM, Simo R, Massana R et al. Bacterial community structure associated with a dimethylsulfoniopropionate-producing North Atlantic algal bloom. *Appl Environ Microbiol* 2000;**66**:4237–46.
- Gralka M, Szabo R, Stocker R et al. Trophic interactions and the drivers of microbial community assembly. *Curr Biol* 2020;**30**:R1176–88.
- Gram L, Grossart H-P, Schlingloff A et al. Possible quorum sensing in marine snow bacteria: production of acylated homoserine lactones by *Roseobacter* strains isolated from marine snow. *Appl Environ Microbiol* 2002;**68**:4111–6.
- Gram L, Melchiorson J, Bruhn JB. Antibacterial activity of marine culturable bacteria collected from a global sampling of ocean surface waters and surface swabs of marine organisms. *Mar Biotechnol* 2010;**12**:439–51.
- Gram L, Rasmussen BB, Wemheuer B et al. *Phaeobacter inhibens* from the *Roseobacter* clade has an environmental niche as a surface colonizer in harbors. *Syst Appl Microbiol* 2015;**38**:483–93.
- Grotkjær T, Bentzon-Tilia M, D'Alvise P et al. *Phaeobacter inhibens* as probiotic bacteria in non-axenic *Artemia* and algae cultures. *Aquaculture* 2016a;**462**:64–9.
- Grotkjær T, Bentzon-Tilia M, D'Alvise PW et al. Isolation of TDA-producing *Phaeobacter* strains from sea bass larval rearing units and their probiotic effect against pathogenic *Vibrio* spp. in *Artemia* cultures. *Syst Appl Microbiol* 2016b;**39**:180–8.
- Harrington C, Reen F, Mooij M et al. Characterisation of non-autoinducing tropodithietic acid (TDA) production from marine sponge *Pseudovibrio* species. *Mar Drugs* 2014;**12**:5960–78.
- Hernandez ME, Newman DK. Extracellular electron transfer. *Cell Mol Life Sci* 2001;**58**:1562–71.
- Hjelm M, Bergh Ø, Riaza A et al. Selection and identification of autochthonous potential probiotic bacteria from turbot larvae (*Scophthalmus maximus*) rearing units. *Syst Appl Microbiol* 2004a;**27**:360–71.
- Hjelm M, Riaza A, Formoso F et al. Seasonal incidence of autochthonous antagonistic *Roseobacter* spp. and *Vibrionaceae* strains in a turbot larva (*Scophthalmus maximus*) rearing system. *Appl Environ Microbiol* 2004b;**70**:7288–94.
- Hoff G, Arguelles-Arias A, Boubsi F et al. Surfactin stimulated by pectin molecular patterns and root exudates acts as a key driver of *Bacillus*-plant mutualistic interaction. *mBio* 2021;**12**:e01774–21.
- Kawano Y, Nagawa Y, Nakanishi H et al. Production of thiotropocin by a marine bacterium, *Caulobacter* sp. and its antimicrobial activities. *J Mar Biotechnol* 1997;**5**:225–9.
- Kiene RP, Linn LJ, Gonzalez J et al. Dimethylsulfoniopropionate and methanethiol are important precursors of methionine and protein-sulfur in marine bacterioplankton. *Appl Environ Microbiol* 1999;**65**:4549–58.
- Kintaka K, Ono H, Tsubotani S et al. Thiotropocin, a new sulfur-containing 7-membered-ring antibiotic produced by a *Pseudomonas* sp. *J Antibiot* 1984;**37**:1294–300.

- Kossel A. Über die chemische Zusammensetzung der zelle. *Archiv für Physiologie*. 1891;181–6. Zeitschrift.
- Lauzon HL, Gudmundsdottir S, Pedersen MH et al. Isolation of putative probiotics from cod rearing environment. *Vet Microbiol* 2008;**132**:328–39.
- Lee M-H, Song E-J, Seo M-J et al. *Phaeobacter marinintestinus* sp. nov., isolated from the intestine of a sea cucumber (*Apostichopus japonicus*). *Antonie Van Leeuwenhoek* 2015;**107**:209–16.
- Li A, Okada BK, Rosen PC et al. Piperacillin triggers virulence factor biosynthesis via the oxidative stress response in *Burkholderia thailandensis*. *Proc Natl Acad Sci* 2021;**118**:e2021483118.
- Li B, Ry RF, Albert AB et al. A backup plan for self-protection: s-methylation of holomycin biosynthetic intermediates in *Streptomyces clavuligerus*. *ChemBioChem* 2013;**13**:2521–6.
- Liang L. Investigation of secondary metabolites of North Sea bacteria: fermentation, isolation, structure elucidation and bioactivity. Ph.D. Thesis, Mathematisch-Naturwissenschaftlichen Fakultäten der Georg-August-Universität Göttingen, 2003.
- Linares JF, Gustafsson I, Baquero F et al. Antibiotics as intermicrobiol signaling agents instead of weapons. *Proc Natl Acad Sci* 2006;**103**:19484–9.
- Lind AL, Wisecaver JH, Lameiras C et al. Drivers of genetic diversity in secondary metabolic gene clusters within a fungal species. Kamoun S (ed.). *PLoS Biol* 2017;**15**:e2003583.
- Liu Z, Wang W, Zhu Y et al. Antibiotics at subinhibitory concentrations improve the quorum sensing behavior of *Chromobacterium violaceum*. *FEMS Microbiol Lett* 2013;**341**:37–44.
- Long RA, Azam F. Antagonistic interactions among marine pelagic bacteria. *Appl Environ Microbiol* 2001;**67**:4975–83.
- López D, Fischbach MA, Chu F et al. Structurally diverse natural products that cause potassium leakage trigger multicellularity in *Bacillus subtilis*. *Proc Natl Acad Sci* 2009;**106**:280–5.
- Lutz C, Thomas T, Steinberg P et al. Effect of interspecific competition on trait variation in *Phaeobacter inhibens* biofilms. *Environ Microbiol* 2016;**18**:1635–45.
- Majzoub ME, Beyersmann PG, Simon M et al. *Phaeobacter inhibens* controls bacterial community assembly on a marine diatom. *FEMS Microbiol Ecol* 2019;**95**:1–12.
- Majzoub ME, McElroy K, Maczka M et al. Causes and consequences of a variant strain of *Phaeobacter inhibens* with reduced competition. *Front Microbiol* 2018;**9**:1–10.
- Majzoub ME, McElroy K, Maczka M et al. Genomic evolution of the marine bacterium *Phaeobacter inhibens* during biofilm growth. *Appl Environ Microbiol* **87**, 2021. DOI: 10.1128/AEM.00769-21.
- Mansson M, Nielsen A, Kjærulff L et al. Inhibition of virulence gene expression in *Staphylococcus aureus* by novel depsipeptides from a marine *Photobacterium* sp. *Mar Drugs* 2011;**9**:2537–52.
- Martens T, Gram L, Grossart HP et al. Bacteria of the *Roseobacter* clade show potential for secondary metabolite production. *Microb Ecol* 2007;**54**:31–42.
- Martens T, Heidorn T, Pukall R et al. Reclassification of *Roseobacter gallaeciensis* Ruiz-Ponte et al. 1998 as *Phaeobacter gallaeciensis* gen. nov., comb. nov., description of *Phaeobacter inhibens* sp. nov., reclassification of *Ruegeria algicola* (Lafay et al. 1995) Uchino et al. 1999 as *Marinovum algicola* gen. nov., comb. nov., and emended descriptions of the genera *Roseobacter*, *Ruegeria* and *Leisingera*. *Int J Syst Evol Microbiol* 2006;**56**:1293–304.
- Medema MH, Cimermancic P, Sali A et al. A systematic computational analysis of biosynthetic gene cluster evolution: lessons for engineering biosynthesis. *PLoS Comput Biol* 2014;**10**:e1004016.
- Miller TR, Belas R. Dimethylsulfoniopropionate metabolism by *Pfesteria*-associated *Roseobacter* spp. *Appl Environ Microbiol* 2004;**70**:3383–91.
- Miller TR, Hnilicka K, Dziedzic A et al. Chemotaxis of *silicibacter* sp. strain TM1040 toward dinoflagellate products. *Appl Environ Microbiol* 2004;**70**:4692–701.
- Moree WJ, Phelan V V, Wu C-H et al. Interkingdom metabolic transformations captured by microbial imaging mass spectrometry. *Proc Natl Acad Sci* 2012;**109**:13811–6.
- Muramatsu Y, Uchino Y, Kasai H et al. *Ruegeria mobilis* sp. nov., a member of the alphaproteobacteria isolated in Japan and Palau. *Int J Syst Evol Microbiol* 2007;**57**:1304–9.
- Nappi J, Soldi E, Egan S. Diversity and distribution of bacteria producing known secondary metabolites. *Microb Ecol* 2019;**78**:885–94.
- Newton RJ, Griffin LE, Bowles KM et al. Genome characteristics of a generalist marine bacterial lineage. *ISME J* 2010;**4**:784–98.
- Paulsen SS, Strube ML, Bech PK et al. Marine chitinolytic *Pseudoalteromonas* represents an untapped reservoir of bioactive potential. *mSystems* 2019;**4**:e00060–19.
- Penesyan A, Tebben J, Lee M et al. Identification of the antibacterial compound produced by the marine epiphytic bacterium *Pseudovibrio* sp. D323 and related sponge-associated bacteria. *Mar Drugs* 2011;**9**:1391–402.
- Pessotti RC, Hansen BL, Traxler MF. In search of model ecological systems for understanding specialized metabolism. *mSystems* 2018;**3**:1–6.
- Petersen J, Frank O, Göker M et al. Extrachromosomal, extraordinary and essential—the plasmids of the *Roseobacter* clade. *Appl Microbiol Biotechnol* 2013;**97**:2805–15.
- Phippen CBW, Jørgensen CM, Bentzon-Tilia M et al. Isolation of methyl troposulfenin from *Phaeobacter inhibens*. *J Nat Prod* 2019;**82**:1387–90.
- Pishchany G, Kolter R. On the possible ecological roles of antimicrobials. *Mol Microbiol* 2020;**113**:580–7.
- Planas M, Pérez-Lorenzo M, Hjelm M et al. Probiotic effect in vivo of *Roseobacter* strain 27-4 against *Vibrio* (*Listonella*) *Anguillarum* infections in turbot (*Scophthalmus maximus* L.) larvae. *Aquaculture* 2006;**255**:323–33.
- Porsby CH, Gram L. *Phaeobacter inhibens* as biocontrol agent against *Vibrio vulnificus* in oyster models. *Food Microbiol* 2016;**57**:63–70.
- Porsby CH, Nielsen KF, Gram L. *Phaeobacter* and *Ruegeria* species of the *Roseobacter* clade colonize separate niches in a danish turbot (*Scophthalmus maximus*)-rearing farm and antagonize *Vibrio anguillarum* under different growth conditions. *Appl Environ Microbiol* 2008;**74**:7356–64.
- Porsby CH, Webber MA, Nielsen KF et al. Resistance and tolerance to tropodithietic acid, an antimicrobial in aquaculture, is hard to select. *Antimicrob Agents Chemother* 2011;**55**:1332–7.
- Prado S, Montes J, Romalde JL et al. Inhibitory activity of *Phaeobacter* strains against aquaculture pathogenic bacteria. *Int Microbiol* 2009;**12**:107–14.
- Price-Whelan A, Dietrich LEP, Newman DK. Rethinking “secondary” metabolism: physiological roles for phenazine antibiotics. *Nat Chem Biol* 2006;**2**:71–8.
- Prol García MJ, D’Alvise PW, Gram L. Disruption of cell-to-cell signaling does not abolish the antagonism of *Phaeobacter gallaeciensis* toward the fish pathogen *Vibrio anguillarum* in algal systems. *Appl Environ Microbiol* 2013;**79**:5414–7.
- Prol García MJ, D’Alvise PW, Rygaard AM et al. Biofilm formation is not a prerequisite for production of the antibacterial compound tropodithietic acid in *Phaeobacter inhibens* DSM17395. *J Appl Microbiol* 2014;**117**:1592–600.

- Rabe P, Klapschinski TA, Brock NL et al. Synthesis and bioactivity of analogues of the marine antibiotic tropodithietic acid. *Beilstein J Org Chem* 2014;**10**:1796–801.
- Raina JB, Tapiolas D, Motti CA et al. Isolation of an antimicrobial compound produced by bacteria associated with reef-building corals. *PeerJ* 2016;**4**:e2275.
- Raina JB, Tapiolas D, Willis BL et al. Coral-associated bacteria and their role in the biogeochemical cycling of sulfur. *Appl Environ Microbiol* 2009;**75**:3492–501.
- Rao D, Webb JS, Holmström C et al. Low densities of epiphytic bacteria from the marine alga *Ulva australis* inhibit settlement of fouling organisms. *Appl Environ Microbiol* 2007;**73**:7844–52.
- Rao D, Webb JS, Kjelleberg S. Microbial colonization and competition on the marine alga *Ulva australis*. *Appl Environ Microbiol* 2006;**72**:5547–55.
- Rasmussen BB, Erner KE, Bentzon-Tilia M et al. Effect of TDA-producing *Phaeobacter inhibens* on the fish pathogen *Vibrio anguillarum* in non-axenic algae and copepod systems. *Microb Biotechnol* 2018;**11**:1070–9.
- Rasmussen BB, Grotkjær T, D'Alvise PW et al. *Vibrio anguillarum* is genetically and phenotypically unaffected by long-term continuous exposure to the antibacterial compound tropodithietic acid. *Appl Environ Microbiol* 2016;**82**:4802–10.
- Rasmussen BB, Kalatzis PG, Middelboe M et al. Combining probiotic *Phaeobacter inhibens* DSM17395 and broad-host-range vibriophage KVP40 against fish pathogenic vibrios. *Aquaculture* 2019;**513**:734415.
- Ringø E. Probiotics in shellfish aquaculture. *Aquacul Fish* 2020;**5**:1–27.
- Romano S, Schulz-Vogt HN, González JM et al. Phosphate limitation induces drastic physiological changes, virulence-related gene expression, and secondary metabolite production in *Pseudovibrio* sp. Strain FO-BEG1. *Appl Environ Microbiol* 2015;**81**:3518–28.
- Romano S. Ecology and biotechnological potential of bacteria belonging to the genus *Pseudovibrio*. *Appl Environ Microbiol* 2018;**84**:1–16.
- Romero D, Traxler MF, López D et al. Antibiotics as signal molecules. *Chem Rev* 2011;**111**:5492–505.
- Ruiz-Ponte C, Cilia V, Lambert C et al. *Roseobacter gallaeciensis* sp. nov., a new marine bacterium isolated from rearings and collectors of the scallop *Pecten maximus*. *Int J Syst Bacteriol* 1998;**48**:537–42.
- Ruiz-Ponte C, Samain JF, Sánchez JL et al. The benefit of a *Roseobacter* species on the survival of scallop larvae. *Mar Biotechnol* 1999;**1**:52–9.
- Russel J, Røder HL, Madsen JS et al. Antagonism correlates with metabolic similarity in diverse bacteria. *Proc Natl Acad Sci* 2017;**114**:10684–8.
- Rypien KL, Ward JR, Azam F. Antagonistic interactions among coral-associated bacteria. *Environ Microbiol* 2010;**12**:28–39.
- Segev E, Wyche TP, Kim KH et al. Dynamic metabolic exchange governs a marine algal-bacterial interaction. *Elife* 2016;**5**:1–28.
- Seyedsayamdost MR, Carr G, Kolter R et al. Roseobactin: small molecule modulators of an algal-bacterial symbiosis. *J Am Chem Soc* 2011a;**133**:18343–9.
- Seyedsayamdost MR, Case RJ, Kolter R et al. The Jekyll-and-Hyde chemistry of *Phaeobacter gallaeciensis*. *Nat Chem* 2011b;**3**:331–5.
- Shnit-Orland M, Kushmaro A. Coral mucus-associated bacteria: a possible first line of defense. *FEMS Microbiol Ecol* 2009;**67**:371–80.
- Simon M, Scheuner C, Meier-Kolthoff JP et al. Phylogenomics of *Rhodobacteraceae* reveals evolutionary adaptation to marine and non-marine habitats. *ISME J* 2017;**11**:1483–99.
- Sonnenschein EC, Jimenez G, Castex M et al. The *Roseobacter*-group bacterium *phaeobacter* as a safe probiotic solution for aquaculture. *Appl Environ Microbiol* 2021;**87**:1–15.
- Sonnenschein EC, Nielsen KF, D'Alvise P et al. Global occurrence and heterogeneity of the *Roseobacter*-clade species *R. uegeria mobilis*. *ISME J* 2017a;**11**:569–83.
- Sonnenschein EC, Phippen CBW, Bentzon-Tilia M et al. Phylogenetic distribution of roseobactin in the *Roseobacter* group and their effect on microalgae. *Environ Microbiol Rep* 2018;**10**:383–93.
- Sonnenschein EC, Phippen CBW, Nielsen KF et al. *Phaeobacter piscinae* sp. nov., a species of the *Roseobacter* group and potential aquaculture probiont. *Int J Syst Evol Microbiol* 2017b;**67**:4559–64.
- Straight PD, Willey JM, Kolter R. Interactions between *Streptomyces coelicolor* and *Bacillus subtilis*: role of surfactants in raising aerial structures. *J Bacteriol* 2006;**188**:4918–25.
- Sunagawa S, Coelho LP, Chaffron S et al. Structure and function of the global ocean microbiome. *Science* 2015;**348**:1261359.
- Tenconi E, Traxler MF, Hoebreck C et al. Production of prodiginines is part of a programmed cell death process in *Streptomyces coelicolor*. *Front Microbiol* 2018;**9**:1742.
- Tesdorpf JE, Geers AU, Strube ML et al. *Roseobacter* group probiotics exhibit differential killing of fish pathogenic *Tenacibaculum* species. *Appl Environ Microbiol* 2022;aem0241821.
- Teufel R, Gantert C, Voss M et al. Studies on the mechanism of ring hydrolysis in phenylacetate degradation. *J Biol Chem* 2011;**286**:11021–34.
- Teufel R, Mascaraque V, Ismail W et al. Bacterial phenylalanine and phenylacetate catabolic pathway revealed. *Proc Natl Acad Sci* 2010;**107**:14390–5.
- Thole S, Kalhoefer D, Voget S et al. *Phaeobacter gallaeciensis* genomes from globally opposite locations reveal high similarity of adaptation to surface life. *ISME J* 2012;**6**:2229–44.
- Trautwein K, Will SE, Hulsch R et al. Native plasmids restrict growth of *Phaeobacter inhibens* DSM 17395: energetic costs of plasmids assessed by quantitative physiological analyses. *Environ Microbiol* 2016;**18**:4817–29.
- Traxler MF, Kolter R. Natural products in soil microbe interactions and evolution. *Nat Prod Rep* 2015;**32**:956–70.
- Tsubotani S, Wada Y, Kamiya K et al. Structure of thiotropocin, a new sulfur-containing 7-membered antibiotic. *Tetrahedron Lett* 1984;**25**:419–22.
- Undabarrena A, Valencia R, Cumsille A et al. *Rhodococcus* comparative genomics reveals a phylogenomic-dependent non-ribosomal peptide synthetase distribution: insights into biosynthetic gene cluster connection to an orphan metabolite. *Microbial Genomics* 2021;**7**:621.
- Vizcaino MI, Johnson WR, Kimes NE et al. Antimicrobial resistance of the coral pathogen *Vibrio coralliilyticus* and Caribbean sister phylotypes isolated from a diseased octocoral. *Microb Ecol* 2010;**59**:646–57.
- Wang R, Gallant É, Seyedsayamdost MR. Investigation of the genetics and biochemistry of roseobactin production in the *Roseobacter* clade bacterium *Phaeobacter inhibens*. *MBio* 2016;**7**:1–10.
- Wang R, Seyedsayamdost MR. Roseochelin b, an algacidal natural product synthesized by the roseobacter *Phaeobacter inhibens* in response to algal sinapic acid. *Org Lett* 2017;**19**:5138–41.
- West NJ, Obernosterer I, Zemb O et al. Major differences of bacterial diversity and activity inside and outside of a natural iron-fertilized phytoplankton bloom in the Southern Ocean. *Environ Microbiol* 2008;**10**:738–56.
- Wichmann H, Brinkhoff T, Simon M et al. Dimethylsulfoniopropionate promotes process outgrowth in neural cells and exerts protective effects against tropodithietic acid. *Mar Drugs* 2016;**14**. DOI: 10.3390/md14050089.

- Wichmann H, Vocke F, Brinkhoff T et al. Cytotoxic effects of tropodithietic acid on mammalian clonal cell lines of neuronal and glial origin. *Mar Drugs* 2015;**13**:7113–23.
- Wietz M, Gram L, Jørgensen B et al. Latitudinal patterns in the abundance of major marine bacterioplankton groups. *Aquat Microb Ecol* 2010;**61**:179–89.
- Will SE, Neumann-Schaal M, Heydorn RL et al. The limits to growth – energetic burden of the endogenous antibiotic tropodithietic acid in *Phaeobacter inhibens* DSM 17395. *PLoS ONE* 2017;**12**:e0177295.
- Wilson MZ, Wang R, Gitai Z et al. Mode of action and resistance studies unveil new roles for tropodithietic acid as an anticancer agent and the γ -glutamyl cycle as a proton sink. *Proc Natl Acad Sci* 2016;**113**:1630–5.
- Wirth JS, Whitman WB. Phylogenomic analyses of a clade within the roseobacter group suggest taxonomic reassignments of species of the genera *Aestuaria*, *Citricella*, *Loktanella*, *Nautella*, *Pelagibaca*, *Ruegeria*, *Thalassobius*, *Thiobacimonas* and *Tropicibacter*, and the proposal of six novel genera. *Int J Syst Evol Microbiol* 2018;**68**: 2393–411.
- Wünsch D, Strijkstra A, Wöhlbrand L et al. Global response of *Phaeobacter inhibens* DSM 17395 to deletion of its 262-kb chromid encoding antibiotic synthesis. *Microb Physiol* 2020;**30**:9–24.
- Yarimizu K, Cruz-López R, García-Mendoza E et al. Distribution of dissolved iron and bacteria producing the photoactive siderophore, vibrioferrin, in waters off Southern California and Northern Baja. *Biometals* 2019;**32**:139–54.
- Yim G, Huimi Wang H, Davies J. Antibiotics as signalling molecules. *Philos Trans R Soc B Biol Sci* 2007;**362**:1195–200.
- Yoon V, Nodwell JR. Activating secondary metabolism with stress and chemicals. *J Ind Microbiol Biotechnol* 2014;**41**:415–24.
- Zan J, Liu Y, Fuqua C et al. Acyl-homoserine lactone quorum sensing in the *Roseobacter* clade. *Int J Mol Sci* 2014;**15**:654–69.
- Zech H, Hensler M, Koßmehl S et al. Dynamics of amino acid utilization in *Phaeobacter inhibens* DSM 17395. *Proteomics* 2013;**13**: 2869–85.
- Zhang S-D, Isbrandt T, Lindqvist LL et al. Holomycin, an antibiotic secondary metabolite, is required for biofilm formation by the native producer *Photobacterium galathea* S2753. *Appl Environ Microbiol* 2021;**87**:e00169–21.
- Zhang W, Liang W, Li C. Inhibition of marine *Vibrio* sp. by pyoverdine from *Pseudomonas aeruginosa* PA1. *J Hazard Mater* 2016;**302**:217–24.
- Zhao W, Dao C, Karim M et al. Contributions of tropodithietic acid and biofilm formation to the probiotic activity of *Phaeobacter inhibens*. *BMC Microbiol* 2016;**16**:1–17.