

Role of Adaptor Complex AP-3 in Targeting Wild-Type and Mutated CD63 to Lysosomes

Brian A. Rous,* Barbara J. Reaves,[†] Gudrun Ihrke,* John A.G. Briggs,*
Sally R. Gray,* David J. Stephens,[‡] George Banting,[‡] and J. Paul Luzio[§]

*University of Cambridge, Department of Clinical Biochemistry, Cambridge Institute for Medical Research, Cambridge, CB2 2XY, United Kingdom; [†]Department of Biology and Biochemistry, University of Bath, Bath BA2 7AY, United Kingdom; and [‡]Department of Biochemistry, School of Medical Sciences, University of Bristol, Bristol BS8 1TD, United Kingdom

Submitted August 15, 2001; Revised December 5, 2001; Accepted December 5, 2001
Monitoring Editor: Juan Bonifacino

CD63 is a lysosomal membrane protein that belongs to the tetraspanin family. Its carboxyterminal cytoplasmic tail sequence contains the lysosomal targeting motif GYEVM. Strong, tyrosine-dependent interaction of the wild-type carboxyterminal tail of CD63 with the AP-3 adaptor subunit $\mu 3$ was observed using a yeast two-hybrid system. The strength of interaction of mutated tail sequences with $\mu 3$ correlated with the degree of lysosomal localization of similarly mutated human CD63 molecules in stably transfected normal rat kidney cells. Mutated CD63 containing the cytosolic tail sequence GYEVI, which interacted strongly with $\mu 3$ but not at all with $\mu 2$ in the yeast two-hybrid system, localized to lysosomes in transfected normal rat kidney and NIH-3T3 cells. In contrast, it localized to the cell surface in transfected cells of *pearl* and *mocha* mice, which have genetic defects in genes encoding subunits of AP-3, but to lysosomes in functionally rescued *mocha* cells expressing the δ subunit of AP-3. Thus, AP-3 is absolutely required for the delivery of this mutated CD63 to lysosomes. Using this AP-3-dependent mutant of CD63, we have shown that AP-3 functions in membrane traffic from the *trans*-Golgi network to lysosomes via an intracellular route that appears to bypass early endosomes.

INTRODUCTION

Lysosomal integral membrane glycoproteins (lysosomal glycoproteins [lgps], lysosomal integral membrane proteins [LIMPs], lysosome-associated membrane proteins [LAMPs]) have short, 10–20 amino acid, cytosolic tails containing tyrosine-based or di-leucine-based motifs that are known, or thought, to determine their lysosomal targeting (Hunziker and Geuze, 1996). The tyrosine-based signals are of the form YXX \emptyset where X is any amino acid and \emptyset is a bulky hydrophobic amino acid, and are clearly related to internalization signals of the same type, which interact with the $\mu 2$ subunit of the AP-2 adaptor found in clathrin-coated pits at the cell surface (Bonifacino and Dell'Angelica, 1999). Lysosomal targeting YXX \emptyset motifs are often located at the carboxy termi-

nus of short cytosolic tails preceded by a glycine. Their effectiveness may be modified by spacing relative to the membrane bilayer (Rohrer *et al.*, 1996), the identity of the carboxyterminal amino acid (Gough and Fambrough, 1997; Gough *et al.*, 1999), or proteolytic modification (Guarnieri *et al.*, 1993; Akasaki *et al.*, 1995). There is evidence that the luminal (Reaves *et al.*, 1998) and transmembrane domains (Wimer-Mackin and Granger, 1996) of some lgps also contain targeting information.

Newly synthesized lgps with GYXX \emptyset motifs in their carboxyterminal cytosolic tails take both direct and indirect (via the plasma membrane) traffic routes from the *trans*-Golgi network (TGN) to lysosomes. The evidence for individual lgps taking mainly one or other of these routes has come from kinetic studies of delivery and from the endocytic uptake of anti-lgp antibodies or surface-labeled lgps from the plasma membrane. Thus, the type I integral membrane proteins LAMP-1 and LAMP-2, which between them account for >50% of the lysosomal content of lgps (Andrejewski *et al.*, 1999), are delivered from the TGN with half-times of 30–90 min, mainly via an intracellular route (Barriocanal *et al.*, 1986; D'Souza and August, 1986; Green *et al.*, 1987; Carlsson and Fukuda, 1992; Akasaki *et al.*, 1995, 1996). In contrast, lysosomal acid phosphatase, also a type I mem-

Article published online ahead of print. Mol. Biol. Cell 10.1091/mbc.01-08-0409. Article and publication date are at www.molbiolcell.org/cgi/doi/10.1091/mbc.01-08-0409.

[§] Corresponding author. E-mail address: jpl10@cam.ac.uk.

Abbreviations used: AP, adaptor protein; 3-AT, 3-amino-1,2,4-triazole; lgp, lysosomal glycoprotein; MPR, mannose 6-phosphate receptor; NRK, normal rat kidney; TGN, *trans*-Golgi network.

brane protein with a cytosolic tail GYXXØ motif, is delivered with a half-time of 5–7 h, mainly via the cell surface (Braun *et al.*, 1989).

The nature of the vesicular carriers used by Igps on the direct route from the TGN to late endocytic compartments is a matter of some dispute. Evidence that the intracellular route taken by some Igps from the TGN to lysosomes involves adaptor protein (AP)-1-associated clathrin-coated vesicles has mainly come from data showing that in vitro the LAMP-1 cytosolic domain interacts with both purified AP-1 and AP-2 and that in vivo, a small fraction, ~3%, of intracellular LAMP-1 is present with mannose 6-phosphate receptors (MPRs) in AP-1-associated clathrin-coated vesicles budding from the TGN (Honing *et al.*, 1996). However, more recent studies showed first, that LAMP-1 and LAMP-2 were mostly in separate TGN-derived vesicles from those containing MPR and AP-1 (Karlsson and Carlsson, 1998), and second, that in μ 1A-deficient cells no change in steady-state distribution or missorting to the cell surface of LAMP-1 occurred, in contrast to the altered MPR distribution (Meyer *et al.*, 2000). The recently discovered clathrin adaptors known as Golgi-localized, γ -ear containing, ARF-binding proteins which are the best candidates for signal-mediated sorting of MPRs at the TGN, appear to play no role in LAMP-1 delivery to lysosomes (Puertollano *et al.*, 2001). Over the past 5 yr considerable, but mostly indirect, evidence has been obtained to suggest that AP-3 plays a role in the targeting of Igps to lysosomes and lysosome-like organelles (reviewed in Hirst and Robinson, 1998; Gu and Gruenberg, 1999; Luzio *et al.*, 2000; Huizing *et al.*, 2001; Mullins and Bonifacino, 2001). In yeast it has been established that two different trafficking routes from the late Golgi to the lysosome exist, which are followed by different newly synthesized proteins, and one of which is AP-3 dependent (Cowles *et al.*, 1997; Piper *et al.*, 1997; Stepp *et al.*, 1997).

To obtain further information about the targeting of Igps to lysosomes and the role of AP-3, we have studied lysosomal targeting of CD63 (LIMP-I, LAMP-3), a member of the tetraspanin family of integral membrane proteins with four predicted transmembrane domains (Metzelaar *et al.*, 1991; Maecker *et al.*, 1997). CD63 was first described as an antigen present on the surface of activated blood platelets after transfer from dense granules (Modderman, 1989). It was later shown to colocalize with other Igps in a variety of cell types (Metzelaar *et al.*, 1991), consistent with its predicted 10-amino acid, cytosolic, carboxyterminal tail ending in the sequence GYEVV. More recently, it has been shown to be enriched on intraluminal vesicles in late endosomes/lysosomes (Escola *et al.*, 1998; Kobayashi *et al.*, 2000; Piper and Luzio, 2001). The kinetics of delivery of newly synthesized CD63 from the TGN to lysosomes has shown a half-time of 2 h, implying that an intracellular delivery route is likely to be important (Barriocanal *et al.*, 1986).

MATERIALS AND METHODS

Reagents

Unless otherwise stated reagents were purchased from Sigma Chemical (Poole, Dorset, United Kingdom). Wortmannin was purchased from Calbiochem (Nottingham, United Kingdom) and was aliquoted at -20°C as a 1 mM stock in dimethyl sulfoxide. Restriction

endonucleases, shrimp alkaline phosphatase, polynucleotide kinase, and DNA polymerases were purchased from New England Biolabs (Hitchin, United Kingdom). Rat CD63 cDNA in pUEX was a gift from Drs. C. Wasmeier and J.C. Hutton (Barbara Davis Center for Childhood Diabetes, University of Colorado, Denver, CO). A human cDNA library from Raji cells (QUICK-Clone cDNA; CLONTECH, Palo Alto, CA) was used as a source of human CD63 cDNA, which was recovered by standard polymerase chain reaction (PCR) techniques. Human CD8 α -chain cDNA in the vector S85 was a gift from Dr. S. Munro (MRC Laboratory of Molecular Biology, Cambridge, United Kingdom), contained an *Afl*III site at the end of the transmembrane region, and was cloned into Δ pMEP4 (Girotti and Banting, 1996) as previously described (Ihrke *et al.*, 2000). Oligonucleotides were from R&D Systems (Abingdon, United Kingdom) and Genosys (Cambridge, United Kingdom).

The species-specific mouse monoclonal antibody (mAb) to rat CD63, designated AD1 (Kitani *et al.*, 1991; Nishikata *et al.*, 1992), was a gift from Dr. R.P. Siraganian (National Institutes of Health, Bethesda, MD). The species-specific mouse mAb to human CD63, designated CLB-gran/12, was purchased from Biotest International (Kennebunk, ME). Fluorescein isothiocyanate-labeled CLB-gran/12 antibody used for fluorescence-activated cell sorting (FACS) analysis was purchased from Immunotech (Marseille, France). Rabbit polyclonal antisera to rat LAMP-2 (lgp110), rat ciMPR, rat TGN38, and mouse mAb to rat TGN38 were as described (Horn and Banting, 1994; Reaves *et al.*, 1996). Rabbit polyclonal anti-rat cathepsin D was a gift from Dr. H.W. Davidson (University of Cambridge, United Kingdom). Rabbit polyclonal anti- γ adaptin and anti- β NAP (β 3B) were gifts from Dr. M.S. Robinson (University of Cambridge). Rat monoclonal antihuman CD8 α (Bindon *et al.*, 1989) was a gift from Dr. G. Hale (University of Oxford, Oxford, United Kingdom). Rabbit polyclonal anti-human cathepsin D was purchased from DAKO (Ely, Cambridgeshire, United Kingdom). The rat mAb to mouse LAMP-1 (1D4B) developed by T. August was from the Developmental Studies Hybridoma Bank (Department of Biological Sciences, University of Iowa, Iowa City, IA). Fluorescein isothiocyanate-labeled goat anti-mouse IgG, Texas Red-labeled goat anti-mouse IgG, and Texas Red-labeled donkey anti-rabbit IgG were obtained from Amersham (Little Chalfont, Buckinghamshire, United Kingdom). Texas Red-X-labeled goat anti-rat IgG was from Molecular Probes (Eugene, OR).

Mocha, *pearl*, and stably transfected *mocha* cells expressing the δ subunit of AP-3 were as described (Peden *et al.*, 2002) and were gifts from Dr. M.S. Robinson (University of Cambridge).

Recombinant DNA Procedures

Standard molecular biology procedures (Sambrook *et al.*, 1989), with the variations previously described (Reaves *et al.*, 1998), were performed unless otherwise stated. Human and rat CD63 molecules containing single amino acid replacements in the cytosolic tail were prepared by mutating cDNA with standard PCR techniques. Details of all primers used are available from the corresponding author on request.

To construct a chimera containing the luminal and transmembrane domains of CD8 and the cytosolic tail of CD63 (CD8-CD63), a corresponding piece of CD63 cDNA was amplified by PCR, introducing an *Afl*III site at the final transmembrane/cytosolic tail boundary. The resulting expressed amino acid sequence after joining the *Afl*III-digested CD8 cDNA and CD63PCR product was . . . TLYCK-RLKSIRSGYEVV, with the underlined 10 residues from CD63 and all other residues from CD8.

The chimera in which the carboxyterminal cytosolic tail of CD63 was replaced by that of TGN38 (CD63-TGN38) was constructed in the same way as the lgp120-TGN38 chimeras described previously (Reaves *et al.*, 1998). A fragment of human CD63 cDNA encoding the entire molecule except the carboxyterminal cytosolic tail, and containing a *Sall* site at the 3' end, was generated by PCR. A piece of TGN38 cDNA encoding the cytosolic tail was amplified by PCR

from Δ pMEP-LTT, one of the Igp120-TGN38 chimeras, so that it contained a *Sall* site at the transmembrane/cytosolic tail boundary. The resulting expressed amino acid sequence after joining the *Sall*-digested CD63 cDNA and TGN38 PCR product was ... AC-CLVDHNRKRKIIAFALEGRKSKVTRRPKASDYQRLNLKL, with the underlined 34 residues from TGN38, the preceding D from the *Sall* cloning site and all other residues from CD63.

All cDNA constructs prepared as described above were ligated into the multiple cloning site in the mammalian expression vector Δ pMEP4 (Girotti and Banting, 1996) and their DNA sequences confirmed by dideoxy chain termination sequencing, with the service provided by the Department of Genetics (University of Cambridge).

Cell Culture and Transfection

Normal rat kidney (NRK), COS-7, NIH-3T3, *pearl*, and *mocha* cells were grown in tissue culture flasks or on glass coverslips as previously described for NRK cells (Reaves *et al.*, 1998). COS-7 cells were transfected by electroporation (Reaves *et al.*, 1998) and *mocha* cells with calcium phosphate coprecipitation (Robinson, 1990), as previously described. NRK cells were transfected using Transfectam reagent (Promega, Southampton, United Kingdom) or FugGENE 6 (Roche Diagnostics, Lewes, East Sussex, United Kingdom), NIH-3T3 fibroblasts with LipofectAMINE (Invitrogen, Carlsbad, CA), and *pearl* cells with Superfect (QIAGEN, Chatsworth, CA), according to the manufacturers' instructions. Selection and growth of stable cell lines and stimulation of protein expression in cells transfected with Δ pMEP4 constructs were as previously described (Reaves *et al.*, 1998). Expression levels of mutant human CD63 in stably transfected NRK cells used for immunofluorescence and fluorescence-activated cell sorting (FACS) analysis were shown by quantitative immunoblotting to be between 80 and 135% of the expression level of wild-type human CD63 in the stably transfected NRK cells.

Immunofluorescence Microscopy

Indirect immunofluorescence microscopy and antibody uptake experiments were carried out as previously described (Reaves *et al.*, 1998).

FACS Analysis

Approximately 10^7 stable transfected NRK cells growing in a 150-cm² tissue culture flask were removed from the culture dish and treated essentially as described by Dell'Angelica *et al.* (1999), except that cells were fixed with 4% paraformaldehyde in phosphate-buffered saline (PBS) for 10 min at 20°C and half was permeabilized with 0.1% Triton X-100 in PBS/1% bovine serum albumin. Intact or permeabilized cells were incubated for 45 min on ice with fluorescein-conjugated CLB-gran12 antibody to human CD63 (10 ng/ml) in 100 μ l PBS/1% bovine serum albumin and subsequently analyzed on a FACScan flow cytometer (BD Biosciences, Franklin Lakes, NJ). Positively fluorescent cells were identified as those with fluorescence intensities >97.5% of nontransfected cells. The ratio of positive cells in intact and permeabilized preparations was used to calculate the proportion of intracellular wild-type or mutant human CD63 in the transfected NRK cells. FACS data from each stably transfected cell line are presented relative to the amount of intracellular CD63 in the cell line expressing the wild-type human protein. This was 36% of total expressed CD63, i.e., at a level similar to that of a transfected chimeric Igp in a previous study where it was shown that Igp trafficking machinery was not saturated (Gough and Fambrough, 1997).

Yeast Two-Hybrid System

The two-hybrid system used to investigate the interactions between AP μ subunits and wild type, and mutated cytosolic domains of CD63 was as previously described (Stephens and Banting, 1998;

Hirst *et al.*, 1999). Constructs encoding μ subunits in the two-hybrid transcriptional activation domain vector pVP16 were as described (Stephens and Banting, 1998). Wild-type (KSIRSGYEVV) and mutant versions of the carboxyterminal cytosolic tail of CD63 were amplified by PCR and subcloned into the two-hybrid DNA-binding domain vector pBTM116. The yeast strain L40, maintained as previously described (Stephens and Banting, 1998), was cotransformed with two plasmids by using a polyethylene glycol-lithium acetate procedure and plated onto selective medium lacking leucine and tryptophan to select for colonies containing both plasmids. After 3–4 d, colonies were incubated in liquid cultures to perform quantitative growth assays. Liquid cultures were set up by inoculating 0.15 OD₆₀₀ units of cells into 2 ml of selective medium lacking leucine, tryptophan, and histidine and were assayed for growth after 0–180 h of incubation at 30°C by measurement of OD₆₀₀. Each time point was assayed in triplicate. Quantitative β -galactosidase assays and determination of inhibition of cell growth with 3-amino-1,2,4-triazole (3-AT) were as previously described (Stephens and Banting, 1998). Growth curves were calculated using SigmaPlot 4.01 (Jandel Scientific, San Rafael, CA).

RESULTS

When cDNAs encoding human CD63, and a chimera of CD8 and the carboxyterminal cytosolic tail of CD63 (CD8-CD63) were transfected into NRK cells, the steady-state localization of expressed protein was essentially indistinguishable from that of endogenous LAMP-2 by fluorescence microscopy (Figure 1, A–D). CD8-CD63 was also observed to colocalize with endogenous rat CD63 (our unpublished data). As a control for CD8-CD63, the localization of expressed CD8 was examined in transfected NRK cells and found to be at the cell surface as expected (Figure 1, E and F). The intracellular localization of CD8-CD63 indicated that the carboxyterminal cytosolic tail of CD63 is sufficient to target the “neutral” plasma membrane reporter protein CD8 to lysosomes. To test whether targeting information is also present in luminal or transmembrane domains, we examined the steady-state localization of CD63-TGN38, a chimera in which the carboxyterminal cytosolic tail of TGN38 replaced that of CD63. This TGN38 tail has been shown to localize a variety of chimeras containing the transmembrane and extracellular domains of neutral plasma membrane proteins to the TGN (Reaves *et al.*, 1998). CD63-TGN38 expressed in transfected NRK cells colocalized with LAMP-2 (Figure 1, G and H) and not with TGN38 (our unpublished data), suggesting that as for LAMP-1 and LAMP-2 (Reaves *et al.*, 1998), the luminal and/or transmembrane domains of CD63 also contain lysosomal targeting information.

To examine the role of the GYXX \emptyset (GYEVM) motif of the carboxyterminal cytosolic tail of CD63 in lysosomal targeting of this protein we carried out alanine scanning mutagenesis of this motif in human CD63. Stably transfected NRK cells expressing the mutant proteins were analyzed by immunofluorescence microscopy to determine the steady-state localization of the different mutated proteins (Figure 2). This showed that mutation of the G, Y, E, or M to A resulted in increased expression on the cell surface, but mutation of V to A had no effect (compare Figure 2, A–J, with wild-type in Figure 1A). Overall, the microscopy showed the order of greatest expression on the cell surface of the mutants was GAEVM > GYEVA > GYAVM > AYEVM > wild-type (GYEVM) = GYEAM. The conservative mutation of Y to F (GFEVM) resulted in increased surface localization of the




Figure 1. Indirect immunofluorescence steady-state localization of wild-type human CD63, CD8, and chimeric proteins in stably transfected NRK cells. NRK cells were transfected with Δ pMEP4 constructs containing cDNA encoding either human CD63 (A and B), CD8 (C and D), CD8-CD63 (E and F), or CD63-TGN38 (G and H). Stably transfected cell lines were treated with $3 \mu\text{M}$ CdCl_2 to induce protein expression and then double labeled with a mouse mAb to either CD63 (A and G) or CD8 (C and E) and a rabbit polyclonal antibody against endogenous LAMP-2 (B, D, F, and H). Bar, $20 \mu\text{m}$.

mutated CD63 (Figure 2, K and L). The effect of mutating E to A (GYAVM) was particularly unexpected in the context of a GYXX \emptyset motif and was therefore further analyzed by FACS analysis, which showed that it resulted in a reduced proportion of the protein being expressed at an intracellular location(s). In contrast, FACS analysis showed that the conservative mutation of M to I (GYEVI) increased intracellular localization (Figure 3), although this was not easily detected by immunofluorescence microscopy (Figure 2, M and N). The data obtained on the localization of alanine scan mutants of human CD63 expressed in NRK cells are not species specific because comparable data were obtained by immunofluorescence microscopy of equivalent alanine scan mutants of rat CD63 expressed in transfected COS-7 cells (our unpublished data). The sequences of the carboxyterminal cytosolic tails of rat and human wild-type CD63 are identical.

The observation that mutation of E to A within the GYEV M motif resulted in greater cell surface localization of




Figure 2. Indirect immunofluorescence steady-state localization of mutated human CD63-LIMP-1 in stably transfected NRK cells. NRK cells were transfected with Δ pMEP4 constructs containing cDNA encoding human CD63 with point mutations in the cytosolic tail, either AYEVM (A and B), GAEVM (C and D), GYAVM (E and F), GYEAM (G and H), GYEVA (I and J), GYEVI (K and L), or GFEVM (M and N). Stably transfected cell lines were treated with $3 \mu\text{M}$ CdCl_2 to induce protein expression and then double labeled with a mouse mAb to CD63 (A, C, E, G, I, K, and M) and a rabbit polyclonal antibody against endogenous LAMP-2 (B, D, F, H, J, L, and N). Bar, $20 \mu\text{m}$.




Figure 3. Fluorescence-activated cell sorting analysis of human CD63-LIMP-I and mutated proteins in stably transfected NRK cells. NRK cells were transfected with Δ pMEP4 constructs containing cDNA encoding human CD63 and constructs with point mutations in the cytosolic tail. After induction of protein expression the distribution of the wild-type and mutated human CD63 proteins between the cell surface and intracellular structures was determined by FACS analysis. (A) Fluorescence intensity for untransfected NRK cells (top left) and cells transfected with wild-type CD63 (GYEVM), and mutated constructs (GYEVI, GAEVM, and GYAVM) are shown. The filled traces represent intact cells and the unfilled traces the permeabilized cells. A larger shift to the right of the unfilled trace compared with the filled trace represents a larger proportion of the CD63 at intracellular sites. (B) Histogram showing the amount of intracellular mutated CD63 in each cell line relative to the amount of intracellular wild-type CD63 in the cell line expressing this construct.

CD63 was initially surprising given the consensus sequence for the motif as GYXX \emptyset . However, data obtained by screening the interaction of adaptor μ subunits with a combinatorial XXXYXX \emptyset library, by using the yeast two-hybrid system, suggested that an E at the Y + 1 position results in a YXX \emptyset motif with preferential binding to μ 3 (Ohno *et al.*, 1998). We therefore decided to investigate the interaction of wild-type and mutant carboxyterminal cytosolic tails of CD63 with μ subunits in the yeast two-hybrid system. The μ subunits were cloned into the two-hybrid transcriptional

activation domain vector pVP16, and wild-type and mutated carboxyterminal cytosolic tails of CD63 were cloned into the two-hybrid DNA-binding domain vector pBTM116. Pairs of constructs were then transformed into yeast cells, and interaction between the cytosolic tails and μ subunits assayed by the ability of cells coexpressing them 1) to grow on agar or in liquid medium lacking histidine; 2) to induce β -galactosidase expression; or, for strong interactors, 3) to grow in the presence of 3-AT, a competitive inhibitor of histidine biosynthesis. By using this two-hybrid system and measuring interaction by growth in medium lacking histidine, it has previously been shown that the wild-type carboxyterminal cytosolic tail of CD63, containing GYEVM, interacts most strongly with μ 3A, weakly with μ 2 and μ 4, and not at all with μ 1 (Hirst *et al.*, 1999). There was an equally strong interaction with the brain-specific isoform μ 3B (our unpublished data). The interaction of mutated CD63 tails with μ 3A and μ 2 was shown by growth on agar lacking histidine (Figure 4). Clearly, the tails containing Y-to-A and M-to-A mutations interacted very poorly with μ 3A because no growth was observed. Interactions with μ 2 occurred only with three constructs (GYEVM, GYAVM, and GYEAM) and were all weak, resulting in poor growth on the agar (Figure 4). To characterize the relative strengths of interaction of the mutated tails with μ 3A in more detail, growth of cotransformed yeast was measured in medium lacking histidine >180 h (Figure 5, A and B). The strengths of interaction of the weakly interacting mutated tails with μ 3A were easily distinguished by this method. The relative strengths of interaction of the strongly interacting wild-type and mutated tails were distinguished better by plotting dose-response curves of 3-AT inhibition of growth of cotransformed yeast in medium lacking histidine (Figure 5C) or measurement of β -galactosidase activity after growth in medium containing histidine (Figure 5D). Taking these data together, the strength of interaction of wild-type and mutated tails of CD63 with μ 3A was determined to be wild-type (GYEVM) = GYEAM > AYEVM > GYAVM \gg GYEVA \gg GAEVM. This order is the exact reverse of that describing cell surface expression of CD63 molecules containing these mutated tails in transfected NRK cells (Figures 2 and 3).

In the yeast two-hybrid experiments we also examined the strength of interaction of μ 3A with a mutated CD63 tail in which M was mutated to I (GYEVI). This construct was chosen because the data from the combinatorial XXXYXX \emptyset library presented by Ohno *et al.* (1998) showed that for interaction with μ 3A, I is favored in the Y + 3 position. We found that the mutated CD63 tail in which M was mutated to I showed the strongest interaction with μ 3A of any tail sequence we examined (Figure 5, A, C, and D) and CD63 containing this mutation had the lowest surface expression in transfected NRK cells (Figures 2K and 3). These data were not due to the creation of a dileucine-type motif of the form VI because CD63 with the mutated tail GAEVI was expressed entirely on the cell surface (our unpublished data). We also observed in the yeast two-hybrid system that the mutated CD63 tail in which M was mutated to I showed no interaction with μ 2 (Figure 4). This implied that any CD63 containing this mutation delivered to the cell surface in transfected cells would be poorly internalized by clathrin-mediated endocytosis. To test this, we investigated anti-




Figure 4. Interaction of the cytosolic domains of CD63 and mutated CD63 with $\mu 2$ and $\mu 3A$ in the yeast two-hybrid system. Yeast cells were transformed with bait constructs (LexA fusion) containing the C-terminal cytosolic domains of CD63, and the mutant forms AYEVM, GAEVM, GYAVM, GYEVA, GYAVM, GYEAM, GYEVA, and GYEVI and a prey construct (either VP16- $\mu 2$, VP16- $\mu 3A$). Transformed yeast were grown on agar plates in the absence of histidine for 3 d ($\mu 3A$) or 10 d ($\mu 2$). All transformed yeast grew in the presence of histidine and no transformed yeast containing "empty" VP16 grew in the absence of histidine.

human CD63 antibody uptake in stable lines of NRK cells transfected with wild-type (GYEVM) and mutated (GYEVI) CD63. Less intracellular accumulation of antibody was observed in cells expressing the mutated (GYEVI) CD63, consistent with inefficient clathrin-mediated uptake (Figure 6).

The experiments described above, together with published data on the kinetics of delivery of wild-type CD63 from TGN to lysosomes, strongly suggest that the wild-type protein traffics to lysosomes from the TGN by an intracellular AP-3-dependent route, but that any CD63 reaching the cell surface is internalized by AP-2-dependent clathrin-mediated endocytosis and delivered by the endocytic pathway to lysosomes. In contrast, mutated (GYEVI) CD63 should only reach lysosomes by an intracellular AP-3-dependent route and any delivered to the cell surface should not be efficiently internalized and delivered by endocytosis. To test this hypothesis we expressed human wild-type and mutated (GYEVI) CD63 in cells from *pearl* and *mocha* mice and, using immunofluorescence microscopy, examined the steady-state distribution of the recombinant expressed protein. As a con-

trol we also transfected wild-type mouse NIH-3T3 fibroblasts. Consistent with our hypothesis, mutated (GYEVI) CD63 was observed almost exclusively on the surface of *pearl* and *mocha* cells but colocalized with an endogenous lgp (LAMP-1) in NIH-3T3 fibroblasts (Figures 7 and 8). In contrast, wild-type (GYEVM) CD63 colocalized with endogenous LAMP-1 in transfected *pearl* and *mocha* cells (Figures 7 and 8). Similar results were obtained from both transiently transfected cells and stable transfected cell lines. To test our hypothesis further, two additional experiments were carried out. In the first, we transfected *mocha* cells that had been rescued by expression of the δ subunit of AP-3 with mutated (GYEVI) CD63. In the rescued cells expressing the δ subunit of AP-3 at a concentration sufficient for detection by immunofluorescence, a significant proportion of the expressed mutated (GYEVI) CD63 colocalized with intracellular, endogenous LAMP-1 (Figure 9). In the second experiment, we further transfected a stable line of *pearl* cells expressing mutated (GYEVI) CD63 with a cDNA construct encoding the $\beta 3B$ subunit of AP-3. Some intracellular localization of mu-




Figure 5. Characterization of the interaction of the cytosolic domains of CD63 and mutated CD63 with $\mu 3A$ in the yeast two-hybrid system. Yeast cells were transformed as described in Figure 4. (A and B) Cultures were set up containing 0.05 OD₆₀₀ units of transfected cells into 2 ml of selective medium lacking histidine and grown at 30°C with shaking. OD₆₀₀ measurements were made up to 180 h (B shows an expanded time scale). The control interaction between the C-terminal cytosolic domain of wild-type CD63 and empty VP16 (filled circles) is shown in each figure. Vector controls for other tails also showed no significant growth. Each point represents the mean \pm SEM of the activity of three separate cultures. (C) Cultures of transformed yeast were established in selective medium lacking histidine, but containing 0–1 M 3-AT, and grown at 30°C with shaking. After incubation for 48 h, OD₆₀₀ was measured and the ratio in the presence versus absence of 3-AT recorded. Each point represents the mean \pm SEM of the activity of three separate cultures. (D) Cultures of transformed yeast were grown in medium containing histidine to an OD₆₀₀ of ~ 1.2 . The β -galactosidase activity of the cultures was then measured. Each bar in the histogram represents the mean \pm SEM of the activity of five separate cultures.




Figure 6. Uptake of anti-CD63 antibody by stably transfected NRK lines expressing either wild-type human CD63 (A and B) or the GYEVI construct (C and D). Protein expression was induced in stably transfected NRK cell lines with $3 \mu\text{M}$ CdCl_2 for 16 h. The cells were then incubated at 37°C for 30 min in the presence of anti-CD63 mAb and then incubated for a further 30 min without antibody (i.e., chased). The cells were then fixed and immunofluorescence localization of the anti-CD63 mAb (A and C) and of endogenous LAMP-2 (B and D) carried out. Bar, $20 \mu\text{m}$.

tated (GYEVI) CD63 was observed in cells expressing $\beta 3\text{B}$ (our unpublished data; $\beta 3\text{B}$ was used in this experiment because of the lack of antibodies to $\beta 3\text{A}$ for immunofluorescence).

The data presented above are consistent with the mutated (GYEVI) CD63 absolutely requiring functional AP-3 for targeting to lysosomes and provided us with a unique reporter protein for this pathway. As discussed in INTRODUCTION it has been reported that AP-3 is associated with endosomes as well as with the TGN (Simpson *et al.*, 1996; Dell'Angelica *et al.*, 1998). Because chloroquine treatment of cells causes swelling of endocytic compartments and blocks traffic through the early endosome (Reaves *et al.*, 1998) we predicted that if mutated (GYEVI) CD63 was trafficking via this compartment it should be trapped there after chloroquine treatment along with TGN38, which normally cycles between the TGN and cell surface/early endosome (Chapman and Munro, 1994; Reaves and Banting, 1994; Banting *et al.*, 1998). However, no mutated (GYEVI) CD63 was observed in the same structures as TGN38 in chloroquine-treated, transfected NRK cells (Figure 10, A and B). In contrast, when CD63-TGN38-expressing NRK cells were treated with chloroquine, CD63-TGN38 was observed in the same swollen structures as endogenous TGN38 (Figure 10, C and D) consistent with this molecule trafficking via an early endosome, i.e., by a different route to the AP-3-dependent route used by mutated (GYEVI) CD63. Other swollen structures depleted of TGN38 also contained CD63-TGN38, consistent with them being late endosomes and lysosomes containing CD63-TGN38 molecules that had arrived before the chloroquine was added. When NRK cells expressing wild-type human CD63 were incubated with chloroquine, this protein was mainly in separate structures to TGN38, although some colocalization was observed that presumably represents the small proportion of wild-type CD63 that traffics via the early endosome (Figure 10, E and F). These data taken together suggest that the AP-3-dependent pathway of membrane




Figure 7. Steady-state localization of wild-type human CD63 and the GYEVI construct in transiently transfected *pearl* cells. *Pearl* cells were transiently transfected with ΔpMEP containing cDNA encoding either wild-type CD63 (A and B) or the GYEVI construct (C and D). The transfected cells were treated with $3 \mu\text{M}$ CdCl_2 to induce protein expression and then double labeled, for indirect immunofluorescence localization, with a mouse mAb to CD63 (A and D) and a rat mAb to endogenous mouse LAMP-1 (B and E). Bar, $20 \mu\text{m}$.

traffic from the TGN to lysosomes does not involve traffic via the early endosomal compartment in which TGN38 accumulates after chloroquine treatment.

DISCUSSION

The increased surface expression of CD63 in which G, Y, or M was mutated to A confirmed the importance of the GYEVV sequence in the localization of this protein, as is the case for the GYXX \emptyset motif in the carboxyterminal cytoplasmic tail of other lgps, including lysosomal acid phosphatase (Peters *et al.*, 1990; Lehmann *et al.*, 1992), LAMP-1 (Williams and Fukuda, 1990; Harter and Mellman, 1992; Guarnieri *et al.*, 1993), and LAMP-2 isoforms (Gough and Fambrough, 1997).

The observation that mutating E to A in the carboxyterminal cytosolic tail of CD63 produced a mutant protein that was more expressed at the cell surface than wild-type protein was unexpected because mutations at the Y + 1 position of the GYXX \emptyset motif have been shown to have no effect on lysosomal targeting of LAMP-1 (Guarnieri *et al.*, 1993; Honing and Hunziker, 1995; Rohrer *et al.*, 1996). It stimulated our use of the yeast two-hybrid system to investigate the interaction of wild-type and mutated cytosolic tails of CD63 with adaptor medium chain (μ) subunits. From their screen of the interactions with μ subunits of a combinatorial peptide li-




Figure 8. Steady-state localization of CD63 and GYEVI constructs in stably transfected *mocha* and 3T3 cell lines. *Mocha* cells (A–D) and 3T3 cells (E–H) were transfected with Δ pMEP constructs containing cDNA encoding wild-type human CD63 (A, B, E, and F) or the GYEVI construct (C, D, G, and H). Stably transfected cell lines were treated with 3 μ M CdCl₂ to induce protein expression and then double labeled, for indirect immunofluorescence localization, with a mouse mAb to CD63 (A, C, E, and G) and a rat mAb to endogenous mouse LAMP-1 (B, D, F, and H). Bar, 20 μ m.

brary, Ohno *et al.* (1998) observed a preference for E at the Y + 1 position of a YXX \emptyset motif for interaction with μ 3A and μ 3B. They pointed out that this was “noteworthy because it is a characteristic of proteins targeted to lysosomes or lysosome-related organelles, such as . . . CD63.” Their data also showed a preference for G rather than A at the Y – 1 position and I rather than M at the Y + 3 position for interaction with μ 3A. Our own yeast two-hybrid data are consistent with the data from Ohno *et al.* (1998), but in our case within the context of the actual cytosolic tail of an lgp. More importantly, we have shown for the first time that the order of cell surface versus lysosomal distribution of GYXX \emptyset mutants of an lgp is accurately predicted by interactions with μ 3A in the yeast two-hybrid system. Two other complete cytosolic tails containing YXX \emptyset motifs have previously been shown to be capable of interacting with adaptor μ chains in the yeast two-hybrid system: those of TGN38




Figure 9. Localization of the GYEVI construct of human CD63 in rescued *mocha* cells expressing the δ subunit of AP-3. A stably transfected *mocha* cell line expressing the δ subunit of AP-3 was transiently transfected with a Δ pMEP construct containing cDNA encoding the human CD63/LIMP-I, GYEVI construct. After treating the transfected cells with 3 μ M CdCl₂ to induce protein expression they were double labeled, for indirect immunofluorescence localization with a mouse mAb to human CD63 (A) and with a rabbit polyclonal antibody to the δ subunit of AP-3 (B). Bar, 20 μ m. (C) Cells that were expressing the GYEVI construct were then counted and grouped according to the presence of intracellular human CD63 or only cell surface human CD63 and the presence or absence of detectable δ subunit of AP-3. The data shown in the histogram represent the mean of two experiments. In the two experiments the counted numbers of human CD63-positive cells showing only surface expression were 208 and 349. The counted numbers of CD63-positive cells also showing intracellular localization of human CD63 were 91 and 44.

and LAMP-1 (Ohno *et al.*, 1995; Stephens *et al.*, 1997; Stephens and Banting, 1998). In both these cases, and for the tail of CD63, interaction is dependent on the Y and the \emptyset in the YXX \emptyset motif.

In our yeast two-hybrid experiments we observed that a mutated CD63 tail in which M was mutated to I (GYEVI) did not interact with μ 2, despite showing the strongest interaction with μ 3A of all the mutated tails examined. Consistent with this observation, examination of the combinatorial XXXYXX \emptyset peptide library screen of Ohno *et al.* (1998) reveals that for interaction with μ 2, I is more disfavored than M at the Y + 3 position. The availability of a mutated CD63 that interacts strongly with μ 3A but not with μ 2 allowed us to test a number of predictions about its behavior, and the role of the AP-3-dependent trafficking pathway, when it was expressed in cultured cells. In particular, we predicted and showed that, in cells lacking functional AP-3, the GYEVI mutant of CD63 was unable to reach lysosomes because it was denied the AP-3-dependent intracellular route and could not be internalized from the cell surface. This route was recovered and the GYEVI mutant correctly delivered to lysosomes in *mocha* cells, which had been rescued by expression of the δ subunit of AP-3. In contrast, wild-type CD63




Figure 10. Indirect immunofluorescence steady-state localization of wild-type human CD63, GYEVI, and CD63-TGN38 constructs in stable lines of transfected NRK cells treated with chloroquine. Stably transfected NRK cell lines expressing either the GYEVI construct (A and B), CD63-TGN38 (C and D), or wild-type CD63 (E and F) were incubated for 2 h in the presence of both 3 μ M CdCl₂ and 100 μ M chloroquine before fixation. The cells were then double labeled, for indirect immunofluorescence localization, with a mouse mAb to human CD63 (A, C, and E) and a rabbit polyclonal antibody to the luminal domain of rat TGN38 (B, D, and F). Structures positive for TGN38, but which do not label with the anti-CD63 antibody, are indicated by single-headed arrows. Structures that label with both the TGN38 antibody and the CD63 antibody are indicated by the double-headed arrows. Bar, 10 μ m.

was still delivered to lysosomes in transfected *pearl* and *mocha* cells, consistent with the ability to be internalized efficiently from the cell surface. Our experiments have identified the first Igp, the GYEVI mutant of CD63, to be grossly

mislocalized in AP-3-deficient cells. They are also consistent with previous experiments where endogenous Igps were observed to traffic to lysosomes via the cell surface in AP-3-deficient cells (Dell'Angelica *et al.*, 1999) or in cells in which functional AP-3 concentrations were reduced by transfecting with antisense oligonucleotides to μ 3A (Le Borgne *et al.*, 1998).

The intracellular site of action of AP-3 in lysosomal targeting has not been well defined. It is unlikely that AP-3 acts at the plasma membrane because it would be difficult to reconcile such a site of action with the observation that increased quantities of Igps traffic via the cell surface in cells deficient in AP-3 (Le Borgne *et al.*, 1998; Dell'Angelica *et al.*, 1999). Moreover, it has been localized by microscopy at or close to the TGN (Simpson *et al.*, 1996) and on endosomes (Dell'Angelica *et al.*, 1998). It thus seems likely that AP-3 functions to traffic between the TGN and lysosomes on a route equivalent to the AP-3-mediated direct trafficking pathway from the Golgi complex to the vacuole proposed in yeast and used by alkaline phosphatase but not carboxypeptidase Y (Cowles *et al.*, 1997). In the present experiments we studied the effects of chloroquine on the distribution of transfected wild-type CD63 and a chimera with the cytoplasmic tail of TGN38 to provide further insights into the intracellular site of action of AP-3. Previous work has shown that endosomal acidification is important for transport from endosomal compartments to both lysosomes and the TGN (Chapman and Munro, 1994; Reaves and Banting, 1994; Van Weert *et al.*, 1995), such that when the pH of endosomes is raised using either a proton pump inhibitor (e.g., bafilomycin A1) or a membrane-permeant weak base (e.g., chloroquine), membrane proteins, including TGN38, furin, LAMP-1, and lysosomal acid phosphatase become trapped in early endosomal compartments (Braun *et al.*, 1989; Chapman and Munro, 1994; Reaves and Banting, 1994; Reaves *et al.*, 1998). In addition, delivery of endocytosed horseradish peroxidase from endosomes to lysosomes is impaired (Van Weert *et al.*, 1995). Thus, in the present study, chloroquine was used to inhibit traffic through early endosomal compartments and to see whether wild-type CD63 or the GYEVI mutant were trapped in early endosomes en route to the lysosomes. Some wild-type CD63 was trapped in early endosomes, suggesting that a proportion traffics through this compartment. In contrast, none of the GYEVI mutant was found in early endosomes, consistent with it trafficking from the TGN to lysosomes by an AP-3-dependent intracellular route that does not involve passage through early endosomes. As a control, we studied the CD63-TGN38 chimera that has a TGN38 cytosolic tail that interacts strongly with μ 2, weakly with μ 1, and barely detectably with μ 3 in the yeast two-hybrid system (Ohno *et al.*, 1995; Stephens and Banting, 1998; our unpublished data). CD63-TGN38 was found in endosomal structures when chloroquine was added to the cells, suggesting that, as predicted, it does indeed traffic through early endosomes.

Why does wild-type CD63 have a GYEV motif rather than a GYEVI motif if the latter is more efficient for AP-3-dependent delivery to the lysosome? One possible explanation is that because of the poor interaction of GYEVI with μ 2 any CD63 with a GYEVI motif that is mistargeted to the cell surface would remain there; this may be detrimental to the cell and therefore unfavorable evolutionarily. Another pos-

sible explanation relates to the possible functions of CD63 at the cell surface, including involvement in cell adhesion (Vischer and Wagner, 1993; Hamamoto *et al.*, 1994) and integrin-mediated cell migration (Berditchevski and Odintsova, 1999). Alterations in the amount of CD63 expressed in melanomas have been implicated in metastatic progression of the tumor, and this may reflect altered cell surface expression (Hara *et al.*, 1994; Radford *et al.*, 1997), suggesting that the amount of CD63 on the cell surface may be capable of modulation. Because the distribution of the wild-type molecule is clearly dependent on interactions of the carboxyterminal cytosolic tail with more than one adaptor protein, regulation of such interactions would provide a means of varying the cell surface concentration and may be a fruitful subject of future investigation.

ACKNOWLEDGMENTS

We thank our colleagues Margaret Robinson, Howard Davidson, Jenny Hirst, Andrew Peden, and Rachel Rudge for AP reagents, cell lines, much valuable discussion, and critical reading of the manuscript. We thank Sean Munro for the gift of a plasmid construct encoding CD8. This work was funded by the Medical Research Council and the Wellcome Trust.

REFERENCES

- Asakaki, K., Michihara, A., Mibuka, K., Fujiwara, Y., and Tsuji, H. (1995). Biosynthetic transport of a major lysosomal membrane glycoprotein, lamp-1: Convergence of biosynthetic and endocytic pathways occurs at three distinctive points. *Exp. Cell Res.* **220**, 464–473.
- Asakaki, K., Michihara, A., Fujiwara, Y., Mibuka, K., and Tsuji, H. (1996). Biosynthetic transport of a major lysosome-associated membrane glycoprotein 2, lamp-2: a significant fraction of newly synthesized lamp-2 is delivered to lysosomes by way of early endosomes. *J. Biochem.* **120**, 1088–1094.
- Andrejewski, N., Punnonen, E.-L., Guhde, G., Tanaka, Y., Lüllmann-Rauch, R., Hartmann, D., von Figura, K., and Saftig, P. (1999). Normal lysosomal morphology and function in LAMP-1-deficient mice. *J. Biol. Chem.* **274**, 12692–12701.
- Banting, G., Maile, R., and Roquemore, E.P. (1998). The steady state distribution of humTGN46 is not significantly altered in cells defective in clathrin-mediated endocytosis. *J. Cell Sci.* **111**, 3451–3458.
- Barriocanal, J.G., Bonifacino, J.S., Yuan, L., and Sandoval, I.V. (1986). Biosynthesis, glycosylation, movement through the Golgi system, and transport to lysosomes by an N-linked carbohydrate-independent mechanism of three lysosomal integral membrane proteins. *J. Biol. Chem.* **261**, 16755–16763.
- Berditchevski, F., and Odintsova, E. (1999). Characterization of integrin-tetraspanin adhesion complexes. Role of tetraspanins in integrin signaling. *J. Cell Biol.* **146**, 477–492.
- Bindon, C.L., Hale, G., and Waldmann, H. (1989). Synergistic complement-mediated cell lysis using pairs of antibodies. In: *Leukocyte Typing IV*, ed. B. Knapp, W.R. Dorken, E.P. Gilks, R.E. Rieber, H. Schmidt, H. Stein, and A.E.G.K. von dem Borne, Oxford, UK: Oxford University Press, 349–350.
- Bonifacino, J.S., and Dell'Angelica, E.C. (1999). Molecular bases for the recognition of tyrosine-based sorting signals. *J. Cell Biol.* **145**, 923–926.
- Braun, M., Waheed, A., and Von Figura, K. (1989). Lysosomal acid phosphatase is transported to lysosomes via the cell surface. *EMBO J.* **8**, 3633–3640.
- Carlsson, S.R., and Fukuda, M. (1992). The lysosomal membrane glycoprotein lamp-1 is transported to lysosomes by two alternative pathways. *Arch. Biochem. Biophys.* **296**, 630–639.
- Chapman, R.E., and Munro, S. (1994). Retrieval of TGN proteins from the cell surface requires endosomal acidification. *EMBO J.* **13**, 2305–2312.
- Cowles, C.R., Odorizzi, G., Payne, G.S., and Emr, S.D. (1997). The AP-3 adaptor complex is essential for cargo-selective transport to the yeast vacuole. *Cell* **91**, 109–118.
- Dell'Angelica, E.C., Klumperman, J., Stoorvogel, W., and Bonifacino, J.S. (1998). Association of the AP-3 adaptor complex with clathrin. *Science* **280**, 431–434.
- Dell'Angelica, E.C., Shotelersuk, V., Aguilar, R.C., Gahl, W.A., and Bonifacino, J.S. (1999). Altered trafficking of lysosomal proteins in Hermansky-Pudlak syndrome due to mutations in the β 3A subunit of the AP-3 adaptor. *Mol. Cell* **3**, 11–21.
- D'Souza, M.P., and August, J.T. (1986). A kinetic analysis of biosynthesis and localization of a lysosome-associated membrane glycoprotein. *Arch. Biochem. Biophys.* **249**, 522–532.
- Escola, J.M., Kleijmeer, M.J., Stoorvogel, W., Griffin, J.M., Yoshie, O., and Geuze, H.J. (1998). Selective enrichment of tetraspan proteins on the internal vesicles of multivesicular endosomes and on exosomes secreted by human B-lymphocytes. *J. Biol. Chem.* **273**, 20121–20127.
- Girotti, M., and Banting, G. (1996). TGN38-green fluorescent protein hybrid proteins expressed in stably transfected eukaryotic cells provide a tool for the real-time, in vivo study of membrane traffic pathways and suggest a possible role for rat TGN38. *J. Cell Sci.* **109**, 2915–2926.
- Gough, N.R., and Fambrough, D.M. (1997). Different steady state subcellular distributions of the three splice variants of lysosome-associated membrane protein LAMP-2 are determined largely by the COOH-terminal amino acid residue. *J. Cell Biol.* **137**, 1161–1169.
- Gough, N.R., Zweifwel, M.E., Martinez-Augustin, O., Aguilar, R.C., Bonifacino, J.S., and Fambrough, D.M. (1999). Utilization of the indirect lysosome targeting pathway by lysosome-associated membrane proteins (LAMPs) is influenced largely by the C-terminal residue of their GYXX Φ targeting signals. *J. Cell Sci.* **112**, 4257–4269.
- Green, S.A., Zimmer, K.P., Griffiths, G., and Mellman, I. (1987). Kinetics of intracellular transport and sorting of lysosomal membrane and plasma membrane proteins. *J. Cell Biol.* **105**, 1227–1240.
- Gu, F., and Gruenberg, J. (1999). Biogenesis of transport intermediates in the endocytic pathway. *FEBS Lett.* **452**, 61–66.
- Guarnieri, F.G., Arterburn, L.M., Penno, M.B., Cha, Y., and August, J.T. (1993). The motif Tyr-X-X-hydrophobic residue mediates lysosomal membrane targeting of lysosome-associated membrane protein 1. *J. Biol. Chem.* **268**, 1941–1946.
- Hamamoto, K., Ohga, S., Nomura, S., and Yasunaga, K. (1994). Cellular distribution of CD63 antigen in platelets and in three megakaryocytic cell lines. *Histochem. J.* **26**, 367–375.
- Hara, H., M.H. Lee, H. Chen, D. Luo, D., and K. Jimbow. (1994). Role of gene expression and protein synthesis of tyrosinase, TRP-1, Lamp-1, and CD63 in UVB-induced melanogenesis in human melanomas. *J. Invest. Dermatol.* **102**, 495–500.
- Harter, C., and Mellman, I. (1992). Transport of the lysosomal membrane glycoprotein lgp120 (lgp-A) to lysosomes does not require appearance on the plasma membrane. *J. Cell Biol.* **117**, 311–325.
- Hirst, J., Bright, N.A., Rous, B., and Robinson, M.S. (1999). Characterization of a fourth adaptor-related protein complex. *Mol. Biol. Cell* **10**, 2787–2802.
- Hirst, J., and Robinson, M.S. (1998). Clathrin and adaptors. *Biochim. Biophys. Acta* **1404**, 173–193.

- Honing, S., Griffith, J., Geuze, H.J., and Hunziker, W. (1996). The tyrosine-based lysosomal targeting signal in lamp-1 mediates sorting into Golgi-derived clathrin-coated vesicles. *EMBO J.* 15, 5230–5239.
- Honing, S., and Hunziker, W. (1995). Cytoplasmic determinants involved in direct lysosomal sorting, endocytosis, and basolateral targeting of rat Igp120 (lamp-I) in MDCK cells. *J. Cell Biol.* 128, 321–332.
- Horn, M., and Banting, G. (1994). Okadaic acid treatment leads to a fragmentation of the trans-Golgi network and an increase in expression of TGN38 at the cell surface. *Biochem. J.* 301, 69–73.
- Huizing, M., Sarangarajan, R., Strovel, E., Zhao, Y., Gahl, W.A., and Boissy, R.E. (2001). AP-3 mediates tyrosinase but not TRP-1 trafficking in human melanocytes. *Mol. Biol. Cell* 12, 2075–2085.
- Hunziker, W., and Geuze, H.J. (1996). Intracellular trafficking of lysosomal membrane proteins. *Bioessays* 18, 379–389.
- Ihrke, G., Gray, S.R., and Luzio, J.P. (2000). Endolyn is a mucin-like type I membrane protein targeted to lysosomes by its cytoplasmic tail. *Biochem. J.* 345, 287–296.
- Karlsson, K., and Carlsson, S.R. (1998). Sorting of lysosomal membrane glycoproteins lamp-1 and lamp-2 into vesicles distinct from mannose 6-phosphate receptor/ γ -adaptin vesicles at the trans-Golgi network. *J. Biol. Chem.* 273, 18966–18973.
- Kitani, S., Berenstein, E., Mergenhagen, S., Tempst, P., and Siraganian, R.P. (1991). A cell surface glycoprotein of rat basophilic leukemia cells close to the high affinity IgE receptor (Fc ϵ RI). Similarity to human melanoma differentiation antigen ME491. *J. Biol. Chem.* 266, 1903–1909.
- Kobayashi T., Vischer, U.M., Rosonoblet, C., Lebrand, C., Lindsay, M., Parton, R.G., Kruihof, E.K., and Gruenberg, J. (2000). The tetraspanin CD63/lamp3 Cycles between endocytic and secretory compartments in human endothelial cells. *Mol. Biol. Cell* 11, 1829–1843.
- Le Borgne, R., Alconada, A., Bauer, U., and Hoflack, B. (1998). The mammalian AP-3 adaptor-like complex mediates the intracellular transport of lysosomal membrane glycoproteins. *J. Biol. Chem.* 273, 29451–29461.
- Lehmann, L.E., Eberle, W., Krull, S., Prill, V., Schmidt, B., Sander, C., Von Figura, K., and Peters, C. (1992). The internalization signal in the cytoplasmic tail of lysosomal acid phosphatase consists of the hexapeptide PGYRHHV. *EMBO J.* 11, 4391–4399.
- Luzio, J.P., Rous, B.A., Bright, N.A., Pryor, P.R., Mullock, B.M., and Piper, R.C. (2000). Lysosome-endosome fusion and lysosome biogenesis. *J. Cell Sci.* 113, 1515–1524.
- Maecker, H.T., Todd, S.C., and Levy, S. (1997). The tetraspanin superfamily: molecular facilitators. *FASEB J.* 11, 428–442.
- Metzelaar, M.J., Wijngaard, P.L., Peters, P.J., Sixma, J.J., Nieuwenhuis, H.K., and Clevers, H.C. (1991). CD63 antigen. A novel lysosomal membrane glycoprotein, cloned by a screening procedure for intracellular antigens in eukaryotic cells. *J. Biol. Chem.* 266, 3239–3245.
- Meyer, C., Zizioli, D., Lausmann, S., Eskelinen, E.-L., Hamann, J., Saftig, P., von Figura, K., and Schu, P. (2000). μ 1A-adaptin-deficient mice: lethality, loss of AP-1 binding and rerouting of mannose 6-phosphate receptors. *EMBO J.* 19, 2193–2203.
- Modderman, P.W. (1989). Cluster report: CD63. In: *Leukocyte Typing IV*, ed. B. Knapp, W.R. Dorken, E.P. Gilks, R.E. Rieber, H. Schmidt, H. Stein, and A.E.G.K. von dem Borne, Oxford, UK: Oxford University Press, 1042.
- Mullins, C., and Bonifacino, J.S. (2001). The molecular machinery for lysosome biogenesis. *BioEssays* 23, 333–343.
- Nishikata, H., Oliver, C., Mergenhagen, S.E., and Siraganian, R.P. (1992). The rat mast cell antigen AD1 (homologue to human CD63 or melanoma antigen ME491) is expressed in other cells in culture. *J. Immunol* 149, 862–870.
- Ohno, H., Aguilar, R.C., Yeh, D., Taura, D., Saito, T., and Bonifacino, J.S. (1998). The medium subunits of adaptor complexes recognize distinct but overlapping sets of tyrosine-based sorting signals. *J. Biol. Chem.* 273, 25915–25921.
- Ohno, H., Stewart, J., Fournier, M.C., Bosshart, H., Rhee, I., Miyatake, S., Saito, T., Gallusser, A., Kirchhausen, T., and Bonifacino, J.S. (1995). Interaction of tyrosine-based sorting signals with clathrin-associated proteins. *Science* 269, 1872–1875.
- Peden, A.A., Rudge, R.E., Lui, W.W.Y., and Robinson, M.S. (2002). Assembly and function of AP-3 complexes in cells expressing mutant subunits. *J. Cell Biol.* (*in press*).
- Peters, C., Braun, M., Weber, B., Wendland, M., Schmidt, B., Pohlmann, R., Waheed, A., and Von Figura, K. (1990). Targeting of a lysosomal membrane protein: a tyrosine-containing endocytosis signal in the cytoplasmic tail of lysosomal acid phosphatase is necessary and sufficient for targeting to lysosomes. *EMBO J.* 9, 3497–3506.
- Piper, R.C., Bryant, N.J., and Stevens, T.H. (1997). The membrane protein alkaline phosphatase is delivered to the vacuole by a route that is distinct from the VPS-dependent pathway. *J. Cell Biol.* 138, 531–545.
- Piper, R.C., and Luzio, J.P. (2001). Late endosomes: sorting and partitioning in multivesicular bodies. *Traffic* 2, 612–621.
- Puertollano, R., Aguilar, R.C., Gorshkova, I., Crouch, R.J., and Bonifacino, J.S. (2001). Sorting of mannose 6-phosphate receptors mediated by the GGAs. *Science* 292, 1712–1716.
- Radford, K.J., Thorne, R.F., and Hersey, P. (1997). Regulation of tumor cell motility and migration by CD63 in a human melanoma cell line. *J. Immunol.* 158, 3353–3358.
- Reaves, B., and Banting, G. (1994). Vacuolar ATPase inactivation blocks recycling to the trans-Golgi network from the plasma membrane. *FEBS Lett.* 345, 61–66.
- Reaves, B.J., Banting, G., and Luzio, J.P. (1998). Lumenal and transmembrane domains play a role in sorting type I membrane proteins on endocytic pathways. *Mol. Biol. Cell* 9, 1107–1122.
- Reaves, B.J., Bright, N.A., Mullock, B.M., and Luzio, J.P. (1996). The effect of wortmannin on the localization of lysosomal type I integral membrane glycoproteins suggests a role for phosphoinositide 3-kinase activity in regulating membrane traffic late in the endocytic pathway. *J. Cell Sci.* 109, 749–762.
- Robinson, M.S. (1990). Cloning and expression of gamma-adaptin, a component of clathrin-coated vesicles associated with the Golgi apparatus. *J. Cell Biol.* 111, 2319–2326.
- Rohrer, J., Schweizer, A., Russell, D., and Kornfeld, S. (1996). The targeting of lamp1 to lysosomes is dependent on the spacing of its cytoplasmic tail tyrosine sorting motif relative to the membrane. *J. Cell Biol.* 132, 565–576.
- Sambrook, J., Fritsch, E.F., and Maniatis, T. (1989). *Molecular Cloning, A Laboratory Manual*, 2nd ed., Cold Spring Harbor, NY: Cold Spring Harbor Laboratory.
- Simpson, F., Bright, N.A., West, M.A., Newman, L.S., Darnell, R.B., and Robinson, M.S. (1996). A novel adaptor-related protein complex. *J. Cell Biol.* 133, 749–760.
- Stephens, D.J., and Banting, G. (1998). Specificity of interaction between adaptor-complex medium chains and the tyrosine-based sorting motifs of TGN38 and Igp120. *Biochem. J.* 335, 567–572.
- Stephens, D.J., Crump, C.M., Clarke, A.R., and Banting, G. (1997). Serine 331 and tyrosine 333 are both involved in the interaction

between the cytosolic domain of TGN38 and the μ 2 subunit of the AP2 clathrin adaptor complex. *J. Biol. Chem.* 272, 14104–14109.

Stepp, J.D., Huang, K., and Lemmon, S.K. (1997). The yeast adaptor protein complex, AP-3, is essential for the efficient delivery of alkaline phosphatase by the alternate pathway to the vacuole. *J. Cell Biol.* 139, 1761–1774.

Van Weert, A.W.M., Dunn, K.W., Geuze, H.J., Maxfield, F.R., and Stoorvogel, W. (1995). Transport from late endosomes to lysosomes, but not sorting of integral membrane proteins in endosomes, depends on the vacuolar proton pump. *J. Cell Biol.* 130, 821–834.

Vischer, U.M., and Wagner, D.D. (1993). CD63 is a component of Weibel-Palade bodies of human endothelial cells. *Blood* 82, 1184–1191.

Williams, M.A., and Fukuda, M. (1990). Accumulation of membrane glycoproteins in lysosomes requires a tyrosine residue at a particular position in the cytoplasmic tail. *J. Cell Biol.* 111, 955–966.

Wimer-Mackin, S., and Granger, B.L. (1996). Transmembrane domain mutations influence the cellular distribution of lysosomal membrane glycoprotein A. *Biochem. Biophys. Res. Commun* 229, 472–478.