

[Click here to view linked References](#)

Title Page

Manuscript title: Role of HIF-1 α and CASPASE-3 in cystogenesis of odontogenic cysts and tumors.

List of authors

Natacha M. M. da Costa, Adriane S. de Siqueira, André L. R. Ribeiro, Maria S. da S. Kataoka, Ruy G. Jaeger, Sérgio M. de Alves-Júnior, Andrew M. Smith, João de J. V. Pinheiro

Author affiliations and addresses

Natacha Malu Miranda da Costa, DDS, MSc

Cell Culture Laboratory, School of Dentistry, Federal University of Pará. Rua Augusto Corrêa, 01, Guamá, 66075110, Belém, PA, Brazil. natacha_malu@hotmail.com

Adriane Sousa de Siqueira, DDS, MSc, PhD

School of Dentistry, Universidade Positivo. Av. Prof. Pedro Viriato Parigot de Souza, 5300, 80740-050, Curitiba, PR, Brazil. adrianess@gmail.com

Andre Luis Ribeiro Ribeiro, DDS, MSc

Clinical lecturer, Department of Oral and Maxillofacial Surgery, University Center of Para - CESUPA, Belem, Brazil. ribeiroalr@ig.com.br

Maria Sueli da Silva Kataoka, DDS, MSc, PhD

Cell Culture Laboratory, School of Dentistry, Federal University of Pará. Rua Augusto Corrêa, 01, Guamá, 66075110, Belém, PA, Brazil. skataoka@ufpa.br

Ruy Gastaldoni Jaeger, DDS, MSc, PhD

Department of Cell and Developmental Biology Institute of Biomedical Sciences, University of São Paulo. Av. Prof. Lineu Prestes, 1524, Ed. Biomédicas 1, 05508-000, São Paulo, SP, Brazil. rgjaeger@usp.br

Sérgio Melo de Alves-Júnior, DDS, MSc, PhD

Cell Culture Laboratory, School of Dentistry, Federal University of Pará. Rua Augusto Corrêa, 01, Guamá, 66075110, Belém, PA, Brazil. sergiomalves@gmail.com

Andrew M. Smith, BSC, PhD

Department of Microbial Diseases, Eastman Dental Institute, University College London, London, UK. The Rayne Building, Centre for Molecular Medicine, 5 University Street, London-UK, WC1E 6JF. andrew.m.smith@ucl.ac.uk

João de Jesus Viana Pinheiro, DDS, MSc, PhD (corresponding author)

Cell Culture Laboratory, School of Dentistry, Federal University of Pará. Rua Augusto Corrêa, 01, Guamá, 66075110, Belém, PA, Brazil. radface@hotmail.com

Corresponding author: João de Jesus Viana Pinheiro

Address: Cell Culture Laboratory, School of Dentistry, Federal University of Pará. Rua Augusto Corrêa, 01, Guamá, 66075110, Belém, PA, Brazil.

Postcode: 66075110

Telephone: +55 91 32590335

+55 91999850765

E-mail: radface@hotmail.com

Acknowledgements

This investigation was supported by the Brazilian National Council for Scientific and Technological Development (CNPq grants 446879/2014; 201261/2015-4).

The author Andre Luis Ribeiro Ribeiro is grateful to CAPES foundation, Ministry of Education of Brazil, for funding his PhD scholarship (grant no. 0698130).

Abstract

1 **Objectives:** Odontogenic cysts and tumors are the most relevant lesions that affect the gnathic bones.
2 These lesions have in common the formation of cystic areas and this common feature may suggest
3 involvement of similar mechanisms. The hypoxia inducible factor 1 alpha (HIF-1 α), a responsive protein
4 to hypoxia and caspase-3, an irreversible apoptosis marker, may contribute to cyst formation. Thus, this
5 study aimed to investigate the immunoexpression of these proteins in odontogenic cysts and tumors.

6 **Material and methods:** Twenty cases of ameloblastoma, keratocystic odontogenic tumor (KOT) (n=20),
7 radicular cyst (RC) (n=18), dentigerous cyst (DC) (n=11), calcifying cystic odontogenic tumor (n=8) and
8 dental follicle (DF) (n=10) were used to investigate HIF-1 α and caspase-3 expression in sequential serial
9 cuts by immunohistochemistry.

10 **Results:** HIF-1 α was overexpressed in RC, DC and ameloblastoma when compared with DF. The basal
11 and sometimes the lower suprabasal layer showed no or very low expression in DC, KOT and
12 ameloblastoma, the last also showing strong expression in solid epithelial areas and initial cystic
13 formation regions. Caspase-3 was found to be overexpressed in all lesions, with the highest expression in
14 odontogenic cysts compared to tumors. HIF-1 α and caspase-3 were localized in similar areas of the same
15 lesions, especially in the epithelium surrounding cystic formations.

16 **Conclusions:** This study showed distinct immunoexpression of HIF-1 α and caspase-3 in odontogenic
17 cyst and tumors, with higher expression observed in odontogenic cysts.

18 **Clinical relevance:** These finding suggesting a possible correlation between hypoxia, apoptosis and
19 cystogenesis, leading to understand the mechanisms responsible to cystic formation in odontogenic
20 lesions.
21
22

23
24 **Keywords:** Odontogenic tumors; Odontogenic cyst; HIF-1 α ; Caspase-3; Cystogenesis
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

INTRODUCTION

1 Odontogenic cysts and tumors are lesions that develop exclusively in or around gnathic bones [1,
2 2]. They are characterized by the presence of a lining epithelium of odontogenic origin and a cystic area
3 containing liquid and/or a fluid material is often observed [1, 2]. Due to their high frequency and/or local
4 aggressiveness, ameloblastoma and keratocystic odontogenic tumor (KOT) represent the most clinical
5 relevant odontogenic tumors, and radicular cyst (RC) and dentigerous cyst (DC) the most important
6 odontogenic cysts. Some odontogenic tumors, such as ameloblastoma, may have cystic and solid areas [2-
7 5]. Little is known regarding the pathogenesis of cyst formation but it is supposed to involve proliferation
8 and death of cells of the lining epithelium [6-8].

14 There are a few proposed theories about the development cystic areas in odontogenic lesions [9,
15 10], but majority of them lack of scientific evidence. The strongest and most accepted theory suggests
16 that the development of odontogenic cysts as a result of an increased epithelial proliferation creating large
17 epithelial island, leading to deficient nutritional supply in central areas. Thus, these centrally located cells
18 undergo to necrosis initiating the cystic cavity and also exposing the hyperosmolar intracellular contents,
19 attracting liquid from the surrounding tissues, contributing to cyst growth [10-12].

23 Odontogenic cysts and tumors are known for a high proliferation rate of the epithelium
24 parenchyma [8, 13-15]. This cellular proliferation will result in an agglomerate of epithelial cells and
25 formation of cellular islands [14, 15]. Because epithelial cells have no direct blood supply, all nutrients
26 required for their metabolism are acquired from the surround connective tissue [16]. This proliferation
27 and cellular aggregation possibly limits the oxygen (O_2) diffusion to the cells in the center of the island,
28 leading to apoptosis and inducing cyst formation [12, 15].

33 Low oxygen (O_2) concentrations ($O_2 \leq 2\%$) [17] are known to create a condition called hypoxia,
34 which is responsible for triggering several cellular events [18]. Different proteins and transcription factors
35 have been associated to hypoxic conditions, which has in the hypoxia inducible factor 1 alpha (HIF-1 α)
36 [19].

40 Hypoxia increases HIF-1 α levels by inhibiting its degradation [20], which induces many different
41 types of cellular responses that include cell proliferation, angiogenesis and apoptosis [21]. Proliferation of
42 the epithelium and apoptosis are fundamental events in cystogenesis [12,22] and have been related to the
43 development of odontogenic cysts and tumors [15, 23]. Hypoxia has already been associated with cyst
44 formation in ameloblastoma [24].

48 Apoptosis has important roles in a variety of physiological and pathological phenomena [25].
49 Apoptosis is a well-regulated process and occurs because of the activation of members of the cysteine-
50 aspartic proteases (caspase) family [26]. Activated caspase-3 is considered as an apoptosis marker, due to
51 its critical and irreversible role in the mechanism of controlled cell death [27].

55 It is believed that cyst formation can be a result of an imbalance between cell proliferation and cell
56 death [28]. Further evidences suggest the modulation of apoptosis by HIF-1 α in pathologic and
57 physiologic conditions [28-30]. During hypoxia and consequent overexpression of HIF-1 α , the cell can
58 get into apoptotic cell death after initiating a caspase-dependent cascade of events [31].

1 Many events can occur and influence the microenvironment that surrounds odontogenic lesions.
2 Thus, this study aimed to evaluate HIF-1 α and caspase-3 immunoexpression in ameloblastoma, KOT,
3 calcifying cystic odontogenic tumor (CCOT), RC, and DC in order to identify the expression of these
4 proteins and their possible association and participation in the formation of cystic areas of these lesions.
5

6 **MATERIALS AND METHODS**

7 **Samples**

8 Tissue microarrays (TMA) containing 20 ameloblastoma cores (Biomax Inc., Rockville, MD,
9 USA), and tissue sections of odontogenic cysts and tumors were used in this study. Cases of KOT (20),
10 RC (18), DC (11), CCOT (8) with 5 μ m thickness completed our sample. Additionally, ten cases of dental
11 follicle (DF) were included as a non-pathological control of normal dental tissue exhibiting neither cystic
12 formation nor neoplastic changes [2]. Samples groups were classified in odontogenic tumors
13 (ameloblastoma, KOT and CCOT) or odontogenic cysts (RC and DC). KOT, RC, DC, CCOT and DF
14 sections were obtained in consecutive cut series from the archives of the department of Oral Pathology of
15 the School of Dentistry of the University Center of Pará (CESUPA, Belém-PA, Brazil). Cystic areas were
16 selected in both odontogenic cysts and tumors. Diagnosis of each lesion was confirmed by a pathologist
17 before experimental procedures. This study was approved by the Human Research Ethics Committee of
18 the Institute of Health Sciences of the Federal University of Pará (CAAE: 36572414.7.0000.0018, nr
19 877.322/2014).
20
21
22
23
24
25
26
27
28

29 **Immunohistochemistry**

30 Histological sections, obtained from the KOT, RC, DC, CCOT and DF samples, were mounted
31 on glass slides treated with 3-Aminopropyl triethoxysilane (Sigma Chemical Corp, St. Louis, MO, USA).
32 TMA cores of ameloblastoma and samples sections were deparaffinized in xylene and hydrated using
33 decreasing concentrations of ethanol. The slides were immersed in a 20% H₂O₂/methanol (1:1) solution
34 for 20 minutes to inhibit endogenous peroxidase activity. Antigen retrieval was performed for 30 seconds
35 in citrate buffer (pH 6.0) using a Pascal chamber (Dako, Carpinteria, CA, USA). Non-specific antibody
36 binding was blocked using 1% bovine serum albumin (BSA, Sigma Chemical Corp.) in a phosphate-
37 buffered saline solution (PBS) for 1 hour.
38
39
40
41
42
43

44 The slides were incubated with primary antibodies anti-HIF-1 α (1:25, clone H1 α 67, Merck
45 Millipore, Darmstadt, Germany) and anti-caspase-3 (1:100, clone CPP32, Diagnostic BioSystems,
46 Pleasanton CA, USA), which reacts against with the active form of this protein, for 1 hour and followed
47 by a 30-minute incubation with EnVision Plus detection system (Dako®). Diaminobenzidine (Dako®)
48 was used as the chromogen. Sections were counterstained with Mayer's hematoxylin (Sigma Chemical
49 Corp.) and mounted using Permount (Fisher Scientific, Fair Lawn, NJ, USA). As a negative control, the
50 primary antibodies were replaced by non-immune serum. Squamous cell carcinoma samples were used as
51 positive control for the immunohistochemical reaction.
52
53
54
55
56

57 **Immunostaining evaluation**

58 The immunostaining evaluation was performed by analyzing the percentage of HIF-1 α and
59 caspase-3 stained area in the epithelial layer in all samples. Brightfield images of five randomly selected
60
61
62
63
64
65

1 areas were acquired for each sample using AxioScope microscope equipped with an AxioCamHRc CCD
2 color camera (Carl Zeiss, Oberkochen, Germany), with a 40× objective. For this step, the sites selected
3 for quantification were from cystic areas, from both odontogenic cists and tumors. Diaminobenzidine
4 stained areas were separated and segmented using the “deconvolution color plug-in” (Gabriel Landini,
5 <http://www.dentistry.bham.ac.uk/landinig/software/software.html>) from ImageJ software (public domain
6 software, NIMH, NIH, Bethesda, MD, USA, <http://rsbweb.nih.gov/ij/>). After image segmentation, the
7 fraction of stained area was measured. Differences between ameloblastoma, KOT, RC, DC, CCOT and
8 DF immunoexpression were then analyzed. The samples were evaluated by one examiner in a blinded
9 study towards diagnosis. The evaluation of expression of HIF-1 α and caspase 3 in odontogenic cysts and
10 tumors was conducted descriptively.
11
12
13
14

15 **Statistical analysis**

16
17 The data was analyzed using GraphPad Prism 5 software (GraphPad Software Inc., San Diego,
18 CA, USA). Following normality analysis, a two-way ANOVA followed by Tukey’s multiple comparison
19 test was used to evaluate the differences in HIF-1 α and caspase-3 stained areas between samples and
20 control.
21
22

23 **RESULTS**

24 **HIF-1 α and caspase-3 were expressed in odontogenic cysts and tumors**

25
26
27 All samples of odontogenic cysts and tumors analyzed in this study expressed HIF-1 α and
28 activated caspase-3 (Figures 1-5). Differences among these lesions were evaluated in consideration of
29 staining intensity, cellular immunolocalization and epithelial layers distribution. Subsequent cuts from the
30 same area were analyzed in parallel for HIF-1 α and caspase-3 immunoexpression.
31
32
33
34

35
36 In ameloblastoma TMA cores, HIF-1 α expression (Figures 1A, 1C, 1E and 1G) was observed as
37 intense nuclear (Figure 1C, arrowhead) and cytoplasmic staining (Figure 1C, arrow) in the epithelial layer
38 near and/or in close contact with the cystic lumen (Figures 1A and 1C, asterisk). On the other hand, the
39 cells of the basal and lower suprabasal layer, showed a weak cytoplasmic staining. There was an intense
40 nuclear and cytoplasmic staining in epithelial islands, where cells presenting large vacuoles areas could
41 be observed (Figure 1E, arrow), and also in cells surrounding initial cystic formation (Figure 1E,
42 asterisk). The cells located in the solid epithelial proliferations of tumor parenchyma displayed intense
43 nuclear and cytoplasmic (Figure 1G, arrow) HIF-1 α staining.
44
45
46
47

48 Caspase-3 analysis (Figures 1B, 1D, 1F e 1H) was carried out using similar regions of the
49 ameloblastoma TMA cores compared to those of HIF-1 α . Caspase-3 immunostaining predominated in the
50 cytoplasm of cells near to cystic areas (Figures 1B and 1D, asterisk) of the epithelial islands and nuclear
51 and cytoplasmic staining in selected cells of the basal layer (Figure 1D, arrow). In tumor parenchyma,
52 areas of cyst formation (Figures 1F, asterisk) and solid areas of epithelial islands showed almost an
53 exclusively cytoplasmic staining for caspase-3 (Figures 1F and 1H, arrow).
54
55
56
57

58 All KOT samples showed colocalized HIF-1 α and caspase-3 staining (Figure 2). HIF-1 α
59 immunoexpression was observed in both cellular compartments, but predominantly located in the
60
61
62
63
64
65

1 parabaasal layer near to the cystic cavity (Figures 2A and C, asterisk) in cytoplasm (Figure 2C, arrow) and
2 nucleus (Figure 2C, arrowhead). No HIF-1 α staining was observed in the basal and lower suprabasal
3 layer. On the other hand, caspase-3 (Figures 2B and 2D) showed cytoplasmic (Figure 2D, arrow) and
4 nuclear (Figure 2D, arrowhead) expression in selected neoplastic cells of all epithelial layers. Expression
5 of caspase-3 appeared to be less intense in the basal layer.
6

7
8 In RC, HIF-1 α (Figures 3A and 3C) and caspase-3 (Figures 3B and 3D) were similarly observed
9 in the cell nucleus (Figures 3C and 3D, arrowhead) and cytoplasm (Figures 3C and 3D, arrow) of cystic
10 epithelium. In cystic stroma, inflammatory cells were positively stained for both proteins (Figures 3E and
11 3F, arrow).
12

13
14 In DC samples, HIF-1 α (Figures 4A and 4C) showed a similar expression to those observed in
15 KOT samples, with cytoplasmic (Figure 4C, arrow) and nuclear staining (Figures 4C, arrowhead) of the
16 suprabasal cells and no staining of the basal layer. However, caspase-3 staining showed a predominantly
17 cytoplasmic immunolocalization especially in the suprabasal layer (Figure 4D), with some basal cells
18 showing both, nuclear and cytoplasmic staining (Figure 4D, arrow).
19

20
21 HIF-1 α and caspase-3 staining showed a similar pattern in CCOT samples. The expression of these
22 proteins was localized in the cytoplasm and distributed similarly in all epithelial layers (Figures 5A and
23 5B). The DF exhibited weak HIF-1 α and moderate caspase-3 staining in the cytoplasm of the epithelial
24 cells (Figures 5C and 5D).
25
26
27
28
29
30

31 **HIF-1 α is highly expressed in ameloblastoma, DC, and comparatively higher in RC when compared** 32 **to odontogenic tumors** 33

34
35 HIF-1 α expression showed variable expression in all samples (Figure 6A). The highest
36 expression was observed in RC (Mean= 58.11; SD=11.61), which was statistically significant when
37 compared to odontogenic tumors (ameloblastoma p=0.026, Mean=44.69, SD=13.87; KOT p<0.0001,;
38 CCOT p<0.0001). Stained area of KOT (Mean=36,20, SD=15.17) and CCOT (Mean=17.60, SD=12.77)
39 was not statistically different from control (Mean=23.07, SD=6.35), and HIF-1 α expression in CCOT was
40 lower when compared with all other lesions (ameloblastoma p=0.0004; KOT p=0.0368; DC p=0.0005,
41 Mean=46,25, SD=13.50; and RC p<0.0001). Although there was not difference between HIF-1 α
42 immunostaining between control and CCOT, CCOT expression was lower than all the other studied
43 lesions.
44
45
46
47
48
49
50

51 **Caspase-3 is highly expressed in odontogenic cysts compared to odontogenic tumors.** 52

53
54 Results of caspase-3 immunostaining quantification demonstrated that odontogenic cysts showed
55 higher percentage of stained areas than odontogenic tumors (Figure 6B). DC (Mean=65,57, SD=13.86)
56 showed the highest expression of caspase-3, followed by RC (Mean=60.23; SD=7.58), ameloblastoma
57 (Mean=50.16, SD=6.63) and KOT (Mean=49.44, SD=8.23). Comparison between odontogenic cysts and
58 tumors revealed higher expression in DC when compared to ameloblastoma (p=0.0005), KOT (p=0.0002)
59
60
61
62
63
64
65

1 and CCOT ($p < 0.0001$, Mean=43.32, SD=13.14) and also higher expression in RC against the same
2 ameloblastoma ($p = 0.0175$), KOT ($p = 0.0086$) and CCOT ($p < 0.0016$). No differences were observed when
3 compared odontogenic cysts or tumors with each other. Caspase-3 expression was significantly higher in
4 all studied samples ($p < 0.0001$) when compared to the non-pathological control of DF (Mean=13.82,
5 SD=10.53).
6

7 DISCUSSION

8
9 Although the pathogenesis of odontogenic lesions may differ in developmental, inflammatory or
10 neoplastic lesions, they can present a common feature that is cystic formation. The reasons why it
11 happens is poorly understood, which arises the question why lesions of different pathogenesis display
12 similar pathological features. It is considered that many of these processes start with proliferation of the
13 odontogenic epithelium [9, 10]. According these cells grow, the central cells start to have more restricted
14 access to oxygen, leading them to hypoxia. In this study was observed a high expression of the hypoxia-
15 related protein HIF-1 α in odontogenic cysts and ameloblastoma and also high immunorexpression of
16 caspase-3 in all odontogenic cysts and tumors.
17
18

19 HIF-1 α is the main cellular protein in response to hypoxic conditions [21]. Although HIF-1 α is
20 normally produced, it is quickly degraded in normoxic condition [19]. HIF-1 α accumulation is only
21 possible at low oxygen concentration levels [19]. Thus, in this context HIF-1 α can be considered a
22 hypoxia marker [32]. In this study, odontogenic cysts showed the highest immunorexpression of HIF-1 α ,
23 suggesting that these lesions undergo to restricted oxygen supply. Ameloblastoma samples also showed
24 increased expression of HIF-1 α in solid areas of epithelial islands, supporting the idea of localized
25 intratumoral hypoxia [24].
26
27

28 KOT, DC and cystic areas of ameloblastoma showed an interesting pattern of HIF-1 α
29 immunostaining. The basal and, in some cases, the suprabasal layer showed no HIF-1 α immunostaining,
30 while the upper layers close to the cystic area and more distant from the connective tissue were positively
31 stained. It strongly suggests a reduced oxygen supply when the cells distantiate from the basal layer,
32 which normally occurs as result of proliferation. This situation leads to stabilization of HIF-1 α activating
33 its response.
34
35

36 Upon activation, HIF-1 α can modulate many cellular responses, including inflammation and
37 apoptosis [32]. This study showed that odontogenic cysts and ameloblastoma have a higher expression of
38 HIF-1 α when compared to DF. It is more evident in RC, which showed an extensive inflammatory
39 infiltrate in the stroma [33].
40
41

42 We then hypothesize that with proliferation of the epithelial cells from basal layer, the cells of
43 the upper layers experience limited O₂ supply, causing a hypoxia-related response overexpressing HIF-1 α .
44 Under certain conditions, the protective mechanism to hypoxia that initiates HIF-1 α response is not
45 enough to maintain cell viability, thus triggering a more extreme response that leads to apoptosis
46 throughout activation of a caspase cascade, analyzed here by the expression of the irreversible apoptosis
47 marker activated caspase-3. Caspase-3 and HIF-1 α were observed in co-localization in this study,
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 suggesting that these processes can be associated and together may play a role in formation of cystic
2 areas.

3 HIF-1 α is known to induce cellular apoptosis [30, 34, 35] and also a key molecule [36, 37] that
4 can overcome stabilization mediated by the p53 suppressor gene [38]. HIF-1 α can promote the release of
5 cytochrome C to the mitochondria and subsequent activation of caspase-3 [31, 39, 40]. This study pointed
6 out that HIF-1 α expression was higher in odontogenic cysts and ameloblastoma. Overexpression of p53
7 suppressor gene is a common found in ameloblastoma [41], which also showed p53 gene mutations and
8 allelic loss [42].

9 Previous studies showed that HIF-1 α knockout suppresses the apoptosis induced by p53 [30] and
10 significantly reduces cyst formation [34, 43]. There is strong evidence of a correlation between HIF-1 α
11 accumulation and caspase-3 activation [44]. In our samples, HIF-1 α and caspase-3 were found to be
12 expressed in the same areas of all lesions, which support the association between those two molecules. An
13 interesting immunolocalization pattern with higher expression of both proteins when the cells were
14 located more distant from the basal layer was observed in cases of ameloblastoma, KOT and DC.

15 Caspase-3 staining in RC and DC was mostly nuclear and cytoplasmic, while in odontogenic
16 tumors was predominantly cytoplasmic. DNA fragmentation is a major step in eliminating the genome of
17 apoptotic cells and subsequent cell death [45]. Hence, intense nuclear staining indicates the beginning of
18 apoptosis in odontogenic cysts. Additionally, apoptosis can also be a protection against oxidative stress or
19 other forms of DNA damage and thereby reduce the risk of neoplastic transformation [46].

20 Ameloblastoma showed a strong nuclear and cytoplasmic staining of HIF-1 α and caspase-3 in
21 specific solid areas of epithelial islands, which are known to present difficulties in O₂ and nutrient
22 diffusion [12, 22], and also in surrounding areas of initial cyst formation. This could resemble the ideal
23 environment were hypoxia is established [47] and apoptosis will take place to form future cystic regions.
24 Some cells in epithelial areas of ameloblastoma presented large vacuoles areas, which is a classical sign
25 of cell death [48, 49]. Low expression of CCOT was observed for both proteins compared to the other
26 lesions. CCOT is a non-aggressive odontogenic neoplasm that shows an indolent behavior and low
27 recurrence rates, which has been widely used as a control group for studying odontogenic tumors [50-54].

28 Although this study can not prove that these events are occurring in consequence of the same
29 cause (hypoxia), the results observed here support this hypothesis and suggest that there might have an
30 association between hypoxia and apoptosis, especially in cases of odontogenic cysts and ameloblastoma.
31 HIF-1 α and caspase-3 immunoexpression and immunolocalization patterns suggest an important
32 association between hypoxia, apoptosis and cyst formation. **Therefore, HIF-1 α and caspase-3 expression
33 labelling in odontogenic cysts and tumors may be important to explain which cellular components are
34 related to the behavior of these lesions. Additionally, these molecules can be used as biological markers
35 in the future, serving as instruments for better diagnosis and treatment planning.**

36 **Compliance with Ethical Standards**

Conflict of Interest: The author Natacha Malu Miranda da Costa declare that she has no conflict of interest. The author Adriane Sousa de Siqueira declare that she has no conflict of interest. The author Andre Luis Ribeiro Ribeiro declare that he has no conflict of interest. The author Maria Sueli da Silva Kataoka declare that she has no conflict of interest. The author Sérgio Melo de Alves-Junior declare that he has no conflict of interest. The author Andrew M. Smith declare that he has no conflict of interest. The author João de Jesus Viana Pinheiro declare that he has no conflict of interest.

Funding: The work was supported by the Department of Oral Pathology of the School of Dentistry of the University Center of Pará (CESUPA, Belém-PA, Brazil) and Cell Culture Laboratory of School of Dentistry of Federal University of Pará (UFPA, Belém-PA, Brazil).

Ethical approval: This article does not contain any studies with human participants or animals performed by any of the authors.

Informed Consent: For this type of study, formal consent is not required.

References

1. Sciubba JJ, Fantasia JE, Kahn LB (2000) Atlas of Tumor Pathology. Tumors and Cyst of Jaw, AFIP, Washington DC.
2. Barnes L, Eveson JW, Reichart P, Sidransky D (2005) World Health Organization classification of tumors. Pathology and genetics of head and neck tumours, 9th edn. IARC Press, Lyon, pp 296–300.
3. Rapidis AD, Andressakis DD, Stavrianos SD, Faratzis G, Arniogiannaki-Liappi N, Lagogiannis GA, Valsamis SV, Apostolikas N (2004) Ameloblastomas of the jaws: clinico-pathological review of 11 patients. *Eur J Surg Oncol* 30:998-1002.
4. Reichart PA, Philipsen HP, Sonner S (1995) Ameloblastoma: biological profile of 3677 cases. *Eur J Cancer B Oral Oncol* 31B:86-99.
5. Chirapathomsakul D, Sastravaha P, Jansisyanont P (2006) A review of odontogenic keratocysts and the behavior of recurrences. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 101:5-9.
6. Shear M, Speight P (2007) Cysts of the Oral and Maxillofacil Regions. Wright, Oxford.
7. Ward JP, Magar V, Franks SJ, Landini G (2004) A mathematical model of the dynamics of odontogenic cyst growth. *Anal Quant Cytol Histol* 26:39-46.
8. Piattelli A, Fioroni M, Santinelli A, Rubini C (1998) Expression of proliferating cell nuclear antigen in ameloblastomas and odontogenic cysts. *Oral Oncol* 34:408-412.
9. Lin LM, Huang GT, Rosenberg PA (2007) Proliferation of epithelial cell rests, formation of apical cysts, and regression of apical cysts after periapical wound healing. *J Endod* 33:908-916.
10. Orstavik Dag, Ford TRP (1998) Essential Endodontology: Prevention And Tratament Of Apical Periodontitis, 1st edn. Blackwell Science, Oxford, pp 68 –105.
11. Ten Cate AR (1972) The epithelial cell rests of Malassez and the genesis of the dental cyst. *Oral Surg Oral Med Oral Pathol* 34:956-964.
12. Toller P (1967) Origin and growth of cysts of the jaws. *Ann R Coll Surg Engl* 40:306-336.
13. Thosaporn W, Iamaroon A, Pongsiriwet S, Ng KH (2004) A comparative study of epithelial cell proliferation between the odontogenic keratocyst, orthokeratinized odontogenic cyst, dentigerous cyst, and ameloblastoma. *Oral Dis* 10:22-26.
14. Seifi S, Shafigh E, Allaie A (2011) Quantitative and qualitative analysis of argyrophilic nuclear organizer regions in follicular cyst, keratocystic odontogenic tumor and ameloblastoma. *J Cancer Res Ther* 7:280-285.
15. Martins CA, Rivero ER, Dufloth RM, Figueiredo CP, Vieira DS (2011) Immunohistochemical detection of factors related to cellular proliferation and apoptosis in radicular and dentigerous cysts. *J Endod* 37:36-39.
16. Kefalides NA, Borel JP (2005) Basement membranes: cell and molecular biology. Academic Press, California.

17. Hammond EM, Asselin MC, Forster D, O'Connor JP, Senra JM, Williams KJ (2014) The meaning, measurement and modification of hypoxia in the laboratory and the clinic. *Clin Oncol* 26:277-288.
18. Kieda C, Greferath R, Crola da Silva C, Fylaktakidou KC, Lehn JM, Nicolau C (2006) Suppression of hypoxia-induced HIF-1 α and of angiogenesis in endothelial cells by myo-inositol trispyrophosphate-treated erythrocytes. *Proc Natl Acad Sci U S A* 103:15576-15581.
19. Lee JW, Bae SH, Jeong JW, Kim SH, Kim KW (2004) Hypoxia-inducible factor (HIF-1) α : its protein stability and biological functions. *Exp Mol Med* 36:1-12.
20. Wang GL, Semenza GL (1995) Purification and characterization of hypoxia-inducible factor 1. *J Biol Chem* 270:1230-1237.
21. Maxwell P (2003) Hif-1: An oxygen response system with special relevance to the kidney. *J Am Soc Nephrol* 14:2712-2722.
22. Edelstein CL (2005) What is the role of tubular epithelial cell apoptosis in polycystic kidney disease (pkd)? *Cell Cycle* 4:1550-1554.
23. Amaral FR, Mateus GC, Bonisson LA, de Andrade BA, Mesquita RA, Horta MC, Marigo Hde A (2012) Cell proliferation and apoptosis in ameloblastomas and keratocystic odontogenic tumors. *Braz Dent J* 23:91-96.
24. Da Costa NM, Fialho AD, Proietti CC, da Silva Kataoka MS, Jaeger RG, de Alves-Junior SM and de Jesus Viana Pinheiro J (2016) Role of hypoxia-related proteins in invasion of ameloblastoma cells: crosstalk between NOTCH1, hypoxia-inducible factor 1 α , a disintegrin and metalloproteinase 12, and heparin-binding epidermal growth factor. *Histopathology* 69:99-106.
25. Oral O, Akkoc Y, Bayraktar O, Gozuacik D (2016) Physiological and pathological significance of the molecular cross-talk between autophagy and apoptosis. *Histol Histopathol* 31:479-498.
26. Shalini S, Dorstyn L, Dawar S, Kumar S (2015) Old, new and emerging functions of caspases. *Cell Death Differ* 22:526-539.
27. Gregoraszczyk EL, Rak-Mardyla A, Rys J, Jakubowicz J, Urbanski K (2015) Effect of chemotherapeutic drugs on caspase-3 activity, as a key biomarker for apoptosis in ovarian tumor cell cultured as monolayer. A pilot study. *Iran J Pharm Res* 14:1153-1161.
28. Edelstein CL (2008) Mammalian target of rapamycin and caspase inhibitors in polycystic kidney disease. *Clin J Am Soc Nephrol* 3:1219-1226.
29. Tao Y, Kim J, Faubel S, Wu JC, Falk SA, Schrier RW, Edelstein CL (2005) Caspase inhibition reduces tubular apoptosis and proliferation and slows disease progression in polycystic kidney disease. *Proc Natl Acad Sci U S A* 102:6954-6959.
30. Suzuki H, Tomida A, Tsuruo T (2001) Dephosphorylated hypoxia-inducible factor 1 α as a mediator of p53-dependent apoptosis during hypoxia. *Oncogene* 20:5779-5788.
31. Greijer AE, Wall EVD (2004) The role of hypoxia inducible factor 1 (hif-1) in hypoxia induced apoptosis. *J Clin Pathol* 57:1009-1014.
32. Palazon A, Goldrath AW, Nizet V, Johnson RS (2014) Hif transcription factors, inflammation, and immunity. *Immunity* 41:518-528.
33. Mourao RV, Junior EC, Barros Silva PG, Turatti E, Mota MR, Alves AP (2016) Study of the relationship between mononuclear inflammatory infiltrate and ki-67 and basement membrane and extracellular matrix protein expression in radicular cysts. *Int Endod* 49:447-453.
34. Belibi F, Zafar I, Ravichandran K, Segvic AB, Jani A, Ljubanovic DG, Edelstein CL (2011) Hypoxia-inducible factor-1 α (HIF-1 α) and autophagy in polycystic kidney disease (PKD). *Am J Physiol Renal Physiol* 300:1235-1243.
35. Bohensky J, Shapiro IM, Leshinsky S, Terkhorn SP, Adams CS, Srinivas V (2007) Hif-1 regulation of chondrocyte apoptosis: Induction of the autophagic pathway. *Autophagy* 3:207-214.
36. Carmeliet P, Dor Y, Herbert JM, Fukumura D, Brusselmans K, Dewerchin M, Neeman M, Bono F, Abramovitch R, Maxwell P, Koch CJ, Ratcliffe P, Moons L, Jain RK, Collen D, Keshert E (1998) Role of HIF-1 α in hypoxia-mediated apoptosis, cell proliferation and tumour angiogenesis. *Nature* 394:485-490.
37. Moritz W, Meier F, Stroka DM, Giuliani M, Kugelmeier P, Nett PC, Lehmann R, Candinas D, Gassmann M, Weber M (2002) Apoptosis in hypoxic human pancreatic islets correlates with HIF-1 α expression. *FASEB J* 16:745-747.

38. Chen D, Li M, Luo J, Gu W (2003) Direct interactions between hif-1 alpha and mdm2 modulate p53 function. *J Biol Chem* 278:13595-13598.
39. Van Hoecke M, Prigent-Tessier AS, Garnier PE, Bertrand NM, Filomenko R, Bettaieb A, Marie C, Beley AG (2007) Evidence of HIF-1 functional binding activity to caspase-3 promoter after photothrombotic cerebral ischemia. *Mol Cell Neurosci* 34:40-47.
40. Cho SJ, Teng IF, Zhang M, Yin T, Jung YS, Zhang J, Chen X (2015) Hypoxia-inducible factor 1 alpha is regulated by RBM38, a RNA-binding protein and a p53 family target, via mRNA translation. *Oncotarget* 6:305-316.
41. Gadbaill AR, Patil R, Chaudhary M (2012) Co-expression of ki-67 and p53 protein in ameloblastoma and keratocystic odontogenic tumor. *Acta Odontol Scand* 70:529-535.
42. Nodit L, Barnes L, Childers E, Finkelstein S, Swalsky P, Hunt J (2004) Allelic loss of tumor suppressor genes in ameloblastic tumors. *Mod Pathol* 17:1062-1067.
43. Buchholz B, Schley G, Faria D, Kroening S, Willam C, Schreiber R, Klanke B, Burzlaff N, Jantsch J, Kunzelmann K, Eckardt KU (2014) Hypoxia-inducible factor-1alpha causes renal cyst expansion through calcium-activated chloride secretion. *J Am Soc Nephrol* 25:465-474.
44. Casely-Hayford MA, Nicholas SA, Sumbayev VV (2009) Azinomycin epoxide induces activation of apoptosis signal-regulating kinase 1 (ask1) and caspase 3 in a hif-1alpha-independent manner in human leukaemia myeloid macrophages. *Eur J Pharmacol* 602:262-267.
45. Zhang JH, Xu M (2000) DNA fragmentation in apoptosis. *Cell Res* 10: 205-211.
46. Torres VE (1999) Apoptosis in cystogenesis: Hands on or hands off? *Kidney Int* 55:334-335.
47. Kohl R, Zhou J, Brune B (2006) Reactive oxygen species attenuate nitric-oxide-mediated hypoxia-inducible factor-1alpha stabilization. *Free Radic Biol Med* 40:1430-1442.
48. Kichi E, Enokiya Y, Muramatsu T, Hashimoto S, Inoue T, Abiko Y, Shimono M (2005) Cell proliferation, apoptosis and apoptosis-related factors in odontogenic keratocysts and in dentigerous cysts. *Oral Pathol Med* 34:280-6.
49. Metgud R, Gupta K (2013) Expression of cell cycle and apoptosis-related proteins in ameloblastoma and keratocystic odontogenic tumor. *Ann Diagn Pathol* 17:518-521.
50. Pinheiro JJ, Freitas VM, Moretti AI, Jorge AG, Jaeger RG (2004) Local invasiveness of ameloblastoma. Role played by matrix metalloproteinases and proliferative activity. *Histopathology* 45:65-72.
51. Siqueira AS, Carvalho MR, Monteiro AC, Freitas VM, Jaeger RG, Pinheiro JJ (2010) Matrix metalloproteinases, timp3 and growth factors regulating ameloblastoma behaviour. *Histopathology* 57:128-137.
52. Ribeiro AL, Nobre RM, Alves-Junior SM, Kataoka MS, Barroso RF, Jaeger RG, Pinheiro JJ (2012) Matrix metalloproteinases, tissue inhibitors of metalloproteinases, and growth factors regulate the aggressiveness and proliferative activity of keratocystic odontogenic tumors. *Oral Surg Oral Med Oral Pathol Oral Radiol* 114:487-496.
53. Ribeiro AL, Nobre RM, Rocha GC, Lobato IH, Alves-Junior SM, Jaeger RG, Pinheiro JJ (2011) Expression of metallothionein in ameloblastoma. A regulatory molecule? *J Oral Pathol Med* 40:516-519.
54. Ribeiro AL, da Costa NM, de Siqueira AS, Brasil da Silva W, da Silva Kataoka MS, Jaeger RG, Alves-Junior SM, Smith AM, Pinheiro JJ (2016) Keratocystic odontogenic tumor overexpresses invadopodia-related proteins, suggesting invadopodia formation. *Oral Surg Oral Med Oral Pathol Oral Radiol* 122:500-508.

FIGURE LEGENDS

Figure 1: HIF-1 α and caspase-3 immunostaining in ameloblastoma. HIF-1 α staining (A, C, E, G) was predominant in the nucleus (C, arrowhead) and cytoplasm (C, arrow) in epithelial layer near cystic lumen (A, C, asterisk). Intense nuclear and cytoplasmic staining in epithelial islands (E and G,

1 arrow), and in cells surrounding initial cystic formation was observed (E, asterisk). Caspase-3
2 immunostaining (B,D,F and H) is predominant in the cytoplasm of cells near cystic areas of epithelial
3 islands (B and D, asterisk). Staining is also detected in selected cells of basal layer (D, arrow). In tumor
4 parenchyma, areas of cyst formation (F, asterisk) and solid areas of epithelial islands showed almost an
5 exclusively cytoplasmic staining for caspase-3 (F and H, arrow). Scale bar: 50 μ m and 20 μ m.
6

7
8 **Figure 2:** HIF-1 α and caspase-3 immunostaining in KOT. HIF-1 α immunoexpression was
9 predominantly located in the parabasal layer near to the cystic cavity (A and C, asterisk) in cytoplasm (C,
10 arrow) and nucleus (C, arrowhead). Caspase-3 (B and D) showed cytoplasmic (D, arrow) and nuclear (D,
11 arrowhead) expression in selected neoplastic cells of all epithelial layers. Expression of caspase-3
12 appeared to be less intense in the basal layer. Immunoperoxidase. Scale bars: 50 μ m and 20 μ m.
13
14
15
16

17
18 **Figure 3:** HIF-1 α and caspase-3 immunostaining in RC. Expression of HIF-1 α (A and C) and
19 caspase-3 (B and D) were similarly observed in the cell nucleus (C and D, arrowhead) and cytoplasm (C
20 and D, arrow) of cystic epithelium. In cystic stroma, inflammatory cells were positively stained for both
21 proteins (E and F, arrow). Immunoperoxidase. Scale bars: 50 μ m and 20 μ m.
22
23
24
25

26
27 **Figure 4:** HIF-1 α and caspase-3 immunostaining in DC. HIF-1 α (A and C) showed a
28 cytoplasmic (C, arrow) and nuclear staining (C, arrowhead) of the suprabasal cells and no staining of the
29 basal layer. Caspase-3 staining showed a predominantly cytoplasmic immunolocalization especially in the
30 suprabasal layer (D). Some basal cells showing both, nuclear and cytoplasmic staining (D, arrow).
31 Immunoperoxidase. Scale bars: 50 μ m and 20 μ m.
32
33
34
35

36
37 **Figure 5:** HIF-1 α and caspase-3 immunolabeling in CCOT and DF. HIF-1 α and caspase-3
38 staining showed a similar pattern in CCOT samples (A and B). The DF exhibited weak HIF-1 α and
39 moderate caspase-3 staining in the cytoplasm of the epithelial cells (C and D). Immunoperoxidase. Scale
40 bars: 50 μ m and 20 μ m.
41
42
43

44
45 **Figure 6.** Comparison of HIF-1 α immunostaining between ameloblastoma, KOT, CCOT, RC
46 and DC (A). Comparison of caspase 3 immunostaining between ameloblastoma, KOT, CCOT, RC and
47 DC (B). Dental follicle was used as control. AME-Ameloblastoma; KOT-Keratocystic odontogenic
48 tumor; CCOT-Calcifying cystic odontogenic tumor; RC-Radicular cyst; DC-Dentigerous cyst; DF-Dental
49 follicle. Significance: *p<0.05; **p<0.01; ***p<0.001.
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65




Figure 2

KOT




Figure 3

RC

HIF-1 α

Caspase-3





Figure 4**DC**




Figure 5

Figure 6

A



B





UNIVERSIDADE FEDERAL DO PARÁ
INSTITUTO DE CIÊNCIAS DA SAÚDE
FACULDADE DE ODONTOLOGIA
LABORATÓRIO DE CULTIVO CELULAR

Belém, February 2th, 2016.

Dr Matthias Hanning
Editor
Clinical Oral Investigations
Dr. Hanning,


Please find enclosed the revised manuscript CLOI-D-16-01334 entitled "RELATIONSHIP AMONG HIF-1 α , CASPASE-3 AND CYSTOGENESIS IN ODONTOGENIC CYSTS AND TUMORS".

We were pleased to see that the reviewer recommended publication, suggesting some revisions of the manuscript. All corrections made are highlighted in the manuscript. We thank the reviewers for the interesting suggestions, which have improved the manuscript.

We have included a point-by-point reply to reviewer's concerns comments below and hope the manuscript will be acceptable for publication in Clinical Oral Investigations.

Hoping to hear from you at your earliest convenience, we thank you for your kind attention

Sincerely,



Joao JV Pinheiro
Corresponding author

Reply to reviewer #1

1. Concern of the reviewer

Very interesting approach to research into cyst Formation. Legend of Figure 1 is much too long, should be shortened.

Our response:

The legend of figure 1 (Figure legends, page 11, last paragraph, page 12 first paragraph) was shortened.

Revised text:

“Figure 1: HIF-1 α and caspase-3 immunostaining in ameloblastoma. HIF-1 α staining (A, C, E, G) was predominant in the nucleus (C, arrowhead) and cytoplasm (C, arrow) in epithelial layer near cystic lumen (A, C, asterisk). Intense nuclear and cytoplasmic staining in epithelial islands (E and G, arrow), and in cells surrounding initial cystic formation was observed (E, asterisk). Caspase-3 immunostaining (B, D, F and H) is predominant in the cytoplasm of cells near cystic areas of epithelial islands (B and D, asterisk). Staining is also detected in selected cells of basal layer (D, arrow). In tumor parenchyma, areas of cyst formation (F, asterisk) and solid areas of epithelial islands showed almost an exclusively cytoplasmic staining for caspase-3 (F and H, arrow). Scale bar: 50 μ m and 20 μ m.”

Reply to reviewer #2

1. Concern of the reviewer

Introduction: Adequate. Though, due to the high number of different lesions, the authors are encouraged to differentiate between odontogenic tumors and cysts at this stage of the manuscript.

Our response:

We included a sentence differentiating odontogenic tumors and cysts in the first paragraph of the Introduction section (page 3).

Revised text:

“Odontogenic cysts and tumors are lesions that develop exclusively in or around gnathic bones [1, 2]. They are characterized by the presence of a lining epithelium of odontogenic origin and a cystic area containing liquid and/or a fluid material is often observed. [1, 2]. Due to their high frequency and/or local aggressiveness, ameloblastoma and keratocystic odontogenic tumor (KOT) represent the most clinically relevant odontogenic tumors, and radicular cyst (RC) and dentigerous cyst (DC) the most important odontogenic cysts. Some odontogenic tumors, such as ameloblastoma, may have cystic and solid areas. [2-5]”

2. Concern of the reviewer

Materials and methods: Was the diagnosis of the lesions confirmed by a pathologist before further evaluation?

Our response:

The diagnosis of each lesion was confirmed by a pathologist before evaluation. We now explain that issue in the Samples section of Materials and methods (page 4, second paragraph).

Revised text:

“Diagnosis of each lesion was confirmed by a pathologist before experimental procedures”

3. Concern of the reviewer

From which site of the lesions the samples were taken?

Our response:

The sites selected for quantification were from cystic areas, both odontogenic cists and tumors. Materials and methods, Samples (page 5, first paragraph).

Revised text:

MATERIALS AND METHODS

Samples

Tissue microarrays (TMA) containing 20 ameloblastoma cores (Biomax Inc., Rockville, MD, USA), and tissue sections of odontogenic cysts and tumors were used in this study. Cases of KOT (20), RC (18), DC (11), CCOT (8) with 5 µm thickness completed our sample. Additionally, ten cases of dental follicle (DF) were included as a non-pathological control of normal dental tissue exhibiting neither cystic formation nor neoplastic changes [2]. Samples groups were classified in odontogenic tumors (ameloblastoma, KOT and CCOT) or odontogenic cysts (RC and DC). KOT, RC, DC, CCOT and DF sections were obtained in consecutive cut series from the archives of the department of Oral Pathology of the School of Dentistry of the University Center of Pará (CESUPA, Belém-PA, Brazil). **Cystic areas were selected in both odontogenic cysts and tumors.** Diagnosis of each lesion was confirmed by a pathologist before experimental procedures.

4. Concern of the reviewer

How many examiners did the evaluation procedures? Were they blinded towards the diagnosis? Did the authors calibrate their parameters such as staining intensity, cellular immunolocalization as well as epithelial layer distribution?

Our response:

The samples were evaluated by one examiner in a blinded study towards diagnosis. We included this information in Immunostaining evaluation section of Materials and methods (page 5, first paragraph).

Revised text: The samples were evaluated by for one examiner in a blinded study towards diagnosis.

5. Concern of the reviewer

For evaluation of expression on cysts and tumors, please add that this was conducted descriptively.

Our response:

We included this information in Immunostaining evaluation section of Material and Methods (page 5, first paragraph).

Revised text:

“The evaluation of expression of HIF-1a and caspase 3 in odontogenic cysts and tumors was conducted descriptively.”

6. Concern of the reviewer

Results: For quantification of expression, please add mean/median values as well as standard deviations to the manuscript.

Our response:

The mean and standard deviations were included into manuscript. The Results section, (page 6, 6th and 7th paragraphs, and page 7, first paragraph).

Revised text:

HIF-1 α is highly expressed in ameloblastoma, DC, and comparatively higher in RC when compared to odontogenic tumors

HIF-1 α expression showed variable expression in all samples (Figure 6A). The highest expression was observed in RC (Mean= 58.11; SD=11.61), which was statistically significant when compared to odontogenic tumors (ameloblastoma p=0.026, Mean=44.69, SD=13.87; KOT p<0.0001,; CCOT p<0.0001). Stained area of KOT (Mean=36,20, SD=15.17) and CCOT (Mean=17.60, SD=12.77) was not statistically different from control (Mean=23.07, SD=6.35), and HIF-1 α expression in CCOT was lower when compared with all other lesions (ameloblastoma p=0.0004; KOT p=0.0368; DC p=0.0005, Mean=46,25, SD=13.50; and RC p<0.0001). Although there was not difference between HIF-1 α immunostaining between control and CCOT, CCOT expression was lower than all the other studied lesions.

Caspase-3 is highly expressed in odontogenic cysts compared to odontogenic tumors.

Results of caspase-3 immunostaining quantification demonstrated that odontogenic cysts showed higher percentage of stained areas than odontogenic tumors (Figure 6B). DC (Mean=65,57, SD=13.86) showed the highest expression of caspase-3, followed by RC (Mean=60.23; SD=7.58), ameloblastoma (Mean=50.16, SD=6.63) and KOT (Mean=49.44, SD=8.23). Comparison between odontogenic cysts and tumors revealed higher expression in DC when compared to ameloblastoma (p=0.0005), KOT (p=0.0002) and CCOT (p<0.0001, Mean=43.32, SD=13.14) and also higher expression in RC against the same ameloblastoma (p=0.0175), KOT (p=0.0086) and CCOT (p<0.0016). No differences were observed when compared odontogenic cysts or tumors with each other. Caspase-3 expression was significantly higher in all studied samples (p<0.0001) when compared to the non-pathological control of DF (Mean=13.82, SD=10.53).

7. Concern of the reviewer

Did the authors try to add mean values from either tumors as well as cysts and compare them to each other?

Our response:

After labeling quantification, we found the mean values for each odontogenic cysts and tumors sample, and compared to each other, as demonstrated on Figure 6. Asterisks in Figure 6 indicate which of these comparisons were significant.

Revised text: Not applicable

8. Concern of the reviewer

Discussion: Adequate. Is there a clinical implication of these findings?

Our response:

We think that there is clinical implications of HIF-1^o and caspase-3 expression in odontogenic cysts and tumors. We commented this issue on the last paragraph of the section.

Revised text:

“Therefore, HIF-1 α and caspase-3 expression in odontogenic cysts and tumors may be important to explain which cellular components are related to the behavior of these lesions. Additionally, these molecules can be used as biological markers in the future, serving as instruments for better diagnosis and treatment planning.”