

Roles of the Host Oxidative Immune Response and Bacterial Antioxidant Rubrerythrin during *Porphyromonas gingivalis* Infection

Piotr Mydel^{1,2}^{✉a}, Yusuke Takahashi^{2,3}^{✉a}, Hiromichi Yumoto^{2,4}, Maryta Sztukowska^{1,5}, Malgorzata Kubica¹, Frank C. Gibson III², Donald M. Kurtz, Jr.⁶, Jim Travis⁵, L. Vincent Collins⁷, Ky-Anh Nguyen⁵, Caroline Attardo Genco²^{✉b*}, Jan Potempa^{1,5}^{✉b*}

1 Department of Microbiology, Faculty of Biochemistry, Biophysics, and Biotechnology, Jagiellonian University, Krakow, Poland, **2** Department of Medicine, Boston University School of Medicine, Boston, Massachusetts, United States of America, **3** Department of Oral Microbiology, Kanagawa Dental College, Yokosuka, Kanagawa, Japan, **4** Department of Conservative Dentistry, The University of Tokushima, School of Dentistry, Tokushima, Japan, **5** Department of Biochemistry and Molecular Biology, University of Georgia, Athens, Georgia, United States of America, **6** Department of Chemistry and Center for Metalloenzyme Studies, University of Georgia, Athens, Georgia, United States of America, **7** Department of Rheumatology and Inflammation Research, University of Göteborg, Göteborg, Sweden

The efficient clearance of microbes by neutrophils requires the concerted action of reactive oxygen species and microbicidal components within leukocyte secretory granules. Rubrerythrin (Rbr) is a nonheme iron protein that protects many air-sensitive bacteria against oxidative stress. Using oxidative burst-knockout (NADPH oxidase-null) mice and an *rbr* gene knockout bacterial strain, we investigated the interplay between the phagocytic oxidative burst of the host and the oxidative stress response of the anaerobic periodontal pathogen *Porphyromonas gingivalis*. Rbr ensured the proliferation of *P. gingivalis* in mice that possessed a fully functional oxidative burst response, but not in NADPH oxidase-null mice. Furthermore, the *in vivo* protection afforded by Rbr was not associated with the oxidative burst responses of isolated neutrophils *in vitro*. Although the phagocyte-derived oxidative burst response was largely ineffective against *P. gingivalis* infection, the corresponding oxidative response to the Rbr-positive microbe contributed to host-induced pathology via potent mobilization and systemic activation of neutrophils. It appeared that Rbr also provided protection against reactive nitrogen species, thereby ensuring the survival of *P. gingivalis* in the infected host. The presence of the *rbr* gene in *P. gingivalis* also led to greater oral bone loss upon infection. Collectively, these results indicate that the host oxidative burst paradoxically enhances the survival of *P. gingivalis* by exacerbating local and systemic inflammation, thereby contributing to the morbidity and mortality associated with infection.

Citation: Mydel P, Takahashi Y, Yumoto H, Sztukowska M, Kubica M, et al. (2006) Roles of the host oxidative immune response and bacterial antioxidant rubrerythrin during *Porphyromonas gingivalis* infection. PLoS Pathog 2(7): e76. DOI: 10.1371/journal.ppat.0020076

Introduction

Phagocytic leukocytes, especially neutrophils, play a critical role in innate immune responses against bacteria, fungi, and other pathogens [1]. Neutrophil-mediated bacterial killing can involve both oxygen-independent and oxygen-dependent processes. Oxygen-independent microbial killing relies on the contents of three cytoplasmic granule subsets, the azurophilic (primary), specific (secondary), and gelatinase granules. After fusing with phagosomes, these granules deliver antimicrobial proteins and peptides, such as defensins, bactericidal/permeability-increasing protein, azurocidin, cathelicidin, and lysozyme, all of which are capable of damaging the bacterial cell envelope. In addition, several proteinases, such as neutrophil elastase and cathepsin G, facilitate bacterial killing by digestion of bacterial outer membrane proteins [2], surface appendages [3], and virulence factors [4]. Oxygen-independent microbial killing also appears to function efficiently extracellularly. Upon activation, neutrophils release granule proteins and chromatin, which together form extracellular fibers. Brinkman et al. [5] have reported that these fibers become saturated with high concentrations of bactericidal peptides, proteins, and proteases, and are able to trap and efficiently kill invading bacteria that become entangled in the fiber meshwork. The

significance of this oxygen-independent mechanism is exemplified by recurrent infections associated with two rare inherited diseases, Chediak-Higashi syndrome and specific granule deficiency, which are characterized by insufficient release of antimicrobial components and the absence of some antimicrobial components in specific and/or azurophilic granules, respectively [6].

Editor: Daniel Portnoy, University of California Berkeley, United States of America

Received: December 7, 2005; **Accepted:** June 21, 2006; **Published:** July 28, 2006

DOI: 10.1371/journal.ppat.0020076

Copyright: © 2006 Mydel et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abbreviations: CGD, chronic granulomatous disease; *Cybb*^{-/-}, NADPH oxidase-null; ELISA, enzyme-linked immunosorbent assay; iNOS, inducible nitric oxide synthase; LPS, lipopolysaccharide; MPO, myeloperoxidase; Rbr, rubrerythrin; RNS, reactive nitrogen species; ROS, reactive oxygen species

* To whom correspondence should be addressed. E-mail: caroline.genco@bmc.org (CAG); potempa@uga.edu (JP)

^{✉a} These authors contributed equally to this work.

^{✉b} These authors contributed equally to this work.

[✉] Current address: Department of Biochemistry and Molecular Biology, University of Georgia, Athens, Georgia, United States of America

Synopsis

The physiological role of neutrophils is to seek out and destroy invading microbes. Professional phagocytes engulf (phagocytose) these organisms and kill them using bactericidal peptides, enzymes, toxic reactive oxygen species, and reactive nitrogen species produced by neutrophils and macrophages. Unfortunately, the reactive oxygen species unleashed in an oxidative burst response can cause considerable collateral damage and are directly responsible for infection-associated tissue injuries, especially if the invaders are protected against killing by neutrophils. The authors investigated the pathogenesis of *Porphyromonas gingivalis*, an anaerobic bacterium that is responsible for human periodontal disease and is protected against oxidative stress by the cytoplasmic protein rubrerythrin. We show that *P. gingivalis* is not only resistant to reactive oxygen species, but that in mice, rubrerythrin shields the bacterium against reactive nitrogen species. These features allow *P. gingivalis* to proliferate in animals that possess a fully functional oxidative burst response. Furthermore, we demonstrate that the neutrophil oxidative burst response, rather than eliminating the bacteria, exacerbates disease by damaging host tissues and facilitating growth and systemic dissemination of the pathogen. Collectively, this study provides important information on how oxygen-dependent killing mechanisms operate during anaerobic infection and on the role of rubrerythrin in protecting against a pathogenic anaerobic organism, while emphasizing the importance of limiting host-mediated tissue injury in inflammatory diseases caused by bacteria.

Oxygen-dependent killing is initiated by the assembly of the NADPH oxidase complex at the phagosome membrane [7]. The NADPH oxidase system is required for immunocompetence against bacteria via the so-called respiratory or oxidative burst, which generates reactive oxygen species (ROS) that are toxic for microbes [8,9]. The importance of this oxygen-dependent killing is exemplified by patients with chronic granulomatous disease (CGD), who carry a dysfunctional NADPH oxidase in their phagocytes and often suffer from recurrent infections [8]. However, patients with CGD rarely have infections that involve anaerobic bacteria [2,6,10,11], which suggests that oxygen-dependent bacterial killing mechanisms are not utilized for the clearance of anaerobic organisms. Furthermore, recent studies have demonstrated a link between the formation of superoxide anion and the activation of granular microbicidal enzymes, which suggests a new paradigm for the molecular mechanisms utilized by neutrophils to kill their targets [12,13]. Thus, the oxidative burst is a prerequisite for the mobilization of cationic neutrophil elastase and cathepsin G, which are stored in granules complexed with the strongly anionic sulfated proteoglycan matrix. In addition, one of the major bactericidal peptides of neutrophils is released from cathelicidins by the action of neutrophil proteases [14]. Thus, it is clear that the elimination of invading bacteria by phagocytes is a complicated and highly coordinated process.

The importance of the orchestrated activation and recruitment of neutrophils for removing bacterial pathogens is clearly evident in the oral cavity during periodontal disease, where the innate immune system responds to an abundant and versatile microflora that matches in mass and complexity those residing in the lower gastrointestinal tract. The tenuous balance between healthy and diseased states within the gingival sulcus is maintained primarily by neutrophils. Even

minor, inherited, or acquired malfunctions of neutrophils, but not of other leukocytes, invariably result in severe inflammation of the gingiva (gingivitis), while more pronounced defects in phagocyte function predispose to rapid destruction of the periodontium (manifested as periodontal disease) and subsequent tooth loss [10,15,16]. The onset and progression of periodontal disease is associated with the proliferation of specific bacterial species, particularly anaerobes, which include *Porphyromonas gingivalis*, *Haemophilus (Actinobacillus) actinomycetemcomitans*, *Treponema denticola*, and *Tannerella (Bacteroides) forsythensis* [17]. *P. gingivalis* has been implicated as the major pathogen associated with adult periodontal disease. In addition, this organism is also associated with localized aggressive periodontitis, which is a more aggressive and destructive form of periodontal disease associated with impaired neutrophil chemotaxis [18]. Patients with localized aggressive periodontitis are otherwise in good general health and are not predisposed to extraoral microbial infections, further supporting the crucial importance of fully functional neutrophils in maintaining homeostasis in the gingival tissues.

P. gingivalis possesses an array of virulence factors, including lipopolysaccharide (LPS), fimbriae, hemagglutinins, hemolysins, and proteolytic enzymes, which facilitate colonization and the initiation of infection [19]. *P. gingivalis* also contains enzymes that have been implicated in protection against oxidative stress, such as an excision/repair system for removing oxidatively damaged DNA bases [20], superoxide dismutase [21], and rubrerythrin (Rbr) [22]. Since the *P. gingivalis* strain that lacks the gene for rubrerythrin (*rbr*⁻) is more dioxygen sensitive and hydrogen peroxide sensitive than the wild-type strain and shows growth inhibition in liquid cultures when exposed to oxidants, it has been proposed that Rbr protects *P. gingivalis* against aerobic stress by functioning as a cytoplasmic peroxidase that reduces hydrogen peroxide to water [22]. Rbr appears to be involved in protection against oxidative stress in several anaerobic bacteria and archaeobacteria, most likely functioning as the terminal component of an NAD(P)H peroxidase [23–25]. In addition, *P. gingivalis* accumulates a hemin layer that is purported to provide oxidative stress protection [26].

The manner in which these bacterial oxidative stress protection systems function to protect anaerobic organisms against the oxygen-dependent bactericidal activity of neutrophils has not been well defined. Using *P. gingivalis* as a model organism for anaerobic infection, we initially investigated the role of NADPH oxidase-dependent mechanisms in the responses of mice to infection with anaerobic bacteria. Subsequently, we examined the contribution of Rbr to the protection of this anaerobic organism against the host oxidative burst. Our results indicate that the host oxygen-dependent bactericidal system is not only ineffective in combating *P. gingivalis* infection, but appears to exacerbate significantly the host tissue damage induced by the infection. Furthermore, we show that Rbr plays an important protective role for the bacteria in the presence of a fully functional host immune response. This function of Rbr is unrelated to the neutrophil oxidative burst and is manifested as the shielding of *P. gingivalis* against reactive nitrogen species (RNS).




Figure 1. Viability of Mice Inoculated with *P. gingivalis*

The data shown are the percentages of mice ($n = 7$ for each group) that were alive on the indicated days after challenge with 100- μ l volume, 1×10^9 CFU *P. gingivalis* wild-type (W83) or Rbr⁻ mutant (Rbr⁻). Control animals were inoculated with sterile broth. Wild-type (C57BL/6J) and Cybb^{-/-} mice were utilized.

DOI: 10.1371/journal.ppat.0020076.g001

Results

Rbr Is Important for *P. gingivalis* Virulence

Since Rbr protects *P. gingivalis* against oxidative stress in vitro [22], we investigated whether it would protect *P. gingivalis* against oxidative stress during infection in an animal model. For these initial studies, we utilized a mouse subcutaneous chamber model in which virulent strains of the bacterium were able to disseminate from an initial infection site and cause high mortality and morbidity [27,28]. Subcutaneous chambers implanted into C57BL/6J mice were inoculated with either a lethal dose (100- μ l volume, 1×10^9 CFU/ml) of *P. gingivalis* W83 or the same dosage of the isogenic Rbr⁻ strain, and the mice were monitored daily for general health condition and the emergence of skin abscesses. Mice infected with the wild-type *P. gingivalis* strain exhibited severe cachexia, with ruffled hair, hunched bodies, and weight loss, as well as severe skin lesions localized on the ventral body surface. All seven of the animals infected with wild-type *P. gingivalis* died by day 8 after infection (Figure 1). In contrast, two of the seven mice that were inoculated with the *P. gingivalis* Rbr⁻ strain succumbed to infection on days 5 and 12 after infection, while the five remaining mice appeared healthy and resembled the unchallenged animals.

To investigate whether the higher survival rates of animals inoculated with the *P. gingivalis* Rbr⁻ strain were the result of altered bacterial growth rates in vivo, the bacterial numbers in the chambers were determined on days 1, 3, 6, and 10 following bacterial challenge. Following infection with the *P. gingivalis* Rbr⁻ strain, the number of organisms recovered from the chamber fluid samples gradually increased to a level on day 10 after inoculation that was approximately 100-fold higher than that of the original inoculum. The numbers of bacteria in the samples collected from the chambers of C57BL/6J mice infected with the wild-type *P. gingivalis* strain increased ~50-fold on day 1, increased further to 3.5×10^{11} CFU/ml on day 3, and reached a level that was 300-fold higher than that administered initially (100- μ l volume, 1×10^9 CFU/ml) (Figure 2). These data indicate that Rbr expression is important for *P. gingivalis* proliferation in subcutaneous chambers.




Figure 2. Bacterial Viability in Chamber Fluid Samples

Subcutaneous chambers were inoculated on day 0 with 100- μ l volume, 1×10^9 CFU/ml of the *P. gingivalis* strains. The CFU/ml values were determined by counting bacterial colonies on anaerobic agar plates. * $p < 0.001$ for data on day 3; Mann-Whitney *U* test ($n = 7$ mice for each group). Increased numbers of *P. gingivalis* were detected on the third day in C57BL/6J mice inoculated with strain W83 (wild-type) as compared to the Cybb^{-/-} mice and challenge with the *P. gingivalis* Rbr⁻ strain. DOI: 10.1371/journal.ppat.0020076.g002

Oxidative Burst Does Not Contribute to the Killing of *P. gingivalis* by Human or Mouse Neutrophils

To investigate the contribution of the neutrophil oxidative burst to the elimination of *P. gingivalis*, we compared the rates of killing of this bacterium by neutrophils that were isolated from wild-type and NADPH oxidase-null (Cybb^{-/-}) mice. Surprisingly, the rate of *P. gingivalis* killing by normal bone marrow murine neutrophils was indistinguishable from that by oxidative burst-deficient neutrophils. The killing by both cell types was relatively slow, with 50% and 10% viable bacterial cells persisting after 45 and 195 min of coinoculation with neutrophils, respectively (Figure 3A). Significantly, except for the shorter periods of incubation, there was no difference in the rates of killing of the wild-type *P. gingivalis* by the normal mouse neutrophils under oxic and anoxic conditions (Figure 3B), where the production of ROS has been reported to be diminished under anoxic conditions [29]. From a comparison of the killing of *P. gingivalis* wild-type and Rbr⁻ strains under aerobic conditions (Figure 3C), it is clear that the elimination of viable bacteria occurs with the same kinetics regardless of the presence or absence of Rbr. The human peripheral blood neutrophils killed *P. gingivalis* more efficiently than the mouse neutrophils, but again, there was no difference between the rates of killing of the wild-type and Rbr⁻ strains (data not shown). Taken together, our results indicate that Rbr does not contribute to the resistance of *P. gingivalis* to the oxygen-dependent bactericidal activity of neutrophils and that the host oxidative burst does not play a major role in the elimination of *P. gingivalis*. These data corroborate the clinical observation that patients with CGD are not unusually susceptible to periodontal diseases despite the lack of oxygen-dependent bactericidal activities of their neutrophils [10], and support the hypothesis that antibacterial peptides and proteins potentially constitute the major armamentarium used by neutrophils to kill *P. gingivalis* [30,31].

Cybb^{-/-} Mice Are Resistant to *P. gingivalis* Infection

To investigate further the involvement of the oxygen-dependent antimicrobial defense mechanism against systemic infection by anaerobic bacteria, we evaluated the suscepti-




Figure 3. *P. gingivalis* Killing by Neutrophils

Murine bone marrow-derived neutrophils isolated from wild-type and *Cybb*^{-/-} animals were incubated with opsonized *P. gingivalis* (wild-type or *Rbr*⁻ strain) at a ratio of 1:5 ratio in an aerobic or anaerobic atmosphere. At the indicated timepoints, samples were withdrawn and plated, and the numbers of bacteria were determined. The CFU value at time zero was taken as 100%.

(A) Wild-type *P. gingivalis* incubated aerobically with neutrophils derived from wild-type (C57BL/6J) and *Cybb*^{-/-} mice.

(B) Wild-type *P. gingivalis* incubated aerobically or anaerobically with neutrophils from wild-type (C57BL/6J) mice.

(C) Wild-type *P. gingivalis* and the *Rbr*⁻ strain incubated aerobically with neutrophils from wild-type (C57BL/6J) mice.

DOI: 10.1371/journal.ppat.0020076.g003

bility of *Cybb*^{-/-} mice to intraperitoneal chamber inoculation with a lethal dose (100- μ l volume, 1×10^9 CFU/ml) of *P. gingivalis*. Interestingly, we found that *Cybb*^{-/-} mice were resistant to infection with either the wild-type or *P. gingivalis* *Rbr*⁻ strains, as evidenced by the survival of all the mice following systemic challenge up to the end of the experiment (Figure 1). Correlating this difference in survival relative to that seen for C57BL/6J mice, a distinct difference in the kinetics of in vivo bacterial growth was observed between the C57BL/6J and *Cybb*^{-/-} mice. In the C57BL/6J mice, an abrupt increase in bacterial numbers occurred 1 d after infection with *P. gingivalis* W83, and the bacterial numbers increased further by day 3 (Figure 2), at which point ~70% of the animals had died (Figure 1). Conversely, in *Cybb*^{-/-} mice,

although the bacterial numbers reached the same level as in the wild-type animals on day 1, the bacterial numbers did not increase during the subsequent days of infection (Figure 2).

The *P. gingivalis* *Rbr*⁻ strain was detected in lower numbers than the wild-type strain in the chamber fluid samples at day 1. Following a significant increase from day 1 to day 3, the number of *Rbr*⁻ bacteria within the chamber fluid samples remained constant over the 10-d infection period (Figure 2). Despite the increased sensitivity of the *Rbr*⁻ strain to oxidative stress in vitro [22], it seems unlikely that the lack of *Rbr* is responsible for the retarded growth of the mutant strain in vivo since *P. gingivalis* is apparently killed by neutrophils in an oxygen-independent manner, for which *Rbr* has no protective function. Furthermore, the presence of *Rbr* does not affect elimination of the bacterium by fully functional neutrophils that respond to the phagocytosed microbes.

RNS Contribute to *P. gingivalis* Killing by Macrophages, and *Rbr* Exerts a Protective Effect against iNOS-Dependent Bactericidal Activity

The different kinetics of *Rbr*⁻ and wild-type *P. gingivalis* growth in vivo (Figure 2) suggests that *Rbr* provides protection against the oxygen-independent antimicrobial activity of the host. This activity is elicited mainly by either bactericidal peptides/proteins or RNS. RNS are generated in macrophages, monocytes, and epithelial cells by inducible nitric oxide synthase (iNOS) in response to bacterial infection. Therefore, we compared the susceptibilities of the *P. gingivalis* wild-type strain and *Rbr*⁻ mutant to killing after phagocytosis by explanted mice peritoneal macrophages in the absence or presence of the specific iNOS inhibitor L-NAME. Of note, L-NAME at the concentration used did not affect proteolytic activity of gingipains R, Arg-Xaa peptide bond-specific proteases, which are considered the major virulence factor of *P. gingivalis*. The significantly higher rates of killing of the wild-type strain (Figure 4A) and the *Rbr*⁻ mutant (Figure 4B) by macrophages with active iNOS underline the importance of RNS in *P. gingivalis* elimination and indicate that *Rbr* provides protection against RNS. This conclusion is validated by a direct comparison of the bacterial CFU in the macrophage lysates after 24 h of phagocytosis (Figure 4C). In summary, this finding corroborates well with the observed in vivo differences in pathological changes in mice that were inoculated with the wild-type strain and *Rbr* mutant in normal and oxidative burst-deficient animals.

Growth of Wild-Type *P. gingivalis* in C57BL/6J Mice Is Associated with the Release of Large Quantities of LPS

The killing of Gram-negative bacteria in vivo may lead to the release of LPS [32], which can cause an aggravated septic shock. Therefore, we compared the levels of soluble LPS in the chamber fluids and sera of mice that were challenged with *P. gingivalis* (Figure 5). In the chamber fluid samples from C57BL/6J mice that were inoculated with the wild-type *P. gingivalis* strain, the concentration of LPS increased from the baseline (10 EU/ml) on day 1 to 600 EU/ml on day 3 (Figure 5A). Only moderately elevated levels of LPS (130 EU/ml) were detected in the chamber fluids from C57BL/6J mice that were infected with the *Rbr*⁻ mutant on days 3 and 6 after inoculation, and these levels increased further to 230 EU/ml




Figure 4. *P. gingivalis* Killing by Murine Macrophages in the Presence and Absence of Nitric Oxide

Murine macrophages were incubated with opsonized *P. gingivalis* (the wild-type or Rbr⁻ strain) at a ratio of 1:10 in the presence or absence of iNOS inhibitor (L-NAME). At the indicated timepoints, samples were withdrawn and plated, and the numbers of bacteria were determined. The CFU value at time zero was taken as 100%.

(A) Wild-type *P. gingivalis* incubated with macrophages in the presence or absence of L-NAME. (B) Rbr⁻ strain incubated with macrophages in the presence or absence of L-NAME.

(C) Representative pictures of *P. gingivalis* W83 (left panel) and Rbr⁻ (right panel) colonies from the 24-h timepoint grown on anaerobic blood agar plates.

In (A) and (B), statistical significance of the differences in the CFU values was calculated using the Mann-Whitney *U* test (**p* < 0.05; ***p* < 0.01).

DOI: 10.1371/journal.ppat.0020076.g004




Figure 5. LPS Levels in Chamber Fluid Samples and/or Sera of C57BL/6J and Cybb^{-/-} Mice during the Course of the Experiment

(A) LPS levels in the chamber fluid samples of C57BL/6J and Cybb^{-/-} mice. On day 0, the two mouse strains were inoculated with 100- μ l volume, 1×10^9 CFU/ml *P. gingivalis* strain W83 or Rbr⁻ mutant strain.

(B) LPS levels in the chamber fluid samples and sera of mice on day 3 after challenge. The mice were challenged with a lethal dose (100- μ l volume, 1×10^9 CFU/ml) of *P. gingivalis* strain W83. The data are presented as the means \pm SD from the results obtained for three mice per group. **p* < 0.05; ***p* < 0.01; as determined by Student's *t* test.

DOI: 10.1371/journal.ppat.0020076.g005




Figure 6. Cytokine Levels in the Sera of C57BL/6J and Cybb^{-/-} Mice after Challenge with the *P. gingivalis* Wild-Type (W83), the Rbr Mutant (Rbr⁻) Strain, and Sterile, Noninoculated Anaerobic Broth (Broth)

The subcutaneous chambers in the wild-type (C57BL/6J) and Cybb^{-/-} mice were inoculated with 100- μ l volume, 1×10^9 CFU/ml *P. gingivalis* or with broth on day 0. The IL-6 (A) and TNF- α (B) levels were measured 1, 3, 6, and 10 d after inoculation. No statistically significant differences in cytokine levels are evident between the wild-type and Cybb^{-/-} mice following *P. gingivalis* challenge. The data shown are the median and interquartile range for each group of mice ($n = 7$) throughout the course of the experiment. Comparisons are made using the Mann-Whitney *U* test ($p < 0.05$). DOI: 10.1371/journal.ppat.0020076.g006

on day 10. The LPS levels in the chamber fluid samples from the oxidative burst-deficient mice increased transiently on day 3 and returned to the baseline level by day 6 (Figure 5A). In addition, high levels of LPS were detected in the sera of C57BL/6J mice inoculated with the wild-type *P. gingivalis* strain (Figure 5B). In the case of Cybb^{-/-} mice inoculated with the wild-type strains of *P. gingivalis* (Figure 5B) or C57BL/6J mice inoculated with the Rbr⁻ strain, the concentrations of LPS in the sera did not differ significantly from the background levels (unpublished data). These results suggest that the respiratory burst is responsible for LPS release from bacteria, both in the chamber fluid samples and in infected tissues, and especially in the blood.

Lack of Cytokines in the Chamber Fluid Samples Following *P. gingivalis* Infection

We next evaluated the levels of the proinflammatory cytokines TNF- α and IL-6 in chamber fluid and serum samples following challenge with *P. gingivalis*. Surprisingly and contrary to a previous report [33], no significant concentrations of these cytokines were detected in the chamber fluid samples following infection with either the wild-type or the Rbr⁻ strain (unpublished data). This discrepancy can be attributed to the degradation of cytokines by cysteine proteases (gingipain R and gingipain K) produced by *P. gingivalis*. Several studies have shown that IL-1 β , TNF- α , and IL-6 are efficiently degraded by gingipains, both in vivo and in vitro [34–36]. This assumption is supported by the high levels of gingipain R and gingipain K activity seen in the chamber fluid samples from *P. gingivalis*-infected mice. We also found that the levels of gingipain activity correlated well ($p < 0.005$) with CFU counts (unpublished data), confirming the production of gingipains in vivo.

Systemic Cytokine Responses to *P. gingivalis* Infection Do Not Correlate with Levels of LPS Release

We measured the levels of specific cytokines in serum samples obtained from *P. gingivalis*-infected mice. With the exception of IL-6 in wild-type mice inoculated with *P. gingivalis*, both the IL-6 and TNF- α levels increased significantly above the baseline levels in mice of either genotype infected with either the wild-type or the Rbr⁻ *P. gingivalis*

strain (Figure 6). However, there was no correlation between the cytokine responses and LPS concentrations (unpublished data). Cybb^{-/-} mice inoculated with the *P. gingivalis* strain or normal mice infected with the Rbr⁻ mutant, although showing a background level of LPS in the serum (Figure 5), elicited similar amounts of IL-6 (Figure 6A) and TNF- α (Figure 6B) compared with normal mice inoculated with the wild-type strain and having a high concentration of LPS in the serum. These results suggest that the increased cytokine production in animals challenged with *P. gingivalis* is at least partially independent of LPS release.

Wild-Type and Rbr⁻ *P. gingivalis* Strains Elicit a Cellular Inflammatory Response

In agreement with previous studies [27,28,33], we observed that challenging mice with *P. gingivalis* via subcutaneously implanted chambers elicited a robust cellular inflammatory response. In addition, we found that the total leukocyte counts recovered in chamber fluid samples were similar and remained at the same level throughout the experiment, regardless of the mouse genotype or the strain of *P. gingivalis* used (Figure 7A). In corroboration of other reports [27,28,33], the differential cell count analysis indicated that the predominant inflammatory cells in the chamber fluid samples following *P. gingivalis* challenge were neutrophils. We did not observe significant differences, either qualitatively or quantitatively, in the levels of infiltration of inflammatory cells into the chambers between C57BL/6J and Cybb^{-/-} mice regardless of whether they were challenged with the wild-type or Rbr⁻ strain (unpublished data). In addition, within each group of animals the numbers of live and dead cells observed 3 days after bacterial challenge were similar, regardless of the strain used for inoculation (Figure 7B). Nonetheless, in the case of C57BL/6J mice inoculated with wild-type *P. gingivalis*, the influx of neutrophils was apparently insufficient to contain the infection.

Since myeloperoxidase (MPO) can be considered a marker of local or systemic neutrophil activation [37–48] and is a good indicator of the rate of neutrophil turnover, we determined the concentration of this enzyme in the chamber fluids and sera of mice inoculated with a lethal dose of wild-type *P. gingivalis*. As expected, MPO was present in the chamber fluid samples collected on day 3 after inoculation, with the enzyme




Figure 7. Inflammatory Cell Infiltrates in Chamber Fluid Samples of Mice Challenged with *P. gingivalis*

(A) The total cell counts are similar between the groups ($n=7$), as determined by trypan blue staining and cell counting ($p > 0.05$ for the course of the experiment; Mann-Whitney *U* test). In both wild-type and *Cybb*^{-/-} mice, the cell numbers in the chambers inoculated with the sterile *P. gingivalis* cultivation broth were similar (0.5×10^7 cell/ml) throughout the course of the experiment (not shown in the figure).

(B) The numbers of live and dead cells in the chamber fluid on day 3 after inoculation.

(C) MPO was measured by ELISA. The chamber fluid and serum MPO levels in the wild-type *P. gingivalis*-infected mice on day 3 after inoculation are approximately four times higher than the corresponding levels in the *P. gingivalis* *Rbr*⁻-inoculated animals ($n = 7$, pooled samples). Control animals inoculated with sterile, noninoculated anaerobic broth had very low levels of MPO (data not shown).

(D) Both the wild-type and *Rbr*-deficient strains of *P. gingivalis* cause strong activation of the NADPH oxidase in the mouse bone marrow neutrophils. No statistically significant differences were noted between the strains.

DOI: 10.1371/journal.ppat.0020076.g007

concentration being four-fold higher in the samples obtained from the wild-type mice than from *Cybb*^{-/-} mice inoculated with wild-type *P. gingivalis*. In the case of inoculation with the *rbr*⁻ strain, the difference in the MPO level in the chamber fluid between wild-type and *Cybb*^{-/-} mice was about two-fold (Figure 7C). It seems likely that the higher level of MPO in the chamber fluid from wild-type mice was not due to increased lysis of the neutrophils, since the numbers of dead leukocytes were much higher in *Cybb*^{-/-} mice than in wild-type mice (Figure 7B). This observation suggests that in the absence of oxidative burst, the increased influx of leukocytes into the chambers is followed by cell death.

Significant quantities of MPO were detected exclusively in the serum samples from wild-type mice inoculated with wild-type *P. gingivalis* (Figure 7C). This observation suggests that systemic activation of neutrophils or increased turnover of these cells is caused only by the wild-type bacteria present in the circulation of wide-type mice. Notably, both the *rbr*⁻ mutant and the parental *P. gingivalis* strain triggered the same level of NADPH oxidase activation in murine bone marrow neutrophils (Figure 7D). Therefore, to elucidate further the profound differences in the host responses to *P. gingivalis* strains, as illustrated by the serum MPO levels, we compared using flow cytometry analysis of the status of neutrophils in the whole blood of C57BL/6J mice that were infected with either the wild-type or the *rbr*⁻ *P. gingivalis* strain. Staining the blood for Ly-6G, which is an antigen that is expressed constitutively on mice leukocytes [49], revealed an approximately 2.5-fold increase in neutrophils in animals infected with *P. gingivalis* W83 compared to the control (uninfected)

mice or those inoculated with the *rbr*⁻ mutant (Figure 8). Furthermore, neutrophils from mice infected with wild-type *P. gingivalis* presented significantly less L-selectin, a marker for neutrophil activation, on their surfaces than neutrophils from the other groups of animals (Figure 8). Collectively, the flow cytometry data corroborate the MPO assay results and indicate that mouse infection with wild-type *P. gingivalis*, but not with the *rbr*⁻ strain, triggers an increase in the number of circulating neutrophils and boosts the systemic activation of these cells. In the bloodstream, the encounter between wild-type *P. gingivalis* and neutrophils that have a functional oxidative burst probably leads to the release of large quantities of LPS (Figure 5). It is tempting to speculate that in addition to disseminated infection and colonization of other organs (see below) systemic activation of neutrophils was responsible for the observed mouse mortality.

Impaired Clearance of *P. gingivalis* from Chambers Leads to Systemic Dissemination of the Infection

We have previously determined that the formation of skin lesions in C57BL/6J mice challenged with *P. gingivalis* indicates systemic spread of the bacterium [28]. This observation was confirmed in a separate experiment in which C57BL/6J and *Cybb*^{-/-} mice were killed on day 3 after inoculation with the wild-type *P. gingivalis* strain, and the numbers of bacteria present in the chamber fluid samples and blood drawn from the heart and extracts of the spleen, liver, lung, and kidney were determined (Figure 9). Every tissue examined from the C57/BL6J mice infected with the wild-type *P. gingivalis* strain was colonized with *P. gingivalis*,




Figure 8. Challenge with *P. gingivalis* W83 Strain Elicits Potent Neutrophil Recruitment and Activation in the Peripheral Blood Compared to the Rbr⁻ Strain

The top row (dotplots) shows the flow cytometric analysis of blood cells stained with Ly-6G and CD62L 3 d after chamber inoculation with (A) PBS, (B) the Rbr⁻ strain, and (C) the W83 strain. In the case of mice infected with the W83 strain, the number of neutrophils increased five-fold compared to the number of neutrophils in mice infected with the Rbr⁻ strain (gated events). The histograms in the bottom row exhibit a two-fold decrease in CD62L expression for peripheral blood neutrophils collected from (F) W83-challenged mice in comparison to (D) control and (E) the *rbr*⁻ mutant strain-infected animals. Black line, isotypic control.

DOI: 10.1371/journal.ppat.0020076.g008




Figure 9. Bacteremia in Wild-Type and *Cybb*^{-/-} Mice 3 d after Challenge with the *P. gingivalis* Wild-Type Strain

(A) Representative pictures of *P. gingivalis* colonies grown on anaerobic blood agar plates were taken using samples from the hearts, spleens, livers, lungs, and kidneys of C57BL/6J and *Cybb*^{-/-} mice, as indicated, 3 d after challenge with 100- μ l volume, 1×10^9 CFU/ml of *P. gingivalis* strain W83. (B) Increased numbers of *P. gingivalis* are present in the chamber fluid samples obtained from the C57BL/6J mice compared to the knockout *Cybb*^{-/-} mice.

(C) In contrast to the C57BL/6J mice, no bacteria are detected in blood samples collected from the hearts of challenged *Cybb*^{-/-} mice. The values (CFU/ml) were determined by counting bacterial colonies on anaerobic agar plates ($n = 3$ mice). The data shown are the median (thick horizontal bars) and interquartile range for each group of mice. * $p < 0.05$; Mann-Whitney *U* test.

DOI: 10.1371/journal.ppat.0020076.g009

indicating that bacteremia led to the colonization of several organs (Figure 9A). The robust proliferation of *P. gingivalis* at these sites may have contributed to the observed mouse mortality following challenge with the wild-type *P. gingivalis* strain. In contrast, in the case of the *Cybb*^{-/-} mice, culturing of the blood, spleen, and liver extracts gave negative results, while only a few sporadic colonies were detected in the lung tissue extract (Figure 9A).

The Exuberant Proliferation of *P. gingivalis* in Wild-Type Mice Is Unrelated to the Generation of Growth-Promoting Nutrients by the Oxidative Burst

To investigate the possibility that the presence of the NADPH oxidase in host cells allows more robust growth of *P. gingivalis* by providing essential growth factors derived from the breakdown products of oxidized proteins and fatty acids (e.g., C5a and arachidonate, respectively), we compared the levels of bacterial growth in media supplemented with the inflammatory exudates elicited from wild-type and *Cybb*^{-/-} mice by thioglycollate treatment. The growth of the wild-type *P. gingivalis* strain was similar in both media (Figure 10), which suggests that the systemic dissemination of *P. gingivalis* in wild-type mice is not due to NADPH oxidase-dependent generation of growth factors.

Rbr Expression by *P. gingivalis* Leads to Increased Oral Bone Loss Following Infection

We also examined the pathogenic potential of the wild-type and *Rbr*⁻ *P. gingivalis* strains in a model of oral infection [50]. As expected, mice that were orally challenged with wild-type *P. gingivalis* exhibited significantly greater alveolar oral bone loss compared to unchallenged mice (Figure 10). In contrast, we did not observe significant changes in oral bone loss in wild-type mice infected with the *P. gingivalis* *Rbr*⁻ strain relative to the unchallenged wild-type mice. Thus, the presence of Rbr promotes increased *P. gingivalis*-induced inflammation and subsequent oral bone loss in this mouse model. However, in this case, and in resemblance to human periodontal disease, the infection was localized to the periodontal tissue. In contrast to the situation in subcutaneous chambers, the local host defense system and/or bacterial growth conditions in the oral cavity prevent systemic dissemination of *P. gingivalis*.

Discussion

In this study, we demonstrate that bacterial Rbr protects in vivo against the host-mediated killing of the oral anaerobe, *P. gingivalis*, but that Rbr does not protect against the neutrophil oxidative burst. These conclusions are based on two observations. First, the wild-type and *Rbr*⁻ *P. gingivalis* strains did not differ in their susceptibilities to killing by normal neutrophils. Second, although the *rbr* mutation hindered the growth and virulence of *P. gingivalis* in mice, no difference was observed in vitro in terms of the killing of *P. gingivalis* between the normal and NADPH oxidase-deficient neutrophils.

Interestingly, the virulence of *Rbr*-deficient *P. gingivalis* was not restored in phagocytic *Cybb*^{-/-} mice. Exactly to the contrary, the genetic abrogation of the oxidative burst in *Cybb*^{-/-} mice was found to protect animals from mortality related to inoculation with wild-type *P. gingivalis*. Signifi-




Figure 10. Kinetics of Wild-Type *P. gingivalis* Growth in Media Supplemented with Inflammatory Exudates Elicited in Wild-Type and *Cybb*^{-/-} Mice with Thioglycollate

DOI: 10.1371/journal.ppat.0020076.g010

cantly, this protection was even more pronounced than in wild-type mice infected with the *P. gingivalis* *Rbr*⁻ mutant strain (Figure 1). As compared to the wild-type *P. gingivalis* strain, the isogenic *Rbr*⁻ mutant strain grew slower, reaching lower numbers in wild-type mice. Furthermore, the *Rbr*⁻ mutant strain was found in higher numbers in wild-type mice than in *Cybb*^{-/-} mice. In *Cybb*^{-/-} mice, the initial growth rate of the *Rbr*⁻ mutant was significantly reduced in comparison to that observed for the wild-type strain (Figure 2).

These interesting observations raise several important questions with regard to how Rbr protects *P. gingivalis*. From the data presented, and in agreement with previous reports showing no clear difference in *P. gingivalis* killing by neutrophils under aerobic and anaerobic conditions [31,51], it is obvious that both in vivo and in vitro Rbr is dispensable for protection against ROS generated by the phagocyte-derived oxidative burst. Physiologically, this is understandable since the ROS-quenching function of Rbr would be redundant within the anaerobic habitat of the subgingival bacteria plaque occupied by *P. gingivalis*, in which the oxygen-dependent bactericidal activities of neutrophils are nonfunctional [29]. In anoxic environments, such as deep wounds, periodontal pockets, and subcutaneous chambers, the major weapons used by phagocytes against invading bacteria are RNS generated by iNOS (NOS2) and/or bactericidal peptides and proteins, which expression is strongly enhanced in hypoxia [52]. The effectiveness of neutrophil-derived bactericidal peptides in the killing of *P. gingivalis* [30] and other bacteria [53] is well documented. Since RNS have been shown to possess strong antimicrobial activities against a broad array of bacteria [54], it is not surprising that RNS participate in *P. gingivalis* killing by murine macrophages (Figure 4). This ex vivo finding corroborates very well the observation that the presence of functional iNOS is essential to attenuate *P. gingivalis* growth in vivo in mice [33]. Significantly, this antibacterial effect was exerted in the anoxic environment of the subcutaneous chambers.

In as much as RNS participation in *P. gingivalis* killing was anticipated, the ability of Rbr to protect against the antibacterial activity of RNS was somewhat unexpected, making this a novel and interesting discovery. Significantly, the dual function of Rbr elicited protection of *P. gingivalis* in

vivo not only against oxygen stress but also against RNS correlates well with the pathological events associated with the infection. Taking into account that mice with the genetically inactivated oxidative burst produce greater amounts of RNS in vivo than normal animals [55–57], our findings explain very well the observed differences in pathology and microbial growth of the *P. gingivalis* strains in normal and *Cybb*^{-/-} mice. In wild-type mice, the initial production of ROS in a chamber, before the environment becomes anoxic, may suppress RNS generation and allow robust growth of the Rbr-protected *P. gingivalis* strain, but not the Rbr⁻ mutant (Figure 2). In *Cybb*^{-/-} animals, the growth of the wild-type *P. gingivalis* strain is controlled by the elevated level of RNS, which overwhelms the protection afforded by Rbr. Finally, in concordance with our findings, the Rbr⁻ *P. gingivalis* strain in the absence of the phagocyte oxidative burst showed the highest level of growth attenuation. In vivo, the predominant sources of antibacterial RNS are stimulated monocytes, macrophages, and epithelial cells, but not neutrophils [58,59]. In this context, the finding that the role of Rbr in *P. gingivalis* infection is related to protection against RNS, but not ROS, explains why the presence or absence of this protein has no effect on bacterium killing by neutrophils in vitro, although Rbr shows protective functions in vivo.

In addition to facilitating the growth of *P. gingivalis*, the combination of the host phagocyte oxidative burst and the expression of bacterial Rbr appears to be instrumental in *P. gingivalis* systemic dissemination from the subcutaneous chambers, which is manifested as bacteremia and heavy colonization of the spleen, liver, lungs, and kidneys (Figure 9). We observed very limited dissemination of the *P. gingivalis* wild-type strain in the *Cybb*^{-/-} mice, apparently due to enhanced production of RNS in the oxidative burst-deficient animals [55–57]. Interestingly, significantly higher numbers of dead cells were found in the chamber fluids recovered from *Cybb*^{-/-} mice than from wild-type animals (Figure 7B). A possible reason for this finding is the very complex intimate relationship between nitric oxide and superoxide in leukocyte migration, cell survival, cell apoptosis, and clearance of apoptotic and necrotic cell bodies [60]. In this context, elimination of the phagocyte NADPH oxidase-dependent regulatory functions may have profound effects on cell fate at the site of a confined local infection [61–66], resulting in disproportionately high numbers of dead cells in the chamber fluids from *Cybb*^{-/-} mice.

The other striking difference in the contents of the chamber fluid samples was the level of MPO, an indicator of neutrophil turnover, which was highest in wild-type mice infected with the *P. gingivalis* wild-type strain (Figure 7C). Furthermore, this particular combination of host and pathogen was accompanied by a significant increase in the numbers of circulating neutrophils and considerable systemic activation of these neutrophils, as illustrated by the two-fold decrease in L-selectin expression on peripheral neutrophils [67] (Figure 8) and the high level of MPO in the serum. Finally, MPO was detected in the serum samples, albeit at very low levels, of *Cybb*^{-/-} mice inoculated with the wild-type strain (Figure 7C). The presence of circulating MPO can be directly linked to the pathology of *P. gingivalis* infection in wild-type mice, since this protein by binding to CD11b/CD18 integrins serves as a powerful autocrine and paracrine stimulator of neutrophil activation, provoking degranulation, expression

of integrins, and oxidative burst discharge [48]. Collectively, these data indicate that the functional oxidative burst response and neutrophil activation are prerequisites for *P. gingivalis* to escape from the chambers and disseminate to the organs. In this context, it is important to reiterate that the differences noted in the growth rate and dissemination of *P. gingivalis* in wild-type and *Cybb*^{-/-} mice were not related to differences in the availability of nutrients or growth factors generated through the oxidation of some host components (Figure 10). These results contrast with those reported previously for aerobic pathogens, such as *Burkholderia cepacia*, *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Escherichia coli*, and *Salmonella typhimurium*, all of which have been shown to exhibit enhanced virulence in genetically abrogated phagocyte oxidative burst animal models [68–72].

The mortality rates of mice infected with *P. gingivalis* correlated with increases in the concentration of LPS (in the serum and chamber fluid samples) and levels of neutrophil activation in the blood samples (Figure 5). The IL-6 and TNF- α concentrations in the sera were unrelated to the LPS levels, either in the chamber fluid samples or in blood (Figure 6), which is not surprising given that *P. gingivalis* LPS is known to be nontoxic and a weak stimulator of cytokine release [73]. Thus, animal mortality following *P. gingivalis* infection was not due to a “cytokine storm,” as in the case of sepsis induced by enterobacterial LPS [74]. These results re-emphasize the self-destructive function of the oxidative burst aimed at *P. gingivalis*. In addition to killing microbes, the oxidative burst may have important regulatory functions, including effects on neutrophil gene transcription [75] and macrophage and lymphocyte activation [76]. A bacterium-induced malfunction of these regulatory pathways and excessive oxidative burst and neutrophil degranulation in aerated tissues, such as the lungs and tissues supplied with oxygen by the blood, which include the spleen, liver, and kidneys, all of which were heavily colonized by *P. gingivalis* in the wild-type mice (Figure 9A), may cause serious loss of function of these organs and animal death. It is tempting to speculate that *P. gingivalis* subverts these signaling pathways to its advantage, such that in *Cybb*^{-/-} mice the virulence of the organism is attenuated.

Impaired neutrophil chemotaxis [77,78] and deficiency of serine proteases [79] or the bactericidal peptide LL-37 [80] in neutrophil granules are associated with aggressive forms of periodontal disease despite the presence of a normal oxidative burst. Conversely, patients with CGD are devoid of the NADPH oxidase activity and seldom suffer from periodontal disease, although they are prone to severe recurrent bacterial infections [10]. In addition, recent evidence indicates that a molecular defect in intracellular lipid signaling contributes to excessive superoxide generation by neutrophils leading to tissue destruction in patients with periodontal disease [81]. These clinical associations, which are in concordance with our data, negate the importance of the oxidative burst in combating periodontal pathogens. Collectively, these results suggest that the host respiratory burst response to acute anaerobic infection may contribute significantly to the morbidity associated with periodontal disease.

The insignificance of the oxygen-dependent bactericidal activity in maintaining periodontal tissue homeostasis following infection with anaerobic pathogens is also clear from our experimental data obtained using the oral model for *P.*




Figure 11. The Presence of the Rbr Gene Increases *P. gingivalis*-Induced Oral Bone Loss

Following a 10-d regimen of oral trimethoprim-sulfamethoxazole and a 3-d rest period, mice were administered 1.0×10^9 CFU of the *P. gingivalis* strains by oral gavage three times over a 1-wk period. One group of animals was not treated, and served as the age-matched controls. Six wk after oral challenge, all of the animals were killed. Linear bone loss measurements ($n = 14$) were obtained from the maxillary molars of each mouse, and the data for linear bone loss (in mm) from the cemento-enamel junction to the alveolar bone crest are presented as the means \pm SEM of the pooled linear measurements obtained for each group. * $p < 0.05$; ** $p < 0.01$; as determined by the Student's *t* test. DOI: 10.1371/journal.ppat.0020076.g011

gingivalis infection (Figure 11). The levels of oral bone loss induced by the wild-type *P. gingivalis* strain were similar in the C57BL/6J and Cybb^{-/-} mice. Unlike subcutaneous chamber inoculation, the presence of the intact oxidative burst system in mice orally infected with the wild-type *P. gingivalis* strain did not result in this model in excessive inflammation and subsequent bone loss or systemic dissemination of infection and high mortality. This is apparently due to a localized type of periodontal infection, in which *P. gingivalis* colonizes only subgingival tooth surfaces, growing in the form of a biofilm separated from the circulation by a thick “wall” of leukocytes [82]. This protection, together with defenses in the blood, is evidently sufficient since human clinical cases of bacteremia caused by *P. gingivalis* are unknown. For *P. gingivalis*, Rbr obviously plays a role in shielding the bacterium against the host immune system at periodontal sites, since the presence of Rbr contributed to the morbidity induced by *P. gingivalis* infection. This is apparent from the fact that in normal mice, oral bone loss was significantly greater upon infection with the wild-type *P. gingivalis* strain as compared to that observed following infection with the Rbr⁻ mutant strain (Figure 11).

The consequences of the harmful interaction between the host oxidative burst response and anaerobic bacteria possessing Rbr are likely to extend beyond infection by the periodontal disease pathogen *P. gingivalis*. A recent BLAST search of the 878 genomic sequences in the National Center for Biotechnology Information Microbial Genomes Database with the *P. gingivalis* *rbr* gene found 114 homologs in 69 organisms, the vast majority being obligate anaerobes or microaerophiles. Although the majority of these air-sensitive bacteria are not known to be harmful to humans, 16 are human commensals or pathogens. These include *B. pseudomallei*, a causative agent of melioidosis and a potential bioterrorist weapon [83], *B. cepacia*, an opportunistic colonizer of the lungs of patients with cystic fibrosis [84], *Campylobacter jejuni*, the most common cause of bacterial

enterocolitis in humans [85], and *Clostridium tetani* and *C. perfringens*, which are the causative organisms of tetanus and gas gangrene, respectively. Furthermore, human commensals that carry the *rbr* gene homolog, such as *Clostridium*, *Bacteroides*, *Prevotella*, *Treponema*, and *Fusobacterium* species, are the most frequently isolated organisms in anaerobic bacteremia, endocarditis, and bone and joint infections, as well as skin and oral abscesses [86]. Thus, our results on *P. gingivalis* may have relevance to oxygen-dependent killing mechanisms operating during anaerobic infections and the contribution of Rbr to protection against molecules of host defense by other pathogenic anaerobic organisms. In the clash between these two systems, an oxidative burst elicited by neutrophils, which are unable to eliminate the bacteria, may be responsible for host tissue damage, either directly through oxidative destruction of macromolecules or indirectly, by activation of other cells. This leads to systemic bacterial dissemination, organ colonization and injury, and eventually death of the host.

Materials and Methods

Mouse strains. All of the animal experiments were performed in accordance with the guidelines of the Institutional Animal Care and Use Committee of Boston University and in conformance with the Standards of Public Health Service Policy on Human Care and Use of Laboratory Animals. Six-week-old female C57BL/6J and phagocytic Cybb^{-/-} mice were purchased from the Jackson Laboratory (Bar Harbor, Maine, United States). All of the animals received the standard laboratory chow diet and were examined daily for any adverse health conditions.

***P. gingivalis* cultures.** *P. gingivalis* strain W83 was grown in anaerobe broth (Difco Laboratories, Sparks, Maryland, United States) in an anaerobic chamber in an atmosphere of 80% N₂, 10% H₂, and 10% CO₂. The isogenic *P. gingivalis* *rbr*-knockout strain (Rbr⁻) [22] was grown under the same conditions with the addition of erythromycin at a concentration of 5 μg/ml. Bacterial cell counts were determined on all the *P. gingivalis* cultures prior to the animal experiments, to confirm viability and to determine the number of viable organisms present at the time of animal challenge.

Chamber implantation. Following anesthesia with isoflurane, sterile wire, coil-shaped stainless steel chambers were implanted subcutaneously in the dorso-lumbar region of each mouse [27]. Incisions were closed using 4.0-G silk sutures and the animals were allowed to rest for 14 d, at which timepoint all the incisions had healed completely.

Chamber inoculation with *P. gingivalis*. C57BL/6J and Cybb^{-/-} animals were each divided into three groups, with $n = 7$ for each group. The first group was inoculated with 0.1 ml of *P. gingivalis* W83, the second group received 0.1 ml of the Rbr⁻ strain (10^9 CFU/ml), and the third group was inoculated with vehicle only. A minimum of 70 μl of chamber fluid was collected from each mouse using a syringe with a 25-G sterile hypodermic needle at 1, 3, 6, and 10 d after challenge.

Chamber fluid analysis. Each chamber fluid sample was divided into several samples to be used for the determination of CFU/ml (10 μl), total inflammatory cell counts (10 μl), and differential inflammatory cell counts (5 μl); the remainder (~40 μl) was stored at -80 °C for subsequent enzyme-linked immunosorbent assay (ELISA) determinations of the IL-6, TNF-α, IL-10, LPS, and MPO levels.

Microbial analysis. Aliquots of chamber fluid from each mouse (10 μl) were serially diluted ten-fold with 1% peptone. Dilutions were plated onto anaerobic blood agar plates (Remel, Lenexa, Kansas, United States) in duplicate and incubated in an anaerobic chamber for 6–7 d. The CFU/ml values were determined by direct colony counting. The presence of *P. gingivalis* was confirmed by Gram staining and colony PCR. The forward (sense) primer 5'-CGTGCCAGCAGCCGCGTAATACG-3' and the reverse (antisense) primer 5'-TACATAGAAGCCCCGAAGGAAGAC-3' were used to amplify the *P. gingivalis* 16S gene. PCR amplification was carried out using the following conditions: 40 cycles of denaturation at 95 °C for 30 sec, annealing at 60 °C for 30 sec, and elongation at 72 °C for 1 min. The PCR products were separated by agarose gel electrophoresis and visualized using a UV transilluminator after staining with ethidium bromide.

Killing of bacteria by murine bone marrow and human peripheral blood neutrophils. Human neutrophils were purified from fresh human blood by dextran sedimentation and centrifugation through Ficoll/Hypaque. The mouse neutrophils were isolated from the bone marrow. Briefly, the bone marrow was collected from the femurs and tibias of normal and *Cybb*^{-/-} mice. The distal and proximal tips were removed and washed with cold KRG buffer (120 mM NaCl, 5 mM KCl, 8.3 mM Na₂HPO₄ [pH 7.3], 10 mM glucose, 1 mM CaCl₂) using a 1-ml syringe and 27-G needle. After dispersing cell clumps using a 1-ml pipette on ice, the suspension was centrifuged (200g, 10 min, 4 °C) and the cell pellet was resuspended in 2 ml of sterile PBS. The cells were carefully layered on the top of a discontinuous Percoll gradient (72%, 63%, 50%; 2 ml each). After centrifugation (500g, 30 min), the lowest band containing neutrophils was collected, residual erythrocytes were removed by hypotonic shock, and the neutrophils were suspended in PBS at a concentration of 2×10^7 cell/ml.

Mouse (three to five animals) and human (one donor) neutrophils (2.5×10^7) in 0.5 ml were mixed with IgG-opsonized *P. gingivalis* (5×10^6 CFU/ml) at a ratio of one target organism to five neutrophils in 0.5 ml PBS and incubated either in an aerobic or anaerobic atmosphere. The rate of bacterial killing was measured as described previously [87], omitting lysostaphin. Cell lysates that contained viable *P. gingivalis* were plated on 10% horse blood agar plates and incubated anaerobically for 5 d. The results were calculated at the mean (\pm SE) from at least two experiments with colony counts performed in triplicate for each sample and expressed as a percentage of the original numbers at time zero.

Killing of *P. gingivalis* W83 and Rbr⁻ strain by murine macrophages in the presence of L-NAME. Mice were injected intraperitoneally with sterile thioglycollate broth (3%; Difco Laboratories) at 3 ml/mice, and inflammatory exudates were recovered after 72 h by peritoneal lavage with sterile RPMI 1640 medium supplemented with 10% FCS. Cells harvested from six animals were pooled, washed in RPMI 1640 with 10% FCS and seeded on 24-well plates (1×10^6 /well; TPP Switzerland, Trasadingen, Switzerland) in 1 ml of RPMI 1640 with 10% of FCS, and divided into two groups. After 6 h, both groups were stimulated to express iNOS by the addition of *E. coli* serotype O26:B6 LPS (400 ng/ml; Sigma, Saint Louis, Missouri, United States). One group was exposed to LPS in the presence of L-NAME (Calbiochem, San Diego, California, United States) at a final concentration of 5 mM. After overnight culture, the nitrite and nitrate levels were determined by the Griess reaction. To synchronise phagocytosis, plates that contained macrophages were inoculated with 5×10^6 fresh, opsonized bacteria per well and centrifuged at 380g for 8 min at 4 °C. At the indicated timepoints, the supernatant and cells were harvested, lysed, and plated onto blood agar plates. Colonies were counted after 5 d of culture, and the percentage of *P. gingivalis* survival was calculated according to the equation $(\text{CFU at } t_1 / \text{CFU at } t_0) \times 100$. This assay measures the total numbers of uningested (in the supernatant) and ingested (in the cell lysate) viable bacteria.

Assessment of ROS production by bone marrow neutrophils exposed to *P. gingivalis*. Bone marrow neutrophils from four mice (1.0×10^7 /ml) suspended in sterile PBS were seeded in 100 μ l into a black 96-well plate (Nunc, Rochester, New York, United States). Then, 100 μ l of serum-opsonized *P. gingivalis* W83 or Rbr⁻ (5.0×10^8 /ml) was added together with 50 μ l of luminol solution (0.53 mg/ml; Sigma) and chemiluminescence was recorded for 120 min using the EG&G Berthold MicroLumet LB 96P luminometer (Berthold Technologies, Bad Wildbad, Germany).

Analysis of *P. gingivalis* growth in media supplemented with an inflammatory exudate from wild-type and *Cybb*^{-/-} mice. Mice were injected intraperitoneally with sterile thioglycollate broth (3%; Difco Laboratories) at 3 ml/mice and inflammatory exudates were recovered by peritoneal lavage with sterile RPMI 1640 medium supplemented with 10% FCS. The lavage was sonicated to disintegrate the cells, mixed 1:1 with a two-fold dilution of Scheadler broth, filter-sterilized (0.4- μ m filter) and inoculated with *P. gingivalis*. Bacterial growth under anaerobic conditions was monitored by measurement of the culture turbidity at 660 nm in a spectrophotometer.

Analysis of blood neutrophil activation. Subcutaneous chambers were implanted in the C57BL/6J mice. Ten d later, the chambers were inoculated with 100 μ l of *P. gingivalis* W83 or Rbr⁻ strain (1×10^9 /ml). As a control, 100 μ l of sterile PBS was inoculated into the chamber. Three d after inoculation, blood samples were collected from the heart into heparinized tubes, and the blood cells were stained with FITC-conjugated Ly-6G monoclonal antibodies and R-phycoerythrin-conjugated CD62L monoclonal antibodies. Rat IgG_{2A} κ monoclonal antibodies were used as the isotype control. The samples were analyzed by flow cytometry in the FACSCalibur (BD Biosciences, Franklin Lakes, New Jersey, United States) using the CellQuest Software (BD Biosciences).

Inflammatory cell analysis. A 10- μ l chamber fluid sample from each mouse was stained with trypan blue, and total and viable inflammatory cell counts were obtained using light microscopy at 40 \times magnification in a hemocytometer chamber. In addition, 5- μ l samples were placed on slides and stained with Giemsa-Wright stain for differential cell counts. The neutrophils were differentiated based on their lobed nuclei.

Quantification of cytokine levels. The serum and chamber fluid levels of TNF- α , IL-6, and IL-10 were determined by ELISA (Pierce Endogen, Rockford, Illinois, United States). For the determination of LPS levels, the QCL-1000 Chromogenic LAL Test Kit (Cambrex, Santa Rosa, California, United States) was used. The levels of MPO in the sera and chamber fluids were determined by ELISA (Calbiochem, La Jolla, California, United States).

Assessment of oral bone loss. Oral bone loss was assessed at defined landmark sites on the maxillary molars of each mouse, as previously described [50]. Using a stereomicroscope, an observer who was blinded to the identity of the groups obtained linear measurements (14 sites) for each animal from the cemento-enamel junction to the alveolar bone crest [88]. Measurements were obtained with a digital camera linked to a computer, and onscreen measurements were converted to micrometers using IPLabs (Scanalytics Inc., Fairfax, Virginia, United States).

Statistical analysis. Statistical comparison analyses were performed using the Mann-Whitney *U* test or Student's *t* test as indicated in the figure legends. Differences in the data were considered significant when the probability value was less than 5.0% ($p < 0.05$).

Supporting Information

Accession Numbers. The National Center for Biotechnology Information (NCBI) (<http://www.ncbi.nlm.nih.gov>) accession number for Rbr is AAK19552.

The Institute for Genomic Research-Comprehensive Microbial Resource database (TIGR-CMR) (<http://cmr.tigr.org/tigr-scripts/CMR/CmrHomePage.cgi>) accession number for the *P. gingivalis* rbr gene is PG 0195.

Acknowledgments

We are indebted to Ms. Paulina Kulig for flow cytometry analyzes.

Author contributions. PM, YT, HY, and JP conceived and designed the experiments. PM, YT, HY, MK, and FCG performed the experiments. PM, YT, HY, MS, MK, FCG, DMK, JT, LVC, KAN, CAG, and JP analyzed the data. MS, CAG, and JP contributed reagents/materials/analysis tools. JP wrote the paper.

Funding. The authors acknowledge the support of the National Institutes of Health through grants DE 09761 (to J.T.), GM040388 (to D.M.K.) and DE 09161 (to C.A.G.), and through grants 3 PO4A 003 24 and 2 P05A 137 24 from the Committee of Scientific Research (KBN, Poland) (to J.P. and P.M., respectively). P.M. was sponsored by a *SUBSYDIUM PROFESORSKIE* award (to J.P.) from the Foundation for Polish Science (FNP, Warszawa, Poland). Financial support from the Swedish Rheumatism Association, Nanna Svartz Foundation, and the King Gustaf V Foundation (to L.V.C.) is also acknowledged.

Competing interests. The authors have declared that no competing interests exist.

References

1. Witko-Sarsat V, Rieu P, Descamps-Latscha B, Lesavre P, Halbwachs-Mecarelli L (2000) Neutrophils: Molecules, functions and pathophysiological aspects. *Lab Invest* 80: 617–653.
2. Belaouaj A, Kim KS, Shapiro SD (2000) Degradation of outer membrane protein A in *Escherichia coli* killing by neutrophil elastase. *Science* 289: 1185–1188.
3. Lopez-Boado YS, Espinola M, Bahr S, Belaouaj A (2004) Neutrophil serine proteinases cleave bacterial flagellin, abrogating its host response-inducing activity. *J Immunol* 172: 509–515.
4. Weinrauch Y, Drujan D, Shapiro SD, Weiss J, Zychlinsky A (2002) Neutrophil elastase targets virulence factors of enterobacteria. *Nature* 417: 91–94.
5. Brinkmann V, Reichard U, Goosmann C, Fauler B, Uhlemann Y, et al. (2004) Neutrophil extracellular traps kill bacteria. *Science* 303: 1532–1535.

6. Lektstrom-Himes JA, Gallin JI (2000) Immunodeficiency diseases caused by defects in phagocytes. *N Engl J Med* 343: 1703–1714.
7. Roos D, van Bruggen R, Meischl C (2003) Oxidative killing of microbes by neutrophils. *Microbes Infect* 5: 1307–1315.
8. Goldblatt D, Thrasher AJ (2000) Chronic granulomatous disease. *Clin Exp Immunol* 122: 1–9.
9. Babior BM (2004) NADPH oxidase. *Curr Opin Immunol* 16: 42–47.
10. Winkelstein JA, Marino MC, Johnston RB, Boyle J, Curnutte, et al. (2000) Chronic granulomatous disease. Report on a national registry of 368 patients. *Medicine (Baltimore)* 79: 155–169.
11. Cohen MS, Leong PA, Simpson DM (1985) Phagocytic cells in periodontal defense. Periodontal status of patients with chronic granulomatous disease of childhood. *J Periodontol* 56: 611–617.
12. Reeves EP, Lu H, Jacobs HL, Messina CG, Bolsover S, et al. (2002) Killing activity of neutrophils is mediated through activation of proteases by K⁺ flux. *Nature* 416: 291–297.
13. Ahluwalia J, Tinker A, Clapp LH, Duchon MR, Abramov AY, et al. (2004) The large-conductance Ca²⁺-activated K⁺ channel is essential for innate immunity. *Nature* 427: 853–858.
14. Nizet V, Gallo RL (2003) Cathelicidins and innate defense against invasive bacterial infection. *Scand J Infect Dis* 35: 670–676.
15. Deas DE, Mackey SA, McDonnell HT (2003) Systemic disease and periodontitis: Manifestations of neutrophil dysfunction. *Periodontol* 2000 32: 82–104.
16. Hart TC, Shapira L, Van Dyke TE (1994) Neutrophil defects as risk factors for periodontal diseases. *J Periodontol* 65: 521–529.
17. Loesche WJ, Grossman NS (2001) Periodontal disease as a specific, albeit chronic, infection: Diagnosis and treatment. *Clin Microbiol Rev* 14: 727–752.
18. Van Dyke TE, Hoop G (1991) Neutrophil function and oral disease. *Crit Rev Oral Biol Med* 1: 117–133.
19. Cutler CW, Kalmal JR, Genco CA (1995) Pathogenic strategies of the oral anaerobe, *Porphyromonas gingivalis*. *Trends Microbiol* 3: 45–51.
20. Johnson NA, McKenzie R, McLean L, Sowers LC, Fletcher HM (2004) 8-oxo-7,8-dihydroguanine is removed by a nucleotide excision repair-like mechanism in *Porphyromonas gingivalis* W83. *J Bacteriol* 186: 7697–7703.
21. Nakayama K (1994) Rapid viability loss on exposure to air in a superoxide dismutase-deficient mutant of *Porphyromonas gingivalis*. *J Bacteriol* 176: 1939–1943.
22. Sztukowska M, Bugno M, Potempa J, Travis J, Kurtz DM Jr (2002) Role of Rbr in the oxidative stress response of *Porphyromonas gingivalis*. *Mol Microbiol* 44: 479–488.
23. Coulter ED, Shenvi NV, Kurtz DM Jr (1999) NADH peroxidase activity of rubrerythrin. *Biochem Biophys Res Commun* 255: 317–323.
24. Lumppio HL, Shenvi NV, Summers AO, Voordouw G, Kurtz DM Jr. (2001) Rubrerythrin and rubredoxin oxidoreductase in *Desulfovibrio vulgaris*. A novel oxidative stress protection system. *J Bacteriol* 183: 101–108.
25. Weinberg WV, Jenney FE Jr, Cui X, Adams MWW (2004) Rubrerythrin from the hyperthermophilic archaeon *Pyrococcus furiosus* is a rubredoxin-dependent iron-containing peroxidase. *J Bacteriol* 186: 7888–7895.
26. Smalley JW, Birss AJ, Silver J (2000) The periodontal pathogen *Porphyromonas gingivalis* harnesses the chemistry of the m-oxo bishaem of iron protophyrin IX to protect against hydrogen peroxide. *FEMS Microbiol Lett* 183: 159–164.
27. Genco CA, Cutler CW, Kapczynski D, Maloney K, Arnold RR (1991) A novel mouse model to study the virulence of and host response to *Porphyromonas (Bacteroides) gingivalis*. *Infect Immun* 59: 1255–1263.
28. Genco CA, Kapczynski DR, Cutler CW, Arko RJ, Arnold RR (1992) Influence of immunization on *Porphyromonas gingivalis* colonization and invasion in the mouse chamber model. *Infect Immun* 60: 1447–1454.
29. Jarstrand C, Wiernik A, Revesz L (2000) Significance of oxygen availability for release of oxygen free radicals and lysozyme by neutrophils. *J Clin Lab Immunol* 32: 37–39.
30. Isogai E, Isogai H, Matuo K, Hirose K, Kowashi Y, et al. (2003) Sensitivity of genera *Porphyromonas* and *Prevotella* to the bactericidal action of C-terminal domain of human CAP18 and its analogues. *Oral Microbiol Immunol* 18: 329–332.
31. Klempner MS (1984) Interactions of polymorphonuclear leukocytes with anaerobic bacteria. *Rev Infect Dis* 6 (Suppl 1): S40–S44.
32. Van Amersfoort ES, Van Berkel TJ, Kuiper J (2003) Receptors, mediators, and mechanisms involved in bacterial sepsis and septic shock. *Clin Microbiol Rev* 16: 379–414.
33. Gyurko R, Boustany G, Huang PL, Kantarci A, Van Dyke TE, et al. (2003) Mice lacking inducible nitric oxide synthase demonstrate impaired killing of *Porphyromonas gingivalis*. *Infect Immun* 71: 4917–4924.
34. Banbula A, Bugno M, Kuster A, Heinrich PC, Travis J, et al. (1999) Rapid and efficient inactivation of IL-6 by gingipains, lysine- and arginine-specific proteinases from *Porphyromonas gingivalis*. *Biochem Biophys Res Commun* 261: 598–602.
35. Calkins CC, Platt K, Potempa J, Travis J (1998) Inactivation of tumor necrosis factor- α by proteinases (gingipains) from the periodontal pathogen, *Porphyromonas gingivalis*. Implications of immune evasion. *J Biol Chem* 273: 6611–6614.
36. Fletcher J, Nair S, Poole S, Henderson B, Wilson M (1998) Cytokine degradation by biofilms of *Porphyromonas gingivalis*. *Curr Microbiol* 36: 216–219.
37. Schmekel B, Karlsson SE, Linden M, Sundstrom C, Tegner H, et al. (1990) Myeloperoxidase in human lung lavage. I. A marker of local neutrophil activity. *Inflammation* 14: 447–454.
38. Dykens JA, Baginski TJ (1998) Urinary 8-hydroxydeoxyguanosine excretion as a non-invasive marker of neutrophil activation in animal models of inflammatory bowel disease. *Scand J Gastroenterol* 33: 628–636.
39. Caimi G, Hoffmann E, Montana M, Incalcaterra E, Dispensa F, et al. (2005) Plasma markers of platelet and polymorphonuclear leukocyte activation in young adults with acute myocardial infarction. *Clin Hemorheol Microcirc* 32: 67–74.
40. Miyasaki KT, Nemirovskiy E (1997) Myeloperoxidase isoform activities released by human neutrophils in response to dental and periodontal bacteria. *Oral Microbiol Immunol* 12: 27–32.
41. Keatings VM, Barnes PJ (1997) Granulocyte activation markers in induced sputum: Comparison between chronic obstructive pulmonary disease, asthma, and normal subjects. *Am J Respir Crit Care Med* 155: 449–453.
42. Holzer W, Petersen F, Strittmatter W, Matzku S, von Hoegen I (1996) A fusion protein of IL-8 and a Fab antibody fragments binds to IL-8 receptors and induces neutrophil activation. *Cytokine* 8: 214–221.
43. Grisham MB, Everse J, Janssen HF (1988) Endotoxemia and neutrophil activation in vivo. *Am J Physiol* 254: H1017–H1022.
44. Yamamoto C, Yoneda T, Yoshikawa M, Fu A, Tokuyama T, et al. (1997) Airway inflammation in COPD assessed by sputum levels of interleukin-8. *Chest* 112: 505–510.
45. Tauber E, Herouy Y, Goetz M, Urbanek R, Hagel E, et al. (1999) Assessment of serum myeloperoxidase in children with bronchial asthma. *Allergy* 54: 177–182.
46. Franck T, Grulke S, Deby-Dupont G, Deby C, Duvivier H, et al. (2005) Development of an enzyme-linked immunosorbent assay for specific equine neutrophil myeloperoxidase measurement in blood. *J Vet Diagn Invest* 17: 412–419.
47. Kristjansson G, Venge P, Wanders A, Loof L, Hallgren R (2004) Clinical and subclinical intestinal inflammation assessed by the mucosal patch technique: Studies of mucosal neutrophil and eosinophil activation in inflammatory bowel diseases and irritable bowel syndrome. *Gut* 53: 1806–1812.
48. Lau D, Mollnau H, Eiserich JP, Freeman BA, Daiber A, et al. (2005) Myeloperoxidase mediates neutrophil activation by association with CD11b/CD18 integrins. *Proc Natl Acad Sci U S A* 102: 431–436.
49. Fleming TJ, Fleming ML, Malek TR (1993) Selective expression of Ly-6G on myeloid lineage cells in mouse bone marrow. RB6-8C5 mAb to granulocyte-differentiation antigen (Gr-1) detects members of the Ly-6 family. *J Immunol* 151: 2399–2408.
50. Gibson FC 3rd, Genco CA (2001) Prevention of *Porphyromonas gingivalis*-induced oral bone loss following immunization with gingipain R1. *Infect Immun* 69: 7959–7963.
51. Thompson HL, Wilton JM (1991) Effects of anaerobiosis and aerobiosis on interactions of human polymorphonuclear leukocytes with the dental plaque bacteria *Streptococcus mutans*, *Capnocytophaga ochracea*, and *Bacteroides gingivalis*. *Infect Immun* 59: 932–940.
52. Peyssonnaud C, Datta V, Cramer T, Doedens A, Theodorakis EA, et al. (2005) HIF-1 α expression regulates the bactericidal capacity of phagocytes. *J Clin Invest* 115: 1806–1815.
53. Mayer-Scholl A, Hurwitz R, Brinkmann V, Schmid M, Jungblut P, et al. (2005) Human neutrophils kill *Bacillus anthracis*. *PLoS Pathog* 1: e23. DOI: 10.1371/journal.ppat.0010023
54. MacMicking J, Xie QW, Nathan C (1997) Nitric oxide and macrophage function. *Annu Rev Immunol* 15: 323–350.
55. van der Veen RC, Dietlin TA, Hofman FM, Pen L, Segal BH, et al. (2000) Superoxide prevents nitric oxide-mediated suppression of helper T lymphocytes: Decreased autoimmune encephalomyelitis in nicotinamide adenine dinucleotide phosphate oxidase knockout mice. *J Immunol* 164: 5177–5183.
56. van der Veen RC, Dietlin TA, Karapetian A, Holland SM, Hofman FM (2004) Extra-cellular superoxide promotes T cell expansion through inactivation of nitric oxide. *J Neuroimmunol* 153: 183–189.
57. Ramsey KH, Sagar IM, Rana SV, Gupta J, Holland SM, et al. (2001) Role for inducible nitric oxide synthase in protection from chronic *Chlamydia trachomatis* urogenital disease in mice and its regulation by oxygen free radicals. *Infect Immun* 69: 7374–7379.
58. Kleinert H, Schwarz PM, Forstermann U (2003) Regulation of the expression of inducible nitric oxide synthase. *Biol Chem* 384: 1343–1364.
59. Aktan F. (2004) iNOS-mediated nitric oxide production and its regulation. *Life Sci* 75: 639–653.
60. Brune B (2005) The intimate relation between nitric oxide and superoxide in apoptosis and cell survival. *Antioxid Redox Signal* 7: 497–507.
61. Nathan C. (2003) Specificity of a third kind: Reactive oxygen and nitrogen intermediates in cell signaling. *J Clin Invest* 111: 769–778.
62. Isenberg JS, Tabatabai N, Spinelli HM (2005) Nitric oxide modulation of low-density mononuclear cell transendothelial migration. *Microsurgery* 25: 452–456.
63. Razavi HM, Wang le F, Weicker S, Rohan M, Law C, et al. (2004) Pulmonary neutrophil infiltration in murine sepsis: Role of inducible nitric oxide synthase. *Am J Respir Crit Care Med* 170: 227–233.
64. Yasui K, Kobayashi N, Yamazaki T, Agematsu K, Matsuzaki S, et al. (2005)

- Superoxide dismutase (SOD) as a potential inhibitory mediator of inflammation via neutrophil apoptosis. *Free Radic Res* 39: 755–762.
65. Ward C, Wong TH, Murray J, Rahman I, Haslett C, et al. (2000) Induction of human neutrophil apoptosis by nitric oxide donors: Evidence for a caspase-dependent, cyclic-GMP-independent, mechanism. *Biochem Pharmacol* 59: 305–314.
 66. Sato E, Simpson KL, Grisham MB, Koyama S, Robbins RA (2000) Reactive nitrogen and oxygen species attenuate interleukin-8-induced neutrophil chemotactic activity in vitro. *J Biol Chem* 275: 10826–10830.
 67. van Pelt LJ, Huisman MV, Weening RS, von dem Borne AE, Roos D, et al. (1996) A single dose of granulocyte-macrophage colony-stimulating factor induces systemic interleukin-8 release and neutrophil activation in healthy volunteers. *Blood* 87: 5305–5313.
 68. Segal BH, Ding L, Holland SM (2003) Phagocyte NADPH oxidase, but not inducible nitric oxide synthase, is essential for early control of *Burkholderia cepacia* and *Chromobacterium violaceum* infection in mice. *Infect Immun* 71: 205–210.
 69. Messina CGM, Reeves EP, Roes J, Segal AW (2002) Catalase negative *Staphylococcus aureus* retain virulence in mouse model of chronic granulomatous disease. *FEBS Lett* 518: 107–110.
 70. Belaouaj A, McCarthy R, Baumann M, Gao Z, Ley TJ, et al. (1998) Mice lacking neutrophil elastase reveal impaired host defence against gram negative bacterial sepsis. *Nat Med* 4: 615–618.
 71. Tkalcevic J, Novelli M, Phylactides M, Iredale JP, Segal AW, et al. (2000) Impaired immunity and enhanced resistance to endotoxin in the absence of neutrophil elastase and cathepsin G. *Immunity* 12: 201–210.
 72. Mastroeni P, Vazquez-Torres A, Fang FC, Xu Y, Khan S, et al. (2000) Antimicrobial actions of the NADPH phagocyte oxidase and inducible nitric oxide synthase in experimental salmonellosis. II. Effects on microbial proliferation and host survival in vivo. *J Exp Med* 192: 237–248.
 73. Ogawa T, Asai Y, Yamamoto H, Taiji Y, Jinno T, et al. (2000) Immunobiological activities of a chemically synthesized lipid A of *Porphyromonas gingivalis*. *FEMS Immunol Med Microbiol* 28: 273–281.
 74. O'Reilly M, Newcomb DE, Remick D (1999) Endotoxin, sepsis, and the primrose path. *Shock* 12: 411–420.
 75. Kobayashi SD, Vovich JM, Braughton KR, Whitney AR, Nauseef WM, et al. (2004) Gene expression profiling provides insight into the pathophysiology of chronic granulomatous disease. *J Immunol* 172: 636–643.
 76. Reth M (2002) Hydrogen peroxide as second messenger in lymphocyte activation. *Nat Immunol* 3: 1129–1134.
 77. Perez HD, Kelly E, Elfman F, Armitage G, Winkler J (1991) Defective polymorphonuclear leukocyte formyl peptide receptor(s) in juvenile periodontitis. *J Clin Invest* 87: 971–976.
 78. Cainciola LJ, Genco RJ, Patters MR, McKenna J, van Oss CJ (1977) Defective polymorphonuclear leukocyte function in a human periodontal disease. *Nature* 265: 445–447.
 79. Hewitt C, McCormick D, Linden G, Turk D, Stern I, et al. (2004) The role of cathepsin C in Papillon-Lefavre syndrome, prepubertal periodontitis, and aggressive periodontitis. *Hum Mutat* 23: 222–228.
 80. Putsep K, Carlsson G, Boma HG, Andersson M (2002) Deficiency of antibacterial peptides in patients with morbus Kostmann: An observation study. *Lancet* 360: 1144–1149.
 81. Gronert K, Kantarci A, Levy BD, Clish CB, Odparlik S, et al. (2004) A molecular defect in intracellular lipid signaling in human neutrophils in localized aggressive periodontal tissue damage. *J Immunol* 172: 1856–1861.
 82. Miyasaki KT. (1991) The neutrophil: mechanisms of controlling periodontal bacteria. *J Periodontol* 62: 761–774.
 83. White NJ (2003) Melioidosis. *Lancet* 361: 1715–1722.
 84. Eberl L, Tummeler B (2004) *Pseudomonas aeruginosa* and *Burkholderia cepacia* in cystic fibrosis: Genome evolution, interactions and adaptation. *Int J Med Microbiol* 294: 123–131.
 85. Crushell E, Harty S, Sharif F, Bourke B (2004) Enteric *Campylobacter*: purging its secrets? *Pediatr Res* 55: 3–12.
 86. Finegold SM (1977) Anaerobic bacteria in human diseases. New York: Academic Press. 710 p.
 87. Segal AW, Geisow M, Garcia R, Harper A, Miller R (1981) The respiratory of phagocytic cells is associated with a rise in vacuolar pH. *Nature* 290: 406–409.
 88. Baker PJ, Evans RT, Roopenian DC (1994) Oral infection with *Porphyromonas gingivalis* and induced alveolar bone loss in immunocompetent and severe combined immunodeficient mice. *Arch Oral Biol* 39: 1035–1040.