

Lawrence Berkeley National Laboratory

Recent Work

Title

Rupturing of Biological Spores As a Source of Secondary Particles in Amazonia.

Permalink

<https://escholarship.org/uc/item/33v7t47h>

Journal

Environmental science & technology, 50(22)

ISSN

0013-936X

Authors

China, Swarup
Wang, Bingbing
Weis, Johannes
et al.

Publication Date

2016-11-01

DOI

10.1021/acs.est.6b02896

Peer reviewed

Rupturing of Biological Spores As a Source of Secondary Particles in Amazonia

Swarup China,[†] Bingbing Wang,^{†,#} Johannes Weis,[‡] Luciana Rizzo,[§] Joel Brito,^{⊥,∇} Glauber G. Cirino,^{||} Libor Kovarik,[†] Paulo Artaxo,[⊥] Mary K. Gilles,[‡] and Alexander Laskin^{*,†}

[†]Environmental Molecular Sciences Laboratory, Pacific Northwest National Laboratory, Richland, Washington 99354, United States

[‡]Chemical Sciences Division, Lawrence Berkeley National Laboratory, Berkeley, California 94720, United States


[§]Federal University of São Paulo, São Paulo - SP, 04021-001, Brazil

^{||}National Institute of Research in Amazonia, Manaus - AM, 69067-375, Brazil

[⊥]Institute of Physics, University of São Paulo, São Paulo - SP, 05508-900, Brazil

Supporting Information

ABSTRACT: Airborne biological particles, such as fungal spores and pollen, are ubiquitous in the Earth's atmosphere and may influence the atmospheric environment and climate, impacting air quality, cloud formation, and the Earth's radiation budget. The atmospheric transformations of airborne biological spores at elevated relative humidity remain poorly understood and their climatic role is uncertain. Using an environmental scanning electron microscope (ESEM), we observed rupturing of Amazonian fungal spores and subsequent release of submicrometer size fragments after exposure to high humidity. We find that fungal fragments contain elements of inorganic salts (e.g., Na and Cl). They are hygroscopic in nature with a growth factor up to 2.3 at 96% relative humidity, thus they may potentially influence cloud formation. Due to their hygroscopic growth, light scattering cross sections of the fragments are enhanced by up to a factor of 10. Furthermore, rupturing of fungal spores at high humidity may explain the bursting events of new particle formation in Amazonia.



INTRODUCTION

Aerosolized biological particles significantly influence the biosphere, atmosphere, and public health.^{1–4} Biological particles impact cloud dynamics and hydrological cycles by forming clouds and ice crystals.^{1,5–7} They influence the Earth's energy budget by scattering and absorbing solar radiation.^{1,8} Primary biological particles, emitted directly from the biosphere, are pollen, fungal spores, bacteria, and fragments of plants and living organisms. High discrepancies exist in estimation of primary biological particle emissions,^{1,9,10} ranging between 56 and 1000 Tg yr⁻¹. Aircraft and balloon measurements suggest that they can be transported to high altitudes and over long distances.^{2,11–13} The physical dimensions of atmospheric biological particles span several orders of magnitude with diameters ranging from nanometers (e.g., viruses) to hundreds of micrometers (e.g., pollen grains, plant debris). Fungal spores and their fragments are one of the most abundant classes of biological particles in various environments.^{2,14–16} In tropical areas, such as the Amazon basin, primary biological particles contribute up to 80% of coarse mode (2–3 μm) particle mass concentration.^{2,6,17} Even at a high altitude site in North America biological particles contribute an average of 40% of the particulate organic carbon mass.¹⁸ The global average loading and emission rates of fungal spores are ~1 μg m⁻³ and ~50 Tg yr⁻¹, respectively.² Fungi can actively discharge their spores via liquid jets into the air, known as actively wet discharged spores (e.g., Ascomycota and

Basidiomycota).^{2,19} Concentrations of these wet discharged spores tend to increase during humid conditions such as during the wet season in the Amazon basin.²

Chemical compositions of primary biological particles are highly variable and remain insufficiently characterized even at a phenomenological level due to difficulties distinguishing between biological and other carbonaceous particles.^{20–22} Therefore, it is currently assumed that actual contributions of primary biological particles to the total atmospheric aerosol are underestimated.^{23,24} Biological materials are primarily carbonaceous and produced from metabolic activity of fungi and bacteria.^{25,26} Sugar alcohols such as mannitol, arabitol, and ergosterol are commonly used as tracers for source apportionment measurements of primary biological particles, such as fungal spores.^{25,26} For example, the average concentrations of mannitol were almost 2–3 times higher in the Amazon basin compared to extratropical locations.²

Previous studies reported that coarse pollen grains (5–150 μm) rupture under high humidity and release cytoplasmic material ranging in size from several nanometers to several micrometers.^{3,4,7,27} However, these submicrometer or nanoparticulates are difficult to detect by traditional bioaerosol

Received: June 9, 2016

Revised: October 4, 2016

Accepted: October 17, 2016

Published: October 17, 2016




Figure 1. Representative SEM images of fungal spores collected in the Amazon basin. Scale bar is 2 μm .

sampling and analytical techniques due to their small size.²⁸ Pollen grains are often covered with amorphous layers and Raman spectroscopy measurements show that the chemical composition of these layers significantly differs between species.²⁹ Many of the fragmented submicrometer particles (subpollen) are starch granules that contribute to atmospheric organic carbon.^{7,29–31} These subpollen particles can act as cloud condensation nuclei⁷ and ice nuclei.^{32,33} In contrast, the environmental impact and cloud formation potential of expelled particles from relatively smaller sizes (1–6 μm) fungal spores has not been recognized.

Because of the smaller size of fungal spores compared to pollens, they have longer atmospheric lifetimes and can be lofted to the mid and upper troposphere. Their abundance depends on environmental factors such as annual season, rain, thunderstorms, wind and temperature.^{1,2,34,35} For example, high ambient concentrations of biological particles are associated with rainfall events.³⁶ Aircraft measurements over central China showed higher concentrations of fungal spore tracers in spring than in summer.³⁷ They have relatively high number concentrations ($\sim 10^4 \text{ m}^{-3}$) in the continental boundary layer.²⁰ Fungal spores and other primary biological particles contribute up to 80% of coarse mode particles in the Amazon basin and significantly impact hydrological cycle.^{2,17}

Here, we show evidence of rupturing of fungal spores collected in the Amazon. The rupturing and release of submicrometer subfungal fragments are observed after exposure to water vapor and subsequent drying. We discuss the previously unexplored climatic implications of these submicrometer fragments and links to new particle burst observations in central Amazonia.

EXPERIMENTAL SECTION

Samples of atmospheric particles were collected during the wet season (January and February, 2015) at the ZF2 Tower (02°35.3517' S, 60 06.8333' W), a pristine rainforest site in Central Amazonia located 40 km North of Manaus margins. Samples were collected from both below ($\sim 2 \text{ m}$ above ground) and above the canopy (39 m above ground). The residence time of air parcel within the forest canopy is on the order of minutes but it can vary depending on the turbulence level. During nighttime turbulence level is low and that limit the transport of air parcel from above to within the canopy. However, during transition of nighttime to daytime, turbulence

progressively increased and reaches the lower half of the canopy $\sim 1.5 \text{ h}$ after sunrise.³⁸ Particles were collected onto 400 mesh transmission electron microscopy (TEM) grids coated with Carbon Type-B films (Ted Pella, Inc.) using 10-stage Micro-Orifice Uniform Deposition Impactors (MOUDI; model 110-R, MSP, Inc.). This study focuses on the samples from stage 4 (size range: 3.2–5.6 μm) where the relative abundance of biological particles is higher than on the other stages. An environmental scanning electron microscope (ESEM, Quanta 3D model, FEI, Inc.) with a Peltier cooling stage was used for water vapor exposure experiments. Fungal spores on the substrate were investigated before water vapor exposure experiments and only intact particles were monitored during experiments. We note that potential caveats of substrate based approach cannot be avoided such as possible damage and morphology modifications of particles upon impaction. A scanning transmission electron detector was used for the ESEM imaging.³⁹ ESEM experiments were conducted at 278–280 K and 0.08–6.50 Torr of water vapor, corresponding to sample exposures of 1–99% relative humidity (RH). Samples were systematically exposed to high RH ($\sim 98\%$) for time periods of 0.5–10 h. Between exposures, samples were dried (to RH $\sim 1\%$) by decreasing the water vapor pressure (to 0.1 Torr) of the ESEM chamber for an interval of 1–2 h. In this study a total of eight samples were investigated. Samples from above and below canopy were collected simultaneously for same duration at the same location. Table S1 (Supporting Information (SI)) shows the sampling time, duration and duty cycle. After prolonged exposure to water vapor, fungal spores ruptured and released submicrometer fragments. We dried the sample inside the ESEM chamber after rupturing of fungal spores. Later we characterized the fragments in dry condition to determine their elemental compositions and morphologies. Furthermore, we conducted dynamic hydration experiments using ESEM to study the hygroscopic behavior and growth of fungal fragment particles. Hydration experiments were conducted up to 96% RH (6.32 Torr of water vapor). Computer-controlled scanning electron microscopy with energy-dispersive X-ray (CCSEM/EDX)⁴⁰ was used to investigate the morphology and elemental composition of biological particles. TEM imaging and electron energy loss spectroscopy (EELS) measurements were performed in scanning mode using an aberration-corrected transmission electron microscope (FEI, Inc. model Titan 80–300) operated at 300 kV. Scanning transmission X-ray




Figure 2. Representative SEM images of ruptured fungal spores and expulsion of submicrometer fragments. Panel (c) shows an example of germination pores of fungal spores as indicated by arrows. Scale bar is 2 μm .

microscopy (STXM)⁴¹ was used to map specific elements (e.g., carbon, sodium) within individual biological particles prior to and post exposure to high relative humidity and subsequent drying.

Furthermore, we used size distribution data from a scanning mobility particle sizer (custom-built SMPS system, Lund University) collected at the ZF2 site during the wet season (Jan–Jun) in the years of 2008–2009, comprising measurements from 133 days⁴² to investigate observed bursting events of nanoparticles in the Amazon basin. Particle size distribution measurements were performed 10 m above the canopy top. All measurements were performed under dry conditions (RH 30–40%), assured by an automatic diffusion dryer.⁴³

RESULTS AND DISCUSSION

Size Distribution and Chemical Composition of Fungal Spores. Typically, fungal spores are a few micrometers in size, often spherical, rod-like or spheroidal (prolate) in shape.^{17,44,45} Figure 1 shows representative morphologies of fungal spores found in the Amazonian samples. Spheroidal fungal spores are 3–7 μm long and 2–4 μm wide. Sizes (area equivalent diameters) of the fungal spore particles range from 1.1 to 5.9 μm with an average diameter of 2.8 μm . Previous studies found presence of several species of fungal spore in the Amazon rainforest canopy with different sizes.^{46,47} These studies applied culture-independent approaches to measure the composition of total and active atmospheric fungal spores over the Amazon forest canopy. An RNA-based approach was also applied to measure metabolically active microbial communities in the atmosphere. Phyla Ascomycota and Basidiomycota were most abundant in total airborne fungal communities with relative abundance of Basidiomycota over 90%. Within the Basidiomycota, Agaricomycetes were most abundant class. However, Ascomycota found to be major fraction (mean relative abundance ~ 80%) within the active community in the atmosphere over the Amazon forest canopy. Sordariomycetes (~27%) and Lecanoromycetes (~18%) were most abundant classes within the Ascomycota community. Overall, these studies found presence of potentially active fungi in the atmosphere, including lichen fungi (class Lecanomyces) and the following genera: Agaricus; Amanita; Aspergillus; Boletus; Cladonia; Lepsita; Mortierella; Puccinia; and Rhizopus.^{46,47}

Elemental composition analysis by CCSEM/EDX shows that fungal spores are primarily composed of C and O. Other frequent elements included N, Na, P, K, S; some fungal spores also contain Cl and Mg. Similar elemental compositions of biological particles have been reported previously.⁴⁸ SI Figure S1 shows an example of EDX elemental maps of a typical fungal spore particle along with its corresponding EDX spectrum. The C map indicates the fungal spore contains significant carbonaceous content. Within this primary unprocessed particle, the elements are relatively homogeneously distributed, consistent with measurements of other fungal spores found in the samples. However, the fractions of different elements vary considerably among fungal spore particles (SI Figure S2).

In this study, fungal spores were identified based on their characteristic shapes and chemical composition, which includes a high carbonaceous component and the presence of other elements such as phosphorus.^{17,42,49} Fungal spore particles contributed up to 56% and 17% of the total number of particles in the size range of 3.2–5.6 μm (aerodynamic diameter) below and above the canopy, respectively.

Rupture of Fungal Spores. Rupturing of fungal spores and subsequent expulsion of fungal fragments was observed after exposure to high RH conditions (~98%) for ~10 h and subsequent drying. Representative examples of fragmented and expelled fungal spores after wet and drying cycles are shown in Figure 2. Substantial rupturing was not observed in the samples exposed to high RH for shorter time periods that were subsequently dried. These results suggest that rupture occurred during the prolonged wet environment, likely repeated several times, and not due to surface tension forces during a single hydration/dehydration cycle.⁷ Rupturing of fungal spores at high humidity may impact the culturability of the fungal spores. Hydrated fungal spores may contain significant amount of osmotically active solutions. At high humidity osmotic pressure can be developed across fungal walls and once it is high enough then spores can rupture.⁵⁰ Figure 3 shows the size (area equivalent diameter) distributions of original fungal spores and expelled fungal fragments after rupture. Here, only individual particles that did not agglomerate during the drying process were used for the analysis. Fungal spores release fungal fragments in a broad range of sizes, from 10s of nanometers up to a micrometer. Fragmented particles smaller than 10 nm may also be present, but they cannot be unambiguously detected due to the imaging limits of ESEM. This study motivates the




Figure 3. Size distribution of fungal spore particles (blue line) and fragmented subfungal spore particles (green line). A total of 711 individual fungal spores and 2041 fragment particles were identified by microscopy and used for the analysis.

need for further investigation of rupturing of airborne biological spores that can provide quantitative information about size and number concentration of fragments, as well as more accurate information about the environmental conditions for the rupturing process. Previous studies found that cytoplasmic material from ruptured pollen grains can be in the range of 0.05–2 μm in size.^{27,34}

The size and chemical composition of fragmented particles significantly influences their cloud activation potential. ESEM images reveal that during rupture the number of particles released ranges from 10 to $\sim 10^3$ individual fragmented particles per fungal spore. Fragmented particles show distinguishable variations in their compositions and morphologies (Figure 2). In the Amazon, the presence of a variety of fungal spores with different compositions and chemical aging results in a wide range of fragmented particles. Figure 4a–c shows representative STXM images at the Na pre edge (1070 eV), peak (1078 eV) and optical density map of biological particle after exposure to high RH. Na pre edge image (Figure 4a) represents the optical density for non-Na elements over the investigated sample area and Na peak represents the maximum Na absorption. The

optical density ($-\ln(I_d)/I_0$) is calculated based on the measured intensity (I_d) using Beer–Lambert’s law.⁵¹ Transmission intensity through a particle free region of the substrate is used to obtain I_0 . The STXM map shows the presence of Na containing fragments and inclusions of fungal spores. Furthermore, EDX spectra of fragmented particles show signatures of inorganic salts, for example, such as Na and Cl (Figure 4d,e) and relatively low C compared to primary spore. We hypothesize that inorganic salts within the primary spore will be in liquid phase at high humidity and these salts can disperse from the primary fragments. This process may result in relatively higher inorganic salt and less carbon in the fragmented spores compared to the primary spore. A previous study reported an elemental composition of C and O in submicrometer (200 nm in diameter) fragmented pollen particles and the presence of carbohydrates and proteins in organic pollen solutions.⁷

Previously, an amorphous film around fragmented pollen was reported.²⁹ Chemical characteristics of the fungal spore walls and their thickness may influence their rupturing process. Water vapor can diffuse and penetrate the film thickness of the wall and facilitate the expansion of the layer, resulting in an increase of the adhesion forces, thus affecting the rupturing process.²⁸ Hence, understanding their wetting behavior requires characterization of the fungal spore walls. For example, pollen walls (exines) are composed of several types of proteins, lipids, and sporopollenin.³⁰ Previous studies found glucuronic acid, uronic acid, and heteropolymers of mannose, galactose, glucose, and glucuronic acid in several kinds of fungal spore walls.^{52,53} These components and their concentrations vary among fungal spore types. The measured width of the film around the spore wall from SEM images ranged from 0.03 to 0.58 μm and increased to 0.11–0.78 μm at elevated relative humidity (SI Figure S4). Saccharide compounds in the fungal spore wall, such as glucose, can lead to the growth of the film at high RH. Similar to other carbonaceous particles, fungal spores contain high carbon, thus TEM/EELS analysis can provide further information on chemical bonding. EELS analysis shows




Figure 4. Representative STXM images at the (a) Na K pre-edge (1070 eV), (b) peak (1078 eV) and (c) optical density map of Na-containing biological particle (optical density difference between Na K peak and pre-edge). SEM image (d) of a ruptured fungal spore particle and its fragments after water exposure; (e) the EDX spectra acquired over different regions of the particle (marked by boxes in the SEM image) show the presence of elemental components representative of salts (i.e. Na and Cl) in the fragmented particles. The Cu peak in the EDX spectrum is from the substrate.

considerable differences in aromatic carbon (exhibited by the intensity of π^* peak) between fungal spores and carbonaceous soot particles (SI Figure S3). Fungal spores show relatively weaker π^* peak compared to soot particles. Furthermore, substantial differences in the intensity of the carbon π^* peak between various fungal spores are also noticeable. For example, our experimental observations suggest that fungal spores with weaker π^* peak (less aromatic) are more susceptible to rupture and release higher number of fragments compared to those with stronger π^* peak (SI Figure S3). Overall, TEM/EELS analysis suggests that structural variability within various fungal spores and their walls may determine the extent and specific conditions for rupturing.

Spore morphology can also influence the rupturing process and the release of materials. For example, elongated (high aspect ratio) fungal spores (e.g., *Aspergillus* and *Penicillium*) are more susceptible to stress compared to spores with less elongated structures (*Cladosporium*).²⁸ Similarly, our limited observations support the hypothesis that elongated particles are more likely to rupture than less elongated particles (Figure 2). Hence, elongated particles release a higher number of submicrometer particles.

Climatic Impacts. Aircraft measurements suggest that fungal spores can be uplifted from the ground surface to the mid-troposphere³⁷ and potentially impact climate. In locations where the ambient relative humidity is high enough (such as the Amazon) fungal spores can rupture and release fungal fragments. Expelled submicrometer particles can potentially be transported to the free troposphere by tropical convection and further impact cloud properties. The number, mass, size, and chemical compositions of fine biological particles are insufficiently quantified and remain highly uncertain for the Amazon basin.^{20–22,54} Our results suggest that expelled fungal fragments may contribute to the fine aerosol fraction. We note that temperature and ambient pollutants (e.g., O_3) during the laboratory experiments were not similar to those in the Amazon basin. Other environmental conditions (e.g., temperature, wind speed and concentrations of ambient pollutants) may also influence rupturing spores under wet conditions in the Amazon basin. However, the release of the small fungal fragments may not be easily detected by in situ techniques such as fluorescence UV-APS due to instrument detection limits.¹⁷

The hygroscopicity of submicrometer fungal fragments was investigated using ESEM. From ESEM images collected in the hydration experiments, area equivalent diameter growth factors ($\text{diameter}_{\text{wet}}/\text{diameter}_{\text{dry}}$) of fungal fragments were estimated in the initial (dry) size range of 150 to 600 nm (SI Figure S5). Since fungal fragments contain inorganic salt elements (e.g., Na and Cl) and fungal spore walls contain sugars, growth factors of fungal fragments are compared with those of NaCl and D-glucose. Figure 5 shows the growth factors of fungal fragments compared with hygroscopic characteristics of pure NaCl and D-glucose submicrometer particles.⁵⁵ Results suggest that the hygroscopicity of fungal fragments lies between NaCl and D-glucose particles within an RH range of 70–96%, indicating that fungal fragments are more hygroscopic than D-glucose particles. The growth factors of fungal fragments approach those of NaCl particles at high RH (~96%). The growth factors of fungal fragments are 1.18, 1.35, 1.64, and 2.31 at RH of 60, 76, 85, and 96%, respectively (Figure 5). As discussed earlier, with increasing RH, the fungal spore film width also increases (SI Figure S4). However, previous studies found aerodynamic diameter growth factors of 1.06–1.27 at




Figure 5. Area equivalent diameter growth factors ($\text{diameter}_{\text{wet}}/\text{diameter}_{\text{dry}}$) of fragmented subfungal spore particles (green line). Gray line indicates the growth factor of the laboratory generated NaCl particles. Black line indicates the growth factor of D-glucose from Mochida and Kawamura⁵⁵ using tandem differential mobility analyzer.

98% for 1.6–3.3 μm fungal spore particles.^{1,56,57} Overall, our results indicate that fungal fragments are hygroscopic in nature and can participate in cloud formation. Similarly, a recent study showed that fragmented pollen grains are active as cloud condensation nuclei, requiring supersaturations of 0.81 and 0.12 for 50 and 200 nm particles, respectively.⁷

The increase in light scattering cross section due to hygroscopic growth of the fragmented particles at elevated RH can be estimated using Mie theory.^{58,59} For these calculations a wavelength of 550 nm was used. The refractive indices of the hydrated particles are derived using the volume mixing rule,⁶⁰ with a real part of the refractive index of 1.40 for biological particles⁴⁸ and 1.33 for water.⁶¹ At 60% RH, the scattering cross section of fungal fragments increased by a factor of 1.2, compared to dry conditions. The scattering cross section of fragments increases with increasing RH, meaning that a larger growth factor leads to a larger scattering cross section. The increases in scattering cross section of fungal fragments are 1.2, 1.9, 4.3, 8.6, and 10.3 at relative humidities of 60, 76, 85, and 96%, respectively (SI Figure S6). These properties could allow fragmented particles to have a direct radiative effect on climate.

Mechanisms of the new particle formation in the Amazon basin remains insufficiently understood and occurs much less frequently than reported in a boreal forest.⁶² Furthermore, the Amazon basin has one of the lowest aerosol concentrations in any continental regions,^{21,42} and as such the release of nano- and submicrometer particles from fungal spores can strongly influence aerosol concentration and thus its role on clouds. Nucleation events and subsequent growth of nucleated nanoparticles (3–10 nm) to larger sizes are regularly observed in other continental locations.^{21,42} Bursting events of nanoparticles (SI Figure S7) in the size range of 10–50 nm occur frequently during the wet season in Amazonia.⁴² However, near surface measurements revealed no significant evidence of regional scale new particle formation from gas-particle nucleation events in the Amazon basin.²¹ A large amount of water vapor emitted from the forest makes the Amazon basin very humid, often >70% RH and in addition deep vertical convection is notorious in this tropical area.⁴² Figure 6 shows the occurrence of bursting events in Amazon basin measured at the ZF2 site during the wet season of 2008–2009. The median diurnal variation of N_{50} concentrations (SI Figure S7 (b)) indicates that particles with diameters less than 50 nm are more frequently observed at nighttime, when ambient relative humidity reaches its highest values close to saturation (SI




Figure 6. Bursting events in the Amazon basin at different RHs.

Figure S7 (c)) and no photochemistry occurs. Figure 6 illustrates that the occurrence of bursting events increases with increasing RH, which is in accordance with our observation of rupturing of spores at high RH. Expulsion of the fine particles from fungal spores under wet conditions in the Amazon basin, and/or outflow from deep convective clouds (in-cloud processing of fungal spores) could be common and may provide insight into new particle formation (see discussion in the SI).

The rupturing process and release of fungal fragments may impact the complex land–atmosphere exchange process (emissions and deposition). The lifetime of the fungal spore particles can increase after their rupturing and release of submicrometer fragments. This process potentially affects the deposition of biological particles. Deposition of biological particles may influence metabolic activity and trigger reproduction of biological spores. These activities can either facilitate or hinder further emission of biological particles, ultimately may affect biogeochemical cycle and related feedback loops.⁶³ Finally, it may alter the feedback loops of the exchange processes between the atmosphere and soil, and long-range transport of particles. Furthermore, release of compounds from the biological particles by a rupturing process is typically ignored,² but can substantially influence budget calculations of biological particles. Future studies should focus on (1) detailed characterization and improved understanding of the chemical composition of different types of fungal spores and the variability of their fragments; (2) systematic evaluation of specific environmental conditions promoting rupturing events (e.g., RH, temperature, atmospheric pressure, humidity cycling rates).

■ ASSOCIATED CONTENT

📄 Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.est.6b02896.

Additional information as noted in the text (PDF)

■ AUTHOR INFORMATION

Corresponding Author

*Phone: +1 509 371-6129; e-mail: alexander.laskin@pnnl.gov .

Present Addresses

[#](B.W.) State Key Laboratory of Marine Environmental Science and College of Ocean and Earth Sciences, Xiamen University, Xiamen, China.

[∇](J.B.) Laboratory for Meteorological Physics, University Blaise Pascal, Clermont-Ferrand, France.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

The Pacific Northwest National Laboratory (PNNL) group acknowledges support from the Chemical Imaging Initiative of the Laboratory Directed Research and Development program at PNNL. The Lawrence Berkeley National Laboratory (LBNL) group acknowledges support from the U.S. Department of Energy's Atmospheric System Research program, an Office of Science, Office of Biological and Environmental Research (OBER). The CCSEM/EDX and ESEM analyses were performed at Environmental Molecular Sciences Laboratory, a national scientific user facility sponsored by OBER at PNNL. PNNL is operated by the U.S. Department of Energy by Battelle Memorial Institute under contract DE-AC06-76RL0. STXM/NEXAFS analysis at beamline 11.0.2 of the Advanced Light Source at Lawrence Berkeley National Laboratory is supported by the Director, Office of Science, Office of Basic Energy Sciences of the U.S. Department of Energy under Contract No. DE-AC02-05CH11231. Beamline 11.0.2 also acknowledges support from the Office of Basic Energy Sciences Division of Chemical Sciences, Geosciences, and Biosciences by the Condensed Phase and Interfacial Molecular Sciences Program of the U.S. Department of Energy. Brazilian group acknowledges support from the FAPESP projects 08/58100-2, 2013/25058-1 and 2013/05014-0. We thank Dr. Erik Swietlicki's group at Lund University for the lending the SMPS system and Simone R. da Silva for assistance with sampling. We also thank INPA (Instituto Nacional de Pesquisas da Amazônia) for the logistical support.

■ REFERENCES

- (1) Després, V. R.; Huffman, J. A.; Burrows, S. M.; Hoose, C.; Safatov, A. S.; Buryak, G.; Fröhlich-Nowoisky, J.; Elbert, W.; Andreae, M. O.; Pöschl, U.; Jaenicke, R., Primary biological aerosol particles in the atmosphere: a review. *Tellus, Ser. B* **2012**, *64*, 10.3402/tellusb.v64i0.15598
- (2) Elbert, W.; Taylor, P. E.; Andreae, M. O.; Pöschl, U. Contribution of fungi to primary biogenic aerosols in the atmosphere: wet and dry discharged spores, carbohydrates, and inorganic ions. *Atmos. Chem. Phys.* **2007**, *7* (17), 4569–4588.
- (3) Grote, M.; Valenta, R.; Reichelt, R. Abortive pollen germination: a mechanism of allergen release in birch, alder, and hazel revealed by immunogold electron microscopy. *J. Allergy Clin. Immunol.* **2003**, *111* (5), 1017–1023.
- (4) Taylor, P. E.; Flagan, R. C.; Valenta, R.; Glovsky, M. M. Release of allergens as respirable aerosols: a link between grass pollen and asthma. *J. Allergy Clin. Immunol.* **2002**, *109* (1), 51–56.
- (5) Murray, B.; Ross, J.; Whale, T.; Price, H.; Atkinson, J.; Umo, N.; Webb, M., The relevance of nanoscale biological fragments for ice nucleation in clouds. *Sci. Rep.* **2015**, 5.808210.1038/srep08082
- (6) Pöschl, U.; Martin, S.; Sinha, B.; Chen, Q.; Gunthe, S.; Huffman, J.; Borrmann, S.; Farmer, D.; Garland, R.; Helas, G. Rainforest aerosols as biogenic nuclei of clouds and precipitation in the Amazon. *Science* **2010**, *329* (5998), 1513–1516.
- (7) Steiner, A. L.; Brooks, S. D.; Deng, C.; Thornton, D. C.; Pendleton, M. W.; Bryant, V. Pollen as atmospheric cloud condensation nuclei. *Geophysical research letters* **2015**, *42*, 3596.
- (8) Guyon, P.; Graham, B.; Roberts, G. C.; Mayol-Bracero, O. L.; Maenhaut, W.; Artaxo, P.; Andreae, M. O. Sources of optically active aerosol particles over the Amazon forest. *Atmos. Environ.* **2004**, *38* (7), 1039–1051.
- (9) Jaenicke, R. Abundance of Cellular Material and Proteins in the Atmosphere. *Science* **2005**, *308* (5718), 73.

- (10) Penner, J. E.; Andreae, M.; Annegarn, H.; Barrie, L.; Feichter, J.; Hegg, D.; Jayaraman, A.; Leaitch, R.; Murphy, D.; Nganga, J., Aerosols, their direct and indirect effects. In *Climate Change 2001: The Scientific Basis. Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change*; Houghton, J. T.; Ding, Y., Griggs, D. J.; Noguer, M.; P. J., van der Linden; Dai, X.; Maskell, K.; Johnson, C. A., Eds.; Cambridge University Press: Cambridge, 2001; pp 289–348.
- (11) Prospero, J. M.; Blades, E.; Mathison, G.; Naidu, R. Interhemispheric transport of viable fungi and bacteria from Africa to the Caribbean with soil dust. *Aerobiologia* **2005**, *21* (1), 1–19.
- (12) Wainwright, M.; Wickramasinghe, N. C.; Narlikar, J.; Rajaratnam, P. Microorganisms cultured from stratospheric air samples obtained at 41 km. *FEMS Microbiol. Lett.* **2003**, *218* (1), 161–165.
- (13) Meier, F. C.; Artschwager, E. Airplane collections of sugar-beet pollen. *Science* **1938**, *88* (2291), 507–508.
- (14) Bauer, H.; Schueller, E.; Weinke, G.; Berger, A.; Hitzemberger, R.; Marr, I. L.; Puxbaum, H. Significant contributions of fungal spores to the organic carbon and to the aerosol mass balance of the urban atmospheric aerosol. *Atmos. Environ.* **2008**, *42* (22), 5542–5549.
- (15) Bowers, R. M.; Clements, N.; Emerson, J. B.; Wiedinmyer, C.; Hannigan, M. P.; Fierer, N. Seasonal Variability in Bacterial and Fungal Diversity of the Near-Surface Atmosphere. *Environ. Sci. Technol.* **2013**, *47* (21), 12097–12106.
- (16) Crawford, C.; Reponen, T.; Lee, T.; Iossifova, Y.; Levin, L.; Adhikari, A.; Grinshpun, S. A. Temporal and spatial variation of indoor and outdoor airborne fungal spores, pollen, and (1→3)- β -D-glucan. *Aerobiologia* **2009**, *25* (3), 147–158.
- (17) Huffman, J. A.; Sinha, B.; Garland, R. M.; Snee-Pollmann, A.; Gunthe, S. S.; Artaxo, P.; Martin, S. T.; Andreae, M. O.; Pöschl, U. Size distributions and temporal variations of biological aerosol particles in the Amazon rainforest characterized by microscopy and real-time UV-APS fluorescence techniques during AMAZE-08. *Atmos. Chem. Phys.* **2012**, *12* (24), 11997–12019.
- (18) Wiedinmyer, C.; Bowers, R. M.; Fierer, N.; Horanyi, E.; Hannigan, M.; Hallar, A. G.; McCubbin, I.; Baustian, K. The contribution of biological particles to observed particulate organic carbon at a remote high altitude site. *Atmos. Environ.* **2009**, *43* (28), 4278–4282.
- (19) Pringle, A.; Patek, S. N.; Fischer, M.; Stolze, J.; Money, N. P. The captured launch of a ballistospore. *Mycologia* **2005**, *97* (4), 866–871.
- (20) Després, V. R.; Nowojsky, J. F.; Klose, M.; Conrad, R.; Andreae, M. O.; Pöschl, U. Characterization of primary biogenic aerosol particles in urban, rural, and high-alpine air by DNA sequence and restriction fragment analysis of ribosomal RNA genes. *Biogeosciences* **2007**, *4* (6), 1127–1141.
- (21) Martin, S. T.; Andreae, M. O.; Artaxo, P.; Baumgardner, D.; Chen, Q.; Goldstein, A. H.; Guenther, A.; Heald, C. L.; Mayol-Bracero, O. L.; McMurry, P. H.; Pauliquevis, T.; Pöschl, U.; Prather, K. A.; Roberts, G. C.; Saleska, S. R.; Silva Dias, M. A.; Spracklen, D. V.; Swietlicki, E.; Trebs, I. Sources and properties of Amazonian aerosol particles. *Rev. Geophys.* **2010**, *48* (2), RG2002.
- (22) Pöschl, U. Atmospheric aerosols: composition, transformation, climate and health effects. *Angew. Chem., Int. Ed.* **2005**, *44* (46), 7520–7540.
- (23) Jaenicke, R.; Matthias-Maser, S.; Gruber, S. Omnipresence of biological material in the atmosphere. *Environmental Chemistry* **2007**, *4* (4), 217–220.
- (24) Hummel, M.; Hoose, C.; Gallagher, M.; Healy, D. A.; Huffman, J. A.; O'Connor, D.; Pöschl, U.; Pöhlker, C.; Robinson, N. H.; Schnaiter, M.; Sodeau, J. R.; Stengel, M.; Toprak, E.; Vogel, H. Regional-scale simulations of fungal spore aerosols using an emission parameterization adapted to local measurements of fluorescent biological aerosol particles. *Atmos. Chem. Phys.* **2015**, *15* (11), 6127–6146.
- (25) Di Filippo, P.; Pomata, D.; Riccardi, C.; Buiarelli, F.; Perrino, C. Fungal contribution to size-segregated aerosol measured through biomarkers. *Atmos. Environ.* **2013**, *64*, 132–140.
- (26) Nirmalkar, J.; Deb, M.; Tsai, Y.; Deshmukh, D. Arabitol and Mannitol as Tracer for Fungal Contribution to Size-Differentiated Particulate Matter of Rural Atmospheric Aerosols. *Int. J. Environ. Sci. Dev.* **2015**, *6* (6), 460.
- (27) Taylor, P.; Flagan, R.; Miguel, A.; Valenta, R.; Glovsky, M. Birch pollen rupture and the release of aerosols of respirable allergens. *Clin. Exp. Allergy* **2004**, *34* (10), 1591–1596.
- (28) Górný, R. L.; Reponen, T.; Willeke, K.; Schmechel, D.; Robine, E.; Boissier, M.; Grinshpun, S. A. Fungal fragments as indoor air biocontaminants. *Appl. Environ. Microbiol.* **2002**, *68* (7), 3522–3531.
- (29) Pummer, B. G.; Bauer, H.; Bernardi, J.; Chazallon, B.; Facq, S.; Lendl, B.; Whitmore, K.; Grothe, H. Chemistry and morphology of dried-up pollen suspension residues. *J. Raman Spectrosc.* **2013**, *44* (12), 1654–1658.
- (30) Laurence, A. R.; Thoms, A. V.; Bryant, V. M.; McDonough, C. Airborne starch granules as a potential contamination source at archaeological sites. *Journal of Ethnobiology* **2011**, *31* (2), 213–232.
- (31) Suphioglu, C.; Singh, M. B.; Taylor, P.; Knox, R.; Bellomo, R.; Holmes, P.; Puy, R. Mechanism of grass-pollen-induced asthma. *Lancet* **1992**, *339* (8793), 569–572.
- (32) O'Sullivan, D.; Murray, B. J.; Ross, J. F.; Whale, T. F.; Price, H. C.; Atkinson, J. D.; Umo, N. S.; Webb, M. E. The relevance of nanoscale biological fragments for ice nucleation in clouds. *Sci. Rep.* **2015**, *5*, 8082.
- (33) Pummer, B. G.; Bauer, H.; Bernardi, J.; Bleicher, S.; Grothe, H. Suspensible macromolecules are responsible for ice nucleation activity of birch and conifer pollen. *Atmos. Chem. Phys.* **2012**, *12* (5), 2541–2550.
- (34) Miguel, A. G.; Taylor, P. E.; House, J.; Glovsky, M. M.; Flagan, R. C. Meteorological influences on respirable fragment release from Chinese elm pollen. *Aerosol Sci. Technol.* **2006**, *40* (9), 690–696.
- (35) Visez, N.; Chassard, G.; Azarkan, N.; Naas, O.; Sénéchal, H.; Sutra, J.-P.; Poncet, P.; Choël, M. Wind-induced mechanical rupture of birch pollen: Potential implications for allergen dispersal. *J. Aerosol Sci.* **2015**, *89*, 77–84.
- (36) Huffman, J. A.; Prenni, A. J.; DeMott, P. J.; Pöhlker, C.; Mason, R. H.; Robinson, N. H.; Fröhlich-Nowojsky, J.; Tobo, Y.; Després, V. R.; Garcia, E.; Gochis, D. J.; Harris, E.; Müller-Germann, I.; Ruzene, C.; Schmer, B.; Sinha, B.; Day, D. A.; Andreae, M. O.; Jimenez, J. L.; Gallagher, M.; Kreidenweis, S. M.; Bertram, A. K.; Pöschl, U. High concentrations of biological aerosol particles and ice nuclei during and after rain. *Atmos. Chem. Phys.* **2013**, *13* (13), 6151–6164.
- (37) Fu, P. Q.; Kawamura, K.; Cheng, Y. F.; Hatakeyama, S.; Takami, A.; Li, H.; Wang, W. Aircraft measurements of polar organic tracer compounds in tropospheric particles (PM10) over central China. *Atmos. Chem. Phys.* **2014**, *14* (8), 4185–4199.
- (38) Fuentes, J. D.; Chamecki, M.; Santos, R. M. N. d.; Randow, C. V.; Stoy, P. C.; Katul, G.; Fitzjarrald, D.; Manzi, A.; Gerken, T.; Trowbridge, A.; Freire, L. S.; Ruiz-Plancarte, J.; Maia, J. M. F.; Tóta, J.; Dias, N.; Fisch, G.; Schumacher, C.; Acevedo, O.; Mercer, J. R., Linking meteorology, turbulence, and air chemistry in the Amazon rainforest. *Bull. Am. Meteorol. Soc.* **2016**, *0*, (0). [10.1175/BAMS-D-15-00152.1](https://doi.org/10.1175/BAMS-D-15-00152.1)
- (39) Laskin, A.; Cowin, J. P.; Iedema, M. J. Analysis of individual environmental particles using modern methods of electron microscopy and X-ray microanalysis. *J. Electron Spectrosc. Relat. Phenom.* **2006**, *150* (2–3), 260–274.
- (40) Laskin, A.; Wietsma, T. W.; Krueger, B. J.; Grassian, V. H., Heterogeneous chemistry of individual mineral dust particles with nitric acid: A combined CCSEM/EDX, ESEM, and ICP-MS study. *J. Geophys. Res.* **2005**, *110*, (D10). [10.1029/2004JD005206](https://doi.org/10.1029/2004JD005206)
- (41) Moffet, R. C.; Tivanski, A. V.; Gilles, M. K. In *Fundamentals and Applications in Aerosol Spectroscopy*; Reid, R. S. J. P., Ed. Taylor and Francis Books, Inc: Boca Raton, FL, 2010; pp 419–462.
- (42) Artaxo, P.; Rizzo, L. V.; Brito, J. F.; Barbosa, H. M.; Arana, A.; Sena, E. T.; Cirino, G. G.; Bastos, W.; Martin, S. T.; Andreae, M. O. Atmospheric aerosols in Amazonia and land use change: from natural biogenic to biomass burning conditions. *Faraday Discuss.* **2013**, *165*, 203–235.

- (43) Tuch, T. M.; Haudek, A.; Müller, T.; Nowak, A.; Wex, H.; Wiedensohler, A. Design and performance of an automatic regenerating adsorption aerosol dryer for continuous operation at monitoring sites. *Atmos. Meas. Tech.* **2009**, *2* (2), 417–422.
- (44) Gabey, A. M.; Gallagher, M. W.; Whitehead, J.; Dorsey, J. R.; Kaye, P. H.; Stanley, W. R. Measurements and comparison of primary biological aerosol above and below a tropical forest canopy using a dual channel fluorescence spectrometer. *Atmos. Chem. Phys.* **2010**, *10* (10), 4453–4466.
- (45) Hussein, T.; Norros, V.; Hakala, J.; Petäjä, T.; Aalto, P. P.; Rannik, Ü.; Vesala, T.; Ovaskainen, O. Species traits and inertial deposition of fungal spores. *J. Aerosol Sci.* **2013**, *61*, 81–98.
- (46) Womack, A. M.; Artaxo, P. E.; Ishida, F. Y.; Mueller, R. C.; Saleska, S. R.; Wiedemann, K. T.; Bohannan, B. J. M.; Green, J. L. Characterization of active and total fungal communities in the atmosphere over the Amazon rainforest. *Biogeosciences* **2015**, *12* (21), 6337–6349.
- (47) Fröhlich-Nowoisky, J.; Burrows, S. M.; Xie, Z.; Engling, G.; Solomon, P. A.; Fraser, M. P.; Mayol-Bracero, O. L.; Artaxo, P.; Begerow, D.; Conrad, R.; Andreae, M. O.; Després, V. R.; Pöschl, U. Biogeography in the air: fungal diversity over land and oceans. *Biogeosciences* **2012**, *9* (3), 1125–1136.
- (48) Ebert, M.; Weinbruch, S.; Hoffmann, P.; Ortner, H. M. The chemical composition and complex refractive index of rural and urban influenced aerosols determined by individual particle analysis. *Atmos. Environ.* **2004**, *38* (38), 6531–6545.
- (49) Artaxo, P.; Martins, J. V.; Yamasoe, M. A.; Procópio, A. S.; Pauliquevis, T. M.; Andreae, M. O.; Guyon, P.; Gatti, L. V.; Leal, A. M. C. Physical and chemical properties of aerosols in the wet and dry seasons in Rondônia, Amazonia. *J. Geophys. Res.* **2002**, *107* (D20), 8081.
- (50) Grote, M.; Vrtala, S.; Niederberger, V.; Wiermann, R.; Valenta, R.; Reichelt, R. Release of allergen-bearing cytoplasm from hydrated pollen: a mechanism common to a variety of grass (Poaceae) species revealed by electron microscopy. *J. Allergy Clin. Immunol.* **2001**, *108* (1), 109–115.
- (51) Moffet, R.; AV, T.; Gilles, M., Scanning X-ray transmission microscopy: applications in atmospheric aerosol research. In *Fundamentals and Applications in Aerosol Spectroscopy*, Signorell, R.; Reid, J., Eds.; Taylor & Francis: New York, 2010; pp 419–462.
- (52) Bartnicki-Garcia, S. Cell wall chemistry, morphogenesis, and taxonomy of fungi. *Annu. Rev. Microbiol.* **1968**, *22* (1), 87–108.
- (53) Brown, R.; Lindberg, B. Polysaccharides from cell walls of *Aureobasidium* (Pullularia) pullulans. II. Heteropolysaccharide. *Acta Chem. Scand.* **1967**, *21* (9), 2383.
- (54) Fuzzi, S.; Decesari, S.; Facchini, M. C.; Cavalli, F.; Emblico, L.; Mircea, M.; Andreae, M. O.; Trebs, I.; Hoffer, A.; Guyon, P.; Artaxo, P.; Rizzo, L. V.; Lara, L. L.; Pauliquevis, T.; Maenhaut, W.; Raes, N.; Chi, X.; Mayol-Bracero, O. L.; Soto-García, L. L.; Claeys, M.; Kourtchev, I.; Rissler, J.; Swietlicki, E.; Tagliavini, E.; Schkolnik, G.; Falkovich, A. H.; Rudich, Y.; Fisch, G.; Gatti, L. V. Overview of the inorganic and organic composition of size-segregated aerosol in Rondônia, Brazil, from the biomass-burning period to the onset of the wet season. *J. Geophys. Res.* **2007**, *112* (D1), D01201.
- (55) Mochida, M.; Kawamura, K. Hygroscopic properties of levoglucosan and related organic compounds characteristic to biomass burning aerosol particles. *Journal of Geophysical Research: Atmospheres* **2004**, *109* (D21), D21202.
- (56) Madelin, T. M.; Johnson, H. E. Fungal and actinomycete spore aerosols measured at different humidities with an aerodynamic particle sizer. *J. Appl. Bacteriol.* **1992**, *72* (5), 400–409.
- (57) Reponen, T.; Willeke, K.; Ulevicius, V.; Reponen, A.; Grinshpun, S. A. Effect of relative humidity on the aerodynamic diameter and respiratory deposition of fungal spores. *Atmos. Environ.* **1996**, *30* (23), 3967–3974.
- (58) Bohren, C. F.; Huffman, D. R. *Absorption and Scattering of Light by Small Particles*; John Wiley & Sons: Hoboken, NJ, 1983.
- (59) Mätzler, C., MATLAB functions for Mie scattering and absorption, version 2. *IAP Res. Rep* 2002, 8.
- (60) Abo Riziq, A.; Erlick, C.; Dinar, E.; Rudich, Y. Optical properties of absorbing and non-absorbing aerosols retrieved by cavity ring down (CRD) spectroscopy. *Atmos. Chem. Phys.* **2007**, *7* (6), 1523–1536.
- (61) Seinfeld, J. H.; Pandis, S. N. *Atmospheric Chemistry and Physics: From Air Pollution to Climate Change*; John Wiley & Sons: Hoboken, NJ, 1998.
- (62) Kulmala, M.; Kontkanen, J.; Junninen, H.; Lehtipalo, K.; Manninen, H. E.; Nieminen, T.; Petäjä, T.; Sipilä, M.; Schobesberger, S.; Rantala, P. Direct observations of atmospheric aerosol nucleation. *Science* **2013**, *339* (6122), 943–946.
- (63) Pöschl, U.; Shiraiwa, M. Multiphase Chemistry at the Atmosphere–Biosphere Interface Influencing Climate and Public Health in the Anthropocene. *Chem. Rev.* **2015**, *115* (10), 4440–4475.