

RUSCHEWEYH'S UNIVALENCE CRITERION AND QUASICONFORMAL EXTENSIONS

IKKEI HOTTA

Abstract

Ruscheweyh extended the work of Becker and Ahlfors on sufficient conditions for a normalized analytic function on the unit disk to be univalent there. In this paper we refine the result to a quasiconformal extension criterion with the help of Becker's method. As an application, a positive answer is given to an open problem proposed by Ruscheweyh.

1. Introduction

Throughout the paper, \mathbf{D} denotes the unit disk $\{|z| < 1\}$ in the complex plane \mathbf{C} and \mathbf{D}^* the exterior domain of \mathbf{D} in the Riemann sphere $\hat{\mathbf{C}} = \mathbf{C} \cup \{\infty\}$.

Let \mathcal{A} be a family of normalized analytic functions $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$ on \mathbf{D} . We say that a sense-preserving homeomorphism f of a plane domain $G \subset \mathbf{C}$ is k -quasiconformal if f is absolutely continuous on almost all lines parallel to the coordinate axes and $|f_{\bar{z}}| \leq k|f_z|$, almost everywhere G , where $f_{\bar{z}} = \partial f / \partial \bar{z}$, $f_z = \partial f / \partial z$ and k is a constant with $0 \leq k < 1$.

Ahlfors [1] has shown that the following condition is sufficient for quasiconformal extensibility of univalent functions as an extension of Becker's univalence condition [2] (see also [7], p. 175);

THEOREM A ([1], [3]). *Let $f \in \mathcal{A}$. If there exists a k , $0 \leq k < 1$, such that for a constant $c \in \mathbf{C}$ satisfying $|c| \leq k$ and all $z \in \mathbf{D}$*

$$(1) \quad \left| c|z|^2 + (1 - |z|^2) \frac{zf''(z)}{f'(z)} \right| \leq k$$

then f has a k -quasiconformal extension to \mathbf{C} .

The limiting case $k \rightarrow 1$ in the above theorem ensures univalence of f in \mathbf{D} . Ruscheweyh [8] extended this univalence condition in the following way;

2000 *Mathematics Subject Classification.* Primary 30C62, Secondary 30C45.

Key words and phrases. Löwner(Loewner) chain, quasiconformal mapping, univalent function.

Received November 19, 2009.

THEOREM B ([8]). *Let $s = a + ib$, $a > 0$, $b \in \mathbf{R}$ and $f \in \mathcal{A}$. Assume that for a constant $c \in \mathbf{C}$ and all $z \in \mathbf{D}$*

$$(2) \quad \left| c|z|^2 + s - a(1 - |z|^2) \left\{ s \left(1 + \frac{zf''(z)}{f'(z)} \right) + (1 - s) \frac{zf'(z)}{f(z)} \right\} \right| \leq M$$

with

$$M = \begin{cases} a|s| + (a - 1)|s + c|, & \text{if } 0 < a \leq 1, \\ |s|, & \text{if } 1 < a, \end{cases}$$

then f is univalent in \mathbf{D} .

The case $s = 1$ with c replaced by $-1 - c$ is the special case of Theorem A.

The purpose of this paper is to refine Ruscheweyh's univalence condition to a quasiconformal extension criterion which includes Theorem A;

THEOREM 1. *Let $s = a + ib$, $a > 0$, $b \in \mathbf{R}$, $k \in [0, 1)$ and $f \in \mathcal{A}$. Assume that for a constant $c \in \mathbf{C}$ and all $z \in \mathbf{D}$*

$$(3) \quad \left| c|z|^2 + s - a(1 - |z|^2) \left\{ s \left(1 + \frac{zf''(z)}{f'(z)} \right) + (1 - s) \frac{zf'(z)}{f(z)} \right\} \right| \leq M$$

with

$$M = \begin{cases} ak|s| + (a - 1)|s + c|, & \text{if } 0 < a \leq 1, \\ k|s|, & \text{if } 1 < a, \end{cases}$$

then f has an l -quasiconformal extension to \mathbf{C} , where

$$(4) \quad l = \frac{2ka + (1 - k^2)|b|}{(1 + k^2)a + (1 - k^2)|s|} < 1.$$

Remark 1.1. If $f \in \mathcal{A}$, then it is easy to verify that there exists a sequence $\{z_n\} \subset \mathbf{D}$ with $|z_n| \rightarrow 1$ such that for each $s \in \{z \in \mathbf{C} : \operatorname{Re} z > 0\}$

$$\sup_n \left| s \left(1 + \frac{z_n f''(z_n)}{f'(z_n)} \right) + (1 - s) \frac{z_n f'(z_n)}{f(z_n)} \right| < \infty$$

which shows that (3) implies the inequality

$$(5) \quad |c + s| \leq M.$$

This inequality is needed for proving that $f(z)$ has no zeros in $0 < |z| < 1$ (see Lemma 7). In [8], it is mentioned that (3) implies $f(z) \neq 0$, $0 < |z| < 1$, without proof. The part of (5) can be found in [8].

Remark 1.2. A similar argument to Remark 1.1 is also valid for Theorem A. It follows that the assumption $|c| \leq k$ is embedded in the inequality (1).

The next application follows from Theorem 1. Let $\alpha > 0$ and $\beta \in \mathbf{R}$. It follows from a result of Sheil-Small [9, Theorem 2] that

$$(6) \quad \operatorname{Re} \left\{ 1 + \frac{zf''(z)}{f'(z)} + (\alpha + i\beta - 1) \frac{zf'(z)}{f(z)} \right\} > 0 \quad (z \in \mathbf{D})$$

is sufficient for $f \in \mathcal{A}$ to be a Bazilevič function of type $(\alpha, \beta)^1$ (see also [5]). Here, a function $f \in \mathcal{A}$ is called *Bazilevič of type (α, β)* if

$$f(z) = \left[(\alpha + i\beta) \int_0^z g(\zeta)^\alpha h(\zeta) \zeta^{i\beta-1} d\zeta \right]^{1/(\alpha+i\beta)}$$

for a starlike univalent function $g \in \mathcal{A}$ and an analytic function h with $h(0) = 1$ satisfying $\operatorname{Re}(e^{i\lambda} h) > 0$ in \mathbf{D} for some $\lambda \in \mathbf{R}$. Together with this fact, the next theorem follows;

THEOREM 2. *Let $\alpha > 0$, $\beta \in \mathbf{R}$ and $k \in [0, 1)$. If $f \in \mathcal{A}$ satisfies*

$$(7) \quad \left| 1 + \frac{zf''(z)}{f'(z)} + (\alpha + i\beta - 1) \frac{zf'(z)}{f(z)} - \frac{\alpha^2 + \beta^2}{\alpha} \right| \leq M$$

for all $z \in \mathbf{D}$ with

$$M = \begin{cases} k & \text{if } \alpha < \alpha^2 + \beta^2, \\ k(\alpha^2 + \beta^2)/\alpha & \text{if } \alpha^2 + \beta^2 \leq \alpha, \end{cases}$$

then f is a Bazilevič function of type (α, β) and can be extended to a \tilde{k} -quasiconformal automorphism of \mathbf{C} , where

$$\tilde{k} = \frac{2k\alpha + (1 - k^2)|\beta|}{(1 + k^2)\alpha + (1 - k^2)\sqrt{\alpha^2 + \beta^2}}.$$

Next, we shall discuss quasiconformal extensibility of functions $g(z) = z + \frac{d}{z} + \dots$ analytic in \mathbf{D}^* .

THEOREM 3. *Let $s = a + ib$, $a \geq 1$, $b \in \mathbf{R}$ and $k \in [0, 1)$ which satisfies $|b/s| \leq k$. Let $g(\zeta) = \zeta + \frac{d}{\zeta} + \dots$ be analytic in \mathbf{D}^* and fulfill*

$$(8) \quad \left| ib + (1 - |\zeta|^2)a \left\{ (1 - s) \left(1 - \frac{\zeta g'(\zeta)}{g(\zeta)} \right) - s \frac{\zeta g''(\zeta)}{g'(\zeta)} \right\} \right| \leq ak|s| - |b|(a - 1)$$

¹The author would like to thank Professor Yong Chan Kim for this remark.

for all $\zeta \in \mathbf{D}^*$. Then g can be extended to an l -quasiconformal automorphism of $\hat{\mathbf{C}}$, where

$$l = \frac{2ka + (1 - k^2)|b|}{(1 + k^2)a + (1 - k^2)|s|}.$$

The case $k \rightarrow 1$ corresponds to a univalence criterion which is due to Ruscheweyh [8].

Theorem 3 yields the following corollary which gives a positive answer to an open problem proposed by Ruscheweyh [8], i.e., whether a function $g(\zeta) = \zeta + d/\zeta + \dots$ with $(|\zeta|^2 - 1)|1 + (\zeta f''(\zeta)/f'(\zeta)) - (\zeta f'(\zeta)/f(\zeta))| \leq k$ for all $\zeta \in \mathbf{D}^*$ admits a quasiconformal extension to \mathbf{C} ;

COROLLARY 4. Let $g(\zeta) = \zeta + \frac{d}{\zeta} + \dots$ be analytic in \mathbf{D}^* . If there exists $k \in [0, 1)$ such that

$$(|\zeta|^2 - 1) \left| 1 + \frac{\zeta g''(\zeta)}{g'(\zeta)} - \frac{\zeta g'(\zeta)}{g(\zeta)} \right| \leq k$$

for all $\zeta \in \mathbf{D}^*$, then g can be extended to a k -quasiconformal automorphism of $\hat{\mathbf{C}} - \{0\}$.

From the above corollary we have another extension criterion for analytic functions on \mathbf{D} ;

COROLLARY 5. Let $f \in \mathcal{A}$ with $f''(0) = 0$. If there exists $k \in [0, 1)$ such that

$$(1 - |z|^2) \left| 1 + \frac{zf''(z)}{f'(z)} - \frac{zf'(z)}{f(z)} \right| \leq k$$

for all $z \in \mathbf{D}$, then f can be extended to a k -quasiconformal automorphism of \mathbf{C} .

2. Preliminaries

Our investigations are based on the theory of Löwner chains. A function $f_t(z) = f(z, t) = a_1(t)z + \sum_{n=2}^{\infty} a_n(t)z^n$, $a_1(t) \neq 0$, defined on $\mathbf{D} \times [0, \infty)$ is called a *Löwner chain* if $f_t(z)$ is holomorphic and univalent in \mathbf{D} for each $t \in [0, \infty)$ and satisfies $f_s(\mathbf{D}) \subsetneq f_t(\mathbf{D})$ and $f(0, s) = f(0, t)$ for $0 \leq s \leq t < \infty$, and if $a_1(t)$ is locally absolutely continuous in $t \in [0, \infty)$ with $\lim_{t \rightarrow \infty} |a_1(t)| = \infty$. Then $f(z, t)$ is absolutely continuous in $t \in [0, \infty)$ for each $z \in \mathbf{D}$ and satisfies the *Löwner differential equation*

$$(9) \quad \dot{f}(z, t) = h(z, t)zf'(z, t)$$

for $z \in \mathbf{D}$ and almost every $t \in [0, \infty)$. Here, $\dot{f}(z, t) = \partial f(z, t)/\partial t$, $f'(z, t) = \partial f(z, t)/\partial z$ and $h(z, t)$ is a function measurable on $t \in [0, \infty)$, holomorphic in $|z| < 1$ and $\text{Re } h(z, t) > 0$ ([6]).

An interesting method connecting the theory of quasiconformal extensions with Löwner chains was obtained by Becker;

THEOREM C ([2], see also [4]). *Suppose that $f(z, t)$ is a Löwner chain for which $h(z, t)$ of (9) satisfies the condition*

$$\left| \frac{h(z, t) - 1}{h(z, t) + 1} \right| \leq k$$

for all $z \in \mathbf{D}$ and almost all $t \in [0, \infty)$. Then $f_t(z)$ admits a continuous extension to $\bar{\mathbf{D}}$ for each $t \geq 0$ and the map defined by

$$\hat{f}(re^{i\theta}) = \begin{cases} f(re^{i\theta}, 0) & \text{if } r < 1, \\ f(e^{i\theta}, \log r) & \text{if } r \geq 1, \end{cases}$$

is a k -quasiconformal extension of f_0 to \mathbf{C} .

3. Proof of Theorem 1

The proof is divided into two parts. The first part of the proof is based on [8].

(i) First we assume that $f(z)/z \neq 0$ for all $z \in \mathbf{D}$. Then we can define

$$f(z, t) = f(e^{-st}z) \left\{ 1 - \frac{a}{c}(e^{2t} - 1) \frac{e^{-st}zf'(e^{-st}z)}{f(e^{-st}z)} \right\}^s$$

and let

$$(10) \quad F(z, t) = f(z, t/|s|).$$

A straightforward calculation shows

$$(11) \quad h(z, t) = \frac{\dot{F}(z, t)}{zF'(z, t)} = \frac{s}{|s|} \cdot \frac{1 + P(e^{-st/|s|}z, t/|s|)}{1 - P(e^{-st/|s|}z, t/|s|)},$$

where

$$P(z, t) = \frac{c}{a}e^{-2t} + 1 + (e^{-2t} - 1)H_s(z)$$

and

$$H_s(z) = s \left(1 + \frac{zf''(z)}{f'(z)} \right) + (1 - s) \frac{zf'(z)}{f(z)}.$$

Since $h(z, t)$ is holomorphic in $z \in \mathbf{D}$ and measurable on $t \in [0, \infty)$, applying Theorem C to (11), we see that the condition

$$\left| \frac{s(1 + P(e^{-st/|s|}z, t/|s|)) - |s|(1 - P(e^{-st/|s|}z, t/|s|))}{s(1 + P(e^{-st/|s|}z, t/|s|)) + |s|(1 - P(e^{-st/|s|}z, t/|s|))} \right| \leq l$$

implies l -quasiconformal extensibility of $f(z)$. This is equivalent to

$$(12) \quad \left| P + \frac{(1+l^2)b}{(1+l^2)a + (1-l^2)|s|} i \right| \leq \frac{2l|s|}{(1+l^2)a + (1-l^2)|s|}.$$

Here, we shall prove the following Lemma;

LEMMA 6. *Under the assumption of Theorem 1, we have*

$$(13) \quad |aP(e^{-st/|s|}z, t/|s|) + ib| < k|s|$$

for $z \in \mathbf{D}$ and $t \in [0, \infty)$.

Proof. We have

$$|aP + ib| \leq m_1 + m_2$$

by triangle inequality, where

$$m_1 = (1 - e^{-2t/|s|}) \left| \frac{ce^{-2at/|s|} + s}{1 - e^{-2at/|s|}} - aH_s(e^{-st/|s|}z) \right|$$

and

$$m_2 = \left| (ce^{-2at/|s|} + s) \frac{1 - e^{-2t/|s|}}{1 - e^{-2at/|s|}} - (ce^{-2t/|s|} + s) \right|.$$

Then it is enough to show that $m_1 + m_2 < k|s|$. (3) implies

$$\left| \frac{c|e^{st/|s|}z|^2 + s}{1 - |e^{st/|s|}z|^2} - aH_s(e^{-st/|s|}z) \right| \leq \frac{M}{1 - |e^{st/|s|}z|^2} \leq \frac{M}{1 - e^{-2at/|s|}}$$

for $z \in \mathbf{D}$. Let $q(t) = (1 - e^{-2t/|s|}) / (1 - e^{-2at/|s|})$. Applying the maximum modulus principle to the function

$$\frac{ce^{-2at/|s|} + s}{1 - e^{-2at/|s|}} - aH_s(e^{-st/|s|}z)$$

we have

$$m_1 \leq q(t)M.$$

On the other hand

$$m_2 \leq |c + s| |1 - q(t)|.$$

Since $1 \leq q(t) < 1/a$ if $0 < a \leq 1$ and $1/a < q(t) \leq 1$ if $1 < a$ for all $t \in [0, \infty)$, we conclude that $m_1 + m_2 < k|s|$ which is our desired inequality. \square

We now let Δ and Δ' be disks which are defined by replacing P in (12) and (13) to a complex variable w . It remains to find the smallest l so that Δ' is

contained by Δ . Note that if $k = l = 1$ then these two disks coincide. The following condition is necessary and sufficient for $\Delta' \subset \Delta$;

$$(14) \quad \left| \frac{(1 + l^2)b}{(1 + l^2)a + (1 - l^2)|s|} - \frac{b}{a} \right| \leq \frac{2l|s|}{(1 + l^2)a + (1 - l^2)|s|} - \frac{k|s|}{a}.$$

Then we conclude

$$l \leq \frac{2ka + (1 - k^2)|b|}{(1 + k^2)a + (1 - k^2)\sqrt{a^2 + b^2}}.$$

which is suitable for our purpose.

(ii) In order to eliminate the additional assumption that $f(z)/z \neq 0$ in \mathbf{D} , we need a sort of stability of the condition (3);

LEMMA 7. *If $f \in \mathcal{A}$ satisfies the assumption of Theorem 1, then so does $f_r(z) = \frac{1}{r}f(rz)$, $r \in (0, 1)$.*

Proof. It follows from the assumption that $aH_s(rz)$ is contained in the disk

$$\Delta = \left\{ w \in \mathbf{C} : \left| w - \frac{cr^2|z|^2 + s}{1 - r^2|z|^2} \right| \leq \frac{M}{1 - r^2|z|^2} \right\}.$$

We want to deduce that $aH_s(rz)$ lies in the disk

$$\Delta' = \left\{ w \in \mathbf{C} : \left| w - \frac{c|z|^2 + s}{1 - |z|^2} \right| \leq \frac{M}{1 - |z|^2} \right\}.$$

Therefore it is enough to see that $\Delta \subset \Delta'$, that is,

$$(15) \quad \left| \frac{c|z|^2 + s}{1 - |z|^2} - \frac{cr^2|z|^2 + s}{1 - r^2|z|^2} \right| \leq \frac{M}{1 - |z|^2} - \frac{M}{1 - r^2|z|^2}.$$

In view of the identity

$$\frac{|z|^2}{1 - |z|^2} - \frac{r^2|z|^2}{1 - |z|^2} = \frac{1}{1 - |z|^2} - \frac{1}{1 - r^2|z|^2},$$

the inequality (15) is equivalent to (5). □

Now we shall show that the condition $f(z)/z \neq 0$ in \mathbf{D} follows from the assumption of Theorem 1. Suppose, to the contrary, that $f(z_0) = 0$ for some $0 < |z_0| < 1$. We may assume that $f(z) \neq 0$ for $0 < |z| < |z_0|$. Then by Lemma 7 we can apply Theorem 1 to the function $f_{r_0}(z) = f(r_0z)/r_0$, $r_0 = |z_0|$ to conclude that f_{r_0} has a quasiconformal extension to \mathbf{C} . In particular, f_{r_0} is injective on $\bar{\mathbf{D}}$. This, however, contradicts the relation $f_{r_0}(z_0/r_0) = f_{r_0}(0) = 0$. □

Remark 3.1. We can replace $|s|$ in (10) to any positive real value and continue our argument. However, it will be found that $|s|$ gives the smallest l by calculations.

Remark 3.2. We have $l \geq k$, where $l = k$ if and only if $b = 0$. Indeed, let $l = l(k)$. Then we have $l'(k) > 0$ and $l''(k) \leq 0$ which imply $l \geq k$. If we suppose $l = k \neq 0$, then the right-hand side of (14) is greater than or equal to 0 only if $b = 0$. In the case $l = k = 0$ we also have $b = 0$ by (14). It easily follows from (4) that $l = k$ if $b = 0$.

4. Proof of Theorem 2

It is easy to see from (6) that f is a Bazilevič function of type (α, β) under our assumption since M is always less than or equal to $(\alpha^2 + \beta^2)/\alpha$.

Let us now prove quasiconformal extensibility of f . Setting $1/s = \alpha + i\beta$ which implies $a = \operatorname{Re} s = \alpha/(\alpha^2 + \beta^2)$ and $b = \operatorname{Im} s = -\beta/(\alpha^2 + \beta^2)$, (7) turns to

$$\left| 1 + \frac{zf''(z)}{f'(z)} + \left(\frac{1}{s} - 1\right) \frac{zf'(z)}{f(z)} - \frac{1}{a} \right| \leq \begin{cases} k, & 0 < a < 1, \\ k/a, & 1 \leq a. \end{cases}$$

Therefore, Theorem 2 follows from Theorem 1 with $c = -s$. □

5. Proof of Theorem 3

First let $s \neq 1$. In that case we may assume $g(\zeta) \neq 0$ for all $\zeta \in \mathbf{D}^*$ because of a similar discussion of the proof of Theorem 1;

LEMMA 8. *Let $g(\zeta) = \zeta + \frac{d}{\zeta} + \dots$ be analytic in \mathbf{D}^* . If g satisfies the same assumption of Theorem 3, then so does $g_R(\zeta) = \frac{1}{R}f(R\zeta)$, $R > 1$.*

Proof. We need to prove

$$\left| \frac{ib}{|\zeta|^2 - 1} - aG_s(R\zeta) \right| \leq \frac{ak|s| - |b|(a - 1)}{|\zeta|^2 - 1}$$

by using

$$\left| \frac{ib}{R^2|\zeta|^2 - 1} - aG_s(R\zeta) \right| \leq \frac{ak|s| - |b|(a - 1)}{R^2|\zeta|^2 - 1},$$

where

$$G_s(\zeta) = (1 - s) \left(\frac{\zeta g'(\zeta)}{g(\zeta)} - 1 \right) + s \frac{\zeta g''(\zeta)}{g'(\zeta)}.$$

In a similar way to the proof of Lemma 7, it suffices to see that

$$\left| \frac{ib}{|\zeta|^2 - 1} - \frac{ib}{R^2|\zeta|^2 - 1} \right| \leq \frac{ak|s| - |b|(a - 1)}{|\zeta|^2 - 1} - \frac{ak|s| - |b|(a - 1)}{R^2|\zeta|^2 - 1}.$$

This is equivalent to $|b| \leq k|s|$. □

Then we let

$$f(1/\zeta, t) = \frac{1}{g(e^{st}\zeta)} \left\{ 1 - (1 - e^{-2t})e^{st}\zeta \frac{g'(e^{st}\zeta)}{g(e^{st}\zeta)} \right\}^{-s}$$

and

$$F(1/\zeta, t) = f(1/\zeta, t/|s|).$$

Since

$$h(1/\zeta, t) = \frac{\dot{F}(1/\zeta, t)}{(1/\zeta)F'(1/\zeta, t)} = \frac{s}{|s|} \cdot \frac{1 + P(e^{st/|s|}\zeta, t/|s|)}{1 - P(e^{st/|s|}\zeta, t/|s|)}$$

where

$$P(\zeta, t) = (e^{2t/|s|} - 1)G_s(\zeta),$$

it is sufficient to see that

$$(16) \quad |aP(e^{st/|s|}\zeta, t/|s|) + ib| < k|s|$$

for all $\zeta \in \mathbf{D}^*$ and $t \in [0, \infty)$ under the assumption of the theorem. By triangle inequality we have

$$|aP + ib| \leq \left| \frac{1 - e^{2t/|s|}}{1 - e^{2at/|s|}} (ib + (1 - e^{2at/|s|})aG_s(e^{st/|s|}\zeta)) \right| + \left| ib \left(1 - \frac{1 - e^{2t/|s|}}{1 - e^{2at/|s|}} \right) \right|$$

for $\zeta \in \mathbf{D}^*$ and $t \in [0, \infty)$. Following the lines of the proof of Lemma 6, one can obtain that (8) implies (16). Therefore, a similar argument of the proof of Theorem 1 implies our assertion. The case $s = 1$ follows from a theorem of Becker [2]. □

6. Proof of Corollary 4 and 5

Proof of Corollary 4. Let $R > 1$ be an arbitrary but fixed number. We would like to show that $g_R(\zeta) = g(R\zeta)/R$ can be extended to a k -quasiconformal mapping of $\hat{\mathbf{C}} - \{0\}$. Since $g(\zeta) \neq 0$ in $\zeta \in \mathbf{D}^*$ from the assumption, there exists a certain constant A such that

$$(|\zeta|^2 - 1) \left| 1 - \frac{\zeta g'_R(\zeta)}{g_R(\zeta)} \right| \leq A < \infty$$

for all $\zeta \in \overline{\mathbf{D}}^*$. We also have

$$\left| 1 - \frac{\zeta g'_R(\zeta)}{g_R(\zeta)} + \frac{\zeta g''_R(\zeta)}{g'_R(\zeta)} \right| \leq \frac{k}{|\zeta R|^2 - 1}$$

for $\zeta \in \mathbf{D}^*$. Thus we obtain with $s = R^2 A/k(R^2 - 1)$

$$(|\zeta|^2 - 1) \left| \frac{1}{s} \left(1 - \frac{\zeta g'_R(\zeta)}{g_R(\zeta)} \right) - 1 - \frac{\zeta g''_R(\zeta)}{g'_R(\zeta)} + \frac{\zeta g'_R(\zeta)}{g_R(\zeta)} \right| \leq \frac{A}{s} + k \frac{|\zeta|^2 - 1}{|\zeta R|^2 - 1} \leq k$$

which implies quasiconformal extensibility of g_R by Theorem 3. A limiting procedure proves Corollary 4. □

Proof of Corollary 5. Note that the function $1 + (zf''(z)/f'(z)) - (zf'(z)/f(z))$ is analytic in \mathbf{D} and has a zero of order 2 at the origin by the condition $f''(0) = 0$. Thus, we obtain from the assumption that

$$\frac{1}{|z|^2} (1 - |z|^2) \left| 1 + \frac{zf''(z)}{f'(z)} - \frac{zf'(z)}{f(z)} \right| \leq k$$

by the maximum modulus principle. Let $g(\zeta)$ be a function defined by

$$g(\zeta) = \frac{1}{f(z)}$$

where $\zeta = 1/z$. Then g is analytic in \mathbf{D}^* and has the form $g(\zeta) = \zeta + d/\zeta + \dots$. From the relations

$$\frac{zf'(z)}{f(z)} = \frac{\zeta g'(\zeta)}{g(\zeta)}$$

and

$$1 + \frac{zf''(z)}{f'(z)} = -1 - \frac{\zeta g''(\zeta)}{g'(\zeta)} + 2 \frac{\zeta g'(\zeta)}{g(\zeta)},$$

we can deduce our assertion by applying Corollary 4. □

Acknowledgement. The author expresses his deep gratitude to Professor Toshiyuki Sugawa for many helpful discussions and his continuous encouragement during this work.

REFERENCES

- [1] L. V. AHLFORS, Sufficient conditions for quasiconformal extension, *Discontinuous groups and Riemann surfaces* **79**, Princeton Univ. Press, 1974, 23–29.
- [2] J. BECKER, Löwnersche Differentialgleichung und quasikonform fortsetzbare schlichte Funktionen, *J. Reine Angew. Math.* **255** (1972), 23–43.

- [3] J. BECKER, Über die Lösungsstruktur einer Differentialgleichung in der konformen Abbildung, *J. Reine Angew. Math.* **285** (1976), 66–74.
- [4] J. BECKER, Conformal mappings with quasiconformal extensions, *Aspects of contemporary complex analysis*, Academic Press, London, 1980, 37–77.
- [5] Y. C. KIM AND T. SUGAWA, A note on Bazilevič functions, *Taiwanese J. Math.* **13** (2009), 1489–1495, MR MR2554472.
- [6] CH. POMMERENKE, Über die Subordination analytischer Funktionen, *J. Reine Angew. Math.* **218** (1965), 159–173.
- [7] CH. POMMERENKE, *Univalent functions*, Vandenhoeck & Ruprecht, Göttingen, 1975.
- [8] S. RUSCHEWEYH, An extension of Becker’s univalence condition, *Math. Ann.* **220** (1976), 285–290.
- [9] T. SHEIL-SMALL, On Bazilevič functions, *Quart. J. Math. Oxford Ser. (2)* **23** (1972), 135–142.

Ikkei Hotta
DIVISION OF MATHEMATICS
GRADUATE SCHOOL OF INFORMATION SCIENCES
TOHOKU UNIVERSITY
6-3-09 ARAMAKI-AZA-AOBA
AOBA-KU, SENDAI 980-8579
JAPAN
E-mail: ikkeihotta@gmail.com